BUB-1 and CENP-C recruit PLK-1 to Control Chromosome Alignment and Segregation

During Meiosis I in *C. elegans* Oocytes

Samuel J.P. Taylor[1,a], Laura Bel Borja[1,a], Flavie Soubigou[1], Dhanya K. Cheerambathur[2], and

Federico Pelisch[1,*]

[1] Centre for Gene Regulation and Expression, Sir James Black Centre, School of Life

Sciences, University of Dundee. Dundee, DD1 5EH, United Kingdom

[2] Wellcome Centre for Cell Biology & Institute of Cell Biology, School of Biological Sciences, The

University of Edinburgh, Edinburgh, EH9 3BF, UK

[a] These authors contributed equally

* Corresponding author: Federico Pelisch (f.pelisch@dundee.ac.uk)

ABSTRACT

Phosphorylation is a key post-translational modification that is utilised in many biological processes for the rapid and reversible regulation of protein localisation and activity. Polo-like kinase 1 (PLK-1) is essential for both mitotic and meiotic cell divisions, with key functions being conserved in eukaryotes. The roles and regulation of PLK-1 during mitosis have been well characterised. However, the discrete roles and regulation of PLK-1 during meiosis have remained obscure. Here, we used *Caenorhabditis elegans (C. elegans)* oocytes to show that PLK-1 plays distinct roles in meiotic spindle assembly/stability, chromosome alignment and segregation, and polar body extrusion during meiosis I. Furthermore, by a combination of live imaging and biochemical analysis we identified the chromosomal recruitment mechanisms of PLK-1 during *C. elegans* oocyte meiosis. The spindle assembly checkpoint kinase BUB-1 directly recruits PLK-1 to the kinetochore and midbivalent while the chromosome arm population of PLK-1 depends on a direct interaction with the centromeric-associated protein CENP-C^{HCP-4}. We found that perturbing both BUB-1 and CENP-C^{HCP-4} recruitment of PLK-1 leads to severe meiotic defects, resulting in highly aneuploid oocytes. Overall, our results shed light on the roles played by PLK-1 during oocyte meiosis and provide a mechanistic understanding of PLK-1 targeting to meiotic chromosomes.

INTRODUCTION

Meiosis consists of two consecutive segregation events following DNA replication – in

meiosis I homologous chromosomes segregate, half of which are then removed in a polar

body, before the remaining sister chromatids segregate in meiosis II to produce haploid

gametes (Marston and Amon, 2004; Ohkura, 2015). Tight spatial and temporal control of

protein localisation and activity is required to ensure chromosome/chromatid alignment and

segregation occur efficiently and the correct number of chromosomes is present in each

gamete (Marston and Wassmann, 2017). Phosphorylation is a key post-translational

modification utilised to regulate protein localisation and activity, which is fundamentally

important for the success of both mitotic and meiotic cell divisions (Marston and Wassmann,

2017; Saurin, 2018). Therefore, the regulation of kinase or phosphatase localisation and

activity is vital for proper cell division, with the balance of their effects determining the

localisation/activity of substrate proteins that play important roles in the cell division process

(Gelens et al., 2018; Novak et al., 2010).

Polo-like kinases (PLKs) are a family of Ser/Thr protein kinases first discovered in

Drosophila (Llamazares et al., 1991; Sunkel and Glover, 1988) and yeast (Kitada et al., 1993;

Ohkura et al., 1995) and later findings showed that PLKs are present in all eukaryotes

(Zitouni et al., 2014). PLK1 is essential for meiotic and mitotic cell divisions, with its

localisation and functions proving well conserved throughout eukaryotic evolution. PLK-1

localises to the centrosomes, kinetochore, central spindle, and midbody during mitosis and is

involved in numerous processes including mitotic entry, spindle assembly, chromosome

alignment, the spindle assembly checkpoint, and cytokinesis (Archambault and Glover, 2009;

Petronczki et al., 2008; Schmucker and Sumara, 2014; Zitouni et al., 2014). During

mammalian meiosis, PLK-1 localises to the chromosomes and spindle poles in prometaphase

and metaphase. In anaphase PLK-1 is localised primarily in the central spindle between the

segregating chromosomes and then the midbody during polar body extrusion (Pahlavan et al.,

2000; Solc et al., 2015; Tong et al., 2002; Wianny et al., 1998). Inhibition of PLK-1 in

mammalian oocytes has indicated roles in many processes including germinal vesicle

breakdown, spindle assembly, chromosome alignment, and polar body extrusion (Solc et al.,

2015; Tong et al., 2002). To understand why PLK-1 is crucial for meiosis it is critical to

characterise precisely where PLK-1 is located and how it is recruited to specific regions,

which will allow further dissection of the discrete roles of PLK-1 during meiosis.

PLK-1 interacts with proteins via its C-terminal polo-binding domain (PBD) (Cheng, 2003;

Elia et al., 2003a). The PBD binds to phosphorylated motifs of the consensus Ser-

phSer/phThr-X, where ph indicates a phosphorylated residue and X indicates any amino acid

(Elia et al., 2003a, 2003b). When X is a proline, phosphorylation of the central Ser/Thr

residue is often mediated by a proline-directed kinase, notably Cdk1:Cyclin B during cell

division (Elowe et al., 2007; Qi et al., 2006) - this motif will henceforth be termed STP motif.

When a non-proline residue occupies position X, PLK-1 itself can phosphorylate the central

Ser/Thr, thereby enhancing its own recruitment (Kang et al., 2006; Neef et al., 2003), referred

to as self-priming. Furthermore, PLK-1 binding to STP motifs via the PBD induces a

conformational change that enhances its kinase activity (Mundt et al., 1997; Xu et al., 2013).

While the mechanism of PLK-1 recruitment to the chromosomes during oocyte meiosis has

not been characterised, recruitment of PLK-1 to the kinetochore during mammalian mitosis

has been investigated. In mammals, the constitutive centromere-associated network (CCAN)

complex of proteins binds to the histone variant CENP-A at centromeres (Foltz et al., 2006;

Izuta et al., 2006; Okada et al., 2006). Outer kinetochore proteins bind to the CCAN and

ultimately mediate chromosome alignment and segregation via interaction with microtubules

(Musacchio and Desai, 2017). Two proteins are primarily responsible for PLK-1 recruitment

to the kinetochore – CCAN component CENP-U (Kang et al., 2011, 2006; Singh et al., 2021)

and the spindle assembly checkpoint kinase BUB-1 (Elowe et al., 2007; Qi et al., 2006; Singh

et al., 2021), both of which directly bind to PLK-1 via STP motifs in a Cdk1-dependent

manner.

In *C. elegans*, PLK-1 localisation in meiosis and mitosis is similar to other organisms (Chase

et al., 2000). In mitosis, roles of PLK-1 include nuclear envelope breakdown (NEBD)

(Martino et al., 2017), merge of parental genomes in the embryo through lamina disassembly

(Rahman et al., 2015; Velez-Aguilera et al., 2020), centrosome maturation (Cabral et al.,

2019; Decker et al., 2011; Ohta et al., 2021; Woodruff et al., 2015), and cytokinesis (Gómez-

Cavazos et al., 2020) – indicating that major mitotic roles of PLK-1 are conserved in *C.*

elegans. However, meiotic roles of PLK-1 in *C. elegans* oocytes have remained obscure, as

PLK-1 depletion results in severely defective NEBD and oocytes with a whole nucleus rather

than condensed chromosomes (Chase et al., 2000). In the PLK-1-depleted oocytes that

'escaped' the NEBD defect, chromosome congression, segregation, and polar body extrusion

were severely disrupted. However it is unclear whether these phenotypes are indirect effects

of the severe early meiotic defects or whether they result from specific functions of PLK-1

throughout meiosis (Chase et al., 2000). Furthermore, while PLK-1 was shown to localise

broadly to chromosomes and the spindle during meiosis (Chase et al., 2000), a more precise

dynamic characterisation of PLK-1 localisation during meiosis is lacking.

Here, by temporally inhibiting an analogue-sensitive PLK-1 mutant we show that PLK-1 is

involved in spindle assembly/stability, chromosome alignment and segregation, and polar

body extrusion in *C. elegans* oocytes. Using live imaging and immunofluorescence, we find

that PLK-1 localises to the spindle poles, chromosome arms, kinetochores, and midbivalent

region between the homologous chromosomes during meiosis I in *C. elegans* oocytes. By a

combination of live imaging and *in vitro* biochemical analysis, we have characterised the full

chromosomal recruitment mechanisms of PLK-1 during meiosis – showing that CENP-C[HCP-4]

directly recruits PLK-1 to the chromosome arms while PLK-1 recruitment to the midbivalent

and kinetochore is mediated by a direct interaction with BUB-1. Furthermore, BUB-1- and

CENP-C-mediated PLK-1 recruitment to chromosomes is essential for meiosis I.

RESULTS

PLK-1 plays roles in spindle stability, chromosome alignment, segregation, and polar

body extrusion in meiosis I

A previous study showed that PLK-1 localises to the meiotic spindle and chromosomes in *C.*

elegans oocytes and depletion of PLK-1 using RNAi led to several defects including defective

NEBD, chromosome segregation, and polar body extrusion (Chase et al., 2000). While this

suggested that PLK-1 plays several roles during meiosis, the use of RNAi presents a

limitation to addressing them independently. In particular, the strong NEBD defect

complicates delineation of the roles of PLK-1 at later stages of meiosis. Therefore, we sought

to understand the distinct localisation and roles of PLK-1 during meiosis I.

To assess PLK-1 localisation during meiosis with high spatial and temporal resolution, we

imaged endogenously tagged sfGFP::PLK-1 in dissected oocytes. PLK-1 localises to the

spindle poles (Figure 1A-C; blue arrows), chromosome arms (Figure 1A-C; yellow arrows),

and midbivalent region between the homologous chromosomes during Prometaphase I

(Figure 1A-C; magenta arrow). This localisation pattern was confirmed with immunostaining

of fixed oocytes using a specific anti-PLK-1 antibody (Figure 1D). As chromosomes begin to

segregate in early anaphase, PLK-1 is mostly observed on chromosomes and, to a lesser

extent, in between the segregating chromosome masses (Figure 1B, yellow and magenta

arrows, respectively; see also Figure S1). During late anaphase, PLK-1 is still detectable on

chromosomes, and is enriched in the central spindle (Figure 1B, green arrow; see also Figure

S1).

Since long-term depletion of PLK-1 leads to severe NEBD defects (Chase et al., 2000, Figure

1E), we used an analogue-sensitive *plk-1* allele (Gómez-Cavazos et al., 2020; Woodruff et al.,

2015) that renders it sensitive to chemically modified derivatives of PP1, a Src family

inhibitor (Bishop et al., 2000). We reasoned that acute PLK-1[as] inhibition for a short period of

time would allow us to study the post-NEBD effects (Figure 1E). We tested a variety of

analogues and all of them led to embryonic lethality of the *plk-1^as* strain without affecting a

wild type strain (Figure S2). We decided to continue our experiments with the 3-substituted

benzyl PP1 derivative 3IB-PP1, which showed the best specificity and potency (Figure S2). A

wild type strain in the presence of 3IB-PP1 and *plk-1^as* in the presence of vehicle control

('EtOH') behaved normally during meiosis (Figure 1F,G & Figure S3). Addition of 10 µM

3IB-PP1 between 5 and 15 min before dissection and imaging of oocytes allowed us to bypass

the NEBD defect and 6 bivalents were easily identifiable within the newly fertilised oocyte

(Figure 1F, yellow arrows). Under these conditions PLK-1^as inhibition led to drastic spindle

defects with no observable bipolar spindle formation and no consequent chromosome

segregation was observed, indicating that PLK-1 is involved in spindle assembly and/or

stability during oocyte meiosis (Figure 1F). We sought to minimise the spindle defects upon

PLK-1 inhibition by reducing the concentration of 3IB-PP1 to 0.1-1 µM and omitting the pre-

treatment step prior to dissection. Under these conditions, ~62% of oocytes had seemingly

bipolar spindles and chromosomes remained somewhat associated with the spindle, although

chromosome alignment was still affected in 56% of oocytes (≥2 misaligned chromosomes,

Figure 1G,H). A more detailed analysis of chromosome dynamics after PLK-1 inhibition is

presented in Figure S3, where individual chromosomes are followed every 20 seconds and, as

opposed to wild type, chromosomes from PLK-1-inhibited oocytes show a highly dynamic

behaviour whereby they seem to briefly align and then become misaligned again (Figure

S3A,B and arrows therein). We then used the pole marker ASPM-1 to allow proper

characterisation of spindle bipolarity under these conditions and confirmed that even when

two ASPM-1 poles are clearly discerned (Figure 1I, blue arrows), chromosome alignment

fails (Figure 1I, yellow arrows). Hence, it appears that PLK-1 participates in chromosome

alignment in a manner that is at least partially independent of its roles in overall spindle

assembly/stability.

BUB-1 recruits PLK-1 to the midbivalent during oocyte meiosis

To further understand the role of PLK-1 during meiosis, we sought to identify the PLK-1

recruitment mechanism(s). In mammalian mitosis, the kinase BUB-1 and its paralog BUBR1

directly recruit PLK-1 to the kinetochore via STP motifs that are phosphorylated by Cdk1

(Elowe et al., 2007; Qi et al., 2006). While the *C. elegans* BUBR1 ortholog MAD3[SAN-1] does

not localise to the chromosomes or spindle during meiosis (Bel Borja et al., 2020), BUB-1

localises to the kinetochores and midbivalent region (Dumont et al., 2010; Monen et al., 2005;

Pelisch et al., 2019, 2017). We investigated whether BUB-1 was involved in PLK-1 targeting

during *C. elegans* meiosis. RNAi-mediated depletion of BUB-1 led to the loss of PLK-1 from

the midbivalent (Figure 2A,B; blue arrows). In contrast, PLK-1 signal at chromosome arms

remained unaffected and what appeared to be a pole signal was also detected (Figure 2A,B;

yellow and green arrows, respectively). Analysis of the BUB-1 protein sequence revealed a

putative polo-docking STP motif in amino-acids 526-528 that is conserved in nematode

species (Figure 2C). Therefore, we sought to identify whether *C. elegans* BUB-1 directly

interacts with PLK-1 to mediate its recruitment to the midbivalent region.

BUB-1 directly interacts with PLK-1 through a Cdk1-dependent STP motif

To test whether BUB-1 directly interacts with PLK-1 *in vitro*, we purified a recombinant

fragment of BUB-1 encompassing the intrinsically disordered region between the TPR and

kinase domains that contains the putative STP motif ('BUB-1[190-628]'). Since the interaction

between STP motifs and the PBD of PLK-1 requires phosphorylation of the central Ser/Thr

residue (Elia et al., 2003a, 2003b), we conducted kinase assays to assess the phosphorylation

of BUB-1$^{190-628}$. Cdk1 and PLK-1 can both phosphorylate BUB-1$^{190-628}$ (Figure S4A,B).

Interestingly, kinase assays conducted with Cdk1 and PLK-1 together produced a prominent

shifted band representing phosphorylated protein that was not present with the individual

kinases (Figure 2D). Since STP motifs are known to be targets of proline-directed kinases

such as Cdk1 (Elowe et al., 2007; Qi et al., 2006), we hypothesised that Cdk1 phosphorylates

the central threonine of the STP motif (T527), allowing PLK-1 to bind directly to BUB-1

resulting in the shifted band representing highly phosphorylated BUB-1. To test this

hypothesis, we mutated T527 in the STP motif to alanine ('BUB-1$^{190-628}$(T527A)') and

assessed the resulting phosphorylation using the phosphoprotein stain Pro-Q^{TM} Diamond.

T527A mutation in BUB-1$^{190-628}$ largely prevented the shift observed in the combined Cdk1

and PLK-1 assay (Figure 2E), indicating that phosphorylation of this residue is essential for

the shifted band observed when Cdk1 and PLK-1 both phosphorylate BUB-1$^{190-628}$.

To determine whether BUB-1 directly binds to PLK-1, we purified a maltose-binding protein

(MBP)-tagged PLK1 PBD (MBP-PLK1^{PBD}) (Singh et al., 2021) and incubated it with

unphosphorylated or Cdk1-phosphorylated BUB-1$^{190-628}$ before assessing complex formation

by size-exclusion chromatography (SEC). Cdk1-phosphorylated BUB-1 formed a stable

complex with MBP-PLK1^{PBD} (Figure 2F). While unphosphorylated BUB-1 showed some

interaction with MBP-PLK1^{PBD}, this complex eluted from the column at a higher volume and

bound to a lower proportion of the MBP-PLK1^{PBD}, suggestive of a weaker interaction and/or

different stoichiometry (Figure 2F). To directly test whether phosphorylation of T527 is

required for PLK-1 to bind to BUB-1, MBP-PLK1^{PBD} was incubated with Cdk1-

phosphorylated BUB-1$^{190-628}$ or BUB-1$^{190-628}$(T527A) and SEC was used to assess complex

formation again. Interestingly, when Cdk1-phosphorylated BUB-1$^{190-628}$(T527A) was

incubated with MBP-PLK1^{PBD}, the resulting elution was reminiscent of the unphosphorylated

wild type BUB-1$^{190-628}$ (Figures 2G). Together, these data indicate that there is a Cdk1

phosphorylation-dependent interaction between BUB-1 and PLK1^{PBD} that requires

phosphorylation of T527 within the STP motif of BUB-1. To further confirm that the STP

motif of BUB-1 can interact with the PBD in a T527 phosphorylation-dependent manner,

fluorescence polarisation assays were conducted using FITC-labelled peptides containing the

BUB-1 STP motif (Figure 2H). The phosphorylated STP motif bound to MBP-PLK1^{PBD} with

high affinity (K_D below 100 nM) while the unphosphorylated peptide did not interact with

MBP-PLK1^{PBD} at the concentrations tested (Figure 2H).

Altogether, these data indicate that *C. elegans* BUB-1 can directly bind to PLK-1 *in vitro* in a

phospho-dependent manner via a newly characterised STP motif.

BUB-1 directly recruits PLK-1 to the midbivalent in vivo

We then sought to determine whether the STP motif in BUB-1 is responsible for PLK-1

recruitment *in vivo.* Using CRISPR-Cas9, we generated the T527A mutation in the

endogenous *bub-1* locus (*bub-1^{T527A}*). *bub-1^{T527A}* mutant worms showed significant embryonic

and larval lethality so we generated a balanced strain in which the *bub-1^{T527A}* allele was

maintained as a heterozygote. Homozygous *bub-1^{T527A}* worms from heterozygous parents

develop to adulthood and produce oocytes that go through meiosis, which allowed us to study

the role of the STP motif in BUB-1 during meiosis. PLK-1 was absent from the midbivalent

in *bub-1^{T527A}* oocytes, reminiscent of the *bub-1(RNAi)* phenotype (Figure 3A, blue arrowheads

and 3B). Importantly, BUB-1 localisation to the midbivalent and kinetochore was maintained

in the *bub-1^{T527A}* strain (Figure 3C), indicating that BUB-1 directly interacts with PLK-1 via

this STP motif *in vivo* to recruit PLK-1 to the midbivalent. To assess the impact of BUB-1

mediated PLK-1 recruitment during meiosis I, we crossed the *bub-1^{T527A}* allele with a strain

expressing GFP-tagged tubulin and mCherry-tagged histone and analysed chromosome

alignment, segregation, and polar body extrusion defects (See Methods). *bub-1^{T527A}* mutant

oocytes displayed chromosome alignment defects in ~62% of the oocytes, with 32% of

oocytes showing severe alignment defects (Figure 3D,E). Additionally, ~1/4 of *bub-1[T527A]*

oocytes showed mild anaphase defects (Figure 3E). Despite these defects, more than 90% of

bub-1[T527A] oocytes show visible separation of two chromosome masses and polar body

extrusion occurred normally at the end of meiosis I (Figure 3E).

Overall, our results show that an STP motif in BUB-1 directly recruits PLK-1 to the

midbivalent during meiosis I and this interaction is primarily important for chromosome

alignment.

CENP-C[HCP-4] recruits PLK-1 to meiotic chromosome arms

PLK-1 localisation to chromosome arms remained unchanged when BUB-1 was depleted

(Figure 2A,B) or the STP motif was mutated (Figure 3A,B), indicating that a different

pathway is required to recruit this population of PLK-1. We therefore sought to identify the

mechanism of PLK-1 recruitment to the chromosome arms. In mammalian mitosis, PLK-1

recruitment to the kinetochore is mediated by BUB-1 and CCAN component CENP-U (Elowe

et al., 2007; Kang et al., 2011, 2006; Qi et al., 2006; Singh et al., 2021). Interestingly, the

CCAN appears to be largely absent in *C. elegans* (Maddox et al., 2012), and kinetochore

assembly depends on the CENP-C orthologue, HCP-4 (hereafter CENP-C[HCP-4]) during mitosis

(Oegema et al., 2001) and on CENP-C[HCP-4] and the nucleoporin ELYS[MEL-28] during meiosis

(Hattersley et al., 2022). CENP-C[HCP-4] localises to chromosomes throughout meiosis I (Figure

4A) (Hattersley et al., 2022; Monen et al., 2005). Like its mammalian counterpart, CENP-

C[HCP-4] is predicted to be mostly disordered and it contains a putative N-terminal STP motif

encompassing amino acids 162-164 that is conserved in nematode species (Figure 4B).

Although CENP-C[HCP-4] depletion does not have a major impact on meiosis I (Hattersley et al.,

2022; Monen et al., 2005), RNAi-mediated depletion of CENP-C[HCP-4] abolished PLK-1

localisation on chromosome arms (Figure 4C,D). PLK-1 is still present in the midbivalent

(Figure 4C,D, blue arrows) which suggests that BUB-1 and CENP-C^{HCP-4} represent

independent pathways for PLK-1 targeting. Additionally, CENP-C^{HCP-4} depletion revealed a

pool of PLK-1 which is kinetochore-associated (Figure 4C,D, yellow arrows). Since the

above data indicated that CENP-C^{HCP-4} is involved in PLK-1 recruitment to chromosome

arms, we sought to determine whether this involved a direct interaction via the putative STP

motif identified in sequence alignments.

CENP-C^{HCP-4} interacts directly with PLK-1 through a Cdk1-dependent STP motif

To investigate whether CENP-C^{HCP-4} directly interacts with PLK-1, we purified a recombinant

N-terminal fragment of CENP-C^{HCP-4} ('CENP-C$^{HCP-4(1-214)}$'). As the putative CENP-C^{HCP-4}

STP motif also contains a proline-directed kinase consensus site, we assessed Cdk1

phosphorylation of the recombinant fragment. Cdk1 kinase assays showed that CENP-C^{HCP-}

$^{4(1-214)}$ can be phosphorylated by Cdk1 and mutation of the putative STP motif threonine to

alanine (T163A) reduced the phosphorylation of the fragment (Figure 5A). To determine

whether this putative STP motif directly binds to PLK-1 in a Cdk1-dependent manner, we

used size exclusion chromatography. Cdk1-phosphorylated CENP-C$^{HCP-4(1-214)}$ forms a stable

complex with MBP-PLK1^{PBD}, while the unphosphorylated protein did not (Figure S5). We

then tested the requirement of T163 phosphorylation on CENP-C$^{HCP-4(1-214)}$ to interact with

MBP-PLK1^{PBD}. Phosphorylated wild type CENP-C$^{HCP-4(1-214)}$ forms a stable complex with

MBP-PLK1^{PBD}; on the contrary, phosphorylated CENP-C$^{HCP-4(1-214)}$ (T163A) did not form a

stable complex with MBP-PLK1^{PBD} (Figure 5B). Together, these data indicate that CENP-

C$^{HCP-4(1-214)}$ phosphorylated by Cdk1 can bind to the PBD and this interaction requires

phosphorylation of T163 within the STP motif. To further confirm this STP motif binds to

MBP-PLK-1^{PBD} in a phospho-dependent manner, we conducted fluorescence polarisation

assays with FITC-labelled CENP-C^{HCP-4} STP motif peptides. The Thr 163 phosphorylated

peptide bound to MBP-PLK1^{PBD} with high affinity (K_D in the low hundreds nM), while the

unphosphorylated peptide did not display binding at the concentrations indicated (Figure 5C).

Collectively, these data indicate that the putative STP motif in CENP-C^{HCP-4} binds to the

PBDs of PLK1 in a Cdk1 phosphorylation-dependent manner, which led us to assess the

importance of this STP motif *in vivo*.

CENP-C^{HCP-4} recruits PLK-1 to chromosome arms in vivo through an STP motif

The T163A mutation in CENP-C^{HCP-4} was generated in the endogenous *hcp-4* locus (*hcp-*

4^{T163A}) which, unlike *bub-1^{T527A}*, did not affect viability. When GFP:PLK-1 was monitored in

dissected oocytes, *hcp-4^{T163A}* recapitulated the full CENP-C^{HCP-4} depletion with PLK-1

localising only to the midbivalent and kinetochore but absent from chromosome arms (Figure

5D,E). This indicates that PLK-1 is targeted to chromosome arms directly through the

phospho-dependent STP motif in CENP-C^{HCP-4}. Importantly, CENP-C^{HCP-4}(T163A) displays

an indistinguishable localisation from wild type CENP-C^{HCP-4} (Figure 5F) and the other PLK-

1 receptor, BUB-1, also localises normally in the *hcp-4^{T163A}* strain (Figure 5G).

These data indicate that CENP-C^{HCP-4} directly recruits PLK-1 to the chromosome arms during

meiosis I via a newly characterised N-terminal STP motif.

Dual PLK-1 recruitment by BUB-1 and CENP-C is essential for meiosis I

Our results so far indicate that PLK-1 recruitment to the midbivalent during meiosis I is

mediated by direct interaction with BUB-1, while PLK-1 localisation to the chromosome

arms requires direct interaction with CENP-C^{HCP-4}. Co-depletion of BUB-1 and CENP-C^{HCP-4}

by RNAi led to complete absence of PLK-1 at the chromosome arms, midbivalent, and

kinetochore (Figure 6A). This confirmed that the BUB-1 and CENP-C^{HCP-4} pathways are the

primary recruiters of PLK-1 to the chromosomes during oocyte meiosis. Additionally, the

kinetochore population of PLK-1 was lost when BUB-1 and HCP-4 were co-depleted,

indicating that BUB-1 is responsible for recruiting PLK-1 to the kinetochore as well as the

midbivalent (Figure 6A). RNAi of BUB-1 causes significant defects in spindle stability,

chromosome alignment, and chromosome segregation during meiosis (Dumont et al., 2010;

Pelisch et al., 2019). However, co-depletion of BUB-1 and CENP-C^{HCP-4} exacerbated the

chromosome alignment and segregation errors and resulted in a significantly higher

proportion of polar body extrusion failures (Figure 6B-C). Therefore, while CENP-C^{HCP-4}

depletion does not have noticeable defects on its own, it enhances the BUB-1 depletion

phenotype. To focus more specifically on the effects of direct PLK-1 recruitment, we depleted

CENP-C^{HCP-4} in *bub-1^{T527A}* mutant oocytes. This abolished PLK-1 localisation in the same

manner as co-depletion of BUB-1 and CENP-C^{HCP-4} (Figure 6D) as well as displaying defects

in chromosome alignment, segregation, and polar body extrusion of a greater severity than

hcp-4(RNAi) or *bub-1^{T527A}* alone (Figure 6E,F). Similar results were obtained when

performing the complementary experiment using *hcp-4^{T163A}/bub-1(RNAi)* oocytes (Figure S6).

These data indicate that recruitment of PLK-1 to the midbivalent and kinetochore by BUB-1

appears to be primarily responsible for the chromosomal roles of PLK-1 during meiosis I, as

disruption of this pathway leads to meiotic defects while perturbing CENP-C^{HCP-4} recruitment

of PLK-1 on its own does not. However, the fact that disruption of both BUB-1 and CENP-

C^{HCP-4} recruitment pathways enhances the severity of the resulting meiotic defects indicates

that the CENP-C^{HPC-4} pathway does still play an active part in the roles of PLK-1 during

meiosis I.

DISCUSION

In the present manuscript we describe specific, post-NEBD roles played by PLK-1 during

oocyte meiosis. PLK-1 is important for spindle assembly/stability, chromosome alignment

and segregation, and polar body extrusion during meiosis I. We found that PLK-1 localises to

spindle poles and chromosomes during metaphase I before localising to the chromosomes and

central spindle in anaphase. Furthermore, we characterised the mechanisms of chromosomal

PLK-1 targeting during oocyte meiosis, which rely on the centromere-associated protein

CENP-C[HCP-4] and the spindle assembly checkpoint kinase BUB-1. While CENP-C[HCP-4]

targets PLK-1 to chromosome arms, BUB-1 directs PLK-1 to the midbivalent and

kinetochores. In both cases, interaction with PLK-1 relies on phosphorylated STP motifs

within predicted disordered regions. While we have not confirmed that these sites are

phosphorylated by Cdk1 *in vivo,* several lines of evidence indicate this is likely the case: 1)

both motifs have Pro at position 3, indicative of potential proline-directed kinase substrates;

2) Cdk1 is the most prominent proline-directed kinase in cell division and known to

phosphorylate STP motifs, including that of mammalian BUB-1 (Qi et al., 2006); 3) both sites

were phosphorylated by Cdk1, but not PLK-1, *in vitro*; and 4) fluorescence polarisation

experiments indicate that phosphorylation of the Thr residues within the STP motifs is

essential for the interaction with the PBD, clearly displaying why the alanine mutants have

such a drastic effect on PLK-1 localisation *in vivo*. We found that mutating the STP motif in

BUB-1 results primarily in chromosome alignment defects. While disrupting the CENP-C[HCP-]

[4]-mediated localisation of PLK-1 to the chromosome arms does not have a significant

phenotypic defect on its own, it does enhance the meiotic defects observed when the BUB-1-

dependent kinetochore and midbivalent populations are disrupted. This suggests that while

BUB-1 recruitment of PLK-1 may mediate the most important functions of PLK-1 during

meiosis I, CENP-C[HCP-4] recruitment of PLK-1 to the chromosome arms also plays an active

role in meiosis.

Expanding on the meiotic roles of PLK-1

By temporally inhibiting an analogue-sensitive PLK-1 mutant during *C. elegans* oocyte

meiosis, we have shown that PLK-1 plays a major role in the regulation of the meiotic

spindle. At high concentrations of analogue this resulted in the complete lack of spindle

bipolarity, and even at low concentrations the majority of oocytes (62%) imaged lacked an

apparent bipolar spindle. While this clearly displays a key role of PLK-1 during meiosis is in

the regulation of the meiotic spindle, we cannot distinguish more specific mechanisms using

our experimental techniques. As a result, we have characterised the phenotype as a lack of

spindle stability throughout this manuscript, but it should be noted that we cannot distinguish

whether these effects are on the spindle assembly process itself or on the maintenance of an

assembled bipolar spindle. Potential mechanisms of meiotic spindle regulation by PLK-1

include the microtubule depolymerase KLP-7, the *C. elegans* ortholog of the MCAK/Kinesin

13 family. There is some evidence to indicate that PLK-1 may regulate the kinesin 13

microtubule depolymerases in other organisms (Jang et al., 2009; Ritter et al., 2014; Sanhaji

et al., 2014; Shao et al., 2015; Zhang et al., 2011) and KLP-7 localises to the chromosomes,

during meiosis I (Connolly et al., 2015; Danlasky et al., 2020; Gigant et al., 2017; Han et al.,

2015), which overlaps with PLK-1 localisation. Furthermore, disrupting KLP-7 function

prevents proper bipolar spindle assembly and results in microtubules protruding out of the

meiotic spindle towards the cytoplasm (Connolly et al., 2015; Gigant et al., 2017), two

phenotypes we also see with PLK-1 inhibition.

Aside from the large-scale spindle defects observed when PLK-1 is inhibited, chromosome

alignment is still disrupted when the structure of the spindle appears bipolar and largely

normal. Since the mechanism of chromosome alignment by acentrosomal meiotic spindles is

not as well characterised as the centrosomal mitotic equivalent, speculating on the

underpinning mechanisms of this phenotype is challenging. However, it should be noted that

we cannot exclude the possibility that this phenotype is also a direct result of the dysregulated

spindle upon PLK-1 inhibition. Indeed, there is some evidence to suggest that a chromosome-

dependent pathway of microtubule formation may be an important aspect of chromosome

alignment and segregation (Conway et al., 2022; Heald et al., 1996; Kiewisz et al., 2022).

Despite these specific hypotheses mentioned above, it is clear that there will be many

different proteins and pathways impacted by PLK-1 phosphorylation throughout meiosis I

that will ultimately contribute to the severe defects we observe upon PLK-1 inhibition. While

a focussed investigation into specific hypotheses would no doubt yield important results, an

unbiased approach to identify the relevant PLK-1 substrates during meiosis would be

particularly useful for investigation of the key effects of PLK-1 during meiosis. There are

obvious technical challenges to overcome before this can be achieved, not least of which

would be isolating a large enough sample of meiotic oocytes to perform robust quantitative

proteomics. The work in this manuscript undertaken to identify the mechanisms of PLK-1

targeting during oocyte meiosis I will be instrumental for a later characterisation of the

localisation and meiotic stage-specific analysis of PLK-1 substrates.

Comparison with dual recruitment in mammals (CENP-U vs CENP-C)

During mammalian mitosis, PLK1 is recruited to kinetochores through BUB1 and CENP-U,

relying on self-priming in addition to Cdk1-mediated priming (Kang et al., 2011, 2006; Qi et

al., 2006; Singh et al., 2021). Our results suggest that Cdk1-mediated priming is the primary

mechanism for PLK-1 recruitment in both BUB-1- and CENP-C-dependent branches.

Additionally, we noted the presence of a putative B56 short linear motif (LxxIxE) 38 aa

downstream of the STP motif in *C. elegans* CENP-C (203-IPTILE-208). This is relevant

because it has been shown that PLK1 and B56 motifs tend to co-exist in close proximity

(Cordeiro et al., 2020; Singh et al., 2021). This makes a putative cross-talk between PLK-1

and PP2A/B56 a worthy avenue to follow-up on our current findings.

Apparent lack of a CCAN network along with retention of crucial roles for PLK1 in species

like *C. elegans* and *Drosophila melanogaster* (*D. melanogaster*) suggest that the CENP-U

pathway could have been replaced by other proteins. CENP-C is a good candidate, as the only

CCAN component in these species. While we confirm that this is the case in *C. elegans*, it is

interesting to note that putative STPs exist in sequences 176-178 and 266-268 in *D.*

melanogaster CENP-C (Uniprot #Q9VHP9). Interestingly, *D. melanogaster* Polo and CENP-

C co-localise and this co-localisation increases by ectopic centromere generation through

CENP-A (CID) over-expression (Heun et al., 2006).

Overall, our results advance our understanding on the roles played by PLK-1 during oocyte

meiosis and provide a mechanistic understanding of PLK-1 targeting to meiotic

chromosomes. The next step will be to identify and characterise PLK-1 meiotic substrates, to

understand exactly how PLK-1 participates in each of its meiotic roles.

ACKNOWLEDGEMENTS

We thank Arshad Desai for sharing strains and antibodies and for helpful discussions and sharing results prior to publication; Pierre Gönczy for the PLK-1 antibody; Lionel Pintard for providing PLK-1 for the initial experiments and the plasmid for expression and purification; Bruce Bowerman for the GFP::ASPM-1 strain; Satpal Virdee for advice on fluorescence polarisation experiments. We also thank Ron Hay for comments on the manuscript. This work was supported by a Career Development Award from the Medical Research Council (grant MR/R008574/1) and an ISSF grant funded by the Wellcome Trust (105606/Z/14/Z). ST is funded by a Medical Research Council Doctoral Training Programme. DKC is supported by a Sir Henry Dale Fellowship from the Wellcome Trust (208833). We acknowledge the FingerPrints Proteomics Facility and the Dundee Imaging Facility, which are supported by a 'Wellcome Trust Technology Platform' award (097945/B/11/Z) and the Tissue Imaging Facility, funded by a Wellcome Trust award (101468/Z/13/Z). Some nematode strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

AUTHOR CONTRIBUTIONS

AUTHOR CONTRIBUTIONS

S.J.P.T. designed, performed and analysed experiments, and wrote the manuscript.

L.B-B. designed, performed and analysed experiments.

F.S. designed, performed and analysed experiments.

D.C. generated novel reagents (nematode strains).

F.P. designed, performed and analysed experiments; acquired funding; wrote the manuscript.

FIGURE LEGENDS

Figure 1. Analysis of PLK-1 localisation and inhibition during meiosis I.

A) Schematic of a *C. elegans* bivalent, highlighting the midbivalent and kinetochore.

B) *In situ* GFP-tagged PLK-1 (Martino et al., 2017) was followed through meiosis I in

live, dissected oocytes. Scale bar, 2 μm. More detailed sequence of events are

displayed in Figure S1. Yellow arrows point to chromosomes, blue arrows indicate

spindle poles, magenta arrows point towards the midbivalent, and green arrow

indicates the central spindle.

C) Line profile analysis of PLK-1::GFP during early Metaphase I, as indicated by the

yellow line in B). Background signal was subtracted and maximum signal for each

channel was set to 1.

D) Immunostaining of (untagged) PLK-1 in fixed oocytes. The insets represent a

magnified image of single bivalents for each channel.

E) Schematic of the last two maturing oocytes and the fertilised egg, highlighting the

difference between PLK-1 depletion and acute PLK-1 inhibition.

F) *plk-1as* worms expressing GFP-tagged tubulin and mCherry-tagged histone were

dissected in medium containing ethanol ('EtOH', control) or the PP1 analogue 3IB-

PP1 (10 μM). Yellow arrows point to the each of the six bivalents. Scale bar, 2 μm.

G) *plk-1as* worms expressing GFP-tagged tubulin and mCherry-tagged histone, were

dissected in medium containing ethanol ('EtOH', control) or 0.1 μM PP1 analogue

3IB-PP1. Scale bar, 2 μm. The panels on the right show specific Z slices to highlight

individual chromosomes. Yellow arrows point to misaligned chromosomes contained

within the spindle, whereas blue arrows indicate chromosomes outside the spindle.

See also Figure S3.

H) Chromosme alignmend defects scored for 0.1-1 µM 3IB-PP1-treated oocytes are

presented in the graph.

I) Worms expressing GFP-tagged ASPM-1 (pole marker) and mCherry-tagged histone

along with analogue-sensitive *plk-1*, were dissected in medium containing ethanol

('EtOH', control) or the PP1 analogue 3IB-PP1 at 1 µM or 10 µM. Scale bar, 2 µm.

Early metaphase I spindles are shown for each condition.

Figure 2. BUB-1 regulates PLK-1 localisation in vivo and directly binds to PLK-1 *in*

***vitro*.**

A) Control (*'wild type'*) and BUB-1 depleted ['*bub-1(RNAi)'*] oocytes expressing sfGFP-

tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded throughout

meiosis I. Prometaphase/Metphase I is shown (before spindle rotation/shortening).

The yellow arrows indicate the bivalent chosen for magnification in each condition.

The blue arrows point to the midbivalent and the green arrow points to the spindle

pole. Left scale bar, 2 µm. Right scale bar, 1 µm.

B) Line profile analysis of PLK-1::GFP during early Metaphase I in *wild type* and *bub-*

1(RNAi) oocytes, as indicated by the yellow lines. Background signal was subtracted

and maximum signal for each channel was set to 1. Blue arrow points to the position

of the midbivalent.

C) Schematic representation of the *C. elegans* BUB-1 protein (top), sequence alignment

of the putative STP motif in nematode species (bottom).

D) Kinase assay of recombinant BUB-1$^{190-628}$ with Cdk1:Cyclin B, PLK-1, and both

kinases combined as indicated. Phosphorylation was assessed using SDS-PAGE

followed by coomassie (total protein) or ProQ diamond (phosphoprotein) staining.

E) Kinase assay of recombinant BUB-1$^{190-628}$ and BUB-1$^{190-628}$(T527A) with Cdk1:Cyclin

B and PLK-1 combined. Proteins were incubated with the kinases for the indicated

time points before phosphorylation was assessed by SDS-PAGE and staining with

either coomassie (total protein) or ProQ diamond (phosphoprotein).

F) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. BUB-1$^{190-628}$ was incubated with MBP-

PLK1^{PBD} at equimolar concentrations before separation by SEC. Binding was

conducted with unphosphorylated or Cdk1:Cyclin B phosphorylated BUB-1$^{190-628}$ as

indicated.

G) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. Wild type or T527A mutant BUB-1$^{190-628}$

was phosphorylated by Cdk1:Cyclin B before incubation with an equimolar

concentration of MBP-PLK1^{PBD}, binding was assessed by SEC.

H) FITC-labelled peptides containing the BUB-1 STP motif were incubated with

increasing concentrations of MBP:PLK1^{PBD} and binding analysed by fluoresence

polarisation. Unphosphorylated versus T527-phosphorylated peptides were compared.

'T^{Ph}' denotes phosphorylated Threonine.

Figure 3. The polo-docking site in BUB-1 is required for PLK-1 targeting and

chromosome alignment.

A) Fixed oocytes were stained with a PLK-1 specific antibody (green) and *bub-1^{T527A}*

heterozygote and homozygote oocytes were compared. DNA is shown in magenta.

The yellow arrow points to the midbivalent magnified on the right in each case and the

blue arrowhead points to the midbivalent.

B) Line profile analysis of PLK-1 localisation in fixed oocytes during early Metaphase I

in *bub-1^{T527A}* heterozygote ('*bub-1^{T527A}/+*') and homozygote ('*bub-1^{T527A}*') oocytes, as

indicated by the yellow lines. Background signal was subtracted and maximum signal

for each channel was set to 1.

C) Fixed oocytes were stained with a BUB-1 specific antibody (green) and *bub-1^{T527A}*

heterozygote and homozygote oocytes were compared. DNA is shown in magenta.

D) *bub-1^{T527A}* heterozygote ('*bub-1^{T527A}/+*') and homozygote ('*bub-1^{T527A}*') oocytes

expressing GFP-tagged tubulin and mCherry-tagged histone were filmed during

meiosis I. Two homozygote ('*bub-1^{T527A}*') oocytes are shown to depict the difference

in severity of the alignment defect. Scale bar, 2 μm.

E) Meiotic defects (as described in the Methods section) were assessed in wild type, *bub-*

1^{T527A} heterozygote ('*bub-1^{T527A}/+*') and homozygote oocytes. Representative images

of the different phenotypes analysed are presented on the right.

Figure 4. CENP-C^{HCP-4} is required for chromosomal PLK-1 targeting.

A) *In situ* GFP-tagged CENP-C^{HCP-4} was followed through meiosis I in live, dissected

oocytes. Scale bar, 2 μm.

B) Schematic representation of the *C. elegans* CENP-C^{HCP-4} (top), sequence alignment of

the putative STP motif in nematode species (bottom).

C) Control (*'wild type'*) and CENP-C^{HCP-4}-depleted ['*hcp-4(RNAi)*'] oocytes expressing

sfGFP-tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded

throughout meiosis I. Prometaphase/Metphase I is shown (before spindle

rotation/shortening). On the right, the blue arrows point to the midbivalent and the

yellow arrow points to the kinetochore. Left scale bar, 2 μm. Right scale bar, 1 μm.

D) Line profile analysis of PLK-1::GFP during early Metaphase I in *wild type* and *hcp-*

4(RNAi) oocytes, as indicated by the yellow lines. Background signal was subtracted

and maximum signal for each channel was set to 1. The blue arrows point to the

midbivalent and the yellow arrow points to the kinetochore

Figure 5. CENP-C^{HCP-4} interacts with PLK-1 in vitro and targets PLK-1 to chromosome

arms in *C. elegans* oocytes through a polo-docking site.

A) Kinase assay of recombinant CENP-C$^{HCP-4(1-214)}$ wild type and T163A proteins with

Cdk1:Cyclin B. Reactions were analysed by SDS-PAGE followed by ProQ diamond

(phosphoprotein) or coomassie (total protein) staining.

B) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. Wild type or T163A mutant CENP-C^{HCP-}

$^{4(1-214)}$ was phosphorylated by Cdk1:Cyclin B before incubation with an equimolar

concentration of MBP-PLK1^{PBD}. Binding was then assessed by SEC.

C) FITC-labelled peptides containing the HCP-4 STP motif were incubated with

increasing concentrations of MBP:PLK1^{PBD} and binding analysed by fluoresence

polarisation. Unphosphorylated versus T163-phosphorylated (T^{ph}) peptides were

compared.

D) Control (*'wild type'*) and CENP-C^{HCP-4} STP mutant [*'hcp-4^{T163A}'*] oocytes expressing sfGFP-tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded throughout meiosis I. Metphase I is shown. Left scale bar, 2 µm. Right scale bar, 1 µm.

E) Line profile analysis of PLK-1::GFP during early Metaphase I in *wild type* and *hcp-4^{T163A}* oocytes, as indicated by the yellow lines. Background signal was subtracted and maximum signal for each channel was set to 1.

F) Fixed oocytes were stained with an HCP-4 specific antibody (green in the merged image). *hcp-4^{T163A}* mutant oocytes were compared to wild type. DNA is shown in magenta in the merged panel.

G) Same as in F) but using a BUB-1 specific antibody to compare BUB-1 localisation in wild type and *hcp-4^{T163A}* mutant oocytes.

Figure 6. Combined disruption of BUB-1- and CENP-C^{HCP-4}-dependent PLK-1 recruitment leads to severe meiotic defects.

A) Control (*'wild type'*), BUB-1-depleted [*'bub-1(RNAi)'*], and CENP-C^{HCP-4}-depleted [*'hcp-4(RNAi)'*] oocytes expressing sfGFP-tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded throughout meiosis I. Prometaphase/Metphase I is shown (before spindle rotation/shortening). Left scale bar, 2 µm. Right scale bar, 1 µm.

B) Control (*'wild type'*), BUB-1-depleted [*'bub-1(RNAi)'*], and CENP-C^{HCP-4}-depleted [*'hcp-4(RNAi)'*] oocytes expressing GFP-tagged tubulin (and mCherry-tagged histone) were dissected and recorded throughout meiosis I. Scale bar, 2 µm.

C) Meiotic defects (as described in the Methods section) were assessed in wild type, *bub-*

1(RNAi), and *hcp-4(RNAi)* oocytes.

D) Fixed wild type, *hcp-4(RNAi)*, *bub-1^{T527A},* and *bub-1^{T527A}+hcp-4(RNAi)* oocytes were

stained with a PLK-1 specific antibody.

E) Control (*'wild type'*), BUB-1^{T527A} ['*bub-1^{T527A}*'], and CENP-C^{HCP-4}-depleted ['*hcp-*

4(RNAi)'] oocytes expressing GFP-tagged tubulin (and mCherry-tagged histone) were

dissected and recorded throughout meiosis I. Scale bar, 2 μm.

F) Meiotic defects (as described in the Methods section) were assessed in wild type, *bub-*

1^{T527A}, *hcp-4(RNAi),* and *bub-1^{T527A}+hcp-4(RNAi)* oocytes.

Figure S1. PLK-1 localisation during oocyte meiosis

In situ GFP-tagged PLK-1 (Martino et al., 2017) was followed through meiosis I in

live, dissected oocytes. Scale bar, 2 μm. This panel shows a more detailed sequence of

events of the same movie presented in Figure 1B.

Figure S2. Embryonic viability assays after PLK-1as inhibition

Three different PP1 analogues were tested: 3IB-PP1, 3MB-PP1, and 1NA-PP1.

Inhibitors were included in the plates at 10, 20, and 50 μM and worms were then

allowed to lay eggs. Viable progeny was assessed 24 and 48 hs later.

Figure S3. Chromosome alignment defects upon acute PLK-1as inhibition

A) *Wild type* or *plk-1as* worms expressing GFP-tagged tubulin and mCherry-tagged

histone, were dissected in medium containing ethanol, 1 μM 3IB-PP1 ('low'), or 10

µM ('high'). Maximum intensity projections are shown and the arrows have different

colours to identify those chromosomes in panel B). Scale bar, 2 µm.

B) Detailed analysis of individual chromosome behaviour using selected Z slices of

+3IB-PP1 'low' from panel A). Highly unstable behaviour of specific chromosomes

(switching back and forth between aligned and misaligned) can be followed by the

different coloured arrows. Panles outside of the green rectangle are timepoints not

shown in A). Scale bar, 2 µm.

Figure S4.

A) Kinase assay of recominant BUB-1$^{190-628}$ and BUB-1$^{190-628}$(T527A) with Cdk1:Cyclin

B. Proteins were incubated with Cdk1:Cyclin B for the indicated time points before

phosphorylation was assessed by SDS-PAGE and staining with either coomassie (total

protein) or ProQ diamond (phosphoprotein).

B) Kinase assay of recominant BUB-1$^{190-628}$ and BUB-1$^{190-628}$(T527A) with PLK-1.

Proteins were incubated with PLK-1 for the indicated time points before

phosphorylation was assessed by SDS-PAGE and staining with either coomassie (total

protein) or ProQ diamond (phosphoprotein).

Figure S5.

Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. CENCP-C$^{HCP-4(1-214)}$ was incubated with

MBP-PLK-1^{PBD} at equimolar concentrations before being analysed by SEC. Binding

was conducted with unphosphorylated or Cdk1:Cyclin B phosphorylated CENP-C^{HCP-}

$^{4(1-214)}$ as indicated.

Figure S6.

A) Control (*'wild type'*), BUB-1-depleted [*'bub-1(RNAi)'*], CENP-C^{HCP-4}(T163A) [*'hcp-*

4^{T163A}'], and *bub-1(RNAi)+hcp-4^{T163A}* oocytes expressing sfGFP-tagged PLK-1 (and

mCherry-tagged histone) were dissected and recorded throughout meiosis I.

Prometaphase/Metphase I is shown (before spindle rotation/shortening). Panels on the

left show the full spindle, while the right-most two columns display single bivalents.

Left scale bar, 2 μm. Right scale bar, 1 μm.

B) Control (*'wild type'*), BUB-1-depleted [*'bub-1(RNAi)'*], CENP-C^{HCP-4}(T163A) [*'hcp-*

4^{T163A}'], and *bub-1(RNAi)+hcp-4^{T163A}* oocytes expressing GFP-tagged tubulin (and

mCherry-tagged histone) were dissected and recorded throughout meiosis I. Scale bar,

2 μm.

C) Meiotic defects (as described in the Methods section) were assessed in wild type, *bub-*

1(RNAi), hcp-4^{T163A}, and bub-1(RNAi)+hcp-4^{T163A} oocytes.

REFERENCES

Archambault V, Glover DM. 2009. Polo-like kinases: conservation and divergence in their

functions and regulation. *Nat Rev Mol Cell Biol* **10**:265–275. doi:10.1038/nrm2653

Arribere JA, Bell RT, Fu BXH, Artiles KL, Hartman PS, Fire AZ. 2014. Efficient Marker-

Free Recovery of Custom Genetic Modifications with CRISPR/Cas9 in

Caenorhabditis elegans. *Genetics* **198**:837–846. doi:10.1534/genetics.114.169730

Bel Borja L, Soubigou F, Taylor SJP, Fraguas Bringas C, Budrewicz J, Lara-Gonzalez P,

Sorensen Turpin CG, Bembenek JN, Cheerambathur DK, Pelisch F. 2020. BUB-1

targets PP2A:B56 to regulate chromosome congression during meiosis I in C. elegans

oocytes. *eLife* **9**:e65307. doi:10.7554/eLife.65307

Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J, Shimizu E, Tsien JZ,

Schultz PG, Rose MD, Wood JL, Morgan DO, Shokat KM. 2000. A chemical switch

for inhibitor-sensitive alleles of any protein kinase. *Nature* **407**:395–401.

doi:10.1038/35030148

Budirahardja Y, Gönczy P. 2008. PLK-1 asymmetry contributes to asynchronous cell division

of *C. elegans* embryos. *Development* **135**:1303–1313. doi:10.1242/dev.019075

Cabral G, Laos T, Dumont J, Dammermann A. 2019. Differential Requirements for

Centrioles in Mitotic Centrosome Growth and Maintenance. *Developmental Cell*

50:355-366.e6. doi:10.1016/j.devcel.2019.06.004

Chase D, Serafinas C, Ashcroft N, Kosinski M, Longo D, Ferris DK, Golden A. 2000. The

polo-like kinase PLK-1 is required for nuclear envelope breakdown and the

completion of meiosis inCaenorhabditis elegans. *genesis* **26**:26–41.

doi:10.1002/(SICI)1526-968X(200001)26:1<26::AID-GENE6>3.0.CO;2-O

Cheerambathur DK, Prevo B, Chow T-L, Hattersley N, Wang S, Zhao Z, Kim T, Gerson-

Gurwitz A, Oegema K, Green R, Desai A. 2019. The Kinetochore-Microtubule

Coupling Machinery Is Repurposed in Sensory Nervous System Morphogenesis.

Developmental Cell **48**:864-872.e7. doi:10.1016/j.devcel.2019.02.002

Cheng K-Y. 2003. The crystal structure of the human polo-like kinase-1 polo box domain and

its phospho-peptide complex. *The EMBO Journal* **22**:5757–5768.

doi:10.1093/emboj/cdg558

Connolly AA, Sugioka K, Chuang C-H, Lowry JB, Bowerman B. 2015. KLP-7 acts through

the Ndc80 complex to limit pole number in C. elegans oocyte meiotic spindle

assembly. *Journal of Cell Biology* **210**:917–932. doi:10.1083/jcb.201412010

Conway W, Kiewisz R, Fabig G, Kelleher CP, Wu H-Y, Anjur-Dietrich M, Müller-Reichert

T, Needleman DJ. 2022. Self-organization of kinetochore-fibers in human mitotic

spindles. *eLife* **11**:e75458. doi:10.7554/eLife.75458

Cordeiro MH, Smith RJ, Saurin AT. 2020. Kinetochore phosphatases suppress autonomous

Polo-like kinase 1 activity to control the mitotic checkpoint. *Journal of Cell Biology*

219:e202002020. doi:10.1083/jcb.202002020

Danlasky BM, Panzica MT, McNally KP, Vargas E, Bailey C, Li W, Gong T, Fishman ES,

Jiang X, McNally FJ. 2020. Evidence for anaphase pulling forces during C. elegans

meiosis. *Journal of Cell Biology* **219**:e202005179. doi:10.1083/jcb.202005179

Decker M, Jaensch S, Pozniakovsky A, Zinke A, O'Connell KF, Zachariae W, Myers E,

Hyman AA. 2011. Limiting Amounts of Centrosome Material Set Centrosome Size in

C. elegans Embryos. *Current Biology* **21**:1259–1267. doi:10.1016/j.cub.2011.06.002

Desai A, Rybina S, Müller-Reichert T, Shevchenko Andrej, Shevchenko Anna, Hyman A,

Oegema K. 2003. KNL-1 directs assembly of the microtubule-binding interface of the

kinetochore in *C. elegans*. *Genes Dev* **17**:2421–2435. doi:10.1101/gad.1126303

Dumont J, Oegema K, Desai A. 2010. A kinetochore-independent mechanism drives

anaphase chromosome separation during acentrosomal meiosis. *Nat Cell Biol* **12**:894–

901. doi:10.1038/ncb2093

Elia AEH, Cantley LC, Yaffe MB. 2003a. Proteomic Screen Finds pSer/pThr-Binding

Domain Localizing Plk1 to Mitotic Substrates. *Science* **299**:1228–1231.

doi:10.1126/science.1079079

Elia AEH, Rellos P, Haire LF, Chao JW, Ivins FJ, Hoepker K, Mohammad D, Cantley LC,

Smerdon SJ, Yaffe MB. 2003b. The Molecular Basis for Phosphodependent Substrate

Targeting and Regulation of Plks by the Polo-Box Domain. *Cell* **115**:83–95.

doi:10.1016/S0092-8674(03)00725-6

Elowe S, Hümmer S, Uldschmid A, Li X, Nigg EA. 2007. Tension-sensitive Plk1

phosphorylation on BubR1 regulates the stability of kinetochore–microtubule

interactions. *Genes Dev* **21**:2205–2219. doi:10.1101/gad.436007

Foltz DR, Jansen LET, Black BE, Bailey AO, Yates JR, Cleveland DW. 2006. The human

CENP-A centromeric nucleosome-associated complex. *Nat Cell Biol* **8**:458–469.

doi:10.1038/ncb1397

Gelens L, Qian J, Bollen M, Saurin AT. 2018. The Importance of Kinase–Phosphatase

Integration: Lessons from Mitosis. *Trends in Cell Biology* **28**:6–21.

doi:10.1016/j.tcb.2017.09.005

Gigant E, Stefanutti M, Laband K, Gluszek-Kustusz A, Edwards F, Lacroix B, Maton G,

Canman JC, Welburn JPI, Dumont J. 2017. Inhibition of ectopic microtubule

assembly by the kinesin-13 KLP-7 prevents chromosome segregation and cytokinesis

defects in oocytes. *Journal of Cell Science* **130**:e1.1-e1.1. doi:10.1242/jcs.205914

Gómez-Cavazos JS, Lee K-Y, Lara-González P, Li Y, Desai A, Shiau AK, Oegema K. 2020.

A Non-canonical BRCT-Phosphopeptide Recognition Mechanism Underlies RhoA

Activation in Cytokinesis. *Current Biology* **30**:3101-3115.e11.

doi:10.1016/j.cub.2020.05.090

Green RA, Mayers JR, Wang S, Lewellyn L, Desai A, Audhya A, Oegema K. 2013. The

midbody ring scaffolds the abscission machinery in the absence of midbody

microtubules. *Journal of Cell Biology* **203**:505–520. doi:10.1083/jcb.201306036

Han X, Adames K, Sykes EME, Srayko M. 2015. The KLP-7 Residue S546 Is a Putative

Aurora Kinase Site Required for Microtubule Regulation at the Centrosome in C.

elegans. *PLoS ONE* **10**:e0132593. doi:10.1371/journal.pone.0132593

Hattersley N, Schlientz AJ, Prevo B, Oegema K, Desai A. 2022. MEL-28/ELYS and CENP-C

coordinately control outer kinetochore assembly and meiotic chromosome-

microtubule interactions. *Current Biology* **32**:2563-2571.e4.

doi:10.1016/j.cub.2022.04.046

Heald R, Tournebize R, Blank T, Sandaltzopoulos R, Becker P, Hyman A, Karsenti E. 1996.

Self-organization of microtubules into bipolar spindles around artificial chromosomes

in Xenopus egg extracts. *Nature* **382**:420–425. doi:10.1038/382420a0

Heun P, Erhardt S, Blower MD, Weiss S, Skora AD, Karpen GH. 2006. Mislocalization of

the Drosophila Centromere-Specific Histone CID Promotes Formation of Functional

Ectopic Kinetochores. *Developmental Cell* **10**:303–315.

doi:10.1016/j.devcel.2006.01.014

Izuta H, Ikeno M, Suzuki N, Tomonaga T, Nozaki N, Obuse C, Kisu Y, Goshima N, Nomura

F, Nomura N, Yoda K. 2006. Comprehensive analysis of the ICEN (Interphase

Centromere Complex) components enriched in the CENP-A chromatin of human

cells. *Genes to Cells* **11**:673–684. doi:10.1111/j.1365-2443.2006.00969.x

Jang C-Y, Coppinger JA, Seki A, Yates JR III, Fang G. 2009. Plk1 and Aurora A regulate the

depolymerase activity and the cellular localization of Kif2a. *Journal of Cell Science*

122:1334–1341. doi:10.1242/jcs.044321

Kang YH, Park CH, Kim T-S, Soung N-K, Bang JK, Kim BY, Park J-E, Lee KS. 2011.

Mammalian Polo-like Kinase 1-dependent Regulation of the PBIP1-CENP-Q

Complex at Kinetochores. *Journal of Biological Chemistry* **286**:19744–19757.

doi:10.1074/jbc.M111.224105

Kang YH, Park J-E, Yu L-R, Soung N-K, Yun S-M, Bang JK, Seong Y-S, Yu H, Garfield S,

Veenstra TD, Lee KS. 2006. Self-Regulated Plk1 Recruitment to Kinetochores by the

Plk1-PBIP1 Interaction Is Critical for Proper Chromosome Segregation. *Molecular*

Cell **24**:409–422. doi:10.1016/j.molcel.2006.10.016

Kiewisz R, Fabig G, Conway W, Baum D, Needleman D, Müller-Reichert T. 2022. Three-

dimensional structure of kinetochore-fibers in human mitotic spindles. *eLife*

11:e75459. doi:10.7554/eLife.75459

Kitada K, Johnson AL, Johnston LH, Sugino A. 1993. A multicopy suppressor gene of the

Saccharomyces cerevisiae G1 cell cycle mutant gene dbf4 encodes a protein kinase

and is identified as CDC5. *Molecular and Cellular Biology* **13**:4445–4457.

doi:10.1128/mcb.13.7.4445-4457.1993

Laband K, Lacroix B, Edwards F, Canman JC, Dumont J. 2018. Chapter 11 - Live imaging of

C. elegans oocytes and early embryos In: Maiato H, Schuh M, editors. Methods in

Cell Biology, Mitosis and Meiosis Part B. Academic Press. pp. 217–236.

doi:10.1016/bs.mcb.2018.03.025

Llamazares S, Moreira A, Tavares A, Girdham C, Spruce BA, Gonzalez C, Karess RE,

Glover DM, Sunkel CE. 1991. polo encodes a protein kinase homolog required for

mitosis in Drosophila. *Genes Dev* **5**:2153–2165. doi:10.1101/gad.5.12a.2153

Maddox PS, Corbett KD, Desai A. 2012. Structure, assembly and reading of centromeric

chromatin. *Current Opinion in Genetics & Development*, Genome architecture and

expression **22**:139–147. doi:10.1016/j.gde.2011.11.005

Marston AL, Amon A. 2004. Meiosis: cell-cycle controls shuffle and deal. *Nat Rev Mol Cell*

Biol **5**:983–997. doi:10.1038/nrm1526

Marston AL, Wassmann K. 2017. Multiple Duties for Spindle Assembly Checkpoint Kinases

in Meiosis. *Frontiers in Cell and Developmental Biology* **5**.

Martino L, Morchoisne-Bolhy S, Cheerambathur DK, Van Hove L, Dumont J, Joly N, Desai

A, Doye V, Pintard L. 2017. Channel Nucleoporins Recruit PLK-1 to Nuclear Pore

Complexes to Direct Nuclear Envelope Breakdown in C. elegans. *Developmental Cell*

43:157-171.e7. doi:10.1016/j.devcel.2017.09.019

Monen J, Maddox PS, Hyndman F, Oegema K, Desai A. 2005. Differential role of CENP-A

in the segregation of holocentric C. elegans chromosomes during meiosis and mitosis.

Nat Cell Biol **7**:1248–1255. doi:10.1038/ncb1331

Mundt KE, Golsteyn RM, Lane HA, Nigg EA. 1997. On the Regulation and Function of

Human Polo-like Kinase 1 (PLK1): Effects of Overexpression on Cell Cycle

Progression. *Biochemical and Biophysical Research Communications* **239**:377–385.

doi:10.1006/bbrc.1997.7378

Musacchio A, Desai A. 2017. A Molecular View of Kinetochore Assembly and Function.

Biology **6**. doi:10.3390/biology6010005

Neef R, Preisinger C, Sutcliffe J, Kopajtich R, Nigg EA, Mayer TU, Barr FA. 2003.

Phosphorylation of mitotic kinesin-like protein 2 by polo-like kinase 1 is required for

cytokinesis. *Journal of Cell Biology* **162**:863–876. doi:10.1083/jcb.200306009

Novak B, Kapuy O, Domingo-Sananes MR, Tyson JJ. 2010. Regulated protein kinases and

phosphatases in cell cycle decisions. *Current Opinion in Cell Biology*, Cell

808   differentiation / Cell division, growth and death **22**:801–808.

809   doi:10.1016/j.ceb.2010.07.001

810 Oegema K, Desai A, Rybina S, Kirkham M, Hyman AA. 2001. Functional Analysis of

811   Kinetochore Assembly in Caenorhabditis elegans. *Journal of Cell Biology* **153**:1209–

812   1226. doi:10.1083/jcb.153.6.1209

813 Ohkura H. 2015. Meiosis: An Overview of Key Differences from Mitosis. *Cold Spring Harb*

814   *Perspect Biol* **7**:a015859. doi:10.1101/cshperspect.a015859

815 Ohkura H, Hagan IM, Glover DM. 1995. The conserved Schizosaccharomyces pombe kinase

816   plo1, required to form a bipolar spindle, the actin ring, and septum, can drive septum

817   formation in G1 and G2 cells. *Genes Dev* **9**:1059–1073. doi:10.1101/gad.9.9.1059

818 Ohta M, Zhao Z, Wu D, Wang S, Harrison JL, Gómez-Cavazos JS, Desai A, Oegema KF.

819   2021. Polo-like kinase 1 independently controls microtubule-nucleating capacity and

820   size of the centrosome. *Journal of Cell Biology* **220**:e202009083.

821   doi:10.1083/jcb.202009083

822 Okada M, Cheeseman IM, Hori T, Okawa K, McLeod IX, Yates JR, Desai A, Fukagawa T.

823   2006. The CENP-H–I complex is required for the efficient incorporation of newly

824   synthesized CENP-A into centromeres. *Nat Cell Biol* **8**:446–457.

825   doi:10.1038/ncb1396

826 Pahlavan G, Polanski Z, Kalab P, Golsteyn R, Nigg EA, Maro B. 2000. Characterization of

827   Polo-like Kinase 1 during Meiotic Maturation of the Mouse Oocyte. *Developmental*

828   *Biology* **220**:392–400. doi:10.1006/dbio.2000.9656

829 Paix A, Folkmann A, Rasoloson D, Seydoux G. 2015. High Efficiency, Homology-Directed

830   Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein

831   Complexes. *Genetics* **201**:47–54. doi:10.1534/genetics.115.179382

Paix A, Folkmann A, Seydoux G. 2017. Precision genome editing using CRISPR-Cas9 and

linear repair templates in C. elegans. *Methods*, CRISPR-Cas systems for Genome

engineering and investigation **121–122**:86–93. doi:10.1016/j.ymeth.2017.03.023

Pelisch F, Bel Borja L, Jaffray EG, Hay RT. 2019. Sumoylation regulates protein dynamics

during meiotic chromosome segregation in *C. elegans* oocytes. *Journal of Cell*

Science jcs.232330. doi:10.1242/jcs.232330

Pelisch F, Tammsalu T, Wang B, Jaffray EG, Gartner A, Hay RT. 2017. A SUMO-Dependent

Protein Network Regulates Chromosome Congression during Oocyte Meiosis.

Molecular Cell **65**:66–77. doi:10.1016/j.molcel.2016.11.001

Petronczki M, Lénárt P, Peters J-M. 2008. Polo on the Rise—from Mitotic Entry to

Cytokinesis with Plk1. *Developmental Cell* **14**:646–659.

doi:10.1016/j.devcel.2008.04.014

Qi W, Tang Z, Yu H. 2006. Phosphorylation- and Polo-Box–dependent Binding of Plk1 to

Bub1 Is Required for the Kinetochore Localization of Plk1. *MBoC* **17**:3705–3716.

doi:10.1091/mbc.e06-03-0240

Rahman MM, Munzig M, Kaneshiro K, Lee B, Strome S, Müller-Reichert T, Cohen-Fix O.

2015. Caenorhabditis elegans polo-like kinase PLK-1 is required for merging parental

genomes into a single nucleus. *Mol Biol Cell* **26**:4718–4735. doi:10.1091/mbc.E15-

04-0244

Ritter A, Sanhaji M, Steinhäuser K, Roth S, Louwen F, Yuan J. 2014. The activity regulation

of the mitotic centromere-associated kinesin by Polo-like kinase 1. *Oncotarget*

6:6641–6655. doi:10.18632/oncotarget.2843

Sanhaji M, Ritter A, Belsham HR, Friel CT, Roth S, Louwen F, Yuan J. 2014. Polo-like

kinase 1 regulates the stability of the mitotic centromere-associated kinesin in mitosis.

Oncotarget **5**:3130–3144. doi:10.18632/oncotarget.1861

Saurin AT. 2018. Kinase and Phosphatase Cross-Talk at the Kinetochore. *Front Cell Dev Biol*

6:62. doi:10.3389/fcell.2018.00062

Schmucker S, Sumara I. 2014. Molecular dynamics of PLK1 during mitosis. *Molecular &*

Cellular Oncology **1**:e954507. doi:10.1080/23723548.2014.954507

Shao H, Huang Y, Zhang L, Yuan K, Chu Y, Dou Z, Jin C, Garcia-Barrio M, Liu X, Yao X.

2015. Spatiotemporal dynamics of Aurora B-PLK1-MCAK signaling axis orchestrates

kinetochore bi-orientation and faithful chromosome segregation. *Sci Rep* **5**:12204.

doi:10.1038/srep12204

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,

Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of

high-quality protein multiple sequence alignments using Clustal Omega. *Molecular*

Systems Biology **7**:539. doi:10.1038/msb.2011.75

Singh P, Pesenti ME, Maffini S, Carmignani S, Hedtfeld M, Petrovic A, Srinivasamani A,

Bange T, Musacchio A. 2021. BUB1 and CENP-U, Primed by CDK1, Are the Main

PLK1 Kinetochore Receptors in Mitosis. *Molecular Cell* **81**:67-87.e9.

doi:10.1016/j.molcel.2020.10.040

Solc P, Kitajima TS, Yoshida S, Brzakova A, Kaido M, Baran V, Mayer A, Samalova P,

Motlik J, Ellenberg J. 2015. Multiple requirements of PLK1 during mouse oocyte

maturation. *PLoS One* **10**:e0116783. doi:10.1371/journal.pone.0116783

Sunkel CE, Glover DM. 1988. polo, a mitotic mutant of Drosophila displaying abnormal

spindle poles. *Journal of Cell Science* **89**:25–38. doi:10.1242/jcs.89.1.25

Tavernier N, Noatynska A, Panbianco C, Martino L, Van Hove L, Schwager F, Léger T,

Gotta M, Pintard L. 2015. Cdk1 phosphorylates SPAT-1/Bora to trigger PLK-1

activation and drive mitotic entry in C. elegans embryos. *Journal of Cell Biology*

208:661–669. doi:10.1083/jcb.201408064

Timmons L, Court DL, Fire A. 2001. Ingestion of bacterially expressed dsRNAs can produce

specific and potent genetic interference in Caenorhabditis elegans. *Gene* **263**:103–112.

doi:10.1016/S0378-1119(00)00579-5

Tong C, Fan H-Y, Lian L, Li S-W, Chen D-Y, Schatten H, Sun Q-Y. 2002. Polo-Like Kinase-

1 Is a Pivotal Regulator of Microtubule Assembly During Mouse Oocyte Meiotic

Maturation, Fertilization, and Early Embryonic Mitosis1. *Biology of Reproduction*

67:546–554. doi:10.1095/biolreprod67.2.546

Velez-Aguilera G, Nkombo Nkoula S, Ossareh-Nazari B, Link J, Paouneskou D, Van Hove

L, Joly N, Tavernier N, Verbavatz J-M, Jantsch V, Pintard L. 2020. PLK-1 promotes

the merger of the parental genome into a single nucleus by triggering lamina

disassembly. *eLife* **9**:e59510. doi:10.7554/eLife.59510

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2—a

multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**:1189–

1191. doi:10.1093/bioinformatics/btp033

Wianny F, Tavares Á, Evans MJ, Glover DM, Zernicka-Goetz M. 1998. Mouse polo-like

kinase 1 associates with the acentriolar spindle poles, meiotic chromosomes and

spindle midzone during oocyte maturation. *Chromosoma* **107**:430–439.

doi:10.1007/s004120050327

Woodruff JB, Wueseke O, Viscardi V, Mahamid J, Ochoa SD, Bunkenborg J, Widlund PO,

Pozniakovsky A, Zanin E, Bahmanyar S, Zinke A, Hong SH, Decker M, Baumeister

W, Andersen JS, Oegema K, Hyman AA. 2015. Regulated assembly of a

supramolecular centrosome scaffold in vitro. *Science* **348**:808–812.

doi:10.1126/science.aaa3923

Xu J, Shen C, Wang T, Quan J. 2013. Structural basis for the inhibition of Polo-like kinase 1.

Nat Struct Mol Biol **20**:1047–1053. doi:10.1038/nsmb.2623

Zhang L, Shao H, Huang Y, Yan F, Chu Y, Hou H, Zhu M, Fu C, Aikhionbare F, Fang G,

Ding X, Yao X. 2011. PLK1 Phosphorylates Mitotic Centromere-associated Kinesin

and Promotes Its Depolymerase Activity*. *Journal of Biological Chemistry* **286**:3033–

3046. doi:10.1074/jbc.M110.165340

Zitouni S, Nabais C, Jana SC, Guerrero A, Bettencourt-Dias M. 2014. Polo-like kinases:

structural variations lead to multiple functions. *Nat Rev Mol Cell Biol* **15**:433–452.

doi:10.1038/nrm3819

METHODS

C. elegans **strains & RNAi**

Strains used in this study were maintained at 20 degrees unless indicated otherwise. For a

complete list of strains, please refer to Table S1.

For RNAi-mediated depletions, the targeting sequence for *bub-1* was 2353-2935 and for *hcp-*

4, 967-2128, both from the first ATG codon. For double depletion, both sequences were

cloned in the same vector. All sequences were inserted into L4440 using the NEBuilder HiFi

DNA Assembly Master Mix (New England Biolabs) and transformed into DH5a bacteria. The

purified plasmids were then transformed into HT115(DE3) bacteria (Timmons et al., 2001).

RNAi clones were picked and grown overnight at 37°C in LB with 100 μg/ml ampicillin.

Saturated cultures were diluted 1:100 and allowed to grow until reaching an OD600 of 0.8–1.

Isopropyl-β-d-thiogalactopyranoside (IPTG) was added to a final concentration of 1 mM, and

cultures were incubated for 1 h at 37°C. Bacteria were then seeded onto NGM plates made

with agarose and 1 mM IPTG and allowed to dry. L4 worms were then plated on RNAi plates

and maintained at 20°C.

CRISPR/Cas9

We used direct injection of in vitro–assembled Cas9-CRISPR RNA (crRNA) trans-activating

crRNA (tracrRNA) ribonucleoprotein complexes (Paix et al., 2017, 2015).

For mutation of Threonine 527 to Alanine in *bub-1*, we used the following crRNA (+ strand):

CCCCGCACAAGGAGTTCATT and repair template (- strand):

acttacTAATTTACTGAAAGTACTGCTGGTTGGAGCAACAAATACTGGAGCTTCCTGT

TCGTGAGTGCTTTCCTCCTCTTTATTTCCGAAATATTCATCAATGTTGACtAAATGA

ACTCCTTGTGCGGGgGcactaGTGACGAAATTACCACGAGACGGTTTGAAAAAGCCA

AACTCGATTTCATCGTCATAAActaaaa

For mutation of Threonine 163 to Alanine in *hcp-4*, we used the following crRNA (- strand):

TGAAATATCAAGCGATCTCA and repair template (+ strand):

taataaatctataatttcagAGTGGAAAAGCTGGATTAAGcTGcagtgCaCCCAAGAGCTCGAGTG

ATACGTCGATGAGGTCtTTGAGATCGCTTGATATTTCACATGTCGTCAATACCGAT

C

Each of the mixes was mixed with dpy-10 crRNA/repair template for screening (Arribere et

al., 2014).

Live imaging of oocytes

A detailed protocol for live imaging of *C. elegans* oocytes was used with minor modifications

(Laband et al., 2018). Fertilized oocytes were dissected and mounted in 5 µl of L-15

blastomere culture medium (0.5 mg/mL Inulin; 25 mM HEPES, pH 7.5 in 60% Leibowitz L-

15 medium and 20% heat-Inactivated FBS) on 24×40 mm #1.5 coverslips. Once dissection

was performed and early oocytes identified using a stereomicroscope, a circle of Vaseline was

laid around the sample, and a custom-made 24×40 mm plastic holder (with a centred window)

was placed on top. The sample was imaged immediately using 488 nm and 561 nm laser

lines. Live imaging was done using a CFI Plan Apochromat Lambda 60X/NA 1.4 oil

objective mounted on a microscope (Nikon Eclipse Ti) equiped with a Prime 95B 22mm

camera (Photometrics), a spinning-disk head (CSU-X1; Yokogawa Electric Corporation).

Acquisition parameters were controlled with NIS software (Nikon). For Figures 1B and 2A,

the microscope used was an IX81 (Olympus) equiped with an EMCCD Cascade II camera

(Photometrics) and a CSU-X1 spinning disk head (Yokogawa). For all live imaging

experiments, partial projections are presented. All files were stored, classified, and managed

using OMERO (Allan et al., 2012). Figures were prepared using OMERO.figure and

assembled using Adobe Illustrator. Representative movies shown in Supplementary material

were assembled using Fiji/ImageJ (Schindelin et al., 2012) custom-made macros.

Immunofluorescence

Worms were placed on 4 µl of M9 worm buffer in a poly-D-lysine (Sigma, P1024)-coated

slide and a 24×24-cm coverslip was gently laid on top. Once the worms extruded the

embryos, slides were placed on a metal block on dry ice for >10 min. The coverslip was then

flicked off with a scalpel blade, and the samples were fixed in methanol at 20°C for 30 min

(except for GFP, where the methanol treatment lasted 5 min). Secondary antibodies were

donkey anti–sheep, goat anti-mouse, or goat anti-rabbit conjugated to Alexa Fluor™ 488,

Alexa Fluor™ 594, and Alexa Fluor™ 647 (1:1,000, Thermo Scientific). Donkey anti-mouse

and donkey anti-rabbit conjugated secondary antibodies were obtained from Jackson

ImmunoReserach. Embryos were mounted in ProLong Diamond antifade mountant (Thermo

Scientific) with DAPI. Primary antibodies were: α-PLK-1 (Budirahardja and Gönczy, 2008),

α-HCP-4 (Oegema et al., 2001), α-BUB-1, purified in house after immunisation of rabbits

using the sequence in (Desai et al., 2003),

Sequence alignments

Sequences shown in Figures 2C and 4B were aligned with Clustal Omega (Sievers et al.,

2011) and visualised with Jalview (Waterhouse et al., 2009).

Protein Purification

GST-BUB-1$^{190-628}$ and GST-HCP-4$^{1-214}$ proteins were expressed in *Escherichia coli* BL21

DE3 bacteria by diluting a saturated culture 1/100 in LB media supplemented with 35 µg/ml

ampicillin and incubating at 37°C/200 rpm until OD_{600} 0.6-0.8 was reached. IPTG was then

added to a final concentration of 100 μM and cultures were incubated at 20°C/200 rpm for 16-

18 h. The bacterial cultures were then centrifuged (20 min/6,250 g/4°C) and pellets

resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5; 150 mM NaCl; 0.5 mM TCEP, 1X

Roche cOmplete protease inhibitors, EDTA free; 30-35 ml/L of culture). The cell suspension

was then sonicated (2 min 40 sec, 20 sec on/40 sec off) before centrifugation (45 min, 27,250

g, 4°C) to remove insoluble material. GSH sepharose beads were washed with 10 column

volumes (CV) of MilliQ water and equilibrated with 10 CV of binding buffer (50 mM Tris-

HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP) before filtered (0.22 μm PES filter) lysate was

added and batch bound at 4°C for 1.5-2 h. The beads were then collected in a column and

washed with at least 10 CV of binding buffer before being transferred to a falcon tube and

incubated overnight with GST-tagged 3C protease. After cleavage, the beads were collected

in a column and the flowthrough containing cleaved protein was concentrated in a Vivaspin

centrifugal concentrator. The protein was then further purified by size exclusion

chromatography in a Superdex 200 10/300 column (run in SEC buffer, see below) and

concentrated before being flash frozen in liquid nitrogen and stored at –80.

MBP-PLK1^{PBD} (6xHIS-MBP tagged human PLK1$^{345-603}$, as per (Singh et al., 2021)) was

expressed and purified in the same manner as the GST tagged proteins with the following

exceptions: filtered lysate was passed through a cobalt-NTA column and washed with at least

10 CV of binding buffer (see buffers below) before elution in 0.5 CV fractions. Fractions

containing the protein were concentrated using Vivapsin centrifugal filters and further

purified by size exclusion chromatography (Superdex 200 10/300) in SEC buffer. Lysis buffer

(50 mM Tris-HCl, pH 7.5; 500 mM NaCl; 10 mM imidazole; 0.5 mM TCEP; 1X Roche

cOmplete protease inhibitors, EDTA free; 50 ml/L of culture); binding buffer (50 mM Tris-

HCl, pH 7.5; 500 mM NaCl; 10 mM imidazole; 0.5 mM TCEP); elution buffer (50 mM Tris-

HCl, pH 7.5; 150 mM NaCl; 200 mM imidazole; 0.5 mM TCEP); SEC buffer (50 mM Tris-

HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP).

1017

Kinase assays

Unless otherwise stated, kinase assays were conducted with 55 nM Cdk1:Cyclin B (Thermo

Scientific) and/or 75 nM PLK-1, produced as described in (Tavernier et al., 2015), at 30°C in

kinase buffer: 50 mM Tris-HCl, pH 7.5; 1 mM ATP; 10 mM $MgCl_2$; 0.5 mM TCEP; 0.1 mM

EDTA. For the assays in Figure 2E and Figure S4; 0.4 µg/µl (8.2 µM) of BUB-1[190-628]

substrate was used and incubated under the above conditions for up to 2 h. The assay in

Figure 5A was conducted in the above conditions for up to 1 h with a substrate concentration

of 0.2 µg/µl (8.2 µM). The assay in Figure 2D was conducted with 165 nM Cdk1:Cyclin B

and 170 nM PLK-1 and incubated for 30 min at 37°C with the following buffer: 40 mM Tris-

HCl, pH 7.5; 100 µM ATP, 10 mM $MgCl_2$. BUB-1[190-628] concentration was 0.2 µg/µl (4.1

µM).

In all assays, aliquots of protein at the indicated timepoints were immediately added to an

equal volume of 2X LDS buffer (Thermo) and incubated at 70 degrees for 15 min. Assays

were assessed by SDS-PAGE combined with ProQ diamond phosphoprotein and coomassie

staining.

1033

***In vitro* Binding Assays**

BUB-1[190-628] or CENP-C[HCP-4(1-214)] recombinant proteins were incubated for 50-60 min at

30°C in the presence or absence of ~100 nM Cdk1:Cyclin B in kinase buffer (see 'kinase

assay' section of methods). The respective proteins were then incubated for 50 min on ice

with MBP-PLK1[PBD] at a concentration of 20 µM in SEC buffer (50 mM Tris-HCl, pH 7.5;

150 mM NaCl; 0.5 mM TCEP). Assays were then centrifuged (13.3k rpm, 10 min, 4°C)

before being loaded onto a Superdex 200 10/300 SEC column. In all assays, 0.2 ml fractions

were collected and selected fractions were assessed by SDS-PAGE with coomassie staining.

More specifically; for the binding assays with BUB-1[190-628] in Figure 2F-G, 200 μl of binding

assay was loaded into a 0.5 ml loop and loaded onto the column with 2 ml of SEC buffer. For

the CENP-C[HCP-4(1-214)] assays in Figure 5B and Figure S5, 100 μl of assay was loaded onto

a100 μl loop and loaded onto the column with 0.4 ml of SEC buffer.

1046

Fluoresence Polarisation

Fluoresence polarisation assays were conducted using FITC-labelled peptides (peptide

sequences indicated in figures). For all assays: a 1:2 dilution series of MBP-PLK1[PBD] was

conducted in FP buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP) with a

constant concentration of 100 nM FITC-labelled peptide. Assays were left for 10-30 min

before briefly centrifuged and 10 μl of each concentration was loaded onto a black 384 well

plate (Greiner) in triplicate. Plates were then centrifuged (2k rpm, 2 min) before being

analysed in a PheraStar FS (BMG Labtech) under the following conditions: excitation: 485

nm, emission: 520 nm, 25°C, positioning delay 0.3 s, 50 flashes per well, 9.3 mm focal

height. Average mP for each triplicate was then plotted against the MBP-PLK1[PBD]

concentration using and a non-linear curve fitted using GraphPad Prism 'One Site – Total'

equation.

1059

Phenotype Analysis

We defined misalignment as metaphase defects. Alignment was counted within the 5 frames

(1 min) before anaphase onset, which was the frame prior to the detection of two separating

chromosome masses. When one bivalent was misaligned (either in angle or distance to

metaphase plate) it was recorded as mild metaphase defect. When two or more bivalents were

misaligned, this was considered as a severe metaphase defect.

The anaphase phenotype was assess during segregation. If lagging chromosome were detected

during chromosome segregation but this was resolved before polar body extrusion, it was

scored as mild anaphase defect. If we could not detect two segregating masses of

chromosomes or if these masses differed in size/intensity, it was quantified as a severe

anaphase defect.

Polar body defect was recorded when no polar body was extruded or if all of the maternal

DNA content was extruded as a polar body, with no maternal DNA remaining in the

cytoplasm.

Graphs were prepared using Graphpad Prism 9.0.

1075

1076   **Table S1**: *C. elegans* strains used in this study

GFP::tubulin; mCherry::histone	OD868	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I*	(Green et al., 2013) PMID: 24217623
GFP::HCP-4	OD3410	*hcp-4(lt72[GFP::hcp-4])I*	(Cheerambathur et al., 2019) PMID: 30827898
PLK-1::sfGFP	OD2425	*plk-1(lt18[plk-1::sGFP]::loxp)III*	(Martino et al., 2017) PMID: 29065307
GFP::ASPM-1	EU2861	*or1935[GFP::aspm-1]I*	(Connolly et al., 2015) PMID: 26370499
PLK-1::sfGFP; mCherry::histone	FGP263	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV*	This study
hcp-4(T163A)	FGP669	*hcp-4(fgp58[hcp-4(T163A)])I*	This study
bub-1(T527A)/hT2	FGP672	*bub-1(fgp4[bub-1(T527A)])I; hT2 [bli-4(e937) let-?(q782) qIs48](I;III)*	This study
plk-1(as)	OD3697	*plk-1((lt106[plk-1 C52V] lt109[plk-1 L115G])III*	(Gómez-Cavazos et al., 2020) PMID: 32619481
GFP::tubulin; mCherry::histone; bub-1(T527A)/hT2	FGP674	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; bub-1(fgp4[bub-1(T527A)])I; hT2 [bli-4(e937) let-?(q782) qIs48](I;III)*	This study

GFP::tubulin; mCherry::histone; plk-1(as)	FGP675	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; plk-1((lt106[plk-1 C52V] lt109[plk-1 L115G])III*	This study
PLK-1::sGFP; mCherry::histone; hcp-4(T163A)	FGP719	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV; hcp-4(fgp58[hcp-4(T163A)])I*	This study
GFP::ASPM-1; mCherry::histone; plk-1(as)	FGP725	*or1935[GFP::aspm-1] I; itIs37[pie-1p::mCherry::H2B::pie-1 3'UTR + unc-119(+)] IV; plk-1((lt106[plk-1 C52V] lt109[plk-1 L115G])III*	This study
GFP::HCP-4; mCherry::histone; TIR1	FGP311	*hcp-4(lt72[GFP::hcp-4])I; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV)I; ieSi65 [sun-1p::TIR1::sun-1 3'UTR + Cbr-unc-119(+)] II; unc-119(ed3)III*	This study

1077

Figure 1

Figure 2

A — PLK-1::sfGFP; mCherry::histone

B — PLK-1, *wild type*, *bub-1(RNAi)*; PLK, histone; position (μm); Relative intensity

C — BUB-1; TPR, CDC-20, MAD-1, Kinase; BUB-3, B56, PLK-1; 1 ... 987

C.japonica	527	TRGNIVTSTPAGSIPFM
C.brenneri	539	SRGMIVTSTPAHNVPYV
C.elegans	519	SRGNFVTSTPAQGVHLV
C.tropicalis	536	PHGMFVTSTPAHNVPFV
C.nigoni	520	KQGSIVTSTPAAGVAFV
C.briggsae	520	THGSIVTSTPAAGVAFV
C.latens	542	TRGGIVTSTPAAGIPFI
C.remanei	541	TRGGIVTSTPAAGIPFI

D — Cdk1/CycB: − + − +; PLK-1: − − + +; Coomassie; Pro-Q™ Diamond

E — **Cdk1/Cyclin B + PLK-1 kinase assay**; BUB-1^190-628; BUB-1^190-628(T527A); t(min): 0 30 60 90 120; Coomassie; Pro-Q™ Diamond

F — S200 10/300; Relative Absorbance; Vol (ml); MBP:PLK1^PBD; BUB-1^190-628 + Cdk1; BUB-1^190-628; BUB-1^190-628 + MBP:PLK1^PBD + Cdk1; BUB-1^190-628 + MBP:PLK1^PBD

G — S200 10/300; Relative Absorbance; Vol (ml); BUB-1^190-628 + MBP:PLK1^PBD + Cdk1; BUB-1^190-628 (T527A)+ MBP:PLK1^PBD + Cdk1

H — RGNFVTSTPAQGVH; RGNFVTS(T^Ph)PAQGVH; Polarisation (mP); MBP:PLK1^PBD (μM)

Figure 3

Figure 4

A

CENP-C^HCP-4   histone

Prometaphase

Metaphase

Anaphase

GFP::CENP-C^HCP-4;
mCherry::histone

B

CENP-C^HCP-4

PLK-1

Mis12
interaction
KMN network

CENP-A
interaction

C.japonica   63 NAGPSCSTPIISGKDA
C.elegans   156 KAGLSCSTPKSSSDT
C.remanei   158 KAGLSCSTPKSGKDV
C.latens   158 KAGLSCSTPKSGNDL
C.brenneri   157 KAGLSCSTPKSANDL
C.briggsae   126 KAGLSCSTPKSANDV

C

	histone	PLK-1
wild type		
hcp-4(RNAi)		

PLK-1::sfGFP; mCherry::histone

D

PLK-1

wild type

— PLK — histone

Relative intensity

hcp-4(RNAi)

Relative intensity

position (μm)

Figure 5

A Cdk1/Cyclin B kinase assay

CENP-C$^{HCP-4(1-214)}$ CENP-C$^{HCP-4(1-214)}$(T163A)

t(min): 0 2 4 8 15 25 35 45 60   0 2 4 8 15 25 35 45 60

Coomassie

Pro-Q™ Diamond

B S200 10/300

MBP:PLK1^{PBD}

CENP-C$^{HCP-4(1-214)}$ + Cdk1

CENP-C$^{HCP-4(1-214)}$ (T163A) + Cdk1

CENP-C$^{HCP-4(1-214)}$ + MBP:PLK1^{PBD} + Cdk1

CENP-C$^{HCP-4(1-214)}$ (T163A) + MBP:PLK1^{PBD} + Cdk1

C KAGLSCSTPKSSSD   KAGLSCS(T^{Ph})PKSSSD

D PLK-1   histone PLK-1   PLK-1

wild type

hcp-4^{T163A}

PLK-1::sfGFP; mCherry::histone

E PLK-1   PLK — DNA

wild type

hcp-4^{T163A}

F HCP-4   DNA HCP-4   DNA

wild type

hcp-4^{T163A}

G wild type   hcp-4^{T163A}

PLK-1

BUB-1   DNA

A

histone
PLK-1

PLK-1

histone
PLK-1

wild type

bub-1(RNAi)

hcp-4(RNAi)

bub-1(RNAi)
hcp-4(RNAi)

PLK-1::sfGFP;
mCherry::histone

B

wild type   hcp-4(RNAi)   bub-1(RNAi)   bub-1(RNAi)
                                                               hcp-4(RNAi)

-60"   -120"   -100"   -80"

0"   0"   0"   0"

100"   100"   100"   100"

200"   200"   200"   200"

420"   420"   400"   620"

GFP::tubulin; mCherry::histone

C

% oocytes

wild type   hcp-4(RNAi)   bub-1(RNAi)   bub-1(RNAi)
                                            hcp-4(RNAi)

Metaphase - Mild
Metaphase - Severe
Anaphase - Mild
Anaphase - Severe
PBE defect

D

PLK-1   DNA

wild type

hcp-4(RNAi)

bub-1^{T527A}

bub-1^{T527A}
hcp-4(RNAi)

E

wild type   hcp-4(RNAi)   bub-1^{T527A} bub-1^{T527A}
                                                                  hcp-4(RNAi)

-280"   -80"   -280"   -160"

-80"   -20"   -100"   -100"

180"   180"   140"   180"

300"   320"   240"   320"

GFP::tubulin; mCherry::histone

F

% oocytes

wild type   hcp-4(RNAi)   bub-1^{T527A} bub-1^{T527A}
                                                hcp-4(RNAi)

Metaphase - Mild
Metaphase - Severe
Anaphase - Mild
Anaphase - Severe
PBE defect

PLK-1   histone

PLK-1::sfGFP; mCherry::histone

A

wild type | *plk-1^as*

1 µM 3-IB-PP1 | 10 µM 3-IB-PP1

Δ20"

GFP::tubulin; mCherry::histone

B

plk-1^as + 1 µM 3IB-PP1

Δ20"

GFP::tubulin; mCherry::histone

A

Cdk1/Cyclin B kinase assay

BUB-1$^{190-628}$ BUB-1$^{190-628}$(T527A)

Time (min):   0   30   60   90   120   0   30   60   90   120

Coomassie

ProQ

B

PLK-1 kinase assay

BUB-1$^{190-628}$ BUB-1$^{190-628}$(T527A)

Time (min):   0   30   60   90   120   0   30   60   90   120

Coomassie

ProQ

A

histone PLK-1 PLK-1 histone PLK-1

wild type

bub-1(RNAi)

hcp-4^{T163A}

bub-1(RNAi) hcp-4^{T163A}

PLK-1::sfGFP; mCherry::histone

B

wild type   *hcp-4^{T163A}*   *bub-1(RNAi)*   *hcp-4^{T163A} bub-1(RNAi)*

GFP::tubulin; mCherry::histone

C

% oocytes

wild type   *bub-1(RNAi)*   *hcp-4^{T163A}*   *hcp-4^{T163A} bub-1(RNAi)*

Metaphase - Mild
Metaphase - Severe

Anaphase - Mild
Anaphase - Severe

PBE defect