BUB-1 and CENP-C recruit PLK-1 to Control Chromosome Alignment and Segregation

During Meiosis I in *C. elegans* Oocytes

Samuel J.P. Taylor[1,a], Laura Bel Borja[1,a], Flavie Soubigou[1], Jack Houston[2], Dhanya K.

Cheerambathur[3], and Federico Pelisch[1,*]

[1] Centre for Gene Regulation and Expression, Sir James Black Centre, School of Life

Sciences, University of Dundee. Dundee, DD1 5EH, United Kingdom

[2] Ludwig Institute for Cancer Research, San Diego Branch, La Jolla, CA 92093, USA

[3] Wellcome Centre for Cell Biology & Institute of Cell Biology, School of Biological

Sciences, The University of Edinburgh, Edinburgh, EH9 3BF, UK

[a] These authors contributed equally.

* Corresponding author: Federico Pelisch (f.pelisch@dundee.ac.uk)

ABSTRACT

Phosphorylation is a key post-translational modification that is utilised in many biological processes for the rapid and reversible regulation of protein localisation and activity. Polo-like kinase 1 (PLK-1) is essential for both mitotic and meiotic cell divisions, with key functions being conserved in eukaryotes. The roles and regulation of PLK-1 during mitosis have been well characterised. However, the discrete roles and regulation of PLK-1 during meiosis have remained obscure. Here, we used *Caenorhabditis elegans (C. elegans)* oocytes to show that PLK-1 plays distinct roles in meiotic spindle assembly and/or stability, chromosome alignment and segregation, and polar body extrusion during meiosis I. Furthermore, by a combination of live imaging and biochemical analysis we identified the chromosomal recruitment mechanisms of PLK-1 during *C. elegans* oocyte meiosis. The spindle assembly checkpoint kinase BUB-1 directly recruits PLK-1 to the kinetochore and midbivalent while the chromosome arm population of PLK-1 depends on a direct interaction with the centromeric-associated protein CENP-C[HCP-4]. We found that perturbing both BUB-1 and CENP-C[HCP-4] recruitment of PLK-1 leads to severe meiotic defects, resulting in highly aneuploid oocytes. Overall, our results shed light on the roles played by PLK-1 during oocyte meiosis and provide a mechanistic understanding of PLK-1 targeting to meiotic chromosomes.

INTRODUCTION

Oocyte meiosis consists of two consecutive segregation events following DNA replication –

in meiosis I homologous chromosomes segregate, half of which are then removed in a polar

body. The remaining sister chromatids segregate in meiosis II, half of which are removed in a

second polar body to produce a haploid gamete (Marston and Amon, 2004; Ohkura, 2015).

Tight spatial and temporal control of protein localisation and activity is required to ensure

chromosome/chromatid alignment and segregation occur efficiently and the correct number of

chromosomes is present in each gamete (Marston and Wassmann, 2017). Phosphorylation is a

key post-translational modification utilised to regulate protein localisation and activity, which

is fundamentally important for the success of both mitotic and meiotic cell divisions (Marston

and Wassmann, 2017; Saurin, 2018). Therefore, the regulation of kinase or phosphatase

localisation and activity is vital for proper cell division, with the balance of their effects

determining the localisation/activity of substrate proteins that play important roles in the cell

division process (Gelens et al., 2018; Novak et al., 2010).

Polo-like kinases (PLKs) are a family of Ser/Thr protein kinases first discovered in

Drosophila (Llamazares et al., 1991; Sunkel and Glover, 1988) and yeast (Kitada et al., 1993;

Ohkura et al., 1995) and later findings showed that PLKs are present in all eukaryotes

(Zitouni et al., 2014). PLK-1 is essential for meiotic and mitotic cell divisions, with its

localisation and functions proving well conserved throughout eukaryotic evolution. PLK-1

localises to the nuclear envelope in prophase (Linder et al., 2017) and to the centrosomes,

kinetochore, central spindle, and midbody during mitosis and is involved in numerous

processes including mitotic entry, spindle assembly, chromosome alignment, the spindle

assembly checkpoint, and cytokinesis (Archambault and Glover, 2009; Petronczki et al.,

2008; Schmucker and Sumara, 2014; Zitouni et al., 2014). During mammalian meiosis, PLK1

localises to the chromosomes and spindle poles in prometaphase and metaphase. In anaphase

PLK1 is localised primarily in the central spindle between the segregating chromosomes and

then the midbody during polar body extrusion (Pahlavan et al., 2000; Solc et al., 2015; Tong

et al., 2002; Wianny et al., 1998). Inhibition of PLK1 in mammalian oocytes has indicated

roles in many processes including germinal vesicle breakdown, spindle assembly,

chromosome alignment, and polar body extrusion (Solc et al., 2015; Tong et al., 2002). To

understand why PLK-1 is crucial for meiosis it is critical to characterise precisely where PLK-

1 is located and how it is recruited to specific regions, which will allow further dissection of

the discrete roles of PLK-1 during meiosis.

PLK-1 interacts with proteins via its C-terminal polo-binding domain (PBD) (Cheng, 2003;

Elia et al., 2003a). The PBD binds to phosphorylated motifs of the consensus Ser-

phSer/phThr-X, where ph indicates a phosphorylated residue and X indicates any amino acid

(Elia et al., 2003a, 2003b). When X is a proline, phosphorylation of the central Ser/Thr

residue is often mediated by a proline-directed kinase, notably Cdk1/Cyclin B during cell

division (Elowe et al., 2007; Qi et al., 2006) - this motif will henceforth be termed STP motif.

When a non-proline residue occupies position X, PLK-1 itself can phosphorylate the central

Ser/Thr, thereby enhancing its own recruitment (Kang et al., 2006; Neef et al., 2003), referred

to as self-priming. Furthermore, PLK-1 binding to STP motifs via the PBD induces a

conformational change that enhances its kinase activity (Mundt et al., 1997; Xu et al., 2013).

While the mechanism of PLK-1 recruitment to the chromosomes during oocyte meiosis has

not been characterised, recruitment of PLK1 to the kinetochore during mammalian mitosis

has been investigated. In mammals, the constitutive centromere-associated network (CCAN)

complex of proteins binds to the histone variant CENP-A at centromeres (Foltz et al., 2006;

Izuta et al., 2006; Okada et al., 2006). Outer kinetochore proteins bind to the CCAN and

ultimately mediate chromosome alignment and segregation via interaction with microtubules

(Musacchio and Desai, 2017). Two proteins are primarily responsible for PLK1 recruitment

to the kinetochore – CCAN component CENP-U (Kang et al., 2006, 2011; Singh et al., 2021)

and the spindle assembly checkpoint kinase BUB-1 (Elowe et al., 2007; Qi et al., 2006a;

Singh et al., 2021), both of which directly bind to PLK1 via STP motifs in a Cdk1-dependent

manner.

In *C. elegans,* PLK-1 localisation in meiosis and mitosis is similar to other organisms (Chase

et al., 2000). In mitosis, roles of PLK-1 include nuclear envelope breakdown (NEBD)

(Martino et al., 2017), merge of parental genomes in the embryo through lamina disassembly

(Rahman et al., 2015; Velez-Aguilera et al., 2020), centrosome maturation (Cabral et al.,

2019; Decker et al., 2011; Ohta et al., 2021; Woodruff et al., 2015), and cytokinesis (Gómez-

Cavazos et al., 2020) – indicating that major mitotic roles of PLK-1 are conserved in *C.*

elegans. However, meiotic roles of PLK-1 in *C. elegans* oocytes have remained obscure, as

PLK-1 depletion results in severely defective NEBD and fertilised oocytes with a whole

nucleus rather than condensed chromosomes (Chase et al., 2000). In the PLK-1-depleted

oocytes that 'escaped' the NEBD defect, chromosome congression, segregation, and polar

body extrusion were severely disrupted. However it is unclear whether these phenotypes are

indirect effects of the severe early meiotic defects or whether they result from specific

functions of PLK-1 throughout meiosis (Chase et al., 2000). Furthermore, while PLK-1 was

shown to localise broadly to chromosomes and the spindle during meiosis (Chase et al.,

2000), a more precise dynamic characterisation of PLK-1 localisation during meiosis is

lacking.

Here, by temporally inhibiting an analogue-sensitive PLK-1 mutant we show that PLK-1 is

involved in spindle assembly and/or stability, chromosome alignment and segregation, and

polar body extrusion in *C. elegans* oocytes. Using live imaging and immunofluorescence, we

find that PLK-1 localises to the spindle poles, chromosome arms, kinetochores, and

midbivalent region between the homologous chromosomes during meiosis I in *C. elegans*

oocytes. By a combination of live imaging and *in vitro* biochemical analysis, we have

characterised the primary chromosomal recruitment mechanisms of PLK-1 during meiosis –

showing that CENP-C[HCP-4] directly recruits PLK-1 to the chromosome arms while PLK-1

recruitment to the midbivalent and kinetochore is mediated by a direct interaction with BUB-

1. Furthermore, BUB-1- and CENP-C-mediated PLK-1 recruitment to chromosomes is

essential for meiosis I.

RESULTS

PLK-1 plays roles in spindle assembly and/or stability, chromosome alignment and segregation, and polar body extrusion in meiosis I

A previous study showed that PLK-1 localises to the meiotic spindle and chromosomes in *C. elegans* oocytes and depletion of PLK-1 using RNAi led to several defects including defective NEBD, chromosome congression, segregation, and polar body extrusion (Chase et al., 2000). While this suggested that PLK-1 plays several roles during meiosis, the use of RNAi presents a limitation to addressing them independently. In particular, the strong NEBD defect complicates delineation of the roles of PLK-1 at later stages of meiosis. Therefore, we sought to understand the distinct localisation and roles of PLK-1 during meiosis I.

To assess PLK-1 localisation during meiosis with high spatial and temporal resolution, we imaged endogenously tagged PLK-1::sfGFP in dissected oocytes. PLK-1 localises to the spindle poles (Figure 1C; blue arrows), chromosome arms (Figure 1A,C; yellow arrows), and midbivalent region between the homologous chromosomes (Figure 1A,C; magenta arrow) during Prometaphase I. This localisation pattern was confirmed with immunostaining of fixed oocytes using a specific anti-PLK-1 antibody (Figure 1B). As chromosomes begin to segregate in early anaphase, PLK-1 is mostly observed on chromosomes and, to a lesser extent, in between the segregating chromosome masses (Figure 1C, yellow and magenta arrows, respectively; see also Figure 1-figure supplement 1). During late anaphase, PLK-1 is still detectable on chromosomes, and is enriched in the central spindle (Figure 1C, green arrow; see also Figure 1-figure supplement 1).

Since long-term depletion of PLK-1 leads to severe NEBD defects (Chase et al., 2000), we used an analogue-sensitive *plk-1* allele (Gómez-Cavazos et al., 2020; Woodruff et al., 2015) that renders it sensitive to chemically modified derivatives of PP1, a Src family inhibitor (Bishop et al., 2000). We reasoned that acute PLK-1[as] inhibition for a short period of time

would allow us to study the post-NEBD effects (Figure 1D). We tested a variety of analogues

and all of them led to embryonic lethality of the *plk-1^{as}* strain without severely affecting a

wild type strain (Figure 1-figure supplement 2). We decided to continue our experiments with

the 3-substituted benzyl PP1 derivative 3IB-PP1 (Figure 1-figure supplement 2). A wild type

strain in the presence of 3IB-PP1 and *plk-1^{as}* in the presence of vehicle control ('EtOH')

behaved normally during meiosis (Figure 1E,F & Figure 1-figure supplement 3). Addition of

10 µM 3IB-PP1 between 5 and 15 min before dissection and imaging of oocytes allowed us to

bypass the NEBD defect and 6 bivalents were easily identifiable within the newly fertilised

oocyte (Figure 1E, yellow arrows). Under these conditions PLK-1^{as} inhibition led to drastic

spindle defects with no observable bipolar spindle formation and no consequent chromosome

segregation was observed, indicating that PLK-1 is involved in spindle assembly and/or

stability during oocyte meiosis (Figure 1E). We sought to minimise the spindle defects upon

PLK-1 inhibition by reducing the concentration of 3IB-PP1 to 1 µM and omitting the pre-

treatment step prior to dissection. Under these conditions, ~62% of oocytes had seemingly

bipolar spindles and chromosomes remained somewhat associated with the spindle, although

chromosome alignment was still affected in 78% of oocytes (≥2 misaligned chromosomes,

Figure 1F,G). A more detailed analysis of chromosome dynamics after PLK-1 inhibition is

presented in Figure 1-figure supplement 3, where individual chromosomes are followed every

20 seconds and, as opposed to wild type, chromosomes from PLK-1-inhibited oocytes show a

highly dynamic behaviour whereby they seem to briefly align and then become misaligned

again (Figure 1-figure supplement 3 and arrows therein). We then used the pole marker

ASPM-1 to allow proper characterisation of spindle bipolarity under these conditions and

confirmed that even when two ASPM-1 poles are clearly discerned (Figure 1H, blue arrows),

chromosome alignment fails (Figure 1H, yellow arrows). Hence, it appears that PLK-1

participates in chromosome alignment in a manner that is at least partially independent of its

roles in overall spindle assembly and/or stability.

BUB-1 recruits PLK-1 to the midbivalent during oocyte meiosis

To further understand the role of PLK-1 during meiosis, we sought to identify the PLK-1

recruitment mechanism(s). In mammalian mitosis, the kinase BUB-1 and its paralog BUBR1

directly recruit PLK-1 to the kinetochore via STP motifs that are phosphorylated by Cdk1

(Elowe et al., 2007; Qi et al., 2006). While the *C. elegans* BUBR1 ortholog MAD3[SAN-1] does

not localise to the chromosomes or spindle during meiosis (Bel Borja et al., 2020), BUB-1

localises to the kinetochores and midbivalent region (Dumont et al., 2010; Monen et al., 2005;

Pelisch et al., 2019, 2017). We investigated whether BUB-1 was involved in PLK-1 targeting

during *C. elegans* meiosis. RNAi-mediated depletion of BUB-1 led to the loss of PLK-1 from

the midbivalent (Figure 2A,B; blue arrows). In contrast, PLK-1 signal at chromosome arms

remained unaffected (Figure 2A,B; yellow arrows). Analysis of the BUB-1 protein sequence

revealed a putative polo-docking STP motif in amino-acids 526-528 that is conserved in

nematode species (Figure 2C). Therefore, we sought to identify whether *C. elegans* BUB-1

directly interacts with PLK-1 to mediate its recruitment to the midbivalent region.

BUB-1 directly interacts with PLK-1 through a Cdk1-dependent STP motif

To test whether BUB-1 directly interacts with PLK-1 *in vitro*, we purified a recombinant

fragment of BUB-1 encompassing the intrinsically disordered region between the TPR and

kinase domains that contains the putative STP motif ('BUB-1[190-628]', Figure 2D). Since the

interaction between STP motifs and the PBD of PLK-1 requires phosphorylation of the

central Ser/Thr residue (Elia et al., 2003a, 2003b), we conducted kinase assays to assess the

phosphorylation of BUB-1[190-628]. Cdk1 and PLK-1 can both phosphorylate BUB-1[190-628] and

Cdk1 can specifically phosphorylate the STP motif at T527 (Figure 2D,E & Figure 2-figure

supplement 1). Interestingly, kinase assays conducted with Cdk1 and PLK-1 together

produced a prominent shifted band representing phosphorylated protein that was not present

with the individual kinases (Figure 2D). Since STP motifs are known to be targets of proline-

directed kinases such as Cdk1 (Elowe et al., 2007; Qi et al., 2006), we hypothesised that

phosphorylation of T527 by Cdk1 enables PLK-1 to bind directly to BUB-1 resulting in the

shifted band representing highly phosphorylated BUB-1. To test this hypothesis, we mutated

T527 in the STP motif to alanine ('BUB-1$^{190\text{-}628}$(T527A)') and assessed the resulting

phosphorylation (Figure 2F). T527A mutation in BUB-1$^{190\text{-}628}$ largely prevented the shift

observed in the combined Cdk1 and PLK-1 assay (Figure 2F), indicating that phosphorylation

of this residue is essential for the shifted band observed.

To determine whether BUB-1 directly binds to PLK-1, we purified a maltose-binding protein

(MBP)-tagged PLK1 PBD (MBP-PLK1^{PBD}) (Singh et al., 2021) and incubated it with

unphosphorylated or Cdk1-phosphorylated BUB-1$^{190\text{-}628}$ before assessing complex formation

by size-exclusion chromatography (SEC). Cdk1-phosphorylated BUB-1 formed a stable

complex with MBP-PLK1^{PBD} (Figure 3A). SEC coupled with multi-angle light scattering

(SEC-MALS) indicated the stoichiometry of this complex is 1:1 (Figure 3B). While

unphosphorylated BUB-1 showed some interaction with MBP-PLK1^{PBD}, this complex eluted

from the column at a higher volume and bound to a lower proportion of MBP-PLK1^{PBD},

suggestive of a weaker interaction (Figure 3A). To directly test whether phosphorylation of

T527 is required for PLK-1 to bind to BUB-1, MBP-PLK1^{PBD} was incubated with Cdk1-

phosphorylated BUB-1$^{190\text{-}628}$ or BUB-1$^{190\text{-}628}$(T527A) and SEC was used to assess complex

formation. Interestingly, when Cdk1-phosphorylated BUB-1$^{190\text{-}628}$(T527A) was incubated

with MBP-PLK1^{PBD} the resulting elution was reminiscent of the unphosphorylated wild type

BUB-1$^{190\text{-}628}$, with a reduced proportion of MBP-PLK1^{PBD} binding and elution occurring at a

higher volume (Figure 3C). Together, these data indicate that there is a Cdk1

phosphorylation-dependent interaction between BUB-1 and PLK1[PBD] that requires

phosphorylation of T527 within the STP motif of BUB-1.

To further confirm that the STP motif of BUB-1 can interact with PLK1[PBD] in a phT527-

dependent manner, FITC-labelled peptides containing the BUB-1 STP motif were incubated

with either wild type MBP-PLK1[PBD] or a mutant that cannot interact with phSTP motifs

('MBP-PLK1[PBD](H538A/K540M)') (Elia et al., 2003b). The phT527 peptide bound to MBP-

PLK1[PBD] with a K_D of 55 ± 3 nM but did not bind to MBP-PLK1[PBD](H538A/K540M), while

the unphosphorylated peptide did not interact with MBP-PLK1[PBD] at the concentrations tested

(Figure 3D).

Altogether, these data indicate that *C. elegans* BUB-1 can directly bind to PLK1[PBD] *in vitro* in

a phospho-dependent manner via a newly characterised STP motif.

BUB-1 directly recruits PLK-1 to the midbivalent *in vivo*

We then sought to determine whether the STP motif in BUB-1 is responsible for PLK-1

recruitment *in vivo.* Using CRISPR-Cas9, we generated the T527A mutation in the

endogenous *bub-1* locus (*bub-1[T527A]*). *bub-1[T527A]* mutant worms showed significant embryonic

and larval lethality so we generated a balanced strain in which the *bub-1[T527A]* allele was

maintained as a heterozygote. Homozygous *bub-1[T527A]* worms from heterozygous parents

develop to adulthood and produce oocytes that go through meiosis, which allowed us to study

the role of the STP motif in BUB-1 during meiosis. PLK-1 was absent from the midbivalent

in *bub-1[T527A]* oocytes, reminiscent of the *bub-1(RNAi)* phenotype (Figure 4A, blue

arrowheads, and 4B). Importantly, BUB-1 localisation to the midbivalent and kinetochore

was maintained in the *bub-1[T527A]* strain (Figure 4C), indicating that BUB-1 directly interacts

with PLK-1 via this STP motif *in vivo* to recruit PLK-1 to the midbivalent. To assess the

impact of BUB-1 mediated PLK-1 recruitment during meiosis I, we crossed the *bub-1^{T527A}*

allele with a strain expressing GFP-tagged tubulin and mCherry-tagged histone and analysed

chromosome alignment, segregation, and polar body extrusion defects (See Methods). *bub-*

1^{T527A} mutant oocytes displayed chromosome alignment defects in ~62% of the oocytes, with

32% of oocytes showing severe alignment defects (Figure 4D,E). Additionally, ~1/4 of *bub-*

1^{T527A} oocytes showed mild anaphase defects (Figure 4E). Despite these defects, more than

90% of *bub-1^{T527A}* oocytes show visible separation of two chromosome masses and polar

body extrusion occurred normally at the end of meiosis I (Figure 4E).

Overall, our results show that an STP motif in BUB-1 directly recruits PLK-1 to the

midbivalent during meiosis I and this interaction is primarily important for chromosome

alignment.

CENP-C^{HCP-4} recruits PLK-1 to meiotic chromosome arms

PLK-1 localisation to chromosome arms remained unchanged when BUB-1 was depleted

(Figure 2A,B) or the STP motif was mutated (Figure 4A,B), indicating that a different

pathway is required to recruit this population of PLK-1. We therefore sought to identify the

mechanism of PLK-1 recruitment to the chromosome arms. In mammalian mitosis, PLK1

recruitment to the kinetochore is mediated by BUB-1 and CCAN component CENP-U (Elowe

et al., 2007; Kang et al., 2006, 2011; Qi et al., 2006b; Singh et al., 2021). Interestingly, the

CCAN appears to be largely absent in *C. elegans* (Maddox et al., 2012), and kinetochore

assembly depends on the CENP-C orthologue, HCP-4 (hereafter CENP-C^{HCP-4}) during mitosis

(Oegema et al., 2001) and on CENP-C^{HCP-4} and the nucleoporin ELYS^{MEL-28} during meiosis

(Hattersley et al., 2022). CENP-C^{HCP-4} localises to chromosomes throughout meiosis I (Figure

5A) (Hattersley et al., 2022; Monen et al., 2005). Like its mammalian counterpart, CENP-

C^{HCP-4} is predicted to be mostly disordered and it contains a putative N-terminal STP motif

encompassing amino acids 162-164 that is conserved in nematode species (Figure 5B).

Although CENP-C[HCP-4] depletion does not have a major impact on meiosis I (Hattersley et al.,

2022; Monen et al., 2005), RNAi-mediated depletion of CENP-C[HCP-4] abolished PLK-1

localisation on chromosome arms (Figure 5C,D). PLK-1 is still present in the midbivalent

(Figure 5C,D, blue arrows) which suggests that BUB-1 and CENP-C[HCP-4] represent

independent pathways for PLK-1 targeting. Additionally, CENP-C[HCP-4] depletion revealed a

pool of PLK-1 which is kinetochore-associated (Figure 5C,D, yellow arrows). To confirm that

PLK-1 localises to the kinetochore, we took advantage of the fact that kinetochore proteins in

C. elegans oocytes are also present in filaments referred to as 'linear elements', which are

spindle- and cortex-associated (Monen et al., 2005). PLK-1 perfectly co-localises with KNL-

3, part of the Mis12 kinetochore complex in spindle- and cortex-associated linear elements

(Figure 5-figure supplement 1). Since the above data indicated that CENP-C[HCP-4] is involved

in PLK-1 recruitment to chromosome arms, we sought to determine whether this involved a

direct interaction via the putative STP motif identified in sequence alignments.

CENP-C[HCP-4] interacts directly with PLK-1 through a Cdk1-dependent STP motif

To investigate whether CENP-C[HCP-4] directly interacts with PLK-1, we purified a recombinant

N-terminal fragment of CENP-C[HCP-4] ('CENP-C[HCP-4(1-214)]', Figure 5E). As the putative

CENP-C[HCP-4] STP motif is also a proline-directed kinase consensus site, we assessed Cdk1

phosphorylation of the recombinant fragment. Cdk1/Cyclin B kinase assays showed that

CENP-C[HCP-4(1-214)] can be phosphorylated by Cdk1 and mutation of the putative STP motif

threonine to alanine (T163A) reduced the phosphorylation of the fragment (Figure 5E,F).

Furthermore, Cdk1 specifically phosphorylated T163 within the putative STP motif (Figure

5G).

To determine whether CENP-C$^{HCP-4(1-214)}$ can directly bind to MBP-PLK1^{PBD}, we used size

exclusion chromatography. CENP-C$^{HCP-4(1-214)}$ formed a stable complex with MBP-PLK1^{PBD}

which was dependent on Cdk1-phosphorylation (Figure 6A). SEC-MALS indicated the

stoichiometry of this interaction was 1:1 (Figure 6B). We then tested whether phosphorylation

of the STP motif is required for the interaction with MBP-PLK1^{PBD}. Mutation of the putative

STP motif (T163A) largely prevented the Cdk1-mediated interaction with MBP-PLK1^{PBD},

indicating phosphorylation of the STP motif at T163 is required for the interaction (Figure

6C).

To further confirm this STP motif binds to MBP-PLK1^{PBD} in a phT163-dependent manner,

we conducted fluorescence polarisation assays with FITC-labelled CENP-C^{HCP-4} STP motif

peptides (Figure 6D). The phT163 peptide bound to MBP-PLK1^{PBD} with a K$_D$ of 104 ± 14

nM but did not interact with MBP-PLK1^{PBD}(H538A/K540M), while the unphosphorylated

peptide did not bind to MBP-PLK1^{PBD} at the concentrations tested (Figure 6D). Collectively,

these data indicate that the putative STP motif in CENP-C^{HCP-4} binds to the PBDs of PLK-1 in

a Cdk1 phosphorylation-dependent manner, which led us to assess the importance of this STP

motif *in vivo*.

CENP-C^{HCP-4} recruits PLK-1 to chromosome arms *in vivo* through an STP motif

The T163A mutation in CENP-C^{HCP-4} was generated in the endogenous *hcp-4* locus (*hcp-*

4^{T163A}) which, unlike *bub-1^{T527A},* did not affect viability. When PLK-1::sfGFP was monitored

in dissected oocytes, *hcp-4^{T163A}* recapitulated the full CENP-C^{HCP-4} depletion with PLK-1

localising to the midbivalent and kinetochore but largely absent from chromosome arms

(Figure 7A,B). This indicates that PLK-1 is targeted to chromosome arms directly through the

phospho-dependent STP motif in CENP-C^{HCP-4}. Importantly, CENP-C^{HCP-4}(T163A) displays

an indistinguishable localisation from wild type CENP-C^{HCP-4} (Figure 7C) and the other PLK-

1 receptor, BUB-1, also localises normally in the *hcp-4^{T163A}* strain (Figure 7D).

These data indicate that CENP-C^{HCP-4} directly recruits PLK-1 to the chromosome arms during

meiosis I via a newly characterised N-terminal STP motif.

Dual PLK-1 recruitment by BUB-1 and CENP-C is essential for meiosis I

Our results so far indicate that PLK-1 recruitment to the midbivalent during meiosis I is

mediated by direct interaction with BUB-1, while PLK-1 localisation to the chromosome

arms requires direct interaction with CENP-C^{HCP-4}. Co-depletion of BUB-1 and CENP-C^{HCP-4}

by RNAi led to a drastic reduction of total PLK-1 signal (15% median intensity of wild type

oocytes; Figure 8A, Figure 8-figure supplement 1). This confirmed that the BUB-1 and

CENP-C^{HCP-4} pathways are the primary recruiters of PLK-1 to the chromosomes during

oocyte meiosis. Additionally, the kinetochore population of PLK-1 was lost when BUB-1 and

HCP-4 were co-depleted, indicating that BUB-1 is responsible for recruiting PLK-1 to the

kinetochore as well as the midbivalent (Figure 8A). RNAi of BUB-1 causes significant

defects in spindle stability, chromosome alignment, and chromosome segregation during

meiosis (Dumont et al., 2010; Pelisch et al., 2019). However, co-depletion of BUB-1 and

CENP-C^{HCP-4} exacerbated the chromosome alignment and segregation errors and resulted in a

significantly higher proportion of polar body extrusion failures (Figure 8B-C). Therefore,

while CENP-C^{HCP-4} depletion does not have noticeable defects on its own, it enhances the

BUB-1 depletion phenotype. To focus more specifically on the effects of direct PLK-1

recruitment, we depleted CENP-C^{HCP-4} in *bub-1^{T527A}* mutant oocytes. This largely abolished

PLK-1 localisation in the same manner as co-depletion of BUB-1 and CENP-C^{HCP-4} (Figure

8D) as well as displaying defects in chromosome alignment, segregation, and polar body

extrusion of a greater severity than *hcp-4(RNAi)* or *bub-1^{T527A}* alone (Figure 8E,F). Similar

results were obtained when performing the complementary experiment using *hcp-4^{T163A}/bub-*

1(RNAi) oocytes (Figure 8-figure supplement 2).

To determine the specific, combined role of the STP motifs in BUB-1 and CENP-C^{HCP-4}, we

used a gene replacement strategy whereby RNAi-resistant wild type or T527A BUB-1

transgenes ('BUB-1^{wt}' and 'BUB-1^{T527A}', respectively) were expressed in the presence of

bub-1(RNAi) to deplete the endogenous protein (Figure 9-figure supplement 1A,B). We then

used this strategy in wild type or *hcp-4^{T163A}* mutant worms ('*hcp-4^{wt}*' and '*hcp-4^{T163A}*',

respectively) to assess the effects of the double STP mutant. PLK-1 localised normally to the

chromosomes and midbivalent in BUB-1^{wt}, but was completely absent from the midbivalent

in BUB-1^{T527A} (Figure 9A; Figure 9-figure supplement 2A, cyan arrows). In the *hcp-4^{T163A}*

mutant PLK-1 was absent from chromosome arms when BUB-1^{wt} was expressed (Figure 9A;

Figure 9-figure supplement 2A, yellow arrows), and it was largely absent from all

chromosome-associated domains in the presence of BUB-1^{T527A} (Figure 9A; Figure 9-figure

supplement 2A).

We then used the same approach to assess the combined effect of the double mutant *bub-*

1^{T527A}/hcp-4^{T163A} on chromosome alignment and segregation, imaging oocytes expressing

GFP-tagged tubulin and mCherry-tagged histone. While mild defects were observed in the

bub-1^{T527A} mutant (and to a lesser extent in the *hcp-4^{T163A}* mutant), the double *bub-1^{T527A}/hcp-*

4^{T163A} mutant displayed severe meiotic defects, with 95% of oocytes displaying severe

metaphase defects, ~70% displaying severe anaphase defects, and ~73% of oocytes failing to

achieve PBE (Figure 9B,C; Figure 9-figure supplement 2).

These data indicate that recruitment of PLK-1 to the midbivalent and kinetochore by BUB-1

appears to be primarily responsible for the chromosomal roles of PLK-1 during meiosis I, as

disruption of this pathway leads to meiotic defects while perturbing CENP-C^{HCP-4} recruitment

of PLK-1 on its own does not. However, the fact that disruption of both BUB-1 and CENP-

C^{HCP-4} recruitment pathways enhances the severity of the resulting meiotic defects indicates

that the CENP-C^{HPC-4} pathway does still play an active part in the roles of PLK-1 during

meiosis I.

DISCUSION

In the present manuscript we describe specific, post-NEBD roles played by PLK-1 during

oocyte meiosis. PLK-1 is important for spindle assembly and/or stability, chromosome

alignment and segregation, and polar body extrusion during meiosis I. We found that PLK-1

localises to spindle poles and chromosomes during metaphase I before localising to the

chromosomes and central spindle in anaphase. Furthermore, we characterised the mechanisms

of chromosomal PLK-1 targeting during oocyte meiosis, which rely on the centromere-

associated protein CENP-C^{HCP-4} and the spindle assembly checkpoint kinase BUB-1. While

CENP-C^{HCP-4} targets PLK-1 to chromosome arms, BUB-1 directs PLK-1 to the midbivalent

and kinetochores. In both cases, interaction with PLK-1 relies on phosphorylated STP motifs

within predicted disordered regions. While we have not confirmed that these sites are

phosphorylated by Cdk1 *in vivo,* several lines of evidence indicate this is likely the case: 1)

both motifs have Pro at position 3, indicative of potential proline-directed kinase substrates;

2) Cdk1 is the most prominent proline-directed kinase in cell division and known to

phosphorylate STP motifs, including that of mammalian BUB1 (Qi et al., 2006); 3) both sites

were phosphorylated by Cdk1 *in vitro*; and 4) fluorescence polarisation experiments indicate

that phosphorylation of the Thr residues within the STP motifs is essential for the interaction

with the PBD, clearly displaying why the alanine mutants have such a drastic effect on PLK-1

localisation *in vivo*. We found that mutating the STP motif in BUB-1 results primarily in

chromosome alignment defects. While disrupting the CENP-C^{HCP-4}-mediated localisation of

PLK-1 to the chromosome arms does not have a significant phenotypic defect on its own, it

does enhance the meiotic defects observed when the BUB-1-dependent kinetochore and

midbivalent populations are disrupted. This suggests that while BUB-1 recruitment of PLK-1

may mediate the most important functions of PLK-1 during meiosis I, CENP-C^{HCP-4}

recruitment of PLK-1 to the chromosome arms also plays an active role in meiosis.

Expanding on the meiotic roles of PLK-1

By temporally inhibiting an analogue-sensitive PLK-1 mutant during *C. elegans* oocyte

meiosis, we have shown that PLK-1 plays a major role in the regulation of the meiotic

spindle. At high concentrations of analogue this resulted in the complete lack of spindle

bipolarity, and even at low concentrations the majority of oocytes (62%) imaged lacked an

apparent bipolar spindle. While this clearly displays a key role of PLK-1 during meiosis is in

the regulation of the meiotic spindle, we cannot distinguish more specific mechanisms using

our experimental techniques. As a result, we have characterised the phenotype as a lack of

spindle stability throughout this manuscript, but it should be noted that we cannot distinguish

whether these effects are on the spindle assembly process itself or on the maintenance of an

assembled bipolar spindle. Potential mechanisms of meiotic spindle regulation by PLK-1

include the microtubule depolymerase KLP-7, the *C. elegans* ortholog of the MCAK/Kinesin

13 family. There is some evidence to indicate that PLK-1 may regulate the kinesin 13

microtubule depolymerases in other organisms (Jang et al., 2009; Ritter et al., 2014; Sanhaji

et al., 2014; Shao et al., 2015; Zhang et al., 2011) and KLP-7 localises to the chromosomes,

during meiosis I (Connolly et al., 2015; Danlasky et al., 2020; Gigant et al., 2017; Han et al.,

2015), which overlaps with PLK-1 localisation. Furthermore, disrupting KLP-7 function

prevents proper bipolar spindle assembly and results in microtubules protruding out of the

meiotic spindle towards the cytoplasm (Connolly et al., 2015; Gigant et al., 2017), two

phenotypes we also see with PLK-1 inhibition.

Aside from the large-scale spindle defects observed when PLK-1 is inhibited, chromosome

alignment is still disrupted when the structure of the spindle appears bipolar and largely

normal. Since the mechanism of chromosome alignment by acentrosomal meiotic spindles is

not as well characterised as the centrosomal mitotic equivalent, speculating on the

underpinning mechanisms of this phenotype is challenging. However, it should be noted that

we cannot exclude the possibility that this phenotype is also a direct result of the dysregulated

spindle upon PLK-1 inhibition. Indeed, there is some evidence to suggest that a chromosome-

dependent pathway of microtubule formation may be an important aspect of chromosome

alignment and segregation (Conway et al., 2022; Heald et al., 1996; Kiewisz et al., 2022).

Despite these specific hypotheses mentioned above, there will be many different proteins and

pathways impacted by PLK-1 phosphorylation throughout meiosis I that will ultimately

contribute to the severe defects we observe upon PLK-1 inhibition. While a focussed

investigation into specific hypotheses would no doubt yield important results, an unbiased

approach to identify the relevant PLK-1 substrates during meiosis would be particularly

useful for investigation of the key effects of PLK-1 during meiosis. There are obvious

technical challenges to overcome before this can be achieved, not least of which would be

isolating a large enough sample of meiotic oocytes to perform robust quantitative proteomics.

The work in this manuscript undertaken to identify the mechanisms of PLK-1 targeting during

oocyte meiosis I will be instrumental for a later characterisation of the localisation and

meiotic stage-specific analysis of PLK-1 substrates.

Comparison with dual recruitment in mammals (CENP-U vs CENP-C)

During mammalian mitosis, PLK1 is recruited to kinetochores through BUB1 and CENP-U,

relying on self-priming in addition to Cdk1-mediated priming (Kang et al., 2006, 2011; Qi et

al., 2006b; Singh et al., 2021). Our results suggest that Cdk1-mediated priming is the primary

mechanism for PLK-1 recruitment in both BUB-1- and CENP-C-dependent branches.

Additionally, we noted the presence of a putative B56 short linear motif (LxxIxE) 38 aa

downstream of the STP motif in *C. elegans* CENP-C (203-IPTILE-208). This is relevant

because it has been shown that PLK1 and B56 motifs tend to co-exist in close proximity

(Cordeiro et al., 2020; Singh et al., 2021). This makes a putative cross-talk between PLK-1

and PP2A/B56 a worthy avenue to follow-up on our current findings.

Apparent lack of a CCAN network along with retention of crucial roles for PLK1 in species

like *C. elegans* and *Drosophila melanogaster* (*D. melanogaster*) suggest that the CENP-U

pathway could have been replaced by other proteins. CENP-C is a good candidate, as the only

CCAN component in these species. While we confirm that this is the case in *C. elegans*, it is

interesting to note that putative STPs exist in sequences 176-178 and 266-268 in *D.*

melanogaster CENP-C (Uniprot #Q9VHP9). Interestingly, *D. melanogaster* Polo and CENP-

C co-localise and this co-localisation increases by ectopic centromere generation through

CENP-A (CID) over-expression (Heun et al., 2006).

Overall, our results advance our understanding on the roles played by PLK-1 during oocyte

meiosis and provide a mechanistic understanding of PLK-1 targeting to meiotic

chromosomes. The next step will be to identify and characterise PLK-1 meiotic substrates, to

understand exactly how PLK-1 participates in each of its meiotic roles.

ACKNOWLEDGEMENTS

We thank Arshad Desai for sharing strains and antibodies and for helpful discussions and sharing results prior to publication; Pierre Gönczy for the PLK-1 antibody; Lionel Pintard for providing PLK-1 for the initial experiments and the plasmid for expression and purification; Bruce Bowerman for the GFP::ASPM-1 strain; Satpal Virdee for advice on fluorescence polarisation experiments. We also thank Ron Hay for comments on the manuscript. This work was supported by a Career Development Award from the Medical Research Council (grant MR/R008574/1) and an ISSF grant funded by the Wellcome Trust (105606/Z/14/Z). ST is funded by a Medical Research Council Doctoral Training Programme. DKC is supported by a Sir Henry Dale Fellowship from the Wellcome Trust (208833). J.H. received support from NSF Graduate Research Fellowship (grant #1650112) and from a grant from the NIH to A. Desai (GM074215). We acknowledge the FingerPrints Proteomics Facility and the Dundee Imaging Facility, which are supported by a 'Wellcome Trust Technology Platform' award (097945/B/11/Z) and the Tissue Imaging Facility, funded by a Wellcome Trust award (101468/Z/13/Z). Some nematode strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

AUTHOR CONTRIBUTIONS

S.J.P.T. designed, performed, and analysed experiments, and wrote the manuscript.

L.B-B. designed, performed, and analysed experiments.

F.S. designed, performed, and analysed experiments.

J.H. generated novel reagents (nematode strains).

D.K.C. generated novel reagents (nematode strains).

F.P. designed, performed, and analysed experiments; acquired funding; wrote the manuscript.

FIGURE LEGENDS

Figure 1. Analysis of PLK-1 localisation and inhibition during meiosis I.

A) Schematic of a *C. elegans* bivalent, highlighting the midbivalent and kinetochore.

B) Immunostaining of (untagged) PLK-1 in fixed oocytes. The insets represent a

magnified image of single bivalents for each channel. Scale bar, 2 µm.

C) *In situ* GFP-tagged PLK-1 (Martino et al., 2017) was followed through meiosis I in

live, dissected oocytes. More detailed sequence of events are displayed in Figure 1-

figure supplement 1. Yellow arrows point to chromosomes, blue arrows indicate

spindle poles, magenta arrows point towards the midbivalent, and green arrow

indicates the central spindle. Scale bars, 2 µm and 1 µm (in zoomed-in bivalents).

D) Schematic of the last two maturing oocytes and the fertilised egg, highlighting the

difference between PLK-1 depletion and acute PLK-1 inhibition.

E) *plk-1as* worms expressing GFP-tagged tubulin and mCherry-tagged histone were

dissected in medium containing ethanol ('EtOH', control) or the PP1 analogue 3IB-

PP1 (10 µM). Yellow arrows point to the each of the six bivalents. Scale bar, 2 µm.

F) *plk-1as* worms expressing GFP-tagged tubulin and mCherry-tagged histone, were

dissected in medium containing ethanol ('EtOH', control) or 0.1 µM PP1 analogue

3IB-PP1. Scale bar, 2 µm. The panels on the right show specific Z slices to highlight

individual chromosomes. Yellow arrows point to misaligned chromosomes contained

within the spindle, whereas blue arrows indicate chromosomes outside the spindle.

See also Figure 1-figure supplement 3.

G) Chromosme alignment defects scored for 1 µM 3IB-PP1-treated oocytes are presented

in the graph.

H) Worms expressing GFP-tagged ASPM-1 (pole marker) and mCherry-tagged histone

along with analogue-sensitive *plk-1*, were dissected in medium containing ethanol

('EtOH', control) or the PP1 analogue 3IB-PP1 at 1 µM or 10 µM. Scale bar, 2 µm.

Early metaphase I spindles are shown for each condition. Cyan arrows point to spindle

poles whereas yellow arrows indicate misaligned chromosomes.

Figure 2. BUB-1 contains a putative STP motif and regulates PLK-1 localisation *in vivo*.

A) Control (*'wild type'*) and BUB-1 depleted ['*bub-1(RNAi)*'] oocytes expressing sfGFP-

tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded throughout

meiosis I. Prometaphase/Metphase I is shown (before spindle rotation/shortening).

The cyan arrows point to the midbivalent and the yellow arrows point to the

chromosome arms. 'N' and 'n' denote the number of oocytes and bivalents analysed,

respectively. Left scale bar, 2 µm. Right scale bar, 1 µm.

B) Line profile analysis of PLK-1::sfGFP during Prometaphase I in *wild type* and *bub-*

1(RNAi) oocytes, as indicated by the yellow lines. Background signal was subtracted

and maximum signal for each channel was set to 1.

C) Schematic representation of the *C. elegans* BUB-1 protein (top) and sequence

alignment of the putative STP motif in nematode species (bottom).

D) (top) Schematic of BUB-1$^{190\text{-}628}$ (bottom) kinase assay of recombinant BUB-1$^{190\text{-}628}$

with Cdk1/Cyclin B, PLK-1, and both kinases combined as indicated. Phosphorylation

was assessed using SDS-PAGE followed by coomassie (total protein) or ProQ^{TM}

diamond (phosphoprotein) staining.

E) Western blot of recombinant BUB-1$^{190\text{-}628}$ Cdk1/Cyclin B kinase assays incubated

with either α-BUB-1 or α-phT527 BUB-1 antibodies. To ensure specifcity, α-phT527

BUB-1 antibody was incubated alone or together with an unphosphorylated or phT527

BUB-1 STP motif peptide, respectively.

F) (top) Schematic of BUB-1$^{190-628}$(T527A) (bottom) kinase assay of recombinant BUB-

1$^{190-628}$ and BUB-1$^{190-628}$(T527A) with Cdk1/Cyclin B and PLK-1 combined. Proteins

were incubated with the kinases for the indicated time points before phosphorylation

was assessed by SDS-PAGE and staining with either coomassie (total protein) or

ProQ^{TM} diamond (phosphoprotein).

Figure 3. BUB-1 directly binds to PLK1 *in vitro* via a Cdk1-phosphorylated STP motif.

A) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. BUB-1$^{190-628}$ was incubated with MBP-

PLK1^{PBD} at equimolar concentrations before separation by SEC. Binding was

conducted with unphosphorylated or Cdk1/Cyclin B phosphorylated BUB-1$^{190-628}$ as

indicated. The colours of the gel borders correspond to the colours of the respective

elution profile.

B) SEC-MALS analysis of BUB-1$^{190-628}$ and MBP-PLK1^{PBD} binding indicates a 1:1

complex stoichiometry. Relative absorbance (dotted lines) and molecular weight

(solid lines) are colour-coded to match the corresponding proteins.

C) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. Wild type or T527A mutant BUB-1$^{190-628}$

was phosphorylated by Cdk1/Cyclin B before incubation with an equimolar

concentration of MBP-PLK1^{PBD}, binding was assessed by SEC. The colours of the gel

borders correspond to the colours of the respective elution profile.

D) FITC-labelled peptides containing the BUB-1 STP motif were incubated with

increasing concentrations of MBP-PLK1^{PBD} or MBP-PLK-1^{PBD}(H538A/K540M) and

Taylor, Bel Borja et al.

binding analysed by fluoresence polarisation. 'T^Ph' denotes phosphorylated Threonine.

The colours of the data points correspond to the colours of the experimental conditions

as indicated.

Figure 4. The BUB-1 STP motif is required for PLK-1 targeting and chromosome

alignment.

A) Fixed oocytes were stained with a PLK-1 specific antibody (green) and *bub-1^T527A*

heterozygote and homozygote oocytes were compared. DNA is shown in magenta.

The yellow arrow points to the midbivalent magnified on the right in each case and the

blue arrowhead points to the midbivalent. Scale bars, 2 µm (left) and 1 µm (right).

B) Line profile analysis of PLK-1 localisation in fixed oocytes during early Metaphase I

in *bub-1^T527A* heterozygote ('*bub-1^T527A/+*') and homozygote ('*bub-1^T527A*') oocytes, as

indicated by the yellow lines. Background signal was subtracted and maximum signal

for each channel was set to 1.

C) Fixed oocytes were stained with a BUB-1 specific antibody (green) and *bub-1^T527A*

heterozygote and homozygote oocytes were compared. DNA is shown in magenta.

Scale bar, 1 µm.

D) *bub-1^T527A* heterozygote ('*bub-1^T527A/+*') and homozygote ('*bub-1^T527A*') oocytes

expressing GFP-tagged tubulin and mCherry-tagged histone were filmed during

meiosis I. Two homozygote ('*bub-1^T527A*') oocytes are shown to depict the difference

in severity of the alignment defect. Scale bar, 2 µm.

E) Meiotic defects (as described in the Methods section) were assessed in wild type, *bub-*

1^T527A heterozygote ('*bub-1^T527A/+*') and homozygote oocytes. Representative images

of the different phenotypes analysed are presented on top (scale bars, 2 µm). 'N'

represents the number of experiments and 'n' denotes the number of oocytes analysed.

Figure 5. CENP-C^{HCP-4} contains a putative STP motif and is required for chromosomal

PLK-1 targeting.

A) *In situ* GFP-tagged CENP-C^{HCP-4} was followed through meiosis I in live, dissected

oocytes. Scale bar, 2 µm.

B) Schematic representation of the *C. elegans* CENP-C^{HCP-4} (top) and sequence

alignment of the putative STP motif in nematode species (bottom).

C) Control (*'wild type'*) and CENP-C^{HCP-4}-depleted [*'hcp-4(RNAi)'*] oocytes expressing

sfGFP-tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded

throughout meiosis I. Prometaphase I is shown (before spindle rotation/shortening).

On the right, the cyan arrows point to the midbivalent and the yellow arrow points to

the kinetochore. 'N' and 'n' denote the number of oocytes and bivalents analysed,

respectively. Left scale bar, 2 µm. Right scale bar, 1 µm.

D) Line profile analysis of PLK-1::sfGFP during early Metaphase I in *wild type* and *hcp-*

4(RNAi) oocytes, as indicated by the yellow lines. Background signal was subtracted

and maximum signal for each channel was set to 1. The blue arrows point to the

midbivalent and the yellow arrow points to the kinetochore

E) Schematic of recombinant CENP-C$^{HCP-4(1-214)}$ proteins.

F) Kinase assay of recombinant CENP-C$^{HCP-4(1-214)}$ wild type and T163A proteins with

Cdk1:Cyclin B. Reactions were analysed by SDS-PAGE followed by ProQ^{TM}

diamond (phosphoprotein) or coomassie (total protein) staining.

G) Western blot of recombinant CENP-C$^{HCP-4(1-214)}$ Cdk1/Cyclin B kinase assays

incubated with either α-HCP-4 or α-phT163 HCP-4 antibodies. To ensure specifcity,

α-phT163 HCP-4 antibody was incubated alone or together with an unphosphorylated

or phT163 HCP-4 STP motif peptide, respectively.

Figure 6. CENP-C^{HCP-4} interacts with PLK-1 *in vitro* in a Cdk1-dependent manner.

A) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. CENCP-C$^{HCP-4(1-214)}$ was incubated with

MBP-PLK1^{PBD} at equimolar concentrations before being analysed by SEC. Binding

was conducted with unphosphorylated or Cdk1:Cyclin B phosphorylated CENP-C^{HCP-}

$^{4(1-214)}$ as indicated. The colours of the gel borders correspond to the colours of the

respective elution profile.

B) SEC-MALS analysis of CENP-C$^{HCP-4(1-214)}$ and MBP-PLK1^{PBD} binding indicates a 1:1

complex stoichiometry. Relative absorbance (dotted lines) and molecular weight

(solid lines) are colour-coded to match the corresponding proteins.

C) Elution profile and coomassie-stained SDS-PAGE gels of representative fractions

from the Superdex 200 10/300 SEC column. Wild type or T163A mutant CENP-C^{HCP-}

$^{4(1-214)}$ was phosphorylated by Cdk1/Cyclin B before incubation with an equimolar

concentration of MBP-PLK1^{PBD}. Binding was then assessed by SEC. The colours of

the gel borders correspond to the colours of the respective elution profile.

D) FITC-labelled peptides containing the HCP-4 STP motif were incubated with

increasing concentrations of MBP-PLK1^{PBD} or MBP-PLK-1^{PBD}(H538A/K540M) and

binding analysed by fluoresence polarisation. Unphosphorylated versus T163-

phosphorylated (T^{ph}) peptides were compared. The colours of the data points

correspond to the colours of the experimental conditions as indicated.

Figure 7. CENP-C^{HCP-4} directly recruits PLK-1 to chromosome arms.

A) Control (*'wild type'*) and CENP-C^{HCP-4} STP mutant ['*hcp-4^{T163A}*'] oocytes expressing

sfGFP-tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded

throughout meiosis I. Metphase I is shown. Left scale bar, 2 µm. Right scale bar, 1

µm.

B) Line profile analysis of PLK-1::sfGFP during early Metaphase I in *wild type* and *hcp-*

4^{T163A} oocytes, as indicated by the yellow lines. Background signal was subtracted and

maximum signal for each channel was set to 1.

C) Fixed oocytes were stained with an HCP-4 specific antibody (green in the merged

image). *hcp-4^{T163A}* mutant oocytes were compared to wild type. DNA is shown in

magenta in the merged panel. Scale bar on the left, 2 µm; scale bar on the right, 1 µm.

D) Same as in C) but using a BUB-1 specific antibody to compare BUB-1 localisation in

wild type and *hcp-4^{T163A}* mutant oocytes. Scale bar, 1 µm.

Figure 8. Combined disruption of BUB-1- and CENP-C^{HCP-4}-dependent PLK-1

recruitment leads to severe meiotic defects.

A) Control (*'wild type'*), BUB-1-depleted ['*bub-1(RNAi)*'], and CENP-C^{HCP-4}-depleted

['*hcp-4(RNAi)*'] oocytes expressing sfGFP-tagged PLK-1 (and mCherry-tagged

histone) were dissected and recorded throughout meiosis I. Prometaphase/Metphase I

is shown (before spindle rotation/shortening). 'N' and 'n' denote the number of

oocytes and bivalents analysed, respectively. Left scale bar, 2 µm. Right scale bar, 1

µm.

B) Control ('wild type'), BUB-1-depleted ['bub-1(RNAi)'], and CENP-C^{HCP-4}-depleted

['hcp-4(RNAi)'] oocytes expressing GFP-tagged tubulin (and mCherry-tagged histone)

were dissected and recorded throughout meiosis I. Scale bar, 2 µm.

C) Meiotic defects (as described in the Methods section) were assessed in wild type, bub-

1(RNAi), and hcp-4(RNAi) oocytes. 'N' and 'n' denote the number of oocytes and

bivalents analysed, respectively.

D) Fixed wild type, hcp-4(RNAi), bub-1^{T527A}, and bub-1^{T527A}+hcp-4(RNAi) oocytes were

stained with a PLK-1 specific antibody. Scale bar, 1 µm.

E) Control ('wild type'), BUB-1^{T527A} ['bub-1^{T527A}'], and CENP-C^{HCP-4}-depleted ['hcp-

4(RNAi)'] oocytes expressing GFP-tagged tubulin (and mCherry-tagged histone) were

dissected and recorded throughout meiosis I. Scale bar, 2 µm.

F) Meiotic defects (as described in the Methods section) were assessed in wild type, bub-

1^{T527A}, hcp-4(RNAi), and bub-1^{T527A}+hcp-4(RNAi) oocytes. 'N' and 'n' denote the

number of oocytes and bivalents analysed, respectively.

Figure 9. Disruption of BUB-1- and CENP-C^{HCP-4} STP motifs leads to severe meiotic

defects.

A) Wild type BUB-1 ('BUB-1^{wt}') and T527A BUB-1 ('BUB-1^{T527A}') were expressed

from transgenes in the absence of endogenous BUB-1, in wild type ('hcp-4^{wt}') or

HCP-4 T163A ('hcp-4^{T163A}') worms. Oocytes expressing sfGFP-tagged PLK-1 (and

mCherry-tagged histone) were dissected and recorded throughout meiosis I. Images

from oocytes in Prometaphase I are shown. Cyan arrow points to the midbivalent and

yellow arrows point to chromosome arms. Scale bar, 2 µm.

B) Oocytes from the same conditions as in A but expressing GFP-tagged tubulin (and

mCherry-tagged histone) were dissected and recorded throughout meiosis I. Scale bar,

2 µm. In the case of the double mutant *hcp-4^{T163A}/bub-1^{T527A}*, two oocytes are shown,

one that achieved (defective) PBE ('oocyte #1') and another in which PBE failed

('oocyte #2').

C) Meiotic defects were assessed and are shown as % of oocytes showing each defect.

See 'Methods' for more details. 'N' and 'n' denote the number of oocytes and

bivalents analysed, respectively.

Figure 1-figure supplement 1. PLK-1 localisation during oocyte meiosis

In situ GFP-tagged PLK-1 (Martino et al., 2017) was followed through meiosis I in

live, dissected oocytes. Scale bar, 2 µm. This panel shows a more detailed sequence of

events of the same movie presented in Figure 1B.

Figure 1-figure supplement 2. Embryonic viability assays after PLK-1as inhibition

Three different PP1 analogues were tested: 3IB-PP1, 3MB-PP1, and 1NA-PP1.

Inhibitors were included in the plates at 10, 20, and 50 µM and worms were then

allowed to lay eggs. Viable progeny was assessed 24 and 48 hs later.

Figure 1-figure supplement 3. Chromosome alignment defects upon acute PLK-1as

inhibition

A) *Wild type* or *plk-1^{as}* worms expressing GFP-tagged tubulin and mCherry-tagged

histone, were dissected in medium containing 1 µM 3IB-PP1 ('low') or 10 µM

('high'). Maximum intensity projections are shown and the arrows have different

colours to identify those chromosomes in panel B). Scale bar, 2 µm.

B) Detailed analysis of individual chromosome behaviour using selected Z slices of

+3IB-PP1 'low' from panel A). Highly unstable behaviour of specific chromosomes

(switching back and forth between aligned and misaligned) can be followed by the

different coloured arrows. Panels outside of the green rectangle are timepoints not

shown in A). Time between frames is 20 seconds. Scale bar, 2 µm.

Figure 2-figure supplement 1. BUB-1 phosphorylation by Cdk1 and PLK-1 in vitro.

A) Kinase assay of recominant BUB-1$^{190-628}$ and BUB-1$^{190-628}$(T527A) with Cdk1/Cyclin

B. Proteins were incubated with Cdk1:Cyclin B for the indicated time points before

phosphorylation was assessed by SDS-PAGE and staining with either coomassie (total

protein) or ProQ^{TM} diamond (phosphoprotein).

B) Kinase assay of recominant BUB-1$^{190-628}$ and BUB-1$^{190-628}$(T527A) with PLK-1.

Proteins were incubated with PLK-1 for the indicated time points before

phosphorylation was assessed by SDS-PAGE and staining with either coomassie (total

protein) or ProQ^{TM} diamond (phosphoprotein).

Figure 5-figure supplement 1. PLK-1 co-localises with the kinetochore component KNL-

3 in non-chromosomal kinetochore assemblies

A) Oocytes co-expressing sfGFP-tagged PLK-1 and mScarlet-tagged KNL-3 were

analysed. The panels show different Z-slices and the panels on the right are higher

magnification images of the delineated areas to better visualise the spindle-associated

PLK-1 kinetochore localisation. Scale bars, 2 µm (left) and 1 µm (right).

B) Same as in A, but with detail on the cortical kinetochore assemblies. Scale bars, 10

µm (left) and 2 µm (right).

Figure 8-figure supplement 1. PLK-1 intensity measurements.

Chromosome-associated PLK-1 intensity was measured (relative to background). The

graph shows the individual values (circles) along with the median of the population

(horizontal black lines).

Figure 8-figure supplement 2. Combined disruption of BUB-1- and CENP-C^{HCP-4}-

dependent PLK-1 recruitment leads to severe meiotic defects.

A) Control (*'wild type'*), BUB-1-depleted ['*bub-1(RNAi)*'], CENP-C^{HCP-4}(T163A) ['*hcp-*

4^{T163A} '], and *bub-1(RNAi)+hcp-4^{T163A}* oocytes expressing sfGFP-tagged PLK-1 (and

mCherry-tagged histone) were dissected and recorded throughout meiosis I.

Prometaphase/Metphase I is shown (before spindle rotation/shortening). Panels on the

left show the full spindle, while the right-most two columns display single bivalents.

Left scale bar, 2 µm. Right scale bar, 1 µm. 'N' and 'n' denote the number of oocytes

and bivalents analysed, respectively.

B) Control (*'wild type'*), BUB-1-depleted ['*bub-1(RNAi)*'], CENP-C^{HCP-4}(T163A) ['*hcp-*

4^{T163A} '], and *bub-1(RNAi)+hcp-4^{T163A}* oocytes expressing GFP-tagged tubulin (and

mCherry-tagged histone) were dissected and recorded throughout meiosis I. Scale bar,

2 µm.

C) Meiotic defects (as described in the Methods section) were assessed in wild type, *bub-*

1(RNAi), *hcp-4^{T163A}, and bub-1(RNAi)+hcp-4^{T163A}* oocytes. 'N' and 'n' denote the

number of oocytes and bivalents analysed, respectively.

Figure 9-figure supplement 1. Strategy for the RNAi/rescue experiments using *bub-1*

transgenes.

A) Schematics depicting the strategy for depletion of endogenous BUB-1 and expression

of RNAi-resistant transgenes.

B) Western blot showing the levels of BUB-1 in the different conditions. All lanes come

from the same blot and the vertical white line indicates that the lanes were not

contiguous in the original blot (available in Figure 9-figure supplement 1-source data).

Figure 9-figure supplement 2. Disruption of BUB-1- and CENP-C^{HCP-4} STP motifs leads

to severe meiotic defects.

A) This panel shows the full set of conditions analysed for the experiment shown in

Figure 9A, fully validating the depletion/rescue strategy. Oocytes expressing sfGFP-

tagged PLK-1 (and mCherry-tagged histone) were dissected and recorded throughout

meiosis I. Images from oocytes in Prometaphase I are shown. Cyan arrows points to

the midbivalent and yellow arrows point to chromosome arms. Scale bar, 2 µm.

B) This panel shows the full set of conditions analysed for the experiment shown in

Figure 9B, fully validating the depletion/rescue strategy. Oocytes from the same

conditions as in A but expressing GFP-tagged tubulin (and mCherry-tagged histone)

were dissected and recorded throughout meiosis I. Scale bar, 2 µm. In the case of the

double mutant $hcp\text{-}4^{T163A}/bub\text{-}1^{T527A}$, two oocytes are shown, one that achieved

(defective) PBE ('oocyte #1') and another in which PBE failed ('oocyte #2').

REFERENCES

Archambault V, Glover DM. 2009. Polo-like kinases: conservation and divergence in their functions and regulation. *Nat Rev Mol Cell Biol* **10**:265–275. doi:10.1038/nrm2653

Arribere JA, Bell RT, Fu BXH, Artiles KL, Hartman PS, Fire AZ. 2014. Efficient Marker-Free Recovery of Custom Genetic Modifications with CRISPR/Cas9 in Caenorhabditis elegans. *Genetics* **198**:837–846. doi:10.1534/genetics.114.169730

Bel Borja L, Soubigou F, Taylor SJP, Fraguas Bringas C, Budrewicz J, Lara-Gonzalez P, Sorensen Turpin CG, Bembenek JN, Cheerambathur DK, Pelisch F. 2020. BUB-1 targets PP2A:B56 to regulate chromosome congression during meiosis I in C. elegans oocytes. *eLife* **9**:e65307. doi:10.7554/eLife.65307

Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J, Shimizu E, Tsien JZ, Schultz PG, Rose MD, Wood JL, Morgan DO, Shokat KM. 2000. A chemical switch for inhibitor-sensitive alleles of any protein kinase. *Nature* **407**:395–401. doi:10.1038/35030148

Budirahardja Y, Gönczy P. 2008. PLK-1 asymmetry contributes to asynchronous cell division of *C. elegans* embryos. *Development* **135**:1303–1313. doi:10.1242/dev.019075

Cabral G, Laos T, Dumont J, Dammermann A. 2019. Differential Requirements for Centrioles in Mitotic Centrosome Growth and Maintenance. *Developmental Cell* **50**:355-366.e6. doi:10.1016/j.devcel.2019.06.004

Chase D, Serafinas C, Ashcroft N, Kosinski M, Longo D, Ferris DK, Golden A. 2000. The polo-like kinase PLK-1 is required for nuclear envelope breakdown and the completion of meiosis inCaenorhabditis elegans. *genesis* **26**:26–41. doi:10.1002/(SICI)1526-968X(200001)26:1<26::AID-GENE6>3.0.CO;2-O

Cheerambathur DK, Prevo B, Chow T-L, Hattersley N, Wang S, Zhao Z, Kim T, Gerson-Gurwitz A, Oegema K, Green R, Desai A. 2019. The Kinetochore-Microtubule Coupling Machinery Is Repurposed in Sensory Nervous System Morphogenesis. *Developmental Cell* **48**:864-872.e7. doi:10.1016/j.devcel.2019.02.002

Cheng K-Y. 2003. The crystal structure of the human polo-like kinase-1 polo box domain and its phospho-peptide complex. *The EMBO Journal* **22**:5757–5768. doi:10.1093/emboj/cdg558

Connolly AA, Sugioka K, Chuang C-H, Lowry JB, Bowerman B. 2015. KLP-7 acts through the Ndc80 complex to limit pole number in C. elegans oocyte meiotic spindle assembly. *Journal of Cell Biology* **210**:917–932. doi:10.1083/jcb.201412010

Conway W, Kiewisz R, Fabig G, Kelleher CP, Wu H-Y, Anjur-Dietrich M, Müller-Reichert T, Needleman DJ. 2022. Self-organization of kinetochore-fibers in human mitotic spindles. *eLife* **11**:e75458. doi:10.7554/eLife.75458

Cordeiro MH, Smith RJ, Saurin AT. 2020. Kinetochore phosphatases suppress autonomous Polo-like kinase 1 activity to control the mitotic checkpoint. *Journal of Cell Biology* **219**:e202002020. doi:10.1083/jcb.202002020

Danlasky BM, Panzica MT, McNally KP, Vargas E, Bailey C, Li W, Gong T, Fishman ES, Jiang X, McNally FJ. 2020. Evidence for anaphase pulling forces during C. elegans meiosis. *Journal of Cell Biology* **219**:e202005179. doi:10.1083/jcb.202005179

Decker M, Jaensch S, Pozniakovsky A, Zinke A, O'Connell KF, Zachariae W, Myers E, Hyman AA. 2011. Limiting Amounts of Centrosome Material Set Centrosome Size

in C. elegans Embryos. *Current Biology* **21**:1259–1267.
doi:10.1016/j.cub.2011.06.002
Desai A, Rybina S, Müller-Reichert T, Shevchenko Andrej, Shevchenko Anna, Hyman A,
Oegema K. 2003. KNL-1 directs assembly of the microtubule-binding interface of
the kinetochore in *C. elegans*. *Genes Dev* **17**:2421–2435.
doi:10.1101/gad.1126303
Dumont J, Oegema K, Desai A. 2010. A kinetochore-independent mechanism drives
anaphase chromosome separation during acentrosomal meiosis. *Nat Cell Biol*
12:894–901. doi:10.1038/ncb2093
Elia AEH, Cantley LC, Yaffe MB. 2003a. Proteomic Screen Finds pSer/pThr-Binding
Domain Localizing Plk1 to Mitotic Substrates. *Science* **299**:1228–1231.
doi:10.1126/science.1079079
Elia AEH, Rellos P, Haire LF, Chao JW, Ivins FJ, Hoepker K, Mohammad D, Cantley LC,
Smerdon SJ, Yaffe MB. 2003b. The Molecular Basis for Phosphodependent
Substrate Targeting and Regulation of Plks by the Polo-Box Domain. *Cell* **115**:83–
95. doi:10.1016/S0092-8674(03)00725-6
Elowe S, Hümmer S, Uldschmid A, Li X, Nigg EA. 2007. Tension-sensitive Plk1
phosphorylation on BubR1 regulates the stability of kinetochore–microtubule
interactions. *Genes Dev* **21**:2205–2219. doi:10.1101/gad.436007
Foltz DR, Jansen LET, Black BE, Bailey AO, Yates JR, Cleveland DW. 2006. The human
CENP-A centromeric nucleosome-associated complex. *Nat Cell Biol* **8**:458–469.
doi:10.1038/ncb1397
Gelens L, Qian J, Bollen M, Saurin AT. 2018. The Importance of Kinase–Phosphatase
Integration: Lessons from Mitosis. *Trends in Cell Biology* **28**:6–21.
doi:10.1016/j.tcb.2017.09.005
Gigant E, Stefanutti M, Laband K, Gluszek-Kustusz A, Edwards F, Lacroix B, Maton G,
Canman JC, Welburn JPI, Dumont J. 2017. Inhibition of ectopic microtubule
assembly by the kinesin-13 KLP-7 prevents chromosome segregation and
cytokinesis defects in oocytes. *Journal of Cell Science* **130**:e1.1-e1.1.
doi:10.1242/jcs.205914
Gómez-Cavazos JS, Lee K-Y, Lara-González P, Li Y, Desai A, Shiau AK, Oegema K. 2020. A
Non-canonical BRCT-Phosphopeptide Recognition Mechanism Underlies RhoA
Activation in Cytokinesis. *Current Biology* **30**:3101-3115.e11.
doi:10.1016/j.cub.2020.05.090
Green RA, Mayers JR, Wang S, Lewellyn L, Desai A, Audhya A, Oegema K. 2013. The
midbody ring scaffolds the abscission machinery in the absence of midbody
microtubules. *Journal of Cell Biology* **203**:505–520. doi:10.1083/jcb.201306036
Han X, Adames K, Sykes EME, Srayko M. 2015. The KLP-7 Residue S546 Is a Putative
Aurora Kinase Site Required for Microtubule Regulation at the Centrosome in C.
elegans. *PLoS ONE* **10**:e0132593. doi:10.1371/journal.pone.0132593
Hattersley N, Schlientz AJ, Prevo B, Oegema K, Desai A. 2022. MEL-28/ELYS and CENP-C
coordinately control outer kinetochore assembly and meiotic chromosome-
microtubule interactions. *Current Biology* **32**:2563-2571.e4.
doi:10.1016/j.cub.2022.04.046
Heald R, Tournebize R, Blank T, Sandaltzopoulos R, Becker P, Hyman A, Karsenti E. 1996.
Self-organization of microtubules into bipolar spindles around artificial
chromosomes in Xenopus egg extracts. *Nature* **382**:420–425.
doi:10.1038/382420a0

Heun P, Erhardt S, Blower MD, Weiss S, Skora AD, Karpen GH. 2006. Mislocalization of
the Drosophila Centromere-Specific Histone CID Promotes Formation of
Functional Ectopic Kinetochores. *Developmental Cell* **10**:303–315.
doi:10.1016/j.devcel.2006.01.014
Izuta H, Ikeno M, Suzuki N, Tomonaga T, Nozaki N, Obuse C, Kisu Y, Goshima N, Nomura
F, Nomura N, Yoda K. 2006. Comprehensive analysis of the ICEN (Interphase
Centromere Complex) components enriched in the CENP-A chromatin of human
cells. *Genes to Cells* **11**:673–684. doi:10.1111/j.1365-2443.2006.00969.x
Jang C-Y, Coppinger JA, Seki A, Yates JR III, Fang G. 2009. Plk1 and Aurora A regulate the
depolymerase activity and the cellular localization of Kif2a. *Journal of Cell Science*
122:1334–1341. doi:10.1242/jcs.044321
Kang YH, Park CH, Kim T-S, Soung N-K, Bang JK, Kim BY, Park J-E, Lee KS. 2011.
Mammalian Polo-like Kinase 1-dependent Regulation of the PBIP1-CENP-Q
Complex at Kinetochores. *Journal of Biological Chemistry* **286**:19744–19757.
doi:10.1074/jbc.M111.224105
Kang YH, Park J-E, Yu L-R, Soung N-K, Yun S-M, Bang JK, Seong Y-S, Yu H, Garfield S,
Veenstra TD, Lee KS. 2006. Self-Regulated Plk1 Recruitment to Kinetochores by
the Plk1-PBIP1 Interaction Is Critical for Proper Chromosome Segregation.
Molecular Cell **24**:409–422. doi:10.1016/j.molcel.2006.10.016
Kiewisz R, Fabig G, Conway W, Baum D, Needleman D, Müller-Reichert T. 2022. Three-
dimensional structure of kinetochore-fibers in human mitotic spindles. *eLife*
11:e75459. doi:10.7554/eLife.75459
Kitada K, Johnson AL, Johnston LH, Sugino A. 1993. A multicopy suppressor gene of the
Saccharomyces cerevisiae G1 cell cycle mutant gene dbf4 encodes a protein
kinase and is identified as CDC5. *Molecular and Cellular Biology* **13**:4445–4457.
doi:10.1128/mcb.13.7.4445-4457.1993
Laband K, Lacroix B, Edwards F, Canman JC, Dumont J. 2018. Chapter 11 - Live imaging
of C. elegans oocytes and early embryos In: Maiato H, Schuh M, editors. Methods
in Cell Biology, Mitosis and Meiosis Part B. Academic Press. pp. 217–236.
doi:10.1016/bs.mcb.2018.03.025
Linder MI, Köhler M, Boersema P, Weberruss M, Wandke C, Marino J, Ashiono C, Picotti
P, Antonin W, Kutay U. 2017. Mitotic Disassembly of Nuclear Pore Complexes
Involves CDK1- and PLK1-Mediated Phosphorylation of Key Interconnecting
Nucleoporins. *Developmental Cell* **43**:141-156.e7.
doi:10.1016/j.devcel.2017.08.020
Llamazares S, Moreira A, Tavares A, Girdham C, Spruce BA, Gonzalez C, Karess RE,
Glover DM, Sunkel CE. 1991. polo encodes a protein kinase homolog required for
mitosis in Drosophila. *Genes Dev* **5**:2153–2165. doi:10.1101/gad.5.12a.2153
Maddox PS, Corbett KD, Desai A. 2012. Structure, assembly and reading of centromeric
chromatin. *Current Opinion in Genetics & Development*, Genome architecture and
expression **22**:139–147. doi:10.1016/j.gde.2011.11.005
Marston AL, Amon A. 2004. Meiosis: cell-cycle controls shuffle and deal. *Nat Rev Mol Cell*
Biol **5**:983–997. doi:10.1038/nrm1526
Marston AL, Wassmann K. 2017. Multiple Duties for Spindle Assembly Checkpoint
Kinases in Meiosis. *Frontiers in Cell and Developmental Biology* **5**.
Martino L, Morchoisne-Bolhy S, Cheerambathur DK, Van Hove L, Dumont J, Joly N, Desai
A, Doye V, Pintard L. 2017. Channel Nucleoporins Recruit PLK-1 to Nuclear Pore
Complexes to Direct Nuclear Envelope Breakdown in C. elegans. *Developmental*
Cell **43**:157-171.e7. doi:10.1016/j.devcel.2017.09.019

Monen J, Maddox PS, Hyndman F, Oegema K, Desai A. 2005. Differential role of CENP-A
in the segregation of holocentric C. elegans chromosomes during meiosis and
mitosis. *Nat Cell Biol* **7**:1248–1255. doi:10.1038/ncb1331

Mundt KE, Golsteyn RM, Lane HA, Nigg EA. 1997. On the Regulation and Function of
Human Polo-like Kinase 1 (PLK1): Effects of Overexpression on Cell Cycle
Progression. *Biochemical and Biophysical Research Communications* **239**:377–
385. doi:10.1006/bbrc.1997.7378

Musacchio A, Desai A. 2017. A Molecular View of Kinetochore Assembly and Function.
Biology **6**. doi:10.3390/biology6010005

Neef R, Preisinger C, Sutcliffe J, Kopajtich R, Nigg EA, Mayer TU, Barr FA. 2003.
Phosphorylation of mitotic kinesin-like protein 2 by polo-like kinase 1 is required
for cytokinesis. *Journal of Cell Biology* **162**:863–876. doi:10.1083/jcb.200306009

Novak B, Kapuy O, Domingo-Sananes MR, Tyson JJ. 2010. Regulated protein kinases and
phosphatases in cell cycle decisions. *Current Opinion in Cell Biology*, Cell
differentiation / Cell division, growth and death **22**:801–808.
doi:10.1016/j.ceb.2010.07.001

Oegema K, Desai A, Rybina S, Kirkham M, Hyman AA. 2001. Functional Analysis of
Kinetochore Assembly in Caenorhabditis elegans. *Journal of Cell Biology*
153:1209–1226. doi:10.1083/jcb.153.6.1209

Ohkura H. 2015. Meiosis: An Overview of Key Differences from Mitosis. *Cold Spring Harb*
Perspect Biol **7**:a015859. doi:10.1101/cshperspect.a015859

Ohkura H, Hagan IM, Glover DM. 1995. The conserved Schizosaccharomyces pombe
kinase plo1, required to form a bipolar spindle, the actin ring, and septum, can
drive septum formation in G1 and G2 cells. *Genes Dev* **9**:1059–1073.
doi:10.1101/gad.9.9.1059

Ohta M, Zhao Z, Wu D, Wang S, Harrison JL, Gómez-Cavazos JS, Desai A, Oegema KF.
2021. Polo-like kinase 1 independently controls microtubule-nucleating capacity
and size of the centrosome. *Journal of Cell Biology* **220**:e202009083.
doi:10.1083/jcb.202009083

Okada M, Cheeseman IM, Hori T, Okawa K, McLeod IX, Yates JR, Desai A, Fukagawa T.
2006. The CENP-H–I complex is required for the efficient incorporation of newly
synthesized CENP-A into centromeres. *Nat Cell Biol* **8**:446–457.
doi:10.1038/ncb1396

Pahlavan G, Polanski Z, Kalab P, Golsteyn R, Nigg EA, Maro B. 2000. Characterization of
Polo-like Kinase 1 during Meiotic Maturation of the Mouse Oocyte. *Developmental*
Biology **220**:392–400. doi:10.1006/dbio.2000.9656

Paix A, Folkmann A, Rasoloson D, Seydoux G. 2015. High Efficiency, Homology-Directed
Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein
Complexes. *Genetics* **201**:47–54. doi:10.1534/genetics.115.179382

Paix A, Folkmann A, Seydoux G. 2017. Precision genome editing using CRISPR-Cas9 and
linear repair templates in C. elegans. *Methods*, CRISPR-Cas systems for Genome
engineering and investigation **121–122**:86–93.
doi:10.1016/j.ymeth.2017.03.023

Pelisch F, Bel Borja L, Jaffray EG, Hay RT. 2019. Sumoylation regulates protein dynamics
during meiotic chromosome segregation in *C. elegans* oocytes. *Journal of Cell*
Science jcs.232330. doi:10.1242/jcs.232330

Pelisch F, Tammsalu T, Wang B, Jaffray EG, Gartner A, Hay RT. 2017. A SUMO-Dependent
Protein Network Regulates Chromosome Congression during Oocyte Meiosis.
Molecular Cell **65**:66–77. doi:10.1016/j.molcel.2016.11.001

Petronczki M, Lénárt P, Peters J-M. 2008. Polo on the Rise—from Mitotic Entry to
Cytokinesis with Plk1. *Developmental Cell* **14**:646–659.
doi:10.1016/j.devcel.2008.04.014
Qi W, Tang Z, Yu H. 2006a. Phosphorylation- and Polo-Box–dependent Binding of Plk1 to
Bub1 Is Required for the Kinetochore Localization of Plk1. *MBoC* **17**:3705–3716.
doi:10.1091/mbc.e06-03-0240
Qi W, Tang Z, Yu H. 2006b. Phosphorylation- and Polo-Box–dependent Binding of Plk1 to
Bub1 Is Required for the Kinetochore Localization of Plk1. *MBoC* **17**:3705–3716.
doi:10.1091/mbc.e06-03-0240
Rahman MM, Munzig M, Kaneshiro K, Lee B, Strome S, Müller-Reichert T, Cohen-Fix O.
2015. Caenorhabditis elegans polo-like kinase PLK-1 is required for merging
parental genomes into a single nucleus. *Mol Biol Cell* **26**:4718–4735.
doi:10.1091/mbc.E15-04-0244
Ritter A, Sanhaji M, Steinhäuser K, Roth S, Louwen F, Yuan J. 2014. The activity
regulation of the mitotic centromere-associated kinesin by Polo-like kinase 1.
Oncotarget **6**:6641–6655. doi:10.18632/oncotarget.2843
Sanhaji M, Ritter A, Belsham HR, Friel CT, Roth S, Louwen F, Yuan J. 2014. Polo-like
kinase 1 regulates the stability of the mitotic centromere-associated kinesin in
mitosis. *Oncotarget* **5**:3130–3144. doi:10.18632/oncotarget.1861
Saurin AT. 2018. Kinase and Phosphatase Cross-Talk at the Kinetochore. *Front Cell Dev*
Biol **6**:62. doi:10.3389/fcell.2018.00062
Schmucker S, Sumara I. 2014. Molecular dynamics of PLK1 during mitosis. *Molecular &*
Cellular Oncology **1**:e954507. doi:10.1080/23723548.2014.954507
Shao H, Huang Y, Zhang L, Yuan K, Chu Y, Dou Z, Jin C, Garcia-Barrio M, Liu X, Yao X.
2015. Spatiotemporal dynamics of Aurora B-PLK1-MCAK signaling axis
orchestrates kinetochore bi-orientation and faithful chromosome segregation. *Sci*
Rep **5**:12204. doi:10.1038/srep12204
Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert
M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-
quality protein multiple sequence alignments using Clustal Omega. *Molecular*
Systems Biology **7**:539. doi:10.1038/msb.2011.75
Singh P, Pesenti ME, Maffini S, Carmignani S, Hedtfeld M, Petrovic A, Srinivasamani A,
Bange T, Musacchio A. 2021. BUB1 and CENP-U, Primed by CDK1, Are the Main
PLK1 Kinetochore Receptors in Mitosis. *Molecular Cell* **81**:67-87.e9.
doi:10.1016/j.molcel.2020.10.040
Solc P, Kitajima TS, Yoshida S, Brzakova A, Kaido M, Baran V, Mayer A, Samalova P,
Motlik J, Ellenberg J. 2015. Multiple requirements of PLK1 during mouse oocyte
maturation. *PLoS One* **10**:e0116783. doi:10.1371/journal.pone.0116783
Sunkel CE, Glover DM. 1988. polo, a mitotic mutant of Drosophila displaying abnormal
spindle poles. *Journal of Cell Science* **89**:25–38. doi:10.1242/jcs.89.1.25
Tavernier N, Noatynska A, Panbianco C, Martino L, Van Hove L, Schwager F, Léger T,
Gotta M, Pintard L. 2015. Cdk1 phosphorylates SPAT-1/Bora to trigger PLK-1
activation and drive mitotic entry in C. elegans embryos. *Journal of Cell Biology*
208:661–669. doi:10.1083/jcb.201408064
Timmons L, Court DL, Fire A. 2001. Ingestion of bacterially expressed dsRNAs can
produce specific and potent genetic interference in Caenorhabditis elegans. *Gene*
263:103–112. doi:10.1016/S0378-1119(00)00579-5
Tong C, Fan H-Y, Lian L, Li S-W, Chen D-Y, Schatten H, Sun Q-Y. 2002. Polo-Like Kinase-1
Is a Pivotal Regulator of Microtubule Assembly During Mouse Oocyte Meiotic

Maturation, Fertilization, and Early Embryonic Mitosis1. *Biology of Reproduction*
67:546–554. doi:10.1095/biolreprod67.2.546
Velez-Aguilera G, Nkombo Nkoula S, Ossareh-Nazari B, Link J, Paouneskou D, Van Hove
L, Joly N, Tavernier N, Verbavatz J-M, Jantsch V, Pintard L. 2020. PLK-1 promotes
the merger of the parental genome into a single nucleus by triggering lamina
disassembly. *eLife* **9**:e59510. doi:10.7554/eLife.59510
Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2—
a multiple sequence alignment editor and analysis workbench. *Bioinformatics*
25:1189–1191. doi:10.1093/bioinformatics/btp033
Wianny F, Tavares Á, Evans MJ, Glover DM, Zernicka-Goetz M. 1998. Mouse polo-like
kinase 1 associates with the acentriolar spindle poles, meiotic chromosomes and
spindle midzone during oocyte maturation. *Chromosoma* **107**:430–439.
doi:10.1007/s004120050327
Woodruff JB, Wueseke O, Viscardi V, Mahamid J, Ochoa SD, Bunkenborg J, Widlund PO,
Pozniakovsky A, Zanin E, Bahmanyar S, Zinke A, Hong SH, Decker M, Baumeister
W, Andersen JS, Oegema K, Hyman AA. 2015a. Regulated assembly of a
supramolecular centrosome scaffold in vitro. *Science* **348**:808–812.
doi:10.1126/science.aaa3923
Woodruff JB, Wueseke O, Viscardi V, Mahamid J, Ochoa SD, Bunkenborg J, Widlund PO,
Pozniakovsky A, Zanin E, Bahmanyar S, Zinke A, Hong SH, Decker M, Baumeister
W, Andersen JS, Oegema K, Hyman AA. 2015b. Regulated assembly of a
supramolecular centrosome scaffold in vitro. *Science* **348**:808–812.
doi:10.1126/science.aaa3923
Xu J, Shen C, Wang T, Quan J. 2013. Structural basis for the inhibition of Polo-like kinase
1. *Nat Struct Mol Biol* **20**:1047–1053. doi:10.1038/nsmb.2623
Zhang L, Shao H, Huang Y, Yan F, Chu Y, Hou H, Zhu M, Fu C, Aikhionbare F, Fang G, Ding
X, Yao X. 2011. PLK1 Phosphorylates Mitotic Centromere-associated Kinesin and
Promotes Its Depolymerase Activity*. *Journal of Biological Chemistry* **286**:3033–
3046. doi:10.1074/jbc.M110.165340
Zitouni S, Nabais C, Jana SC, Guerrero A, Bettencourt-Dias M. 2014. Polo-like kinases:
structural variations lead to multiple functions. *Nat Rev Mol Cell Biol* **15**:433–452.
doi:10.1038/nrm3819
1053

1054

METHODS

***C. elegans* strains & RNAi**

Strains used in this study were maintained at 20 degrees unless indicated otherwise. For a

complete list of strains, please refer to Table S1.

For RNAi-mediated depletions, the targeting sequence for *bub-1* was 2353-2935 and for *hcp-*

4, 967-2128, both from the first ATG codon. For double depletion, both sequences were

cloned in the same vector. All sequences were inserted into L4440 using the NEBuilder HiFi

DNA Assembly Master Mix (New England Biolabs) and transformed into DH5a bacteria. The

purified plasmids were then transformed into HT115(DE3) bacteria (Timmons et al., 2001).

RNAi clones were picked and grown overnight at 37°C in LB with 100 μg/ml ampicillin.

Saturated cultures were diluted 1:100 and allowed to grow until reaching an OD600 of 0.8–1.

Isopropyl-β-d-thiogalactopyranoside (IPTG) was added to a final concentration of 1 mM, and

cultures were incubated for 1 h at 37°C. Bacteria were then seeded onto NGM plates made

with agarose and 1 mM IPTG and allowed to dry. L4 worms were then plated on RNAi plates

and maintained at 20°C.

1070

CRISPR/Cas9

We used direct injection of in vitro–assembled Cas9-CRISPR RNA (crRNA) trans-activating

crRNA (tracrRNA) ribonucleoprotein complexes (Paix et al., 2017, 2015).

For mutation of Threonine 527 to Alanine in *bub-1*, we used the following crRNA (+ strand):

CCCCGCACAAGGAGTTCATT and repair template (- strand):

acttacTAATTTACTGAAAGTACTGCTGGTTGGAGCAACAAATACTGGAGCTTCCTGT

TCGTGAGTGCTTTCCTCCTCTTTATTTCCGAAATATTCATCAATGTTGACtAAATGA

ACTCCTTGTGCGGGgGcactaGTGACGAAATTACCACGAGACGGTTTGAAAAAGCCA

AACTCGATTTCATCGTCATAAActaaaa

1080

For mutation of Threonine 163 to Alanine in *hcp-4*, we used the following crRNA (- strand):

TGAAATATCAAGCGATCTCA and repair template (+ strand):

taataaatctataatttcagAGTGGAAAAGCTGGATTAAGcTGcagtgCaCCCAAGAGCTCGAGTG

ATACGTCGATGAGGTCtTTGAGATCGCTTGATATTTCACATGTCGTCAATACCGAT

C.

Each of the mixes was mixed with dpy-10 crRNA/repair template for screening (Arribere et

al., 2014).

1088

Live imaging of oocytes

A detailed protocol for live imaging of *C. elegans* oocytes was used with minor modifications

(Laband et al., 2018). Fertilized oocytes were dissected and mounted in 5 µl of L-15

blastomere culture medium (0.5 mg/mL Inulin; 25 mM HEPES, pH 7.5 in 60% Leibowitz L-

15 medium and 20% heat-Inactivated FBS) on 24×40 mm #1.5 coverslips. In experiments

where the PP1 analogue was used, 3IB-PP1 was diluted to the desired concentration in L-15

media from a 10 mM stock solution in ethanol. Control experiments were conducted with the

equivalent dilution of ethanol. The incubation time while dissection took place was typically

between 3-5 minutes, unless otherwise stated. Once dissection was performed and early

oocytes identified using a stereomicroscope, a circle of Vaseline was laid around the sample,

and a custom-made 24×40 mm plastic holder (with a centred window) was placed on top. The

sample was imaged immediately using 488 nm and 561 nm laser lines. Live imaging was

done using a CFI Plan Apochromat Lambda 60X/NA 1.4 oil objective mounted on a

microscope (Nikon Eclipse Ti) equiped with a Prime 95B 22mm camera (Photometrics), a

spinning-disk head (CSU-X1; Yokogawa Electric Corporation). Acquisition parameters were

controlled with NIS software (Nikon). For all live imaging experiments, partial projections

are presented. All files were stored, classified, and managed using OMERO (Allan et al.,

2012). Figures were prepared using OMERO.figure and assembled using Adobe Illustrator.

Representative movies shown in Supplementary material were assembled using Fiji/ImageJ

(Schindelin et al., 2012) custom-made macros.

1109

Viability assay

Three different PP1 analogues were tested for their effect on embryo viability. Three worms

were placed in NGM coated wells of 24-well plates containing vehicle ('control') or the

indicated concentration of each analogue, in triplicate wells per condition. Once ≥200

embryos had been laid per well (with the exception of 50 µM analogue, where ≥100 embryos

were analysed), adult worms were removed, and the exact number of embryos recorded.

Seventy two hours later, viable worms were counted. In experiments where the PP1 analogue

was used, 3IB-PP1 was diluted to the desired concentration in L-15 media from a 10 mM

stock solution in ethanol. Control experiments were conducted with the equivalent dilution of

ethanol. The incubation time while dissection took place was typically between 3-5 minutes,

unless otherwise stated.

1121

Immunofluorescence

Worms were placed on 4 µl of M9 worm buffer in a poly-D-lysine (Sigma, P1024)-coated

slide and a 24×24-cm coverslip was gently laid on top. Worms were dissected to release the

embryos, slides were placed on a metal block on dry ice for >10 min. The coverslip was then

flicked off with a scalpel blade, and the samples were fixed in methanol at 20°C for 30 min.

Secondary antibodies were donkey anti–sheep, goat anti-mouse, or goat anti-rabbit conjugated

to Alexa Fluor™ 488, Alexa Fluor™ 594, and Alexa Fluor™ 647 (1:1,000, Thermo

Scientific). Donkey anti-mouse and donkey anti-rabbit conjugated secondary antibodies were

obtained from Jackson ImmunoReserach. Embryos were mounted in ProLong Diamond

antifade mountant (Thermo Scientific) with DAPI. Primary antibodies were: α-PLK-1

(Budirahardja and Gönczy, 2008), α-HCP-4 (Oegema et al., 2001), α-BUB-1, purified in

house after immunisation of rabbits using the sequence in (Desai et al., 2003),

1134

1135 **Sequence alignments**

1136 Sequences shown in Figures 2C and 5B were aligned with Clustal Omega (Sievers et al.,

1137 2011) and visualised with Jalview (Waterhouse et al., 2009).

1138

Protein Purification

GST-BUB-1$^{190\text{-}628}$ and GST-HCP-4$^{1\text{-}214}$ proteins were expressed in *Escherichia coli* BL21

DE3 bacteria by diluting a saturated culture 1/100 in LB media supplemented with 35 µg/ml

ampicillin and incubating at 37°C/200 rpm until OD_{600} 0.6-0.8 was reached. IPTG was then

added to a final concentration of 100 µM and cultures were incubated at 20°C/200 rpm for 16-

18 h. The bacterial cultures were then centrifuged (20 min/6,250 g/4°C) and pellets

resuspended in lysis buffer (50 mM Tris-HCl, pH 7.5; 150 mM NaCl; 0.5 mM TCEP, 1X

Roche cOmplete protease inhibitors, EDTA free; 30-35 ml/L of culture). The cell suspension

was then sonicated (2 min 40 sec, 20 sec on/40 sec off) before centrifugation (45 min, 27,250

g, 4°C) to remove insoluble material. GSH sepharose beads were washed with 10 column

volumes (CV) of MilliQ water and equilibrated with 10 CV of binding buffer (50 mM Tris-

HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP) before filtered (0.22 µm PES filter) lysate was

added and batch bound at 4°C for 1.5-2 h. The beads were then collected in a column and

washed with at least 10 CV of binding buffer before being transferred to a falcon tube and

incubated overnight with GST-tagged 3C protease. After cleavage, the beads were collected

in a column and the flowthrough containing cleaved protein was concentrated in a Vivaspin

centrifugal concentrator. The protein was then further purified by size exclusion

chromatography in a Superdex 200 10/300 column (run in SEC buffer, see below) and

concentrated before being flash frozen in liquid nitrogen and stored at –80.

MBP-PLK1[PBD] (6xHIS-MBP tagged human PLK1[345-603], as per (Singh et al., 2021)) was

expressed and purified in the same manner as the GST tagged proteins with the following

exceptions: filtered lysate was passed through a cobalt-NTA column and washed with at least

10 CV of binding buffer (see buffers below) before elution in 0.5 CV fractions. Fractions

containing the protein were concentrated using Vivapsin centrifugal filters and further

purified by size exclusion chromatography (Superdex 200 10/300) in SEC buffer. Lysis buffer

(50 mM Tris-HCl, pH 7.5; 500 mM NaCl; 10 mM imidazole; 0.5 mM TCEP; 1X Roche

cOmplete protease inhibitors, EDTA free; 50 ml/L of culture); binding buffer (50 mM Tris-

HCl, pH 7.5; 500 mM NaCl; 10 mM imidazole; 0.5 mM TCEP); elution buffer (50 mM Tris-

HCl, pH 7.5; 150 mM NaCl; 200 mM imidazole; 0.5 mM TCEP); SEC buffer (50 mM Tris-

HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP).

1169

Kinase assays

Unless otherwise stated, kinase assays were conducted with 55 nM Cdk1:Cyclin B (Thermo

Scientific) and/or 75 nM PLK-1, produced as described in (Tavernier et al., 2015), at 30ºC in

kinase buffer: 50 mM Tris-HCl, pH 7.5; 1 mM ATP; 10 mM $MgCl_2$; 0.5 mM TCEP; 0.1 mM

EDTA. For the assays in Figure 2E,F and Figure 2-figure supplement 1; 0.4 μg/μl (8.2 μM) of

BUB-1[190-628] substrate was used and incubated under the above conditions for 1 h (Figure 2E)

or up to 2 h (Figure 2F, Figure 2-figure supplement 1). The assays in Figure 5F,G were

conducted in the above conditions for up to 1 h with a substrate concentration of 0.2 μg/μl

(8.2 μM). The assay in Figure 2D was conducted with 165 nM Cdk1:Cyclin B and 170 nM

PLK-1 and incubated for 30 min at 37°C with the following buffer: 40 mM Tris-HCl, pH 7.5;

100 μM ATP, 10 mM MgCl$_2$. BUB-1$^{190-628}$ concentration was 0.2 μg/μl (4.1 μM).

In all assays, aliquots of protein at the indicated timepoints were immediately added to an

equal volume of 2X LDS buffer (Thermo) and incubated at 70 degrees for 15 min. Assays

were assessed by SDS-PAGE combined with ProQ diamond phosphoprotein and coomassie

staining.

1185

***In vitro* Binding Assays**

BUB-1$^{190-628}$ or CENP-C$^{HCP-4(1-214)}$ recombinant proteins were incubated for 50-60 min at

30°C in the presence or absence of ~100 nM Cdk1:Cyclin B in kinase buffer (see 'kinase

assay' section of methods). The respective proteins were then incubated for 50 min on ice

with MBP-PLK1^{PBD} at a concentration of 20 μM in SEC buffer (50 mM Tris-HCl, pH 7.5;

150 mM NaCl; 0.5 mM TCEP). Assays were then centrifuged (13.3k rpm, 10 min, 4°C)

before being loaded onto a Superdex 200 10/300 SEC column. In all assays, 0.2 ml fractions

were collected and selected fractions were assessed by SDS-PAGE with coomassie staining.

More specifically; for the binding assays with BUB-1$^{190-628}$ in Figure 3A,C, 200 μl of binding

assay was loaded into a 0.5 ml loop and loaded onto the column with 2 ml of SEC buffer. For

the CENP-C$^{HCP-4(1-214)}$ assays in Figure 6A,C, 100 μl of assay was loaded onto a 100 μl loop

and loaded onto the column with 0.4 ml of SEC buffer.

1198

Size-exclusion chromatography multi-angle light scattering (SEC-MALS)

An Agilent 1260 Infinity II HPLC system was used to inject 50 μl of protein (ranging from

0.5-2.5 mg/ml) onto a Superdex 200 10/300 gel filtration column (Cytiva) coupled to a Wyatt

miniDawn multi-angle light scattering system and Wyatt Optilab differential refractive index

(dRI) detector. A dn/dc value of 0.185 was used, UV extinction coefficients were calculated

from RI and molecular weights were calculated using Zimm plot extrapolation. Astra v8.1

was used for data analysis. The following buffer was used in all experiments: 50 mM Tris-

HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP.

1207

Western Blot

In vitro kinase assays were conducted as described above and separated by SDS-PAGE on 4-

12% Bis-Tris gels (Thermo) with ~20 ng of protein loaded per well for the α-BUB-1 or α-

HCP-4 blots and either 80 ng (BUB-1$^{190-628}$) or 75 ng (CENP-C$^{HCP-4(1-214)}$) loaded per lane for

the phosphospecific antibodies. Transfer was conducted for 90 min at 90 V in transfer buffer

(25 mM Tris, 200 mM glycine, 20% methanol) before membrane was washed in PBS-T (PBS

with 0.1% tween-20) and blocked with Intercept PBS blocking buffer (Licor). Membranes

were washed in PBS-T before incubation with primary antibodies overnight at 4°C. For the

peptide competition blots, peptide or phosphopeptides were added to a final concentration of

1 μM alongside the primary antibodies. The next day, membranes were washed in PBS-T

before incubation with Licor secondary antibodies (1:25000) in 5% milk for 1 h. Membranes

were then washed once more in PBS-T before scanning in a Licor Odyssey CLx.

Worm extract western blots (Figure 9-figure supplement 1) worms were grown and

transferred to RNAi plates as described above. 50 worms per lane were picked and washed

with M9 + 0.1% triton (x2) and M9 (x1). LDS buffer was then added to a final concentration

of 1X and samples were heated (70°C, 5 min), sonicated (Diagenode cell disruptor, high

power, 10 cycles, 30s ON/30s OFF), and heated again (70°C, 10 min) before western blots

were conducted following the protocol described above.

1226

Fluorescence Polarisation

Fluoresence polarisation assays were conducted using FITC-labelled peptides (peptide

sequences indicated in figures). For all assays: a 1:2 dilution series of MBP-PLK1^{PBD} was

conducted in FP buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.5 mM TCEP) with a

constant concentration of 100 nM FITC-labelled peptide. Assays were left for 20-60 min

before being centrifuged and 10 μl of each concentration was loaded onto a black 384 well

plate (Greiner) in triplicate. Plates were then centrifuged (2k rpm, 2 min) before being

analysed in a PheraStar FS (BMG Labtech) under the following conditions: excitation: 485

nm, emission: 520 nm, 25°C, settling time 0.2 s, 50 flashes per well, gain and focal height

automatically adjusted prior to measurement. Mean mP for each triplicate was then used as

the value for each independent experiment. Data represent the average of 3 independent

experiments with K$_D$ values being estimated using GraphPad Prism 9 "Nonlinear regression,

One Site – Total" equation. The uncertainity indicated represents the standard deviation of the

calculated K$_D$ from each independent replicate.

1241

Phenotype Analysis

We defined misalignment as metaphase defects. Alignment was counted within the 5 frames

(1 min) before anaphase onset, which was the frame prior to the detection of two separating

chromosome masses. When one bivalent was misaligned (either in angle or distance to

metaphase plate) it was recorded as mild metaphase defect. When two or more bivalents were

misaligned, this was considered as a severe metaphase defect. The anaphase phenotype was

assess during segregation. If lagging chromosome were detected during chromosome

segregation but this was resolved before polar body extrusion, it was scored as mild anaphase

defect. If we could not detect two segregating masses of chromosomes or if these masses

differed in size/intensity, it was quantified as a severe anaphase defect. Polar body defect was

recorded when no polar body was extruded or if all of the maternal DNA content was

extruded as a polar body, with no maternal DNA remaining in the cytoplasm. Graphs were

prepared using Graphpad Prism 9.0.

1255

1256   **Table S1**: *C. elegans* strains used in this study

PLK-1::sfGFP; mCherry::histone	FGP263	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV*	This study
GFP::HCP-4; mCherry::histone; TIR1	FGP311	*hcp-4(lt72[GFP::hcp-4])I; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV)I; ieSi65 [sun-1p::TIR1::sun-1 3'UTR + Cbr-unc-119(+)] II; unc-119(ed3)III*	This study
hcp-4(T163A)	FGP669	*hcp-4(fgp58[hcp-4(T163A)])I*	This study
bub-1(T527A)/hT2	FGP672	*bub-1(fgp4[bub-1(T527A)])I; hT2 [bli-4(e937) let-?(q782) qIs48](I;III)*	This study
GFP::tubulin; mCherry::histone; bub-1(T527A)/hT2	FGP674	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; bub-1(fgp4[bub-1(T527A)])I; hT2 [bli-4(e937) let-?(q782) qIs48](I;III)*	This study
GFP::tubulin; mCherry::histone; plk-1(as)	FGP675	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; plk-1((lt106[plk-1 C52V] lt109[plk-1 L115G])III*	This study
mScarlet::KNL-3; PLK-1::sGFP	FGP718	*knl-3[dha19(mScarlet:: 3XFLAG::knl-3)]V; plk-1(lt18[plk-1::sGFP]::loxp)III.*	This study
PLK-1::sGFP; mCherry::histone; hcp-4(T163A)	FGP719	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV; hcp-4(fgp58[hcp-4(T163A)])I*	This study
mScarlet::KNL-3; PLK-1::sGFP; hcp-4(T163A)	FGP721	*knl-3[dha19(mScarlet:: 3XFLAG::knl-3)]V; plk-1(lt18[plk-1::sGFP]::loxp)III; hcp-4(fgp58[hcp-4(T163A)])I*	This study

GFP::ASPM-1; mCherry::histone; plk-1(as)	FGP725	*or1935[GFP::aspm-1] I; itIs37[pie-1p::mCherry::H2B::pie-1 3'UTR + unc-119(+)] IV; plk-1((lt106[plk-1 C52V] lt109[plk-1 L115G])III*	This study
GFP::tubulin; mCherry::histone; bub-1(wt)	FGP731	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	This study
GFP::tubulin; mCherry::histone; hcp-4(T163A); bub-1(wt)	FGP732	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; hcp-4(fgp58[hcp-4(T163A)])I; unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	This study
GFP::tubulin; mCherry::histone; bub-1(T527A)	FGP733	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	This study
GFP::tubulin; mCherry::histone; hcp-4(T163A); bub-1(T527A)	FGP734	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I; hcp-4(fgp58[hcp-4(T163A)])I; unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	This study
PLK-1::sGFP; mCherry::histone; bub-1(wt)	FGP735	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37[pAA64; pie-1/mCHERRY::his-58; unc-119(+)]IV; unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	This study
PLK-1::sGFP; mCherry::histone; hcp-4(T163A); bub-1(wt)	FGP736	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37[pAA64; pie-1/mCHERRY::his-58; unc-119(+)]IV; hcp-4(fgp58[hcp-4(T163A)])I; unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	This study
PLK-1::sGFP; mCherry::histone; bub-1(T527A)	FGP737	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37[pAA64; pie-1/mCHERRY::his-58; unc-119(+)]IV; unc-119(ed3)III;ltSi1346[pTK070;Pbub-1::BUB-1 T527A reencoded; cb-unc-119(+)]II*	This study

PLK-1::sGFP; mCherry::histone; hcp-4(T163A); bub-1(T527A)	FGP738	*plk-1(lt18[plk-1::sGFP]::loxp)III; ltIs37 [pAA64; pie-1/mCHERRY::his-58; unc-119 (+)]IV; hcp-4(fgp58[hcp-4(T163A)])I; unc-119(ed3)III;ltSi1346[pTK070;Pbub-1::BUB-1 T527A reencoded; cb-unc-119(+)]II*	This study
GFP::tubulin; mCherry::histone	OD868	*ltSi220[pOD1249/pSW077; Pmex-5::GFP::tbb-2::operon_linker::mCHerry::his-11; cb-unc-119(+)]I*	(Green et al., 2013) PMID: 24217623
bub-1 Mos Chr II	OD991	*unc-119(ed3)III;ltSi268[pOD/pTK013; Pbub-1::Bub1 reencoded; cb-unc-119(+)]II*	(Houston et al., 2022) doi.org/10.1101/2022.10.07.511323
PLK-1::sfGFP	OD2425	*plk-1(lt18[plk-1::sGFP]::loxp)III*	(Martino et al., 2017) PMID: 29065307
bub-1(T527A) Mos Chr II	OD2721	*unc-119(ed3)III;ltSi1346[pTK070;Pbub-1::BUB-1 T527A reencoded; cb-unc-119(+)]II*	(Houston et al., 2022) doi.org/10.1101/2022.10.07.511323
GFP::HCP-4	OD3410	*hcp-4(lt72[GFP::hcp-4])I*	(Cheerambathur et al., 2019) PMID: 30827898
plk-1(as)	OD3697	*plk-1((lt106[plk-1 C52V] lt109[plk-1 L115G])III*	(Gómez-Cavazos et al., 2020) PMID: 32619481
GFP::ASPM-1	EU2861	*or1935[GFP::aspm-1]I*	(Connolly et al., 2015) PMID: 26370499
mScarlet::KNL-3	DKC78	*knl-3[dha19(mScarlet::3XFLAG::knl-3)]V*	This study

1257

Figure 1

Figure 2

Figure 3

Figure 4

Figure 6

Figure 9