The plastidial protein MRC promotes starch granule initiation in wheat leaves but

delays B-type granule initiation in the endosperm

Jiawen Chen[1], Yi Chen[1], Alexander Watson-Lazowski[1,4], Erica Hawkins[1], J. Elaine Barclay[1],

Brendan Fahy[1], Robin Denley Bowers[1], Kendall Corbin[2,5], Frederick J. Warren[2], Andreas

Blennow[3], Cristobal Uauy[1], David Seung[1]

[1.] John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK

[2.] Quadram Institute, Norwich Research Park, Norwich, NR4 7UQ, UK

[3.] University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg, Copenhagen,

Denmark

[4.] Current Address: Harper Adams University, Newport, Shropshire, TF10 8NB, UK

[5.] Current Address: Department of Horticulture, College of Agriculture, Food and

Environment, University of Kentucky, Lexington, Kentucky, USA

Address correspondence to: David.Seung@jic.ac.uk

Short title: Tissue-specific roles of MRC in granule initiation in wheat

Abstract

The spatial and temporal patterns by which starch granules initiate vary greatly between

species and organs, but molecular factors that contribute to these diverse patterns are poorly

understood. We reveal distinct organ-specific roles of the MYOSIN-RESEMBLING

CHLOROPLAST PROTEIN (MRC) in regulating granule initiation in the endosperm and

leaves of wheat. We isolated three independent TILLING mutants of tetraploid wheat

(*Triticum turgidum* cv. Kronos) with premature stop or missense mutations in the A-genome

homeolog, which we showed to be the only active homeolog in tetraploid wheat due to a

disruption of the B-genome homeolog. Wheat endosperm contains both large A-type granules

initiated during early grain development, and small B-type granules that initiate about $10 - 15$

days later. The *mrc* mutants had significantly smaller A-type granules and a higher relative

volume of B-type granules in the endosperm than the wild type. Whereas B-type granules

initiated 15 - 20 days post anthesis (dpa) in the wild-type, they appeared as early as 10 dpa in

the *mrc-1* mutant, suggesting a role for MRC in suppressing B-type granule initiation during

early grain development. By contrast, MRC promotes granule initiation in leaves: mutants

carrying premature stop mutations in *MRC* had fewer granules per chloroplast than the wild

type. These contrasting roles of MRC among wheat organs provide new insight into

functional diversification of granule initiation proteins, and suggest that they may facilitate

the diverse patterns of granule initiation observed across species and organs.

Introduction

Starch is the major storage carbohydrate in leaves, seeds and storage organs of most plants. It is synthesised in plastids as insoluble granules composed of two glucose polymers: amylopectin and amylose. Amylopectin is a branched polymer with linear α-1,4-glucan chains and α-1,6-branch points, and forms the semi-crystalline starch granule matrix - typically constituting 70-90% *w/w* of starch (Smith and Zeeman, 2020). Amylose constitutes 10 - 30% *w/w* of starch and is composed primarily of long α-1,4-linked linear chains (Seung, 2020). The biosynthesis of the starch polymers is relatively well understood at the molecular level, and is generally conserved between leaf and storage starches (Smith and Zeeman, 2020). However, we are only beginning to understand how starch granule formation is initiated, and the factors underpinning the vast diversity in granule initiation patterns observed between different organs and species (Tetlow and Emes, 2017; Seung and Smith, 2019; Chen et al., 2021).

A prime example of diverse granule initiation patterns between species can be observed in the seed endosperms of grasses (Matsushima et al., 2013). Grass species of the Triticeae, including important cereal crops such as wheat, barley and rye, have a unique bimodal size distribution of starch granules in the grain endosperm – containing large flattened A-type granules (20-30 μm in diameter) and small round B-type granules (2-7 μm in diameter) (Howard et al., 2011). The initiation of these two different types of granules is both spatially and temporally separated: A-type granules initiate in amyloplasts as early as 4 days post anthesis (dpa), whereas B-type granules initiate 10-15 days after the A-type granules (i.e., around 15-20 dpa) and at least partly within stromules that emanate from the amyloplast (Parker, 1985; Bechtel et al., 1990; Langeveld et al., 2000; Howard et al., 2011). This is distinct from most other grass species, which produce "compound" starch granules – where multiple granules initiate early during grain development in each amyloplast and eventually fuse (e.g., in rice) (Matsushima et al., 2013). Our recent work has identified proteins that are important for determining the initiation of bimodal starch granules in wheat. These include B-GRANULE CONTENT1 (BGC1) in wheat/*Aegilops* - which is orthologous to FLOURY ENDOSPERM6 (FLO6) in barley and rice and PROTEIN TARGETING TO STARCH 2 (PTST2) in Arabidopsis and *Brachypodium* (Peng et al., 2014; Saito et al., 2017; Seung et al., 2017; Chia et al., 2020; Watson-Lazowski et al., 2022). BGC1 in wheat has a dose-dependent effect on granule initiation, where partial reductions in gene dosage can almost eliminate B-type granules without affecting A-type granule formation, whereas complete loss of function

also causes defective A-type granule formation, including the formation of some

compound/semi-compound granules that arise from multiple initiations (Howard et al., 2011;

Chia et al., 2020; Saccomanno et al., 2022). STARCH SYNTHASE 4 (SS4) is also required

for normal A-type granule formation. In its absence, compound granules form in place of

most A-type granules (Hawkins et al., 2021). The increased number of initiations per

amyloplast that led to compound granule formation in these mutants was unexpected since in

Arabidopsis leaves both SS4 and PTST2 promote granule initiation, and mutants lacking

either protein have reduced numbers of starch granules per chloroplast (Roldán et al., 2007;

Seung et al., 2017). These observations suggest that the proteins involved in granule initiation

are to some extent conserved between species and organs, but they can act differently

depending on the patterns of granule initiation in the species/tissue.

To gain further insight into such differences, we explored the function of wheat MYOSIN-

RESEMBLING CHLOROPLAST PROTEIN (MRC, also known as PROTEIN INVOLVED

IN STARCH INITIATION, PII1) in both chloroplasts of leaves and amyloplasts of the

endosperm. MRC is a long coiled-coil protein that promotes granule initiation in Arabidopsis

leaves, as most chloroplasts in the *mrc* mutant contain only a single granule (Seung et al.,

2018; Vandromme et al., 2019). The exact function of MRC is unknown, but it is proposed to

act via an interaction with SS4 and PTST2 (Seung et al., 2018; Vandromme et al., 2019).

Consistent with previous work in Arabidopsis, we found that MRC promotes starch granule

initiation in wheat leaves, but we discovered an unexpected, distinct role for MRC in the

temporal control of B-type granule initiation in the wheat endosperm. MRC is expressed at

the early stages of grain development, and wheat *mrc* mutants have severe alterations in the

starch granule size distribution relative to the wild type, with smaller A-type granules and a

higher relative volume of B-type granules. We demonstrate that this phenotype arises from

the early initiation of B-type granules in the mutant, suggesting MRC represses B-type

granule initiation during early grain development. This role of MRC in the wheat endosperm

demonstrates how the function of granule initiation proteins can be adapted to mediate

specific patterns of granule initiation among different species/tissues.

Results

The wheat orthologs of MRC are encoded on chromosomes 6A and 6D

The starch granule initiation protein MRC is highly conserved among land plants (Seung et al., 2018). To determine the role of MRC in wheat, we searched the wheat genome for genes encoding MRC orthologs. We ran a BLASTp search using the amino acid sequence of Arabidopsis MRC (*At*MRC, At4g32190) against the protein sequences from both tetraploid durum wheat (Svevo v1.1)(Maccaferri et al., 2019) and hexaploid bread wheat (IWGSG Chinese Spring)(Appels et al., 2018) genomes on Ensembl Plants. For hexaploid wheat (*Triticum aestivum*), the two top protein hits were TraesCS6A02G180500.1 (encoded on chromosome 6A) and TraesCS6D02G164600.1 (encoded on chromosome 6D), which shared 95% sequence identity with each other and were predicted as homeologs on Ensembl. Both genes had a two-exon structure like the Arabidopsis gene (Seung et al., 2018), and were in syntenic positions on the A and D genomes (Figure 1). For tetraploid durum wheat (*Triticum turgidum*), the top protein hit was TRITD6Av1G081580.1 (encoded on chromosome 6A), which was identical in nucleotide and amino acid sequence to TraesCS6A02G180500.1. To determine whether these proteins were true orthologs of *At*MRC, we repeated the phylogenetic analyses of MRC homologs from our previous study (Seung et al., 2018) with the wheat protein sequences included. The 6A and 6D proteins clustered together on the tree, distinctly within the grass clade containing the rice and maize sequences (Supplemental Figure 1A). This confirms that the proteins are the wheat orthologs of MRC. They will hereafter be referred to as *Ta*MRC-A1 (TraesCS6A02G180500) or *Tt*MRC-A1 (TRITD6Av1G081580), and *Ta*MRC-D1 (TraesCS6D02G164600). Notably, we did not find a full gene model for MRC on chromosome 6B, or anywhere else on the B genome, either in the durum or bread wheat genome.

To investigate why no homeolog was detected on chromosome 6B, we looked at the syntenic region of chromosome 6B in Chinese Spring, where there was a stretch of sequence that had homology to exon 2 and the beginning of the 3'UTR (Figure 1A). Interestingly, around 14kb downstream of that, there was a region highly similar to the end of the 3' UTR of *TaMRC-A1*. We looked at the transposable element annotation of the wheat reference genome around the exon 2 fragment (Daron et al., 2014), and a complete *gypsy* retrotransposon was annotated between the two 3'UTR fragments (Figure 1B). Further, we identified the 5 bp target site duplication (GAGAT, which is part of the 3'UTR) and the inverted terminal repeat

(TGTAA and TTACA at the start and end of the retrotransposon, respectively) characteristic

of retrotransposon insertions. The distance between the 5' end of the exon 2 fragment and its

upstream neighbouring gene (TraesCS6B02G20500, a respiratory burst oxidase homolog)

was much larger than the distance between *Ta*MRC and the homeologs of the same

neighbouring gene on 6A and 6D, indicating a large insertion in this 6B region (Figure 1A).

Indeed, a fragment of another *gypsy* retrotransposon was found ca. 16 kb upstream of the

exon 2 fragment. Additionally, there was a sequence with homology to the 5' UTR of

TaMRC-A1 (85% identity over 243 bp) just 3 bp upstream of the exon 2 fragment, which

suggested that a ~1.3 kbp deletion (based on A-genome distances) removed some of the 5'

UTR, all of exon 1, intron 1 and the start of exon 2 of *TaMRC-B1*. Similar to Chinese Spring,

we found identical disruptions in *TaMRC-B1* sequences with retrotransposon insertions and

deletions in ten additional wheat genome assemblies (Walkowiak et al., 2020). Overall, it

appears that a deletion and a series of transposon insertions severely disrupted *MRC* on

chromosome 6B in bread wheat.

Since a B-genome copy was also absent from durum wheat, it is likely that the disruption of

MRC on chromosome 6B preceded the second hybridisation that resulted in hexaploid wheat.

To further investigate when the disruption of *MRC-B1* occurred, we looked for *MRC-B1* in

more tetraploid wheat accessions by aligning genome sequencing reads from *Triticum*

dicoccoides (wild emmer) (n=10) and *Triticum turgidum* ssp. *durum* (pasta wheat) (n=12)

against the A and B genomes of Chinese Spring (Zhou et al., 2020) (Supplemental Table 1).

The exon 1 deletion and the retrotransposon insertion at the 3' end were detected in all

accessions (except for a few lines that had poor sequencing depth in the region), suggesting

that *TaMRC* on 6B was disrupted before or immediately after the hybridization of diploid

ancestors carrying the A and B genomes. To distinguish these possibilities, we examined

MRC in *Aegilops speltoides*, the diploid species thought to be most closely related to the

progenitor species of the wheat B genome. We ran a BLASTn search on the genome

assembly of the *Aegilops speltoides* accession TS01 using the coding sequence of *TaMRC-A1*

(Li et al., 2022) (Supplemental File 1). The top hit was on chromosome 6S (homologous to

chromosome 6B in wheat) with intact exons one and two (97.5% sequence identity), and the

translated protein sequence had 96% identity to *Ta*MRC-A1 with BLASTp. Thus, it is likely

that *MRC* is intact in *Aegilops speltoides*, suggesting that the loss of the B-homeolog

occurred shortly after the hybridisation that gave rise to tetraploid wheat (Figure 1C). It is

therefore expected that all tetraploid wheats have one *MRC* homeolog (on chromosome 6A),

and all hexaploids have two (on chromosomes 6A and 6D).

MRC is expressed in leaves and developing endosperm

Using the hexaploid wheat expression browser (Borrill et al., 2016; Ramirez-Gonzalez et al.,

2018), we found that transcripts of the 6A and 6D homeologs were present in both leaves and

grains, suggesting that MRC plays a role in both these tissues (Figure 2A). To obtain

temporal information on MRC expression in the endosperm during grain development, we

performed RNAseq analysis on dissected endosperms of durum wheat (*Triticum turgidum* cv.

Kronos) throughout grain development. The full dataset is available in our accompanying

paper (Chen et al., 2022). The Kronos variety was chosen as it is the genetic background of

the tetraploid wheat TILLING mutants characterised below. *TtMRC-A1*

(TRITD6Av1G081580.1) showed a peak of expression during early grain development (8

dpa), but strongly decreased in expression between 10 – 20 dpa (Figure 2B). It therefore

appears that *MRC* is expressed in the wheat endosperm almost exclusively during early grain

development.

Loss of MRC does not affect the growth of wheat plants, or grain development

To study the function of MRC in wheat, we obtained mutants in durum wheat (*Triticum*

turgidum cv. Kronos) defective in *MRC*. We used the wheat *in silico* TILLING mutant

resource, which has an EMS-mutagenised population of Kronos with exome-capture

sequencing data for identification of lines with mutations of interest (Krasileva et al., 2017).

We obtained three mutants that were likely to cause a loss of function in *TtMRC-6A* (Figure

1B). The K3272 and K4681 lines contained premature stop codons in place of codons for the

258[th] and 550[th] amino acids respectively. In addition, we obtained a third line that contained a

missense Leu289Phe mutation, which was predicted to be deleterious to protein function by

SIFT scoring (Ng and Henikoff, 2006). The Leu[289] residue is highly conserved in all MRC

orthologs, and its mutation to a Phe residue is predicted to disrupt coiled coil formation in the

region of the residue (Supplemental Figure 1B). Since the 6B homeolog of MRC has likely

become a pseudogene, we predicted that *TtMRC-A1* would be the only functional *MRC*

homeolog in tetraploid wheat. However, to rule out the possibility that the fragment of exon 2

on chromosome 6B affects MRC function, we also obtained the K4305 and K3078 lines

which contain two different premature stop codon mutations in the putative reading frame of

the exon. We generated the *mrc-1* lines by crossing K3272 and K3078, and the *mrc-2* lines by

crossing K4681 and K4305, where we isolated lines homozygous for either the 6A (F_2 *aa*BB)

or 6B (F_2 AA*bb*) mutation, or both (F_2 *aabb*). The *mrc-1* double mutant line was backcrossed

twice to wild type (WT), and the wild-type segregant (BC_2F_2 AABB) and the homozygous

double mutant (BC_2F_2 *aabb*) were selected. These will be hereafter referred to as BC2 AABB

and BC2 *aabb*. No backcrossing was done for the *mrc-2* line. The *mrc-3* line contained the

K598 missense mutation, and no crossing was done.

There was no consistent effect of the independent *mrc* mutations on plant growth or grain

development under our growth conditions. None of the *mrc* mutant plants appeared different

from WT with respect to growth or the number of tillers per plant (Figure 3A, B,

Supplemental Table 2A). The number of grains per plant also did not differ in the mutants,

except for a slight decrease in *mrc-3* and the wild-type segregant (*mrc-1* BC2 AABB)

compared to the WT (Figure 3C, Supplemental Table 2B). The morphology of the mature

grains of the mutants was indistinguishable from the WT (Figure 3D), and there were no

differences in thousand grain weight (TGW) and grain size between the WT and any of the

three mutants *mrc-1, mrc-2, mrc-3*, or between the WT and the backcrossed *mrc-1* (*mrc-1*

BC2 *aabb*); but the wild-type segregant (*mrc-1* BC2 AABB) had a slightly higher TGW and

grain size compared to WT (Figure 3E, F, Supplemental Table 2C,D). This suggests that

some of the background mutations in the wild-type segregant may have affected grain

development, but these effects are small.

Loss of MRC greatly alters starch granule size distributions in the endosperm

To determine the effect of the *mrc* mutations on starch synthesis in grains, we first measured

total starch content of mature grains. Starch content was largely similar between WT and the

mrc mutants. Although some pairwise comparisons showed $p<0.05$ (WT with *mrc-2* and *mrc-*

3), the confidence intervals of the difference in means was still close to zero, suggesting that

any effect of the mutations was small (Figure 4A). Coulter counter measurements revealed

that there were some minor differences between genotypes in the total number of granules

relative to grain weight (granules/mg grain) (Figure 4B) but overall there was not a strong

effect of loss of MRC. The backcrossed *mrc-1* mutant had more starch granules per unit grain

weight than both the WT and the wild-type segregant, but the non-backcrossed *mrc-1* mutant

was not significantly different to the WT. The *mrc-2* mutant also had relatively more granules

than the WT, but *mrc-3* did not.

Granule size distributions were determined from the Coulter counter data by plotting the

percentage of starch volume in each size bin relative to the total volume of starch measured.

We observed clear bimodal distributions for all genotypes, with a peak corresponding to A-

type granules (~ 18 – 25 μm) and a peak corresponding to B-type granules (~3 – 10 μm).

However, the distribution profiles in the mutants were very different from the WT, with the

peak corresponding to B-type granules being more prominent in the mutants (Figure 5A). The

profiles of the backcrossed *mrc-1* lines looked similar to their non-backcrossed equivalents.

We fitted a bimodal log-normal distribution curve to the profiles of each sample to estimate

the total volume percentage of B-type granules and the mean sizes of A- and B-type granules.

Comparing the means of these extracted values between genotypes showed a higher B-type

granule percentage (by volume) for all three mutants (*mrc-1, mrc-2* and *mrc-3*) compared to

the WT (Figure 5B). The strongest increase was seen for *mrc-1*, and this increase was

consistent when comparing the double backcrossed *mrc-1* (*mrc-1* BC2 *aabb*) with WT and

the wild-type segregant (*mrc-1* BC2 AABB). There was a small increase in the volume

percentage of B-type granules in the wild-type segregant compared to the WT, but the

difference was much smaller than between the other genotypes. The higher B-type granule

volume percentage in *mrc* mutants could be due to an increase in both B-type granule size

and number. The mean B-type granule size was larger than WT for *mrc-1* (and *mrc-1* BC2

aabb) and *mrc-2*, but not *mrc-3* (Figure 5D). By contrast, the mean A-type granule diameter

was smaller for all three mutants than for WT (Figure 5C). Thus, the increased proportion of

B-type granule volume may be due to a combination of smaller A-type granules and larger

and/or more numerous B-type granules.

We also explored whether the *mrc* mutations affected starch granule shape. Examination of

iodine-stained thin sections of mature grains using light microscopy showed that, like the

WT, all mutants had flattened A-type and round B-type granules (Figure 6A). Similarly, no

defects in A- or B-type granule shape were observed in the mutants using scanning electron

microscopy (SEM) (Figure 6B). Starch polymer structure and granule composition was not

affected by loss of functional MRC. The *mrc-1* mutant, with the strongest alteration in

granule size distribution, had normal amylopectin structure and amylose content

(Supplemental Figure 2).

We sought experimental evidence about the contribution of the fragment of exon 2 on

chromosome 6B to the observed differences in granule size distribution between genotypes

(Supplemental Figure 3). Quantification of starch granule size distribution in the full set of

homozygous genotypes resulting from the crosses that yielded the *mrc-1* and *mrc-2* mutants

[indicated as *aa*BB (6A mutant), AA*bb* (6B mutant) and *aabb* (6A and 6B double mutant)]

showed that WT and AA*bb* genotypes had identical granule distributions. The *aa*BB and

aabb had different distributions from the WT but were identical to each other. These data

showed that the fragment of exon 2 on chromosome 6B has no influence on granule size

distribution. They are consistent with the suggestion that this is a pseudogene, and hence that

the 6A copy of *MRC* is the only functional homeolog in tetraploid wheat.

Overall, these data suggest that MRC is required for the normal size distribution of starch

granules in wheat endosperm. In tetraploid wheat, mutants lacking in the 6A copy of *MRC*

consistently had a higher relative volume of B-type granules in the endosperm than the WT,

and smaller A-type granules. This change in granule size distribution occurred without

accompanying changes in total starch content, starch granule shape, amylose content or

amylopectin structure.

Loss of MRC results in the early initiation of B-type granules

To understand how MRC affects the size distribution of endosperm starch granules, and its

specific effects on A-type or B-type granules, we investigated granule initiation during grain

development in the *mrc* mutant with the strongest phenotype, *mrc-1*. We measured the total

starch content and number of starch granules in dissected endosperms of developing grains

harvested 8, 14, 20 and 30 days post anthesis (dpa). The total starch content of the endosperm

increased between each time point, and there was no significant difference between the

mutant and the WT at any time point (Figure 7A). At the 8 dpa timepoint, the mutant and the

WT contained similar numbers of starch granules. Interestingly, for the two subsequent time

points (14 dpa and 20 dpa), the mutant endosperms contained almost twice as many starch

granules as the WT, despite similar starch contents (Figure 7B). The largest increase in

granule number during grain filling was observed between the 20 and 30 dpa timepoints in

the WT, and between the 14 and 20 dpa timepoints in the mutant. At the 30 dpa timepoint,

the difference in granule number between the mutant and WT decreased. We also noted that

in both the WT and mutant, the number of starch granules decreased between the 8 and 14

dpa timepoints. The reason for this is unknown, but it has also been observed in *Aegilops*

species – which are close relatives of wheat (Howard et al., 2011).

In WT endosperm, there was a unimodal distribution of starch granule sizes at the 8 and 14

dpa timepoints, and only A-type granules with their characteristic flattened morphology were

observed using SEM (Figure 8). The A-type granules grew substantially in size between the

two timepoints, seen as a shift in the granule size distribution peak. B-type granules only

became prominent at the 20 dpa timepoint. In the *mrc-1* mutant A-type granules were initially

the same size as those of WT (at 8 dpa), but subsequently grew more slowly than wild-type

granules. By contrast, B-type granules were already present at 14 dpa in the *mrc-1* mutant,

(seen as a distinct shoulder that appeared in the granule size distribution), considerably earlier

than in the WT. Taken together, these data suggest that the larger number of granules

observed between 14-20 dpa in the *mrc-1* endosperm compared to the WT (observed in

Figure 7B) is due to the early initiation of B-type granules in the mutant.

B-type granules typically initiate in close proximity to each other– appearing as 'clusters' in

between the A-type granules – and at least some B-type granules form in amyloplast

stromules (Parker, 1985; Langeveld et al., 2000). Given the unusual timing of B-type granule

initiation in *mrc-1*, we explored whether the loss of MRC also affected the location of B-type

granule initiation. First, we harvested grains during their development (10, 15, and 20 dpa),

subjected them to critical point drying, and imaged the cut face of sections through the

endosperm using SEM. Consistent with the findings from the purified starch granules (Figure

8), B-type granules were already present at 10 dpa in the mutant, whereas they only became

prominent after 20 dpa in the WT (Figure 9A). The B-type granules occurred in clusters in

the mutant that resembled those of the WT. Secondly, we examined sections off developing

grains using light and electron microscopy. For light microscopy, sections were stained with

toluidine blue (a negative stain for starch). At 15 dpa, most starch granules in the wild-type

endosperm were flattened A-type granules, and very few B-type granules were visible

(Figure 9B). However, in the endosperm of the *mrc-1* mutant, many clusters of B-type

granules were present. Using transmission electron microscopy (TEM), we investigated

whether these clusters of B-type granules occurred within single amyloplasts, and in

particular in stromules. Indeed, multiple B-type granules were enclosed within amyloplasts,

and the elongated morphology of these amyloplast regions strongly suggested that they are

stromules (Figure 9C). It is difficult to determine the exact percentage of B-type granules in

stromules, because stromules are difficult to observe in two-dimensional sections. However,

apart from their earlier occurrence in the mutant, we did not observe anything unusual about

the location of B-type granules in the *mrc-1* mutant.

In conclusion, MRC is required for the temporal control of B-type granule initiation during

wheat grain development. It is expressed during early grain development, and its loss leads to

the early initiation of B-type granules. We therefore propose that MRC acts as a repressor of

B-type granule formation in the developing wheat endosperm during early grain

development.

TaMRC promotes granule initiation in wheat leaves

Considering the role of MRC in promoting granule initiation in Arabidopsis leaves (Seung et

al., 2018; Vandromme et al., 2019), it was surprising that MRC repressed B-type granule

formation in wheat endosperm. We therefore investigated whether this repressive role also

applies to granule initiation in wheat leaves, which would suggest a divergence in MRC

function between wheat and Arabidopsis; or whether the role of MRC in wheat leaves is the

same as in Arabidopsis but it has a distinct function in granule initiation in the endosperm.

We found that *mrc-1* and *mrc-2* had fewer granules per chloroplast than the WT (Figure 10

A, C, Table 1). As in the endosperm, the *mrc-1* mutant showed the strongest effect, having

almost 50% fewer granules per chloroplast than the WT (Table 1). Interestingly, there was no

effect of *mrc-3*, suggesting that the L289F mutation does not affect MRC function in leaves

(Figure 10 A, C, Table 1). To assess granule number, shape and size in leaf chloroplasts of

our wheat mutants, leaf tissue was harvested from the middle of the older of two leaves of 10-

day-old wheat seedlings and sections were imaged with light microscopy (Figure 10A, B).

The number of granules per chloroplast was quantified from these images. We measured

three individual plants per genotype for each experiment and compared the mean granules per

chloroplast between genotypes using a negative binomial mixed effects model with

individual biological replicates as random effect. We first compared all three *mrc* mutants

with the WT (Figure 10C, Table 1), then we compared the backcrossed *mrc-1* lines with the

WT (Figure 10D, Table 1).

The backcrossed *mrc-1* (*mrc*-1 BC2 *aabb*) had fewer granules per chloroplast than WT,

similar to the non-backcrossed *mrc-1*. However, we were not able to make meaningful

comparisons with the wild-type segregant (*mrc*-1 BC2 AABB) because – as for comparisons

of TGW and grain size above – values for the wild-type segregant differed from those of the

WT, with significantly fewer granules per chloroplast than WT. Overall, the smaller number

of granules in *mrc* chloroplasts than in WT chloroplasts suggests that MRC promotes granule

initiation in wheat leaves, pointing to distinct roles of MRC in different organs.

While the effect of the *mrc-1* mutation on starch granule number per chloroplast was

consistent, two other aspects of leaf starch – granule size and total starch content – appeared

to be interdependent and varied between our experiments. We quantified the size of granules

from these images by measuring the area of granules in the sections. The distributions

showed a trend towards larger granule sizes in the *mrc-1* mutant (Figure 10E, F). However,

after fitting a linear mixed effects model with individual biological replicates as random

effect (Table 2), pairwise comparisons showed no differences between any of the non-

backcrossed *mrc* mutants and WT. However, in the experiment comparing the backcrossed

mrc-1 mutants to WT and wild-type segregant, the granules were larger in both backcrossed

and non-backcrossed *mrc-1* mutants. Aside from the differences in size, there was no visible

difference in starch granule shape in any of the mutants compared to WT.

Similar variability between experiments was seen in total starch content, which was measured

in both leaves of 10-day-old wheat seedlings at the end of the day, when maximum starch is

expected. Our first three experiments compared the genotypes WT, *mrc-1, mrc-2* and *mrc-3*

(Experiments 1, 2, 3; Supplemental Table 4), and the pooled data showed a lower starch

content in all *mrc* mutants compared to WT (Figure 11A). Our next two experiments

(Experiments 4 and 5, Supplemental Table 4) compared end of day starch content in WT,

mrc-1, mrc-1 BC2 *aabb* and *mrc-1* BC2 AABB. By contrast, these experiments showed no

difference in starch content between *mrc-1* and WT (Figure 11B). In fact, when comparing

mrc1 BC2 *aabb* with *mrc-1* BC2 AABB or with the WT, there was a small increase in starch

content in the backcrossed mutant. The reason for the discrepancy between the results from

experiments 1 – 3 compared to experiments 4 and 5 is unknown, but it may indicate that total

starch content in wheat leaves is particularly variable and dependent on many environmental

and physiological factors.

Most interestingly, lower starch content in *mrc* mutants compared to the WT was observed in

the same batch of plants where we did not observe a significant difference in granule size

between *mrc* mutants (Experiment 2, Figure 10E), and there was identical (or greater) starch

content observed for *mrc-1* mutants in the same batch of plants where we observed larger

granules for these mutants (Experiment 5, Figure 10F). Since the reduction in granule number

described above was consistently observed in all experiments, it is plausible that the effect on
granule size depends on starch content, such that the *mrc-1* mutant only produces larger
granules under conditions where total starch content is equal. Pooled data from plants
harvested at the end of the night (quantified in experiments 1 and 2) showed very low starch
content in all three *mrc* mutants, indicating substantial nocturnal starch turnover in all
genotypes, and no differences between WT and mutants (Figure 11C).
418

Discussion
A novel role for MRC in repressing B-type granule initiation during endosperm starch
synthesis
Starch granule initiation remains the most enigmatic part of the starch synthesis process,
where we understand little about how the diverse numbers and morphologies of starch
granules are determined in our most important crops (Seung and Smith, 2019; Abt and
Zeeman, 2020; Chen et al., 2021). Here, we shed new light on the temporal regulation of
granule initiation in wheat endosperm, by demonstrating the unique role of the protein MRC
in the repression of B-type granule initiation during early grain development.
428

The expression pattern of MRC during grain development is consistent with the change in
onset of B-type granule initiation observed in the mutant. MRC is expressed in the
endosperm between 6–10 dpa but decreases rapidly in expression between 10–15 dpa,
remaining low after 15 dpa (Figure 2). This decline in expression coincides with when B-type
granules start to form in the WT (between 14 –20 dpa; Figures 7 - 9). In the *mrc-1* mutant, B-
type granules were initiated earlier than in the WT, already at 10 dpa (Figures 8 and 9), which
could be due to the loss of B-type granule repression by MRC in early grain development.
There was an increased number of starch granules in the *mrc-1* mutant compared to WT from
14 to 30 dpa (Figure 7), and this increase remained apparent in the mature grains of the
backcrossed *mrc-1* mutant (Figure 4).
439

We propose that the early initiation of B-type granules results in drastically altered starch
granule size distributions in the endosperm. The early appearance of B-type granules in the
mrc-1 mutant could present competition with A-type granules for substrates for granule
growth (i.e., ADP-glucose) from an earlier stage of grain development than in the WT,
resulting in a higher volume of B-type granules and lower volume of A-type granules in the

mature mutant grains (Figure 5). At 8 dpa, before the appearance of B-type granules, the size

distribution curves of A-type granules were almost identical between mutant and WT (Figure

8), and it was only at the later stages of grain development following B-type granule initiation

that the A-type granules became smaller in the mutant compared to WT. The lack of

difference in A-type granule number and size at 8 dpa suggests that the *mrc-1* mutation did

not affect A-type granule initiation.

Our data show that the early B-type granule initiation alone could result in the increased

proportion of B-type granule volume in mature grains, but it is difficult to make conclusions

about the effect of MRC on B-type granule number specifically. The higher B-type granule

volume could be caused by a combination of smaller A-type granules and larger and perhaps

relatively more numerous B-type granules (Figure 12). We measured a decrease in A-type

granule size and an increase in B-type granule size in our *mrc-1* and *mrc-2* mutants (Figure

5), and in some lines there was also a slight increase in total granule number (Figure 4B).

However, it should be noted that there is currently no method for specifically quantifying the

number of B-type granules, considering our definition of B-type granules comes from curve

fitting volume/size distribution graphs from the Coulter counter, and small differences in B-

type granule number may be difficult to detect as the B-type granules already make up a

much larger proportion of the total granule number in WT. For *mrc-3,* a higher B-type

granule volume percentage was observed (Figure 5B) without a measurable increase in B-

type granule size or in total granule number (Figure 4B), but it is still possible that there is a

relative increase in B-type compared to A-type granule number.

Despite all three *mrc* mutants having an increased proportion of B-type granule volume, they

differed in severity (Figure 5). The strongest effect on B-type granule percentage was seen in

mrc-1, and since the phenotypes of the *mrc-1* backcrossed and non-backcrossed lines were

similar, the severity of this line is not due to background mutations. The *mrc-1* line may have

the strongest phenotype because the premature stop codon occurs earlier in the coding

sequence than in *mrc-2*, and it is possible that the truncated protein in *mrc-2* is partially

functional. The *mrc-3* mutant had the weakest phenotypes, suggesting the Leu289Phe

mutation might also produce a partially functional protein.

The *mrc-1* wild-type segregant likely has some background mutations, as several of its

phenotypes differed from the WT. The presence of background mutations is not unusual in

these EMS-mutagenised wheat TILLING lines (Uauy et al., 2017), and importantly, the
effects observed in the wild-type segregant were minor compared to the effects observed in
the mutants.
482

MRC has tissue-specific roles in promoting or repressing granule initiation
Since wheat produces starch in both leaves and endosperm, the effects of *mrc* mutations
could be explored in both organs. We discovered that MRC has contrasting functions in the
endosperm compared to the leaves. Rather than repressing granule initiation, MRC promotes
granule initiation in the leaves, as in Arabidopsis leaves, although the mutant phenotype in
wheat appears to be less severe than in Arabidopsis. In Arabidopsis *mrc* mutants, there is one
large granule per chloroplast rather than the multiple smaller granules in wild-type
chloroplasts (Seung et al., 2018; Vandromme et al., 2019). Wheat *mrc* chloroplasts have
fewer granules per chloroplast, but there are still multiple granules in each chloroplast (Figure
10). This difference in phenotype severity is similar to those observed for the orthologs of
other granule initiation proteins of Arabidopsis (BGC1 and SS4) in wheat (Hawkins et al.,
2021; Watson-Lazowski et al., 2022). The *mrc-1* and *mrc-2* mutants both have fewer granules
per chloroplast compared to WT, with *mrc-1* having the strongest effect, consistent with the
endosperm phenotype.
497

The primary effect of the *mrc* mutations in leaves was the reduction in granule number per
chloroplast, but we also saw varying effects on granule size. In one set of experiments, we
detected larger granules in *mrc-1* compared to WT, although in another set we did not (Figure
10). This appears to be linked to whether a change in total leaf starch content was observed or
not, since in both experiments, there was a reduced number of starch granules in the mutants.
In the experiment where there was no difference in total starch at the end of the day, the
individual (fewer) granules were larger in the mutant, as would be expected. However, in the
experiment where the total starch content was lower in the *mrc-1* mutant compared to WT,
the increase in granule size was not observed.
507

508 In Arabidopsis, lack of MRC does not reduce the total end of day leaf starch content; in fact,
509 this is slightly higher in the mutant compared to the WT (Seung et al., 2018). Overall, in
510 wheat leaves the effect of *mrc* mutations on the total end of day starch content was variable in
511 our experiments (Figure 11). This could reflect differences in the regulation of starch
512 turnover in wheat vs. Arabidopsis leaves, since soluble sugars make up the majority of the

wheat leaf carbohydrate content and are thought to be more important as a storage reserve than starch (Nie et al., 1995; Müller et al., 2018; Watson-Lazowski et al., 2022). These differences may also contribute to the less severe effect on granule number in wheat *mrc* mutants compared to Arabidopsis leaves. The effect of MRC on leaf starch granule initiation is likely to differ depending on the developmental stage, so future studies should focus on the biochemistry of starch granule initiation across leaf and plant development to reveal a complete picture of the role of MRC and other granule initiation proteins in the wheat chloroplast. In Arabidopsis, reduced *At*SS4 abundance has a much stronger effect on granule formation in younger leaves compared to mature leaves (Crumpton-Taylor et al., 2013).

Exploring the biochemical basis of the contrasting roles of MRC between organs
Since MRC is a long coiled-coil protein with no known enzymatic domains, it is possible that it can exert opposite effects on granule initiation by interacting with different interaction partners. The Arabidopsis *At*MRC can interact directly with *At*SS4 in a yeast two hybrid experiment (Vandromme et al., 2019), and in Arabidopsis leaves *At*MRC was also pulled down in association with *At*PTST2, the ortholog of wheat BGC1 (Seung et al., 2018). Therefore, it is possible that in wheat, protein-protein interactions also play an important role in the function of MRC, although the nature of the interactions may be quite different. Recent work has begun to uncover the distinct roles of Arabidopsis starch granule initiation orthologs in wheat, and as for MRC, they are all quite different from their role in Arabidopsis. In the endosperm, SS4 restricts granule initiation during early grain development to ensure proper A-type granule formation and mutants defective in SS4 form compound granules (Hawkins et al., 2021). Similar, but less severe, is the *bgc1* mutant in wheat which also forms irregular compound-like granules but has a higher proportion of normal-looking A-type granules than *ss4* mutants (Chia et al., 2020; Hawkins et al., 2021). BGC1 not only represses granule initiation during A-type granule formation, but also promotes B-type granule formation, as mutants with reduced gene dosage of BGC1 have fewer B-type granules than WT, with no apparent impact on A-type granules (Chia et al., 2017; Chia et al., 2020).

However, a B-type granule suppressing mechanism during early grain development, like we report here for MRC, has not previously been described, and suggests that MRC and BGC1 might have opposing roles in the wheat endosperm, despite their similar initiation promoting roles seen in leaves of Arabidopsis (Seung et al., 2017; Seung et al., 2018) and wheat

(Watson-Lazowski et al., 2022). In support of this, our recent RNA-sequencing analysis of

the developing wheat endosperm shows opposite expression patterns for MRC and BGC1

throughout grain development (Chen et al., 2022). Wheat SS4 also promotes leaf granule

initiation, as mutants had mostly starchless chloroplasts in the leaves (Hawkins et al., 2021).

Together with our finding for MRC, this suggests that there may be more distinct differences

for granule initiation proteins between different organs of the same species in comparison to

the same organs from different species. The Arabidopsis granule initiation proteins localise to

distinct puncta in the chloroplast, but we do not yet know anything about the localisation of

these wheat orthologs in the chloroplast or amyloplast. In the future, exploring the protein

localisation, protein-protein interactions, and the connection between the two in both tissues

could reveal what underpins these distinct functions of MRC.

The specific effect of the L289F mutation in the *mrc-3* mutant on endosperm starch may

provide an important clue in these studies, since *mrc-3* had increased B-type granule

percentage in the endosperm, but no differences in granule number in leaves. Determining

whether the L289F mutation affects protein conformation, interactions and localisation might

help dissect the differences in MRC function in leaves versus endosperm.

MRC as a gene target for biotechnological modification of starch granule size

There is significant industrial interest in manipulating starch granule size in crop species, as

granule size affects the physico-chemical properties of starch and digestibility (Jobling, 2004;

Lindeboom et al., 2004; Chen et al., 2021; Li et al., 2021). Our results establish MRC as a

promising gene target for modifying starch granule size distribution in wheat, specifically to

achieve smaller starch granules and a narrower granule size distribution range than

conventional cultivars. Small granules are more efficiently digested *in vitro* than large

granules, due to their larger surface area to volume ratio (Dhital et al., 2010). Possible uses

for wheat *mrc* starch within the food industry include pasta making, where more B-type

granules positively affect pasta quality due to their higher rate of water absorption (Soh et al.,

2006). Wheat *mrc* starch may also be useful in industrial applications like papermaking and

biodegradable plastics, where small granules are desirable (Lindeboom et al., 2004).

Functional tests can be directly performed on our material to provide proof of concept that

the altered granule size distribution in the *mrc* mutants improves grain/starch quality.

MRC is a promising target because different granule size can be achieved without

accompanying effects on overall plant growth (Figure 3), on grain weight and total starch

content (Figure 4), on starch granule shape (Figure 6), or on amylopectin structure and

amylose content (Supplemental Figure 3). It is also ideal that the B-genome homeolog has

become a pseudogene (Figure 1), meaning that only one or two homeologs need to be

mutated in durum and bread wheat respectively to achieve an effect on granule size. Further,

we have also demonstrated that different mutations in MRC can be used to fine-tune the

volume of B-type granules in endosperm starch, such as those in *mrc-2* and *mrc-3* to achieve

moderate increases, and *mrc-1* to achieve larger increases. Along this line, we are currently

investigating whether overexpression of MRC can be used to reduce the volume of B-type

granules.

Since the repression of B-type granule initiation is likely to be a role specific to Triticeae

species carrying a bimodal size distribution of endosperm starch, it remains to be determined

what the role of MRC is in cereal species that do not have a bimodal distribution of starch

granules, such as those that have compound granules (e.g. in rice). Also, oats have a bimodal

distribution of starch granules, with large compound granules and smaller simple granules.

However, in oat, the smaller granules initiate at the same time as the larger compound

granules (Saccomanno et al., 2017), and it would therefore be interesting to determine if

differences in MRC function play a role in timing the initiation of the small granules during

oat endosperm development. Exploring the role of MRC in multiple crop species would

therefore not only reveal the molecular differences that result in distinct spatiotemporal

patterns of granule initiation between species, but also potentially increase its

biotechnological potential.

Materials and Methods

TaMRC bioinformatic analysis

To characterize *TtMRC-B1* in tetraploid wheat, we aligned the whole genome-sequencing

reads of *Triticum dicoccoides* (n = 10) and *Triticum turgidum ssp. durum* (n = 10) from Zhou

et al., 2020 against the 'tetraploid' version of the Ref-Seqv1.0 Chinese Spring assembly

(Appels et al., 2018). We used HiSat2-v-2.1.0 (Kim et al., 2019) with the default settings and

visualized the read alignments on the genetic signatures of retrotransposon insertion on *MRC-*

B1 (Figure 1B) using Integrated Genomics Viewer (Robinson et al., 2011). Phylogenetic

analyses were performed as described in Seung et al., (2018).

Plant materials and growth

EMS mutants of tetraploid wheat (*Triticum turgidum* cv. Kronos) carrying mutations in

TtMRC-A1 and the chromosome 6B pseudogene were identified from the wheat *in silico*

TILLING database (http://www.wheat-tilling.com; Krasileva et al., 2017) and obtained from

the John Innes Centre Germplasm Resource Unit. The selected mutants for *TtMRC-A1* were

Kronos3272(K3272), Kronos598(K598) and Kronos4681(K4681); while

Kronos4305(K4305) and Kronos3078(K3078) were selected for the 6B pseudogene. From

these mutants, we generated three different sets of lines. The *mrc-1* lines descend from a

cross between K3272 and K3078, while the *mrc-2* lines descend from a cross between K4681

and K4305. For both crosses, *aa*BB, AA*bb* and *aabb* genotypes were obtained in the F2

generation. The *mrc-3* lines are the original K598 mutants. The KASP markers used to

genotype the mutations are provided in Supplemental Table 3.

For experiments on grains and leaves, plants were grown in soil in a controlled environment

room with fluorescent lamps supplemented with LED panels. The chambers were set to

provide a 16-h light at 300 µmol photons m^{-2} s^{-1} and 20°C , and 8-h dark at 16°C, with

relative humidity of 60%. Grains from the first three tillers were harvested from mature, dry

spikes (approximately 4 months after sowing). Leaves were harvested 10 days after

germination, when two leaves were present. The two leaves were pooled for starch

quantification, and a section from the middle of the older leaf was used for light microscopy.

Grain morphometrics

The number of grains harvested per plant, plus grain area and thousand grain weight were

quantified using the MARViN seed analyser (Marvitech GmbH, Wittenburg). Multiple grains

from each plant (15 - 88 individual grains per plant) were measured for grain area to calculate

the average value for each plant, and these values were used in the plots of Figure 3F and the

analysis of Supplemental Table 2D.

Starch purification from mature grains or developing endosperm

Starch was purified from mature grains using 3-6 grains per extraction. Dry grains were

soaked overnight at 4°C in 5 mL of sterile water. The softened grains were homogenised in

10 mL sterile water using a mortar and pestle, and the homogenate was filtered through a 100

µm mesh. The starch was pelleted by centrifugation at 3,000*g* for 5 minutes, and resuspended

in 2 mL of water. The resuspended starch was loaded on top of a 5 mL 90% Percoll (Sigma)

cushion buffered with 50 mM Tris-HCl, pH 8, and was spun at 2,500*g* for 15 minutes. We

verified that no intact granules were left in the Percoll interface after the spin. The starch

pellet was washed twice with wash buffer (50 mM Tris-HCl, pH 6.8; 10 mM EDTA; 4%

SDS; and 10 mM DTT), then three times with water, followed by a final wash in absolute

ethanol. The starch was then air dried overnight.

For starch extraction from developing endosperm, developing grains were harvested at the

indicated timepoints and were snap frozen in liquid nitrogen and stored at -80°C until

analysis. Each grain was thawed just prior to extraction and the endosperm was carefully

dissected and placed into a chilled tube and weighed. The tissue was then homogenised in

sterile water with a pestle, then filtered through a 60 µm mesh. The pellet was washed three

times in 90% Percoll (Sigma) buffered with 50 mM Tris-HCl, pH 8, then three times with

wash buffer (as above), followed by three times with water.

Coulter counter analysis of starch granule size and number

For profiles of granule size distribution, purified starch was suspended in Isoton II diluent

(Beckman Coulter) and analysed with a Multisizer 4e Coulter counter fitted with a 70 µm

aperture (Beckman Coulter). Granules were counted either in volumetric mode, measuring 1

mL from a total 100 mL volume preparation containing 20 µL of purified starch or set to

count at least 100,000 granules. For calculations of granule counts in volumetric mode, the

number of granules per mg grain weight was back calculated to the total starting grain

weight. The granules were sized, with the Coulter counter collecting the data using

logarithmic bins for the granule diameter (standard settings).

For each plant, to calculate the mean A- and B-type granule size, as well as relative B-type

granule volume, we fitted a mixed bimodal gaussian curve to the distribution using R

(https://github.com/JIC-CSB/coulter_counter_fitting). As the data collection on the Coulter

counter is set to logarithmic bins on the x-axis, for these calculations and for the traces in

Figures 5A, 8A, and Supplemental Figure 3, we transformed the x-axis to even bins, by

changing the y-axis to volume percentage density (volume percentage for each bin divided by

bin width). For each of the extracted phenotypes of mean A- and B-type granule size and

relative B-type granule volume, we fitted individual linear models and performed a one-way

ANOVA and Tukey post-hoc tests for pairwise comparisons of the genotypes, using the lm()

and emmeans() functions in R, from the 'stats' and 'emmeans' packages. Nine individual

plants per genotype were used for this experiment.

Light and electron microscopy

For light microscopy of starch granules in leaf chloroplasts, leaf material (1.5 mm x 1.5 mm

squares) was harvested and fixed in 2.5% glutaraldehyde in 0.05 M sodium cacodylate, pH

7.4, which was vacuum infiltrated into the leaf. The segments were post-fixed with osmium,

then dehydrated in an ascending ethanol series, and embedded in LR white resin using EM

TP embedding machine (Leica). Semi-thin sections (0.5 μm thick) were produced from the

embedded leaves using a glass knife and were dried onto PTFE-coated slides. Chloroplasts

and cell walls were stained using toluidine blue stain (0.5% toluidine blue 'O', 0.5% sodium

borate) for 1 min. Starch was stained using reagents from the Periodic Acid-Schiff staining

kit (Abcam): using a 30 min incubation with periodic acid solution, followed by 5 min with

Schiff's solution. Then, chloroplasts and cell walls were again stained using toluidine blue

stain for 1 min. The sections were mounted with Histomount (National Diagnostics) and

imaged on a DM6000 microscope with 63X oil immersion lens (Leica).

For light microscopy of endosperm sections from mature grains, thin sections (1 μm thick) of

mature grains were made using a microtome fitted with a glass knife. Sections were mounted

onto a glass slide and stained with 3% Lugol's iodine solution (Sigma) prior to imaging on a

DM6000 (Leica) or AxioObserver Z1 (Zeiss) microscope.

For light/electron microscopy of developing endosperm tissue, developing grains (15 dpa)

were harvested into 4% paraformaldehyde, 2.5% glutaraldehyde in 0.05 M sodium

cacodylate, pH 7.4. The osmium post-fixation, dehydration and embedding into LR white

resin was done as described above for leaves. For light microscopy, semi-thin sections were

stained with toluidine blue and imaged as described above.

For transmission electron microscopy (TEM), ultra-thin sections (approximately 90 nm) were

produced from the embedded grains by sectioning with a diamond knife using a Leica UC7

ultramicrotome (Leica, Milton Keynes). The sections were picked up on 200 mesh copper

grids which had been formvar and carbon coated, then stained with 2% (w/v) uranyl acetate

for 1hr and 1% (w/v) lead citrate for 1 minute, washed in distilled water and air dried. The

grids were viewed in a FEI Talos 200C transmission electron microscope (FEI UK Ltd,

Cambridge, UK) at 200kV and imaged using a Gatan OneView 4K x 4K digital camera

(Gatan, Cambridge, UK) to record DM4 files.

For scanning electron microscopy (SEM): For imaging starch granules, a drop of purified

starch suspended in water (5 mg/mL) was air-dried onto a glass coverslip attached onto an

SEM stub. For imaging sections through developing endosperm, harvested grains were fixed

in 2.5% glutaraldehyde in 0.05 M sodium cacodylate, pH 7.4. The fixative was removed by

washing with 0.05 M sodium cacodylate, pH 7.4, after which the grains were dehydrated in

an ascending ethanol series, and then subjected to critical point drying in a CPD300

instrument (Leica) according to the manufacturer's instructions. Thick transverse sections

were produced from the dried grains and were glued onto SEM stubs. All stubs were sputter

coated with gold and observed using either a Supra 55 VPFEG (Zeiss) or Nova NanoSEM

450 (FEI) SEM instrument.

Analysis of leaf light microscopy images

For quantification of the number of granules per chloroplast, light microscopy images from

three individual plants of each of experiments 2 and 5 were used (Supplemental Table 4).

Using the cell counter plug-in in ImageJ, the number of granules per chloroplast were

counted manually, until 200 – 250 chloroplasts were reached. Cells were chosen across the

section, where most of the granules were stained well and clearly visible, and cells directly

next to vasculature were avoided. It should be noted that these are counts of granule sections,

rather than the actual numbers of granules per chloroplast, as the actual number cannot be

determined from two-dimensional sections. As much as possible, all chloroplasts in a cell

were counted to avoid bias, although occasionally a few chloroplasts where the granules were

unclear had to be omitted. Both lighter and darker stained granules were counted, but

occasionally light-staining patches that were difficult to discern as granules were observed in

all samples, and these were not included in the count.

For quantification of the granule size, we analysed the same samples as for the granules per

chloroplast, using the 'analyse particles' function in ImageJ, and manually adjusted the

thresholding in an image until the dark granules in the images were correctly separated. This

was done for several images across the section, until 100 – 400 granules were reached for

each plant. As much as possible, we removed instances where the thresholding had fused

multiple granules in proximity by checking obvious outliers against the original images.

For the statistics of the quantification of the number of granules per chloroplast, we consulted

the Statistical Services Centre Ltd (Reading, UK), who wrote the statistical analysis script in

R for us, available at

[https://github.com/Jiaawen/2022_MRC_wheat_Rscripts/tree/main/Fig10]. We used a

negative binomial mixed effects model, since we were analysing count data and wanted to

account for both the random effect of biological replicate and the frequency distribution of

number of granules per chloroplast. We used the mixed_model() function with

'family=negative.binomial' from the 'GLMMadaptive' package, and the emmeans() function

from the 'emmeans' package for pairwise comparisons in R. Our data had overdispersion

when fitting a Poisson model but not when using a negative binomial. We used individual

plants as the random effect (for both experiments: 3 plants for each of 4 genotypes – 12

plants total), and genotype as fixed effect.

For the statistics of the granule area, we used a linear mixed effects model, also with the

individual plants as random effect and genotype as fixed effect. We used the lmer() function

from the 'lmerTest' package, and the emmeans() function from the 'emmeans' package for

pairwise comparisons.

Quantification of starch content in leaves and endosperm

Starch was quantified in leaf tissue as previously described (Smith and Zeeman, 2006).

Briefly, frozen leaf tissue was ground into a powder with a ball mill and then extracted with

perchloric acid. Starch in the insoluble fraction of the extraction was gelatinised at 95°C and

digested to glucose with α-amylase (Megazyme) and amyloglucosidase (Roche). The glucose

released was measured using the hexokinase/glucose-6-phosphate dehydrogenase method

(Roche). Starch content (in glucose equivalents) was calculated relative to the original dry

weight of the analysed grains.

Five individual experiments were done to quantify the total end of day leaf starch content

(Supplemental Table 4). One set of genotypes was measured in experiments 1 – 3: WT, *mrc-*

1, mrc-2, mrc-3. Another set was measured in experiments 4 and 5: WT, *mrc-1,* the double

backcrossed *mrc-1* BC2 *aabb* and the wild-type segregant from that backcross, *mrc-1* BC2

AABB. Therefore, in the statistical analysis experiments 1, 2, 3 were pooled together and

experiments 4 and 5 were pooled together. An ANOVA using a fixed effects model showed

no interaction between the experiment effect and genotype effect, and experiment and

genotype were set as fixed effects in our linear model. We fitted individual linear models for

end of day starch in experiments 1 – 3, end of day starch in experiments 4 & 5 and end of

night starch in experiments 1 & 2. Then we performed ANOVA and Tukey post-hoc tests

using these linear models for pairwise comparisons of the genotypes. For the details of

number of replicates, see Supplemental Table 4. We used the lm() function from the 'stats'

package in R for the linear models, and used the emmeans() function from the 'emmeans'

package for calculating adjusted means and pairwise comparisons.

A similar method to leaves was used for starch quantification in grains. Mature grains (5-6

grains) were soaked overnight at 4°C in 5 mL of sterile water and were homogenised using a

mortar and pestle. Developing endosperm tissue was extracted in 1 mL of sterile water with

the pestle. Insoluble material in an aliquot of the homogenate was collected by centrifugation

at 5,000g for 5 mins, then washed once in 0.7 M perchloric acid, once in sterile water, then

three times in 80% ethanol. The pellet was then resuspended in water. Starch in the pellet was

gelatinised by heating at 95°C for 15 min, then digested using α-amylase (Megazyme) and

amyloglucosidase (Roche).

Analysis of amylopectin structure and amylose content

Amylopectin structure and amylose content were analysed using purified starch. Amylopectin

structure in terms of chain length distribution was quantified using High Performance Anion

Exchange Chromatography with Pulsed Amperometric Detection (HPAEC-PAD) (Blennow

et al., 1998). For amylose content, granules were dispersed in DMSO and quantified using an

iodine-binding method (Warren et al., 2016).

Statistics

All statistical analyses were done in R version 4. Overall, we used the emmeans() function

from the 'emmeans' package throughout for pairwise comparisons. For most experiments, we

used linear models using the lm() function from the 'stats' package and did one-way

ANOVAs with Tukey post-hoc tests. Details for individual experiments where we used other

models with more than a single fixed effect are described above for the relevant sections.

Tables 1, 2 and Supplemental Table 2 also describe details of statistics for individual
experiments.
817
818 *Accession numbers*
819 The accession numbers corresponding to the genes investigated in this study are*: Tt*MRC-A1
820 (TRITD6Av1G081580)*, Ta*MRC-A1 (TraesCS6A02G180500)*, Ta*MRC-D1
821 (TraesCS6D02G164600).
822
823

Acknowledgements
The authors thank the John Innes Centre (JIC) Horticultural Services for providing growth
facilities and maintenance of plant material, and JIC Bioimaging for providing access to
microscopes. We thank Prof. Alison Smith for critically reading this manuscript. This work
was funded through a John Innes Foundation (JIF) Chris J. Leaver Fellowship (to D.S), a
Biotechnology and Biological Sciences Research Council (BBSRC, UK) Future Leader
Fellowship BB/P010814/1 (to D.S.), a JIF Rotation Ph.D. studentship (to J.C. and Y.C.) and
BBSRC Institute Strategic Programme grants BBS/E/J/000PR9790, BBS/E/J/000PR9799 and
BB/P016855/1 (to the John Innes Centre).

833
834

835 **Author contributions**
836 J.C. and D.S. conceived the study. J.C., Y.C., A.W-L., F.J.W., A.B., C.U., and D.S. designed
837 the research. J.C., Y.C., A.W-L., E.H., J.E.B., B.F., R.D.B., K.C., F.J.W., A.B., C.U., and
838 D.S. performed the research and analysed data; J.C. and D.S. wrote the article with input
839 from all authors.

840
841

References

Abt, M.R., and Zeeman, S.C. (2020). Evolutionary innovations in starch metabolism. Current Opinion in Plant Biology **55,** 109-117.

Appels, R., Eversole, K., Stein, N., Feuillet, C., Keller, B., Rogers, J., Pozniak, C.J., Choulet, F., Distelfeld, A., Poland, J., Ronen, G., Sharpe, A.G., Barad, O., Baruch, K., Keeble-Gagnère, G., Mascher, M., Ben-Zvi, G., Josselin, A.-A., Himmelbach, A., Balfourier, F., Gutierrez-Gonzalez, J., Hayden, M., Koh, C., Muehlbauer, G., Pasam, R.K., Paux, E., Rigault, P., Tibbits, J., Tiwari, V., Spannagl, M., Lang, D., Gundlach, H., Haberer, G., Mayer, K.F.X., Ormanbekova, D., Prade, V., Šimková, H., Wicker, T., Swarbreck, D., Rimbert, H., Felder, M., Guilhot, N., Kaithakottil, G., Keilwagen, J., Leroy, P., Lux, T., Twardziok, S., Venturini, L., Juhász, A., Abrouk, M., Fischer, I., Uauy, C., Borrill, P., Ramirez-Gonzalez, R.H., Arnaud, D., Chalabi, S., Chalhoub, B., Cory, A., Datla, R., Davey, M.W., Jacobs, J., Robinson, S.J., Steuernagel, B., van Ex, F., Wulff, B.B.H., Benhamed, M., Bendahmane, A., Concia, L., Latrasse, D., Bartoš, J., Bellec, A., Berges, H., Doležel, J., Frenkel, Z., Gill, B., Korol, A., Letellier, T., Olsen, O.-A., Singh, K., Valárik, M., van der Vossen, E., Vautrin, S., Weining, S., Fahima, T., Glikson, V., Raats, D., Číhalíková, J., Toegelová, H., Vrána, J., Sourdille, P., Darrier, B., Barabaschi, D., Cattivelli, L., Hernandez, P., Galvez, S., Budak, H., Jones, J.D.G., Witek, K., Yu, G., Small, I., Melonek, J., Zhou, R., Belova, T., Kanyuka, K., King, R., Nilsen, K., Walkowiak, S., Cuthbert, R., Knox, R., Wiebe, K., Xiang, D., Rohde, A., Golds, T., Čížková, J., Akpinar, B.A., Biyiklioglu, S., Gao, L., N'Daiye, A., Kubaláková, M., Šafář, J., Alfama, F., Adam-Blondon, A.-F., Flores, R., Guerche, C., Loaec, M., Quesneville, H., Condie, J., Ens, J., Maclachlan, R., Tan, Y., Alberti, A., Aury, J.-M., Barbe, V., Couloux, A., Cruaud, C., Labadie, K., Mangenot, S., Wincker, P., Kaur, G., Luo, M., Sehgal, S., Chhuneja, P., Gupta, O.P., Jindal, S., Kaur, P., Malik, P., Sharma, P., Yadav, B., Singh, N.K., Khurana, J.P., Chaudhary, C., Khurana, P., Kumar, V., Mahato, A., Mathur, S., Sevanthi, A., Sharma, N., Tomar, R.S., Holušová, K., Plíhal, O., Clark, M.D., Heavens, D., Kettleborough, G., Wright, J., Balcárková, B., Hu, Y., Salina, E., Ravin, N., Skryabin, K., Beletsky, A., Kadnikov, V., Mardanov, A., Nesterov, M., Rakitin, A., Sergeeva, E., Handa, H., Kanamori, H., Katagiri, S., Kobayashi, F., Nasuda, S., Tanaka, T., Wu, J., Cattonaro, F., Jiumeng, M., Kugler, K., Pfeifer, M., Sandve, S., Xun, X., Zhan, B., Batley, J., Bayer, P.E., Edwards, D., Hayashi, S., Tulpová, Z., Visendi, P., Cui, L., Du, X., Feng, K., Nie, X., Tong, W., and Wang, L. (2018). Shifting the limits in wheat research and breeding using a fully annotated reference genome. Science **361,** eaar7191.

Bechtel, D.B., Zayas, I., Kaleikau, L., and Pomeranz, Y. (1990). Size-distribution of wheat starch granules during endosperm development. Cereal Chemistry **67,** 59-63.

Blennow, A., Bay-Smidt, A.M., Wischmann, B., Olsen, C.E., and Møller, B.L. (1998). The degree of starch phosphorylation is related to the chain length distribution of the neutral and the phosphorylated chains of amylopectin. Carbohydrate Research **307,** 45-54.

Borrill, P., Ramirez-Gonzalez, R., and Uauy, C. (2016). expVIP: a customizable RNA-seq data analysis and visualization platform. Plant Physiology **170,** 2172-2186.

Chen, J., Hawkins, E., and Seung, D. (2021). Towards targeted starch modification in plants. Current Opinion in Plant Biology **60,** 102013.

Chen, J., Watson-Lazowski, A., Vickers, M., and Seung, D. (2022). Gene expression profile of the developing endosperm in durum wheat provides insight into starch biosynthesis. BioRxiv (https://doi.org/10.1101/2022.10.21.513215).

Chia, T., Adamski, N.M., Saccomanno, B., Greenland, A., Nash, A., Uauy, C., and Trafford, K. (2017). Transfer of a starch phenotype from wild wheat to bread wheat by deletion of a locus controlling B-type starch granule content. Journal of Experimental Botany **68,** 5497-5509.

Chia, T., Chirico, M., King, R., Ramirez-Gonzalez, R., Saccomanno, B., Seung, D., Simmonds, J., Trick, M., Uauy, C., Verhoeven, T., and Trafford, K. (2020). A carbohydrate-binding protein, B-GRANULE CONTENT 1, influences starch granule size distribution in a dose-dependent manner in polyploid wheat. Journal of Experimental Botany **71,** 105-115.

Crumpton-Taylor, M., Pike, M., Lu, K.J., Hylton, C.M., Feil, R., Eicke, S., Lunn, J.E., Zeeman, S.C., and Smith, A.M. (2013). Starch synthase 4 is essential for coordination of starch granule formation with chloroplast division during Arabidopsis leaf expansion. New Phytologist **200,** 1064-1075.

Daron, J., Glover, N., Pingault, L., Theil, S., Jamilloux, V., Paux, E., Barbe, V., Mangenot, S., Alberti, A., Wincker, P., Quesneville, H., Feuillet, C., and Choulet, F. (2014). Organization and evolution of transposable elements along the bread wheat chromosome 3B. Genome Biol **15,** 546.

Dhital, S., Shrestha, A.K., and Gidley, M.J. (2010). Relationship between granule size and in vitro digestibility of maize and potato starches. Carbohydrate Polymers **82,** 480-488.

Hawkins, E., Chen, J., Watson-Lazowski, A., Ahn-Jarvis, J., Barclay, J.E., Fahy, B., Hartley, M., Warren, F.J., and Seung, D. (2021). STARCH SYNTHASE 4 is required for normal starch granule initiation in amyloplasts of wheat endosperm. New Phytologist **230,** 2371-2386.

Howard, T., Rejab, N.A., Griffiths, S., Leigh, F., Leverington-Waite, M., Simmonds, J., Uauy, C., and Trafford, K. (2011). Identification of a major QTL controlling the content of B-type starch granules in Aegilops. Journal of Experimental Botany **62,** 2217-2228.

Jobling, S. (2004). Improving starch for food and industrial applications. Current Opinion in Plant Biology **7,** 210-218.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol **37,** 907-915.

Krasileva, K.V., Vasquez-Gross, H.A., Howell, T., Bailey, P., Paraiso, F., Clissold, L., Simmonds, J., Ramirez-Gonzalez, R.H., Wang, X., Borrill, P., Fosker, C., Ayling, S., Phillips, A.L., Uauy, C., and Dubcovsky, J. (2017). Uncovering hidden variation in polyploid wheat. Proceedings of the National Academy of Sciences **114,** E913-E921.

Langeveld, S.M.J., Van wijk, R., Stuurman, N., Kijne, J.W., and de Pater, S. (2000). B-type granule containing protrusions and interconnections between amyloplasts in developing wheat endosperm revealed by transmission electron microscopy and GFP expression. Journal of Experimental Botany **51,** 1357-1361.

Li, L.F., Zhang, Z.B., Wang, Z.H., Li, N., Sha, Y., Wang, X.F., Ding, N., Li, Y., Zhao, J., Wu, Y., Gong, L., Mafessoni, F., Levy, A.A., and Liu, B. (2022). Genome sequences of five Sitopsis species of Aegilops and the origin of polyploid wheat B subgenome. Mol Plant **15,** 488-503.

Li, M., Daygon, V.D., Solah, V., and Dhital, S. (2021). Starch granule size: Does it matter?
Critical Reviews in Food Science and Nutrition **0,** 1-21.
Lindeboom, N., Chang, P.R., and Tyler, R.T. (2004). Analytical, biochemical and
physicochemical aspects of starch granule size, with emphasis on small granule
starches: a review. Starch - Stärke **56,** 89-99.
Maccaferri, M., Harris, N.S., Twardziok, S.O., Pasam, R.K., Gundlach, H., Spannagl,
M., Ormanbekova, D., Lux, T., Prade, V.M., Milner, S.G., Himmelbach, A.,
Mascher, M., Bagnaresi, P., Faccioli, P., Cozzi, P., Lauria, M., Lazzari, B., Stella,
A., Manconi, A., Gnocchi, M., Moscatelli, M., Avni, R., Deek, J., Biyiklioglu, S.,
Frascaroli, E., Corneti, S., Salvi, S., Sonnante, G., Desiderio, F., Marè, C.,
Crosatti, C., Mica, E., Özkan, H., Kilian, B., De Vita, P., Marone, D., Joukhadar,
R., Mazzucotelli, E., Nigro, D., Gadaleta, A., Chao, S., Faris, J.D., Melo, A.T.O.,
Pumphrey, M., Pecchioni, N., Milanesi, L., Wiebe, K., Ens, J., MacLachlan, R.P.,
Clarke, J.M., Sharpe, A.G., Koh, C.S., Liang, K.Y.H., Taylor, G.J., Knox, R.,
Budak, H., Mastrangelo, A.M., Xu, S.S., Stein, N., Hale, I., Distelfeld, A.,
Hayden, M.J., Tuberosa, R., Walkowiak, S., Mayer, K.F.X., Ceriotti, A.,
Pozniak, C.J., and Cattivelli, L. (2019). Durum wheat genome highlights past
domestication signatures and future improvement targets. Nature Genetics **51,** 885-
895.
Matsushima, R., Yamashita, J., Kariyama, S., Enomoto, T., and Sakamoto, W. (2013). A
phylogenetic re-evaluation of morphological variations of starch grains among
Poaceae species. Journal of Applied Glycoscience **60,** 131-135.
Müller, L.M., Gol, L., Jeon, J.-S., Weber, A.P.M., Davis, S.J., and von Korff, M. (2018).
Temperature but not the circadian clock determines nocturnal carbohydrate
availability for growth in cereals. bioRxiv.
Ng, P.C., and Henikoff, S. (2006). Predicting the effects of amino acid substitutions on
protein function. Annu Rev Genomics Hum Genet **7,** 61-80.
Nie, G., Hendrix, D.L., Webber, A.N., Kimball, B.A., and Long, S.P. (1995). Increased
accumulation of carbohydrates and decreased photosynthetic gene transcript levels in
wheat grown at an elevated CO_2 concentration in the field. Plant Physiol **108,** 975-
983.
Parker, M.L. (1985). The relationship between A-type and B-type starch granules in the
developing endosperm of wheat. Journal of Cereal Science **3,** 271-278.
Peng, C., Wang, Y., Liu, F., Ren, Y., Zhou, K., Lv, J., Zheng, M., Zhao, S., Zhang, L.,
Wang, C., Jiang, L., Zhang, X., Guo, X., Bao, Y., and Wan, J. (2014). FLOURY
ENDOSPERM6 encodes a CBM48 domain-containing protein involved in compound
granule formation and starch synthesis in rice endosperm. Plant Journal **77,** 917-930.
Ramirez-Gonzalez, R.H., Borrill, P., Lang, D., Harrington, S.A., Brinton, J., Venturini,
L., Davey, M., Jacobs, J., van Ex, F., Pasha, A., Khedikar, Y., Robinson, S.J.,
Cory, A.T., Florio, T., Concia, L., Juery, C., Schoonbeek, H., Steuernagel, B.,
Xiang, D., Ridout, C.J., Chalhoub, B., Mayer, K.F.X., Benhamed, M., Latrasse,
D., Bendahmane, A., International Wheat Genome Sequencing, C., Wulff,
B.B.H., Appels, R., Tiwari, V., Datla, R., Choulet, F., Pozniak, C.J., Provart,
N.J., Sharpe, A.G., Paux, E., Spannagl, M., Brautigam, A., and Uauy, C. (2018).
The transcriptional landscape of polyploid wheat. Science **361.**
Robinson, J.T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G.,
and Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol **29,** 24-26.
Roldán, I., Wattebled, F., Mercedes Lucas, M., Delvallé, D., Planchot, V., Jiménez, S.,
Pérez, R., Ball, S., D'Hulst, C., and Mérida, A. (2007). The phenotype of soluble
starch synthase IV defective mutants of Arabidopsis thaliana suggests a novel

989 function of elongation enzymes in the control of starch granule formation. Plant
990  Journal **49,** 492-504.

991 **Saccomanno, B., Chambers, A.H., Hayes, A., Mackay, I., McWilliam, S.C., and**
992  **Trafford, K.** (2017). Starch granule morphology in oat endosperm. Journal of Cereal
993  Science **73,** 46-54.

994 **Saccomanno, B., Berbezy, P., Findlay, K., Shoesmith, J., Uauy, C., Viallis, B., and**
995  **Trafford, K.** (2022). Characterization of wheat lacking B-type starch granules.
996  Journal of Cereal Science **104.**

997 **Saito, M., Tanaka, T., Sato, K., Vrinten, P., and Nakamura, T.** (2017). A single
998  nucleotide polymorphism in the "Fra" gene results in fractured starch granules in
999  barley. Theoretical and Applied Genetics **131,** 353-364.

1000 **Seung, D.** (2020). Amylose in starch: towards an understanding of biosynthesis, structure and
1001  function. New Phytologist **228,** 1490-1504.

1002 **Seung, D., and Smith, A.M.** (2019). Starch granule initiation and morphogenesis – progress
1003  in Arabidopsis and cereals. Journal of Experimental Botany **70,** 771-784.

1004 **Seung, D., Schreier, T.B., Bürgy, L., Eicke, S., and Zeeman, S.C.** (2018). Two plastidial
1005  coiled-coil proteins are essential for normal starch granule initiation in Arabidopsis.
1006  Plant Cell **30,** 1523-1542.

1007 **Seung, D., Boudet, J., Monroe, J.D., Schreier, T.B., David, L.C., Abt, M., Lu, K.-J.,**
1008  **Zanella, M., and Zeeman, S.C.** (2017). Homologs of PROTEIN TARGETING TO
1009  STARCH control starch granule initiation in Arabidopsis leaves. Plant Cell **29,** 1657-
1010  1677.

1011 **Smith, A.M., and Zeeman, S.C.** (2006). Quantification of starch in plant tissues. Nature
1012  Protocols **1,** 1342-1345.

1013 **Smith, A.M., and Zeeman, S.C.** (2020). Starch: A flexible, adaptable carbon store coupled
1014  to plant growth. Annual Review of Plant Biology **71,** 217-245.

1015 **Soh, H.N., Sissons, M.J., and Turner, M.A.** (2006). Effect of starch granule size
1016  distribution and elevated amylose content on durum dough rheology and spaghetti
1017  cooking quality. Cereal Chemistry **83,** 513-519.

1018 **Tetlow, I., and Emes, M.** (2017). Starch biosynthesis in the developing endosperms of
1019  grasses and cereals. Agronomy **7,** 81.

1020 **Uauy, C., Wulff, B.B.H., and Dubcovsky, J.** (2017). Combining traditional mutagenesis
1021  with new high-throughput sequencing and genome editing to reveal hidden variation
1022  in polyploid wheat. Annu Rev Genet **51,** 435-454.

1023 **Vandromme, C., Spriet, C., Dauvillée, D., Courseaux, A., Putaux, J.-l., Wychowski, A.,**
1024  **Krzewinski, F., Facon, M., D'Hulst, C., and Wattebled, F.** (2019). PII1: a protein
1025  involved in starch initiation that determines granule number and size in Arabidopsis
1026  chloroplast. New Phytologist **221,** 356-370.

1027 **Walkowiak, S., Gao, L., Monat, C., Haberer, G., Kassa, M.T., Brinton, J., Ramirez-**
1028  **Gonzalez, R.H., Kolodziej, M.C., Delorean, E., Thambugala, D., Klymiuk, V.,**
1029  **Byrns, B., Gundlach, H., Bandi, V., Siri, J.N., Nilsen, K., Aquino, C.,**
1030  **Himmelbach, A., Copetti, D., Ban, T., Venturini, L., Bevan, M., Clavijo, B., Koo,**
1031  **D.H., Ens, J., Wiebe, K., N'Diaye, A., Fritz, A.K., Gutwin, C., Fiebig, A., Fosker,**
1032  **C., Fu, B.X., Accinelli, G.G., Gardner, K.A., Fradgley, N., Gutierrez-Gonzalez,**
1033  **J., Halstead-Nussloch, G., Hatakeyama, M., Koh, C.S., Deek, J., Costamagna,**
1034  **A.C., Fobert, P., Heavens, D., Kanamori, H., Kawaura, K., Kobayashi, F.,**
1035  **Krasileva, K., Kuo, T., McKenzie, N., Murata, K., Nabeka, Y., Paape, T.,**
1036  **Padmarasu, S., Percival-Alwyn, L., Kagale, S., Scholz, U., Sese, J., Juliana, P.,**
1037  **Singh, R., Shimizu-Inatsugi, R., Swarbreck, D., Cockram, J., Budak, H.,**
1038  **Tameshige, T., Tanaka, T., Tsuji, H., Wright, J., Wu, J., Steuernagel, B., Small,**

I., Cloutier, S., Keeble-Gagnere, G., Muehlbauer, G., Tibbets, J., Nasuda, S.,
Melonek, J., Hucl, P.J., Sharpe, A.G., Clark, M., Legg, E., Bharti, A., Langridge,
P., Hall, A., Uauy, C., Mascher, M., Krattinger, S.G., Handa, H., Shimizu, K.K.,
Distelfeld, A., Chalmers, K., Keller, B., Mayer, K.F.X., Poland, J., Stein, N.,
McCartney, C.A., Spannagl, M., Wicker, T., and Pozniak, C.J. (2020). Multiple
wheat genomes reveal global variation in modern breeding. Nature **588,** 277-283.
Warren, F.J., Gidley, M.J., and Flanagan, B.M. (2016). Infrared spectroscopy as a tool to
characterise starch ordered structure - A joint FTIR-ATR, NMR, XRD and DSC
study. Carbohydrate Polymers **139,** 35-42.
Watson-Lazowski, A., Raven, E., Feike, D., Hill, L., Elaine Barclay, J., Smith, A.M., and
Seung, D. (2022). Loss of PROTEIN TARGETING TO STARCH 2 has variable
effects on starch synthesis across organs and species. J Exp Bot.
Zhou, Y., Zhao, X., Li, Y., Xu, J., Bi, A., Kang, L., Xu, D., Chen, H., Wang, Y., Wang,
Y.-g., Liu, S., Jiao, C., Lu, H., Wang, J., Yin, C., Jiao, Y., and Lu, F. (2020).
Triticum population sequencing provides insights into wheat adaptation. Nat Genet
52, 1412-1422.

1055

Tables

Table 1. Statistics of pairwise comparisons of leaf granule per chloroplast in young wheat leaves, from negative binomial mixed effect models with biological replicate (3 of each genotype) as random effect, using Tukey post-hoc tests. Individual models for each experiment were used. SE = standard error, 95% CI = 95% confidence interval.

contrast	Experiment	ratio	SE	- 95% CI	+ 95 CI	z.ratio	p.value
WT / *mrc-1*	2	1.997	0.190	1.564	2.550	7.271	2.17E-12
WT / *mrc-2*	2	1.390	0.131	1.091	1.769	3.497	0.00265
WT / mrc-3	2	1.091	0.102	0.858	1.387	0.929	0.789273
mrc-1 / mrc-2	2	0.696	0.067	0.544	0.890	-3.783	0.000892
mrc-1 / mrc-3	2	0.546	0.052	0.427	0.698	-6.336	1.41E-09
mrc-2 / mrc-3	2	0.785	0.074	0.616	1.000	-2.564	0.05063
mrc-1* BC2 *aabb / mrc-1	5	1.030	0.097	0.810	1.311	0.318	0.988897
mrc-1* BC2 *AABB / mrc-1	5	1.234	0.115	0.971	1.569	2.256	0.108551
mrc-1* BC2 *AABB / mrc-1* BC2 *aabb	5	1.198	0.111	0.943	1.522	1.943	0.210022
WT / *mrc-1*	5	1.635	0.152	1.287	2.077	5.285	7.52E-07
WT / *mrc-1* BC2 *aabb*	5	1.587	0.147	1.251	2.014	4.982	3.76E-06
WT / *mrc-1* BC2 *AABB*	5	1.325	0.122	1.045	1.679	3.048	0.012347

Table 2. Statistics of pairwise comparisons of granule size in young wheat leaves, from linear mixed effect models with biological replicate (3 of each genotype) as random effect, using Tukey post-hoc tests and Satterthwaite degrees of freedom calculation. Individual models for each experiment were used. SE = standard error, df = degrees of freedom, 95% CI = 95% confidence interval.

contrast	Experiment	difference	SE	- 95% CI	+ 95 CI	df	t.ratio	p.value
WT – *mrc-1*	2	-0.304	0.125	-0.706	0.0975	7.998	-2.42484	0.149207
WT – *mrc-2*	2	-0.087	0.125	-0.488	0.3150	7.979	-0.69054	0.89788
WT – *mrc-3*	2	0.045	0.125	-0.357	0.4461	7.941	0.356537	0.983352
mrc-2 – mrc-1	2	-0.218	0.126	-0.619	0.1841	8.055	-1.73159	0.368589
mrc-3 – mrc-1	2	-0.349	0.125	-0.750	0.0529	8.017	-2.77906	0.090736
mrc-3 – mrc-2	2	-0.131	0.125	-0.533	0.2704	7.998	-1.04602	0.729078
mrc-1* BC2 *aabb - mrc-1	5	-0.034	0.0394	-0.160	0.0924	7.990	-0.85905	0.825337
mrc-1* BC2 *AABB - mrc-1	5	-0.255	0.0398	-0.381	-0.1289	8.317	-6.41375	0.000799
mrc-1* BC2 *AABB - mrc-1* BC2 *aabb	5	-0.221	0.0400	-0.348	-0.0949	8.470	-5.5362	0.002041
WT - *mrc-1*	5	-0.260	0.0396	-0.386	-0.1335	8.122	-6.56687	0.000748
WT - *mrc-1* BC2 *aabb*	5	-0.226	0.0397	-0.352	-0.0996	8.274	-5.68343	0.001851
WT - *mrc-1* BC2 *AABB*	5	-0.005	0.0401	-0.131	0.1218	8.607	-0.11473	0.999416

Figure Legends

Figure 1. *MRC* homeologs in wheat are encoded on chromosomes 6A and 6D, with a disruption of the 6B homeolog. A) Location of *TaMRC* homeologs on chromosome 6A and 6D. The pink boxes represent *TaMRC* homeologs, while homeologs of the adjacent genes are shown in green (cytochrome P450 family protein), purple (respiratory burst oxidase homolog) and blue (uncharacterised protein). Arrowheads on the boxes indicate direction of transcription. The syntenic region on chromosome 6B has a large insertion, depicted with a black arrowhead. The diagram is drawn to scale, and chromosome coordinates of the region are indicated. **B)** Gene models of the *TaMRC-A1* and *–D1* homeologs and 6B pseudogene. Exons are represented with pink boxes, while light pink boxes represent the 5' and 3' UTRs. On the 6B region, areas with sequence similarity to exon 2 and UTRs of *TaMRC-A1* are indicated, as well as the location of *gypsy* retrotransposons. The locations of the mutations in the *mrc* mutants are depicted with red arrows, and the mutated codons/amino acids are shown in red letters. Large black arrowheads show where the sequence has been truncated for illustration - the length of truncated sequence is indicated above. **C)** Summary of species analysed for the loss of *MRC-B1* during wheat hybridisation

Figure 2. *MRC* is expressed in wheat leaves and wheat endosperm during early grain development. A) Expression of *TaMRC* homeologs between between different tissue types in *Triticum aestivum*. Data were obtained from the wheat expression browser, and values represent transcript per million (TPM) ± SEM from n = 89 (roots), 48 (leaves/shoots), 280 (spike), 166 (grain) samples from different experiments. **B)** Expression of *TtMRC-A1* (TRITD6Av1G081580.1) in the starchy endosperm during grain development of *Triticum turgidum* (variety Kronos). Data are from an RNAseq experiment described in Chen *et al.* (2022). Values represent TPM ± SEM from n = 3 for all timepoints.

Figure 3. Mutations in *MRC* do not affect plant growth or the number and size of grains. A) Photographs of 7-week-old wild type (WT) and *mrc* mutant plants. Bar = 9 cm. **B)** Number of tillers per plant. **C)** Number of grains per plant. **D)** Photographs of mature grains. Bar = 1 cm. **E)** Thousand Grain Weight (TGW) of mature grains. **F)** Grain size, measured as the average area of individual mature grains from each plant. For statistical analysis of B, C, E, F, see Supplemental Table 1. For all boxplots, each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. Dots are measurements from individual plants, with the same group of plants measured for all phenotypes and n = 9 – 10 individual plants.

Figure 4. Mutations in *MRC* have variable effects on total starch in mature grains. A) Total endosperm starch content in mature grains in WT and *mrc* mutants. B) Number of starch granules per mg of grain, counted with a Coulter counter. For A and B, dots on the boxplots indicate individual plants, with n = 8 - 9. Each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. All statistical analyses were performed using a linear model, with a one-way ANOVA and Tukey post-hoc test. Panels on the right indicate differences in means between genotypes from pairwise comparisons based on these models. The difference in means is indicated by a dot, with whiskers showing the 95% confidence interval (CI) of this difference, with the corresponding p-value. Grey indicates the WT or *mrc-1* mutant compared to the backcrossed line with the equivalent genotype at the *MRC* loci, and blue indicates all other pairwise comparisons.

Figure 5. Endosperm starch in mature grains of *mrc* mutants have altered granule size distribution. A) Coulter counter traces with evenly binned x-axes show a bimodal distribution of granule sizes from purified wheat endosperm starch. Data points are mean values from 9 individual plants of each genotype (3 grains from each plant), with the standard error of the mean shown as a shaded ribbon. WT and *mrc-1* in the left and right panel are the same data. **B, C, D)** Mean B-type granule percentage volume (B), A-type granule diameter (C) and B-type granule diameter (D) values extracted from bimodal log-normal distribution curves fitted to Coulter counter traces of individual plants. Extracted values and boxplots are shown on the left, where dots indicate mean values from individual plants ($n = 9$). Each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. All statistical analyses were performed using linear models for each panel, with a one-way ANOVA and Tukey post-hoc test. Panels on the right indicate differences in means between genotypes from pairwise comparisons based on these models. The difference in means is indicated by a dot, with whiskers showing the 95% confidence interval (CI) of this difference, with the corresponding p-value. Grey indicates the WT or *mrc-1* mutant compared to the backcrossed line with the equivalent genotype at the *MRC* loci, and blue indicates all other pairwise comparisons.

Figure 6. Endosperm starch granules of mature grains in three wheat *mrc* mutants are similar in shape compared to wild type. A) Thin sections of mature endosperm tissue were stained with Lugol's solution and imaged using light microscopy. Bar = 40 µm. **B)** Purified endosperm starch granules were observed using scanning electron microscopy (SEM). Bar = 20 µm.

Figure 7. In developing endosperm, granule number increases in *mrc-1* compared to WT despite having similar starch content. The endosperm was dissected from developing grains of WT and *mrc-1*, harvested at 8, 14, 20 and 30 dpa, with $n = 3 - 4$ individual plants for each genotype per time point, indicated by coloured dots in the upper panels, with the mean ± 95% confidence interval (CI) in black dots and whiskers. **A)** Starch content of the endosperm. Values are expressed relative to the fresh weight of the dissected endosperm. **B)** Starch granule number in the endosperm. Starch was purified from dissected endosperm and the number of granules was determined using a Coulter counter. Values are expressed relative to the fresh weight of the dissected endosperm. For A and B, individual linear models were fitted to the data of each time point, with a one-way ANOVA and Tukey post-hoc test to compare the means of WT and *mrc-1*. Panels on the bottom summarise the differences in means from these linear models, indicated by a dot, with whiskers showing the 95% CI of this difference, with the corresponding p-value.

Figure 8. The *mrc-1* mutant initiates B-type granules earlier in grain development than in the wild type (WT). The endosperm was dissected from developing grains of WT and *mrc-1* harvested at 8, 14, 20 and 30 dpa. **A)** Coulter counter traces with evenly binned x-axes show the starch granule size distribution of endosperm starch. Distributions are the average of $n = 4$ measurements, each carried out on grains harvested from a different plant (three grains per measurement). Data points are mean values from these 4 measurements, with the standard error of the mean shown as a shaded ribbon. The B-type granule peak in the *mrc-1* mutant at 14 dpa is indicated with a blue arrow. **B)** Endosperm starch granules were observed using scanning electron microscopy (SEM). Bars = 20 µm. Examples of A-type granules and B-type granules are marked with red and blue arrows respectively.

Figure 9. The early initiating B-type granules in *mrc-1* occur at least partially in stromules. A) Scanning electron micrographs of developing endosperm tissue subjected to critical point drying. Grains were harvested from WT and *mrc-1* at 10, 15 and 20 days post anthesis (dpa). A representative region of each panel has been pseudo-coloured, with A-type granules in red shading and B-type granules in blue shading. Bars = 50 μm. **B)** Light micrographs of endosperm sections. Semi-thin sections of embedded developing grains (15 dpa) were stained with toluidine blue as a negative stain for starch granules. Examples of A-type granules and B-type granules are marked with red and blue arrows respectively. Bars = 50 μm. **C)** Endosperm sections observed using transmission electron microscopy, using the same samples as in B. The periphery of amyloplast membranes, where visible, are indicated with a dotted white line. Bars = 1 μm.

Figure 10. Chloroplasts of *mrc-1* mutants have fewer and larger granules than the wild type. A, B) Representative light microscopy images of thin sections (500 nm) from the middle of the older of two leaves in 10-day old wheat seedlings, collected at the end of day. Sections were stained for starch using periodic acid and Schiff staining. Scale bar = 10 μm. **C, D)** Distributions of counts of granule sections per chloroplast section, counted from light microscopy images as in A and B. The three different colours correspond to three individual plants of each genotype, per experiment. For each sample, 200 – 250 chloroplasts were counted, with absolute values varying between samples. Dots represent individual chloroplasts. **E, F)** Distributions of starch granule sizes measured as the area in the light microscopy images as in A and B. The three different colours correspond to three individual plants of each genotype, per experiment. 100 – 400 granules were measured in each sample. Dots represent individual granules. For all boxplots, each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. Measurements were made on the same batch of plants as Figure 11, experiment 2 (panels A, C, E here) and experiment 5 (panels B, D, F here).

Figure 11. Loss of MRC has variable effects on the total end of day (ED) leaf starch content in wheat leaves. A) Pooled starch content data of wheat leaves harvested at the end of day (ED) in experiments 1 (dark brown, $n = 5 – 6$ per genotype), 2 (light brown, $n = 5 – 6$ per genotype) and 3 (orange, $n = 16 – 19$ per genotype). **B)** Pooled starch content data from wheat leaves in experiments 4 (dark brown, $n = 10 – 15$ per genotype) and 5 (light brown, $n = 10 – 12$ per genotype) harvested at ED. **C)** Pooled starch content data of wheat leaves harvested at the end of night (EN) in experiments 1 (dark brown, $n = 4 – 6$ per genotype) and 2 (light brown $n = 4 – 6$ per genotype) **For all panels**, raw data and boxplots are shown on the left, where dots indicate values from individual wheat plants, with colours indicating each experiment. Each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. All statistical analyses were performed using a linear model with genotype and experiment as fixed effects for each panel, ANOVA and Tukey post-hoc tests. Panels on the right indicate differences in adjusted means of total starch content based on these models and pairwise comparisons of the genotypes. The difference in means is indicated by a dot, with whiskers showing the 95% confidence interval (CI) of this difference, with the corresponding p-value. For A and C, blue indicates comparisons between mutant and wild type, while grey indicates comparisons between mutants. For B, grey indicates the WT or *mrc-1* mutant compared to the backcrossed line with the equivalent genotype at the *MRC* loci, and blue indicates all other pairwise comparisons.

Figure 12. Model of MRC function in wheat developing endosperm. MRC is required for the control of the timing of B-type granule initiation during early grain development. In wild type, A-type granules initiate around 4 days post anthesis (dpa), and B-type granules initiate around 15 – 20 dpa. We propose that the expression of MRC during early endosperm development prevents the B-type granule formation, and B-type granule volume increases as MRC expression decreases after 10 dpa. This process is disrupted in mutants lacking a functional MRC protein, and granules initiate already from 10 dpa. The early initiation of B-type granules provides them with more time and substrate to grow, leading to higher volume of B-type granules in the mutant at grain maturity compared to the wild type, and a concomitant decrease in A-type granule size in the mutant.

Supplemental data

Supplemental File 1. *Aegilops speltoides* MRC gene model

Supplemental Figure 1. Phylogenetic analysis of wheat MRC sequences.

Supplemental Figure 2. Chain length distribution and amylose content of the *mrc* mutants.

Supplemental Figure 3. The 6B pseudogene does not contribute to granule size distribution in wheat endosperm starch.

Supplemental Table 1. Reads mapped to retrotransposon insertion genetic signatures on *MRC-B1*.

Supplemental Table 2. Pairwise comparisons of wheat growth phenotypes.

Supplemental Table 3. KASP markers for genotyping the wheat mutants.

Supplemental Table 4. Summary of leaf starch quantification experiments.

Figure 1. *MRC* **homeologs in wheat are encoded on chromosomes 6A and 6D, with a disruption of the 6B homeolog. A)** Location of *TaMRC* homeologs on chromosome 6A and 6D. The pink boxes represent *TaMRC* homeologs, while homeologs of the adjacent genes are shown in green (cytochrome P450 family protein), purple (respiratory burst oxidase homolog) and blue (uncharacterised protein). Arrowheads on the boxes indicate direction of transcription. The syntenic region on chromosome 6B has a large insertion, depicted with a black arrowhead. The diagram is drawn to scale, and chromosome coordinates of the region are indicated. **B)** Gene models of the *TaMRC-A1* and *–D1* homeologs and 6B pseudogene. Exons are represented with pink boxes, while light pink boxes represent the 5' and 3' UTRs. On the 6B region, areas with sequence similarity to exon 2 and UTRs of *TaMRC-A1* are indicated, as well as the location of *gypsy* retrotransposons. The locations of the mutations in the *mrc* mutants are depicted with red arrows, and the mutated codons/amino acids are shown in red letters. Large black arrowheads show where the sequence has been truncated for illustration - the length of truncated sequence is indicated above. **C)** Summary of species analysed for the loss of *MRC-B1* during wheat hybridisation

Figure 2. *MRC* **is expressed in wheat leaves and wheat endosperm during early grain development. A)** Expression of *TaMRC* homeologs between between different tissue types in *Triticum aestivum*. Data were obtained from the wheat expression browser, and values represent transcript per million (TPM) ± SEM from n = 89 (roots), 48 (leaves/shoots), 280 (spike), 166 (grain) samples from different experiments. **B)** Expression of *TtMRC-A1* (TRITD6Av1G081580.1) in the starchy endosperm during grain development of *Triticum turgidum* (variety Kronos). Data are from an RNAseq experiment described in Chen *et al.* (2022). Values represent TPM ± SEM from n = 3 for all timepoints.

Figure 3. Mutations in *MRC* do not affect plant growth or the number and size of grains. A) Photographs of 7-week-old wild type (WT) and *mrc* mutant plants. Bar = 9 cm. **B)** Number of tillers per plant. **C)** Number of grains per plant. **D)** Photographs of mature grains. Bar = 1 cm. **E)** Thousand Grain Weight (TGW) of mature grains. **F)** Grain size, measured as the average area of individual mature grains (15 – 88 grains) from each plant. For statistical analysis of B, C, E, F, see Supplemental Table 2. For all boxplots, each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. Dots are measurements from individual plants, with the same group of plants measured for all phenotypes and *n* = 9 – 10 individual plants.

Figure 4. Mutations in *MRC* have variable effects on total starch in mature grains. A) Total endosperm starch content in mature grains in WT and *mrc* mutants. B) Number of starch granules per mg of grain, counted with a Coulter counter. For A and B, dots on the boxplots indicate individual plants, with *n* = 8 - 9. Each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. All statistical analyses were performed using a linear model, with a one-way ANOVA and Tukey post-hoc test. Panels on the right indicate differences in means between genotypes from pairwise comparisons based on these models. The difference in means is indicated by a dot, with whiskers showing the 95% confidence interval (CI) of this difference, with the corresponding p-value. Grey indicates the WT or *mrc-1* mutant compared to the backcrossed line with the equivalent genotype at the *MRC* loci, and blue indicates all other pairwise comparisons.

Figure 5. Endosperm starch in mature grains of *mrc* mutants have altered granule size distribution. A) Coulter counter traces with evenly binned x-axes show a bimodal distribution of granule sizes from purified wheat endosperm starch. Data points are mean values from 9 individual plants of each genotype (3 grains from each plant), with the standard error of the mean shown as a shaded ribbon. WT and *mrc-1* in the left and right panel are the same data. **B, C, D)** Mean B-type granule percentage volume (B), A-type granule diameter (C) and B-type granule diameter (D) values extracted from bimodal log-normal distribution curves fitted to Coulter counter traces of individual plants. Extracted values and boxplots are shown on the left, where dots indicate mean values from individual plants (*n* = 9). Each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. All statistical analyses were performed using linear models for each panel, with a one-way ANOVA and Tukey post-hoc test. Panels on the right indicate differences in means between genotypes from pairwise comparisons based on these models. The difference in means is indicated by a dot, with whiskers showing the 95% confidence interval (CI) of this difference, with the corresponding p-value. Grey indicates the WT or *mrc-1* mutant compared to the backcrossed line with the equivalent genotype at the *MRC* loci, and blue indicates all other pairwise comparisons.

Figure 6. Endosperm starch granules of mature grains in three wheat *mrc* mutants are similar in shape compared to wild type. A) Thin sections of mature endosperm tissue were stained with Lugol's solution and imaged using light microscopy. Bar = 40 μm. **B)** Purified endosperm starch granules were observed using scanning electron microscopy (SEM). Bar = 20 μm.

Figure 7. In developing endosperm, granule number increases in *mrc-1* compared to WT despite having similar starch content. The endosperm was dissected from developing grains of WT and *mrc-1*, harvested at 8, 14, 20 and 30 dpa, with *n* = 3 - 4 individual plants for each genotype per time point, indicated by coloured dots in the upper panels, with the mean ± 95% confidence interval (CI) in black dots and whiskers. **A)** Starch content of the endosperm. Values are expressed relative to the fresh weight of the dissected endosperm. **B)** Starch granule number in the endosperm. Starch was purified from dissected endosperm and the number of granules was determined using a Coulter counter. Values are expressed relative to the fresh weight of the dissected endosperm. For A and B, individual linear models were fitted to the data of each time point, with a one-way ANOVA and Tukey post-hoc test to compare the means of WT and *mrc-1*. Panels on the bottom summarise the differences in means from these linear models, indicated by a dot, with whiskers showing the 95% CI of this difference, with the corresponding p-value.

Figure 8. The *mrc-1* mutant initiates B-type granules earlier in grain development than in the wild type (WT). The endosperm was dissected from developing grains of WT and *mrc-1* harvested at 8, 14, 20 and 30 dpa. **A)** Coulter counter traces with evenly binned x-axes show the starch granule size distribution of endosperm starch. Distributions are the average of *n* = 4 measurements, each carried out on grains harvested from a different plant (three grains per measurement). Data points are mean values from these 4 measurements, with the standard error of the mean shown as a shaded ribbon. The B-type granule peak in the *mrc-1* mutant at 14 dpa is indicated with a blue arrow. **B)** Endosperm starch granules were observed using scanning electron microscopy (SEM). Bars = 20 μm. Examples of A-type granules and B-type granules are marked with red and blue arrows respectively.

Figure 9. The early initiating B-type granules in *mrc-1* occur at least partially in stromules. **A)** Scanning electron micrographs of developing endosperm tissue subjected to critical point drying. Grains were harvested from WT and *mrc-1* at 10, 15 and 20 days post anthesis (dpa). A representative region of each panel has been pseudo-coloured, with A-type granules in red shading and B-type granules in blue shading. Bars = 50 μm. **B)** Light micrographs of endosperm sections. Semi-thin sections of embedded developing grains (15 dpa) were stained with toluidine blue as a negative stain for starch granules. Examples of A-type granules and B-type granules are marked with red and blue arrows respectively. Bars = 50 μm. **C)** Endosperm sections observed using transmission electron microscopy, using the same samples as in B. The periphery of amyloplast membranes, where visible, are indicated with a dotted white line. Bars = 1 μm.

Figure 10. Chloroplasts of *mrc-1* mutants have fewer and larger granules than the wild type. A, B) Representative light microscopy images of thin sections (500 nm) from the middle of the older of two leaves in 10-day old wheat seedlings, collected at the end of day. Sections were stained for starch using periodic acid and Schiff staining. Scale bar = 10 μm. **C, D)** Distributions of counts of granule sections per chloroplast section, counted from light microscopy images as in A and B. The three different colours correspond to three individual plants of each genotype, per experiment. For each sample, 200 – 250 chloroplasts were counted, with absolute values varying between samples. Dots represent individual chloroplasts. **E, F)** Distributions of starch granule sizes measured as the area in the light microscopy images as in A and B. The three different colours correspond to three individual plants of each genotype, per experiment. 100 – 400 granules were measured in each sample. Dots represent individual granules. For all boxplots, each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. Measurements were made on the same batch of plants as Figure 11, experiment 2 (panels A, C, E here) and experiment 5 (panels B, D, F here).

Figure 11. Loss of MRC has variable effects on the total end of day (ED) leaf starch content in wheat leaves. A) Pooled starch content data of wheat leaves harvested at the end of day (ED) in experiments 1 (dark brown, $n = 5 - 6$ per genotype), 2 (light brown, $n = 5 - 6$ per genotype) and 3 (orange, $n = 16 - 19$ per genotype). **B)** Pooled starch content data from wheat leaves in experiments 4 (dark brown, $n = 10 - 15$ per genotype) and 5 (light brown, $n = 10 - 12$ per genotype) harvested at ED. **C)** Pooled starch content data of wheat leaves harvested at the end of night (EN) in experiments 1 (dark brown, $n = 4 - 6$ per genotype) and 2 (light brown $n = 4 - 6$ per genotype) **For all panels**, raw data and boxplots are shown on the left, where dots indicate values from individual wheat plants, with colours indicating each experiment. Each box encloses the middle 50% of the distribution, the middle line is the median and the whiskers are the minimum and maximum values within 1.5 of the interquartile range. All statistical analyses were performed using a linear model with genotype and experiment as fixed effects for each panel, ANOVA and Tukey post-hoc tests. Panels on the right indicate differences in adjusted means of total starch content based on these models and pairwise comparisons of the genotypes. The difference in means is indicated by a dot, with whiskers showing the 95% confidence interval (CI) of this difference, with the corresponding p-value. For A and C, blue indicates comparisons between mutant and wild type, while grey indicates comparisons between mutants. For B, grey indicates the WT or *mrc-1* mutant compared to the backcrossed line with the equivalent genotype at the *MRC* loci, and blue indicates all other pairwise comparisons.

Figure 12. Model of MRC function in wheat developing endosperm. MRC is required for the control of the timing of B-type granule initiation during early grain development. In wild type, A-type granules initiate around 4 days post anthesis (dpa), and B-type granules initiate around 15 – 20 dpa. We propose that the expression of MRC during early endosperm development prevents the B-type granule formation, and B-type granule volume increases as MRC expression decreases after 10 dpa. This process is disrupted in mutants lacking a functional MRC protein, and granules initiate already from 10 dpa. The early initiation of B-type granules provides them with more time and substrate to grow, leading to higher volume of B-type granules in the mutant at grain maturity compared to the wild type, and a concomitant decrease in A-type granule size in the mutant.