Altered GC- and AT-biased genotypes of *Ophiocordyceps sinensis* in the stromal fertile portions and ascospores of natural *Cordyceps sinensis*

Yu-Ling Li[1][¶], Ling Gao[2][¶], Yi-Sang Yao[2][¶], Xiu-Zhang Li[1], Zi-Mei Wu[3], Ning-Zhi Tan[2], Zhou-Qing Luo[3][#a], Wei-Dong Xie[2], Jian-Yong Wu[4,5], Jia-Shi Zhu[1,2]*

[1] State Key Laboratory of Plateau Ecology and Agriculture, Qinghai Academy of Animal and Veterinary Sciences, Qinghai University, Xining 810016, Qinghai, China

[2] Shenzhen Key Laboratory of Health Science and Technology, Institute of Biopharmaceutical and Health Engineering, Shenzhen International Graduate School, Tsinghua University, Shenzhen 518055, Guangdong, China

[3] School of Life Sciences, Tsinghua University, Beijing 100084, China

[4] State Key Laboratory of Chinese Medicine and Molecular Pharmacology, Shenzhen 518057, Guangdong, China

[5] Department of Applied Biology and Chemistry Technology, The Hong Kong Polytechnic University,

Hong Kong

[#a] Current address: State Key Laboratory of Cellular Stress Biology, School of Life Sciences, Faculty of

Medicine and Life Sciences, Xiamen University, Xiamen 361102, China

* Corresponding author

E-mail: zhujosh@163.com (JSZ)

¶ These authors contributed equally to this work.

Short title:

Altered GC- and AT-biased genotypes of *Ophiocordyceps sinensis* in natural *Cordyceps sinensis*

Abstract:

Objective: To examine multiple genotypes of *Ophiocordyceps sinensis* in a semi-quantitative manner in the stromal fertile portion (SFP) densely covered with numerous ascocarps and ascospores of natural *Cordyceps sinensis* and to outline the dynamic alterations of the coexisting *O. sinensis* genotypes in different developmental phases. **Methods:** Mature *Cordyceps sinensis* specimens were harvested and continuously cultivated in our laboratory (altitude 2,254 m). The SFPs (with ascocarps) and fully and semi-ejected ascospores were collected for histological and molecular examinations. Biochip-based single nucleotide polymorphism (SNP) MALDI-TOF mass spectrometry (MS) was used to genotype multiple *O. sinensis* mutants in the SFPs and ascospores. **Results:** Microscopic analysis revealed distinct morphologies of the SFPs (with ascocarps) before and after ascospore ejection and SFP of developmental failure, which, along with the fully and semi-ejected ascospores, were subjected to SNP MS genotyping analysis. Mass spectra showed the coexistence of GC- and AT-biased genotypes of *O. sinensis* that were genetically and phylogenetically distinct in the SFPs before and after ejection and of developmental failure and in fully and semi-ejected ascospores. The intensity ratios of MS peaks were dynamically altered in the SFPs and the fully and semi-ejected ascospores. Mass spectra also showed transversion mutation alleles of unknown upstream and downstream sequences with altered intensities in the SFPs and ascospores. Genotype #5 of AT-biased Cluster-A maintained a high intensity in all SFPs and ascospores. An MS peak with a high intensity containing AT-biased Genotypes #6 and #15 in pre-ejection SFPs was significantly attenuated after ascospore ejection. The abundance of Genotypes #5–6 and #16 of AT-biased Cluster-A was differentially altered in the fully and semi-ejected ascospores that were collected from the same *Cordyceps sinensis* specimens. **Conclusion:** Multiple *O. sinensis* genotypes coexisted in different combinations with altered abundances in the SFPs prior to and after ejection, the SFP of developmental failure, and the two types of ascospores of *Cordyceps sinensis*, demonstrating their genomic independence. Metagenomic fungal members present in different combinations and with dynamic alterations play symbiotic roles in different compartments of natural *Cordyceps sinensis*.

Keywords: natural *Cordyceps sinensis*; stromal fertile portion densely covered with numerous ascocarps; ascospores; SNP MALDI-TOF mass spectrometry genotyping; GC- and AT-biased genotypes of *Ophiocordyceps sinensis*; transition point mutation; transversion point mutation

Introduction

Natural *Cordyceps sinensis* is one of the most highly valued therapeutic agents in traditional Chinese medicine (TCM), with a centuries-long history of clinical use as a folk medicine administered for "Yin-Yang" double invigoration, health maintenance, disease amelioration, and post-disease and post-surgery recovery [1–2]. The therapeutic profiles of *Cordyceps sinensis* for respiratory, cardiovascular, endocrine, kidney, and liver systems, etc., have been studied using modern pharmacological approaches and summarized [2–3]. The lifespan-extending properties of natural *Cordyceps sinensis* and its mycelial fermentation products were validated in mice in a 4-year pharmacology study, and the molecular mechanisms have been explored using a whole-gene expression biochip technique [4–5].

The Chinese Pharmacopeia defines natural *Cordyceps sinensis* as an insect-fungal complex consisting of the *Ophiocordyceps sinensis* fruiting body and a dead Hepialidae moth larva, *i.e.*, natural *Cordyceps sinensis* is not equal to the *O. sinensis* fungus [6–11]. Anatomy studies have reported that the caterpillar body of *Cordyceps sinensis* contains an intact larval intestine, an intact and thick larval body wall with numerous bristles, head tissues, and fragments of other larval tissues [7,10–11]. Yao and Zhu [6] and Li et al. [7] addressed the controversy surrounding the indiscriminate use of Latin names for natural insect-fungal complexes and multiple anamorphic-teleomorphic fungi. In this context, we temporarily refer to the fungus/fungi as *O. sinensis* and continue the customary use of the name *Cordyceps sinensis* to refer to the wild or cultivated insect-fungal complex, although this practice will very likely be replaced by the discriminative use of exclusive Latin names for different biological entities in the future.

Natural *Cordyceps sinensis* grows only in alpine areas above 3,000–3,500 m on the Qinghai-Tibetan Plateau and has a complex lifecycle [9–11]. Its maturation stages greatly impact its compartmental mycobiota profile [12–18], competitive proliferation of *Hirsutella sinensis*-like fungus/fungi [14], metagenomic polymorphism and diversity [19–26], differential coexistence and dynamic alterations of multiple genotypes of *O. sinensis* [13,19–23], metatranscriptomic and proteomic expression [27–28], chemical constituent fingerprint [14], and therapeutic efficacy and potency of the caterpillar body and stroma of *Cordyceps sinensis* as a natural therapeutic agent [29]. Mycologists have identified 22 species from 13 fungal genera in natural *Cordyceps sinensis* [30], and molecular methodologies have revealed >90 species from >37 fungal genera [15–16] and 17 genotypes of *O. sinensis* [7,9–11,31] and demonstrated the predominance of different fungi and metagenomic fungal diversity in the stroma and caterpillar body of natural and cultivated *Cordyceps sinensis* [15–16,19–23,31].

Wei et al. [32] hypothesized that *H. sinensis* is the sole anamorph of *O. sinensis* based on the collection of 3 lines of evidence: (1) frequent isolation and mycological identification according to sporulation, conidial morphology and growth characteristics [30]; (2) microcycle conidiation of the ascospores [33–35]; and (3) systematic molecular analyses based on internal transcribed spacer (ITS) sequencing and random molecular marker polymorphism assays [32,36–39]. The sole *O. sinensis* anamorph hypothesis for *H. sinensis* [32] has been widely accepted because the 3 lines of evidence meet the first and second criteria of Koch's postulates. Wei et al. [40] published a successful industrial artificial cultivation project reportedly using 3 GC-biased Genotype #1 *H. sinensis* strains as inoculants, seemingly satisfying the third criterion of Koch's postulates. However, the authors reported the detection of a single teleomorphic *O. sinensis* fungus in the fruiting body of cultivated *Cordyceps sinensis*, the sequence of which was reported to be identical to AT-biased Genotype #4 AB067749. Many studies have proven that the sequences of all AT-biased *O. sinensis* genotypes are not present in the 5 genome assemblies of GC-biased Genotype #1 *H. sinensis* Strains 1229, CC1406-203, Co18, IOZ07, and ZJB12195 [41–45] but instead belong to the genomes of independent *O. sinensis* fungi [7–11,13,31]. Thus, the sequence mismatching between the inoculants and teleomorphic *O. sinensis* in cultivated *Cordyceps sinensis* products reported by Wei et al. [40] disproves the sole anamorph hypothesis for *H. sinensis* according to the fourth criterion of Koch's postulates.

Teleomorphs of *O. sinensis* found in natural *Cordyceps sinensis* specimens collected from different geographic areas have been reported to belong to Genotypes #1 [37–38,40], #3 [46], #4 [40], either #4 or #5 [37,47] and #7 [48] without coexisting with other genotypes of *O. sinensis*. Wei et al. [40] reported teleomorphs of GC-biased Genotype #1 and AT-biased Genotype #4 of *O. sinensis* in natural and cultivated *Cordyceps sinensis*, respectively. Li et al. [49] reported the detection of both GC-biased Genotype #1 and AT-biased Genotype #5 in 25-day cultures of teleomorphic monoascospores of natural *Cordyceps sinensis*. Bushley et al. [50] reported multicellular heterokaryotic mono-, bi-, and tri-nucleated structures of *Cordyceps sinensis* hyphae and ascospores. Li et al. [31] reported the differential coexistence of multiple GC- and AT-biased genotypes of *O. sinensis* in the immature and mature stromata, stromal fertile portion (SFP) that was densely covered with numerous ascocarps, and ascospores of natural *Cordyceps sinensis*, which were altered during *Cordyceps sinensis* maturation. These findings regarding teleomorphic *O. sinensis* indicate that the independent metagenomic members of natural *Cordyceps sinensis* alternate dynamically and play symbiotic roles in different developmental and maturational stages of the *Cordyceps sinensis* lifecycle, constituting a natural integrated microecosystem [29].

In this study, we examined the microscopic morphology of SFPs prior to and after ascospore ejection and SFP of developmental failure. We also monitored the dynamic alterations of *O. sinensis* genotypes in a semiquantitative manner in the SFP samples and in the fully and semi-ejected ascospores using the biochip-based MassARRAY single nucleotide polymorphism (SNP) MALDI-TOF mass spectrometry (MS) genotyping technique. We observed differential co-occurrence and dynamic alterations of *O. sinensis* genotypes in the SFPs and ascospores of *Cordyceps sinensis*.

Materials and Methods

Collection of *Cordyceps sinensis* specimens

Cordyceps sinensis specimens were purchased in local markets from the Hualong (located at 36°13'N, 102°19'E; altitude 3553.1 m) and Yushu areas (located at 33°01'N, 96°48'E; altitude 4457.6 m) of Qinghai Province, China. Governmental permission was not required for *Cordyceps sinensis* purchases in local markets, and the collections of *Cordyceps sinensis* specimens from sales by local farmers fall under the governmental regulations for traditional Chinese herbal products.

Mature *Cordyceps sinensis* specimens were collected in mid-June. These specimens showed a plump caterpillar body and a long stroma (>5.0 cm) with the formation of an expanded fertile portion close to the stromal tip (*cf*. Fig 3.1 below), which was densely covered with ascocarps (*cf*. Fig 3.2) [25,31].

Some specimens used in the histology examination were washed thoroughly on site in running water with gentle brushing, soaked in 0.1% mercuric chloride for 10 min for surface sterilization and washed 3 times with sterile water. The thoroughly cleaned specimens were immediately frozen in liquid nitrogen on site and kept frozen during transportation to the laboratory and during storage prior to further processing [14,19,31].

Continued cultivation of mature *Cordyceps sinensis* specimens

Some mature *Cordyceps sinensis* specimens were harvested along with the outer mycelial cortices and soil surrounding the caterpillar body, replanted in paper cups in soil obtained from *Cordyceps sinensis* production areas and cultivated in our laboratory (altitude 2,254 m; longitude and latitude of 101º45' E 36º40' N) in Xining City, Qinghai Province of China (*cf*. Fig 1 of [31] and Supplementary SR1 Fig).

Because of the phototropism of natural *Cordyceps sinensis*, we kept the windows fully open, allowing sufficient sunshine and a natural plateau breeze blowing over the cultivated specimens in the paper cups. The room temperature was maintained naturally, fluctuating with the lowest temperature at 18−19°C during the night and the highest temperature at 22−23°C in the early afternoon. The relative humidity of our laboratory was maintained at approximately 30%–40%, and the absolute humidity on the surface of *Cordyceps sinensis* stromata was maintained at up to 90%–100% by spraying water using an atomizer twice a day in the morning and evening [31].

Collection and preparation of *Cordyceps sinensis* ascospores

Fig 1 of [31] (Supplementary SR1 Fig) shows a collection of the fully ejected ascospores of *Cordyceps sinensis* using double layers of autoclaved weighing paper. It also showed numerous ascospores adhered to the outer surface of asci during massive ascospore ejection, which failed to be brushed off for collection using an autoclaved brush; hence, these ascospores were instead gently scratched off using a disinfected inoculation shovel or ring and referred to as semi-ejected ascospores.

The fully and semi-ejected ascospores were cleaned separately by 2 washes with 10% and 20% sucrose solutions and 10-min centrifugation at 1,000 rpm (desktop centrifuge, Eppendorf, Germany); the supernatant was discarded after each centrifugation. The pellets (ascospores) were subsequently washed with 50% sucrose solution and centrifuged for 30 min, and the ascospores that floated on the liquid were collected [31,51]. The 2 types of ascospores were stored in a -80°C freezer prior to further processing.

Collection of *Cordyceps sinensis* SFPs

The SFPs (*cf.* Fig 3.1 below) were sampled prior to and after ascospore ejection. The SFP of developmental failure that showed no sign of ascospore ejection during prolonged cultivation were also sampled. These SFP samples were stored in an -80°C freezer prior to further processing for DNA extraction.

Reagents

Laboratory common reagents, such as ethanol, sucrose, paraformaldehyde, hematoxylin, eosin, agarose and electrophoresis reagents, were purchased from Beijing Bioland Technology Company. The Universal DNA Purification kit was a product of the TIANGEN Biotech Company, China. The DNeasy Plant Mini Kit was a product of Qiagen Company, Germany. The *Taq* PCR reagent kit and Vector NTI

Advance 9 software were purchased from Invitrogen, United States.

Reagents and biochips required for MassARRAY SNP MALDI-TOF MS genotyping were from Sequenom, USA (acquired by Agena Bioscience Company). The SNP MALDI-TOF MS genotyping experiments were performed at CapitalBio Co., Beijing, using a 384-well biochip (SpectroCHIP, Sequenom) and a commercial "iLEX Gold Cocktail" (Sequenom; exact formula unavailable) containing all necessary PCR reagents, including polymerase and dNTPs, and dideoxy A, T, C, and G terminator nucleotides (ddNTPs) to control DNA chain elongation.

Histological examination of the SFP, ascocarps and ascospores of *Cordyceps sinensis*

Some of the mature *Cordyceps sinensis* stromal specimens collected prior to and after the massive ejection of ascospores and of developmental failure were immersed in 10% formalin for fixation and subjected to dehydration in 50%, 70% and 95% ethanol for 1 h each [52]. The SFP tissues (with numerous ascocarps) were embedded in paraffin and sliced to 5-μm thickness (Model TN7000 Microtome, Tanner Scientific, Germany) [31,52]. The ascus slices were stained with hematoxylin-eosin (HE) and observed under optical and confocal microscopes (Model Primo Star and Model LSM 780, ZEISS, Germany).

Sample preparation and extraction of genomic DNA

The frozen SFP samples were individually ground to a powder in liquid nitrogen, including the SFPs (with ascocarps) (1) prior to ascospore ejection, (2) after ascospore ejection, and (3) of developmental failure, each of which was a combination of 5 specimens. The ascospores (4) fully ejected and (5) semi-ejected were collected from 5 *Cordyceps sinensis* specimens and treated in a lysate solution before extracting genomic DNA [31]. Genomic DNA was individually extracted from these SFP powders and lysed ascospores using the DNeasy Plant Mini Kit (Qiagen, Germany) according to the manufacturer's manual [14,25,31] and quantified by measuring the absorbance at UV_{260} (Eppendorf BioSpectrometer®; Hamburg, Germany).

Primer design for the first-step PCR in SNP genotyping analysis

The primers *Hsprp1*, *Hsprp3*, *ITS4*, *P1*, *P2* and *P4*, which were designed, verified, used, and reported previously [14,19–23], were used in the first-step PCR (Table 1, Fig 1) to amplify the ITS1-5.8S-ITS2 segments according to the following protocol: (1) 95°C for 5 min; (2) 35 cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 1 min; (3) 72°C for 10 min; with final holding at 4°C [14,19]. The primer pairs *Hsprp1/3* and *Hsprp1/ITS4* were favorable for amplifying the ITS sequences of GC-biased genotypes, and the primer pairs *P1/P2*, *P2/P1* and *P2/P4* were designed to amplify the sequences of AT-biased genotypes (the GC and AT content information of the 17 *O. sinensis* genotypes is listed in Supplementary S1 Table). All primers were synthesized by Invitrogen Beijing Lab. or Beijing Bomaide Technology Co. Amplicons from the first-step PCR were quantified by measuring the absorbance at UV_{260} and examined by agarose gel electrophoresis.

Table 1. "Universal" primer and genotype-specific primers for the first-step PCR amplification of ITS segments.

Primer Name	Direction	Primer sequence
For amplification of the ITS sequences of GC-biased Genotype #1		
Hsprp1	Forward	ATTATCGAGTCACCACTCCCAAACCCCC
Hsprp3	Reverse	CGAGGTTCTCAGCGAGCTACT
ITS4	Reverse	TCCTCCGCTTATTGATATGC
For amplification of the ITS sequences of AT-biased genotypes		
P1 (AB067740$_{260–279}$, AB067744$_{256–275}$)	Forward	ACGCAGCGAAATACGATAAG
P2 (AB067740$_{356–338}$, AB067744$_{352–334}$)	Reverse	CATGCCCGCTAGAGTGCTA
P4 (AB067740$_{260–279}$, AB067744$_{256–267}$)	Forward	ACGCAGCGAAATGCAATAAG

Fig 1. Alignment of ITS sequences of GC- and AT-biased genotypes of *O. sinensis* and primer locations. GT refers to genotype. Genotypes #1–2 and #13–14 are GC-biased *O. sinensis* genotypes, and Genotypes #4–6 and #15–17 are AT-biased *O. sinensis* genotypes. The GC and AT contents of the 17 *O. sinensis* genotypes are shown in Supplementary S1 Table. The underlined sequences in black represent primers for the first-step PCR. Extension primers 067721-477, 067721-531, 067740-324, 067744-324 (RC, reverse complement) and 067740-328 (RC) in brown were used for SNP mass spectrometry genotyping to distinguish between genotypes of *O. sinensis*. Arrows indicate the directions of the primer extension reactions toward the SNP alleles in brown. "-" refers to an identical base, and spaces refer to unmatched sequence gaps.

S1 Table. Percent contents of GC and AT bases in the ITS1-5.8S-ITS2 sequences of multiple genotypes of *O. sinensis*.

	Genotype	Representative sequence	% GC content	% AT content
GC-biased	#1	AB067721	63.3%	36.7%
	#2	MG770309	58.1%	41.9%
	#3	HM595984	63.3%	36.7%
	#7	AJ488254	65.3%	34.7%
	#8	GU246286	59.5%	40.5%
	#9	GU246288	64.4%	35.6%
	#10	GU246287	59.8%	40.2%
	#11	JQ695935	56.3%	43.7%
	#12	GU246296	63.7%	36.3%
	#13	KT339190	58.5%	41.5%
	#14	KT339178	58.1%	41.9%
AT-biased	#4	AB067744	52.2%	47.8%
	#5	AB067740	48.1%	51.9%
	#6	EU555436	51.3%	48.7%
	#15	KT232017	53.1%	46.9%
	#16	KT232019	50.9%	49.1%
	#17	KT232010	51.2%	48.8%

Phylogenetic analysis of multiple genotypes of *O. sinensis*

GenBank accession numbers of 17 genotypes of *O. sinensis* are listed in Supplementary S2 Table with hyperlinks to the GenBank database and were analyzed by conferring a phylogenetic tree using the maximum likelihood (ML) algorithm with MEGA software (Fig 2) [53] to assist in the design and analysis

of the SNP genotyping experiments. The ITS1-5.8S-ITS2 segment sequence ranges in 5 genomes of

Genotype #1 *H. sinensis* Strains 1229, CC1406-203, Co18, IOZ07, and ZJB12195 [41–45], either a single

ITS copy or multiple repetitive ITS copies (Supplementary S2 Table), were entered into the ML

phylogenetic analysis.

S2 Table. GenBank accession numbers that are hyperlinked to the GenBank database, *H.*

***sinensis* strain, *O. sinensis* isolate, amplicon clone information, and sequence ranges for the**

52 ITS sequences analyzed phylogenetically in Fig 2.

GenBank Accession#	*H. sinensis* strain	*O. sinensis* isolate or amplicon clone	Sequence range
LWBQ01000008	ZJB12195		991797→992287
NGJJ01000753	CC1406-203		14824→14113
NGJJ01000852	CC1406-203		10703→11192
ANOV01021709	Co18		896→1386
JAAVMX010000002	IOZ07		18688917→18689407
JAAVMX010000008	IOZ07		1199→1687
JAAVMX010000008	IOZ07		13823→14313
JAAVMX010000017	IOZ07		47097→47569
JAAVMX010000018	IOZ07		700→1186
JAAVMX010000017	IOZ07		9147→9637
JAAVMX010000017	IOZ07		21791→22281
JAAVMX010000018	IOZ07		51467→51958
JAAVMX010000018	IOZ07		38771→39261
JAAVMX010000018	IOZ07		26076→26566
JAAVMX010000018	IOZ07		13381→13871
JAAVMX010000019	IOZ07		44729→45251
JAAVMX010000019	IOZ07		332048→32538
JAAVMX010000019	IOZ07		19404→19894
LKHE01000582	1229		2132→2622
NGJJ01000796	CC1406-203		1514→2004
NGJJ01000799	CC1406-203		1666→17156
GU246296		Isolate ISO-29	6→509
JAAVMX010000017	IOZ07		34435→34925
KJ175199	Haplotype 7		5→492
HM595984	CUHK:CDC2-2		33→519
AB067721		Isolate GYOKUJU	59→549
AB067715		Isolate DERUGE	59→549
NGJJ01000798	CC1406-203		1177→1665
JQ695935	unknown	unknown	12→530
NGJJ01000573	CC1406-203		1→260
JAAVMX010000002	IOZ07		18702095→18702586
NGJJ01000573	CC1406-203		13133→13623
AJ488254		Isolate H1023	1←457
KT339190	Clone 2-7-13-14	Clone 2-7-13-14	37→537

GU246288		Isolate ISO-21	6→497
GU246287		Isolate ISO-20	6→497
JAAVMX010000019	IOZ07		6233→6733
KT232017		Clone 6-17-9	37→522
KT232018		Clone 6-17-10	37→522
AB067743		Isolate Yagyusan	55→544
AB067744		Isolate Satoku	54→544
KT720572		Isolate SANBI5308	11→324
KT218215	unknown	unknown	11→324
KT232019		Clone 6-17-11	37→525
AB067739		Isolate Nyaramu-1	59→544
AB067740		Isolate Nyaramu-2	59→544
KT232028		Clone 6-17-20	37→525
KT232010		Clone 6-17-2	37→525
KT232012		Clone 6-17-4	37→525
MG770309	unknown	unknown	1→301
GU246286		Isolate ISO9	6→497
KT339178		Clone 1-7-13-4	56→537

Extension primer design for MassARRAY SNP MS genotyping analysis based on the phylogenetic relationships of multiple genotypes of *O. sinensis*

In addition to the GC-biased genotype clade (side-noted in blue in Fig 2), the AT-biased genotype clade (side-noted in red) contains 2 AT-biased branches [7–11,31]: AT-biased Cluster-A consists of Genotypes #5–6 and #16–17, and AT-biased Cluster-B consists of Genotypes #4 and #15. The representative sequences of GC-biased Genotype #1 AB067721, Genotype #5 AB067740 within AT-biased Cluster-A, and Genotype #4 AB067744 within AT-biased Cluster-B were used as the reference sequences for designing SNP extension primers (Table 2) for biochip-based MassARRAY SNP MALDI-TOF MS genotyping. Extension primers 067721-477 and 067721-531 were used to distinguish between the GC- and AT-biased genotypes but not between the multiple AT-biased genotypes (*cf.* Fig 1) [19,21–22]. Extension primers 067740-324, 067744-324 (reverse complement), and 067740-328 (reverse complement) were used to distinguish between the multiple AT-biased genotypes using a stratification genotyping strategy.

Ophiocordyceps sinensis strain ZJB12195 LWBQ01000008.1 991797 to 992287
Ophiocordyceps sinensis strain CC1406-203 NGJJ01000573.1 14624 to 15113
Ophiocordyceps sinensis strain CC1406-203 NGJJ01000582.1 10703 to 11192
Ophiocordyceps sinensis CO18 ANOV01021709.1 896 to 1386
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000002.1 18688917 to 18689407
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000008.1 1199 to 1687
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000008.1 13823 to 14313
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000017.1 47079 to 47569
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000018.1 700 to 1186
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000017.1 9147 to 9637
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000017.1 21791 to 22281
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000018.1 51467 to 51958
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000018.1 38771 to 39261
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000018.1 26076 to 26566
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000018.1 13381 to 13871
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000019.1 44729 to 45251
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000019.1 32048 to 32538
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000019.1 19404 to 19894
Ophiocordyceps sinensis strain 1229 LKHE01000582.1 2132 to 2622
Ophiocordyceps sinensis strain CC1406-203 GJJ01000796.1 1514 to 2004
Ophiocordyceps sinensis strain CC1406-203 GJJ01000799.1 1666 to 17156

#1 Genome ITS sequences & genome repetitive ITS copies

71
GU246296 6-509 **#12**
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000017.1 34435 to 34925 **#1 Genome ITS copy**
KJ175199 5-492
HM595984 33-519 **#3**
AB067721 59-549
AB067715 59-549 **#1**
Ophiocordyceps sinensis strain CC1406-203 NGJJ01000798.1 1177 to 1665 **#1 Genome ITS copy**
JQ695935 12-530 **#11**
62 Ophiocordyceps sinensis strain CC1406-203 NGJJ01000573.1 1 to 260
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000002.1 18702095 to 18702586
63 Ophiocordyceps sinensis strain CC1406-203 NGJJ01000573.1 13133 to 13623

#1 Genome repetitive ITS copies

60 AJ488254 457-1 **#7**
KT339190 37-537 **#13**
GU246288 6-497 **#9**
76
67 GU246287 6-497 **#10**
Ophiocordyceps sinensis strain IOZ07 JAAVMX010000019.1 6233 to 6733 **#1 Genome ITS copy**

GC-biased sequences of *O. sinensis*

AT-biased Cluster B
KT232017 37-522
69
KT232018 37-522 **#15**
84
AB067743 55-544
AB067744 54-544 **#4**

AT-biased genotypes of *O. sinensis*
100
99 KJ720572 11-324
KX218215 11-324 **#6**
KT232019 37-525 **#16**
71
AT-biased Cluster A
88 AB067739 59-544
AB067740 59-544 **#5**
93
84 KT232028 37-525
KT232010 37-522 **#17**
KT232012 37-522

MG770309 1-301 **#2**
GU246286 6-497 **#8**
KT339178 56-537 **#14**

0.02

Fig 2. A maximum likelihood (ML) phylogenetic tree. The ML tree was inferred using MEGA software [53], containing 25 ITS sequences (representing 17 genotypes of *O. sinensis*) and 27 genomic ITS sequences (including multiple repetitive ITS copies) of Genotype #1 *H. sinensis* Strain 1229, CC1406-203, Co18, IOZ07, and ZJB12195. Supplementary S2 Table lists the GenBank accession numbers that are hyperlinked to GenBank database, strain or isolate information and ITS1-5.8S-ITS2 sequence range for the 52 ITS sequences that were analyzed phylogenetically in Fig 2. The phylogenetic topology of the ML tree was verified according to the topologies of Bayesian majority-rule consensus phylogenetic trees [7,10–11,31].

Table 2. SNP extension primers for MassARRAY MALDI-TOF MS genotyping.

Primer Name	Primer sequence
Used in the second-step PCR designed based on the AB067721 sequence to amplify the GC-biased sequences of *O. sinensis* genotypes	
067721-477	CGCCGCGGCTCCCCT
067721-531	AGGCGGTCACGCCGT
Used in the second-step PCR designed based on both AB067744 and AB067740 sequences to amplify the AT-biased sequences of *O. sinensis* genotypes	
067740-324	GTAAACTATCGAATCTTTAAACG
067744-324 (reverse complement)	GCTAGCGGGCGTAGTAT
067740-328 (reverse complement)	GTGCTAGCGGGCGTA

Biochip-based MassARRAY SNP MALDI-TOF MS genotyping

ITS sequences were amplified using the aforementioned primer pairs from genomic DNA prepared from SFPs prior to and after ejection and of developmental failure and fully and semi-ejected ascospores. The amplicons obtained from the first-step PCR were subjected to neutralization of the unincorporated dNTPs with shrimp alkaline phosphatase and used as templates for single nucleotide extension toward the selected SNP sites shown in Fig 1. These amplicons were mixed with an iLEX Gold cocktail (Sequenom) containing one of the extension primers listed in Table 2 per reaction and amplified according to the following protocol: (1) 94°C for 30 s; (2) 94°C for 5 s; (3) 52°C for 5 s; (4) 80°C for 5 s; (5) return to (3) for 4 more times (for a total of 5 "smaller" cycles); (6) return to (2) for 39 more times (for a total of 40 "larger" cycles); (7) 72°C for 3 min; and (8) 4°C indefinitely [19,21–22]. After purification, the extended products with one of the mass-modified nucleotides were analyzed using a 384-well SpectroCHIP (Sequenom), and their masses were examined using a MassARRAY analyzer to determine heterogeneous genotypes [19,21–22,54]. Each primer-extension MS genotyping test for the combined 5 samples was

289 repeated at least twice, and the genotyping results were analyzed using software TYPER 4.0 (Sequenom).

290 The MassARRAY SNP MALDI-TOF MS genotyping experiments were performed by a service company,

291 CapitalBio Co., in Beijing, China.

Results

Microscopy observations of histological structures of the SFPs

Fig 3.1 shows a mature *Cordyceps sinensis* specimen labeled for the SFP (the stromal fertile portion),

and Fig 3.2 shows a confocal image (bar: 1 mm) of a transverse section of the SFP of *Cordyceps sinensis*,

which is densely covered by numerous ascocarps (with a red circle in Fig 3.2 indicating one of the multiple

ascocarps). Fig 3.3 shows an optical microscopy image (40x) of an HE-stained *Cordyceps sinensis* ascocarp,

which contains multiple ascospores prior to ejection. Fig 3.4 shows a confocal image (bar: 100 μm) of 2

ascocarps; the upper ascocarp is a maturing ascocarp with not yet opened perithecium containing multiple

ascospores prior to ejection, and the lower ascocarp is a fully mature ascocarp during ascospore ejection

with multiple ascospores aggregating toward the opening of the perithecium. Fig 3.5 shows a confocal

image close to the opening of the perithecium during ascospore ejection with an arrow pointing to a semi-

ejected ascospore hanging out of the opening of the perithecium. Fig 3.6 shows an optical microscopy

image (10x) of HE-stained SFP after ascospore ejection. Figs 3.7 (10x) and 3.8 (40x) show optical images

of HE-stained SFPs of developmental failure with abnormal structures of ascocarps (compared to normal

structures shown in Figs 3.3–3.4) and containing no normally developed ascospores.

In addition, Fig 1 of [31] (reproduced and shown in Supplementary SR1 Fig) shows the numerous

semi-ejected ascospores that tightly adhered to the outer surface of asci. Fig 3 of [31] shows the

microscopic images of fully ejected *Cordyceps sinensis* ascospores with or without calmodulin fluorescent

staining.

3.2

1 mm

3.3

3.4

100 nm

3.5

20 nm

3.6

3.7

318
Fig 3. Microscopic images of the SFPs (with ascocarps) of natural *Cordyceps sinensis*. Panel 3.1 shows a
photo of a mature *Cordyceps sinensis* specimen labeled with stromal fertile portion (SFP). Panel 3.2 shows a
confocal image (bar: 1 mm) of a transverse section of the *Cordyceps sinensis* SFP densely covered with numerous
ascocarps prior to ascospore ejection, with a red circle indicating one of the multiple ascocarps densely covering
the SFP. Panel 3.3 shows an optical close-up image (40x) of an HE-stained ascocarp containing multiple
ascospores of multicellular ascosporic cells. Panel 3.4 shows a confocal image (bar: 100 μm) of 2 ascocarps; the
upper one is a maturing ascocarp, and the lower one is a mature ascocarp containing multiple ascospores
aggregating toward the opening of the perithecium. Panel 3.5 shows a confocal image close to the opening of the
perithecium and a semi-ejected ascospore hanging out of the opening of the perithecium. Panel 3.6 shows an
optical image (10x) of HE-stained SFPs after ascospore ejection. Panels 3.7 (10x) and 3.8 (40x) show optical
images of HE-stained SFPs of developmental failure. The confocal images shown in Panels 3.2 and 3.4 are large
images created by stitching together tile images automatically by the computer and software of the ZEISS LSM
780 confocal microscope, which allow extremely high resolution from designated large fields of view, because of
limited scanning areas of the ZEISS LSM 780 confocal microscope.

Phylogenetic analysis of 17 genotypes of *O. sinensis* and multiple repetitive

ITS copies in the genome of Genotype #1 *H. sinensis*

The ML phylogenetic tree (*cf*. Fig 2) shows the phylogeny of the 17 GC- and AT-biased genotypes
of *O. sinensis* (the GC and AT contents of the genotypes are shown in Supplementary S1 Table) and the
genome ITS sequences of Genotype #1 *H. sinensis*, including multiple repetitive ITS copies in the genome

assemblies JAAVMX000000000 and NGJJ00000000 and a single copy each in ANOV00000000, LKHE00000000 and LWBQ00000000 of Genotype #1 *H. sinensis* Strains IOZ07, CC1406-203, Co18, 1229 and ZJB12195, respectively [41–45]. The topology of the ML tree was verified with Bayesian majority rule consensus trees [7,11,31,55], which were inferred using MrBayes v3.2.7 software (the Markov chain Monte Carlo [MCMC] algorithm) [56]. Notably, each of the genome assemblies ANOV00000000, LKHE00000000 and LWBQ00000000 of *H. sinensis* Strains Co18, 1229 and ZJB12195, respectively [41–43], contains only one ITS copy, indicating that other repetitive ITS copies might have been discarded during assembly of genome shotgun sequences. All these genomic ITS sequences are GC biases and were clustered in the GC clade along with all GC-biased genotypes labeled in blue alongside Fig 2, but very distant from the AT-biased clade labeled in red alongside. The genome repetitive ITS copies contain more or less multiple scattered insertion/deletion and transversion point mutations, but no or only a few transition point mutations [57]. The genetic and phylogenetic findings indicate that the genomic repetitive ITS copies were not the mutagenic targets of Repeat-Induced Point mutation (RIP) causing cytosine-to-thymine transition. All AT-biased genotypes feature multiple scattered transition point mutations (*cf*. Fig 1), among which Genotypes #5–6 and #16–17 were clustered into AT-biased Cluster-A and Genotypes #4 and #15 in AT-biased Cluster-B (*cf*. Fig 2). The genome assemblies of GC-biased Genotype #1 *H. sinensis* strains do not contain any AT-biased genotype sequences, indicating genomic independence of the *O. sinensis* genotypes.

SNP genotyping to distinguish between GC- and AT-biased *O. sinensis* genotypes in SFPs before and after ascospore ejection and of developmental failure

Li et al. [31] reported the co-occurrence of GC-biased Genotype #1 *H. sinensis* and AT-biased Genotypes #4–6 and #15 of *O. sinensis* in SFPs based on the application of several pairs of genotype-specific primers and cloning-based amplicon sequencing approaches. The biochip-based MassARRAY SNP MALDI-TOF MS genotyping technique was used to examine the dynamic alterations of the *O. sinensis* genotypes in SFPs prior to and after ascospore ejection and of developmental failure in a semiquantitative manner. Equal amounts of genomic DNA (calculated based on the absorbance at UV_{260}) were applied to the first-step PCR using the primer pairs *Hsprp1/3* and *Hsprp1/ITS4* to amplify the ITS

sequences of GC-biased genotypes in greater quantities than those of AT-biased genotypes, and the primer pairs *P1/P2*, *P2/P1* and *P2/P4* were used to amplify the sequences of AT-biased genotypes (*cf*. Table 1 and Fig 1).

Extension primers 067721-477 and 067721-531 (*cf*. Table 2) were used in the second-step PCR, and the reactions extended the extension primers with a single nucleotide toward the SNPs at positions 477 and 531 in the AB067721 sequence (*cf*. Fig 1). Mass spectra showed 4 allelic peaks, Peaks A, C, G, and T, representing the extension reactions of the primers 067721-477 and 067721-531 with a single extended nucleotide adenine, cytosine, guanine, and thymine, respectively.

SNP genotyping using extension primer 067721-477

Fig 4.1 shows a mass spectrum of extension primer 067721-447 (*cf*. Table 2) for the SFP prior to ascospore ejection. Peak G with the highest MS intensity represents the GC-biased Genotype #1 *H. sinensis*. Peaks C, A, and T possess much lower intensities. Peak A represents AT-biased genotypes with transition point mutations (*cf*. Fig 1), and Peaks C and T represent 2 transversion point mutation genotypes of unknown upstream and downstream sequences of the allelic peaks. The primers employed for the first-step PCR preferentially amplified GC-biased genotype sequences, resulting in a greater quantity of GC-biased amplicon copies that served as the templates for the second-step PCR. Table 3 shows the intensity ratios of the MS peaks: G:A, G:C and G:T.

Fig 4.2 shows a mass spectrum for the postejection SFP. The intensities of Peaks A and G appeared to increase proportionally compared with the corresponding peaks shown in Fig 4.1, resulting in a similar G:A ratio (Table 3). In contrast, Peaks C and T were greatly attenuated, resulting in significant elevations in the intensity ratios of G:C and G:T (Table 3). The ejection-related increase in the GC- and AT-biased genotypes of *O. sinensis*, coupled with declines in the 2 unknown transversion mutation genotypes, indicate that these genotypes play different roles in SFPs prior to, during and after ascospore ejection.

Fig 4.3 shows the overlapping mass spectra of Figs 4.1 and 4.2 after adjusting and aligning the vertical axis for intensity and visualizes the dynamic alteration of the allelic peaks in the SFPs from pre-ejection to post-ejection.

Fig 4. MALDI-TOF mass spectra (MS) of extension primer 067721-477 to distinguish between GC- and AT-genotypes in the SFPs (with ascocarps) prior to and after ascospore ejection. The extension reaction proceeded from extension primer 067721-477 toward the SNP at position 477 in the AB067721 sequence (*cf*. Fig 1 for the location). In the allelic peaks, "C" denotes extension of the primer with an extended cytosine; "A" indicates an extended adenine, "G" represents an extended guanine, and "T" refers to an extended thymine. Panels 4.1 and 4.2 show the mass spectra for the SFPs prior to and after ascospore ejection, respectively. Panel 4.3 shows the overlapping mass spectra for Panels 4.1 (black tracing) and 4.2 (red tracing) with alignment of the horizontal and vertical axes of the panels.

Supplementary S1 Fig shows a mass spectrum for an SFP of developmental failure. Compared with Peaks G and A shown in Fig 4.1, the intensity of Peak G in Supplementary S1 Fig was elevated, while Peak A remained unchanged, resulting in a small elevation of the G:A intensity ratio (Table 3). Peak C was absent, while Peak T of the transversion point mutation genotype was lower than that for the pre-

408   ejection SFP shown in Fig 4.1, resulting in a significant elevation of the G:T ratio (Table 3).

409
S1 Fig. MALDI-TOF mass spectra of extension primer 067721-477 to distinguish between GC- and AT-

biased genotypes in the SFP (with ascocarps) of developmental failure. The extension reaction proceeded

from extension primer 067721-477 toward the SNP at position 477 in the AB067721 sequence (*cf*. Fig 1 for the

location). In the allelic peaks, "C" denotes extension of the primer with an extended cytosine; "A" indicates an

extended adenine, "G" represents an extended guanine, and "T" refers to an extended thymine.

Table 3. Mass intensity ratios of the SNP peaks of transition and transversion mutation genotypes in the SFPs prior
to and after ascospore ejection and of developmental failure.

Extension primer	Allelic ratio	Intensity ratio		
		Prior-ejection SFP (*cf*. Fig 4.1)	**Post-ejection SFP** (*cf*. Fig 4.2)	**SFP of developmental failure** (*cf*. Supplementary S1 Fig)
067721-477	G:A	**3.3** (5.3÷1.6)	**3.0** (↔; 8.4÷2.8)	**3.9** (↑; 7.0÷1.8)
	G:C	**4.4** (5.3÷1.2)	**21.0** (↑↑↑↑; 8.4÷0.4)	—
	G:T	**3.3** (5.3÷1.6)	**9.3** (↑↑; 8.4÷0.9)	**10.0** (↑↑↑; 7.0÷0.7)

Note: Peak G in Figs 4 and Supplementary S1 Fig represents GC-biased Genotype #1 *H. sinensis*; Peak A indicates
AT-biased genotypes of *O. sinensis* (*cf*. Fig 1). Peaks C and T denote 2 transversion mutation genotypes of unknown
upstream and downstream sequences. "↔" indicates no significant change (within 20% variation), and "↑" (increase
less than two-fold), "↑↑" (increase greater than two-fold but less than three-fold), "↑↑↑" (increase greater than three-
fold but less than four-fold) and "↑↑↑↑" (increase greater than four-fold) denote significant increases in intensity
ratios compared to that in the pre-ejection SFP. "—" means that one of the allelic peaks was missing, and no ratio
could be calculated.

SNP genotyping using extension primer 067721-531

Supplementary S2.1 Fig shows a mass spectrum of extension primer 067721-531 (Table 2) for the

pre-ejection SFP, displaying allelic Peaks A and G (representing AT-biased genotypes of *O. sinensis* and

GC-biased Genotype #1, respectively) with comparable intensities and resulting in a G:A intensity ratio

of 1.06 (Supplementary S3 Table). Because the primers employed for the first-step PCR preferentially

amplified the sequences of GC-biased genotypes, the comparable intensities of Peaks A and G might

indicate a greater quantity of natural abundance of AT-biased genotypes than GC-biased Genotype #1 *H.*

sinensis in the genomic DNA pool extracted from the pre-ejection SFP. While Peak C was absent, Peak T

showed a very low intensity, representing a transversion mutation genotype of unknown upstream and

downstream sequences and resulting in a high G:T ratio of 12.7 (Supplementary S3 Table).

S2 Fig. MALDI-TOF mass spectra of extension primer 067721-531 to distinguish between the GC- and AT-

genotypes in the SFPs (with ascocarps) before and after ascospore ejection and the SFP that failed to

develop and eject the ascospores. The extension primer 067721-531 was extended to the SNP at position 531 in

the AB067721 sequence (*cf*. Fig 1 for the location). The allele peaks are marked: "C" denotes the primer with an

extended cytosine; "A" indicates an extended adenine, "G" represents an extended guanine, and "T" refers to an

extended thymine. Panel S2.1 shows the mass spectrum for the SFP before ascospore ejection. Panel S2.2 shows

the mass spectrum for the SFP after ascospore ejection. Panel S2.3 shows the mass spectrum for the SFP that

failed to develop and eject the ascospore. Panel S2.4 shows the overlapping mass spectra for Panels S2.1 (black

tracing) and S2.2 (red tracing) with alignment of the horizontal and vertical axes of the panels.

S3 Table. Mass intensity ratios of the SNP peaks of transition and transversion mutation genotypes

in the SFPs prior to and after ascospore ejection and of developmental failure.

Extension primer	Allelic ratio	Intensity ratio		
		Pre-ejection SFP (*cf*. S2.1 Fig)	Post-ejection SFP (*cf*. S2.2 Fig)	SFP of developmental failure (*cf*. S2.3 Fig)
067721-531	G:A	1.06 (3.8÷3.6)	19.3 (↑↑↑↑; 5.5÷0.3)	13.5 (↑↑↑↑; 5.4÷0.4)
	G:C	—	—	—
	G:T	12.7 (3.8÷0.3)	—	—

Note: Peak G represents GC-biased Genotype #1 *H. sinensis*; Peak A indicates AT-biased genotypes of

O. sinensis (Fig 1). Peaks C and T denote 2 transversion mutation genotypes of unknown upstream and

downstream sequences. "↑↑↑↑" denotes significant increases in intensity ratios greater than four-fold

compared to that in the pre-ejection SFP. "—" means that one of the allelic peaks was missing and no

ratio could be calculated.

Supplementary S2.2 and S2.3 Figs show similar topologies of the mass spectra for the SFPs after

ascospore ejection and of developmental failure, respectively, displaying an elevated Peak G and a considerably attenuated Peak A relative to those shown in Supplementary S2.1 Fig. These results indicated significant decreases in the amount of AT-biased genotypes of *O. sinensis* in the SFPs after ascospore ejection and of developmental failure, while GC-biased Genotype #1 *H. sinensis* was relatively increased. The dynamic bi-directional changes resulted in increases in the G:A intensity ratios to 19.3 and 13.5 for the SFPs after ejection and of developmental failure, respectively (Supplementary S3 Table), compared with a G:A ratio of 1.06 obtained for the pre-ejection SFP. Peaks C and T are absent (Supplementary S2.2 and S2.3 Figs).

Supplementary S2.4 Fig shows the overlapping mass spectra of Supplementary S2.1 and S2.2 Figs after adjusting and aligning the vertical axis for intensity and visualizes the dynamic alteration of the allelic peaks in the SFPs from pre-ejection to post-ejection.

Stratified SNP genotyping to distinguish between the AT-biased genotypes of *O. sinensis* in the SFPs prior to and after ascospore ejection and of developmental failure

Li et al. [31] reported the detection of Genotypes #5–6 of AT-biased Cluster-A and Genotypes #4 and #15 of AT-biased Cluster-B in the SFP (*cf*. Fig 2). According to the sequence alignment shown in Fig 1, SNP MS genotyping was performed *via* a stratification strategy using extension primers 067740-324, 067744-324 (reverse complement) and 067740-328 (reverse complement) (*cf*. Table 2 and Fig 1) to distinguish between Genotypes #4–6 and #15. The templates of the extension reactions were the amplicons obtained from the first-step PCR using primer pairs *P1/P2*, *P2/P1* and *P2/P4* (*cf*. Table 1), which amplify the AT-biased sequences of *O. sinensis*.

Using extension primer 067740-328 to distinguish between AT-biased Genotype #4 and other AT-biased genotypes

Supplementary S3.1 Fig shows a mass spectrum of extension primer 067740-328 (reverse complement; *cf*. Fig 1 and Table 2) for the SFP prior to ascospore ejection. The major Peak T represents the sum of the mass intensities of Genotypes #5–6 and #15. Although the extension reaction using primer 067740-328 could also produce an extended thymine on the DNA templates of Genotypes #16–17 (*cf*. Fig

1), these genotypes were absent in the SFPs of natural *Cordyceps sinensis* according to the results obtained

using genotype-specific primers and PCR amplicon-cloning techniques [31]. Peak C was buried in a

descending MS loop, representing a very low level of Genotype #4 (*cf*. Fig 1), resulting in a T:C intensity

ratio of 14.1 (Supplementary S4 Table). Peak G was nearly flat and negligible. Peak A with a low intensity

represented a co-occurring transversion mutation genotype of unknown upstream and downstream

sequences, resulting in a T:A ratio of 11.1 (Supplementary S4 Table).

S3 Fig. MALDI-TOF mass spectra of extension primer 067740-328 (reverse complement) to distinguish

between the AT-genotypes in the SFPs (with ascocarps) prior to and after ascospore ejection and the SFP

of developmental failure. The extension reaction proceeded from extension primer 067740-328 (reverse

complement) toward the SNP at position 328 in the AB067740 sequence (*cf*. Fig 1 for the location). In the allelic peaks, "G" represents the primer with an extended reverse complement cytosine (guanine in the sense chain; *cf*. Fig 1); "T" represents an extended reverse complement adenine (thymine in the sense chain); "C" denotes an extended reverse complement guanine (cytosine in the sense chain); and "A" indicates an extended reverse complement thymine (adenine in the sense chain). Panel S3.1 shows the mass spectrum of the pre-ejection SFP. Panel S3.2 shows the mass spectrum of the post-ejection SFP. Panel S3.3 shows the mass spectrum of the SFP of developmental failure.

S4 Table. Mass intensity ratios of the SNP allelic peaks of transition and transversion mutant genotypes in SFPs prior to and after ascospore ejection and of developmental failure.

Extension primer	Allelic ratio	Intensity ratio		
		Pre-ejection SFP (*cf*. S3.1 Fig)	**Post-ejection SFP** (cf. S3.2 Fig)	**SFP of developmental failure** (*cf*. S3.3 Fig)
067740-328	T:C	14.1 (52.0÷3.7)	16.5 (↔; 61.0÷3.7)	19.3 (↑; 65.5÷3.4)
	T:G	—	—	—
	T:A	11.1 (52.0÷4.7)	20.3 (↑; 61.0÷3.0)	17.7 (↑; 65.5÷3.7)

Note: Peak T represents AT-biased Genotypes #5–6 and #15 of *O. sinensis,* and Peak C indicates AT-biased Genotype #4 of *O. sinensis* (*cf*. Fig 1). Peak G was nearly flat and negligible. Peak A denotes a transversion mutation genotype of unknown upstream and downstream sequences. "↔" indicates no significant change (within 20% variation), and "↑" denotes increases in intensity ratios less than two-fold compared to that for the pre-ejection SFP. "—" means that one of the allelic peaks was missing, and no ratio could be calculated.

Supplementary S3.2 and S3.3 Figs show mass spectra for the SFPs after ascospore ejection and of developmental failure with similar MS topologies to that shown in Supplementary S3.1 Fig. Compared with Supplementary S3.1 Fig, Peak T (Genotypes #5–6 and #15) remained the major allele peak with increased intensities in both Supplementary S3.2 and S3.3 Figs. Peak C (Genotype #4) was still buried in the descending MS loop, resulting in a similar (Supplementary S3.2 Fig) or an increased (Supplementary S3.3 Fig) T:C ratio (Supplementary S4 Table). Peak A (a transversion point mutation genotype of unknown upstream and downstream sequences) was attenuated in Supplementary S3.2 and S3.3 Figs relative to that shown in Supplementary S3.1 Fig, resulting in increases in the T:A ratio (Supplementary S4 Table).

Peak C with extremely low intensities shown in Supplementary S3.1–S3.3 Figs for all 3 SFP samples indicated a consistently low quantity of Genotype #4 of AT-biased Cluster-B, consistent with the report by

518 Li et al. [31] based on the use of genotype-specific primers and cloning-based PCR amplicon sequencing
519 techniques.

Using extension primer 067744-324 to distinguish between AT-biased Genotypes #5 and #15

After confirming consistently extremely low intensities of Peak C (Genotype #4) in all SFP
specimens shown in Supplementary S3.1–S3.3 Figs, the second step of the stratified genotyping strategy
was then applied using extension primer 067744-324 (*cf*. Table 2) to further analyze the extremely high-
intensity Peak T shown in Supplementary S3.1–S3.3 Figs, representing Genotypes #5–6 of AT-biased
Cluster-A and Genotype #15 of AT-biased Cluster-B (*cf*. Figure 2).

Fig 5.1 shows a mass spectrum of extension primer 067744-324 (reverse complement) (*cf*. Table 2)
for the pre-ejection SFP. Peak T possessing extremely high intensity represented Genotypes #6 and #15
(*cf*. Fig 1), while the component Genotype #4 was attributed only to a very small portion of Peak T, as
demonstrated in Supplementary S3.1 Fig. Peak C possessing the second highest intensity represented
Genotype #5. Although an extended cytosine may be produced in the extension reaction on the DNA
templates of Genotypes #16–17 using primer 067744-324 (*cf*. Fig 1), these 2 genotypes were absent in the
Cordyceps sinensis SFP when using genotype-specific primers and PCR amplicon-cloning techniques [31].
Peaks C and T presented a C:T intensity ratio of 0.71 (Table 4). Peak G was absent. Peak A showed a low
intensity, resulting in a C:A intensity ratio of 4.56 (Table 4), representing a co-occurring transversion point
mutation genotype of unknown upstream and downstream sequences.

Fig 5.2 shows a mass spectrum for the post-ejection SFP, displaying a high intensity of Peak C
(Genotype #5). Peak T, representing Genotypes #6 and #15, appeared in the mass spectrum as a shoulder
of the acceding MS loop and was dramatically attenuated compared to that shown in Fig 5.1, resulting in
a significantly elevated C:T ratio of 10.6 (Table 4). Peak G showed a low intensity, resulting in a C:G ratio
of 53, while Peak A remained at a low level with an elevated C:A ratio of 7.57 (Table 4).

Fig 5.3 shows the overlapping mass spectra of Figs 5.1–5.2 after adjusting and aligning the vertical
axis for intensity and visualizes the dynamic alteration of the allelic peaks in the SFPs from pre-ejection
to post-ejection.

Fig 5. MALDI-TOF mass spectra of extension primer 067744-324 (reverse complement) to distinguish between the AT-biased genotypes in the SFPs (with ascocarps) prior to and after ascospore ejection. The extension reaction proceeded from extension primer 067744-324 (reverse complement) toward the SNP at position 324 in the AB067740 sequence (*cf*. Fig 1 for the location). In the allelic peaks, "G" represents the primer with an extended reverse complement cytosine (guanine in the sense chain; *cf*. Fig 1); "T" represents an extended reverse complement adenine (thymine in the sense chain); "C" denotes an extended reverse complement guanine (cytosine in sense chain); and "A" indicates an extended reverse complement thymine (adenine in the sense chain). Panel 5.1 shows the mass spectrum of the pre-ejection SFP. Panel 5.2 shows the mass spectrum of the post-ejection SFP. Panel 5.3 shows the overlapping mass spectra for Panels 5.1 (tracing in black) and 5.2 (tracing in red) with alignment of the horizontal and vertical axes of the panels.

557 **Table 4. Mass intensity ratios of the SNP allelic peaks of transition and transversion mutant genotypes in**
558 **SFPs prior to and after ascospore ejection and of developmental failure**

Extension primer	Allelic ratio	Intensity ratio		
		Pre-ejection SFP (*cf.* Fig 5.1)	Post-ejection SFP (*cf.* Fig 5.2)	SFP of developmental failure (*cf.* Supplementary S4 Fig)
067744-324	C:T	0.71 (41÷58)	10.6 (↑↑↑↑; 53÷5.0)	15.1 (↑↑↑↑; 68÷4.5)
	C:G	—	53.0 (53÷1.0)	—
	C:A	4.56 (41÷9.0)	7.57 (↑; 53÷7.0)	8.50 (↑; 68÷8.0)

Note: Peak C in Figs 5 and Supplementary S4 Fig represents AT-biased Genotype #5 of *O. sinensis,* and Peak T
represents AT-biased Genotypes #6 and #15 of *O. sinensis* and an extremely low amount of Genotype #4. Peaks G
and A denote 2 transversion mutation genotypes of unknown upstream and downstream sequences. "↑" (increase
less than two-fold) and "↑↑↑↑" (increase greater than four-fold) denote significant increases in intensity ratios
compared to that for the pre-ejection SFP. "—" means that one of the allelic peaks was missing, and no ratio could
be calculated.

Supplementary S4 Fig shows a mass spectrum for the SFP of developmental failure. The mass

spectrum topology was similar to that shown in Fig 5.2, except that the intensity of Peak C (Genotype #5)

was further elevated, resulting in a further elevated C:T ratio (Table 4). Peak G was nearly flat and

negligible, while Peak A remained at a low intensity, with an elevated C:A ratio of 8.50 (Table 4).

569
S4 Fig. MALDI-TOF mass spectra of extension primer 067744-324 (reverse complement) to distinguish

between the AT-biased genotypes in the SFP (with ascocarps) of developmental failure. The extension

reaction proceeded from extension primer 067744-324 (reverse complement) toward the SNP at position 324 in

the AB067740 sequence (*cf.* Fig 1 for the location). In the allelic peaks, "G" represents the primer with an

extended reverse complement cytosine (guanine in the sense chain; *cf.* Fig 1); "T" represents an extended reverse

complement adenine (thymine in the sense chain); "C" denotes an extended reverse complement guanine

(cytosine in sense chain); and "A" indicates an extended reverse complement thymine (adenine in the sense

chain). This shows the mass spectrum of the post-ejection SFP.

In contrast to the consistently low levels of Genotype #4 in all 3 SFP samples shown in

Supplementary S3.1–S3.3 Figs, the second step of the stratified analysis indicated dramatic alterations of

Peak T, which showed a very high level in the pre-ejection SFP (Fig 5.1) but presented significant

attenuation in the SFPs after ejection (Fig 5.2) and of developmental failure (Supplementary S4 Fig). Peak

T represents the sum of intensities of Genotypes #6 and #15, as well as a very small portion of Genotype

#4. Both Genotypes #4 and #15 of AT-biased Cluster-B were present in the *Cordyceps sinensis* SFP but

absent in ascospore samples, as demonstrated by Li et al. [31,49]. In contrast, Genotype #5 of AT-biased

Cluster-A, which was represented by Peak C in Figs 5 and Supplementary S4 Fig, remained at a

consistently high intensity in all SFP samples. Transversion mutation genotypes of unknown upstream and

downstream sequences either remained at a constantly low level (Peak A) or were absent (Peak G) in the

SFP samples. Genotype #6 of AT-biased Cluster-A underwent dynamic quantitative alteration along with

Genotype #15 of AT-biased Cluster-B when comparing Peak T in Figs 5.1 and 5.2. Genotype #6 was

clustered phylogenetically in the clade of AT-biased Cluster-A, as shown in Fig 2, but it was closer to the

clade of AT-biased Cluster-B than other genotypes in AT-biased Cluster-A. The results may indicate that

Genotype #6 is on an evolutionary branch different from other genotypes of AT-biased Clusters A and B

(*cf*. Figure 2) and functions symbiotically in the *Cordyceps sinensis* SFPs.

SNP genotyping analysis to distinguish between the GC- and AT-biased genotypes of *O. sinensis* in the fully and semi-ejected ascospores

Li et al. [31] reported the detection of Genotype #13 in semi-ejected ascospores but not in fully

ejected ascospores and the detection of Genotype #14 in fully ejected ascospores but not in semi-ejected

ascospores. These 2 offspring genotypes of *O. sinensis* were characterized by large DNA segment

reciprocal substitutions and genetic material recombination between 2 parental fungi (Group A *H. sinensis*

and an AB067719-type Group E fungus [7,9–11,31]). The ITS1-5.8S-ITS2 sequences of Genotypes #13–

14 were alternatively segmentally inherited from their parental fungi [31]. The segment sequences of

Genotypes #13–14 are either completely identical to or far different from those of Genotype #1, and the

primer pairs for the first-step PCR listed in Table 1 are unfavorable for amplifying the ITS sequences of

Genotypes #13–14. Thus, the genotyping study presented herein does not cover the analysis of GC-biased

Genotypes #13–14.

Fig 6.1 shows a mass spectrum of extension primer 067721-477 (*cf*. Table 2) for fully ejected

ascospores. Peak G represents GC-biased Genotype #1 (*cf.* Fig 1). Peak A represents transition point
mutation AT-biased genotypes. Peak C was absent. Peak T had a very low intensity, representing a
transversion point mutation genotype of unknown upstream and downstream sequences. The primers
employed for the first-step PCR preferentially amplified the sequences of GC-biased genotypes, producing
an amplicon pool containing sequences of GC-biased genotypes in greater quantities than the AT-biased
sequences. The amplicon pool was used as the template for the second-step PCR, showing the higher-
intensity Peak G and the lower-intensity Peak A. The G:A and G:T ratios were 2.5 and 15.0, respectively
(Table 5).

Fig 6.2 shows a mass spectrum for the semi-ejected ascospores with a similar MS topology to that
shown in Fig 6.1. Peak G represents GC-biased Genotype #1, and Peak A represents AT-biased genotypes.
Peak C was absent, and Peak T had a very low intensity. There were no significant changes in the intensity
ratios of G:A (2.2) and G:T (13.8) relative to those for the fully ejected ascospores (Table 5).

Fig 6. MALDI-TOF mass spectra of extension primer 067721-477 to distinguish between GC- and AT-
genotypes in fully and semi-ejected ascospores. The extension reaction proceeded from extension primer
067721-477 toward the SNP at position 477 in the AB067721 sequence (*cf.* Fig 1 for the location). In the allelic
peaks, "C" denotes the primer with an extended cytosine; "A" indicates an extended adenine; "G" represents an
extended guanine; and "T" represents an extended thymine. Panel 6.1 shows the mass spectrum of fully ejected
ascospores. Panel 6.2 shows the mass spectrum of semi-ejected ascospores.

Table 5. Mass intensity ratios of the SNP allelic peaks of transition and transversion mutant genotypes in fully and semi-ejected ascospores

Extension primer	Allelic ratio	Intensity ratio	
		Fully ejected ascospores	**Semi-ejected ascospores**
		(*cf.* Fig 6.1)	(*cf.* Fig 6.2)
067721-477	G:A	**2.5** (7.5÷3.0)	**2.2** (↔; 6.7÷3.2)
	G:C	—	—
	G:T	**15.0** (7.5÷0.5)	**13.8** (↔; 6.9÷0.5)

Note: Peak G represents GC-biased Genotype #1 *H. sinensis*; Peak A indicates AT-biased genotypes of *O. sinensis* (*cf.* Fig 1). Peaks C and T denote 2 transversion mutation genotypes of unknown upstream and downstream sequences. "↔" indicates no significant change (within 20% variation) compared to the intensity ratios for fully ejected ascospores. "—" means that one of the allelic peaks was missing, and no ratio could be calculated.

The extension primer 067721-531 (*cf.* Table 2) was also used to distinguish between the GC- and AT-biased genotypes of *O. sinensis* in the fully and semi-ejected ascospores. Mass spectra (not shown) showed the same topologies of Peaks A and G in both panels of Fig 6, except for the absence of both Peaks C and T representing transversion alleles. There was no meaningful change in the G:A intensity ratio (Supplementary S5 Table).

S5 Table. Mass intensity ratios of the SNP allelic peaks of transition and transversion mutant genotypes in fully and semi-ejected ascospores.

Extension primer	Allelic ratio	Intensity ratio	
		Fully ejected ascospores (Mass spectrum not shown)	Semi-ejected ascospores (Mass spectrum not shown)
067721-531	G:A	18.3 (5.5÷0.3)	16.0 (↔; 8.0÷0.5)
	G:C	—	—
	G:T	—	—

Note: Peak G represents GC-biased Genotype #1 *H. sinensis*; Peak A indicates AT-biased genotypes of *O. sinensis* (*cf.* Fig 1). Peaks C and T denote 2 transversion mutation genotypes of unknown upstream and downstream sequences. "↔" indicates no significant change (within 20% variation) compared to the intensity ratios for fully ejected ascospores. "—" means that one of the allelic peaks was missing, and no ratio could be calculated.

SNP genotyping analysis to distinguish between the AT-biased genotypes of

O. sinensis in fully and semi-ejected ascospores

Li et al. [31] reported the absence of genotypes of AT-biased Cluster-B (Genotypes #4 and #15; *cf.* Fig 2) in the 2 types of ascospores. SNP genotyping using extension primer 067740-328 (reverse complement) (*cf.* Table 2) confirmed that both mass spectra (not shown) for the fully and semi-ejected ascospores showed a single Peak T of high intensity, representing Genotypes #5–6 and #16 of AT-biased Cluster-A (*cf.* Fig 2) [31]. Peaks C (Genotype #4 of AT-biased Cluster-B) and G (a transversion allele of unknown upstream and downstream sequences) are nearly flat. Peak A with very low intensities indicated the co-occurrence of a transversion mutation genotype of unknown upstream and downstream sequences in both ascosporic samples. The increase in the T:A ratio from 14.0 in fully ejected ascospores to 22.0 (Supplementary S6 Table) in semi-ejected ascospores was due to the considerably increased intensity of Peak T, the genotypes of AT-biased Cluster-A.

S6 Table. Mass intensity ratios for the SNP allelic peaks for multiple transition and transversion mutation genotypes in the fully and semi-ejected ascospores.

Extension primer	Allelic ratio	Intensity ratio	
		Fully ejected ascospores (Mass spectrum not shown)	Semi-ejected ascospores (Mass spectrum not shown)
067740-328	T:C	—	—
	T:A	**14.0** (42.0÷3.0)	**22.0** (↑; 66÷3.0)
	T:G	—	—

Note: Peak C represents AT-biased Genotype #5 of *O. sinensis,* and Peak T indicates other AT-biased genotypes of *O. sinensis* (*cf.* Fig 1). Peaks G and A denote 2 transversion mutation genotypes of unknown upstream and downstream sequences. "↑" indicates a significant increase in the intensity ratio less than two-fold compared to that of the fully ejected ascospores. "—" means missing one of the allele peaks, and no ratio could be calculated.

Fig 7.1 shows a mass spectrum obtained using extension primer 067740-324 (*cf.* Table 2) from the fully ejected ascospores. Peak C represented AT-biased Genotype #5 (*cf.* Fig 1), while AT-biased Genotype #17, with the same extended cytosine, was absent in the ascospores based on the use of genotype-specific primers and cloning-based amplicon sequencing [31]. Peak T represented AT-biased Genotypes #6 and #16 of AT-biased Cluster-A, while Genotypes #4 and #15 of AT-biased Cluster-B, with the same extended thymine, were absent in the ascospores [31]. The lower intensity of Peak C and the higher intensity of

673 Peak T resulted in a C:T ratio of 0.69 (Table 6). Peaks A and G represent extremely low-intensity

674 transversion point mutation genotypes of unknown upstream and downstream sequences, with C:A and

675 C:G ratios of 9.06 and 15.4, respectively (Table 6).

Fig 7. MALDI-TOF mass spectra of extension primer 067740-324 to distinguish between AT-biased genotypes in the fully and semi-ejected ascospores. The extension reaction proceeded from extension primer 067740-324 toward the SNP at position 324 in the AB067740 sequence (*cf*. Fig 1 for the location). In the allelic peaks, "C" denotes the primer with an extended cytosine; "A" indicates an extended adenine; "G" represents an extended guanine; and "T" indicates an extended thymine. Panel 7.1 shows the mass spectrum of fully ejected ascospores. Panel 7.2 shows the mass spectrum of semi-ejected ascospores. Panel 7.3 shows the overlapping mass spectra for Panels 7.1 (black tracing) and 7.2 (red tracing) with alignment of the horizontal and vertical axes of the panels.

Fig 7.2 shows a mass spectrum for semi-ejected ascospores. The intensity of Peak C (Genotype #5)

was elevated more than two-fold, from 15.4 in fully ejected ascospores to 34.8 in semi-ejected ascospores,

as visualized in Fig 7.3 after adjusting and aligning the vertical axis for intensities for the overlapping

mass spectra of Figs 7.1 and 7.2. Peak T (Genotypes #6 and #16) was slightly elevated from 22.3 to 26.8

(Figs 7.1–7.2). The asynchronous, disproportional increases in intensity (Fig 7.3) resulted in an elevation

of the C:T ratio from 0.69 to 1.30 from the fully ejected ascospores to the semi-ejected ascospores (Table

6). The intensities of Peaks A and G were slightly increased. The C:A and C:G intensity ratios were

increased from 9.06 to 14.5 and from 15.4 to 18.3, respectively (Table 6).

Table 6. Mass intensity ratios for the SNP allelic peaks for multiple transition and transversion mutation

genotypes in the fully and semi-ejected ascospores

Extension primer	Allelic ratio	Intensity ratio	
		Fully ejected ascospores (*cf*. Fig 7.1)	**Semi-ejected ascospores** (*cf*. Fig 7.2)
067740-324	C:T	**0.69** (15.4÷22.3)	**1.30** (↑; 34.8÷26.8)
	C:A	**9.06** (15.4÷1.7)	**14.5** (↑; 34.8÷2.4)
	C:G	**15.4** (15.4÷1.0)	**18.3** (↑; 34.8÷1.9)

Note: In the 067740-328 primer section (upper panel), Peak T represents AT-biased Genotype #4 of *O. sinensis,* and
Peak T indicates other AT-biased genotypes of *O. sinensis* (*cf*. Fig 1). In the 067740-324 primer section (lower panel),
Peak C represents AT-biased Genotype #5 of *O. sinensis,* and Peak T indicates other AT-biased genotypes of *O.*
sinensis (*cf*. Fig 1). Peaks G and A in both panels denote 2 transversion mutation genotypes of unknown upstream
and downstream sequences. "↑" indicates a significant increase (less than two-fold) in the intensity ratio compared to
that of the fully ejected ascospores. "—" means missing one of the allele peaks, and no ratio could be calculated.

Discussion

Biochip-based MassARRAY MS SNP genotyping technique for semi-quantitative analysis

Natural *Cordyceps sinensis* is an integrated microecosystem of the insect-fungal complex [6–7], and

its fungal members are differentially alternated in different combinations and play different biological

roles in its complex lifecycle (personal communication with Prof. Liang Z-Q). Along with >90 fungal

species from at least 37 genera [15–16,30], 17 genotypes of *O. sinensis* of genetically and phylogenetically

distinct (*cf*. Figs 1–2) have been identified, which differentially co-occur in the stroma and caterpillar body

[6–7,13,19–23,48], SFP (with ascocarps) and ascospores of natural *Cordyceps sinensis* [31,49] or in entire

Cordyceps sinensis [36–37,47], as previously summarized [6–11]. The nature of *Cordyceps sinensis*

causes technical difficulties and debates in detection of fungal members and in monitoring their dynamic changes during development and maturation of the natural product.

The stroma and caterpillar body of *Cordyceps sinensis* have been examined using the Southern blotting technique, and the natural abundance (without DNA amplification) of GC- and AT-biased genotypes of *O. sinensis* have been found to exhibit dynamic alterations in an asynchronized and disproportional manner in the compartments of natural *Cordyceps sinensis* during maturation [19,23]. Although Southern blotting analysis is not believed to be supersensitive, it is able to demonstrate the natural abundance of DNA species in test materials, reflecting the natural biomasses of the target DNA moieties. However, the Southern blotting technique can only detect *O. sinensis* genotypes as GC- and AT-biased groups after the samples are pre-treated with endonucleases [19,23], not individually. On the contrary, PCR-based techniques are considered sensitive molecular approaches, they may lead to disproportional exponential amplifications and inaccurate quantifications of various metagenomic components in complex DNA samples because of the involvement of multiple technical factors influencing the PCR process and sometimes cause failure of DNA amplification and sequencing for various reasons [8]. Restriction fragment length polymorphism (RFLP) techniques combined with cloning-based amplicon sequencing has also been employed to monitor the dynamic changes of GC-biased Genotypes #1–2 of *O. sinensis* during *Cordyceps sinensis* maturation [19–20]. This technique has limitations in quantitative analysis of multiple *O. sinensis* genotypes. Technical difficulties are encountered when using quantitative real-time PCR (qrPCR) to quantify and monitor the dynamic changes of metagenomic members in multi-fungal natural products, such as *Cordyceps sinensis*, because considerable sequence differences of fungi prevent balanced design of primers and probes for unbiased amplifications of the sequences of target intrinsic fungi. The current study used biochip-based MassARRAY MOLDI-TOF mass spectrometry SNP genotyping to examine teleomorphic genotypes of *O. sinensis* and monitor the dynamic alterations of the individual genotypes in the SFPs and ascospores of natural *Cordyceps sinensis*. This technique was used previously to monitor the dynamic alterations of coexisting *O. sinensis* genotypes in the stromata and caterpillar bodies of *Cordyceps sinensis* during maturation [19,21–23].

We designed two types of PCR-based experiments in this research project: (1) qualitative examinations using multiple sets of species- and genotype-specific primers for touch-down PCR and cloning-based amplicon sequencing with collection and sequencing of more than 30 white colonies to profile the genotypic components of the metagenomic DNA extracted from the immature and mature

stromata, SFPs with ascocarps, and 2 types of ascospores of natural *Cordyceps sinensis*, as presented in the companion paper [31]; and (2) a semiquantitative analysis using SNP mass spectrometry genotyping to monitor the dynamic alterations of the various genotypes in the SFPs densely covered with numerous ascocarps prior to and after ascospore ejection and of developmental failure and in the 2 types of ascospores, as presented herein.

Secondary structures/conformation in ITS1-5.8S-ITS2 sequences of *O. sinensis* genotypes have been recognized previously and greatly impact and even perplex PCR amplification and amplicon sequencing, often causing false negative results and consequently incorrect conclusions [8–11,18,49]. MassARRAY SNP MALDI-TOF MS genotyping represents a useful technique to overcome the shortcomings of regular PCR identification [54]. It is a two-step PCR-mass spectrometry technique involving a 384-well biochip [19–22,54,58]. Sequences containing various SNP alleles of different genotypes may not be amplified at the same rate because of (1) different amounts of DNA templates (*i.e.*, the amplicons obtained from the first-step PCR) in the samples; (2) variable DNA sequences and related secondary structures; (3) different primer binding rates and DNA chain extension rates, causing the MS intensities of the SNP alleles to not directly reflect the copy numbers of the genotypic sequences; and (4) perhaps other reasons in the natural world [8]. However, under the same PCR conditions using the same pair of primers in the first-step PCR, the same extension primer in the second-step PCR, and the same batch of reaction buffer run according to the same PCR settings and extension reaction on a single 384-well chip, the sequences of genotypes will be amplified at comparable amplification rates from one reaction well to another in a single chip.

Although direct quantitative comparisons of the intensities of the same allelic peaks in different mass spectra may not be accurate, it is reasonable to select the intensity of one SNP allelic peak in the mass spectra as the internal reference for intensity ratio computation (Tables 3–6 and Supplementary S3–S6 Tables), which provides legitimate information for semiquantitative comparisons of the relative abundances of genotypes in different mass spectra and semiquantitative monitoring of the alterations of the genotypes among the study samples. Such semiquantitative comparisons of intensity ratios are logical and valid within a systematic error, reflecting the relative alterations of GC- and AT-biased genotypes of *O. sinensis* and the transversion mutation genotypes of unknown upstream and downstream sequences in the SFPs prior to and after ascospore ejection and of developmental failure and in fully and semi-ejected ascospores collected from the same *Cordyceps sinensis* specimens.

Dynamic alterations of the genotypes of *O. sinensis*

Southern blotting, restriction fragment length polymorphism (RFLP) analysis combined with amplicon sequencing with or without amplicon cloning, SNP genotyping, and other molecular examinations demonstrated that GC-biased Genotype #1 *H. sinensis* was never the dominant fungal species in the *Cordyceps sinensis* stroma and that AT-biased genotypes were the dominant species [7,9–11,19–23,31]. Li et al. [49] proposed that all AT-biased genotypes are "ITS pseudogenes … in a single genome" of Genotype #1 *H. sinensis*. However, the 5 genome assemblies (ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, and NGJJ00000000) of *H. sinensis* Strains Co18, IOZ07, 1229, ZJB12195, and CC1406-203, respectively [41–45], do not contain the sequences of Genotypes #2–17, which belong to the genomes of independent *O. sinensis* fungi [6–11,13,19–23,31]. Multiple repetitive ITS copies were identified in the genome assemblies JAAVMX000000000 and NGJJ00000000 of Genotype #1 *H. sinensis* Strains IOZ07 and CC1406-203, respectively (*cf*. Fig 2) [44–45]. These multiple genomic ITS repeats are GC-biased and feature multiple scattered insertion/deletion and transversion point mutations, which might not be induced through RIP mutagenesis that theoretically causes cytosine-to-thymine transition mutations [11,57]. The ML phylogenetic tree shown in Fig 2 of the current paper and a Bayesian majority rule consensus tree shown in Fig 2 of [57] demonstrated that all sequences of repetitive ITS copies in JAAVMX000000000 and NGJJ00000000 were clustered into the GC-biased genotype clade along with GC-biased genotypes, far distant from the AT-biased genotype clade. The dynamic alterations of GC- and AT-biased genotypes of *O. sinensis* in an asynchronous, disproportional manner observed in this study (*cf*. Figs 4–7 and Supplementary S1–S4 Figs) and previous studies [19–23] further confirmed the genomic independence of the multiple genotypes, providing evidence of the coexistence of interindividual *O. sinensis* fungi [13,19–23]. These results thus disproved the hypothesis of "ITS pseudogenes … in a single genome", which was based on the misinterpretation of genomic residency data for the multiple AT-biased mutant genotypes in the genome of GC-biased Genotype #1 *H. sinensis* [6–11,13,31].

In addition to the companion paper [31] that reported the differential co-occurrence of multiple genotypes of *O. sinensis* in different combinations in the immature and mature stroma, SFPs with ascocarps, and ascospores of natural *Cordyceps sinensis* using qualitative molecular techniques, the current paper also revealed the dynamic alterations of GC- and AT-biased genotypes in the SFPs (with

ascocarps) and ascospores through a semiquantitative SNP genotyping approach (*cf.* Figs 4–7 and

Supplementary S1-S4 Figs). For instance, Supplementary S2.1 Fig demonstrates the nearly identical

intensities of Peaks A and G, representing AT- and GC- biased genotypes of *O. sinensis*, in the pre-ejection

SFP, and Supplementary S2.2 Fig shows a significant reduction of Peak A (AT-biased genotypes) after

ascospore ejection, resulting in a dramatic increase in the G:A intensity ratio (*cf.* Supplementary S3 Table).

Figs 7.1–7.3 demonstrate that Peak C (AT-biased Genotype #5) with slightly lower intensity (than that for

Peak T) in the fully ejected ascospores increased more than two-fold in its peak height in the semi-ejected

ascospores collected from the same pieces of natural *Cordyceps sinensis* specimens, compared with Peak

T (AT-biased Genotypes #6 and #16) with very high intensity in the fully ejected ascospores but only a

disproportional minimal increase in the semi-ejected ascospores (*cf.* Fig 7 and Table 6). Genotypes #5–6

and #16 were phylogenetically clustered into AT-biased Cluster-A (*cf.* Fig 2). Significant changes in the

intensities of AT-biased genotypes are also shown in other figures and tables, indicating that the sequences

of AT-biased genotypes belong to the independent genomes of interindividual *O. sinensis* fungi.

The dynamic alterations of the intensities of the allelic peaks suggested the genomic independence

of GC- and AT-biased *O. sinensis* genotypes, inconsistent with the hypothesis that multiple AT-biased "ITS

pseudogene" components of *O. sinensis* belong to "a single genome" of Genotype #1 *H. sinensis* [49].

Notably, the intensities of the allelic peaks and their alterations shown in Figs 4 and 6 and Supplementary

S1–S2 Figs do not necessarily reflect the natural biomasses of the GC- and AT-biased genotypes of *O.*

sinensis because the primers employed for the first-step PCR preferentially amplified GC-biased

genotypes over AT-biased genotypes, producing a greater amount of GC-biased amplicons that served as

the templates for primer extension reactions in the second-step PCR.

Mass spectra obtained using extension primers 067721-447 and 067721-531 demonstrated the co-

occurrence of GC- and AT-biased *O. sinensis* genotypes in fully and semi-ejected ascospores, resulting in

essentially unchanged G:A ratios (*cf.* Fig 6, Table 5, and Supplementary S5 Table).

GC-biased Genotype #1 *H. sinensis* is postulated to be an anamorph of *O. sinensis* [32], although it

may not be the sole anamorph [11,31,49,59–60]. In addition, GC-biased Genotype #1 was reported to be

the sole teleomorph of *O. sinensis* in natural *Cordyceps sinensis* [32,40], and AT-biased Genotype #4 was

the sole teleomorph of *O. sinensis* in cultivated *Cordyceps sinensis* [40]. Our findings presented in this

paper and the companion paper [31] indicate that multiple co-occurring teleomorphic AT-biased genotypes

may play critical symbiotic roles in the development and maturation of SFPs with ascocarps and

ascospores in the sexual life stage of *Cordyceps sinensis*. Without certain AT-biased *O. sinensis* genotypes

in various combinations, ascocarps may not be able to normally develop and mature to support the

development, maturation, and ejection of the *Cordyceps sinensis* ascospores (*cf*. Figs 3–5 and

Supplementary S1–S4 Figs), especially lacking high intensities of Genotype #6 (AT-biased Cluster-A) and

Genotype #15 (AT-biased Cluster-B) in the SFP of developmental failure, as shown in Supplementary S4

Fig. We also noticed the coexistence of transversion mutated alleles of unknown upstream and downstream

sequences that were altered in the SFPs and ascospores (Figs 4–7, and Supplementary S1–S4 Figs; Tables

3–6, and Supplementary S4 and S6 Tables), suggesting that these co-occurring transversion genotypes

may also participate symbiotically in the development and maturation of ascocarps and ascospores.

Stratification strategy for genotyping AT-biased genotypes

The stratification genotyping strategy used in this study profiled in a step-by-step manner the

multiple genotypic components of AT-biased Clusters A and B in the *Cordyceps sinensis* SFPs. The first

step of stratifying assays used extension primer 067740-328. The mass spectra of the SFPs prior to and

after ascospore ejection and of developmental failure demonstrated the existence of an extremely small

quantity of Genotype #4 of AT-biased Cluster-B that is represented by Peak C in Supplementary S3.1–

S3.3 Figs, in addition to large amounts of Genotypes #5–6 of AT-biased Cluster-A and Genotype #15 of

AT-biased Cluster-B. In contrast to the dominance of Genotype #4 of AT-biased Cluster-B in the stroma

of immature *Cordyceps sinensis* [21–22], in its asexual growth stage, the abundance of Genotype #4

dramatically declined in the stroma of mature *Cordyceps sinensis* [19,21–22] and retained a very low level

in the SFPs (with ascocarps) in the sexual stage of the *Cordyceps sinensis* lifecycle [31].

The second step of the stratification analysis involved the extension primer 067744-324. The mass

spectrum (*cf*. Fig 5.1) demonstrated a higher intensity of Peak T (sum of a minimal quantity of Genotype

#4 and a large amount of Genotypes #6 and #15) and a slightly lower intensity of Peak C (Genotype #5)

in the pre-ejection SFP, resulting in a C:T ratio of 0.71 (*cf*. Table 4). In contrast to retaining its high

intensity of Peak C (Genotype #5 of AT-biased Cluster-A) in the post-ejection SFP, Peak T was

significantly attenuated in the SFPs after ascospore ejection (*cf*. Figs 5.2), representing a dramatic

reduction in Genotype #6 (AT-biased Cluster-A) and Genotype #15 (AT-biased Cluster-B) along with the

extremely small quantity of component Genotype #4 in the *Cordyceps sinensis* SFP after ascospore

ejection; this caused a significant increase in the C:T ratio (*cf*. Table 4). Peak T (Genotypes #6 and #15,

and minimal amount of Genotype #4) also showed extremely low intensity in the SFP of developmental

failure (*cf*. Supplementary S4 Fig), resulting in an extremely high C:T ratio (*cf*. Table 4). These findings

indicated that AT-biased Genotypes #6 and #15 may play critical roles in the development and maturation

of *Cordyceps sinensis* ascocarps and that insufficient Genotypes #6 and #15 in the SFP may cause

developmental failure of *Cordyceps sinensis* ascocarps (comparing Fig 5.1 and Supplementary S4 Fig),

while AT-biased Genotype #4 may play an important role in the early asexual developmental stage of

natural *Cordyceps sinensis* [21–22,31]. Although both Genotypes #4 and #15 were phylogenetically

clustered into AT-biased Cluster-B, they play distinct symbiotic roles in different maturation stages of

Cordyceps sinensis.

Genotype #6 was clustered into AT-biased Cluster-A, but its phylogenetic leaf was closer to Cluster-

B than other Cluster-A genotypes (*cf*. Fig 2), indicating that Genotype #6 may play different symbiotic

roles from Genotypes #5 and #16 of AT-biased Cluster-A. We also noticed that all Genotype #6 sequences

uploaded in GenBank are short, missing small segments at the 5' end of ITS1 and the 3' end of ITS2,

indicating that unique secondary conformation may exist in these segments that prevents PCR detection

of Genotype #6 when using the so called "universal" primers (primers *ITS1–ITS5*) and that design of

specific primers may need to detect Genotype #6 sequence.

Genotypes #4 and #15 of AT-biased Cluster-B were absent in the ascospores of *Cordyceps sinensis*

[31,49], although Wei et al. [40] reported that Genotype #4 was the sole teleomorph of *O. sinensis* in

cultivated *Cordyceps sinensis*. Genotype #5 (Peak C in Fig 7) and Genotypes #6 and #16 (components of

Peak T in Fig 7) were phylogenetically clustered into AT-biased Cluster-A (*cf*. Fig 2) and play symbiotic

roles in assisting the function of GC-biased Genotype #1 *H. sinensis* in the sexual life stage of natural

Cordyceps sinensis. Fungal symbiosis is critical for the development, maturation, and ejection of

Cordyceps sinensis ascospores.

This paper and its companion paper [31] demonstrated the symbiotic co-occurrence of GC-biased

Genotype #1 and AT-biased Genotypes #5–6 and #16 of *O. sinensis* in *Cordyceps sinensis* ascospores.

However, Li et al. [49] reported the detection of GC-biased Genotype #1 and AT-biased Genotype #5 in

8 of 15 clones in 25-day cultures generated from mono-ascospores. Apparently, the culture-dependent

study by Li et al. [49] overlooked AT-biased Genotypes #6 and #16 as well as GC-biased Genotype #14

in fully ejected ascospores [31], which might be attributed to either nonculturability of these overlooked

genotypes of *O. sinensis* fungi under the 25-day *in vitro* culture conditions set by [49] or a failure of

detection due to an inappropriate design of the molecular methodology. As shown in Fig 1, the sequences
of AT-biased Genotypes #6 (EU555436) and #16 (KT232019) are both short compared with that of
Genotype #1 *H. sinensis*; Genotype #6 EU555436 is especially short, lacking short DNA segments close
to its 5' and 3' ends. There have been no reports of the detection of Genotypes #6 and #16 using "universal"
primers. The molecular features (secondary structure/conformation) of these genotypes may prevent their
detection through PCR when using "universal" primers [6–11,19,21].

Li et al. [31] reported the detection of Genotype #13 in semi-ejected ascospores and Genotype #14
in fully ejected ascospores. These 2 GC-biased genetically variant genotypes feature large DNA segment
reciprocal substitutions and genetic material recombination between the 2 parental fungi (Group-A
Genotype #1 *H. sinensis* and a Group-E AB067719-type fungus), regardless of whether Genotypes #13–
14 are the offspring derived from fungal hybridization or parasexual reproduction [7,11,59]. Using
genotype-specific primers and cloning-based amplicon sequencing techniques is an appropriate approach
to identify these 2 offspring genotypes, as well as other AT-biased genotypes [31]. However, the
MassARRAY SNP mass spectrometry genotyping technique using the primer pairs and extension primers
designed in this study is not applicable to identify ascosporic Genotypes #13–14 because the ITS1-5.8S-
ITS2 sequences of Genotypes #13–14 were alternatively segmentally inherited from their parental fungi
[31], either completely identical to or far different from those of Genotype #1 and the primer pairs for the
first-step PCR (*cf*. Fig 1, Table 1). Thus, the semiquantitative study presented in this paper does not cover
the analysis of GC-biased Genotypes #13–14.

True causative fungus/fungi of natural *Cordyceps sinensis*

Zhang et al. [63] summarized nearly 40 years of mycological experience involving the failed
induction of the fruiting body and ascospore production in research-oriented laboratory settings, either in
fungal cultures or after infecting insects with fungal inoculants. Hu et al. [41] reported success in inducing
infection-mortality-mummification in 40 larvae of *Hepialus* sp. in an experiment using a mycelial mixture
of anamorphic *H. sinensis* Strains Co18 and QH195-2 as inoculants but failed to induce the production of
fruiting bodies.

Li et al. [58] demonstrated that a mixture of wild-type Strains CH1 and CH2 induced a favorable
infection-mortality-mummification rate of 55.2±4.4% in 100 larvae of *Hepialus armoricanus*, which
represents a 15–39-fold greater inoculation potency (P<0.001) than the rates (1.4–3.5%) achieved after

inoculation with the conidia or mycelia of *H. sinensis* or pure ascospores of *Cordyceps sinensis* (n=100

larvae in each inoculation experiment). The wild-type CH1 and CH2 strains exhibit *H. sinensis*-like

morphological and growth characteristics but contain GC-biased Genotype #1 *H. sinensis*, AT-biased

Genotypes #4–6 of *O. sinensis*, and *Paecilomyces hepiali*.

Wei et al. [40] reported an industrial artificial inoculation project and demonstrated a species

contradiction between the anamorphic inoculant GC-biased Genotype #1 *H. sinensis* (Strains 130508-KD-

2B, 20110514 and H01-20140924-03) and the sole teleomorph of AT-biased Genotype #4 of *O. sinensis*

in cultivated *Cordyceps sinensis*. In addition to reporting the sole teleomorph of AT-biased Genotype #4

in cultivated *Cordyceps sinensis*, Wei et al. [40] also reported the sole teleomorph of GC-biased Genotype

#1 in natural *Cordyceps sinensis*, reporting at least 2 teleomorphs of *O. sinensis*.

We reported the co-occurrence and dynamic alterations of multiple genotypes of *O. sinensis* in the

teleomorphic SFPs and ascospores of natural *Cordyceps sinensis* in this study and the companion study

[31], suggesting that Genotype #1 is not the sole teleomorph of *O. sinensis* and that Genotype #1 *H.*

sinensis may be among the true causative fungi for natural and cultivated *Cordyceps sinensis*. Our findings

of multiple teleomorphs of *O. sinensis* may explain the 40-year failure of induction of the fruiting body

and ascospore production in research-orientated academic settings, as summarized by Zhang et al. [63].

The evidence from the studies [31,40–41,58,63] suggested that the synergistic inoculation of multiple

fungal species is required to initiate the development of the stromal primordia and fruiting bodies of

natural and cultivated *Cordyceps sinensis* and to subsequently achieve the biological transition from the

initial asexual growth to sexual reproduction in the later maturational stages and complete the *O. sinensis*

infection cycle and the natural *Cordyceps sinensis* lifecycle.

Inability of self-fertilization reproductive behavior in *H. sinensis*

The reproductive behavior of ascomycetes is controlled by transcription factors encoded at the

mating-type (*MAT*) locus [60–62]. The expression of mating-type genes is controlled at the genomic,

epigenetic, transcriptomic, post-transcriptive, translational, post-translational, and proteomic levels [59].

Zhang & Zhang [64] reported the differential occurrence of mating-type genes of *MAT1-1* and *MAT1-2*

idiomorphs in many *Cordyceps sinensis* specimens and proposed a facultative hybridization hypothesis

for *O. sinensis*. Li et al. [59] analyzed transcriptome assemblies of *H. sinensis* strains focusing on the

mating-type genes of *MAT1-1* and *MAT1-2* idiomorphs and reported the differential transcription of mating-type genes in Genotype #1 *H. sinensis* strains, in addition to the differential occurrence of the mating-type genes in 237 *H. sinensis* strains.

Liu et al. [65] reported the transcriptome assembly GCQL00000000 of the *H. sinensis* Strain L0106, in which the MAT1-1-1 transcript was absent, but the MAT1-2-1 transcript was present [59]. Although Bushley et al. [50] reported the detection of transcripts of mating-type genes of both *MAT1-1* and *MAT1-2* idiomorphs in *H. sinensis* Strain 1229, the MAT1-2-1 transcript contained nonspliced intron I that contains 3 stop codons. The MAT1-2-1 transcript would then produce a largely truncated protein, lacking the major portion of the protein encoded by exons II and III of MAT1-2-1 gene [59], inevitably resulting in biological dysfunction of the truncated MAT1-2-1 protein. Thus, the evidence of the differential transcription of mating-type genes indicated the self-sterilization of Genotype #1 *H. sinensis* [59] and the inability to carry out self-fertilization under homothallic or pseudohomothallic reproduction [41,50].

The evidence of genetic, transcriptional, and coupled transcriptional-translational controls of *O. sinensis* sexual reproduction indicated the requirement of mating partners to carry out sexual reproduction in the lifecycle of natural *Cordyceps sinensis*. The potential sexual partners in physiological heterothallism may be coexisting intraspecific GC-biased Genotypes #1 of *O. sinensis*, monoecious or dioecious [11,59]. For instance, *H. sinensis* Strains 1229 and L0106 may be able to produce complementary mating proteins and potentially act as mating partners in physiological heterothallic reproduction [11,41,59,65].

The potential mating partners may also be heterospecific fungal species co-colonized in natural and cultivated *Cordyceps sinensis* if the fungi are able to break through the interspecific reproductive isolation barrier to achieve hybridization. Notably, the taxonomic positions of GC-biased Genotypes #2–3 and #7–14 and AT-biased Genotypes #4–6 and #15–17 have not been determined, probably due to their *H. sinensis*-like morphological and growth characteristics and nonculturability *in vitro* leading to technical difficulties in fungal isolation and purification [7,9–11,37,47]. However, they most likely belong to independent *O. sinensis* fungi because their sequences do not reside in the genome of Genotype #1 *H. sinensis* [6–11,41–46] and because of the phylogenetic analysis and conclusion by Stensrud et al. [55] that a large sequence variation in the conservative 5.8S gene "far exceeds what is normally observed in fungi … even at higher taxonomic levels (genera and family)." Studies have reported a close association of GC-biased Genotypes #1 and #2 and AT-biased genotypes of *O. sinensis* with dynamic genotypic alterations in the stroma of natural *Cordyceps sinensis* [5–11,19–23], especially in the SFPs and ascospores as coexisting teleomorphs

in the sexual life stages presented in this study and other studies [31,49]. Hopefully, the multiple genotypes

of *O. sinensis* reported in this study and the companion study [31] may potentially serve as sexual partners

for hybridization if they are finally determined to be independent fungal species. Coincidentally,

transcriptome assemblies of *H. sinensis* Strain L0106 and *P. hepiali* Strain Feng showed differential

transcription of mating-type genes that were complementary to each other [59,65–66]. Barseghyan et al.

[67] reported that *H. sinensis* and *Tolypocladium sinensis* were anamorphs of *O. sinensis*; however, the

mating-type genes and their transcripts for *T. sinensis* have not been documented.

These findings provide genetic and transcriptomic evidence for the possible outcrossing of *O.*

sinensis fungi and encourage further investigation of their reproductive physiology. In addition, Li et al.

[7,11,31,59] provided evidence of the potential hybridization or parasexual reproduction of Group-A *H.*

sinensis (Genotype #1 of *O. sinensis*) and a Group-E AB067719-type fungus in *Cordyceps sinensis*

ascospores, producing genetically variant offspring, Genotypes #13–14 of *O. sinensis*, each in the semi-

or fully ejected ascospores. The evidence of complex reproduction behaviors of *O. sinensis* supports the

facultative hybridization hypothesis for *O. sinensis* proposed by Zhang & Zhang [64].

Conclusions

We have demonstrated in this paper the histology of the SFPs (densely covered with numerous

ascocarps) prior to and after ascospore ejection and of developmental failure and differential cooccurrence

and dynamic alterations of multiple teleomorphic genotypes of *O. sinensis* in different combinations in

the SFPs and ascospores of natural *Cordyceps sinensis*. These findings in regard to the genital cells

(ascospores) and their production organ (the ascocarps that densely cover the SFPs) provided evidence of

the genomic independence of the *O. sinensis* genotypes, which do not reside in a single genome of

Genotype #1 *H. sinensis* but instead belong to interindividual fungi. Metagenome members, including

several genotypes of *O. sinensis*, play symbiotic roles in various developmental and maturational stages

in different compartments of natural *Cordyceps sinensis* to accomplish sexual reproduction of *O. sinensis*

during the lifecycle of natural *Cordyceps sinensis*.

1009

1010

Acknowledgments

The authors are grateful to Prof. Mu Zang, Prof. Zong-Qi Liang, Prof. Ru-Qin Dai, Prof. Ying-Lan Guo, Prof. Ping Zhu, Prof. Zhao-Lan Li, Prof. Yu-Guo Zheng, Dr. Jia-Gang Zhao and Dr. Yan-Jiao Zhou for consultations, and Prof. Xin Liu, Prof. Hai-Feng Xu, Ms. Shao-Li Ma, Ms. Ming Yang, Mr. Wei Chen, Mr. Tao-Ye Zheng, Mr. Jin-Jin Li, and Mr. Yu-Chun Zhou for their assistance.

[REFERENCES]

[1] Song L-R, Hong X, Ding X-L, Zang Z-Y. *Monographs of Modern Pharmacy of Chinese Traditional Medicine*. People′s Health Publishing House, Beijing 2001. pp. 733–737.

[2] Zhu J-S, Halpern GM, Jones K. The scientific rediscovery of an ancient Chinese herbal medicine: *Cordyceps sinensis*: Part I. J. Altern. Complem. Med. 1998a;4(3): 289–303. DOI: 10.1089/acm.1998.4.3-289

[3] Zhu J-S, Halpern GM, Jones K. The scientific rediscovery of an ancient Chinese herbal medicine: *Cordyceps sinensis*: Part II. J. Altern. Complem. Med. 1998b;4(4): 429–457. DOI: 10.1089/acm.1998.4.429

[4] Li C-L, Tan N-Z, Berger JL, Ferguson SB, Zhang Y, Prolla TA, et al. The Combined use of ageLOC whole-gene expression profiling technology and mouse lifespan test in anti-aging herbal product study. Proc. 2011 New TCM Prod. Innovat. Indust. Devel. Summit, Hangzhou, China, 2011, pp. 443–448.

[5] Tan N-Z, Berger JL, Zhang Y, Ferguson SB, Wu Z-M, Prolla TA, et al. The lifespan-prolonging effect of *Cordyceps sinensis* Cs-4 in normal mice and its molecular mechanisms. FASEB J. 2011;25(1): 599.1. https://doi.org/10.1096/fasebj.25.1_supplement.599.1

[6] Yao Y-S, Zhu J-S. Indiscriminative use of the Latin name for natural *Cordyceps sinensis* insect-fungi complex and multiple *Ophiocordyceps sinensis* fungi. Chin. J. Chin. Mater Med. 2016;41(7): 1361−1366. DOI: 10.4268/cjcmm20160733

[7] Li X-Z, Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. Molecular identification of *Ophiocordyceps sinensis* genotypes and the indiscriminate use of the Latin name for the multiple genotypes and the natural insect-fungi complex. Am J BioMed Sci. 2022;14(3): 115−135. doi:10.5099/aj220300115

[8] Zhu J-S, Wu J-Y. Genetic heterogeneity of natural *Cordyceps sinensis* with co-existence of
multiple fungi. Chin. J. Cell Biol. 2015;37(2): 284–298. DOI: 10.11844/cjcb.2015.02.0284

[9] Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. Molecular heterogeneity of natural *Cordyceps sinensis* with
multiple *Ophiocordyceps sinensis* fungi challenges the anamorph-teleomorph connection
hypotheses and implementation of the Amsterdam Declaration of IMA. Am. J. Biomed. Sci.
2016;8(2): 123−159. doi: 10.5099/aj160100123

[10] Zhu J-S, Li Y-L. A Precious Transitional Chinese Medicine, *Cordyceps sinensis*: Multiple
heterogeneous *Ophiocordyceps sinensis* in the insect-fungi complex. Lambert Academic
Publishing, Saarbrüchen. Germany 2017.

[11] Li Y-L, Li X-Z, Zhu J-S. *Cordyceps sinensis* Fungal Heterokaryons. Chemistry Industry Press Co.,
Beijing, China, 2022.

[12] Yang J-L, Xiao W, He H-H, Zhu H-X, Wang S-F, Cheng K-D, et al. Molecular phylogene
tic analysis of *Paecilomyces hepiali* and *Cordyceps sinensis*. Acta Pharmaceut. Sinica. 200
8;43(4): 421–426. DOI: 10.16438/j.0513-4870.2008.04.001

[13] Xiao W, Yang J-P, Zhu P, Cheng K-D, He H-X, Zhu H-X, et al. Non-support of species complex
hypothesis of *Cordyceps sinensis* by targeted rDNA-ITS sequence analysis. Mycosystema.
2009;28(6): 724–730.

[14] Zhu J-S, Guo Y-L, Yao Y-S, Zhou Y-J, Lu J-H, Qi Y, et al. Maturation of *Cordyceps sinensis*
associates with co-existence of *Hirsutella sinensis* and *Paecilomyces hepiali* DNA and dynamic
changes in fungal competitive proliferation predominance and chemical profiles. J. Fungal Res.
2007;5(4): 214–224.

[15] Zhang Y-J, Sun B-D, Zhang S, Wàngmǔ, Liu X-Z, Gong W-F. Mycobiotal investigation of
natural *Ophiocordyceps sinensis* based on culture-dependent investigation. Mycosistema. 20
10;29(4): 518–527. DOI: 10.13346/j.mycosystema.2010.04.012

[16] Xia F, Liu Y, Shen G-L, Guo L-X, Zhou X-W. Investigation and analysis of microbiological
communities in natural *Ophiocordyceps sinensis*. Can. J. Microbiol. 2015;61: 104–111. DOI:
10.1139/cjm-2014-0610

[17] He S-Q, Jin X-L, Luo J-C, Wang C-M, Wang S-R. Morphological characteristics and submerged
culture medium screening of a strain of *Geomyces pannorum* isolation from *Cordyceps sinensis*.

Chin. J. Grassland. 2010;32(z1): 70–75.

[18]   He S-Q, Jin X-L, Luo J-C, Wang C-M. Morphological characteristics and submerged cultur
e medium screening of *Pseudogymnoascus roseus*. Microbiol. Chin. 2011;38(9): 1371−1376.
DOI: 10.13344/j.microbiol.china.2011.09.016

[19]   Zhu J-S, Gao L, Li X-H, Yao Y-S, Zhou Y-J, Zhao J-Q, et al. Maturational alterations of oppositely
orientated rDNA and differential proliferations of CG:AT-biased genotypes of *Cordyceps sinensis*
fungi and *Paecilomyces hepiali* in natural *C. sinensis*. Am. J. Biomed. Sci. 2010;2(3): 217–238.
doi: 10.5099/aj100300217

[20]   Zhu J-S, Zhao J-G, Gao L, Li X-H, Zhao J-Q, Lu J-H. Dynamically altered expressions o
f at least 6 *Ophiocordyceps sinensis* mutants in the stroma of *Cordyceps sinensis*. J. Funga
l Res. 2012;10(2): 100–112. DOI: 10.13341/j.jfr.2012.02.007

[21]   Gao L, Li X-H, Zhao J-Q, Lu J-H, Zhu J-S. Detection of multiple *Ophiocordyceps sinensis*
mutants in premature stroma of *Cordyceps sinensis* by MassARRAY SNP MALDI-TOF mass
spectrum genotyping. J. Peking Univ. (Health Sci.). 2011;43(2): 259–266.

[22]   Gao L, Li X-H, Zhao J-Q, Lu J-H, Zhao J-G, Zhu J-S. Maturation of *Cordyceps sinensis* associates
with alterations of fungal expressions of multiple *Ophiocordyceps sinensis* mutants with transition
and transversion point mutations in stroma of *Cordyceps sinensis*. J. Peking Univ. (Health Sci.).
2012;44(3): 454–463.

[23]   Yao Y-S, Zhou Y-J, Gao L, Lu J-H, Zhu J-S. Dynamic alterations of the differential funga
l expressions of *Ophiocordyceps sinensis* and its mutant genotypes in stroma and caterpilla
r during maturation of natural *Cordyceps sinensis*. J. Fungal Res. 2011;9(1): 37–49,53. DO
I: 10.13341/j.jfr.2011.01.007

[24]   Xiao Y-Y, Cao Y-M, Zhang L, Li C-R. Study on Genetic Diversity of Anamorphic Isolates from
Different Sources of *Ophiocordyceps sinensis* by ISSR. J. Fungal Res. 2014;12(3): 154–159.

[25]   Yao Y-S, Gao L, Li Y-L, Ma S-L, Wu Z-M, Tan N-Z, et al. Amplicon density-weighted a
lgorithms analyze dissimilarity and dynamic alterations of RAPD polymorphisms in the inte
grated micro-ecosystem *Cordyceps sinensis*. J. Peking Univ. (Health Sci.). 2014;46(4): 618–
628. DOI: 10.13341/j.jfr.2014.0025

[26]   Ni L-Q, Yao Y-S, Gao L, Wu Z-M, Tan N-Z, Wu J-Y, et al. Density-weighted algorithms for

similarity computation and cluster tree construction in the RAPD analysis of a micro-ecosystem
Cordyceps sinensis. Am. J. Biomed. Sci. 2014;6(2): 82–104. doi: 10.5099/aj140200082

[27] Liu Y-N, Jin Y-S, Liu Z, Han Y-N. *Cordyceps sinensis* cDNA library construction and
identification. Chin. J. Informat. TCM 2006;13(9): 43−45.

[28] Dong Y-Z, Zhang L-J, Wu Z-M, Gao L, Yao Y-S, Tan N-Z, et al. Altered proteomic polymorphisms
in the caterpillar body and stroma of natural *Cordyceps sinensis* during maturation. PLoS ONE.
2014;9(10): e109083. doi: 10.1371/journal.pone.0109083

[29] Liang Z-Q, Han Y-F, Liang J-D, Dong X, Du W. Issues of concern in the studies of *Oph*
iocordyceps sinensis. Microbiol. Chin. 2010;37(11): 1692–1697. DOI: 10.13344/j.microbiol.c
hina.2010.11.023

[30] Jiang Y, Yao Y-J. A review for the debating studies on the anamorph of *Cordyceps sinensis*.
Mycosistema. 2003;22(1): 161–176.

[31] Li Y-L, Li X-Z, Yao Y-S, Wu Z-M, Lou Z-Q, Xie W-D, et al. Differential cooccurrence of multiple
genotypes of *Ophiocordyceps sinensis* in the stromata, stromal fertile portion (ascocarps) and
ascospores of natural *Cordyceps sinensis*. PLoS ONE. 2023;18(3): e0270776. Doi:
10.1371/journal.pone.0270776

[32] Wei X-L, Yin X-C, Guo Y-L, Shen N-Y, Wei J-C. Analyses of molecular systematics on *Cordyceps*
sinensis and its related taxa. Mycosystema. 2006;25(2): 192–202.

[33] Mo M-H, Chi S-Q, Zhang K-Q. Microcycle conidiation of *Cordyceps sinensis* and anamorph
isolation. Mycosystema. 2001;20: 482−485.

[34] Liu Z-Y, Liang Z-Q, Liu A-Y. Investigation on microcycle conidiation of ascospores and
conidiogenous structures of anamorph of *Cordyceps sinensis*. Guizhou Agricult. Sci. 2003;31(1):
3–5.

[35] Xiao Y-Y, Chen C, Dong J-F, Li C-R, Fan M-Z. Morphological observation of ascospores
of *Ophiocordyceps sinensis* and its anamorph in growth process. J. Anhui Agricult. Univ.
2011;38(4): 587–591. DOI: 10.13610/j.cnki.1672-352x.2011.04.022

[36] Zhao J, Wang N, Chen Y-Q, Li T-H, Qu L-H. Molecular identification for the asexual stage of
Cordyceps sinensis. Acta Sci. Nat. Univ. Sunyatseni. 1999;38(1): 122–123.

[37] Kinjo N, Zang M. Morphological and phylogenetic studies on *Cordyceps sinensis* distributed in
southwestern China. Mycoscience. 2001;42: 567–574.

[38] Liu Z-Y, Yao Y-J, Liang Z-Q, Liu A-Y, Pegler DN, Chase M-W. Molecular evidence for the
anamorph-teleomorph connection in *Cordyceps sinensis*. Mycol. Res. 2001;105(7): 827–832.

[39] Li Z-Z, Huang B, Li C-R, Fan M-Z. Molecular evidence for anamorph determination of *Cordyceps*
sinensis (Berk.) Sacc. Mycosystema. 2000;9(1): 60–64.

[40] Wei J-C, Wei X-L, Zheng W-F, Guo W, Liu R-D. Species identification and component d
etection of *Ophiocordyceps sinensis* cultivated by modern industry. Mycosystema. 2016;35
(4): 404–410. DOI: 10.13346/j.mycosystema.160045

[41] Hu X, Zhang Y-J, Xiao G-H, Zheng P, Xia Y-L, Zhang X-Y, et al. Genome survey uncovers the
secrets of sex and lifestyle in caterpillar fungus. Chin. Sci. Bull. 2013;58: 2846–2854.
https://doi.org/10.1007/s11434-013-5929-5

[42] Li Y, Hsiang T, Yang R-H, Hu X-D, Wang K, Wang W-J, et al. Comparison of different sequencing
and assembly strategies for a repeat-rich fungal genome, *Ophiocordyceps sinensis*. J. Microbiol.
Methods. 2016;128: 1−6. DOI: 10.1016/j.mimet.2016.06.025

[43] Jin L-Q, Xu Z-W, Zhang B, Yi M, Weng C-Y, Lin S, et al. Genome sequencing and analysis of
fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*. AMB Expr. 2020;10: 105. doi:
10.1186/s13568-020-01039-x

[44] Liu J, Guo L-N, Li Z-W, Zhou Z, Li Z, Li Q, et al. Genomic analyses reveal evolutionary and
geologic context for the plateau fungus *Ophiocordyceps sinensis*. Clin. Med. 2020;15: 107–119.
DOI: 10.1186/s13020-020-00365-3

[45] Shu R-H, Zhang J-H, Meng Q, Zhang H, Zhou G-L, Li M-M, et al. A new high-quality draft
genome assembly of the Chinese cordyceps *Ophiocordyceps sinensis*. Genome Biol. Evol.
2020;12(7): 1074−1079. doi: 10.1093/gbe/evaa112

[46] Zhang Y-J, Xu L-L, Zhang S, Liu X-Z, An Z-Q, Wàngmǔ, et al. Genetic diversity of
Ophiocordyceps sinensis, a medicinal fungus endemic to the Tibetan Plateau: implications for its
evolution and conservation. BMC Evol. Biol. 2009;9: 290. doi:10.1186/1471-2148-9-290

[47] Mao X-M, Zhao S-M, Cao L, Yan X, Han R-C. The morphology observation of anamorph of
Ophiocordyceps sinensis from different origins. J Environ. Entomol. 2013;35(3): 343–353.

[48] Chen C-S, Hseu R-S, Huang C-T. Quality control of *Cordyceps sinensis* teleomorph, anamorph,
and Its products. Chapter 12, *in* (Shoyama, Y., Ed.) *Quality Control of Herbal Medicines and*
Related Areas. InTech, Rijeka, Croatia, 2011. www.intechopen.com

[49] Li Y, Jiao L, Yao Y-J. Non-concerted ITS evolution in fungi, as revealed from the important
medicinal fungus *Ophiocordyceps sinensis*. Mol. Phylogenet. Evol. 2013;68: 373−379. doi:
10.1016/j.ympev.2013.04.010

[50] Bushley KE, Li Y, Wang W-J, Wang X-L, Jiao L, Spatafora JW, et al. Isolation of the MAT1-1
mating type idiomorph and evidence for selfing in the Chinese medicinal fungus *Ophiocordyceps*
sinensis. Fungal Biol. 2013;117(9): 599−610. doi: 10.1016/j.funbio.2013.06.001

[51] Li Y-L. Preliminary study on the ascospores of *Cordyceps sinensis*. Edible Fungi Chin.
2002;21(4): 9−10.

[52] Lin W-C, Wu Y-W, Xie M-C, Zhu J-S. Reduced liver necrosis and fibrosis and improved liver
functions in a CCl4-induced hepatitis model by ReishiMax supplementation. Gastroenterol.
2004;126(4II): A431.

[53] Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular Evolutionary
Genetics Analysis version 6.0. Molecular Biology Evolution 2013;30: 2725−2729. doi:
10.1093/molbev/mst197

[54] Buetow KH, Edmonson M, MacDonald R, Clifford R, Yip P, Kelley J, et al. High-throughput
development and characterization of a genome wide collection of gene-based single nucleotide
polymorphism markers by chip-based matrix-assisted laser desorption/ionization time-of-flight
mass spectrometry. Proc. Natl. Acad. Sci. USA. 2001;98(2): 581–584.
https://doi.org/10.1073/pnas.98.2.581

[55] Stensrud Ø, Schumacher T, Shalchian-Tabrizi K, Svegardenib IB, Kauserud H. Accelerated
nrDNA evolution and profound AT bias in the medicinal fungus *Cordyceps sinensis*. Mycol. Res.
2007;111: 409–415. doi: 10.1016/j.mycres.2007.01.015

[56] Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformat.
2001;17(8): 754−755. doi: 10.1093/bioinformatics/17.8.754

[57] Li X-Z, Li Y-L, Wang Y-N, Zhu J-S. Multiple genomically independent genotypes of
Ophiocordyceps sinensis may not be targets of RIP mutagenesis causing cytosine-to-thymine

transitions. G3 Genes|Genomes|Genetics 2023 (in peer review). BIORXIV Preprint

https://www.biorxiv.org/content/10.1101/2022.12.09.519819v2

[58] Li Y-L, Yao Y-S, Zhang Z-H, Xu H-F, Liu X, Ma S-L, et al. Synergy of fungal complex

es isolated from the intestines of *Hepialus lagii* larvae in increasing infection potency. J. F

ungal Res. 2016;14: 96−112. DOI: 10.13341/j.jfr.2014.1067

[59] Li X-Z, Li Y-L, Zhu J-S. Differential transcription of the mating-type genes during the sexual

reproduction of natural *Cordyceps sinensis*. Chin J Chin Mater Med 2023, 48 (in press). DOI:

10.19540/j.cnki.cjcmm.20230213.102

[60] Debuchy R, Turgeon BG. Mating-Type Structure, Evolution, and Function in Euascomycetes. In

Kües U & Fischer R. (eds.) Growth, Differentiation and Sexuality. The Mycota, vol 1. Springer,

Berlin, Heidelberg. 2006. pp. 293–323. https://doi.org/10.1007/3-540-28135-5_15

[61] Jones SK, Bennett RJ. Fungal mating pheromones: choreographing the dating game. Fungal Genet

Biol. 2011;48(7): 668–676. doi: 10.1016/j.fgb.2011.04.001

[62] Wilson AM, Wilken PM, van der Nest MA, Steenkamp ET, Wingfield MJ, Wingfield BD.

Homothallism: an umbrella term for describing diverse sexual behaviours. IMA Fungus 2015;6(1):

207–214. doi: 10.5598/imafungus.2015.06.01.13

[63] Zhang S, Zhang Y-J, Shrestha B, Xu J-P, Wang C-S, Liu X-Z. *Ophiocordyceps sinensis* and

Cordyceps militaris: research advances, issues and perspectives. Mycosystema. 2013;32: 577−597.

[64] Zhang S, Zhang Y-J. Molecular evolution of three protein-coding genes in the Chinese cat

erpillar fungus *Ophiocordyceps sinensis*. Microbiol. China. 2015;42(8): 1549−1560. DOI: 1

0.13346/j.mycosystema.2013.04.001

[65] Liu Z-Q, Lin S, Baker PJ, Wu L-F, Wang X-R, Wu H, et al. Transcriptome sequencing and analysis

of the entomopathogenic fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*. BMC

Genomics. 2015;16: 106–123. doi: 10.1186/s12864-015-1269-y

[66] Pang F, Wang L-P, Jin Y, Guo L-P, Song L-P, Liu G-M, et al. Transcriptome analysis of

Paecilomyces hepiali at different growth stages and culture additives to reveal putative genes in

cordycepin biosynthesis. Genomics 2018;110: 162–170. doi: 10.1016/j.ygeno.2017.09.008

[67]   Barseghyan GS, Holliday JC, Price TC, Madison LM, Wasser SP. Growth and cultural-

1212   morphological characteristics of vegetative mycelia of medicinal caterpillar fungus

1213   *Ophiocordyceps sinensis* G.H. Sung et al. (Ascomycetes) Isolates from Tibetan Plateau

1214   (P.R.China). Intl. J. Med. Mushrooms. 2011;13(6): 565−581. DOI:

1215   10.1615/intjmedmushr.v13.i6.90

1216