Dissecting infant leukemia developmental origins with a hemogenic gastruloid model

Denise Ragusa[1,2], Chun-Wai Suen[3*], Gabriel Torregrosa Cortés[4*], Liza Dijkhuis[,1,9], Connor Byrne[1],*

Giulia-Andreea Ionescu[1,10], Joana Cerveira[5], Kamil R. Kranc[6], Anna Bigas[7,8], Jordi Garcia-Ojalvo[4],

Alfonso Martinez Arias[4], Cristina Pina[1,2¶]

[1]College of Health, Medicine and Life Sciences, [2]Centre for Genome Engineering and Maintenance, Brunel

University London, UK; [3]Department of Genetics, University of Cambridge, UK; [4]Department of Medicine and

Life Sciences, Universitat Pompeu Fabra, Spain; [5]Department of Pathology, University of Cambridge, UK;

[6]Centre for Haemato-Oncology, Barts Cancer Institute, Queen Mary University of London, UK; [7]Program in

Cancer Research, Institut Hospital del Mar d'Investigacions Mèdiques, CIBERONC; [8]Josep Carreras

Leukemia Research Institute, Barcelona, Spain

[9]Current address: Sanquin Research Laboratory, Department of Hematopoiesis, Amsterdam, The Netherlands

[10]Current address: Centre for Immunology, Blizard Institute, Queen Mary University of London, UK

*These authors contributed equally to this work

¶Lead contact. Correspondence: cristina.pina@brunel.ac.uk. Brunel University London, Kingston Lane,

Uxbridge UB8 3PH, United Kingdom.

ABSTRACT

Current *in vitro* models of developmental blood formation lack spatiotemporal coherence and

weakly replicate the hematopoietic microenvironment. Developmentally-appropriate models

can enhance understanding of infant acute myeloid leukemia (infAML), which putatively

originates *in utero* and has 50% age-unique genetic events, suggesting unique biology. The

commonest genetic abnormality unique to infants involves homeobox gene *MNX1*, whose

leukemogenic mechanisms remain unknown. Recently, 3D self-organising embryonic stem

cell (SC)-based gastruloids have shown promise in recapitulating embryonic events with

time/space precision. Herein, we report a hemogenic gastruloid (haemGx) system that

captures multi-wave blood formation, progenitor specification from hemogenic endothelium

(HE), and approximates generation of hematopoietic SC precursors. Enforced *MNX1*

expression in haemGx promotes HE formation, perturbs endothelial-to-hemogenic transition, and critically achieves transformation, generating myeloid colonies which display *MNX1* AML signatures. By combining functional assays with single-cell transcriptomics, we establish the haemGx as a new model of normal and leukemic embryonic hematopoiesis amenable to mechanistic exploration.

KEYWORDS

Hematopoiesis; Developmental hematopoiesis; Leukemia; Infant Leukemia; Acute Myeloid Leukemia; MNX1; t(7;12); gastruloid; organoid; single-cell RNA-sequencing

INTRODUCTION

The development of organoid systems revolutionized fundamental and translational biology in the last decade (Anonymous, 2018). Organoids are three-dimensional (3D) cultures of primary, or primary-derived cells, which self-organize under defined culture conditions, to recapitulate key structural features of the tissue from which they originate (Clevers, H., 2016; Kim, J. et al., 2020; Lancaster and Knoblich, 2014). In recapitulating structure, organoids recreate elements of their organismal niche, and faithfully establish some of the regulatory physiology that maintains the tissues of origin in their native, *in vivo* environment (Huch et al., 2017). Organoids have been established from a multitude of tissues, primarily neuronal and epithelial (Barker et al., 2010; Gotoh et al., 2014; Greggio et al., 2014; Huch, Dorrell et al., 2013; Huch, Bonfanti et al., 2013; Lancaster et al., 2013; Sato et al., 2009). They are routinely used as 'avatars' of their tissues of origin to test the consequences of genetic perturbation or drug treatment, and can be superior to classical animal pre-clinical models in anticipating therapeutic effect or toxicity (Clevers, Hans C., 2019).

Another category of organoids uses embryonic or pluripotent stem cells to recapitulate developmental processes in a 3D *in vitro* or *ex vivo* space (van den Brink, S C and van Oudenaarden, 2021). Blastoid-type organoids focus on the initial zygotic divisions and can progress through

cleavage-like and blastula-like formation (Rivron et al., 2018; Yu et al., 2021). Gastruloids, on the other hand, display elongation and symmetry breaking to mimic gastrulation and early tissue territorialisation for primordial organogenesis (Arias et al., 2022). Multiple variations of gastruloid protocols have captured mesendodermal specification (Turner et al., 2017; van den Brink, S C et al., 2014), posterior and anterior neurulation (Girgin et al., 2021), somitogenesis (van den Brink, S C et al., 2020), as well as fetal heart organogenesis (Rossi et al., 2021) with varying efficiencies. Tissue-specific topography of molecular programs is broadly maintained (Beccari et al., 2018; Moris, Naomi et al., 2020), making gastruloid models superior to embryoid bodies in capturing embryonic organisation *in vitro*.

Developmental haematopoiesis progresses through 3 waves of cell specification which produce blood cells in reverse hierarchy, with early emergence of unipotent cells and late stem cell specification (Costa et al., 2012; Dzierzak and Bigas, 2018; Lacaud and Kouskoff, 2017; Medvinsky, Alexander et al., 2011). Embryonic red blood cells and macrophages are initially produced in the yolk sac of the mouse embryo at E7.5 and are generated from angioblasts (Lacaud and Kouskoff, 2017). Specification of these early blood cell types does not rely on bilineage or multilineage progenitors, and the red blood cells produced are nucleated and express embryonic globins (Kingsley et al., 2006). Around E8.25, a second, pre-definitive wave of blood production generates erythro-myeloid-megakaryocytic progenitors (EMPs), which eventually differentiate into enucleated erythrocytes, granulocytes, monocytes and megakaryocytes / platelets (McGrath et al., 2015). EMPs, like the subsequent intra-embryonic wave of blood production, are specified from a specialised endothelium – haemogenic endothelium (HE) (Marcelo et al., 2013) – through columnar remodelling and intra-luminal budding of haematopoietic cells, a process known as endothelial-haematopoietic transition (EHT) (Lacaud and Kouskoff, 2017). The third wave of blood production is also HE-based and occurs in the ventral wall of the dorsal aorta and adjacent vessels between E9.5-E11.5. It produces myelo-lymphoid (MLP) and multipotent progenitors (MPP) (Zhu et al., 2020), and a small number of haematopoietic stem cells (HSC) (Medvinsky, A. and Dzierzak, 1996; Medvinsky, A. L. et al., 1993), which expand in the fetal liver and eventually migrate to the bone marrow, where

they mature to sustain blood production postnatally and throughout adult life (Hall et al., 2022). EHT

generates characteristic clusters of cells, which are a hallmark of definitive and intra-embryonic

hematopoiesis (Lacaud and Kouskoff, 2017; Medvinsky et al., 2011).

In vitro specification of hematopoiesis from embryonic and induced pluripotent stem cells under

specified growth factor and stromal co-culture conditions efficiently generate granulocytic-monocytic

and erythroid cells, and to a lesser extent, lymphocytes (Sroczynska et al., 2009). However, they

cannot produce self-renewing, re-populating HSCs in the absence of genetic manipulation, and even

in this case, efficiency and reproducibility are variable (Lis et al., 2017; Sugimura et al., 2017).

In vitro recapitulation of developmental hematopoiesis has the potential to shed light on the biology

of those hematological malignancies that are initiated *in utero* and potentially transform an embryonic

cell stage and/or require the embryonic environment for transformation (Cazzola et al., 2021).

Indeed, by using an iPS-based model, it has been possible to identify the transient embryonic cell of

origin of the most common pediatric malignancy, the pro-B acute lymphoblastic leukemia (ALL)

associated with t(11;21) translocation, which generates the *ETV6-RUNX1* fusion (Boiers et al.,

2018). ALL is the most common hematological malignancy in children and young adults, while Acute

Myeloid Leukemia (AML) dominates in elderly individuals (Britten et al., 2019). However, in the first

year of life, AML is at least as frequent as ALL, and is characterized by a distinct set of chromosomal

abnormalities, 50% of which are exclusive to this age group (Balgobind et al., 2011; Fornerod et al.,

2021). The most common of these is the translocation t(7;12)(q36;p13), a deadly form of AML

molecularly characterized by ectopic activation of the *MNX1* gene at 7q36 (Espersen et al., 2018).

Its overexpression does not result in leukemic transformation of neonatal cord blood cells or adult

mouse bone marrow, but blocks erythroid differentiation and results in cellular senescence

(Ingenhag et al., 2019; Waraky et al., 2022; Wildenhain et al., 2010; Wildenhain et al., 2012).

Engineering of the t(7;12) translocation in human iPS cells captures some of the transcriptional

characteristics of t(7;12) patients and enhances erythroid and myeloid differentiation, but has not

been shown to result in transformation (Nilsson et al., 2022). Interestingly, the translocation

significantly depletes megakaryocytic signatures in iPS cells, suggesting that this model may not

fully capture the spectrum of *MNX1*-rearrangement leukemias, which are mostly undifferentiated or

poorly differentiated AML (FAB M0-M2, 70%), but include 15% of megakaryoblastic leukemia

(Espersen et al., 2018; Taketani et al., 2008).

In this study, we adapt the three-dimensional gastruloid model of mammalian development to

capture multi-wave establishment of blood formation in the embryo. Through cytokine-driven

maturation of self-organising gastruloids over a 216 hour-period, we observe sequential specification

of endothelium, HE, erythro-myeloid progenitor (EMP) and myelo-lymphoid progenitor (MLP)

programs and recapitulate the EHT topography with generation of hematopoietic clusters.

Interestingly, the gastruloid model can be transformed by introduction of *MNX1*, with sustained serial

re-plating of colony-forming cells. By systematically contrasting the gene expression profile of *MNX1*-

overexpressing hemogenic gastruloids and MNX1-rearranged leukemias with single-cell signatures

obtained from hemogenic gastruloid differentiation, we position the cells targeted by *MNX1* at the

HE-to-EMP transition and provide a mechanistic explanation for the strict developmental association

of MNX1 with infant AML.

RESULTS

The hemogenic gastruloid protocol captures cellularity and topography of developmental

blood formation

The original 120h-gastruloid protocol (van den Brink, S C et al., 2014) matches molecular and

organizational aspects of embryo development up to day 8 (E8.0) and captures incipient endothelial

and erythroid-biased transcriptional signatures which are expected to correspond to yolk-sac based

EMP production (Beccari et al., 2018). More recent adaptations of the protocol have used

mechanical agitation, ultra-low adherence plates, and/or matrices to extend the gastruloid life to

168h (Rossi et al., 2021; van den Brink, S C et al., 2020), or E9.5-10.0, and recapitulate later

developmental events such as cardiac specification and somitogenesis, both of which are critically

associated with maturation of the intra-embryonic hematopoietic system. We used ultralow

adherence multi-well plates (see STAR Methods) and sought to promote hemato-endothelial

specification by combining the axes-inducing WNT pulse with the BMP pathway inducer activin (Fig.

1A). The addition of activin was required for activation of the endothelial *Vegfr2/Kdr/Flk-1* locus (Fig.

S1A), a step which we consistently noted as critical for the subsequent detection of hematopoietic

cells. We extended the culture beyond 168h to 216h (Fig. 1A) to attempt to capture HSC-producing

AGM hematopoiesis at E10.5-E11.0. We used Flk1-GFP mouse ES cells (Jakobsson et al., 2010)

to visually monitor endothelial specification (Fig. 1B) and added VEGF and basic FGF (FGF-2) after

the activin pulse to promote development of HE (Sroczynska et al., 2009). Based on paracrine

signals critical to AGM maturation (Souilhol et al., 2016), we tested the time-dependent effect of Shh

in the cultures and observed that its addition in a 24h-pulse at 144h only, increased the fraction of

endothelial VE-Cadherin (CD144)$^+$ and candidate VE-Cadherin$^+$CD45$^+$ pre-HSC (Rybtsov, S. et al.,

2011; Taoudi et al., 2008) at 192h (Fig. S1B-C), suggestive of a recapitulation of the *in vivo* effect

on pre-HSC specification from HE. Addition of SCF, Flt3-ligand and TPO in the last 48h of the

gastruloid culture increased the fraction of hematopoietic cells at end-point, as measured by %CD45$^+$

cells (Fig. S1D). The presence of FGF-2 in the same period did not affect the final %CD45$^+$ cells

(Fig. S1E), and we omitted the cytokine from 168h onwards to simplify the protocol. We typically

cultured *Flk1-GFP* mES cells in serum with leukemia inhibiting factor (LIF) and pre-treated the cells

in '2i' (GSK3b inhibitor – Chiron – and MEK inhibitor – PD) + LIF prior to gastruloid assembly (Fig.

1A). However, the pre-treatment step may be dispensable if the mES cultures have a compact

pluripotent morphology, with minimum cell differentiation.

We used flow cytometry to monitor the timing of emergence of phenotypic endothelium, HE,

hemogenic progenitors, and candidate pre-HSC/HSC on the basis of Flk1-GFP (Fehling et al., 2003),

c-Kit (Marcelo et al., 2013), CD41 (Mikkola et al., 2003) and CD45 (Rybtsov et al., 2011), respectively

(Fig. 1C). The first endothelial cells were detectable at 96h (Fig. 1C), with c-Kit$^+$ HE first apparent at

120h (Fig. 1C). A transient wave of CD41$^+$ cells followed at 144h (Fig. 1C and S1F), which also

included double CD41$^+$CD43$^+$ cells (Fig. S1G-H), compatible with early pre-HSC (Rybtsov, Stanislav

et al., 2014). This was followed by the emergence of CD45$^+$ cells at 192h, which was consolidated

at 216h (Fig. 1C and S1I). We confirmed that sequential emergence of CD41$^+$ and CD45$^+$ cells could

be observed in the more widely used E14TG2a (E14) mES cells (Hooper et al., 1987) (Fig. S2A-C),

with end-point gastruloids initiated with mES cells from different genetic backgrounds generating

similar levels of CD45+ cells (Fig. S2D). From 120h, gastruloids contained cells with hematopoietic

progenitor potential, evidenced in colony-forming cell (CFC) assays (Fig. 1D). Erythro-myeloid

progenitor numbers peaked at 144-168h, with a downward trend at the 2 latest timepoints, which

also showed a bias towards myeloid colonies (Fig. 1D). Notably, CD45+ cells at 216h (Fig. 1E and

S2E) were observed in small clusters budding from a Flk1-GFP / CD31 endothelium reminiscent of

hematopoietic emergence in the dorsal aorta. Together with the ordered emergence of HE, CD41+

progenitors, and candidate CD45+ pre-HSC, the cluster-like arrangement of hematopoietic cells

configures the hemogenic gastruloid as a faithful *in vivo* model of developmental hematopoiesis

amenable to cellular and molecular dissection of blood cell specification, including the role of the

hematopoietic niche.

Hemogenic gastruloid single-cell trajectories reveal pre-definitive and definitive waves of

hematopoiesis and support generation of pre-HSC

In order to characterize the extent and progression of developmental hematopoiesis in the

hemogenic gastruloid model, we performed a single-cell RNA-sequencing (scRNA-seq) time-course

analysis of gastruloid cells in reference to the mouse embryo. As summarized in Fig. 2A, we sorted

cells from 2 independent gastruloid cultures at 120, 144, 168, 192 and 216h and profiled a total of

846 cells using the Smart-Seq2 protocol (Picelli et al., 2014). In line with the flow cytometric

phenotyping, we sorted c-Kit+ cells at 144 and 192h to capture endothelial and HE cells at 2 distinct

points of blood production, as well as 144h-enriched CD41+ cells, and the CD45+ cells emergent at

192 and 216h. We also profiled live single cells obtained at the different timepoints without selection

on hematopoietic markers, with the goal of understanding the microenvironment in which gastruloid

hematopoiesis is specified. Library preparation and sequencing generated an average of 120000

reads/cell, which were mapped to an average of 4000 genes/cell, with almost no cells showing signs

of stress or dying as seen by the mitochondrial DNA fraction (Fig. S3A). Read and gene counts were

Figure 1 – Hemogenic gastruloids produced from mES cells promote hemato-endothelial specification with spatio-temporally accurate ontogeny. (A) Timeline of mES cells assembly and culture into gastruloids over a 216h time period with the addition of appropriate factors for the promotion of hemato-endothelial specification. **(B)** Imaging of hemogenic gastruloids over time from 72h to 216h at 10x magnification, showing the assembly and growth of the 3D structures and the polarization of the Flk-1-GFP marker from 96h; scale bar: 100μm **(C)** Flow cytometry analysis of hemogenic gastruloids for expression of c-Kit, CD41, and CD45 markers from 96h to 216h, assessing the emergence of c-Kit-endothelium, hemogenic progenitors, and pre-HSC/HSC at each time point. **(D)** Colony-forming unit (CFU) assay of disassembled gastruloids assessing the ability to form hematopoietic colonies in multipotential methylcellulose-based medium. GEMM: granulocyte-erythroid-monocyte-megakaryocyte; GM: granulocyte-monocyte; M: monocyte; E: erythroid; BFU-E: burst-forming unit erythroid. CFU frequency of 5 gastruloids, n=3, mean±SD **(E)** Immunostaining of whole individual gastruloids at 216h showing the localized expression of Flk-1 (green), c-Kit (yellow), and CD45 (red) and DAPI (blue) nuclear staining, in a topologically accurate configuration; scale bar: 50μm.

similar between biological replicates, cell types and at different time points, with the sole exception

of the 120h unsorted sample which was sequenced to same depth (similar average read count), but

mapped to twice the number of genes (Fig. S3B), potentially reflecting multi-gene program priming

at the onset of hemogenic specification. We selected highly varying genes before principal

component analysis (PCA) dimensionality reduction and retained the most relevant dimensions. In

the PCA reduced space we constructed a KNN graph and used uniform manifold approximation and

projection (UMAP) to visualize the data on 2-dimensions. We looked for cell communities using

Leiden clustering on the graph (Fig. 2B). We made use of the transcriptional identity of the cells to

describe chronological and biological aspects of multi-step hemogenic specification. We identified

12 cell clusters of which 2 (clusters 0 and 5) almost exactly mapped to c-Kit+ cells, 2 (clusters 1 and

8) contained CD45+ sorted cells, and cluster 4 uniquely captured the CD41+ population of cells

observed at 144h. Although some unsorted cells could be observed overlapping with the hemogenic

cell clusters (Fig. 2B), most cells occupied different transcriptional spaces, in line with the relatively

low frequency of hemogenic cells, particularly at the later time points (Fig. 1C).

To explore the tissue or lineage affiliation of the different clusters, we performed differential gene

expression between each cluster and the remainder cells using Wilcoxon ranking test, and

established classifier gene lists for each cluster (Supplemental File S1). The gene lists were

compared with the PanglaoDB (Franzen et al., 2019) (Fig. 2C) and Descartes (Cao et al., 2020) (Fig.

S3C) repositories of scRNA-seq expression profiles through the EnrichR gene set enrichment

analysis tool suite (https://maayanlab.cloud/Enrichr/) (Chen, E. Y. et al., 2013; Kuleshov et al., 2016)

to identify enriched cell type representation within the clusters (Supplemental File S2). Clusters 5

and 0, which were mostly populated by c-Kit+ cells at 144 and 192h, respectively (Fig. 2B and S3D)

had widespread expression of *Cdh5* (VE-cadherin), with some cells also expressing *Gata2* or *Runx1*,

compatible with hematopoietic specification from HE (Fig. 2D) (Chen, M. J. et al., 2009; de Pater et

al., 2013). Accordingly, their gene expression signatures captured the program of endothelial cells,

including those of the aorta, with some evidence of HSC-enriched genes (Fig. 2C). The cells

configure distinct clusters at the individual time points (Fig. S3D: 144h cluster 1; 192h cluster 5) (also

Figure 2 – Time-resolved analysis of scRNA-seq of hemogenic gastruloids captures successive waves of hematopoietic specification. (A) Summary of plating strategy for scRNA-seq analysis of gastruloids at 120h, 144h, 168h, 192h, and 216h without selection of surface markers ('all') or on the basis of expression of c-Kit/ScaI (green shading), CD41 (blue), or CD45 (orange); P = plate. **(B)** UMAP projection of all sequenced cells colored by annotated clusters (left) and by positivity to sorting markers c-Kit/ScaI, CD41, and CD45, or mapping to the unselected sort ('whole' corresponding to 'all' in panel A). **(C)** Cell type enrichment analysis of cluster classifier genes, extracted by differential expression in comparison to all other clusters, using the PanglaoDB and Descartes repositories. The statistical power of representation of individual cell types is expressed in EnrichR combined score with a p-value threshold of < 0.05. **(D)** Expression of key lineage markers *Cdh5* (VE-Cadherin), *Gata2*, *Runx1*, and *Pdgfra* in individual cells projected onto UMAP plots of global analysis shown in **A**. Colour scale represents Z-score gene expression from 0 (blue, bottom) to maximum (orange, top).

Supplemental File S1), and occupy minimally overlapping transcriptional spaces, suggesting

different waves of HE specification or maturation. Compatible with wave-like specification of

hematopoietic cell types from HE, 144h-enriched HE cluster 5 was immediately adjacent to CD41$^+$-

enriched (Fig. 2B) *Runx1$^+$Cdh5$^-$* (Fig. 2D) cluster 4, which exhibits an erythroid/megakaryocytic cell

signature suggestive of pre-definitive EMPs (Supplemental File S1) (Fig. 2C and S3D). The EMP-

like space was devoid of cells, including hematopoietic CD45$^+$ cells (Fig. 2B), at the later 192 and

216h-time points (Fig. S3D), which instead occupied a second discrete *Runx1$^+$Cdh5$^-$* transcriptional

space (Fig. 2D) of myeloid-lymphoid affiliation (Fig. 2C and S3D) (Supplemental File S1), compatible

with MLP-like cells. We sought to align gastruloid-derived hemogenic cells with developmental

hematopoiesis in the embryo by projecting our data onto recent scRNA-seq studies which dissected

endothelial maturation and HE-to-hematopoietic transitions (Fadlullah et al., 2022), and pre-HSC

and HSC emergence (Vink et al., 2020) in the AGM. We reanalyzed the scRNA-seq datasets using

our own pipeline. UMAP-based clustering and dimensionality reduction representations broadly

recapitulated the clustering and representations of the original publications (Fig. 3A-C, and S4A-B),

with the exception of minor clusters (clusters 6, 9, and 10, Fig. 3A; 4, 5, 7, and 9, Fig. 3B), which are

not represented or are captured distinctly in the original probably due to filtering criteria, but which

configure distinct hematopoietic lineage affiliations (Supplemental File S3) (Fig. S4A-B). Compatible

with our cell type classification, gastruloid endothelial clusters 0 and 5 projected onto embryonic

clusters of endothelial, aortic endothelial, and HE affiliation (Fig. 3D-E), with rare cells exhibiting pre-

HSC characteristics (Fig. 3E). Candidate EMP-like (gastruloid cluster 4, Fig. 2B) and MLP-like

(gastruloid cluster 8, Fig. 2B) clusters, on the other hand, projected onto clusters of similar affiliation

in the embryo datasets (Fig. 3A, D, cluster 6; Fig. 3C, E, clusters 4 and 5). Interestingly, some of the

cells occupied the transcriptional space of HE-to-hematopoietic (gastruloid cluster 4, Fig. 3A, D

cluster 7), and pre-HSC-to-HSC transition (gastruloid cluster 8, Fig. 3C, E clusters 0 and 6), in

support of incipient definitive hematopoiesis representation in the gastruloid model. Importantly, the

gastruloid model also recapitulates differentiation of stromal / niche elements critical for

hematopoiesis specification in the embryo. These include: *Pdgfra$^+$* stromal cells (gastruloid cluster

1, Fig. 2B and Fig. 2D), which project onto the PDGFRa⁺ mesenchyme in Fadlullah et al. (2022)

(Fadlullah et al., 2022) (Fig. 3A,D, embryo cluster 5); and cells with candidate autonomic nervous

system identity (gastruloid clusters 3 and 9, Fig. 2B) (Fig. 2C and S3C) which are not captured in

the Fadlullah et al. (2022) and the Vink et al. (2020) studies, but have been reported to sustain HSC

specification (Fitch et al., 2012; Kapeni et al., 2022). Both these niche cell types are specifically

present at 192h (Fig. S3D, 192h clusters 1-3) and probably emerge 1 day earlier (Fig. S3D, 168h),

thus configuring a timeframe compatible with their reported support of pre-HSC. The remaining

Figure 3 – Single cell profiles of hemogenic gastruloids project onto hemato-endothelial differentiation programs
in the mouse AGM. (A-B) Re-analyzed UMAP projection of scRNA-seq dataset from Fadlullah et al. (2020) showing
annotated clusters **(A)** and sequenced conditions **(B)** through endothelial to hematopoietic maturation. **(C)** Re-analyzed
UMAP projection and cluster annotation of scRNA-seq dataset from Vink et al. (2020) capturing pre-HSC and HSC
emergence (clusters 0 and 1), endothelial cells (clusters 2 and 3), and other hematopoietic-affiliated populations (clusters
4-9). **(D-E)** Projection of hemogenic gastruloids single-cell profiles from this study (colored according to their annotated
clusters in Fig. 2B) over the UMAP from Fadlullah et al. (2020) **(D)** and Vink et al. (2020) **(E)**.

gastruloid clusters have unclear lineage affiliations (Supplemental File S2). They do not project onto AGM-derived cells (Fig. 3D-E), probably reflecting their early gastruloid time signatures (Fig. S3D, 120-144h) or a distinct topography which is not strictly hemogenic.

Altogether, the data support the notion that the gastruloid model captures developmental blood formation with cellular and temporal precision, and faithfully recapitulates the multi-wave nature of pre-definitive and definitive hematopoiesis. Projection onto the embryo places the 216h-end of the gastruloid protocol at the pre-HSC transition, suggesting that it constitutes a good model to understand intrinsic and extrinsic regulation of definitive hematopoiesis.

Infant leukemia gene *MNX1* favours early steps of hemogenic specification

We explored the utility of the hemogenic gastruloid model in understanding the cellular consequences of oncogenic events putatively initiated during development. We elected to investigate *MNX1*-driven AML, the most common of infant-unique forms of AML, which we and others (Ingenhag et al., 2019; Ragusa et al., 2022; Waraky et al., 2022) have shown to be incapable of transforming adult cells.

We overexpressed *MNX1* (MNX1-OE) in *FLK1-GFP* mouse ES cells by lentiviral transduction (Fig. S5A) and compared MNX1-OE and control (empty vector, EV) cell performance in the hemogenic gastruloid model (Fig. 4A). We used a human *MNX1* cDNA (Fig. 4B) to distinguish from the endogenous gene, but the degree of homology is nevertheless high (84%), supporting functional equivalence. MNX1-OE gastruloids activated polarised *FLK1-GFP* expression and elongated with similar kinetics to EV (Fig. 4A), but consistently produced larger gastruloids (Fig. 4C) denoting increased cellularity (Fig. S5B). From 192h onwards, MNX1-OE gastruloids had a higher frequency of spontaneously contractile structures (Fig. 4D and Supplemental Movie), compatible with mesodermal cardiac specification. We interrogated the cellularity of gastruloids at the critical hemogenic timepoints of 144h and 192h using markers c-Kit, CD41 and CD45 in flow cytometry. We observed a significant expansion of the c-Kit+ compartment specifically at 144h (Fig. 4E and S5C), with relative reduction of CD41+ cells (Fig. 4F and Fig. S5D) at the same time point. No changes in

proportion of any of the cell types were observed at 192h (Fig. 4E-G and S5C-D), although absolute

cell numbers may be higher in MNX1-OE gastruloids (Fig. S5B).

In order to understand the molecular programs downstream of *MNX1* which promote cardiogenic,

endothelial and potentially hematogenic fates, we initially performed bulk RNA-sequencing of end-

point gastruloids and their starting mouse ES cells with and without *MNX1* overexpression (Fig. S6A-

B). We verified expression of the human *MNX1* transgene in the sequenced reads (Fig. S6C).

Interestingly, we could not detect endogenous *Mnx1* in any of the samples (Fig. S6C), suggesting

that it does not normally play a role at the hematogenic stage of embryonic development, or indeed

in pluripotent cells. We detected 464 differentially overexpressed genes in MNX1-OE gastruloids (p-

value<0.05, FDR adjusted p-value<0.10; Supplemental File S4), and used cell type enrichment

(PanglaoDB, Fig. 5A) and GO Biological Process (Fig. 5B-C and S6D) to identify enriched functional

categories. MNX1-OE hemogenic gastruloids had enrichment in cardiogenic programs (Fig. 5A-C;

5A – cardiomyocytes; 5B-C – muscle contraction, heart development, muscle organ development),

in agreement with our observations of autonomous contractile activity (Fig. 4D). There was also

enrichment of programs associated with endodermal derivatives, including the gut (Fig. 5A –

enterocytes, which may further enhance heart development (Silva et al., 2021). Other enriched

functional categories were predominantly metabolic, including lipid metabolism (Fig. 5B, D; S6D-E),

reminiscent of some of the transcriptional consequences of *MNX1* in the context of leukemia

(Ragusa et al., 2022). Compatible with the absence of relative enrichment of hemogenic cells in late

gastruloids, we did not find an enrichment in endothelial or hematopoietic-affiliated programs in

MNX1-OE gastruloids. In contrast, *MNX1*-expressing mouse ES cells had clear evidence of

activation of blood (Fig. 5D – platelets) and angiogenic signatures (Fig. 5E-F – angiogenesis),

including key markers and regulators such as *c-Kit*, *Vegfa*, *Sox17* (Clarke et al., 2013; Nobuhisa et

al., 2014), and *Runx1* (Fig. 5F) (Supplemental File S4), which are required for the establishment of

definitive hematopoiesis. Likewise, incipient cardiogenic programs are also enriched in MNX1-OE

Figure 4 – Hemogenic gastruloids with *MNX1* overexpression have increased cellularity and enhanced cardiogenic and hemogenic endothelial potential. (A) Imaging of hemogenic gastruloids with *MNX1* overexpression and EV controls at 10x magnification, showing appropriate assembly and polarization of the Flk1-GFP marker. scale bar: 300μm. **(B)** Quantitative (q)RT-PCR confirming *MNX1* overexpression in hemogenic gastruloids at endpoint (216h). Fold change in expression was calculated by normalization to *HPRT1*. Statistical difference was calculated by Student's t-test, p < 0.001 (**). Error bars indicate ± SD of 3 replicates. **(C)** Size of gastruloids at each timepoint, determined by the distance of the longest extremes in μm. Statistical difference was calculated by Student's t-test, p value < 0.05 (*), 0.001 (**), 0.0001 (***), and 0.00001 (****). Error bars represent standard deviation (SD) of n=3. **(D)** Proportion of gastruloids exhibiting spontaneous contraction at 192h; error bars show SD of n=3. **(E-G)** Flow cytometry quantification of positive MNX1 and EV gastruloids cells for c-Kit **(E)**, CD41 **(F)**, and CD45 **(G)** at 144h and 192h. Statistical difference was calculated by Student's t-test, p < 0.05 (*).

mouse ES cells (Fig. 5D – cardiac precursor cells) prior to cardiac cell specification in gastruloids. It is important to note that MNX1-OE mouse ES cells can be maintained as pluripotent cells in the presence of LIF and do not exhibit morphological signs of spontaneous differentiation, suggesting that the enhancement of hemogenic and cardiogenic signatures denotes priming rather than full-scale activation of lateral mesoderm lineage programs.

Taken together, functional and transcriptional analysis of MNX1-OE gastruloids suggests that *MNX1* ectopically promotes the specification of lateral mesodermal lineages, including activation of hemato-endothelial programs and early expansion of hemogenic endothelium, compatible with the hypothesis of an early developmental origin of *MNX1* infant leukemia.

***MNX1* targets the HE-to-EMP transition to initiate leukemic transformation**

In an attempt to deconvolute the cellular heterogeneity underlying bulk RNA-seq differential gene expression of MNX1-OE hemogenic gastruloids, we interrogated the cluster-specific signatures inferred from our single-cell time-course analysis of gastruloid differentiation (Fig. 6A; refer to Fig. 2B), including the clustering structure at individual timepoints (Fig. 6B; refer to Fig. S3D), to infer differential cell type representation. In support of enhanced hemato-endothelial specification driven by *MNX1*, we observed enrichment of HE cluster signatures, comprising both clusters 5 and 0 (144 and 192h, respectively), as well as of EMP-like cluster 4 (144h) and MLP/pre-HSC-like cluster 8 (192/216h) signatures (Fig. 6A), altogether compatible with increased representation of HE and hematopoietic cells in MNX1-OE gastruloids. The signature for clusters 10 (120h), which has less clear lineage-affiliations and include pluripotency genes such as *Zfp42*, *Nanog*, and *Sall4* (Papatsenko et al., 2015), was also enriched, which could suggest persistence of primed ES cells. Significantly, clusters 5 and 0 HE signatures were also enriched in RNA-seq data of infAML patients (Balgobind et al., 2011) (TARGET, https://ocg.cancer.gov/programs/target) carrying an *MNX1*-overexpressing chromosomal rearrangement (MNX1-r) [t(7;12) translocation] (Fig. 6A-B), thus implicating a characteristic hemogenic developmental stage in infAML biology, and validating the mechanistic role of *MNX1* in this form of leukemia. It is noteworthy that, unlike HE signatures, EMP and MLP/pre-HSC signatures were not enriched in MNX1-r/t(7;12) infAML (Fig. 6A-B).

The t(7;12) RNA-seq signature was calculated in comparison with other forms of pediatric leukemia (Supplemental File S5) (see STAR Methods for detail), which could dilute hematopoietic progenitor signatures. However, RNA-seq data for the most common form of infAML, driven by the t(9;11) *KMT2A-MLLT3* fusion (https://ocg.cancer.gov/programs/target), shows a significant enrichment of

Figure 5 – *MNX1* **overexpression promotes lateral mesodermal expansion in hemogenic gastruloids and primes angio- and hemogenic programs in mES. (A)** Enrichment of Gene Ontology (GO) terms in differentially upregulated genes in GX-MNX1 compared to GX-EV (filtered by FDR ≤ 0.1) relating to biological processed (PANTHER database). The grey intercept shows p = 0.05 threshold in −log10. **(B)** Cell type analysis of differentially upregulated genes in GX-MNX1 using the Panglao DB 2021 database. **(C)** Bipartite network plot of upregulated genes in GX-MNX1 mapping to PANTHER biological processes filtered, constructed on ExpressAnalyst (www.expressanalyst.ca). Full network shown in Supplementary Figure S6D. **(D)** Biological processes (PANTHER) enriched in differentially upregulated genes in overexpressing mES compared to EV control. **(E)** Cell type analysis corresponding to upregulated genes in MNX1 overexpressing mES. **(F)** Bipartite network of upregulated genes in MNX1-mES mapping to biological processes from PANTHER. Full network in Supplemental Figure S6E.

the MLP/pre-HSC-like cluster 8 (Fig. S7A-B), compatible with the myeloid progenitor affiliation of

KMT2A-rearranged AML. This re-enforces the notion that the HE stage may indeed be critical in

t(7;12) *MNX1* leukemia biology. To verify the significance of the individual genes enriched in each of

the signatures, we considered the leading-edge genes (LEGs) in MNX1-OE and t(7;12) enrichments

(Fig. 6C and Supplemental File S5). Over 40% of t(7;12) LEGs overlapped with MNX1-OE gastruloid

LEGs, with less reverse overlap as expected, given the heterogenous cellular composition of

gastruloids (Fig. 6C). Again, common LEGs, as well as a subset t(7;12)-unique genes aligned with

endothelial cells (Fig. 6D and S7C), firming the link with HE. Similar to cluster enrichments (Fig. 6A-

B), association with erythroid cell types was exclusive to MNX1-OE (Fig. 6D), putatively separating leukemia and non-leukemia-specific roles of MNX1. Conversely, a t(7;12)-unique association with mesodermal / stromal cell types (Fig. 6D and Fig. S7C), encapsulated by the *Pdgfra*⁺ cluster 1 (Fig. 6A and 2D), is not captured in MNX1-OE and EV gastruloids (Fig. 6B).

Although heterogeneous in terms of hematopoietic-lineage affiliation, *MNX1* t(7;12) leukemias are most often of moderately-differentiated myeloid morphology (Espersen et al., 2018; Wildenhain et al., 2010), suggesting progression beyond the HE stage. To more closely investigate later hematopoietic stages, we performed scRNA-seq on MNX1-OE and EV cells from 216h gastruloids broadly sorted on CD41⁺ and/or CD45⁺ gates to encompass all hematopoietic cells. The sequencing quality control confirmed robust detection of an average of 120000 reads/cell, corresponding to 4000 aligned genes/cell (EV and MNX1-OE, respectively), with no stress signatures. Clustering analysis (Fig. S7D and Supplemental File S6) distinguished 6 clusters with representation of both EV and MNX1-OE cells, albeit with enrichment of MNX1-OE cells in cluster 0 and of EV cells in cluster 1 (Fig. S7D-E). Separate clustering and projection of each condition identified 3 clusters each in EV and MNX1-OE cells, which were more clearly discrete in EV compared to MNX1-OE gastruloid cells (Fig. 6E), putatively suggesting distinct degrees of cell specification and/or lineage resolution. Accordingly, cluster-defining molecular signatures were more statistically robust in EV cells (Supplemental File S6). More than half of EV cells were found in cluster 1, which has a gene expression signature enriched for targets of key hematopoietic regulators PU.1 (SPI1), RUNX1 and GATA2 (Fig. 6F), with evidence of myeloid, lymphoid and erythroid lineage programs (Fig. 6G) (Supplemental File 6). Hematopoietic affiliations are more evenly spread across MNX1-OE clusters, particularly 0 and 2, which capture enrichments for GATA2 and PU.1 targets, respectively (Fig. 6F). Accordingly, lineage-affiliated programs were present in the 3 clusters (Fig. 6G), with a bias towards endothelial or hemato-endothelial signatures enriched in cluster 0, and lymphoid and myelo-lymphoid signatures in cluster 2, consistent with the enriched transcriptional regulators (Fig. 6H-I). Relative to EV, MNX1-OE has more extensive erythroid priming, particularly of hemoglobin chains (Fig. 6G), as well as of endothelial signatures, including co-expression with erythro-myeloid

Figure 6 – Hemogenic gastruloids overexpressing MNX1 capture transcriptional features of MNX1-r AML. (A) UMAP of time-resolved global clustering of scRNA-seq of hemogenic gastruloids showing the enriched clusters in GX-MNX1 (blue boxes) and MNX1-r AML (red boxes) determined by GSEA. Cluster numbers as per Fig. 2B. **(B)** Bubble plot of GSEA NES values and statistical significance by –log10(FDR) of enrichments in specific time clusters and corresponding global clusters in GX-MNX1 compared to GX-EV, and MNX1-r AML samples compared to other pediatric AML. **(C)** Intersect between GSEA leading edge genes (LEGs) enriched in MNX1-r AML and GX-MNX1 against time-resolved scRNA-seq gastruloids clusters. **(D)** Radar chart mapping enriched cell types (analyzed in EnrichR using the Panglao DB database) using LEGs shown in **C. (E)** UMAP of scRNA-seq of sorted CD45+ and/or CD41+ MNX1 and EV gastruloids cells at 216h. **(F)** Transcription factor binding site enrichment on top 100 scRNA-seq cluster identifier genes (filtered by p value ≤ 0.05) using the ENCODE and ChEA Consensus TFs from ChIP database on EnrichR. **(G)** Heatmap showing the expression of hematopoietic lineage markers (listed in Methods) for each cell in CD45+ and/or CD41+ sorted GX-EV and GX-MNX1 at 216h. **(H)** Percentage of cells for each lineage identity (determined by co-expression of at least 2 lineage genes) by cluster in CD45+/CD41+ EV (left) and MNX1 (right) gastruloids at 216h. **(I)** Percentage of multi-lineage priming (determined by co-existence of lineage signatures) by cluster in CD45+/CD41+ EV (left) and MNX1 (right) gastruloids at 216h.

signatures (Fig. 6I). In contrast, EV cells had a higher frequency of expression of recombinant Ig

chains, as well as of hematopoietic progenitor regulatory signatures (Fig. 6G). Altogether, the data

are suggestive of perturbed hematopoietic progression upon MNX1-OE, with persistence or

expansion of HE cells at the HE-to-EMP transition, and relative depletion of more progressed myelo-

lymphoid progenitors and pre-HSC. Crucially, MNX1-OE cells in cluster 2 were singularly enriched

in t(7;12) AML patient-signature genes (Fig. 7A), suggesting that this cluster may capture myeloid

leukemia transformation. To test transformation potential of MNX1-OE cells, we dissociated EV and

Figure 7 – MNX1 overexpression transforms hemogenic gastruloids to gain re-plating capacity of myeloid
colonies. (A) Enrichment of MNX1-r AML patient signatures from Ragusa et al. (2022) in clusters of CD45+/CD41+ MNX1
and EV gastruloids, determined by GSEA by comparison of each cluster against the others ('rest'). Statistical significance
is by FDR; *p<0.05. **(B)** Serial colony-forming assay (CFC) of disassembled unsorted MNX1 and EV gastruloids at 216h
on methylcellulose-based medium. 'P' indicates the order of plating. Mean±SD of n=2-4 replicate experiments; *p<0.05,
unpaired t-test with Welch correction. Underneath, pie charts indicate the proportion of colony types at each replating. **(C)**
Representative images of colony types identified as 'compact' (top) and 'diffuse' (bottom) at 20x magnification. Scale bar
= 1000 μm. **(D)** Representative images of methylcellulose plates at early (P2-P3) and late platings (P7). **(E)** Flow cytometry
plot of late plating colonies at P8 for the myeloid marker CD11b; data are overlayed with the respective unstained control
sample, with normalized counts.

MNX1-OE gastruloids, placed them in CFC progenitor assays containing multi-lineage cytokines,

and assessed colony frequency and serial re-plating capacity as an *in vitro* measure of

transformation. From the initial plating, MNX1-OE cells generated more colonies (Fig. 7B), with

progressive selection of a characteristic compact colony morphology depicted in Fig. 7C. EV-derived

colonies were extinguished after 3 platings, while MNX1-OE colony-forming cells persisted at even

frequencies for at least 5 additional platings (Fig. 7D), and generated cells with a myeloid surface

phenotype, as per flow cytometry detection of early myelo-monocytic marker CD11b (Fig. 7E).

Altogether, the data indicate that overexpression of *MNX1* during developmental hematopoietic

specification, expands HE and favours a distinct and potentially heterogenous pathway of myeloid

differentiation with leukemic self-renewal potential.

DISCUSSION

In this study, we have developed and explored a 3-dimensional gastruloid model of hemogenic

mouse development. Through sequential utilisation of extrinsic cues for symmetry breaking,

mesodermal induction, vascular development, and hematopoietic maturation, we extended the self-

organising properties of the gastruloid system to recapitulate key aspects of blood formation with

spatiotemporal accuracy. Namely, we observed the successive specification of hemogenic

endothelium (HE), and of pre-definitive erythro-myeloid progenitors (EMP), definitive myelo-

lymphoid progenitors (MLP) and pre-HSC in coherent temporal progression, and we captured the

formation of discrete micro-aggregates of hematopoietic cells in intimate association with

endothelial-like lumina, reminiscent of mid-gestation HSC-generating aortic clusters. Hemato-

endothelial development was accompanied by generation of stromal components critical for HSC

emergence, including PDGFRA+ mesenchyme (Chandrakanthan et al., 2022) and sympathetic

neurons (Fitch et al., 2012; Kapeni et al., 2022), suggesting coordinated organisation of elements

of a hemogenic niche. Importantly, the *in vitro* gastruloid system responded to a relevant oncogenic

event unique to infant AML (infAML) and putatively characteristic of the developmental period, i.e.

MNX1 overexpression (MNX1-OE). MNX1-OE gastruloids displayed perturbation of hemogenic trajectories and selection of a transformed myeloid progenitor, which not only recapitulated the molecular signatures of human patients, but critically shed light on previously uncharacterised mechanisms of transformation associated with *MNX1*. Thus, recapitulation of physiological stages of blood development and clinical-relevant responsiveness to a pathological leukemia-initiating event uniquely associated with embryonic development, together configure the hemogenic gastruloid as a faithful, robust, and accessible *in vitro* model to dissect blood formation.

Our hemogenic gastruloid system extends recent findings by (Rossi et al., 2022), who observed early erythro-myeloid specification in a gastruloid protocol previously developed for modelling of primordial heart development (Rossi et al., 2021). Together, our data highlight the power of gastruloid models in faithfully capturing mid-development specification of lateral mesoderm-derived programs. They also suggest that modelling of post-gastrulation mesodermal lineage development progresses modularly, with protocol optimization preferentially supporting individual cell fates. Our protocol is not without its limitations, namely (1) it does not organise a networked vascular system, and (2) there is no production of functional hematopoietic stem cells (HSC). In line with the notion of modularity of lineage development, our protocol does not consistently produce spontaneously contractile structures, i.e. a heart primordium, a process intimately associated with HSC formation. Despite the ability of gastruloids to execute transcriptional programs and configure temporally-accurate regulatory networks that sustain specification of cell identity and enable the correct assembly of network-directed micro-topographies, they are unable to fully capture the mechanical and chemical cues that direct and remodel macro-topographies and ultimately specify organismal physiology. Likely, their optimal cell number at initiation gives gastruloids an advantage relative to, for example, much larger embryoid bodies in which micro-topographies may quickly become redundant and disorganised, and, in the case of blood formation, require cell dissociation and additional stages of cell culture to reach similar stages of hematopoietic specification, with loss of the temporal resolution (Garcia-Alegria et al., 2018; Pearson et al., 2015; Sugimura et al., 2017). At a fundamental level, gastruloid models can be seen as capturing the interface between intrinsic and

extrinsic regulation, and become limiting at the critical transition where intrinsic cell identity is superseded by extrinsic tissue or organ coordination enabled by long-distance chemical and physical signalling. Other single-tissue organoid models employ matrices that provide the extracellular support required to organise a degree of macro-topography. Gastruloids themselves, elegantly organise somitogenesis (van den Brink, S C et al., 2020), or approximate a central nervous system (Girgin et al., 2021), in the presence of matrices, but precise combinations of chemical and mechanical cues are tissue-specific and successful 3-dimensional *in vitro* organisation are necessarily modular. In the case of blood, different cell identities are specified independently, at different times and in different places, during embryonic development (Costa et al., 2012; Dzierzak and Bigas, 2018; Lacaud and Kouskoff, 2017; Medvinsky et al., 2011), and the micro-topographical accuracy of the gastruloid model assists in faithfully capturing the successive hematopoietic waves in intimate association with hemogenic endothelium and with contemporaneous specification of at least some relevant stromal cells. Absence of *bona fide* HSC, i.e. transplantable cells capable of long-term reconstitution of the hematopoietic system may be due to insufficient developmental time at the end of the protocol – 216h may be short of HSC emergence – or insufficient extrinsic macro-topographical regulation – with the disorganised vascular network and the absence of coordinated contractile activity key omissions. The protocols that have achieved production of *bona fide* HSC from pluripotent sources (Sugimura et al., 2017) required ectopic expression of multiple transgenes to enforce key transcriptional networks and signalling pathways, and have been difficult to reproduce in other laboratories, suggesting a requirement of yet undefined factors. It has also been proposed that circulating blood flow contributes to HSC specification and shedding from the intra-aortic clusters (North et al., 2009), compatible with the sequential nature of initiation of heart contractility and blood flow, and intra-embryonic re-localization of blood formation and HSC production. Protocols with reported success at HSC formation (Lis et al., 2017; Sugimura et al., 2017) generate HSC at the stromal co-culture step which may expose the culture surface to a certain level of haphazard flow stress. In a proportion of gastruloids, we have observed the generation of cluster-like structures at the surface, rather than internally, re-enforcing the notion of a contribution from mechanical cues.

Several recent reports have demonstrated the utility of organ-on-a-chip (Ooc) solutions for the recreation of adult bone marrow environments in physiological (Sugimura et al., 2020; Torisawa et al., 2014) and pathological (Abdullah Obaid Khan et al., 2022; Chou et al., 2020), i.e. leukemia settings. In the future, it is worth considering adapting Ooc platforms to support gastruloid growth, providing exposure to bespoke matrices and levels of flow at specific stages, to promote HSC development.

The hemogenic gastruloid protocol described in this study does nevertheless faithfully capture pre-definitive EMP-like cells and definitive MLP-like cells and putative pre-HSC which are transcriptionally aligned with E8.5-E10.5 stages of mouse development. Interestingly, the gastruloid hemogenic endothelial cells that temporally associate with EMP-like and definitive hematopoietic progenitors occupy adjacent but largely non-overlapping clusters, in agreement with independent waves of blood formation. The physiological alignment of gastruloid HE is highlighted by the successful modelling of *MNX1*-rearranged (*MNX1*-r) AML, which had so far remained elusive. Several reports by us and others have used ectopic *MNX1-ETV6* cDNA (Wildenhain et al., 2010), *MNX1* overexpression (Ingenhag et al., 2019), or genomic engineering of the t(7;12) rearrangement (Nilsson et al., 2022; Ragusa et al., 2022) in adult mouse or human HSC and progenitors in an attempt to model *MNX1*-r leukemia, but consistently failed to achieve transformation *in vitro* or *in vivo*. Similar to this study, (Ingenhag et al., 2019) observed a relative expansion of erythroblasts upon *MNX1* overexpression, consistently supporting a unique effect in the erythroid lineage, but one which is independent of the potential for transformation. While these observations could bring into question the value of *MNX1* overexpression as a surrogate of *MNX1*-r, the recent engineering of the t(7;12) rearrangement in human iPS cells (Nilsson et al., 2022) also captured an expansion of erythroid progenitors, re-aligning the functional consequences of *MNX1* and its targeting translocation. Interestingly, t(7;12) engineering in iPS cells also promoted differentiation of myeloid progenitors, providing a more accurate alignment with the lineage phenotype of the leukemia, putatively favoured by the expression of embryonic transcriptional programs, although with no evidence of transformation (Nilsson et al., 2022). Systematic comparison of *MNX1*-r / t(7;12) with

other pediatric AML (Balgobind et al., 2011; Fornerod et al., 2021; Ragusa et al., 2022) identified a

unique transcriptional program enriched for lipid metabolism and cell adhesion signatures, but with

limited affiliation to hematopoietic lineages. Instead, t(7;12) signatures include endothelial genes

such as *KDR*, *VWF* or *EDIL3*, as well as *LIN28B*, a negative regulator of fetal-to-adult HSC

maturation via repression of *let-7* (Wang et al., 2022). Detailed mapping of gastruloid hematopoietic

development against *MNX1*-r / t(7;12) leukemia signatures unveils their affiliation with HE, which we

found to be expanded by MNX1-OE at the EMP-like stage suggestive of an early, putatively pre-

definitive, target cell of *MNX1* ectopic expression. The later HE signature at 192h was also enriched

in both end-point MNX1-OE gastruloids and t(7;12) AML samples, although no relative expansion of

Kit+ cells was observed at that time point of gastruloid differentiation, suggesting that the 192h

enrichment may convey the HE lineage affiliation rather than precisely reflect the developmental

time. Indeed, the dispersed profiles of MNX1-OE hemogenic cells by scRNA-seq are compatible with

differentiation heterogeneity, asynchrony and/or delayed resolution of developmental programs, also

apparent through detection of myeloid differentiation signatures in 2 neighbouring clusters, 0 and 2,

which reflect distinct weak GATA2-centric and PU.1 / SPI1-centric regulatory networks. The latter

likely corresponds to the early-transformed cell, which we were able to select through serial replating

and is phenotypically aligned with the myeloid classification of *MNX1*-r / t(7;12) AML. Our study also

detects the erythroid affiliation bias previously described in *MNX1* overexpressing BM cells, which

diverges from t(7;12) AML-associated signatures. Significantly, this erythroid bias does not configure

a distinct MNX1-OE gastruloid cluster upon single-cell analysis, but instead corresponds to pervasive

priming of erythroid signatures, which co-exist with or are superimposed on, endothelial, myeloid,

and lymphoid signatures. Notably, the endothelial priming is more extensive, but the lymphoid

programs are less progressed than in control cells, and there is lower frequency of expression of

pre-HSC associated transcription factors. This suggests expansion and/or selection of a cell at the

HE-to-hematopoietic transition, which is developmentally closer to the pre-definitive EMP stage

and/or uniquely tolerant of ectopic erythroid expression, a property not shared by adult BM cells,

which couple MNX1-expanded erythroid expression with senescence. In contrast, MNX1-OE

gastruloid cells persist and expand, and can transform into a leukemia which is myeloid in nature,

and carries unique residual endothelial, but not erythroid, programs, effectively dissociating

transformation from erythroid specification or bias. It is possible that the MNX1-driven oncogenic

program requires a cell that is tolerant of erythroid gene priming, but does not progress through

hematopoietic lineage differentiation, and consequently dispenses full activation, or full repression

of the erythroid signature. Such tolerance may be explained by chromatin plasticity and/or an

adaptive checkpoint response, either as a consequence of MNX1 activity, or as a result of additional

genetic events. In t(7;12) AML patients, additional copies of chromosomes 19 or 8 are frequently

encountered (Espersen et al., 2018), whose origin and significance are currently unexplored. Further

exploration of the gastruloid model and dissection of the corresponding stages in the embryo will

refine these findings, and pinpoint the cell-of-origin and precise developmental window in which

MNX1 is active, as well as suggest specific vulnerabilities that can be targeted therapeutically.

In conclusion, the hemogenic gastruloid model constitutes a faithful *in vitro* model of embryonic and

fetal blood specification and development. Its cellular and temporal resolution has the potential to

add to the mechanistic understanding of developmental blood disorders, offering a medium-

throughput platform for therapeutic compound testing. Refinement of the mouse gastruloid model

and extension to the human will further strengthen the translatability of its findings.

ACKNOWLEDGEMENTS

This project was funded by a start-up grant and a BRIEF award from Brunel University London to

CP, and by ERC Advanced Grant (MiniEmbryoBlueprint 834580) to AMA. GTC was funded by grant

FPU18/05091 from the Spanish Ministry of Universities. CP was also funded by a KKLF Intermediate

Fellowship (KL888), a Leuka John Goldman Fellowship for Future Science (2017-2019), and a

Wellcome Trust / ISSF Bridge Funding award at the University of Cambridge (2019). JGO

acknowledges financial support from the Spanish Ministry of Science and Innovation and FEDER

(grant PGC2018-101251-B-I00), by the Maria de Maeztu Programme for Units of Excellence in R&D

(grant CEX2018-000792-M), and by the Generalitat de Catalunya (ICREA Academia programme).

Library preparation and next-generation sequencing for single-cell RNA-seq analysis were

performed by the Single Cell Genomics Group at the National Centre for Genomic Analysis – Centre

for Genomic Regulation (CNAG-CRG), Barcelona. The Authors wish to thank Susanne van den Brink

(UPF, Barcelona) for helpful discussions. The Authors also wish to acknowledge Tina Balayo, Ana

Filipa Domingues, Oliver Davies, Kristen Place and Remisha Gurung's technical support at different

stages of the project.

AUTHOR CONTRIBUTIONS

Conceptualization: CP, AMA; Methodology: CWS, DR, GTC, KRK, JGO, AMA, CP; Software: GTC,

JGO; Validation: CWS, DR, CP; Investigation: CWS, DR, GTC, LD, CB, G-AI, CP; Formal Analysis:

GTC, DR, JGO, CP; Resources: GTC, JC, JGO, AMA; Data curation: DR, GTC, JGO; Writing –

Original Draft: CP, DR, GTC; Writing – Review and editing: CP, DR, GTC, AB, JGO, AMA;

Visualisation: DR, GTC, CWS, CP; Supervision: CP; Project administration: CP, AMA, JGO; Funding

acquisition: CP, AMA.

DECLARATION OF INTERESTS

AMA and CP are co-inventors in the patent application PCT/GB2019/052668: Polarised Three-

Dimensional Cellular Aggregates. The other authors have no interests to declare.

REFERENCES

Anonymous (2018). Method of the Year 2017: Organoids. Nature Methods *15,* 1.

Abdullah Obaid Khan, Michela Colombo, Jasmeet S Reyat, Guanlin Wang, Antonio Rodriguez-Romera, Wei Xiong Wen,
Lauren Murphy, Beata Grygielska, Christopher Mahony, Andrew Stone*, et al.* (2022). Human bone marrow organoids for
disease modelling, discovery and validation of therapeutic targets in hematological malignancies. bioRxiv

Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput sequencing data.
Bioinformatics *31,* 166-169.

Arias, A.M., Marikawa, Y., and Moris, N. (2022). Gastruloids: Pluripotent stem cell models of mammalian gastrulation and
embryo engineering. Dev. Biol. *488,* 35-46.

Baillie-Johnson, P., van den Brink, S C, Balayo, T., Turner, D.A., and Martinez Arias, A. (2015). Generation of Aggregates
of Mouse Embryonic Stem Cells that Show Symmetry Breaking, Polarization and Emergent Collective Behaviour In Vitro.
J. Vis. Exp. *(105). doi,* 10.3791/53252.

Balgobind, B.V., Van den Heuvel-Eibrink, M M, De Menezes, R.X., Reinhardt, D., Hollink, I.H., Arentsen-Peters, S.T., van
Wering, E.R., Kaspers, G.J., Cloos, J., de Bont, E.S*., et al.* (2011). Evaluation of gene expression signatures predictive of
cytogenetic and molecular subtypes of pediatric acute myeloid leukemia. Haematologica *96,* 221-230.

Barker, N., Huch, M., Kujala, P., van de Wetering, M., Snippert, H.J., van Es, J.H., Sato, T., Stange, D.E., Begthel, H., van
den Born, M*., et al.* (2010). Lgr5(+ve) stem cells drive self-renewal in the stomach and build long-lived gastric units in vitro.
Cell. Stem Cell. *6,* 25-36.

Beccari, L., Moris, N., Girgin, M., Turner, D.A., Baillie-Johnson, P., Cossy, A.C., Lutolf, M.P., Duboule, D., and Arias, A.M.
(2018). Multi-axial self-organization properties of mouse embryonic stem cells into gastruloids. Nature *562,* 272-276.

Boiers, C., Richardson, S.E., Laycock, E., Zriwil, A., Turati, V.A., Brown, J., Wray, J.P., Wang, D., James, C., Herrero, J*.,*
et al. (2018). A Human IPS Model Implicates Embryonic B-Myeloid Fate Restriction as Developmental Susceptibility to B
Acute Lymphoblastic Leukemia-Associated ETV6-RUNX1. Dev. Cell. *44,* 362-377.e7.

Britten, O., Ragusa, D., Tosi, S., and Kamel, Y.M. (2019). MLL -Rearranged Acute Leukemia with t(4;11)(q21;q23)-Current
Treatment Options. Is There a Role for CAR-T Cell Therapy? Cells *8,* 1341.

Cao, J., O'Day, D.R., Pliner, H.A., Kingsley, P.D., Deng, M., Daza, R.M., Zager, M.A., Aldinger, K.A., Blecher-Gonen, R.,
Zhang, F*., et al.* (2020). A human cell atlas of fetal gene expression. Science *370,* 10.1126/science.aba7721.

Cazzola, A., Cazzaniga, G., Biondi, A., Meneveri, R., Brunelli, S., and Azzoni, E. (2021). Prenatal Origin of Pediatric
Leukemia: Lessons From Hematopoietic Development. Front. Cell. Dev. Biol. *8,* 618164.

Chandrakanthan, V., Rorimpandey, P., Zanini, F., Chacon, D., Olivier, J., Joshi, S., Kang, Y.C., Knezevic, K., Huang, Y.,
Qiao, Q*., et al.* (2022). Mesoderm-derived PDGFRA(+) cells regulate the emergence of hematopoietic stem cells in the
dorsal aorta. Nat. Cell Biol. *24,* 1211-1225.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and Ma'ayan, A. (2013). Enrichr:
interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics *14,* 128-128.

Chen, M.J., Yokomizo, T., Zeigler, B.M., Dzierzak, E., and Speck, N.A. (2009). Runx1 is required for the endothelial to
haematopoietic cell transition but not thereafter. Nature *457,* 887-891.

Chou, D.B., Frismantas, V., Milton, Y., David, R., Pop-Damkov, P., Ferguson, D., MacDonald, A., Vargel Bolukbasi, O.,
Joyce, C.E., Moreira Teixeira, L.S*., et al.* (2020). On-chip recapitulation of clinical bone marrow toxicities and patient-
specific pathophysiology. Nat. Biomed. Eng. *4,* 394-406.

Clarke, R.L., Yzaguirre, A.D., Yashiro-Ohtani, Y., Bondue, A., Blanpain, C., Pear, W.S., Speck, N.A., and Keller, G. (2013).
The expression of Sox17 identifies and regulates haemogenic endothelium. Nature Cell Biology *15,* 502-510.

Clevers, H. (2016). Modeling Development and Disease with Organoids. Cell *165,* 1586-1597.

Clevers, H.C. (2019). Organoids: Avatars for Personalized Medicine. Keio Journal of Medicine *68,* 95.

Costa, G., Kouskoff, V., and Lacaud, G. (2012). Origin of blood cells and HSC production in the embryo. Trends in
Immunology *33,* 215-223.

Dalby, A., Ballester-Beltran, J., Lincetto, C., Mueller, A., Foad, N., Evans, A., Baye, J., Turro, E., Moreau, T., Tijssen, M.R.,
and Ghevaert, C. (2018). Transcription Factor Levels after Forward Programming of Human Pluripotent Stem Cells with
GATA1, FLI1, and TAL1 Determine Megakaryocyte versus Erythroid Cell Fate Decision. Stem Cell. Reports *11,* 1462-
1478.

de Pater, E., Kaimakis, P., Vink, C.S., Yokomizo, T., Yamada-Inagawa, T., van der Linden, R., Kartalaei, P.S., Camper, S.A., Speck, N., and Dzierzak, E. (2013). Gata2 is required for HSC generation and survival. J. Exp. Med. *210,* 2843-2850.

Dzierzak, E., and Bigas, A. (2018). Blood Development: Hematopoietic Stem Cell Dependence and Independence. Cell. Stem Cell. *22,* 639-651.

Espersen, A.D.L., Noren-Nystrom, U., Abrahamsson, J., Ha, S.Y., Pronk, C.J., Jahnukainen, K., Jonsson, O.G., Lausen, B., Palle, J., Zeller, B., Palmqvist, L., and Hasle, H. (2018). Acute myeloid leukemia (AML) with t(7;12)(q36;p13) is associated with infancy and trisomy 19: Data from Nordic Society for Pediatric Hematology and Oncology (NOPHO-AML) and review of the literature. Genes Chromosomes Cancer *57,* 359-365.

Fadlullah, M.Z.H., Neo, W.H., Lie-A-Ling, M., Thambyrajah, R., Patel, R., Mevel, R., Aksoy, I., Do Khoa, N., Savatier, P., Fontenille, L.*, et al.* (2022). Murine AGM single-cell profiling identifies a continuum of hemogenic endothelium differentiation marked by ACE. Blood *139,* 343-356.

Fehling, H.J., Lacaud, G., Kubo, A., Kennedy, M., Robertson, S., Keller, G., and Kouskoff, V. (2003). Tracking mesoderm induction and its specification to the hemangioblast during embryonic stem cell differentiation. Development *130,* 4217-4227.

Fitch, S.R., Kimber, G.M., Wilson, N.K., Parker, A., Mirshekar-Syahkal, B., Gottgens, B., Medvinsky, A., Dzierzak, E., and Ottersbach, K. (2012). Signaling from the sympathetic nervous system regulates hematopoietic stem cell emergence during embryogenesis. Cell. Stem Cell. *11,* 554-566.

Fornerod, M., Ma, J., Noort, S., Liu, Y., Walsh, M.P., Shi, L., Nance, S., Liu, Y., Wang, Y., Song, G.*, et al.* (2021). Integrative Genomic Analysis of Pediatric Myeloid-Related Acute Leukemias Identifies Novel Subtypes and Prognostic Indicators. Blood Cancer. Discov. *2,* 586-599.

Frankish, A., Diekhans, M., Ferreira, A.M., Johnson, R., Jungreis, I., Loveland, J., Mudge, J.M., Sisu, C., Wright, J., Armstrong, J.*, et al.* (2019). GENCODE reference annotation for the human and mouse genomes. Nucleic Acids Res. *47,* D766-D773.

Franzen, O., Gan, L.M., and Bjorkegren, J.L.M. (2019). PanglaoDB: a web server for exploration of mouse and human single-cell RNA sequencing data. Database (Oxford) *2019,* 10.1093/database/baz046.

Garcia-Alegria, E., Menegatti, S., Fadlullah, M.Z.H., Menendez, P., Lacaud, G., and Kouskoff, V. (2018). Early Human Hemogenic Endothelium Generates Primitive and Definitive Hematopoiesis In Vitro. Stem Cell. Reports *11,* 1061-1074.

Girgin, M.U., Broguiere, N., Mattolini, L., and Lutolf, M.P. (2021). Gastruloids generated without exogenous Wnt activation develop anterior neural tissues. Stem Cell. Reports *16,* 1143-1155.

Gotoh, S., Ito, I., Nagasaki, T., Yamamoto, Y., Konishi, S., Korogi, Y., Matsumoto, H., Muro, S., Hirai, T., Funato, M.*, et al.* (2014). Generation of alveolar epithelial spheroids via isolated progenitor cells from human pluripotent stem cells. Stem Cell. Reports *3,* 394-403.

Greggio, C., De Franceschi, F., Figueiredo-Larsen, M., and Grapin-Botton, A. (2014). In vitro pancreas organogenesis from dispersed mouse embryonic progenitors. J. Vis. Exp. *(89). doi,* 10.3791/51725.

Guibentif, C., Rönn, R.E., Böiers, C., Lang, S., Saxena, S., Soneji, S., Enver, T., Karlsson, G., and Woods, N. (2017). Single-Cell Analysis Identifies Distinct Stages of Human Endothelial-to-Hematopoietic Transition. Cell Reports *19,* 10-19.

Gulino, G.M., Bruno, F., Sturiale, V., Brancato, D., Ragusa, D., Tosi, S., Saccone, S., and Federico, C. (2021). From FISH to Hi-C: The Chromatin Architecture of the Chromosomal Region 7q36.3, Frequently Rearranged in Leukemic Cells, Is Evolutionary Conserved. Int. J. Mol. Sci. *22,* 10.3390/ijms22052338.

Hall, T.D., Kim, H., Dabbah, M., Myers, J.A., Crawford, J.C., Morales-Hernandez, A., Caprio, C.E., Sriram, P., Kooienga, E., Derecka, M.*, et al.* (2022). Murine fetal bone marrow does not support functional hematopoietic stem and progenitor cells until birth. Nature Communications *13,* 5403.

Hooper, M., Hardy, K., Handyside, A., Hunter, S., and Monk, M. (1987). HPRT-deficient (Lesch-Nyhan) mouse embryos derived from germline colonization by cultured cells. Nature (London) *326,* 292-295.

Huch, M., Bonfanti, P., Boj, S.F., Sato, T., Loomans, C.J., van de Wetering, M., Sojoodi, M., Li, V.S., Schuijers, J., Gracanin,
A., *et al.* (2013). Unlimited in vitro expansion of adult bi-potent pancreas progenitors through the Lgr5/R-spondin axis.
Embo J. *32,* 2708-2721.

Huch, M., Dorrell, C., Boj, S.F., van Es, J.H., Li, V.S., van de Wetering, M., Sato, T., Hamer, K., Sasaki, N., Finegold, M.J.,
et al. (2013). In vitro expansion of single Lgr5+ liver stem cells induced by Wnt-driven regeneration. Nature *494,* 247-250.

Huch, M., Knoblich, J.A., Lutolf, M.P., and Martinez-Arias, A. (2017). The hope and the hype of organoid research.
Development *144,* 938-941.

Ingenhag, D., Reister, S., Auer, F., Bhatia, S., Wildenhain, S., Picard, D., Remke, M., Hoell, J.I., Kloetgen, A., Sohn, D., *et*
al. (2019). The homeobox transcription factor HB9 induces senescence and blocks differentiation in hematopoietic stem
and progenitor cells. Haematologica *104,* 35-46.

Jakobsson, L., Franco, C.A., Bentley, K., Collins, R.T., Ponsioen, B., Aspalter, I.M., Rosewell, I., Busse, M., Thurston, G.,
Medvinsky, A., Schulte-Merker, S., and Gerhardt, H. (2010). Endothelial cells dynamically compete for the tip cell position
during angiogenic sprouting. Nat. Cell Biol. *12,* 943-953.

Kapeni, C., Nitsche, L., Kilpatrick, A.M., Wilson, N.K., Xia, K., Mirshekar-Syahkal, B., Chandrakanthan, V., Malouf, C.,
Pimanda, J.E., Gottgens, B., *et al.* (2022). p57Kip2 regulates embryonic blood stem cells by controlling sympathoadrenal
progenitor expansion. Blood *140,* 464-477.

Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S.L. (2013). TopHat2: accurate alignment of
transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. *14,* R36-r36.

Kim, J., Koo, B.K., and Knoblich, J.A. (2020). Human organoids: model systems for human biology and medicine. Nat.
Rev. Mol. Cell Biol. *21,* 571-584.

Kingsley, P.D., Malik, J., Emerson, R.L., Bushnell, T.P., McGrath, K.E., Bloedorn, L.A., Bulger, M., and Palis, J. (2006).
"Maturational" globin switching in primary primitive erythroid cells. Blood *107,* 1665-1672.

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S.L., Jagodnik,
K.M., Lachmann, A., *et al.* (2016). Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic
Acids Res. *44,* 90.

Lacaud, G., and Kouskoff, V. (2017). Hemangioblast, hemogenic endothelium, and primitive versus definitive
hematopoiesis. Exp. Hematol. *49,* 19-24.

Lancaster, M.A., and Knoblich, J.A. (2014). Organogenesis in a dish: modeling development and disease using organoid
technologies. Science *345,* 1247125.

Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E., Homfray, T., Penninger, J.M., Jackson,
A.P., and Knoblich, J.A. (2013). Cerebral organoids model human brain development and microcephaly. Nature *501,* 373-
379.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat. Methods *9,* 357-359.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and 1000 Genome
Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25,* 2078-
2079.

Lis, R., Karrasch, C.C., Poulos, M.G., Kunar, B., Redmond, D., Duran, J.G.B., Badwe, C.R., Schachterle, W., Ginsberg,
M., Xiang, J., *et al.* (2017). Conversion of adult endothelium to immunocompetent haematopoietic stem cells. Nature *545,*
439-445.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. Genome Biol. *15,* 550-8.

Marcelo, K.L., Sills, T.M., Coskun, S., Vasavada, H., Sanglikar, S., Goldie, L.C., and Hirschi, K.K. (2013). Hemogenic
endothelial cell specification requires c-Kit, Notch signaling, and p27-mediated cell-cycle control. Dev. Cell. *27,* 504-515.

McGrath, K.E., Frame, J.M., Fegan, K.H., Bowen, J.R., Conway, S.J., Catherman, S.C., Kingsley, P.D., Koniski, A.D., and
Palis, J. (2015). Distinct Sources of Hematopoietic Progenitors Emerge before HSCs and Provide Functional Blood Cells
in the Mammalian Embryo. Cell. Rep. *11,* 1892-1904.

Medvinsky, A.L., Samoylina, N.L., Muller, A.M., and Dzierzak, E.A. (1993). An early pre-liver intraembryonic source of
CFU-S in the developing mouse. Nature *364,* 64-67.

Medvinsky, A., and Dzierzak, E. (1996). Definitive hematopoiesis is autonomously initiated by the AGM region. Cell *86,*
897-906.

Medvinsky, A., Rybtsov, S., and Taoudi, S. (2011). Embryonic origin of the adult hematopoietic system: advances and
questions. Development (Cambridge) *138,* 1017-1031.

Mikkola, H.K., Fujiwara, Y., Schlaeger, T.M., Traver, D., and Orkin, S.H. (2003). Expression of CD41 marks the initiation
of definitive hematopoiesis in the mouse embryo. Blood *101,* 508-516.

Moris, N., Edri, S., Seyres, D., Kulkarni, R., Domingues, A.F., Balayo, T., Frontini, M., and Pina, C. (2018). Histone
Acetyltransferase KAT2A Stabilizes Pluripotency with Control of Transcriptional Heterogeneity. Stem Cells *36,* 1828-1838.

Moris, N., Anlas, K., van den Brink, Susanne C, Alemany, A., Schröder, J., Ghimire, S., Balayo, T., van Oudenaarden, A.,
and Martinez Arias, A. (2020). An in vitro model of early anteroposterior organization during human development. Nature
(London) *582,* 410-415.

Niakan, K.K., Ji, H., Maehr, R., Vokes, S.A., Rodolfa, K.T., Sherwood, R.I., Yamaki, M., Dimos, J.T., Chen, A.E., Melton,
D.A., McMahon, A.P., and Eggan, K. (2010). Sox17 promotes differentiation in mouse embryonic stem cells by directly
regulating extraembryonic gene expression and indirectly antagonizing self-renewal. Genes Dev. *24,* 312-326.

Nilsson, T., Waraky, A., Ostlund, A., Li, S., Staffas, A., Asp, J., Fogelstrand, L., Abrahamsson, J., and Palmqvist, L. (2022).
An induced pluripotent stem cell t(7;12)(q36;p13) acute myeloid leukemia model shows high expression of MNX1 and a
block in differentiation of the erythroid and megakaryocytic lineages. Int. J. Cancer *151,* 770-782.

Nobuhisa, I., Osawa, M., Uemura, M., Kishikawa, Y., Anani, M., Harada, K., Takagi, H., Saito, K., Kanai-Azuma, M., Kanai,
Y., Iwama, A., and Taga, T. (2014). Sox17-Mediated Maintenance of Fetal Intra-Aortic Hematopoietic Cell Clusters.
Molecular and Cellular Biology *34,* 1976-1990.

North, T.E., Goessling, W., Peeters, M., Li, P., Ceol, C., Lord, A.M., Weber, G.J., Harris, J., Cutting, C.C., Huang, P.,
Dzierzak, E., and Zon, L.I. (2009). Hematopoietic Stem Cell Development Is Dependent on Blood Flow. Cell *137,* 736-748.

Papatsenko, D., Darr, H., Kulakovskiy, I., Waghray, A., Makeev, V., MacArthur, B., and Lemischka, I. (2015). Single-Cell
Analyses of ESCs Reveal Alternative Pluripotent Cell States and Molecular Mechanisms that Control Self-Renewal. Stem
Cell Reports *5,* 207-220.

Pearson, S., Cuvertino, S., Fleury, M., Lacaud, G., and Kouskoff, V. (2015). In vivo repopulating activity emerges at the
onset of hematopoietic specification during embryonic stem cell differentiation. Stem Cell. Reports *4,* 431-444.

Picelli, S., Faridani, O.R., Bjorklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R. (2014). Full-length RNA-seq from
single cells using Smart-seq2. Nat. Protoc. *9,* 171-181.

Pina, C., May, G., Soneji, S., Hong, D., and Enver, T. (2008). MLLT3 regulates early human erythroid and megakaryocytic
cell fate. Cell. Stem Cell. *2,* 264-273.

Pina, C., Fugazza, C., Tipping, A.J., Brown, J., Soneji, S., Teles, J., Peterson, C., and Enver, T. (2012). Inferring rules of
lineage commitment in haematopoiesis. Nature Cell Biology *14,* 287-294.

Pina, C., Teles, J., Fugazza, C., May, G., Wang, D., Guo, Y., Soneji, S., Brown, J., Edén, P., Ohlsson, M., Peterson, C.,
and Enver, T. (2015). Single-Cell Network Analysis Identifies DDIT3 as a Nodal Lineage Regulator in Hematopoiesis. Cell
Reports *11,* 1503-1510.

Putri, G.H., Anders, S., Pyl, P.T., Pimanda, J.E., and Zanini, F. (2022). Analysing high-throughput sequencing data in
Python with HTSeq 2.0. Bioinformatics *38,* 2943-2945.

Ragusa, D., Ciciro, Y., Federico, C., Saccone, S., Bruno, F., Saeedi, R., Sisu, C., Pina, C., Sala, A., and Tosi, S. (2022).
Engineered model of t(7;12)(q36;p13) AML recapitulates patient-specific features and gene expression profiles.
Oncogenesis *11,* 50-2.

Rivron, N.C., Frias-Aldeguer, J., Vrij, E.J., Boisset, J.C., Korving, J., Vivie, J., Truckenmuller, R.K., van Oudenaarden, A.,
van Blitterswijk, C.A., and Geijsen, N. (2018). Blastocyst-like structures generated solely from stem cells. Nature *557,* 106-
111.

Rossi, G., Broguiere, N., Miyamoto, M., Boni, A., Guiet, R., Girgin, M., Kelly, R.G., Kwon, C., and Lutolf, M.P. (2021).
Capturing Cardiogenesis in Gastruloids. Cell. Stem Cell. *28,* 230-240.e6.

Rossi, G., Giger, S., Hubscher, T., and Lutolf, M.P. (2022). Gastruloids as in vitro models of embryonic blood development
with spatial and temporal resolution. Sci. Rep. *12,* 13380-1.

Rybtsov, S., Sobiesiak, M., Taoudi, S., Souilhol, C., Senserrich, J., Liakhovitskaia, A., Ivanovs, A., Frampton, J., Zhao, S.,
and Medvinsky, A. (2011). Hierarchical organization and early hematopoietic specification of the developing HSC lineage
in the AGM region. J. Exp. Med. *208,* 1305-1315.

Rybtsov, S., Batsivari, A., Bilotkach, K., Paruzina, D., Senserrich, J., Nerushev, O., and Medvinsky, A. (2014). Tracing the
Origin of the HSC Hierarchy Reveals an SCF-Dependent, IL-3-Independent CD43– Embryonic Precursor. Stem Cell
Reports *3,* 489-501.

Sato, T., Vries, R.G., Snippert, H.J., van de Wetering, M., Barker, N., Stange, D.E., van Es, J.H., Abo, A., Kujala, P., Peters,
P.J., and Clevers, H. (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche.
Nature *459,* 262-265.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. Nat. Methods
9, 671-675.

Silva, A.C., Matthys, O.B., Joy, D.A., Kauss, M.A., Natarajan, V., Lai, M.H., Turaga, D., Blair, A.P., Alexanian, M., Bruneau,
B.G., and McDevitt, T.C. (2021). Co-emergence of cardiac and gut tissues promotes cardiomyocyte maturation within
human iPSC-derived organoids. Cell. Stem Cell. *28,* 2137-2152.e6.

Souilhol, C., Gonneau, C., Lendinez, J.G., Batsivari, A., Rybtsov, S., Wilson, H., Morgado-Palacin, L., Hills, D., Taoudi, S.,
Antonchuk, J., Zhao, S., and Medvinsky, A. (2016). Inductive interactions mediated by interplay of asymmetric signalling
underlie development of adult haematopoietic stem cells. Nat. Commun. *7,* 10784.

Sroczynska, P., Lancrin, C., Pearson, S., Kouskoff, V., and Lacaud, G. (2009). In vitro differentiation of mouse embryonic
stem cells as a model of early hematopoietic development. Methods Mol. Biol. *538,* 317-334.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub,
T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting
genome-wide expression profiles. Proc. Natl. Acad. Sci. U. S. A. *102,* 15545-15550.

Sugimura, R., Jha, D.K., Han, A., Soria-Valles, C., da Rocha, E.L., Lu, Y.F., Goettel, J.A., Serrao, E., Rowe, R.G.,
Malleshaiah, M.*, et al.* (2017). Haematopoietic stem and progenitor cells from human pluripotent stem cells. Nature *545,*
432-438.

Sugimura, R., Ohta, R., Mori, C., Li, A., Mano, T., Sano, E., Kosugi, K., Nakahata, T., Niwa, A., Saito, M.K., and Torisawa,
Y.S. (2020). Biomimetic aorta-gonad-Mesonephros-on-a-Chip to study human developmental hematopoiesis. Biomed.
Microdevices *22,* 34-2.

Swiers, G., Baumann, C., O'Rourke, J., Giannoulatou, E., Taylor, S., Joshi, A., Moignard, V., Pina, C., Bee, T., Kokkaliaris,
K.D.*, et al.* (2013). Early dynamic fate changes in haemogenic endothelium characterized at the single-cell level. Nature
Communications *4,* 2924.

Taketani, T., Taki, T., Sako, M., Ishii, T., Yamaguchi, S., and Hayashi, Y. (2008). MNX1-ETV6 fusion gene in an acute
megakaryoblastic leukemia and expression of the MNX1 gene in leukemia and normal B cell lines. Cancer Genet.
Cytogenet. *186,* 115-119.

Taoudi, S., Gonneau, C., Moore, K., Sheridan, J.M., Blackburn, C.C., Taylor, E., and Medvinsky, A. (2008). Extensive Hematopoietic Stem Cell Generation in the AGM Region via Maturation of VE-Cadherin+CD45+ Pre-Definitive HSCs. Cell Stem Cell *3,* 99-108.

Thomas, P.D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L., and Mi, H. (2022). PANTHER: Making genome-scale phylogenetics accessible to all. Protein Science *31,* 8-22.

Torisawa, Y.S., Spina, C.S., Mammoto, T., Mammoto, A., Weaver, J.C., Tat, T., Collins, J.J., and Ingber, D.E. (2014). Bone marrow-on-a-chip replicates hematopoietic niche physiology in vitro. Nat. Methods *11,* 663-669.

Turner, D.A., Girgin, M., Alonso-Crisostomo, L., Trivedi, V., Baillie-Johnson, P., Glodowski, C.R., Hayward, P.C., Collignon, J., Gustavsen, C., Serup, P.*, et al.* (2017). Anteroposterior polarity and elongation in the absence of extraembryonic tissues and spatially localised signalling in Gastruloids , mammalian embryonic organoids. Development (Cambridge) *144,* 3894-3906.

van den Brink, S C, Alemany, A., van Batenburg, V., Moris, N., Blotenburg, M., Vivie, J., Baillie-Johnson, P., Nichols, J., Sonnen, K.F., Martinez Arias, A., and van Oudenaarden, A. (2020). Single-cell and spatial transcriptomics reveal somitogenesis in gastruloids. Nature *582,* 405-409.

van den Brink, S C, Baillie-Johnson, P., Balayo, T., Hadjantonakis, A.K., Nowotschin, S., Turner, D.A., and Martinez Arias, A. (2014). Symmetry breaking, germ layer specification and axial organisation in aggregates of mouse embryonic stem cells. Development *141,* 4231-4242.

van den Brink, S C, and van Oudenaarden, A. (2021). 3D gastruloids: a novel frontier in stem cell-based in vitro modeling of mammalian gastrulation. Trends Cell Biol. *31,* 747-759.

Vink, C.S., Calero-Nieto, F.J., Wang, X., Maglitto, A., Mariani, S.A., Jawaid, W., Gottgens, B., and Dzierzak, E. (2020). Iterative Single-Cell Analyses Define the Transcriptome of the First Functional Hematopoietic Stem Cells. Cell. Rep. *31,* 107627.

Wang, D., Tanaka-Yano, M., Meader, E., Kinney, M.A., Morris, V., Lummertz da Rocha, E., Liu, N., Liu, T., Zhu, Q., Orkin, S.H.*, et al.* (2022). Developmental maturation of the hematopoietic system controlled by a Lin28b-let-7-Cbx2 axis. Cell Reports (Cambridge) *39,* 110587.

Waraky, A., Ostlund, A., Nilsson, T., Weichenhan, D., Lutsik, P., Bahr, M., Hey, J., Adamsson, J., Hamdy, M., Morsy, A.*, et al.* (2022). Mnx1 Induces Leukemia Transformation Through Altering Histone Methylation in a Model of Pediatric t(7;12) Acute Myeloid Leukemia. BiorXiv

Wildenhain, S., Ingenhag, D., Ruckert, C., Degistirici, O., Dugas, M., Meisel, R., Hauer, J., and Borkhardt, A. (2012). Homeobox protein HB9 binds to the prostaglandin E receptor 2 promoter and inhibits intracellular cAMP mobilization in leukemic cells. J. Biol. Chem. *287,* 40703-40712.

Wildenhain, S., Ruckert, C., Rottgers, S., Harbott, J., Ludwig WD FASchuster, F R, Schuster, F.R., Beldjord, K., Binder, V., Slany, R., Hauer, J., and Borkhardt, A. (2010). Expression of cell-cell interacting genes distinguishes HLXB9/TEL from MLL-positive childhood acute myeloid leukemia. Leukemia *24,* 1657-1660.

Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., Jeon, M., and Ma'ayan, A. (2021). Gene Set Knowledge Discovery with Enrichr. Curr. Protoc. *1,* e90.

Yu, L., Wei, Y., Duan, J., Schmitz, D.A., Sakurai, M., Wang, L., Wang, K., Zhao, S., Hon, G.C., and Wu, J. (2021). Blastocyst-like structures generated from human pluripotent stem cells. Nature *591,* 620-626.

Zhu, Q., Gao, P., Tober, J., Bennett, L., Chen, C., Uzun, Y., Li, Y., Howell, E.D., Mumau, M., Yu, W.*, et al.* (2020). Developmental trajectory of prehematopoietic stem cell formation from endothelium. Blood *136,* 845-856.

908 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-mouse c-Kit (CD117) clone 2B8 - APC-Cy7	Biolegend	Cat. #105825
Anti-mouse CD41 clone MWReg30 - PE-Dazzle594	Biolegend	Cat. #133935
Anti-mouse CD45 clone 30F11 - APC	Biolegend	Cat. #103111
Anti-mouse CD144 (VE-cadherin) clone BV13 - APC	Biolegend	Cat. #138011
Anti-mouse CD43 clone S11 - PE	Biolegend	Cat. #143205
CD11b anti-mouse/human clone M1/70 - biotin	Biolegend	Cat. #101203
Mouse Anti-Mouse CD45.2 clone 104 - PE	BD Biosciences	Cat. #560695
Rat Anti-Mouse CD31 clone MEC13.3 - biotin	BD Biosciences	Cat. #553371
Goat Anti-Mouse c-Kit/CD117	R&D Systems	Cat. #AF1356SP
Bacterial and virus strains		
E. coli: NEB 5-alpha Competent *E. coli*	New England Biolabs (NEB)	Cat. #C2987H
Chemicals, peptides, and recombinant proteins		
Murine LIF	Peprotech	Cat. #250-02
StemMACS PD0325901	Miltenyi Biotec	Cat. #130-106-5411
Chiron (CHIR99021)	Biogems	Cat. #2520691
Activin A PLUS	QKine	Cat. #Qk005
Murine VEGF 165	Peprotech	Cat. #450-32
Murine FGF-basic	Peprotech	Cat. #450-33
Murine Sonic Hedgehog (Shh)	Peprotech	Cat. #315-22
Murine TPO	Peprotech	Cat. #315-14
Murine Flt3-Ligand	Peprotech	Cat. #250-31L
Murine SCF	Peprotech	Cat. #250-03
N2B27 medium (NDiff 227)	Takara Bio	Cat. #Y40002
Critical commercial assays		
Mouse Methylcellulose Complete Media	R&D Systems	Cat. #HSC007
Deposited data		
Single cell RNA-seq of hemogenic gastruloids at timepoints	This paper	ArrayExpress: E-MTAB-12148
Bulk RNA-seq of hemogenic gastruloids with MNX1 overexpression at 216 h	This paper	ArrayExpress: E-MTAB-12173
Single cell RNA-seq of hemogenic gastruloids with MNX1 overexpression at 216 h, sorted CD45+/CD41+	This paper	ArrayExpress: E-MTAB-12149
Experimental models: Cell lines		
Mus musculus: Flk-1-GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	(Jakobsson et al., 2010)
Mus musculus: Sox17-GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	(Niakan et al., 2010)
Mus musculus: T/Bra::GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	(Fehling et al., 2003)
Mus musculus: E14Tg2a mouse embryonic stem cell line	MMRRC, University of California Davis, US	(Hooper et al., 1987)
Homo sapiens: HEK293T	ATCC	CRL-3216
Oligonucleotides		
MNX1 (forward 5-GTTCAAGCTCAACAAGTACC-3; reverse 5- GGTTCTGGAACCAAATCTTC-3)	(Gulino et al., 2021)	n/a
Ppia (forward 5-TTACCCATCAAACCATTCCTTCTG-3; reverse 5-AACCCAAAGAACTTCAGTGAGAGC-3)	(Moris, N. et al., 2018)	n/a
Recombinant DNA		

pWPT-LSSmOrange-PQR	Ghevaert Lab, WT-MRC Cambridge Stem Cell Institute, UK	(Dalby et al., 2018)
pWPT-LSSmOrange-PQR-MNX1	Cloned by Biomatik Corporation (Kitchener, Canada)	n/a
Software and algorithms		
GSEA software v4.2.3	(Subramanian et al., 2005)	https://www.gsea-msigdb.org/gsea/index.jsp
ImageJ	(Schneider et al., 2012)	https://imagej.nih.gov/ij/
Bowtie2	(Langmead and Salzberg, 2012)	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
Samtools	(Li et al., 2009)	http://samtools.sourceforge.net/
Tophat2	(Kim, D. et al., 2013)	https://ccb.jhu.edu/software/tophat/index.shtml
HTseq	(Anders et al., 2015)	https://htseq.readthedocs.io/en/master/
DEseq2	(Love et al., 2014)	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
ExpressAnalyst	n/a	https://www.expressanalyst.ca/
PANTHER	(Thomas et al., 2022)	http://www.pantherdb.org/
EnrichR	(Chen et al., 2013; Kuleshov et al., 2016; Xie et al., 2021)	https://maayanlab.cloud/Enrichr/
Panglao BP	(Franzen et al., 2019)	https://panglaodb.se
Descartes	(Cao et al., 2020)	https://descartes.brotmanbaty.org/
TARGET (Therapeutically Applicable Research to Generate Effective Treatments) Database	https://ocg.cancer.gov/programs/target https://xenabrowser.net/	GDC TARGET-AML
Single-cell RNA sequencing analysis scripts	DockerHub: dsblab/single_cell_analysis:0.5	https://github.com/dsb-lab/blood_gastruloids
Other		
CELLSTAR cell-repellent cell culture plate, 96 well, U-bottom	Greiner (Bio-One)	Cat. #650970

METHODS

Cell culture

Flk-1-GFP (Jakobsson et al., 2010), Sox17-GFP (Niakan et al., 2010), T/Bra::GFP (Fehling et al.,

2003), and E14Tg2A (Hooper et al., 1987) mouse embryonic stem cells (mES) lines were cultured

in ES+LIF medium in gelatinized (0.1% gelatin) with daily medium change, as previously described

(Turner et al., 2017). The ES+LIF medium contained 500 ml Glasgow MEM BHK-21 (Gibco), 50 ml

of fetal bovine serum (FBS, Embryonic Stem Cells tested, biosera, Nuaillé, France), 5 ml GlutaMAX

supplement (Gibco), 5 ml MEM Non-Essential Amino Acids (Gibco), 5 ml Sodium pyruvate solution

(100 mM, Gibco), and 1 ml 2-Mercaptoethanol (50mM, Gibco). Murine LIF (Peprotech) was added

at 1000 U/ml. HEK293T cells were grown in Dulbecco's modified Eagle medium (Gibco)

supplemented with 10% FBS (Gibco) and 1% penicillin/streptomycin (Gibco). All cultures were kept

at 37°C and 5% CO2.

Hemogenic gastruloids assembly, culture, and dissociation

mES were maintained in ES+LIF medium and transferred to 2i+LIF (containing Chiron and MEK

inhibitor PD03) for 24 hours prior to the assembly into gastruloids. 250 cells were seeded in each

well of a U-bottom, cell-repellent 96-well plate (Greiner Bio-One, Stonehouse, UK) in 40 µl of N2B27

medium (Takara Bio). The plate was centrifuged at 750 rpm for 2 minutes to promote deposition and

aggregation of the cells and was then incubated at 37°C, 5% CO_2 for 48 hours. After 48 hours, 150

µl of N2B27 medium supplemented with 100 ng/ml Activin A (QKine, Cambridge, UK) and 3 µM

chiron (Peprotech) was added to each well. At 72 hours, 150 µl of medium were removed, without

disrupting the gastruloids in the wells. 100 µl of N2B27 with 5 ng/ml of Vegf and Fgf2 each

(Peprotech) were added to each well. From 72 h to 144 h, each day 100 µl of medium were removed

and replaced with N2B27 + Vegf + Fgf2. At 144 h, the medium was further supplemented with Shh

at 20 ng/ml. From 168 h to 216 h, the medium was N2B27 + 5 ng/ml Vegf, plus 20 ng/ml mTpo, 100

ng/ml mFlt3l, and 100 ng/ml mScf (Peprotech). To dissociate cells from the gastruloid structures,

medium was removed and individual gastruloids were collected using a pipette and precipitated at

the bottom of a microcentrifuge tube. The remaining medium was aspirated and the bulk of gastruloids was washed in PBS. 50 µl of TrypLE express was added to pelleted gastruloids to be incubated at 37°C for 2 minutes to dissociate cells.

Methylcellulose colony forming assays (CFC)

Disassembled gastruloids cells were plated on Mouse Methylcellulose Complete Media (R&D Systems). Cells were first suspended in Iscove's Modified Dulbecco's Medium (IMDM, Gibco) with 20% FBS (Gibco) before addition to the methylcellulose medium. Cells were plated in duplicate 35 mm dishes with $2x10^5$ cells/plate. Plates were incubated at 37°C and 5% CO2 for 10 days, when colonies were scored. For serial replating experiments, cells in methylcellulose were collected and washed in phosphate buffer saline (PBS) to achieve single-cell suspensions and replated as described above.

Immunofluorescence

Immunostaining of whole gastruloids was performed as described before (Baillie-Johnson et al., 2015). Briefly, gastruloids were fixed in 4% paraformaldehyde (PFA) dissolved in PBS for 4 hours at 4°C on orbital shaking and permeabilised in PBSFT (10% FBS and 0.2% Triton X-100), followed by one hour blocking in PBSFT at 4°C on orbital shaking. Antibody dilutions were made in PBSFT at 1:200 for primary and 1:500 for secondary antibodies. Antibody incubations were performed overnight at 4°C on orbital shaking, and subjected to optical clearing overnight in ScaleS4 clearing solution. Individual gastruloids were then mounted on glass coverslips by pipetting as droplets in ScaleS4 and DAPI nuclear stain.

Imaging

Images of cultured gastruloids and CFC plates were captured using the Cytation 5 Cell Imaging Multi-Mode Reader (Biotek) plate reader using bright field and FITC channels. ImageJ (Schneider et al., 2012) was used for gastruloid size quantification. Confocal microscopy was performed on

LSM700 on a Zeiss Axiovert 200 M with Zeiss EC Plan-Neofluar 10x/0.30 M27 and Zeiss LD Plan-

Neofluar 20x/0.4 M27 objective lens.

Lentiviral vector packaging and transduction

The lentiviral overexpression vector pWPT-LSSmOrange-PQR was used to clone the *MNX1* gene

cDNA. The viral packaging vectors pCMV and pMD2.G, described in (Pina, C. et al., 2008), were

used to assemble lentiviral particles using HEK293T cells via transfection using TurboFect Reagent

(Invitrogen). Transduction of mES cells was performed overnight by addition of lentivirus to culture

medium and washed the following day (Moris et al., 2018). Transduced cells were sorted for positivity

to LSSmOrange.

Flow cytometry

Surface cell marker analysis was performed by staining using the antibodies listed in Key Resources

Table. Disassembled gastruloids cells were resuspended in PBS, 2% FBS and 0.5 mM EDTA and

stained at a dilution of 1:100 for primary antibodies for 20 minutes at 4°C. When indicated,

streptavidin was added at a dilution of 1:200. Analysis was performed on ACEA Novocyte (Agilent)

or AttuneNxT (Thermo) analyzers, using the respective software packages. Cell sorting was

performed using a CS&T calibrated BDFACS Aria III system (488 nm 40 mW, 633 nm 20 mW, 405nm

30 mW, and 561 nm 50 mW), set with the 100μm nozzle at 20PSI and a 4-way purity mask. Single-

cell deposition in 96-well plates was performed using single-cell sorting mode. Intact cells were gated

on FSC-A vs SSC-A plot, followed by doublet exclusion on FSC-A vs FSC-H and SSC-A vs SSC-H,

prior to gating on fluorescent parameters for the markers described in the results.

Single cell RNA sequencing of time-resolved hemogenic gastruloids

Gastruloids were collected at different timepoints of the protocol, disassembled and FACS deposited

into 96-well plates, either as unsorted (global) or sorted by CD45, CD41 and c-Kit/ScaI markers

(sorted). RNA from the cells were extracted from single cells using Smart-seq2 technology at 500Kb

and depth of 151 bases. Sequencing reads were quality-checked using FastQC (v0.11.9). We

trimmed the samples using trimGalore! (0.6.6) with a cutoff of 30, clipping 15 base pairs and retaining

reads of more than 100 bases. Alignment was performed on STAR (2.7.8a) and *Mus muluscus*

annotations from GENCODE vM26. Aligned BAM files were annotated using featureCounts (v2.0.1)

and count matrices were computed in python by directly accessing the BAM files with gtfparse

packages and collapsing lanes.

Single cell RNA sequencing of MNX1 overexpressing hemogenic gastruloids at endpoint

MNX1 overexpressing gastruloids and empty vector (EV) control gastruloids were collected at 216

h, sorted by positivity to CD45 or CD41, and deposited into 96-well plates for sequencing. Processing

of reads was performed as described above, but retaining reads over 75 bases to account for the

sequencing depth of 101 bases.

1003

Single cell RNA sequencing analysis

For quality control, based on the histogram of counts and multimodality distributions, we set a

minimum count threshold of 200000 counts, a minimum threshold of 1000 expressed genes, and a

maximum threshold of 20% mitochondrial fraction per cell. We performed the same procedure of the

second dataset with a more restrictive threshold of 400000 counts and similar expressed genes and

mitochondrial thresholds. Cells that did not pass the quality control metrics were omitted from

analysis. We normalized the cells to the mean count number per dataset and applied a plus-one-log

transformation of the data before proceeding to the downstream analysis.

Dimensionality reduction was performed on feature selection of the gene space using the function

scanpy.highly_varying_genes with default parameters. Selection of principal components in principal

component analysis (PCA) was performed by heuristic elbow method. Nearest neighbor analysis

was constructed by KNN graphs using a correlation metric and 10 nearest neighbors. Data was

projected for low dimensional visualization using the UMAP algorithm with default parameters as

implemented in *scanpy.tl.umap*. We used the leiden algorithm as implemented in *scanpy.tl.leiden* to

partition the data into clusters. In order to assess the election of the resolution parameter we used

Newman-Girvan modularity as a metric of clustering quality. Differential expressed genes were

computed comparing each cluster against the rest using the Wilcoxon test with Benjamini-Hochberg

correction.

1022

Annotation and projection to additional datasets

Raw counts matrices from Fadlullah et al. (2022) and Vink et al. (2020) were downloaded and

processed following the same pipeline as described above for scRNA-seq of gastruloids to generate

UMAP and clustering. We implemented the scmap algorithm to compare our scRNA sequencing of

gastruloids with the available datasets. We reduced the dimensionality of the space by selecting

highly varying genes from the annotated dataset. Then, we constructed a KNN classifier with

correlation metric and computed the nearest neighbors of the target data. If neighbors with

correlation metrics below 0.7 default standards, the projected cells were not projected onto any cell

from the annotated dataset. To visualize the cells over the UMAP plots of the other datasets, we

constructed a KNN regressor with three neighbors and a correlation metric.

1033

Bulk RNA sequencing

Total RNA was extracted from disassembled gastruloid cells at 216 hours. Sequencing was

performed on NovaSeq PE150 platform, at 20M paired-end reads per sample. Tophat2 with Bowtie2

were used to map paired-end reads to the reference *Mus musculus* genome build GRCm39 (Kim et

al., 2013; Langmead and Salzberg, 2012) GENCODE Release M30 (Frankish et al., 2019) was used

as the reference mouse genome annotation. Aligned reads were filtered by quality using samtools

(Li et al., 2009) with a minimum threshold set at 30 (q30). Transcript assembly and quantification

was achieved using htseq (Putri et al., 2022). Differential expression between sample and control

was performed by collapsing technical replicates for each condition using Deseq2 (Love et al., 2014)

in R environment (*Deseq2* library v 1.32.0). The differential expression was expressed in the form of

log2 fold change and filtered by false discovery rate (FDR) of 0.1.

1045

Real-time polymerase chain reaction (qPCR)

Extracted RNA was reverse-transcribed into complementary DNA (cDNA) using High-Capacity RNA-to-cDNA Kit (Applied Biosystems). QPCR was performed using FastGene 2x IC Green Universal qPCR Mix (Nippon Genetics, Duren, Germany) using primers for human *MNX1* (forward 5-GTTCAAGCTCAACAAGTACC-3; reverse 5- GGTTCTGGAACCAAATCTTC-3) (Gulino et al., 2021) and *Ppia* for endogenous control (forward 5- TTACCCATCAAACCATTCCTTCTG-3; reverse 5-AACCCAAAGAACTTCAGTGAGAGC-3) (Moris et al., 2018). Differential gene expression was calculated using the delta delta Ct ($\Delta\Delta$Ct) method.

Gene list enrichment analyses

Gene ontology (GO) analysis was performed in ExpressAnalyst (available at www.expressanalyst.ca) using the PANTHER Biological Process (BP) repository. GO terms and pathways were filtered by false discovery rate (FDR) with a cut-off of ≤ 0.1 for meaningful association. EnrichR (Chen et al., 2013; Kuleshov et al., 2016; Xie et al., 2021) was used for cell type analysis using the Panglao DB (Franzen et al., 2019) and Descartes (Cao et al., 2020) databases using a FDR threshold of ≤ 0.1 or p value ≤ 0.01, where specified, as well as transcription factor (TF) binding site enrichment using the ENCODE and ChEA Consensus TFs from ChIP database.

Gene Set Enrichment Analysis (GSEA)

Custom gene signatures were used as gene sets for GSEA analysis (Subramanian et al., 2005) on the GSEA software v4.2.3 on RNA sequencing expression values in counts units GSEA was ran in 10000 permutations on gene set using the weighted Signal2Noise metric. Enrichment metrics are shown as normalized enrichment score (NES) and filtered by FDR ≤ 0.05. Leading edge genes (LEGs) are genes with a "Yes" values for core enrichment. For AML patient analysis, clinical phenotype and expression data (in counts units) were extracted from the GDC TARGET-AML cohorts in the Therapeutically Applicable Research to Generate Effective Treatments project (TARGET, https://ocg.cancer.gov/programs/target), downloaded from the University of California

1073   Santa Cruz (UCSC) Xena public repository (last accessed 31st August 2022). Patient samples were

1074   selected according to the reported karyotype to include t(7;12), inv(16), *MLL*, normal karyotype, and

1075   t(8;21). GSEA was performed comparing RNA sequencing counts of t(7;12) samples against pooled

1076   AML subtypes (inv(16), MLL, normal karyotype and t(8;21)) as "other AML".

1077

Lineage analysis from scRNA-seq

Genes with detectable expression in 216-hour CD41/CD45+ single MNX-OE or EV gastruloid cells

were manually scanned for lineage-affiliated markers and regulators conventionally associated with

the endothelial and hematopoietic lineages (Guibentif et al., 2017; Pina, Cristina et al., 2012; Pina,

Cristina et al., 2015; Swiers et al., 2013). Erythroid genes were: *Gata1*, *Klf1*, *Zfpm1*, *Tal1*, *Gfi1b* and

Nfe2 (transcription factors, TF), *Epor* (growth factor receptor, GFR), and *Epb42*, *Slc4a1*/Band3,

Gypa and *Gypc* (membrane-associated proteins, MAP), in addition to alpha (*Hba*) and beta globin

(*Hbb*) chains. Myeloid genes were *Spi1*, *Cebpa*, *Cebpe* and *Egr2* (TF), *Csf1r*, *Csf2ra*, *Csf2rb*, *Csf3r*

and *Il3ra* (GFR), *Fcgr1*, *Fcgr2b*, *Anpep*/Cd13, *Cd14*, *Cd33*, *Ly6c*, *Ly6g* and *Ly6a* (MAP), and

enzymes *Mpo*, *Lyz1* and *Lyz2*. Lymphoid genes were *Notch1*, *Gata3*, *Ebf1*, *Ebf2* and *Ikfz1* (TF),

Il2ra, *Il2rg* and *Il7r* (GFR), *Cd5*, *Cd7*, *Cd8a* and *Cd79a* (MBP), as well as surrogate (*Igll1*, *Vpreb1*)

and mature Ig chains (*Ighm*, *Ighd*, *IghJ*); variable regions of *Igl* and *Igh* chains were considered

separately as present or absent. Endothelial (*Flt1*, *Kdr*/Flt2, *Flt4*, *Cdh5*, *Esam*, *Epas1* and *Sox17*),

stem/progenitor (*Runx1*, *Cbfb*, *Myb* and *Gata2*, in addition to the HSC/erythroid-associated *Mllt3*),

and megakaryocytic genes (*Mpl*, *Fli1*, *Itga2b*/Cd41, *Itgb3*/Cd61) were also annotated. Lineage

signatures were called if 2 or more lineage genes were expressed. Erythroid lineage was separated

into marker/regulator (Reg) and Hemoglobin signatures, the latter requiring at least one *Hbb* and

one *Hba* chain. Priming for multiple lineages (erythroid, myeloid or lymphoid ± endothelial) was

quantified separately based on co-existence of lineage signatures. Megakaryocytic lineages were

not considered for multi-lineage priming due to systematic co-expression of markers in progenitor

and erythroid cells. Co-expression of endothelial and progenitor signatures were called endoHaem

if no or a single lineage signature were present.

1100

Statistical analysis

Experiments were performed at least in triplicates, unless specified otherwise. Data are plotted to include standard deviation (+/- SD) between replicates. Statistical significance was set at a threshold of p value < 0.05. Statistical analysis was performed in R environment (version 4.1.3) or using GraphPad Prism 8.0 software.

Data availability

Raw data as well as processed count matrices and post-processed files from single-cell RNA-seq for the time-resolved data is available at E-MTAB-12148. Single-cell RNA-seq for the MNX1 overexpression experiment is available at Array Express with accession code E-MTAB-12149. Bulk RNA-seq of MNX1 overexpressing gastruloids is available at Array Express with accession code E-MTAB-12173. The post-processing was performed in Python on DockerHub: dsblab/single_cell_analysis:0.5. Scripts are available in https://github.com/dsb-lab/blood_gastruloids and Zenodo (https://doi.org/10.5281/zenodo.7053423). The results published here are partly based upon data generated by the Therapeutically Applicable Research to Generate Effective Treatments (TARGET) (https://ocg.cancer.gov/programs/target) initiative, of the Acute Myeloid Leukemia (AML) cohort GDC TARGET-AML. The data used for this analysis are available at https://portal.gdc.cancer.gov/projects and https://xenabrowser.net/.