Dissecting infant leukemia developmental origins with a hemogenic gastruloid

model

Denise Ragusa[1,2], Chun-Wai Suen[3*], Gabriel Torregrosa-Cortés[4*], Fabio Pastorino[5*],*

Ayona Johns[1,2], Ylenia Cicirò[1,2], Liza Dijkhuis[1,12], Susanne van den Brink[6,7], Michele

Cilli[8], Connor Byrne[1,13], Giulia-Andreea Ionescu[1,14], Joana Cerveira[9], Kamil R.

Kranc[10,11], Victor Hernandez-Hernandez[1,2], Mirco Ponzoni[5], Anna Bigas[6,7], Jordi

Garcia-Ojalvo[4], Alfonso Martinez Arias[4], Cristina Pina[1,2¶]

[1]College of Health, Medicine and Life Sciences, [2]Centre for Genome Engineering and Maintenance,

Brunel University London, UK; [3]Department of Genetics, University of Cambridge, UK; [4]Department of

Medicine and Life Sciences, Universitat Pompeu Fabra, Spain; [5]Laboratory of Experimental Therapies

in Oncology, IRCCS Istituto Giannina Gaslini, Genova, Italy; [6]Program in Cancer Research, Hospital

de Mar Research Institute, Barcelona, Spain; [7]Josep Carreras Leukemia Research Institute, Barcelona,

Spain; [8]Animal Facility, IRCCS Policlinico San Martino, Genova, Italy; [9]Department of Pathology,

University of Cambridge, UK; [10]Centre for Haemato-Oncology, Barts Cancer Institute, Queen Mary

University of London, UK; [11]Centre for In Vivo Modelling, Institute of Cancer Research, Sutton, UK.

[12]Current address: Sanquin Research Laboratory, Department of Hematopoiesis, Amsterdam, The

Netherlands

[13]Current address: Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

[14]Current address: Division of Infection and Immunity, Cardiff University School of Medicine, Cardiff,

Henry Wellcome building, CF14 4XN

*These authors contributed equally to this work

¶Lead contact. Correspondence: cristina.pina@brunel.ac.uk. Brunel University London, Kingston Lane,

Uxbridge UB8 3PH, United Kingdom.

ABSTRACT

Current in vitro models of developmental blood formation lack spatio-temporal accuracy and weakly replicate successive waves of hematopoiesis. Herein, we describe a mouse embryonic stem cell (SC)-derived 3D hemogenic gastruloid (hGx) that captures multi-wave blood formation, progenitor specification from hemogenic endothelium (HE), and generates hematopoietic SC precursors capable of short-term engraftment of immunodeficient mice upon maturation in an adrenal niche. We took advantage of the hGx model to interrogate the origins of infant acute myeloid leukemia (infAML). We focused on MNX1-driven leukemia, representing the commonest genetic abnormality unique to the infant group. Enforced MNX1 expression in hGx promotes the expansion and in vitro transformation of yolk sac-like erythroid-myeloid progenitors (EMP) at the HE-to-hematopoietic transition to faithfully recapitulate patient transcriptional signatures. By combining phenotypic, functional and transcriptional profiling, including at the single-cell level, we establish the hGx as a useful new model for the study of normal and leukemic embryonic hematopoiesis.

INTRODUCTION

Blood formation in the embryo is a multi-stage process that develops across multiple

cellular niches, reflecting a complex hierarchy of tissue interactions. It involves

sequential production of distinct cell types in so-called hematopoietic waves,

separated in time and space (Lacaud and Kouskoff, 2017; Costa, Kouskoff and

Lacaud, 2012; Medvinsky, Rybtsov and Taoudi, 2011; Dzierzak and Bigas, 2018). The

first wave of hematopoiesis produces unipotent red blood cell and macrophage

precursors in the yolk sac (YS) from an angioblast precursor which can also form

endothelium; it occurs at mouse embryo day (E)7-E7.5 (Fujimoto *et al.,* 2001; McGrath

and Palis, 2005; Moore and Metcalf, 1970). The subsequent wave, known as

definitive, produces: first, oligopotent erythro-myeloid progenitors (EMPs) in the YS

(E8-E8.5); and later myelo-lymphoid progenitors (MLPs – E9.5-E10), multipotent

progenitors (MPPs – E10-E11.5), and hematopoietic stem cells (HSCs – E10.5-

E11.5), in the aorta-gonad-mesonephros (AGM) region of the embryo proper (Ivanovs

et al., 2011; McGrath *et al.,* 2015; Palis *et al.,* 1999; Medvinsky and Dzierzak, 1996;

Medvinsky, Rybtsov and Taoudi, 2011). Definitive blood waves are characteristically

specified from a specialised hemogenic endothelium (HE) through a process of

endothelial-to-hematopoietic transition (EHT) (Zovein *et al.,* 2008; Li, W. *et al.,* 2005).

HSC specification from HE is the last event in hematopoietic development and occurs

in characteristic cell clusters found on the ventral wall of the dorsal aorta (Ivanovs *et*

al., 2011; Boisset *et al.,* 2010). All blood cell types enter circulation from their original

locations and migrate to the fetal liver (FL) (Ciriza *et al.,* 2013; Ghiaur *et al.,* 2008).

Most embryonic blood progenitors are transient and sustain blood production

exclusively during embryonic development, through differentiation. In contrast, HSCs

make little or no contribution to embryonic and fetal blood: instead, they proliferate (i.e.

self-renew) to expand their numbers. At least a subset of HSC migrates to the bone

marrow (BM) niche at the end of gestation and balances self-renewal and

differentiation to maintain blood production throughout the entire post-natal life (Bowie

et al., 2006; Kikuchi and Kondo, 2006; Ganuza *et al.,* 2022; Yokomizo *et al.,* 2022).

Blood progenitors and HSCs can be targeted by genetic alterations that confer clonal

advantage and/or drive initiation and development of leukemia. Leukemia is the most

common malignancy and the most common cause of cancer-related death in the

paediatric age group (Steliarova-Foucher *et al.,* 2017; Dong *et al.,* 2020). Several

paediatric leukemias originate *in utero*, and a subset of these are driven by mutations

exclusive to the paediatric group (Masetti *et al.,* 2015; Cazzola *et al.,* 2021). Such

pediatric-exclusive mutations target blood cell types restricted to development or

depend on signals that are specific to embryonic hematopoietic niches, effectively

defining distinct biological entities to adult malignancies (Cazzola *et al.,* 2021; Bolouri

et al., 2018; Chaudhury *et al.,* 2018). Dissection of these forms of developmental

leukemia is confounded by the inability to robustly identify and isolate their embryonic

cells of origin, and/or to fully recapitulate the disease in post-natal cells, which leads

to failure in generating relevant targeted therapies (Alexander and Mullighan, 2021).

One such example is t(7;12)(q36;p13) acute myeloid leukemia (AML) driven by

rearrangement and overexpression of the *MNX1* gene locus: *MNX1*-rearranged

(MNX1-*r*) AML is restricted to infants under the age of 2 and carries a high risk of

relapse and poor survival (Ragusa *et al.,* 2023). *MNX1* is a developmental gene

required for specification of motor neurons and pancreatic acinar cells (Thaler *et al.,*

1999; Harrison *et al.,* 1999). MNX1 is not known to participate in embryonic

haematopoiesis, however it has been discordantly described to be expressed in some

subsets of adult haematopoietic cells (Deguchi and Kehrl, 1991; Nagel *et al.,* 2005;

Wildenhain *et al.,* 2012; Taketani *et al.,* 2008). It is ectopically expressed in AML cells

carrying one of a variable set of t(7;12) translocations which place the normally silent

MNX1 gene under the control of *ETV6* regulatory elements (Weichenhan *et al.,* 2023;

Bousquets-Muñoz *et al.,* 2024; Ragusa *et al.,* 2023; Ballabio *et al.,* 2009). *ETV6* is

required for hematopoietic cell specification from HE via VEGF signalling (Ciau-Uitz *et*

al., 2010), potentially positioning *MNX1* within a critical embryonic hemogenic

regulatory network.

Faithful *in vitro* recapitulation of developmental blood cell types and their embryonic

niches has the potential to dissect the cellular and molecular mechanisms presiding

at initiation of developmental leukemia, reveal prognostic biomarkers, and indicate

therapeutic vulnerabilities which are otherwise challenging to test in the embryo.

Despite widespread use of embryonic stem (ES) cell and induced pluripotent stem

(iPS) cell-based models of blood specification, a system which recapitulates the spatial

and temporal complexity of embryonic blood formation within the respective time-

dependent niches is still lacking (Dijkhuis *et al.,* 2023). In the last few years, gastruloid
models have emerged as robust avatars of early development, self-organising
processes of symmetry breaking, elongation, multi-axis formation, somitogenesis and
early organogenesis, with remarkable similarity to embryo processes in space and
time (van den Brink, Susanne C and van Oudenaarden, 2021; Beccari *et al.,* 2018;
van den Brink, Susanne C *et al.,* 2020). Gastruloids are grown individually in a multi-
well format that facilitates tracking and screening of developmental processes.
Relevant to blood formation, a variant gastruloid culture system, which successful
organises a heart primordium (Rossi *et al.,* 2021), was also able to recapitulate YS
blood formation with erythroid and myeloid cellularity (Rossi *et al.,* 2021; Rossi *et al.,*
2022).

Herein, we describe a new protocol of gastruloid formation, i.e. hemogenic gastruloids
(hGx), that captures YS and AGM-like waves of definitive blood progenitor
specification from HE. The AGM-like wave includes cells capable of short-term
hematopoietic engraftment of the spleen and BM of immunodeficient mice, overall
suggesting nearly full EMP-to-MPP/HSC recapitulation of embryonic blood
development. We harnessed the embryonic haematopoietic context provided by hGx
for modelling MNX1-*r* infant leukemia by proxy of *MNX1* overexpression. MNX1-
overexpressing hGx showed expansion of cells at the HE-to-EMP transition, which
became susceptible to transformation *in vitro* and recapitulated MNX1-*r* AML patient
signatures, placing *MNX1*'s leukemogenic effects at a very early stage of
developmental blood formation.

RESULTS

HGxs capture cellularity and topography of developmental blood formation

With the aim of recapitulating embryonic haematopoiesis with spatial and temporal
accuracy, we modified existing gastruloid formation protocols to promote the
specification of hemogenic mesoderm derivatives and extend developmental time
beyond the E8.0-equivalent of early developmental gastruloids (Van den Brink,
Susanne C *et al.,* 2014) and the E9.0 timepoint of cardiogenic gastruloids (Rossi *et*
al., 2022). We established a 216h hGx protocol (**Fig. 1A**) using a *Flk1-GFP* mouse ES

cell line (Jakobsson *et al.,* 2010), which allows the tracking of early hemato-endothelial

specification by expression of the *Flk1* (i.e. *Kdr*) *locus*. Activation of the *Flk1*

expression, visible by fluorescence of the GFP tag, necessitated induction of the TGF-

β signalling pathway by Activin A, in addition to the characteristic gastruloid patterning

via WNT activation by CHI99021 supplementation at 48h (**Fig. S1A**). We sought to

promote hemato-endothelial programmes through addition of VEGF and FGF2 at 72h

(Sroczynska *et al.,* 2009) (**Fig. 1A**), and observed extension of a polarised *Flk1-GFP*-

expressing cellular network (**Fig. 1B**), which included VE-cadherin$^+$ C-Kit$^+$ cells

suggestive of EHT (**Fig. 1C-D**). This process of HE specification and EHT peaked at

120-144h; it was followed by a surge in CD41$^+$ candidate hematopoietic cells at 144h

(**Fig. 1D**). In our experience, observation of early polarised activation of the *Flk1 locus*

at 96h (**Fig. 1B**) was predictive of CD41$^+$ detection 2 days later, suggesting that TGF-

β activation in early patterning is required for later hematopoietic development,

putatively by modifying mesoderm specification. We extended the protocol by

incorporation of a 24-hour pulse of sonic hedgehog (Shh) between 144-168h to mimic

the aortic patterning that precedes cluster formation (**Fig. 1A**) (Rybtsov *et al.,* 2014).

Continued exposure to VEGF after the Shh pulse resulted in specification of CD45$^+$

hematopoietic cells progenitors (**Fig. 1D-E**), which increased after 168h. The

proportion of CD45$^+$ cells was enhanced by the addition of SCF, FLT3L and TPO (**Fig.**

S1B-C), a combination of cytokines normally used for HSC maintenance; conversely,

it did not require continued addition of FGF2 after 168h (**Fig. S1D**), which was omitted.

CD45$^+$ cells appeared in a small number of 'cluster'-like structures adjacent to CD31$^+$

endothelium (**Fig. 1E**), which were reminiscent of HSC and progenitor cell budding

from the aortic HE, and suggested recapitulation of hemogenic tissue architecture.

Time-dependent specification of C-Kit$^+$, CD41$^+$ and CD45$^+$ cells in hGxs could be

obtained with other mouse ES cell lines, including commonly used E14Tg2a (E14)

cells (**Fig. S1E-G**), with successful specification of comparable frequencies of CD45$^+$

progenitors (**Fig. S1H**), attesting to the reproducibility of the protocol.

Overall, our extended hGx model could capture successive specification of HE, CD41$^+$

and CD45$^+$ cells, with temporal and spatial congruence relative to the embryo, thus

showing initial promise as a tool for the exploration of normal and perturbed

developmental hematopoiesis.

Figure 1

HGx capture time-dependent emergence of YS-like and AGM-like progenitors from hemogenic endothelium and contain elements of hemogenic niches

To better characterize the extent and progression of developmental hematopoiesis in hGx, we performed single-cell RNA-sequencing (scRNA-seq) time-course analysis of gastruloid cell specification. We sorted cells from 2 independent hGx cultures at 120, 144, 168, 192 and 216h, and profiled a total of 846 cells using the Smart-Seq2 protocol (Picelli *et al.,* 2014) (**Fig. S2A**). We analysed unfractionated live single hGx cells at 120-192h and enriched the dataset with sorted CD41$^+$ cells at their peak time of 144h, and sorted CD45$^+$ cells at 192 and 216h. At timepoints of significant enrichment in CD41$^+$ (144h) and CD45$^+$ cells (192h) (**Fig. 1D**), we also sorted C-Kit$^+$ cells. Library preparation and sequencing generated an average of 120000 reads/cell, which were mapped to an average of 4000 genes/cell, with minimal signs of cell stress / death as measured by mitochondrial DNA fraction (**Fig. S2B**). Read and gene counts were similar between biological replicates, cell types and at different time points (**Fig. S2C**). The exception was 120h unfractionated hGx cells, which despite similar sequencing depth (average read counts), were mapped to twice the number of genes (**Fig. S2C**), possibly reflecting multi-gene program priming at the onset of hemogenic specification. We selected highly varying genes (HVG) before principal component analysis (PCA) dimensionality reduction and retained the most relevant dimensions. In the PCA reduced space, we constructed a KNN graph and used uniform manifold approximation and projection (UMAP) to visualize the data on 2 dimensions, looking for cell communities using Leiden clustering (**Fig. 2A**). We identified 12 cell clusters of which 2 (clusters 0 and 5) almost exactly mapped to C-Kit$^+$ cells (also Sca-1$^+$), 2 (clusters 1 and 8) contained CD45$^+$ sorted cells, and cluster 4 uniquely captured the CD41$^+$ cells observed at 144h (**Fig. 2A**). Although some unfractionated cells overlapped with the hemogenic cell clusters (**Fig. 2A** - center), most occupied different transcriptional spaces, reflecting the frequency of hematopoietic cells and suggesting the presence of potential hemogenic niches.

To explore tissue and lineage affiliation of the different clusters, we performed differential gene expression against all other cells using Wilcoxon ranking test, and established cluster classifier gene lists (**Supplemental File S1**). Top classifier genes

for clusters 0 and 5 had endothelial affiliation (**Supplemental File S1**), and the cellular

composition of the clusters broadly reflected their time signature (**Fig. 2A**): cluster 5

comprised 144h cells and cluster 0 included most C-Kit$^+$ cells at 192h. Cluster 5

occupied the vicinity of 144h-CD41$^+$ cells (**Fig. 2A**), which carried an erythroid-biased

signature (**Supplemental File S1**). On the other hand, 192/216h-CD45$^+$ cells in cluster

8 expressed myeloid and lymphoid genes (**Supplemental File S1**). The remaining

clusters largely corresponded to time-restricted unfractionated cells (**Fig. 2A** - right),

with clusters of cells present at later time-points matching mesenchymal stromal cell

and autonomic neuron signatures (**Fig. S2D**), potentially configuring incipient

formation of niches relevant to production of HSC (Kapeni *et al.,* 2022; Fitch *et al.,*

2012).

To further characterise the identity of hGx cells, we interrogated lineage priming in

individual hemato-endothelial cells (**Fig. 2B-C**), to probe the presence of putative

multilineage YS (EMP) and AGM (MLP/MPP) progenitors, and HE cells. The

heatmaps (**Fig. 2B-C**) show expression of lineage markers and regulators, with

unsupervised clustering of cells (**Fig. S2E-F**) and genes. The gene list used was

identical in both analyses, with only expressed genes shown. Concordance between

sorting antigens and expression of the respective genes (*Kit, Itga2b* [CD41] and *Ptprc*

[CD45]) was partial potentially reflecting low levels of transcript expression in the hGx

system. Summary of endothelial, co-expressing erythro-myeloid (EM) and co-

expressing myelo-lymphoid (ML) programs (**Fig. 2D**) clearly associates the first with

C-Kit$^+$ and the latter with CD45$^+$ cells; CD41$^+$ cells capture a heterogeneous

population, which includes a proportion of cells co-expressing erythroid and myeloid

genes that constitute candidate EMPs.

Clustering of cells sorted as C-Kit$^+$ did not separate the 144h and 192h time-points

(**Figs. 2B** and **S2E**), indicating that the relative separation on the global UMAP was

not based on endothelial identity genes. We observed widespread expression of

arterial genes (e.g. *Flt1, Efnb2*) with relative reciprocity with venous endothelial genes

(e.g. *Flt4*) supporting specification of both types of endothelia (**Fig. 2B**). There was

limited co-expression of hematopoietic genes, but genes associated with endothelial-

to-hematopoietic transition (EHT), e.g. *Gata2, Notch1,* and *Procr,* were significantly

more frequent within the C-Kit⁺ population (**Fig. 2E**). Specific associations of *Cd44*

and *Procr* with the 192h time-point (**Fig. 2F**) putatively represent aorta-like EHT.

Clustering of CD41⁺ and CD45⁺ cells (**Fig. 2C** and **S2F**) more clearly separates unique

hematopoietic signatures. A subset of cells, predominantly CD41⁺, expresses high

levels of globin genes. While embryonic β-globin chains are highly expressed, there

is a gain in fetal *Hbb-y* and adult *Hbb-bt* and *Hbb-bs* (C57Bl/6 strain-specific β major

and β minor, respectively) in CD45⁺ cells at 216h, indicative of AGM-like progenitor

identity (**Fig. 2C**). 192 and 216h cells with high expression of *Ptprc* have an MLP profile

(**Fig. 2C-D**), co-expressing myeloid and lymphoid genes including immunoglobulin

chains. There is progression of the lymphoid signature between 192 and 216h (**Fig.**

2D and **S2G**). Other CD45⁺ cells have low-frequency expression of HSC-associated

transcription factors, including *Myb, Hlf* and *Hoxa9* (**Fig. 2G** and **S2I-J**), suggesting

incipient specification of AGM-like MPP and pre-HSC, albeit with incomplete

programming.

Altogether, single-cell RNA-seq data suggest that hGx specifies YS-like and AGM-like

progenitors in a time-dependent manner, with concomitant emergence of

hematopoietic-supporting niche cells. The data suggested putative incipient

specification of rare candidate MPP or pre-HSC, which we investigated further in

comparison with mouse AGM datasets and at a functional level.

252   Figure 2

253
Late-stage hGx cells transcriptionally resemble mouse AGM progenitors

We made use of a dataset recently published by one of us (Thambyrajah *et al.,* 2024)

that interrogates the mouse AGM at the point of HSC emergence from the dorsal aorta

HE. In this study, a combination of surface markers and *Gfi1* locus reporting of EHT,

tested functionally through transplantation, identified HE and HSC-enriched clusters

(**Fig. S3A** - clusters 1 and 2, respectively, with cluster 4 a putatively heterogenous

cluster at the transition between the 2 states). We projected cell transcription profiles

from hGx to AGM using a version of scmap algorithm (Kiselev, Yiu and Hemberg,

2018) with 1-cell and 3-cell neighbourhoods for lower and higher levels of confidence

on similarity projections (respectively gray and black circles on **Fig. S3A** - left). The

numbers of cluster-specific hGx cells projected onto AGM clusters is quantified in **Fig.**

S3B. Top differentially expressed genes (DEG) between AGM HE and HSC-projected

hGx cells are represented in Fig. **S3C** and **Supplemental File S2**.

With respect to HE specification, we observed that a subset of hGx C-Kit$^+$ cells present

in endothelial clusters 5 and 0 (**Fig. 2A**) projected onto the AGM HE-identity cluster 1

(**Fig. S3A**), with more frequent similarity amongst hGx 192h cluster 0 (41%) than 144h

cluster 5 (21%). Overall, this supports the notion that specification of endothelial and

HE cells in hGx follows with time-dependent developmental progression into putative

AGM-like HE. Lineage identity genes supported the nature of HE-projected hGx cells

(**Fig. S3D**), which expressed *Gata2* but not *Runx1*, *Myb*, or *Gfi1b*, indicating that EHT

occurs downstream of these cells. We then asked if any of the hGx cells in hemopoietic

clusters bore resemblance to cells from the HSC-enriched AGM cluster 2. We

observed that most (85%) hGx CD45$^+$ cells in cluster 8 (**Fig. 2A**) projected onto the

HSC-enriched cluster 2, which also showed similarity with 76% of CD41$^+$-enriched

hGx cluster 4 cells. The relative positioning of hGx cells cluster 4 and cluster 8

progressively further away from the AGM HE cluster was in line with the time-

dependent specification in the hGx. However, the lineage marker programs (Fig. **S3D**)

expressed by the hGx cells, although definitive in nature (e.g. there is expression of

immunoglobulin chains and adult β-globins), lacked detection of HSC genes like *Hlf*,

Gfi1, or *Hoxa9*. On the other hand, *Mllt3* was present (Calvanese *et al.,* 2019), even

in the absence of an erythroid signature (Pina *et al.,* 2008), suggesting a developing

pre-HSC program compatible with the co-expression of multi-lineage markers (**Fig.**
S3D).

Late-stage hGx cells can engraft hematopoietic tissues upon maturation

We inspected the cellularity of hGx at the 216h time-point more closely through a
combination of multi-parameter flow cytometry (**Fig. 3A-B**), colony-forming progenitor
assays (CFC) (**Fig. 3C-D**) and *in vivo* engraftment of haematopoietic tissues (**Fig. 3E-**
G). In order to facilitate the analysis of engrafted animals, we engineered constitutive
expression of BFP from the *Rosa26* locus in the *Flk1-GFP* mES cell line (**Fig. S4A**),
which we also used in some of the *in vitro* analyses. Phenotypic characterisation of
216h-hGx (4 replicates, 1 with *Rosa26-BFP::Flk1-GFP* line) (**Fig. 3A**, **S4B**) showed
the presence of pre-HSC-like CD45$^+$CD41^{lo}c-Kit$^+$VE-Cadherin$^+$ cells (1.8% of all
CD45$^+$CD41^{lo} cells) (**Fig. 3B**), corresponding to an estimated 8 cells per hGx. Analysis
of the hGx progenitor content using multipotential CFC assays (**Fig. 3C-D**) revealed
the presence of >50 CFC/hGx, of which an average of 7 have mixed-lineage
granulocytic-monocytic-erythroid (GEM) potential, numerically matching the estimated
number of pre-HSC. Other progenitors were predominantly granulo-monocytic (GM)
and erythroid (E) progenitors (**Fig. 3C-D**). Differentiated cells of G, M and E lineages
(**Fig. S4C-D**) were also detectable in 216h hGx, putatively reflecting the progeny of
EMP-like cells specified earlier at the 144h timepoint.

As an ultimate test of the presence of pre-HSC, we tested whether late-stage 216h
hGx contained cells capable of hematopoietic engraftment of immunodeficient mice.
In early experiments, we transplanted dissociated hGx cells into irradiated
immunocompetent C57Bl/6 recipients, sub-lethally irradiated immunodeficient NSG
mice, or non-conditioned NSGW41 hosts, but failed to detect signs of engraftment of
peripheral blood, spleen (Spl) or bone marrow (BM) at 12 days (including absence of
spleen colony formation), at 4-6 weeks (short-term engraftment), or beyond 8-10
weeks (long-term engraftment) by either flow cytometry detection of CD45.2 or by
genomic DNA (gDNA) PCR analysis of the *Flk1-GFP* locus. However, given the
absence of clear HSC-like transcriptional programs in 216h CD45$^+$ cells, as analysed
by scRNA-seq (**Fig. 2** and **S2**), and in light of the late specification of mesenchymal or

sympathetic neuronal cells (**Fig. 2A** and **S2D**) putatively supportive of HSC formation,

we hypothesized that hGx pre-HSC required additional maturation in a relevant

supportive niche. We thus tested implantation of undissociated hGx in the adrenal

gland of *Nude* mice (**Fig. 3E**), a topography commonly used to support tumor

development, and recently shown to support adult haematopoietic development

(Schyrr *et al.,* 2023). Given the coincidence of CD45 isoform between *Flk1-GFP*

mouse ES cells and *Nude* mice, we made use of the *Rosa26-BFP::Flk1-GFP* mES

cell line (**Fig. S3A**). We implanted 3 hGx each unilaterally in the adrenal gland of

unconditioned recipient mice. Control mice were injected with an equivalent volume of

PBS. Animals were sacrificed at 24 hours, 12 days, and 4 weeks after implantation,

with 5 experimental animals and 1 control collected at each timepoint. We checked

Spl and BM BFP engraftment at the different timepoints and analyzed the morphology

of adrenal glands receiving implants (**Fig. 3F**). We did not observe BFP engraftment,

spleen enlargement or colony development, or macroscopic changes in the adrenal

gland at the earlier timepoints. However, 3 of 5 adrenal glands were enlarged, and

their histology suggested *in situ* development of the hGx (**Fig. 3F**), particularly

mesodermal tissues – somite derivatives (**Fig. 3F** - left, red box), and lateral plate and

intermediate mesoderm derivatives, respectively blood cells and renal tubules (**Fig.**

3F - left, green box). The control, PBS-injected glands showed normal tissue

architecture (**Fig. 3F** - right). It should be noted that none of the specimens contained

glomeruli in addition to the renal tubules, pointing to an origin from the hGx, and not

from the adjacent kidney tissue. BFP engraftment of the Spl and BM by flow cytometry

was very low level albeit consistently above control (**Fig. S4E**), and we opted for

analysing engraftment by PCR analysis of gDNA using primers directed against the

BFP insert (**Fig. S4A**). We measured the sensitivity of the amplification through serial

dilution of *Flk1-GFP Rosa26-BFP* mES cell gDNA into that of human K562 cells, and

determined detection of 10pg/100ng gDNA, equivalent to 2-3 cells (**Fig. S4F**). PCR

analysis of spleen and BM gDNA detected *BFP* amplicon in both tissues in 3 of the 5

samples (#11, 12 and 15) (**Fig. 3G**). We sampled the gDNA of each recipient 4-5 times

and detected the presence of the amplicon in only a fraction of the replicate samples

of the positive animals, reflecting the low-level detection by flow cytometry. Replicate

sampling of the control was consistently negative (**Fig. 3G**). Sorting of erythroid

Ter119$^+$ and lympho-myeloid CD45$^+$ cells (**Fig. S4G**) showed PCR engraftment in both

lineage fractions (**Fig. S4H**).

Altogether, the data suggest that 216h hGx generate AGM-like pre-HSC capable of at

least short-term multilineage engraftment upon maturation. The low-level production

of engrafting cells recapitulates their rarity *in vivo*, in agreement with the embryo-like

qualities of the gastruloid system, which accommodates modular qualitative bias, but

not quantitative changes in tissue specification.

356   **Figure 3**

HGx resolve the early hemogenic ontogeny targeted by infant AML gene *MNX1*

The ability of hGx to reconstitute YS and AGM-like HE and hemopoietic progenitors with temporal coherence lends itself as an attractive model to understand developmental leukemia. In particular, forms of AML unique to the first 2 years of life – infant (inf)AML – have been shown to transform fetal liver (FL) but not adult haematopoietic cells (Waraky *et al.,* 2024; Mercher *et al.,* 2009; Chen, W. *et al.,* 2011; Ragusa *et al.,* 2023), compatible with targeting of a transient embryonic cell. In the case of t(7;12) AML (herein MNX1-*r*) (Ragusa *et al.,* 2023), which results in a transformation-driving ectopic expression of *MNX1* (Waraky *et al.,* 2024), analysis of patient transcriptional signatures (Ragusa *et al.,* 2022) for over-representation of cell atlas-defined affiliations, indicates enrichment in cardiomyogenic, endothelial, HSC/progenitor and mast cells (**Fig. 4A**). These cell enrichments are compatible with an early hemogenic cell derivative, capturable in the hGx model. Other cell type-enrichments reflect MNX1 functions in pancreatic and neuronal development (Ragusa *et al.,* 2022) (**Fig. S5A**).

We overexpressed *MNX1* (MNX1-OE) in *Flk1-GFP* mouse ES cells by lentiviral transduction (**Fig. 4B**) and analyzed progression of hGx generation (**Fig. 4C**). We used human *MNX1* cDNA to distinguish from the endogenous gene, but the degree of homology is nevertheless high (84%), supporting functional equivalence. We confirmed that *MNX1* overexpression in hGx is maintained at endpoint (**Fig S5B**). MNX1-OE hGx activated polarized *Flk1-GFP* expression and elongated with similar kinetics to empty-vector (EV) control (**Fig. 4C**), but consistently produced larger hGx (**Fig. 4D**) denoting increased cellularity (**Fig. S5C**). From 192h onwards, MNX1-OE hGx had a higher frequency of spontaneously contractile structures (**Fig. S5D**), compatible with the cardiogenic cell association of patients' transcriptomes (**Fig. 4A**). We interrogated the time-dependent hemogenic cell composition of hGx by flow cytometry, quantifying C-Kit, CD41 and CD45 (**Fig. 4E-G** and **S5E-F**). We observed a significant expansion of the C-Kit$^+$ compartment specifically at 144h (**Fig. 4E** and **S5E**). All cell markers were relatively unchanged at later timepoints, although absolute cell numbers were higher in MNX1-OE hGx (**Fig. S5C**).

To better understand the consequences of MNX1-OE on hemogenic development, we performed RNA-sequencing (RNA-seq) of hGx at 144h and 216h, matching YS and AGM-like hemopoietic waves, respectively (**Fig. 4H**). We confirmed expression of the human *MNX1* transgene in the sequenced reads (**Fig. S6A**); in contrast, we could not detect endogenous mouse *Mnx1* in any of the samples (**Fig. S6A**), confirming that it does not normally play a role in hemogenic development. DEGs between MNX1-OE and EV (**Supplemental File S3**) configured distinct MNX1-driven programs at 144h and 216h, with >80% genes timepoint-specific (**Fig. 4H**). Both programs include up-regulation of hematopoietic-associated genes, capturing EHT and progenitor regulators (e.g. *Runx1*, and *Epor* and *Zfpm1*, respectively) at the earlier timepoint, and more differentiated effectors, e.g. *Hba-a1*, *Hbb-bh1* and *Il3ra* at 216h, compatible with selective targeting and/or expansion of newly specified progenitors at the YS-like stage (**Fig. 4H**). An overview of transcriptional regulatory programs at both timepoints denotes a clearer hemogenic enrichment at 144h, with over-representation of GATA2, GATA1 and RUNX1 binding targets (**Fig. 4I**), putatively capturing an expansion of the HE-to-EMP transition, also represented in enriched GO categories related to VEGF signalling and epithelial-to-mesenchymal transition (EMT) (**Fig. 4J**). Compatible with the increase in contractile foci in MNX1-OE late hGx, there was an enrichment in cardiogenic gene ontologies (GO) at the 216h timepoint (**Fig. 4J**). No hemogenic transcription factors (TF)-ChIP or GO categories were enriched in MNX1-OE hGx at 216h (**Fig. 4J-I**).

We attempted to deconvolute the cellular heterogeneity underlying bulk RNA-seq DEGs by interrogating the cluster signatures obtained from the single-cell analysis of hGx development (**Fig. S6B-C**; refer to **Fig. 2A**). In support of enhanced YS-stage hemato-endothelial specification, MNX1-OE transcriptomes showed enrichment of hGx HE cluster 5 and 0 signatures, and of the EMP-like cluster 4 (**Fig. S6B**). MNX1-OE DEGs at both timepoints also enriched for the relatively less differentiated cluster 10, which predominantly captures unfractionated hGx cells at 120h (**Fig. S6B-C**; refer to **Fig. 2A**). Significantly, clusters 5 and 0 HE signatures were enriched in *MNX1*-r infAML patient signatures (Ragusa et al., 2022) (**Fig. S5B-C**), positioning MNX1-OE at early stages of hemogenic specification. MNX1-OE DEGs at 216h show enrichment in late-stage clusters 7 and 8 (**Fig. S6B-C**), denoting enhancement of hGx

development, including MLP/MPP in cluster 8, but diverging from *MNX1*-r infAML

signatures. Projection of infAML subtype-specific gene signatures onto hGx

development separates the lineage affiliations of *MNX1*-r from, for example, the most

frequent infAML driver – *KMT2A (MLL)* rearrangements (**Fig. S6B-C**), which carries

myelo-monocytic and/or mixed-lineage affiliation, highlighting the utility of the hGx

system for interrogating infAML biology.

Figure 4

Serial replating of MNX1-OE hGx cells identifies a candidate C-Kit⁺ leukemia-propagating cell more susceptible to transformation in the YS period

We further interrogated the *MNX1*-r-like leukemia-initiating potential of MNX1-OE hGx through serial replating of CFC assays, a classical *in vitro* assay of leukemia transformation. We compared 144h and 216h hGx to match functional and transcriptome data, and plated unfractionated hGx, given the transient nature of C-Kit⁺ enrichment. Both 144h (**Fig. 5A-B**) and 216h (**Fig. 5C-D**) MNX1-OE hGx replated for at least 5 plates, with significant differences to control (EV) (**Fig. 5B,D**). EV-hGx replating could not be sustained beyond plate 3 in most experiments started from 216h (**Fig. 5C**); 144h EV-hGx exhibited some low-level replating until plate 5 (**Fig. 5A**), but the colonies were formed by few scattered cells clearly distinct from the well-demarcated colonies obtained from MNX1-OE hGx. We analyzed early-stage (plate 1) and late-stage (plate 5) colonies by flow cytometry to identify the nature of the replating cells. Cells selected through replating were C-Kit⁺ (**Fig. 5E**), matching the cell phenotype transiently enriched at 144h. Early replating of MNX1-OE hGx also enriched for erythroid-affiliated Ter119⁺ cells particularly from the 144h time-point (**Fig. 5F**), reflecting the differential erythroid potential of EMP and MLP/MPP at 144h and 216h, respectively. This enrichment is not sustained upon replating, compatible with the undifferentiated nature of *MNX1*-r infAML, which is also reflected by the low levels of CD11b, a marker affiliated with the myelo-monocytic lineages (**Fig. 5E**). Notably, the cells obtained are not CD45⁺ (**Fig. 5F**), suggesting perpetuation of an early hemogenic state.

We performed RNA-seq analysis of MNX1-OE hGx CFC cells obtained through re-

plating and integrated the data with the 144h and 216h hGx time-points to identify

genes enriched through the process of transformation. We observed relative

enrichment in Hs*MNX1* reads in MNX1-OE CFC samples, supporting selective

expansion of transduced cells (**Fig. 6A**). Hierarchical clustering of all DEGs in pairwise

comparisons (**Fig. 6B**), identified a subgroup of genes up-regulated upon CFC

replating which was also up-regulated at MNX1-OE hGx at 144h (K14), thus matching

the putative time-dependent enrichment of the MNX1-OE propagating population. To

understand the relevance of these genes in *MNX1*-r leukemia, we performed Gene

Set Enrichment Analysis (GSEA) of *MNX1*-r patients in comparison to other AML in

the same infant (0-2) age group. K14-genes were enriched in MNX1-*r* all comparisons

(**Fig. 6C**), with considerable overlap between leading-edge (i.e. significantly-enriched)

genes (**Fig. S6D**). Interrogation of cell-type-specific signatures within the K14 MNX1-

r leading edge genes (**Fig. 6D**) against the single-cell PanglaoDB atlas captured

similarities to hemogenic and non-hemogenic cell types, including interneuron, and

other neural, signatures, which reflect MNX1 biological functions (Harrison *et al.,* 1999;

Thaler *et al.,* 1999; Ragusa *et al.,* 2022). The most highly significant hemogenic

signatures capture early developmental components, such as 'endothelial cells', which

were specifically up-regulated within MNX1-OE hGx (**Fig. S6B**). 'Erythroid cell

precursors' also reflect MNX1-OE hGx effects (**Fig. 4E-J**). We were intrigued by the

'mast cells' affiliation of *MNX1*-r AML signatures within the hGx K14 cluster, which was

more specifically aligned with MNX1-OE hGx isolated at 144h (**Fig. S6E**). Given the

characteristic morphology of these cells, we analysed Giemsa-Wright-stained

cytospins of CFC replating assays (**Fig. 6E** and **S6F**). Consecutive plates of 144h

MNX1-OE-initiated *in vitro* transformation showed a shift from blast-like at plate 3 to

precursor cell morphology with some differentiated mast cells at plate 5 (**Fig. S6F**).

However, the extent of differentiation was greater, and apparent earlier, in 216h-

MNX1-OE initiated colonies (**Fig. 6E**). This suggests that the MNX1-OE-responsive

hemogenic C-Kit$^+$ progenitor may experience maturation between 144h and 216h, and

is more likely targeted for transformation at the YS EMP stage. The data support the

utility of the hGx in dissecting ontogeny of embryonic origin in hematological cancers,

allowing us to specifically narrow down the initiating cellular targets of MNX1-*r* AML.

490   **Figure 6**

DISCUSSION

In this study, we developed and characterized a hGx model from mouse ES cells, which successfully recapitulates YS-like and AGM-like waves of definitive blood progenitor specification, and can engraft adult hematopoietic tissues upon *in vivo* maturation. We took advantage of the hGx's nearly full recapitulation of embryonic hemogenic specification, including critical early stages of HE production and EHT to EMP and MLP/MPPs, to interrogate the biology of *MNX1*-r AML, a poorly characterized form of leukemia exclusive to the infant age group (Ragusa *et al.,* 2023). We showed that MNX1-OE transiently expanded a C-Kit$^+$ cell with hemato-endothelial characteristics at the YS-like stage, and could selectively sustain the phenotype through serial replating of CFC assays. Replating CFCs could be initiated beyond the YS-like stage, albeit at the relative expense of an early differentiation state and of the similarity to patient transcriptome, putatively defining a window of susceptibility to MNX1-driven transformation. The hGx MNX1-OE propagating cell bears transcriptional resemblance to patient *MNX1*-r AML, overall positioning the hGx as an attractive tractable model in which to study developmental leukemias.

MNX1-r has been difficult to characterize due to a combination of rarity, diagnostic difficulty (the underlying t(7;12) can be missed if not specifically looked for) (Tosi *et al.,* 2015), and the lack of a fusion protein that can be ectopically expressed and tracked through development, as is the case for other translocations unique to infAML (Alexander and Mullighan, 2021). However, a recent paper using MNX1-OE has shown that MNX1 can initiate a transplantable leukemia in immunodeficient mice by targeting FL, but not BM, cells (Waraky *et al.,* 2024). This suggests that ectopic expression of *MNX1* transforms an embryonic cell, rather than depends on an embryonic niche. It also suggests that MNX1-OE can cause leukemia as a single genetic hit, although it cannot be excluded that other mutations are required for, and may indeed have contributed to, full transformation *in vivo*, as observed in a recent report confirming the *in utero* origin of the leukemia (Bousquets-Muñoz *et al.,* 2024). However, the mouse model does not address the nature of the embryonic cell targeted by MNX1-OE. Like the MNX1-OE hGx cells selected through serial replating, MNX1-OE mouse leukemia are also C-Kit$^+$ (Waraky *et al.,* 2024). Interestingly, clusters of genes up-regulated in the mouse MNX1-OE leukemia (**Fig. S7A** – K1, 2, 5, 6, and 9)

do not robustly separate *MNX1*-r from other infAML by GSEA analysis (**Fig. S7B**), and have diverged more from the patient data in their cell-type signature analysis than the original MNX1-OE cells in the FL (**Fig. S7C**), putatively selecting a more generic program of leukemia transformation. We propose that the divergence from *MNX1*-r infAML signatures increases from MNX1-OE hGx 144h (YS-like), to 216h (AGM-like) to FL, positioning the origin of the AML at the YS stage, likely at the HE-to-EMP transition. This cell may persist in the hGx culture system to the 216h AGM-like time-point, and migrate to the FL in the embryo, eventually extinguishing through differentiation, as suggested by the MNX1-OE hGx CFC assays. Persistence of the original MNX1-OE target cell will determine the sensitivity to MNX1-OE, and can be addressed in the hGx system through lineage tracing. Whether additional mutations, namely a reported association between *MNX1*-r and tri(19) (Espersen *et al.,* 2018; Ragusa *et al.,* 2023), contribute to prolonging the undifferentiated state of the candidate YS target cell with increased probability of transformation, or otherwise expand the targeted cell in a more classical second-hit fashion, can potentially also be screened in the hGx system, in combination with transplantation experiments. Finally, although our study has positioned the MNX1-OE target cell within the YS-EMP stage, the exact nature of the cell requires better definition. The mast cell affiliation, transcriptional and by CFC morphology, is more likely to reflect the period of susceptibility to MNX1 and the multilineage potential of the target cell, than the lineage commitment of the latter. Indeed, blast cells in the leukemia mouse model are morphologically undifferentiated (Waraky *et al.,* 2024) and a mast cell morphology has not been described in patients (Ragusa *et al.,* 2023; Espersen *et al.,* 2018). Although adult HSC have mast cell potential *in vitro*, this is not employed *in vivo* under physiological conditions (Chia *et al.,* 2023). Tissue resident mast cells are formed in the embryo at 2 periods: the YS EMP stage, which we propose to have targeted with MNX1-OE in the hGx model; and a transient AGM FL-HSC population which does not persist in post-natal life (Chia *et al.,* 2023; Yoshimoto *et al.,* 2022), and that may be incipiently specified at the end of the current hGx protocol. Prospective isolation of these cell types from embryos and over-expression of *MNX1* followed by transformation *in vitro* and *in vivo*, will precisely define the embryonic time-window of susceptibility to *MNX1*-r and the nature the leukemia-initiating cell. Harnessing the

high-throughput nature of the hGx culture system will help identifying therapeutic
vulnerabilities that can improve the prognosis of this often-fatal disease.

The hGx model furthermore constitutes an advanced platform to probe physiological
development of the hematopoietic system. Currently, *in vitro* platforms do not
recapitulate *in vivo* blood formation sufficiently faithfully (Dijkhuis *et al.,* 2023; Fidanza
and Forrester, 2021). Establishment of a tractable *in vitro* model of hematopoietic
development which faithfully recapitulates time-dependent emergence of
hematopoietic progenitors in their physiological niche would propel translational
applications by enabling high-throughput studies of HSC emergence. Most ESC-
based models of blood specification do not achieve concomitant recapitulation of the
surrounding microenvironment. Protocols typically remove ESC-derived hemogenic
cells from an early embryonic-like niche to direct differentiation through genetic
manipulation (Sugimura *et al.,* 2017) and/or empirically-determined growth factor
addition (Lis *et al.,* 2017; Vo and Daley, 2015; Fowler *et al.,* 2024; Nafria *et al.,* 2020;
Sroczynska *et al.,* 2009; Piau *et al.,* 2023), with minimal probing of intermediate steps.
When hematopoietic waves have been specifically inspected, emergence of YS and
AGM-like stages were conflated in time (Pearson *et al.,* 2015; Murry and Keller, 2008),
suggesting a bias towards intrinsic hematopoietic cell potential, at the expense of cell
fate modulation by time-coherent extrinsic physiological cues, affecting the nature of
cellular output. The present hGx system, in contrast, successfully deconvolutes YS-
like and AGM-like stages of blood production, including capturing differences in the
HE generation, suggesting a level of organisation amenable to further development
into an integrated hemogenic niche. Although the system is shy of achieving HSC
maturation *in situ*, it can mature in a supportive microenvironment, in this case
provided by the adrenal gland (Schyrr *et al.,* 2023), to produce engrafting multilineage
progenitors, and potentially HSCs. This suggests that access to the relevant
hematopoietic molecular programs is preserved. Incipient specification of non-
hematopoietic tissues shown to support HSC emergence, including PDGFRA[+]
mesenchymal (Chandrakanthan *et al.,* 2022) and neuro-mesenchymal (Miladinovic *et*
al., 2024) cells, and neural-crest derivatives (Kapeni *et al.,* 2022; Fitch *et al.,* 2012),
occurs in sync with development of AGM-like hematopoiesis, to indicate the presence

of cues that with further maturation may facilitate HSPC specification at the
appropriate developmental time.

We have observed that the nature of the symmetry-breaking pulse affects the
specification of hemato-endothelial programs, with a more robust *Flk1*/*Kdr* activation
in the presence of Activin A, potentially balancing mesodermal derivatives and the
anterior-posterior axis. The hematopoietic output of cardiogenic gastruloids (Rossi *et*
al., 2022) does not seem to require Activin A, but their blood formation is of lower
efficiency, at least at the comparable timepoints of 144 and 168h, and apparently
biased towards erythroid formation, reflecting its YS, and potentially primitive, nature.
In the absence of additional cytokines after the CHI99021 pulse, standard gastruloid
protocols (Beccari *et al.,* 2018) capture a similar transcriptional erythrogenic wave.
Although Rossi *et al.* (2022) reported the formation of spleen colonies, these are
difficult to ascertain from the data published, and the engraftment capacity of
cardiogenic gastruloid-derived blood remains uncertain. Our hGx system has more
robust CD41+ cell production at 144-168h, which may reflect distinct tissue induction
by the CHI99021+Activin A pulse. Interestingly, through extending the hGx culture,
and in addition to AGM-like hematopoiesis, we also captured the potential to form renal
tubular structures upon *in vivo* maturation, suggesting induction of lateral plate and
intermediate mesoderm from which the AGM region derives, and putative
reconstitution of the intra-embryonic hematopoietic niche. An additional challenge is
tissue organisation: non-hematopoietic gastruloid models benefitted from mechanical,
or chemo-mechanical, support of artificial ECM embedding (Veenvliet *et al.,* 2020; van
den Brink, Susanne C *et al.,* 2020), to match temporal coherence and tissue
polarisation, with detailed tissue architecture illustrated by somite formation. Prior to
HSC emergence in the AGM, critical signals such as SCF and Shh, which we have
delivered to the cultures, are secreted by different cells on opposite dorsal-ventral
locations (Souilhol *et al.,* 2016), indicating the potential relevance of tissue
organisation to improve and potentially extend hematopoietic formation. Moreover,
HSPC formation in the AGM is facilitated by circulatory shear stress (North *et al.,* 2009;
Diaz *et al.,* 2015; Lundin *et al.,* 2020), suggesting that introduction of flow may improve
hemato-endothelial organization and boost blood production.

Over the last decade, the effort to achieve robust hematopoietic production from PSC,

mouse and human, has been centred on the generation of HSC, a critical translational

need (Chang *et al.,* 2023; Dijkhuis *et al.,* 2023; Fidanza and Forrester, 2021). There

are good indications that this is achievable, both with (Vo and Daley, 2015; Dang *et*

al., 2002) and without (Piau *et al.,* 2023) genetic manipulation. However, success

remains episodic, mainly because mechanistic understanding of processes

underpinning blood generation during development is limiting, exacerbated by limiting

access to human embryonic and fetal material. On-going optimization of gastruloid

models and the success at integration of organoid cultures in microfluidic systems

(Saorin, Caligiuri and Rizzolio, 2023; Quintard *et al.,* 2024) will allow us to further

optimize hGx culture. In conclusion, the hGx represents a faithful *in vitro* reference to

replace and complement embryo studies, holding promise to understand the biology

of healthy and malignant developmental haematopoiesis and advance blood

production for clinical use.

EXPERIMENTAL PROCEDURES

MATERIALS

	SOURCE	IDENTIFIER
Antibodies		
Anti-mouse c-Kit (CD117) clone 2B8 - APC-Cy7	Biolegend	Cat. #105825
Anti-mouse CD41 clone MWReg30 - PE-Dazzle594	Biolegend	Cat. #133935
Anti-mouse CD45 clone 30F11 - APC	Biolegend	Cat. #103111
Anti-mouse CD144 (VE-cadherin) clone BV13 - APC	Biolegend	Cat. #138011
Anti-mouse CD43 clone S11 - PE	Biolegend	Cat. #143205
Anti-mouse Ter119 – PC7	Biolegend	Cat. #116223
CD11b anti-mouse/human clone M1/70 - biotin	Biolegend	Cat. #101203
Mouse Anti-Mouse CD45.2 clone 104 - PE	BD Biosciences	Cat. #560695
Rat Anti-Mouse CD31 clone MEC13.3 - biotin	BD Biosciences	Cat. #553371
Goat Anti-Mouse c-Kit/CD117	R&D Systems	Cat. #AF1356SP
Bacterial and virus strains		
E. coli: NEB 5-alpha Competent *E. coli*	New England Biolabs (NEB)	Cat. #C2987H
Chemicals, peptides, and recombinant proteins		
Murine LIF	Peprotech	Cat. #250-02
StemMACS PD0325901	Miltenyi Biotec	Cat. #130-106-5411
Chiron (CHIR99021)	Biogems	Cat. #2520691
Activin A PLUS	QKine	Cat. #Qk005
Murine VEGF 165	Peprotech	Cat. #450-32
Murine FGF-basic	Peprotech	Cat. #450-33

Murine Sonic Hedgehog (Shh)	Peprotech	Cat. #315-22
Murine TPO	Peprotech	Cat. #315-14
Murine Flt3-Ligand	Peprotech	Cat. #250-31L
Murine SCF	Peprotech	Cat. #250-03
N2B27 medium (NDiff 227)	Takara Bio	Cat. #Y40002
DMEM/F-12	Gibco	Cat. #11320033
Neurobasal medium	Gibco	Cat. #21103049
N-2 Supplement	Gibco	Cat. #17502048
B-27 Supplement	Gibco	Cat. #17504044
Critical commercial assays		
Mouse Methylcellulose Complete Media	R&D Systems	Cat. #HSC007
Deposited data		
Single cell RNA-seq of hemogenic gastruloids at timepoints	This paper	ArrayExpress: E-MTAB-12148
Bulk RNA-seq of hemogenic gastruloids with MNX1 overexpression at 144h and 216 h	This paper	ArrayExpress: E-MTAB-12173
Experimental models: Cell lines		
Mus musculus: Flk-1-GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	Jakobsson et al., 2010
Mus musculus: Sox17-GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	Niakan et al., 2010
Mus musculus: T/Bra::GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	Fehling et al., 2003
Mus musculus: E14Tg2a mouse embryonic stem cell line	MMRRC, University of California Davis, US	Hooper et al., 1987
Mus musculus: Flk-1-GFP::Rosa26-BFP mouse embryonic stem cell line	This paper	n/a
Homo sapiens: HEK293T	ATCC	CRL-3216
Oligonucleotides		
MNX1 (forward 5-GTTCAAGCTCAACAAGTACC-3; reverse 5- GGTTCTGGAACCAAATCTTC-3)	Gulino et al., 2021	n/a
Ppia (forward 5- TTACCCATCAAACCATTCCTTCTG-3; reverse 5- AACCCAAAGAACTTCAGTGAGAGC-3)	Moris et al., 2018	n/a
BFP (forward 5-GCACCGTGGACAACCATCACTT-3; reverse 5-CAGTTTGCTAGGGAGGTCGC-3)	This paper	n/a
Recombinant DNA		
pWPT-LSSmOrange-PQR	Ghevaert Lab, WT-MRC Cambridge Stem Cell Institute, UK	Dalby et al., 2018
pWPT-LSSmOrange-PQR-MNX1	Cloned by Biomatik Corporation (Kitchener, Canada)	n/a
pcdna3-rosa26-rosa26-bfp	This paper	n/a
Software and algorithms		
GSEA software v4.2.3	Subramanian et al., 2005	https://www.gsea-msigdb.org/gsea/index.jsp
ImageJ	Schneider et al., 2012	https://imagej.nih.gov/ij/
Bowtie2	Langmead and Salzberg, 2012	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
Samtools	Li et al., 2009	http://samtools.sourceforge.net/

Tophat2	Kim et al., 2013	https://ccb.jhu.edu/software/tophat/index.shtml
HTseq	Anders et al., 2015	https://htseq.readthedocs.io/en/master/
DEseq2	Love et al., 2014	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
ExpressAnalyst	n/a	https://www.expressanalyst.ca/
PANTHER	Thomas et al., 2022	http://www.pantherdb.org/
EnrichR	Chen et al., 2013; Kuleshov et al., 2016; Xie et al., 2021	https://maayanlab.cloud/Enrichr/
Panglao BP	Franzen et al., 2019	https://panglaodb.se
TARGET (Therapeutically Applicable Research to Generate Effective Treatments) Database	https://ocg.cancer.gov/programs/target https://xenabrowser.net/	GDC TARGET-AML
Single-cell RNA sequencing analysis scripts	DockerHub: dsblab/single_cell_analysis:0.5	https://github.com/dsb-lab/blood_gastruloids
Other		
CELLSTAR cell-repellent cell culture plate, 96 well, U-bottom	Greiner (Bio-One)	Cat. #650970

636
637

METHODS

Cell culture

Flk-1-GFP (Jakobsson *et al.,* 2010), Flk-1-GFP::Rosa26-BFP, Sox17-GFP (Niakan *et al.,* 2010), T/Bra::GFP (Fehling *et al.,* 2003), and E14Tg2A (Hooper *et al.,* 1987) mouse embryonic stem cells (mES) lines were cultured in ES+LIF medium in gelatinized (0.1% gelatin) with daily medium change, as previously described (Turner *et al.,* 2017). The ES+LIF medium contained 500 ml Glasgow MEM BHK-21 (Gibco), 50 ml of fetal bovine serum (FBS, Embryonic Stem Cells tested, biosera, Nuaillé, France), 5 ml GlutaMAX supplement (Gibco), 5 ml MEM Non-Essential Amino Acids (Gibco), 5 ml Sodium pyruvate solution (100 mM, Gibco), and 1 ml 2-Mercaptoethanol (50mM, Gibco). Murine LIF (Peprotech) was added at 1000 U/ml. HEK293T cells were grown in Dulbecco's modified Eagle medium (Gibco) supplemented with 10% FBS

650 (Gibco) and 1% penicillin/streptomycin (Gibco). All cultures were kept at 37°C and 5%

651 CO_2.

Hemogenic gastruloids assembly, culture, and dissociation

mES were maintained in ES+LIF medium and transferred to 2i+LIF (containing Chiron

and MEK inhibitor PD03) for 24 hours prior to the assembly into gastruloids. 250

untransduced mES cells (or 400 for GX-MNX1 and GX-EV) were seeded in each well

of a U-bottom, cell-repellent 96-well plate (Greiner Bio-One, Stonehouse, UK) in 40 µl

of N2B27 medium [Takara Bio or home-made as Cold Spring Harbour Protocols

'N2B27 Medium' (2017)]. The plate was centrifuged at 750 rpm for 2 minutes to

promote deposition and aggregation of the cells and was then incubated at 37°C, 5%

CO2 for 48 hours. After 48 hours, 150 µl of N2B27 medium supplemented with 100

ng/ml Activin A (QKine, Cambridge, UK) and 3 µM chiron (Peprotech) was added to

each well. At 72 hours, 150 µl of medium were removed, without disrupting the

gastruloids in the wells. 100 µl of N2B27 with 5 ng/ml of Vegf and Fgf2 each

(Peprotech) were added to each well. From 72 h to 144 h, each day 100 µl of medium

were removed and replaced with N2B27 + Vegf + Fgf2. At 144 h, the medium was

further supplemented with Shh at 20 ng/ml. From 168 h to 216 h, the medium was

N2B27 + 5 ng/ml Vegf, plus 20 ng/ml mTpo, 100 ng/ml mFlt3l, and 100 ng/ml mScf

(Peprotech). To dissociate cells from the gastruloid structures, medium was removed

and individual gastruloids were collected using a pipette and precipitated at the bottom

of a microcentrifuge tube. The remaining medium was aspirated and the bulk of

gastruloids was washed in PBS. 50 µl of TrypLE express was added to pelleted

gastruloids to be incubated at 37°C for 2 minutes to dissociate cells.

Animals and implantation

Female athymic nude-Foxn1[nu] (*nu/nu*) mice (Envigo, Bresso, Italy) were housed under

pathogen-free conditions. In accordance with the "3Rs policy", experiments were

reviewed and approved by the Animal Welfare Body (OPBA) of IRCCS Ospedale

Policlinico San Martino and by the Italian Ministry of Health (n. 883/2020-PR).

Intact hGx (3 per mouse in 10 µL of PBS) were inoculated in the adrenal gland of five-

week-old mice. Briefly, mice were anesthetized with a mixture of xylazine (ROMPUN,

Bayer) and ketamine (LOBOTOR, Acme S.r.l.) and injected with hGx, after

laparotomy, in the capsule of the left adrenal gland, as previously described (Brignole

et al., 2023; Pastorino *et al.,* 2003). All mice survived surgery. Control mice (CTR)

received vehicle only. Mice body weight and general physical conditions were

recorded daily. One month after hGx inoculation, mice were sacrificed and spleen and

bone marrow (BM) from femurs collected. BM cells and spleens, mechanically reduced

to cell suspension, were stored at -80°C until use. Adrenal glands were paraffin-

embedded for subsequent IHC studies.

Methylcellulose colony forming assays (CFC)

Disassembled gastruloids cells were plated on Mouse Methylcellulose Complete

Media (R&D Systems). Cells were first suspended in Iscove's Modified Dulbecco's

Medium (IMDM, Gibco) with 20% FBS (Gibco) before addition to the methylcellulose

medium. Cells were plated in duplicate 35 mm dishes with 2×10^5 cells/plate. Plates

were incubated at 37°C and 5% CO2 for 10 days, when colonies were scored. For

serial replating experiments, cells in methylcellulose were collected and washed in

phosphate buffer saline (PBS) to achieve single-cell suspensions and replated as

described above. Dissociated gastruloid cells from CFC were collected as cell

suspensions in PBS and centrifuged onto slides (Shandon Cytospin 2 Cytocentrifuge)

at 350rpm for 5-7min, with a PBS-only pre-spin at 1000rpm, 2min. Slides were stained

in Giemsa-Wright stain for 30 seconds followed by addition of phosphate buffer pH 6.5

for 5 min.

Immunofluorescence staining

Immunostaining of whole gastruloids was performed as described before (Baillie-

Johnson et al., 2015). Briefly, gastruloids were fixed in 4% paraformaldehyde (PFA)

dissolved in PBS for 4 hours at 4°C on orbital shaking and permeabilised in PBSFT

(10% FBS and 0.2% Triton X-100), followed by one hour blocking in PBSFT at 4°C on

orbital shaking. Antibody dilutions were made in PBSFT at 1:200 for primary and 1:500

for secondary antibodies. Antibody incubations were performed overnight at 4°C on

orbital shaking, and subjected to optical clearing overnight in ScaleS4 clearing

solution. Individual gastruloids were then mounted on glass coverslips by pipetting as

droplets in ScaleS4 and DAPI nuclear stain.

Histology and H&E staining

Samples were fixed with 10% formalin at room temperature for 48 hours followed by

an ethanol series of ethanol and paraffin; ethanol 70% (90 minutes), ethanol 80% (120

minutes), ethanol 95% (180 minutes), 3 times ethanol 100% of 120, 120 and 180

minutes, 50% ethanol/paraffin (3x120 minutes) and 3 times paraffin (60, 90 and 120

minutes) before embedding in histological cassettes. Blocks were sectioned with a

HistoCore BIOCUT Microtome (Leica) at 10 mm and slides were stained with

Haematoxylin/Eosin.

Imaging

Images of cultured gastruloids and CFC plates were captured using the Cytation 5 Cell

Imaging Multi-Mode Reader (Biotek) plate reader using bright field and FITC channels.

ImageJ (Schneider et al., 2012) was used for gastruloid size quantification. Confocal

microscopy was performed on LSM700 on a Zeiss Axiovert 200 M with Zeiss EC Plan-

Neofluar 10x/0.30 M27 and Zeiss LD Plan-Neofluar 20x/0.4 M27 objective lens. Tissue

sections were imaged with a THUNDER Imager 3D microscope system (Leica).

Lentiviral vector packaging and transduction

The lentiviral overexpression vector pWPT-LSSmOrange-PQR was used to clone the

MNX1 gene cDNA. The viral packaging vectors pCMV and pMD2.G, described in Pina

et al. (2008), were used to assemble lentiviral particles using HEK293T cells via

transfection using TurboFect Reagent (Invitrogen). Transduction of mES cells was

performed overnight by addition of lentivirus to culture medium and washed the

following day (Moris *et al.,* 2018). Transduced cells were sorted for positivity to
LSSmOrange.

Flow cytometry

Surface cell marker analysis was performed by staining using the antibodies listed in
Key Resources Table. Disassembled gastruloids cells were resuspended in PBS, 2%
FBS and 0.5 mM EDTA and stained at a dilution of 1:100 for primary antibodies for 20
minutes at 4°C. When indicated, streptavidin was added at a dilution of 1:200. Analysis
was performed on ACEA Novocyte (Agilent) or AttuneNxT (Thermo) analyzers, using
the respective software packages. Cell sorting was performed using a CS&T calibrated
BDFACS Aria III system (488nm 40 mW, 633nm 20 mW, 405nm 30 mW, and 561nm
50 mW), set with the 100µm nozzle at 20PSI and a 4-way purity mask or, in the case
of hGx adrenal implants, a Beckman Coulter CytoFlex SRT (488nm 50 mW, 633nm
100 mW, 405nm 90 mW, and 561nm 30 mW), set with 100µm nozzle at 15PSI, also
using 4-way purity. Single-cell deposition in 96-well plates was performed using single-
cell sorting mode. Intact cells were gated on FSC-A vs SSC-A plot, followed by doublet
exclusion on FSC-A vs FSC-H and SSC-A vs SSC-H, prior to gating on fluorescent
parameters for the markers described in the results.

Single cell RNA sequencing of time-resolved hemogenic gastruloids

Gastruloids were collected at different timepoints of the protocol, disassembled and
FACS deposited into 96-well plates, either as unsorted (global) or sorted by CD45,
CD41 and C-Kit/ScaI markers (sorted). RNA from the cells were extracted from single
cells using Smart-seq2 technology at 500Kb and depth of 151 bases. Sequencing
reads were quality-checked using FastQC (v0.11.9). We trimmed the samples using
trimGalore! (0.6.6) with a cutoff of 30, clipping 15 base pairs and retaining reads of
more than 100 bases. Alignment was performed on STAR (2.7.8a) and Mus musculus
annotations from GENCODE vM26. Aligned BAM files were annotated using
featureCounts (v2.0.1) and count matrices were computed in python by directly
accessing the BAM files with gtfparse packages and collapsing lanes.

For quality control, based on the histogram of counts and multimodality distributions,

we set a minimum count threshold of 200000 counts, a minimum threshold of 1000

expressed genes, and a maximum threshold of 20% mitochondrial fraction per cell.

We performed the same procedure of the second dataset with a more restrictive

threshold of 400000 counts and similar expressed genes and mitochondrial

thresholds. Cells that did not pass the quality control metrics were omitted from

analysis. We normalized the cells to the mean count number per dataset and applied

a plus-one-log transformation of the data before proceeding to the downstream

analysis.

Dimensionality reduction was performed on feature selection of the gene space using

the function scanpy.highly_varying_genes with default parameters. Selection of

principal components in principal component analysis (PCA) was performed by

heuristic elbow method. Nearest neighbor analysis was constructed by KNN graphs

using a correlation metric and 10 nearest neighbors. Data was projected for low

dimensional visualization using the UMAP algorithm with default parameters as

implemented in scanpy.tl.umap. We used the leiden algorithm as implemented in

scanpy.tl.leiden to partition the data into clusters. In order to assess the election of the

resolution parameter we used Newman-Girvan modularity as a metric of clustering

quality. Differential expressed genes were computed comparing each cluster against

the rest using the Wilcoxon test with Benjamini-Hochberg correction.

Annotation and projection to additional datasets

Raw counts matrices from Thambyrajah *et al.* (2024) were downloaded and processed

following the same pipeline as described above for scRNA-seq of gastruloids to

generate UMAP and clustering. We implemented the scmap algorithm to compare our

scRNA sequencing of gastruloids with the available datasets. We reduced the

dimensionality of the space by selecting highly varying genes from the annotated

dataset. Then, we constructed a KNN classifier with correlation metric and computed

the nearest neighbors of the target data. If neighbors with correlation metrics below

0.7 default standards, the projected cells were not projected onto any cell from the

annotated dataset. To visualize the cells over the UMAP plots of the other datasets,

we constructed a KNN regressor with three neighbors and a correlation metric.
Confusion matrices were used to visualize the overlap between gastruloid clusters and
the embryo dataset assigned clusters using scmap.

Bulk RNA sequencing

Total RNA was extracted from disassembled gastruloid cells at 216 hours. Sequencing
was performed on NovaSeq PE150 platform, at 20M paired-end reads per sample.
Tophat2 with Bowtie2 were used to map paired-end reads to the reference Mus
musculus genome build GRCm39. GENCODE Release M30 (Frankish *et al.,* 2019)
was used as the reference mouse genome annotation. Aligned reads were filtered by
quality using samtools (Li, H. *et al.,* 2009) with a minimum threshold set at 30 (q30).
Transcript assembly and quantification was achieved using htseq (Putri et al., 2022).
Differential expression between sample and control was performed by collapsing
technical replicates for each condition using Deseq2 (Love, Huber and Anders, 2014)
in R environment (Deseq2 library v 1.32.0). The differential expression was expressed
in the form of log2 fold change and filtered by false discovery rate (FDR) of 0.05.

Gene list enrichment analyses

Gene ontology (GO) analysis was performed in ExpressAnalyst (available at
www.expressanalyst.ca) using the PANTHER or GO Biological Process (BP)
repository. GO terms and pathways were filtered by false discovery rate (FDR) with a
cut-off of ≤ 0.1 for meaningful association. EnrichR (Xie *et al.,* 2021; Kuleshov *et al.,*
2016; Chen, E. Y. *et al.,* 2013) was used for cell type analysis using the Panglao DB
(Franzén, Gan and Björkegren, 2019) databases using a FDR threshold of ≤ 0.1 or p
value ≤ 0.01, where specified, as well as transcription factor (TF) binding site
enrichment using the ENCODE and ChEA Consensus TFs from ChIP database.

Gene Set Enrichment Analysis (GSEA)

Custom gene signatures were used as gene sets for GSEA analysis (Subramanian *et*

al., 2005) on the GSEA software v4.2.3 on RNA sequencing expression values in

counts units. GSEA was ran in 10000 permutations on gene set using the weighted

Signal2Noise metric. Enrichment metrics are shown as normalized enrichment score

(NES) and filtered by FDR ≤ 0.05. Leading edge genes (LEGs) are genes with a "Yes"

values for core enrichment. For AML patient analysis, clinical phenotype and

expression data (in counts units) were extracted from the GDC TARGET-AML cohorts

in the Therapeutically Applicable Research to Generate Effective Treatments project

(TARGET, https://ocg.cancer.gov/programs/target), downloaded from the University

of California Santa Cruz (UCSC) Xena public repository (last accessed 31st August

2022). Patient samples were selected according to the reported karyotype to include

t(7;12), inv(16), MLL, normal karyotype, and t(8;21). GSEA was performed comparing

RNA sequencing counts of t(7;12) samples against pooled AML subtypes (inv(16),

MLL, normal karyotype and t(8;21)) as "other AML".

Real-time polymerase chain reaction (qPCR)

Extracted RNA was reverse-transcribed into complementary DNA (cDNA) using High-

Capacity RNA-to-cDNA Kit (Applied Biosystems). QPCR was performed using

FastGene 2x IC Green Universal qPCR Mix (Nippon Genetics, Duren, Germany) using

primers for human MNX1 (forward 5-GTTCAAGCTCAACAAGTACC-3; reverse 5-

GGTTCTGGAACCAAATCTTC-3) (Gulino et al., 2021) and Ppia for endogenous

control   (forward   5-TTACCCATCAAACCATTCCTTCTG-3;   reverse   5-

AACCCAAAGAACTTCAGTGAGAGC-3) (Moris *et al.,* 2018). Differential gene

expression was calculated using the delta delta Ct (ΔΔCt) method.

Polymerase chain reaction (PCR)

Genomic DNA was extracted using Monarch Genomic DNA Purification Kit (New

England Biolabs, Hitchin, UK) using manufacturer's instructions. PCR on genomic

DNA was performed using Phire PCR Master Mix (Thermo Scientific) to amplify the

BFP locus using forward primer 5'-GCACCGTGGACAACCATCACTT-3' and reverse

primer 5'-CAGTTTGCTAGGGAGGTCGC-3'.

Statistical analysis

Experiments were performed at least in triplicates, unless specified otherwise. Data

are plotted to include standard deviation (+/-SD) between replicates. Statistical

significance was set at a threshold of p value < 0.05. Statistical analysis was performed

in R environment (version 4.1.3) or using GraphPad Prism 8.0 software and is detailed

in respective figure legends.

Data availability

Raw data as well as processed count matrices and post-processed files from single-

cell RNA-seq for the time-resolved data is available at E-MTAB-12148. Bulk RNA-seq

of MNX1 overexpressing gastruloids is available at Array Express with accession code

E-MTAB-12173. The post-processing was performed in Python on DockerHub:

dsblab/single_cell_analysis:0.5. Scripts are available in https://github.com/dsb-

lab/blood_gastruloids and Zenodo (https://doi.org/10.5281/zenodo.7053423). The

results published here are partly based upon data generated by the Therapeutically

Applicable   Research   to   Generate   Effective   Treatments   (TARGET)

(https://ocg.cancer.gov/programs/target) initiative, of the Acute Myeloid Leukemia

(AML) cohort GDC TARGET-AML. The data used for this analysis are available at

https://portal.gdc.cancer.gov/projects and https://xenabrowser.net/.

ACKNOWLEDGEMENTS

This project was funded by a start-up grant and a BRIEF award from Brunel University

London to CP, by a Little Princess Trust (LPT) Project Grant (CCLGA 2023 22 Pina)

to CP, and by an ERC Advanced Grant (MiniEmbryoBlueprint 834580) to AMA. LPT

research is funded in partnership with CCLG through the CCLG Charity Research
Network. DR received funding from the Royal Society of Biology (MRSB Travel Grant).
GTC was funded by grant FPU18/05091 from the Spanish Ministry of Universities. AJ
is funded by a Lady Tata Memorial Trust International Scholarship (2022). SvdB was
funded by an EMBO Postdoctoral Fellowship (ALTF 195-2021) and is in receipt of a
HFSP Postdoctoral Fellowship (LT0047/2022-L). Work in the CP lab was also funded
by a KKLF Intermediate Fellowship (KL888), a Leuka John Goldman Fellowship for
Future Science (2017-2019), and a Wellcome Trust / ISSF Bridge Funding award at
the University of Cambridge (2019). JGO acknowledges financial support from the
Spanish Ministry of Science and Innovation and FEDER (grant PGC2018-101251-B-
I00), by the Maria de Maeztu Programme for Units of Excellence in R&D (grant
CEX2018-000792-M), and by the Generalitat de Catalunya (ICREA Academia
programme). MP acknowledges funding from the Italian Ministry of Health (Ricerca
Finalizzata 5 per mille and Ricerca Corrente to MP). Library preparation and next-
generation sequencing for single-cell RNA-seq analysis were performed by the Single
Cell Genomics Group at the National Centre for Genomic Analysis – Centre for
Genomic Regulation (CNAG-CRG), Barcelona. The Authors wish to acknowledge
Tina Balayo, Ana Filipa Domingues, Shikha Gupta, Oliver Davies, Kristen Place and
Remisha Gurung's technical support at different stages of the project.

AUTHOR CONTRIBUTIONS

Conceptualization: CP, AMA; Methodology: CWS, DR, GTC, FP, SvdB, MC,MP, KRK,
VH-H, AB, JGO, AMA, CP; Software: GTC, JGO; Validation: CWS, DR, AJ, YC, SvdB,
CP; Investigation: CWS, DR, GTC, FP, AJ, YC, LD, CB, G-AI, CP; Formal Analysis:
GTC, DR, JGO, CP; Resources: GTC, JC, MP, JGO, AMA; Data curation: DR, GTC,
JGO; Writing – Original Draft: CP, DR, GTC; Writing – Review and editing: CP, DR,
GTC, AB, JGO, AMA; Visualisation: DR, GTC, CWS, CP; Supervision: CP; Project
administration: CP, AMA, JGO; Funding acquisition: CP, AMA.

DECLARATION OF INTERESTS

917 AMA and CP are co-inventors in the patent application PCT/GB2019/052668:
918 Polarised Three-Dimensional Cellular Aggregates. The other authors have no
919 interests to declare.

SUPPLEMENTAL INFORMATION

923 Supplemental Figures S1-S7

924 Supplemental Files S1-S3 (Excel files)

 FIGURE LEGENDS

Figure 1 – Hemogenic gastruloids (hGx) produced from mES cells promote hemato-
endothelial specification with spatio-temporally accurate ontogeny. (A) Timeline of mES
cells assembly and culture into gastruloids over a 216h period with the addition of specified
factors for the promotion of hemato-endothelial specification. The 24-hour pre-treatment in
2i+LIF was omitted when mES cultures showed no signs of differentiation. **(B)** Imaging of hGx
over time from 72h to 216h at 10x magnification, showing the assembly and growth of the 3D
structures and the polarization of the Flk-1-GFP marker from 96h; scale bar: 100mm. (**C)** Flow
cytometry timecourse analysis of hGx for the presence of CD144 (VE-cadherin), C-Kit, CD41,
and CD45 markers; representative plots. **(D)** Quantification of analysis in (D). Violin plots with
median and interquartile range of up to 26 replicates; Kruskal-Wallis CD144 (p=0.0755), C-Kit
(p<0.0001), CD41 (p<0.0342), CD45 (p<0.0001). Shown are significant Dunn's multiple
comparisons tests; adjusted q-value. **(E)** Immunostaining of whole individual gastruloids at
216h showing the localized expression of CD31 (yellow), C-Kit (turquoise), and CD45 (pink)
markers; scale bar: 100μm.

Figure 2 – Time-resolved analysis of scRNA-seq of hGxs captures successive waves of
hematopoietic specification. (A) UMAP projection of all sequenced cells colored by
annotated clusters (left), by sorting markers C-Kit/ScaI, CD41, and CD45, or unfractionated
cells ('whole' corresponding to 'all' in panel A) (center), and by hGx culture time (right). **(B)**
Heatmap representation of the expression of endothelial and hematopoietic marker genes in
individual hGx C-Kit$^+$ cells sorted at 144 and 192h. Cells and genes ordered by unsupervised
hierarchical clustering; cell cluster dendrogram in Fig. S2E. **(C)** Heatmap representation of the
expression of endothelial and hematopoietic marker genes in individual hGx CD41$^+$ and
CD45$^+$ cells sorted at 144 and 192/216h, respectively. Cells and genes ordered by
unsupervised hierarchical clustering; cell cluster dendrogram in Fig. S2F. **(D)** Summary
representation of the proportion of hGx cells expressing endothelial (endo), erythroid/myeloid
(Ery+My), myeloid/lymphoid (My+Ly) and HSPC gene signatures. Endo: *Kdr*; Ery+My:
Epor±Gata1±Klf1±Hbb±Hba±Hbg + Spi1±Mpo±Anpep±Csf1/2/3r); My+Ly: *Spi1±Csf1/2a/3r +*
Ikzf1±Ighm/d±Igk); HSPC: *Ptprc + Myb*). **(E-G)** Cell type-associations of selected surface
markers and transcription factors affiliated with the HE-to-hematopoietic transition (*CD44*,
Procr, *Etv6*, *Gata2*, *Notch1*, *Tal1*) and with hematopoietic (*Gfi1b*, *Mllt3*, *Runx1*) and definitive
hematopoietic (*Ikzf1*, *Hlf*, *Hoxa9*, *Myb*) gene expression programs with **(E)** C-Kit$^+$ (endo-haem)
and CD41$^+$/CD45$^+$ (haem), **(F)** endo-haem cells obtained at 144h or 192h, and **(G)** CD41$^+$ or

CD45+ cells. Gene expression analysed as present (non-zero expression) or absent (zero-
expression). Odds ratio with significance ($-\log_{10}p<1$) estimated by Fisher's test.

Figure 3 – HGxs specify cells with pre-HSC characteristics. (A) Flow cytometry analysis
of 216h *Rosa26-BFP::Flk1-GFP* hGx cells for CD41, CD45, c-Kit and CD144 (VE-Cadherin).
(B) Quantification of pre-HSC-like c-Kit+CD144+ cells within the CD45+CD41lo fraction of 216h-
hGx; mean of 4 independent replicate experiments. **(C)** Colony-forming cell (CFC) assay of
216h-hGx in multipotential methylcellulose-based medium. GEM: granulocyte-erythroid-
monocyte; GM: granulocyte-monocyte; M: monocyte; E: erythroid. Please note that the
medium does not include TPO and does not assess the presence of megakaryocytic
progenitors. CFC frequency in 3hGx, n=3, mean±SD. **(D)** Representative photographs of
colonies quantified in C; scale bar=2mm. **(E)** Schematic representation of the experimental
workflow for execution and analysis of hGx implantation in the adrenal gland of
immunodeficient mice. **(E)** Histology of implanted and control adrenal glands; paraffin sections
stained with hematoxylin & eosin (H&E). Inserts in implanted animal highlight neural (orange)
and hematopoietic and renal tubular epithelium (green); inserts in control animal highlight
normal cortical (blue) and medullar (red) adrenal architecture, absent in implanted adrenal
glands. **(F)** PCR detection of hGx genomic (g)DNA in the bone marrow (BM) and spleen (Spl)
of immunodeficient mice 1 month after unilateral adrenal implantation of 3 gastruloids/gland.
Analysis of 5 replicates of 100ng gDNA /recipient tissue; control animal was injected
unilaterally with PBS in an adrenal gland in parallel with experimental implantation. Reaction
positive control used 100ng of gDNA from *Rosa26-BFP*::*Flkj1-GFP* mES cells (mES) used to
generate hGxs. NTC: no template control. See Fig. S4F-H for additional details.

Figure 4 – HGxs with MNX1 overexpression have increased cellularity and enhanced
hemogenic endothelial potential. (A) Cell type enrichment analysis in MNX1-r AML
transcriptomes, compared to normal paediatric bone marrow (BM) or other paediatric AML
from the TARGET database. GSEA used representative cell type gene sets from the 2021 DB
database; bubble plot shows NES scores (color gradient) and statistical significance (–
log10(FDR), bubble size). **(B)** Schematic representation of generation of *MNX1-*
overexpressing and empty vector (EV) mES cells by lentiviral transduction, with subsequent
assembly into hGxs (haemGX). **(C)** Imaging of hGxs with *MNX1* overexpression and EV
controls at 10x magnification, showing appropriate assembly and polarization of the Flk1-GFP
marker. scale bar: 300mm. **(D)** Size of hGx at each timepoint, determined by the distance
between the furthest extreme points in μm. Mean ± SD of 3 replicate experiments; 2-tailed t-

test, p< 0.05 (*), 0.001 (**), 0.0001 (***), and 0.00001 (****). **(E-G)** Flow cytometry timecourse

analysis of **(E)** C-Kit$^+$, **(F)** CD41$^+$, and **(G)** CD45$^+$ cell abundance in MNX1 and EV hGxs (120-

216h). Mean ± standard deviation of 3-7 independent experiments; 2-way ANOVA and Sidak's

multiple comparison test significant at p<0.05 for C-Kit$^+$ cells only (construct contribution to

variance p=0.0191; 144h comparison *p=0.0190). **(H)** Volcano plots of differentially expressed

genes (DEGs) by bulk RNA-seq between MNX1 and EV hGxs at 144h and 216h. Intersection

of statistically significant DEGs between 144 h and 216 h (shown in Venn diagram on top right)

identified unique DEGs for each condition (labelled in red for 144 h and blue for 216 h). **(J)**

Gene Ontology (GO) term enrichment for differentially upregulated genes between MNX1 and

EV hGx at 144h (red) and 216h (blue), computed in EnrichR using the GO Biological Process

repository. **(I)** Transcription factor (TF) binding site enrichment on differentially upregulated

genes between MNX1 and EV hGx at 144h (red) and 216 hr (blue) using the ENCODE and

ChEA Consensus TFs from ChIP database on EnrichR.

1008

Figure 5 – MNX1 selects C-Kit$^+$ clonogenic cells and selectively transforms end-stage

hGxs. (A) Representative photographs of serial replating of colony-forming cell assays

initiated from EV or MNX1 cells obtained at 144h of the hGx protocol. **(B)** Quantification of

colony-replating efficiency of EV and MNX1 144h-hGx cells. Mean+SD of n>3 replicates;

Kruskal-Wallis with Dunn's multiple comparison testing at significant q<0.05. **(C)**

Representative photographs of serial replating of colony-forming cell assays initiated from EV

or MNX1 cells obtained at 216h of the hGx protocol. **(D)** Quantification of colony-replating

efficiency of EV and MNX1 216h-hGx cells. Mean+SD of n=5-8 replicates; Kruskal-Wallis with

Dunn's multiple comparison testing at significant q<0.05. **(E)** Flow cytometry analysis of

hemogenic progenitor C-Kit$^+$ cells and myeloid-affiliated CD11b$^+$ cells at first round of plating

of 144h and 216h-hGxs initiated from EV and MNX1 cells; representative plots. **(F)**

Representative flow cytometry plots of serially re-plating MNX1 cells from 144h and 216h-

gastruloids (plate 5 cells).

1022

Figure 6 – HGxs with *MNX1* overexpression recapitulate MNX1-Acute Myeloid Leukemia

patient signatures. (A) Quantification of human *MNX1* transcripts in FPKM units from RNA-

seq of GX-EV, GX-MNX1, and CFC at 144 hr and 216 hr. Bars represent mean of replicates.

(B) Heatmap comparing the expression of all differentially expressed genes (DEGs) between

all conditions (GX-MNX1 144 hr vs GX-EV 144 hr; GX-MNX1 216 hr vs GX-EV 216 hr; CFC

MNX1 144 hr vs GX MNX1 144 hr; CFC MNX1 216 hr vs GX MNX1 216 hr) as Z score.

Hierarchical clustering by Ward D method on Euclidean distances identifies 14 clusters (k).

(C) GSEA plots for gene set extracted from k14 from **(B)** against MNX1-r patients RNA-seq

counts vs MLL, core-binding factors (CBF), or other pediatric AML. **(D)** Cell type analysis of

the intersect of leading edge genes LEGs (n=476) from Fig. S5D in MNX1-r patients (k14

LEGs) **(B)** using the Panglao DB 2021 database. **(E)** Representative Giemsa-Wright stained

dissociated CFC replating MNX1 hGx cells from 144h and 216h. Solid black arrows, mast cell

precursors; dashed black arrows, mast cells.

REFERENCES

Alexander, T.B. and Mullighan, C.G. (2021) 'Molecular biology of childhood leukemia',
Annual Review of Cancer Biology, 5, pp. 95-117.

Ballabio, E., Cantarella, C.D., Federico, C., Di Mare, P., Hall, G., Harbott, J., Hughes, J.,
Saccone, S. and Tosi, S. (2009) 'Ectopic expression of the HLXB9 gene is associated with
an altered nuclear position in t (7; 12) leukaemias', *Leukemia,* 23(6), pp. 1179-1183.

Beccari, L., Moris, N., Girgin, M., Turner, D.A., Baillie-Johnson, P., Cossy, A., Lutolf, M.P.,
Duboule, D. and Arias, A.M. (2018) 'Multi-axial self-organization properties of mouse
embryonic stem cells into gastruloids', *Nature,* 562(7726), pp. 272-276.

Boisset, J., van Cappellen, W., Andrieu-Soler, C., Galjart, N., Dzierzak, E. and Robin, C.
(2010) 'In vivo imaging of haematopoietic cells emerging from the mouse aortic
endothelium', *Nature,* 464(7285), pp. 116-120.

Bolouri, H., Farrar, J.E., Triche Jr, T., Ries, R.E., Lim, E.L., Alonzo, T.A., Ma, Y., Moore, R.,
Mungall, A.J. and Marra, M.A. (2018) 'The molecular landscape of pediatric acute myeloid
leukemia reveals recurrent structural alterations and age-specific mutational interactions',
Nature medicine, 24(1), pp. 103.

Bousquets-Muñoz, P., Molina, O., Varela, I., Álvarez-Eguiluz, Á, Fernández-Mateos, J.,
Gómez, A., Sánchez, E.G., Balbín, M., Ruano, D. and Ramírez-Orellana, M. (2024)
'Backtracking NOM1:: ETV6 fusion to neonatal pathogenesis of t (7; 12)(q36; p13) infant
AML', *Leukemia,* , pp. 1-5.

Bowie, M.B., McKnight, K.D., Kent, D.G., McCaffrey, L., Hoodless, P.A. and Eaves, C.J.
(2006) 'Hematopoietic stem cells proliferate until after birth and show a reversible phase-
specific engraftment defect', *The Journal of clinical investigation,* 116(10), pp. 2808-2816.

Brignole, C., Calarco, E., Bensa, V., Giusto, E., Perri, P., Ciampi, E., Corrias, M.V.,
Astigiano, S., Cilli, M. and Loo, D. (2023) 'Antitumor activity of the investigational B7-H3
antibody-drug conjugate, vobramitamab duocarmazine, in preclinical models of
neuroblastoma', *Journal for Immunotherapy of Cancer,* 11(9).

Calvanese, V., Nguyen, A.T., Bolan, T.J., Vavilina, A., Su, T., Lee, L.K., Wang, Y., Lay, F.D.,
Magnusson, M. and Crooks, G.M. (2019) 'MLLT3 governs human haematopoietic stem-cell
self-renewal and engraftment', *Nature,* 576(7786), pp. 281-286.

Cazzola, A., Cazzaniga, G., Biondi, A., Meneveri, R., Brunelli, S. and Azzoni, E. (2021)
'Prenatal origin of pediatric leukemia: lessons from hematopoietic development', *Frontiers in*
cell and developmental biology, 8, pp. 618164.

Chandrakanthan, V., Rorimpandey, P., Zanini, F., Chacon, D., Olivier, J., Joshi, S., Kang,
Y.C., Knezevic, K., Huang, Y. and Qiao, Q. (2022) 'Mesoderm-derived PDGFRA cells
regulate the emergence of hematopoietic stem cells in the dorsal aorta', *Nature cell biology,*
24(8), pp. 1211-1225.

Chang, Y., Hummel, S.N., Jung, J., Jin, G., Deng, Q. and Bao, X. (2023) 'Engineered hematopoietic and immune cells derived from human pluripotent stem cells', *Experimental hematology,* .

Chaudhury, S., O'Connor, C., Cañete, A., Bittencourt-Silvestre, J., Sarrou, E., Prendergast, Á, Choi, J., Johnston, P., Wells, C.A. and Gibson, B. (2018) 'Age-specific biological and molecular profiling distinguishes paediatric from adult acute myeloid leukaemias', *Nature communications,* 9(1), pp. 5280.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V. and Clark, N.R. (2013) 'A. 1126 Ma'ayan, Enrichr: Interactive and collaborative HTML5 gene list enrichment analysis tool. BMC 1127', *Bioinformatics,* .

Chen, W., O'Sullivan, M.G., Hudson, W. and Kersey, J. (2011) 'Modeling human infant MLL leukemia in mice: leukemia from fetal liver differs from that originating in postnatal marrow', *Blood,* 117(12), pp. 3474.

Chia, S.L., Kapoor, S., Carvalho, C., Bajénoff, M. and Gentek, R. (2023) 'Mast cell ontogeny: From fetal development to life-long health and disease', *Immunological reviews,* 315(1), pp. 31-53.

Ciau-Uitz, A., Pinheiro, P., Gupta, R., Enver, T. and Patient, R. (2010) 'Tel1/ETV6 specifies blood stem cells through the agency of VEGF signaling', *Developmental cell,* 18(4), pp. 569-578.

Ciriza, J., Thompson, H., Petrosian, R., Manilay, J.O. and García-Ojeda, M.E. (2013) 'The migration of hematopoietic progenitors from the fetal liver to the fetal bone marrow: lessons learned and possible clinical applications', *Experimental hematology,* 41(5), pp. 411-423.

Costa, G., Kouskoff, V. and Lacaud, G. (2012) 'Origin of blood cells and HSC production in the embryo', *Trends in immunology,* 33(5), pp. 215-223.

Dang, S.M., Kyba, M., Perlingeiro, R., Daley, G.Q. and Zandstra, P.W. (2002) 'Efficiency of embryoid body formation and hematopoietic development from embryonic stem cells in different culture systems', *Biotechnology and bioengineering,* 78(4), pp. 442-453.

Deguchi, Y. and Kehrl, J.H. (1991) 'Selective expression of two homeobox genes in CD34-positive cells from human bone marrow', *Blood,* 78(2), pp. 323-328.

Diaz, M.F., Li, N., Lee, H.J., Adamo, L., Evans, S.M., Willey, H.E., Arora, N., Torisawa, Y., Vickers, D.A. and Morris, S.A. (2015) 'Biomechanical forces promote blood development through prostaglandin E2 and the cAMP–PKA signaling axis', *Journal of Experimental Medicine,* 212(5), pp. 665-680.

Dijkhuis, L., Johns, A., Ragusa, D., van den Brink, Susanne C and Pina, C. (2023) 'Haematopoietic development and HSC formation in vitro: promise and limitations of gastruloid models', *Emerging Topics in Life Sciences,* 7(4), pp. 439-454.

Dong, Y., Shi, O., Zeng, Q., Lu, X., Wang, W., Li, Y. and Wang, Q. (2020) 'Leukemia incidence trends at the global, regional, and national level between 1990 and 2017', *Experimental hematology & oncology,* 9, pp. 1-11.

Dzierzak, E. and Bigas, A. (2018) 'Blood development: hematopoietic stem cell dependence and independence', *Cell stem cell,* 22(5), pp. 639-651.

Espersen, A.D.L., Noren-Nyström, U., Abrahamsson, J., Ha, S., Pronk, C.J., Jahnukainen, K., Jónsson, ÓG., Lausen, B., Palle, J. and Zeller, B. (2018) 'Acute myeloid leukemia (AML) with t (7; 12)(q36; p13) is associated with infancy and trisomy 19: Data from Nordic Society for Pediatric Hematology and Oncology (NOPHO-AML) and review of the literature', *Genes, Chromosomes and Cancer,* 57(7), pp. 359-365.

Fehling, H.J., Lacaud, G., Kubo, A., Kennedy, M., Robertson, S., Keller, G. and Kouskoff, V. (2003) 'Tracking mesoderm induction and its specification to the hemangioblast during embryonic stem cell differentiation', .

Fidanza, A. and Forrester, L.M. (2021) 'Progress in the production of haematopoietic stem and progenitor cells from human pluripotent stem cells', *Journal of Immunology and Regenerative Medicine,* 13, pp. 100050.

Fitch, S.R., Kimber, G.M., Wilson, N.K., Parker, A., Mirshekar-Syahkal, B., Göttgens, B., Medvinsky, A., Dzierzak, E. and Ottersbach, K. (2012) 'Signaling from the sympathetic nervous system regulates hematopoietic stem cell emergence during embryogenesis', *Cell stem cell,* 11(4), pp. 554-566.

Fowler, J.L., Zheng, S.L., Nguyen, A., Chen, A., Xiong, X., Chai, T., Chen, J.Y., Karigane, D., Banuelos, A.M. and Niizuma, K. (2024) 'Lineage-tracing hematopoietic stem cell origins in vivo to efficiently make human HLF HOXA hematopoietic progenitors from pluripotent stem cells', *Developmental Cell,* 59(9), pp. 1110-1131. e22.

Frankish, A., Diekhans, M., Ferreira, A., Johnson, R., Jungreis, I., Loveland, J., Mudge, J.M., Sisu, C., Wright, J. and Armstrong, J. (2019) 'GENCODE reference annotation for the human and mouse genomes', *Nucleic acids research,* 47(D1), pp. D766-D773.

Franzén, O., Gan, L. and Björkegren, J.L. (2019) 'PanglaoDB: a web server for exploration of mouse and human single-cell RNA sequencing data', *Database,* 2019, pp. baz046.

Fujimoto, T., Ogawa, M., Minegishi, N., Yoshida, H., Yokomizo, T., Yamamoto, M. and Nishikawa, S. (2001) 'Step-wise divergence of primitive and definitive haematopoietic and endothelial cell lineages during embryonic stem cell differentiation', *Genes to Cells,* 6(12), pp. 1113-1127.

Ganuza, M., Hall, T., Myers, J., Nevitt, C., Sánchez-Lanzas, R., Chabot, A., Ding, J., Kooienga, E., Caprio, C. and Finkelstein, D. (2022) 'Murine foetal liver supports limited detectable expansion of life-long haematopoietic progenitors', *Nature cell biology,* 24(10), pp. 1475-1486.

Ghiaur, G., Ferkowicz, M.J., Milsom, M.D., Bailey, J., Witte, D., Cancelas, J.A., Yoder, M.C. and Williams, D.A. (2008) 'Rac1 is essential for intraembryonic hematopoiesis and for the initial seeding of fetal liver with definitive hematopoietic progenitor cells', *Blood, The Journal of the American Society of Hematology,* 111(7), pp. 3313-3321.

Harrison, K.A., Thaler, J., Pfaff, S.L., Gu, H. and Kehrl, J.H. (1999) 'Pancreas dorsal lobe agenesis and abnormal islets of Langerhans in Hlxb9-deficient mice', *Nature genetics,* 23(1), pp. 71-75.

Hooper, M., Hardy, K., Handyside, A., Hunter, S. and Monk, M. (1987) 'HPRT-deficient (Lesch–Nyhan) mouse embryos derived from germline colonization by cultured cells', *Nature,* 326(6110), pp. 292-295.

Ivanovs, A., Rybtsov, S., Welch, L., Anderson, R.A., Turner, M.L. and Medvinsky, A. (2011) 'Highly potent human hematopoietic stem cells first emerge in the intraembryonic aorta-gonad-mesonephros region', *Journal of Experimental Medicine,* 208(12), pp. 2417-2427.

Jakobsson, L., Franco, C.A., Bentley, K., Collins, R.T., Ponsioen, B., Aspalter, I.M., Rosewell, I., Busse, M., Thurston, G. and Medvinsky, A. (2010) 'Endothelial cells dynamically compete for the tip cell position during angiogenic sprouting', *Nature cell biology,* 12(10), pp. 943-953.

Kapeni, C., Nitsche, L., Kilpatrick, A.M., Wilson, N.K., Xia, K., Mirshekar-Syahkal, B., Chandrakanthan, V., Malouf, C., Pimanda, J.E. and Göttgens, B. (2022) 'p57Kip2 regulates embryonic blood stem cells by controlling sympathoadrenal progenitor expansion', *Blood, The Journal of the American Society of Hematology,* 140(5), pp. 464-477.

Kikuchi, K. and Kondo, M. (2006) 'Developmental switch of mouse hematopoietic stem cells from fetal to adult type occurs in bone marrow after birth', *Proceedings of the National Academy of Sciences,* 103(47), pp. 17852-17857.

Kiselev, V.Y., Yiu, A. and Hemberg, M. (2018) 'scmap: projection of single-cell RNA-seq data across data sets', *Nature methods,* 15(5), pp. 359-362.

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S.L., Jagodnik, K.M. and Lachmann, A. (2016) 'Enrichr: a comprehensive gene set enrichment analysis web server 2016 update', *Nucleic acids research,* 44(W1), pp. W90-W97.

Lacaud, G. and Kouskoff, V. (2017) 'Hemangioblast, hemogenic endothelium, and primitive versus definitive hematopoiesis', *Experimental hematology,* 49, pp. 19-24.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R. and 1000 Genome Project Data Processing Subgroup (2009) 'The sequence alignment/map format and SAMtools', *bioinformatics,* 25(16), pp. 2078-2079.

Li, W., Ferkowicz, M.J., Johnson, S.A., Shelley, W.C. and Yoder, M.C. (2005) 'Endothelial cells in the early murine yolk sac give rise to CD41-expressing hematopoietic cells', *Stem cells and development,* 14(1), pp. 44-54.

Lis, R., Karrasch, C.C., Poulos, M.G., Kunar, B., Redmond, D., Duran, J.G.B., Badwe, C.R., Schachterle, W., Ginsberg, M. and Xiang, J. (2017) 'Conversion of adult endothelium to immunocompetent haematopoietic stem cells', *Nature,* 545(7655), pp. 439-445.

Love, M., Huber, W. and Anders, S. (2014) 'No title', *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2 Genome Biology 15 (12), 550,* .

Lundin, V., Sugden, W.W., Theodore, L.N., Sousa, P.M., Han, A., Chou, S., Wrighton, P.J., Cox, A.G., Ingber, D.E. and Goessling, W. (2020) 'YAP regulates hematopoietic stem cell formation in response to the biomechanical forces of blood flow', *Developmental cell,* 52(4), pp. 446-460. e5.

Masetti, R., Vendemini, F., Zama, D., Biagi, C., Pession, A. and Locatelli, F. (2015) 'Acute myeloid leukemia in infants: biology and treatment', *Frontiers in pediatrics,* 3, pp. 37.

McGrath, K.E., Frame, J.M., Fegan, K.H., Bowen, J.R., Conway, S.J., Catherman, S.C., Kingsley, P.D., Koniski, A.D. and Palis, J. (2015) 'Distinct sources of hematopoietic progenitors emerge before HSCs and provide functional blood cells in the mammalian embryo', *Cell reports,* 11(12), pp. 1892-1904.

McGrath, K.E. and Palis, J. (2005) 'Hematopoiesis in the yolk sac: more than meets the eye', *Experimental hematology,* 33(9), pp. 1021-1028.

Medvinsky, A. and Dzierzak, E. (1996) 'Definitive hematopoiesis is autonomously initiated by the AGM region', *Cell,* 86(6), pp. 897-906.

Medvinsky, A., Rybtsov, S. and Taoudi, S. (2011) 'Embryonic origin of the adult hematopoietic system: advances and questions', *Development,* 138(6), pp. 1017-1031.

Mercher, T., Raffel, G.D., Moore, S.A., Cornejo, M.G., Baudry-Bluteau, D., Cagnard, N., Jesneck, J.L., Pikman, Y., Cullen, D. and Williams, I.R. (2009) 'The OTT-MAL fusion oncogene activates RBPJ-mediated transcription and induces acute megakaryoblastic leukemia in a knockin mouse model', *The Journal of clinical investigation,* 119(4), pp. 852-864.

Miladinovic, O., Canto, P., Pouget, C., Piau, O., Radic, N., Freschu, P., Megherbi, A., Brujas Prats, C., Jacques, S. and Hirsinger, E. (2024) 'A multistep computational approach reveals a neuro-mesenchymal cell population in the embryonic hematopoietic stem cell niche', *Development,* 151(7), pp. dev202614.

Moore, M.A. and Metcalf, D. (1970) 'Ontogeny of the haemopoietic system: yolk sac origin of in vivo and in vitro colony forming cells in the developing mouse embryo', *British journal of haematology,* 18(3), pp. 279-296.

Moris, N., Edri, S., Seyres, D., Kulkarni, R., Domingues, A.F., Balayo, T., Frontini, M. and Pina, C. (2018) 'Histone acetyltransferase KAT2A stabilizes pluripotency with control of transcriptional heterogeneity', *Stem cells,* 36(12), pp. 1828-1838.

Murry, C.E. and Keller, G. (2008) 'Differentiation of embryonic stem cells to clinically relevant populations: lessons from embryonic development', *Cell,* 132(4), pp. 661-680.

Nafria, M., Bonifer, C., Stanley, E.G., Ng, E.S. and Elefanty, A.G. (2020) 'Protocol for the generation of definitive hematopoietic progenitors from human pluripotent stem cells', *STAR protocols,* 1(3), pp. 100130.

Nagel, S., Scherr, M., Quentmeier, H., Kaufmann, M., Zaborski, M., Drexler, H.G. and MacLeod, R. (2005) 'HLXB9 activates IL6 in Hodgkin lymphoma cell lines and is regulated by PI3K signalling involving E2F3', *Leukemia,* 19(5), pp. 841.

Niakan, K.K., Ji, H., Maehr, R., Vokes, S.A., Rodolfa, K.T., Sherwood, R.I., Yamaki, M., Dimos, J.T., Chen, A.E. and Melton, D.A. (2010) 'Sox17 promotes differentiation in mouse embryonic stem cells by directly regulating extraembryonic gene expression and indirectly antagonizing self-renewal', *Genes & development,* 24(3), pp. 312-326.

North, T.E., Goessling, W., Peeters, M., Li, P., Ceol, C., Lord, A.M., Weber, G.J., Harris, J., Cutting, C.C. and Huang, P. (2009) 'Hematopoietic stem cell development is dependent on blood flow', *Cell,* 137(4), pp. 736-748.

Palis, J., Robertson, S., Kennedy, M., Wall, C. and Keller, G. (1999) 'Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse', *Development,* 126(22), pp. 5073-5084.

Pastorino, F., Brignole, C., Marimpietri, D., Cilli, M., Gambini, C., Ribatti, D., Longhi, R., Allen, T.M., Corti, A. and Ponzoni, M. (2003) 'Vascular damage and anti-angiogenic effects of tumor vessel-targeted liposomal chemotherapy', *Cancer research,* 63(21), pp. 7400-7409.

Pearson, S., Cuvertino, S., Fleury, M., Lacaud, G. and Kouskoff, V. (2015) 'In vivo repopulating activity emerges at the onset of hematopoietic specification during embryonic stem cell differentiation', *Stem Cell Reports,* 4(3), pp. 431-444.

Piau, O., Brunet-Manquat, M., L'homme, B., Petit, L., Birebent, B., Linard, C., Moeckes, L., Zuliani, T., Lapillonne, H. and Benderitter, M. (2023) 'Generation of transgene-free hematopoietic stem cells from human induced pluripotent stem cells', *Cell Stem Cell,* 30(12), pp. 1610-1623. e7.

Picelli, S., Faridani, O.R., Björklund, ÅK., Winberg, G., Sagasser, S. and Sandberg, R. (2014) 'Full-length RNA-seq from single cells using Smart-seq2', *Nature protocols,* 9(1), pp. 171-181.

Pina, C., May, G., Soneji, S., Hong, D. and Enver, T. (2008) 'MLLT3 regulates early human erythroid and megakaryocytic cell fate', *Cell stem cell,* 2(3), pp. 264-273.

Quintard, C., Tubbs, E., Jonsson, G., Jiao, J., Wang, J., Werschler, N., Laporte, C., Pitaval, A., Bah, T. and Pomeranz, G. (2024) 'A microfluidic platform integrating functional vascularized organoids-on-chip', *Nature Communications,* 15(1), pp. 1452.

Ragusa, D., Cicirò, Y., Federico, C., Saccone, S., Bruno, F., Saeedi, R., Sisu, C., Pina, C., Sala, A. and Tosi, S. (2022) 'Engineered model of t (7; 12)(q36; p13) AML recapitulates patient-specific features and gene expression profiles', *Oncogenesis,* 11(1), pp. 50.

Ragusa, D., Dijkhuis, L., Pina, C. and Tosi, S. (2023) 'Mechanisms associated with t (7; 12) acute myeloid leukaemia: from genetics to potential treatment targets', *Bioscience reports,* 43(1), pp. BSR20220489.

Rossi, G., Broguiere, N., Miyamoto, M., Boni, A., Guiet, R., Girgin, M., Kelly, R.G., Kwon, C. and Lutolf, M.P. (2021) 'Capturing cardiogenesis in gastruloids', *Cell stem cell,* 28(2), pp. 230-240. e6.

Rossi, G., Giger, S., Hübscher, T. and Lutolf, M.P. (2022) 'Gastruloids as in vitro models of embryonic blood development with spatial and temporal resolution', *Scientific Reports,* 12(1), pp. 13380.

Rybtsov, S., Batsivari, A., Bilotkach, K., Paruzina, D., Senserrich, J., Nerushev, O. and Medvinsky, A. (2014) 'Tracing the origin of the HSC hierarchy reveals an SCF-dependent, IL-3-independent CD43– embryonic precursor', *Stem cell reports,* 3(3), pp. 489-501.

Saorin, G., Caligiuri, I. and Rizzolio, F. (2023) *Microfluidic organoids-on-a-chip: The future of human models.* Elsevier, pp. 41.

Schyrr, F., Alonso-Calleja, A., Vijaykumar, A., Gebhard, S., Sarkis, R., Lopes, S.F., Oggier, A., De Leval, L., Nombela-Arrieta, C. and Naveiras, O. (2023) 'Adrenal extramedullary hematopoiesis as an inducible model of the adult hematopoietic niche', *bioRxiv,* , pp. 2023.03. 15.531679.

Souilhol, C., Gonneau, C., Lendinez, J.G., Batsivari, A., Rybtsov, S., Wilson, H., Morgado-Palacin, L., Hills, D., Taoudi, S. and Antonchuk, J. (2016) 'Inductive interactions mediated by interplay of asymmetric signalling underlie development of adult haematopoietic stem cells', *Nature communications,* 7(1), pp. 10784.

Sroczynska, P., Lancrin, C., Pearson, S., Kouskoff, V. and Lacaud, G. (2009) 'In vitro differentiation of embryonic stem cells as a model of early hematopoietic development', *Leukemia: Methods and Protocols,* , pp. 317-334.

Steliarova-Foucher, E., Colombet, M., Ries, L.A., Moreno, F., Dolya, A., Bray, F., Hesseling, P., Shin, H.Y. and Stiller, C.A. (2017) 'International incidence of childhood cancer, 2001–10: a population-based registry study', *The Lancet Oncology,* 18(6), pp. 719-731.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R. and Lander, E.S. (2005) 'Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles', *Proceedings of the National Academy of Sciences,* 102(43), pp. 15545-15550.

Sugimura, R., Jha, D.K., Han, A., Soria-Valles, C., Da Rocha, E.L., Lu, Y., Goettel, J.A., Serrao, E., Rowe, R.G. and Malleshaiah, M. (2017) 'Haematopoietic stem and progenitor cells from human pluripotent stem cells', *Nature,* 545(7655), pp. 432-438.

Taketani, T., Taki, T., Sako, M., Ishii, T., Yamaguchi, S. and Hayashi, Y. (2008) 'MNX1–ETV6 fusion gene in an acute megakaryoblastic leukemia and expression of the MNX1 gene in leukemia and normal B cell lines', *Cancer genetics and cytogenetics,* 186(2), pp. 115-119.

Thaler, J., Harrison, K., Sharma, K., Lettieri, K., Kehrl, J. and Pfaff, S.L. (1999) 'Active suppression of interneuron programs within developing motor neurons revealed by analysis of homeodomain factor HB9', *Neuron,* 23(4), pp. 675-687.

Thambyrajah, R., Maqueda, M., Neo, W.H., Imbach, K., Guillén, Y., Grases, D., Fadlullah, Z., Gambera, S., Matteini, F. and Wang, X. (2024) 'Cis inhibition of NOTCH1 through JAGGED1 sustains embryonic hematopoietic stem cell fate', *Nature Communications,* 15(1), pp. 1604.

Tosi, S., Kamel, Y.M., Owoka, T., Federico, C., Truong, T.H. and Saccone, S. (2015) 'Paediatric acute myeloid leukaemia with the t (7; 12)(q36; p13) rearrangement: a review of the biological and clinical management aspects', *Biomarker research,* 3(1), pp. 1.

Turner, D.A., Girgin, M., Alonso-Crisostomo, L., Trivedi, V., Baillie-Johnson, P., Glodowski, C.R., Hayward, P.C., Collignon, J., Gustavsen, C. and Serup, P. (2017) 'Anteroposterior polarity and elongation in the absence of extra-embryonic tissues and of spatially localised signalling in gastruloids: mammalian embryonic organoids', *Development,* 144(21), pp. 3894-3906.

van den Brink, Susanne C, Alemany, A., van Batenburg, V., Moris, N., Blotenburg, M., Vivie, J., Baillie-Johnson, P., Nichols, J., Sonnen, K.F. and Martinez Arias, A. (2020) 'Single-cell and spatial transcriptomics reveal somitogenesis in gastruloids', *Nature,* 582(7812), pp. 405-409.

Van den Brink, Susanne C, Baillie-Johnson, P., Balayo, T., Hadjantonakis, A., Nowotschin, S., Turner, D.A. and Martinez Arias, A. (2014) 'Symmetry breaking, germ layer specification and axial organisation in aggregates of mouse embryonic stem cells', *Development,* 141(22), pp. 4231-4242.

van den Brink, Susanne C and van Oudenaarden, A. (2021) '3D gastruloids: A novel frontier in stem cell-based in vitro modeling of mammalian gastrulation', *Trends in cell biology,* 31(9), pp. 747-759.

Veenvliet, J.V., Bolondi, A., Kretzmer, H., Haut, L., Scholze-Wittler, M., Schifferl, D., Koch, F., Guignard, L., Kumar, A.S. and Pustet, M. (2020) 'Mouse embryonic stem cells self-organize into trunk-like structures with neural tube and somites', *Science,* 370(6522), pp. eaba4937.

Vo, L.T. and Daley, G.Q. (2015) 'De novo generation of HSCs from somatic and pluripotent stem cell sources', *Blood, The Journal of the American Society of Hematology,* 125(17), pp. 2641-2648.

Waraky, A., Östlund, A., Nilsson, T., Weichenhan, D., Lutsik, P., Bähr, M., Hey, J., Tunali, G., Adamsson, J. and Jacobsson, S. (2024) 'Aberrant MNX1 expression associated with t (7; 12)(q36; p13) pediatric acute myeloid leukemia induces the disease through altering histone methylation', *Haematologica,* 109(3), pp. 725.

Weichenhan, D., Riedel, A., Sollier, E., Toprak, U.H., Hey, J., Breuer, K., Wierzbinska, J.A., Touzart, A., Lutsik, P. and Bähr, M. (2023) 'Altered enhancer-promoter interaction leads to MNX1 expression in pediatric acute myeloid leukemia with t (7; 12)(q36; p13)', *bioRxiv,* , pp. 2023.09. 13.557546.

Wildenhain, S., Ingenhag, D., Ruckert, C., Degistirici, O., Dugas, M., Meisel, R., Hauer, J. and Borkhardt, A. (2012) 'Homeobox protein HB9 binds to the prostaglandin E receptor 2 promoter and inhibits intracellular cAMP mobilization in leukemic cells', *The Journal of biological chemistry,* 287(48), pp. 40703-40712 Available at: 10.1074/jbc.M111.308965 [doi].

Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J., Evangelista, J.E., Jenkins, S.L., Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E. and Jagodnik, K.M. (2021) 'Gene set knowledge discovery with Enrichr', *Current protocols,* 1(3), pp. e90.

Yokomizo, T., Ideue, T., Morino-Koga, S., Tham, C.Y., Sato, T., Takeda, N., Kubota, Y., Kurokawa, M., Komatsu, N. and Ogawa, M. (2022) 'Independent origins of fetal liver haematopoietic stem and progenitor cells', *Nature,* 609(7928), pp. 779-784.

Yoshimoto, M., Kosters, A., Cornelius, S., Valiente, N., Cheng, H., Latorre, A., Nishida, C., Ghosn, E.E. and Kobayashi, M. (2022) 'Mast cell repopulating ability is lost during the transition from pre-HSC to FL HSC', *Frontiers in Immunology,* 13, pp. 896396.

1351   Zovein, A.C., Hofmann, J.J., Lynch, M., French, W.J., Turlo, K.A., Yang, Y., Becker, M.S.,
1352   Zanetta, L., Dejana, E. and Gasson, J.C. (2008) 'Fate tracing reveals the endothelial origin
1353   of hematopoietic stem cells', *Cell stem cell,* 3(6), pp. 625-636.

1354