Dissecting infant leukemia developmental origins with a hemogenic gastruloid

model

Denise Ragusa[1,2]*, Chun-Wai Suen[3]*, Gabriel Torregrosa-Cortés[4]*, Fabio Pastorino[5]*,

Ayona Johns[1,2], Ylenia Cicirò[1,2], Liza Dijkhuis[1,12], Susanne van den Brink[6,7], Michele

Cilli[8], Connor Byrne[1,13], Giulia-Andreea Ionescu[1,14], Joana Cerveira[9], Kamil R.

Kranc[10,11], Victor Hernandez-Hernandez[1,2], Mirco Ponzoni[5], Anna Bigas[6,7], Jordi

Garcia-Ojalvo[4], Alfonso Martinez Arias[4], Cristina Pina[1,2,15]¶

[1]College of Health, Medicine and Life Sciences, [2]Centre for Genome Engineering and Maintenance,

Brunel University London, UK; [3]Department of Genetics, University of Cambridge, UK; [4]Department of

Medicine and Life Sciences, Universitat Pompeu Fabra, Spain; [5]Laboratory of Experimental Therapies

in Oncology, IRCCS Istituto Giannina Gaslini, Genova, Italy; [6]Program in Cancer Research, Hospital de

Mar Research Institute, CIBERONC, Barcelona, Spain; [7]Josep Carreras Leukemia Research Institute,

Barcelona, Spain; [8]Animal Facility, IRCCS Policlinico San Martino, Genova, Italy; [9]Department of

Pathology, University of Cambridge, UK; [10]Centre for Haemato-Oncology, Barts Cancer Institute, Queen

Mary University of London, UK; [11]Centre for In Vivo Modelling, Institute of Cancer Research, Sutton,

UK.

[12]Current address: Sanquin Research, Landsteiner Laboratory, Amsterdam University Medical Center,

Amsterdam, The Netherlands

[13]Current address: Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

[14]Current address: Division of Infection and Immunity, Cardiff University School of Medicine, Cardiff,

Henry Wellcome building, CF14 4XN, UK

[15]Current address: Sanquin Research, Amsterdam, The Netherlands

*These authors contributed equally to this work

¶Lead contact. Correspondence: c.pina@sanquin.nl. Sanquin Research, Plesmanlaan 125, 1066 CX,

Amsterdam, The Netherlands.

ABSTRACT

Current in vitro models of developmental blood formation lack spatio-temporal
accuracy and weakly replicate successive waves of hematopoiesis. Herein, we
describe a mouse embryonic stem cell (SC)-derived 3D hemogenic gastruloid
(haemGx) that captures multi-wave blood formation, progenitor specification from
hemogenic endothelium (HE), and generates hematopoietic progenitors capable of
short-term engraftment of immunodeficient mice upon maturation in an in vivo niche.
We took advantage of the haemGx model to interrogate the origins of infant acute
myeloid leukemia (infAML). We focused on MNX1-driven leukemia, representing the
commonest genetic abnormality unique to the infant group. Enforced MNX1
expression in haemGx promotes the expansion and in vitro transformation of yolk sac-
like erythroid-myeloid progenitors at the HE-to-hematopoietic transition to faithfully
recapitulate patient transcriptional signatures. By combining phenotypic, functional
and transcriptional profiling, including at the single-cell level, we establish the haemGx
as a useful new model for the study of normal and leukemic embryonic hematopoiesis.

INTRODUCTION

Blood formation in the embryo is a multi-stage process that develops across multiple cellular niches, reflecting a complex hierarchy of tissue interactions. It involves sequential production of distinct cell types in so-called hematopoietic waves, separated in time and space (Lacaud and Kouskoff, 2017; Costa, Kouskoff and Lacaud, 2012; Medvinsky, Rybtsov and Taoudi, 2011; Dzierzak and Bigas, 2018). The first wave of hematopoiesis produces unipotent red blood cell and macrophage precursors in the yolk sac (YS) from an angioblast precursor which can also form endothelium; it occurs at mouse embryo day (E)7-E7.5 (Fujimoto *et al.,* 2001; McGrath and Palis, 2005; Moore and Metcalf, 1970). The subsequent wave, known as definitive, produces: first, oligopotent erythro-myeloid progenitors (EMPs) in the YS (E8-E8.5); and later myelo-lymphoid progenitors (MLPs – E9.5-E10), multipotent progenitors (MPPs – E10-E11.5), and hematopoietic stem cells (HSCs – E10.5-E11.5), in the aorta-gonad-mesonephros (AGM) region of the embryo proper (Ivanovs *et al.,* 2011; McGrath *et al.,* 2015; Palis *et al.,* 1999; Medvinsky and Dzierzak, 1996; Medvinsky, Rybtsov and Taoudi, 2011). Definitive blood waves are characteristically specified from a specialised hemogenic endothelium (HE) through a process of endothelial-to-hematopoietic transition (EHT) (Zovein *et al.,* 2008; Li, W. *et al.,* 2005). HSC specification from HE is the last event in hematopoietic development and occurs in characteristic cell clusters found on the ventral wall of the dorsal aorta (Ivanovs *et al.,* 2011; Boisset *et al.,* 2010). All blood cell types enter circulation from their original locations and migrate to the fetal liver (FL) (Ciriza *et al.,* 2013; Ghiaur *et al.,* 2008). Most embryonic blood progenitors are transient and sustain blood production exclusively during embryonic development, through differentiation. In contrast, HSCs make little or no contribution to embryonic and fetal blood: instead, they proliferate (i.e. self-renew) to expand their numbers. At least a subset of HSC migrates to the bone marrow (BM) niche at the end of gestation and balances self-renewal and differentiation to maintain blood production throughout the entire post-natal life (Bowie *et al.,* 2006; Kikuchi and Kondo, 2006; Ganuza *et al.,* 2022; Yokomizo *et al.,* 2022).

Blood progenitors and HSCs can be targeted by genetic alterations that confer clonal advantage and/or drive initiation and development of leukemia. Leukemia is the most common malignancy and the most common cause of cancer-related death in the paediatric age group (Steliarova-Foucher *et al.,* 2017; Dong *et al.,* 2020). Several

paediatric leukemias originate *in utero*, and a subset of these are driven by mutations

exclusive to the paediatric group (Masetti *et al.,* 2015; Cazzola *et al.,* 2021). Such

paediatric-exclusive mutations target blood cell types restricted to development or

depend on signals that are specific to embryonic hematopoietic niches, effectively

defining distinct biological entities to adult malignancies (Cazzola *et al.,* 2021; Bolouri

et al., 2018; Chaudhury *et al.,* 2018). Dissection of these forms of developmental

leukemia is confounded by the inability to robustly identify and isolate their embryonic

cells of origin, and/or to fully recapitulate the disease in post-natal cells, which leads

to failure in generating relevant targeted therapies (Alexander and Mullighan, 2021).

One such example is t(7;12)(q36;p13) acute myeloid leukemia (AML) driven by

rearrangement and overexpression of the *MNX1* gene locus: *MNX1*-rearranged

(MNX1-*r*) AML is restricted to infants under the age of 2 and carries a high risk of

relapse and poor survival (Ragusa *et al.,* 2023). *MNX1* is a developmental gene

required for specification of motor neurons and pancreatic acinar cells (Thaler *et al.,*

1999; Harrison *et al.,* 1999). MNX1 is not known to participate in embryonic

hematopoiesis, however it has been discordantly described to be expressed in some

subsets of adult hematopoietic cells (Deguchi and Kehrl, 1991; Nagel *et al.,* 2005;

Wildenhain *et al.,* 2012; Taketani *et al.,* 2008). It is ectopically expressed in AML cells

carrying one of a variable set of t(7;12) translocations which place the normally silent

MNX1 gene under the control of *ETV6* regulatory elements (Weichenhan *et al.,* 2023;

Bousquets-Muñoz *et al.,* 2024; Ragusa *et al.,* 2023; Ballabio *et al.,* 2009). *ETV6* is

required for hematopoietic cell specification from HE via VEGF signalling (Ciau-Uitz *et*

al., 2010), potentially positioning *MNX1* within a critical embryonic hemogenic

regulatory network.

Faithful *in vitro* recapitulation of developmental blood cell types and their embryonic

niches has the potential to dissect the cellular and molecular mechanisms presiding

at initiation of developmental leukemia, reveal prognostic biomarkers, and indicate

therapeutic vulnerabilities which are otherwise challenging to test in the embryo.

Despite widespread use of embryonic stem (ES) cell and induced pluripotent stem

(iPS) cell-based models of blood specification, a system which recapitulates the spatial

and temporal complexity of embryonic blood formation within the respective time-

dependent niches is still lacking (Dijkhuis *et al.,* 2023). In the last few years, gastruloid

models have emerged as robust avatars of early development, self-organising

processes of symmetry breaking, elongation, multi-axis formation, somitogenesis and early organogenesis, with remarkable similarity to embryo processes in space and time (van den Brink, Susanne C and van Oudenaarden, 2021; Beccari *et al.,* 2018; van den Brink, Susanne C *et al.,* 2020). Gastruloids are grown individually in a multi-well format that facilitates tracking and screening of developmental processes. Relevant to blood formation, a variant gastruloid culture system, which successful organises a heart primordium (Rossi *et al.,* 2021), was also able to recapitulate YS blood formation with erythroid and myeloid cellularity (Rossi *et al.,* 2021; Rossi *et al.,* 2022).

Herein, we describe a new protocol of gastruloid formation, i.e. hemogenic gastruloids (haemGx), that captures putative YS-like and AGM-like waves of definitive blood progenitor specification from HE. The earlier wave of putative YS-like hematopoiesis includes EMP-like cells which further mature in the gastruloid culture. HaemGx AGM-like hematopoiesis captures the emergence of intra-aortic cluster-like cells, including cells capable of short-term multi-lineage hematopoietic engraftment of the spleen and BM of immunodeficient mice. We harnessed the embryonic hematopoietic context provided by haemGx to attest its utility in modelling leukemias with developmental origins, specifically by recapitulating MNX1-*r* infant leukemia by proxy of *MNX1* overexpression. MNX1-overexpressing haemGx showed expansion of cells at the HE-to-EMP transition, which became susceptible to transformation *in vitro* and recapitulated MNX1-*r* AML patient signatures, placing *MNX1*'s leukemogenic effects at a very early stage of developmental blood formation.

RESULTS

HaemGxs capture time-dependent emergence of endothelium and hematopoietic cells

With the aim of recapitulating embryonic hematopoiesis with spatial and temporal accuracy, we modified existing gastruloid formation protocols to (1) promote the specification of hemogenic mesoderm derivatives, and (2) extend developmental time beyond the E8.0-equivalent of early developmental gastruloids (Van den Brink, Susanne C *et al.,* 2014) and the E9.0 timepoint of cardiogenic gastruloids (Rossi *et al.,* 2022). We established a 216h haemGx protocol (**Fig. 1A**) using a *Flk1-GFP* mouse ES cell line (Jakobsson *et al.,* 2010), which allows the tracking of early hemato-

endothelial specification by expression of the *Flk1* (i.e. *Kdr*) *locus*. The haemGx
protocol includes an indispensable pulse of Activin A to induce hemato-endothelial
programmes via TGF-β signalling (visible by proxy of GFP-tagged *Flk1* activation), in
addition of the characteristic gastruloid patterning via WNT activation by CHI99021
supplementation at 48h (**Fig. 1S1A-B**). We promoted hemato-endothelial programs
through addition of VEGF and FGF2 at 72h (Sroczynska *et al.,* 2009) (**Fig. 1A**). We
observed extension of a polarised *Flk1-GFP*-expressing branched endothelial network
(**Fig. 1B**). Flow cytometry analysis revealed the onset of VE-cadherin$^+$ C-Kit$^+$ cells
suggestive of EHT, with a peak at 120-144h (**Fig. 1C-D**). This was followed by a surge
in CD41$^+$ candidate hematopoietic cells at 144h (**Fig. 1C-D**). In our experience,
observation of early polarised activation of the *Flk1 locus* at 96h (**Fig. 1B**) was
predictive of CD41$^+$ detection 2 days later, suggesting that TGF-β activation in early
patterning is required for later hematopoietic development, putatively by modifying
mesoderm specification. We incorporated a 24-hour pulse of sonic hedgehog (Shh)
between 144-168h to mimic the aortic patterning that precedes cluster formation (**Fig.**
1A) (Rybtsov *et al.,* 2014). Continued exposure to VEGF after the Shh pulse resulted
in specification of CD45$^+$ hematopoietic cells progenitors (**Fig. 1C-E**), which were
apparent after 168h. The proportion of CD45$^+$ cells was enhanced by the addition of
SCF, FLT3L and TPO (**Fig. 1S1C-D**), a combination of cytokines routinely used for
HSC maintenance; conversely, it did not require continued addition of FGF2 after 168h
(**Fig. 1S1E**), which was omitted. CD45$^+$ cells, some of which co-expressed C-Kit, were
present in a small number of 'cluster'-like structures adjacent to Flk1-GFP$^+$ Kit$^+$
endothelium (**Fig. 1E**). The process is overall reminiscent of hematopoietic progenitor
cell budding from the aortic HE, suggesting recapitulation of hemogenic tissue
architecture. Time-dependent specification of C-Kit$^+$, CD41$^+$ and CD45$^+$ cells in
haemGx could be obtained with other mouse ES cell lines, including commonly used
E14Tg2a (E14) cells (**Fig. 1S1F-H**), with successful specification of comparable
frequencies of CD45$^+$ progenitors (**Fig. 1S1I**), attesting to the reproducibility of the
protocol across cell lines of different genetic backgrounds.

Overall, our extended haemGx model could capture time-dependent specification of
CD41$^+$ and CD45$^+$ hematopoietic cells, with recapitulation of the cluster-like spatial
arrangement of CD45$^+$ cell emergence from HE, thus showing initial promise as a tool
for the exploration of normal and perturbed developmental hematopoiesis.

HaemGx specify cells with phenotypic and functional hematopoietic progenitor characteristics

We confirmed the induction of endothelial patterning from the Flk1-GFP population by immunofluorescence staining showing the branched vascular-like network of Flk1 co-expressing CD31 (**Fig. 2A**). Endothelial identity of Flk1-GFP$^+$ populations was also detectable by flow cytometry through the presence of endothelial markers CD31 and CD34 (**Fig. 2S1A-B**). Given the co-expression of CD31 and CD34 with hematopoietic progenitors emerging from HE, we inspected CD41$^+$ and CD45$^+$ cells for the presence of these markers. Indeed, we found that subsets of CD41$^+$ and CD45$^+$ hematopoietic cells co-express CD31 and CD34 (**Fig. 2B, Fig. 2S1C-D**), with increased relative representation of hemato-endothelial markers within CD41$^+$ cells at 144h and CD45$^+$ at 216h (**Fig. 2C**), consistent with two successive waves of hemogenic emergence from HE. CD45$^+$ cells from 216h haemGx also co-expressed C-Kit, Flk1-GFP and VE-cadherin, a phenotype enriched in emergent hematopoietic stem and progenitor cells (HSPC) in the AGM (**Fig. 2S1E-F**). The phenotype is detected in 1.8% of CD45$^+$ cells on average (**Fig. 2S1G**), corresponding to an estimated 8 cells per haemGx. In agreement, analysis of the haemGx progenitor content at 216h using multipotential CFC assays (**Fig. 2D-E**) revealed the presence of >50 CFC/haemGx, of which an average of 7 have mixed-lineage granulocytic-monocytic-erythroid (GEM) potential, numerically matching the estimated number of HSPC. Other progenitors were predominantly granulo-monocytic (GM) and erythroid (E) progenitors (**Fig. 2D-E**). Analysis of the cellularity of dissociated haemGx at the 216h timepoint using Giemsa-Wright stained cytospins revealed differentiated cells of G, M, E and megakaryocytic lineages (**Fig. 2S2A**), putatively reflecting the progeny of EMP-like cells.

We observed temporal progression of expression of key markers by real-time quantitative PCR (qPCR), configuring a sustained hemato-endothelial program from 120h through 216h (*Dll4* and *Mecom*), and the generation of hematopoietic populations based on the time-dependent patterns of *Myb, Runx1*, *Hlf* and *Hoxa9* (**Fig. 2F**). We note that *Myb* peaks at 144h, coincident with the emergence of CD41$^+$ cells,

which we postulate to have YS-like EMP characteristics. On the other hand, *Runx1* has a bimodal pattern peaking at 144h, and later at 216h, suggesting the presence of two distinct hematopoietic waves. The later timepoint of 216h has significantly higher expression of both *Hoxa9* and *Hlf* raising the possibility that the second wave indeed corresponds to the specification of AGM-like HSPC. In order to further test the capture of YS-like and AGM-like hematopoietic waves, given the lack of markers that specifically associate with AGM hematopoiesis, we took advantage of the distinct regulatory requirements of polycomb-associated EZH2, which is necessary for EMP differentiation in the YS, but not for AGM-derived hematopoiesis or primitive erythroid progenitors (Neo *et al.*, 2018; Neo *et al.*, 2021). Upon EZH2 inhibition via treatment with GSK216 from 120h onwards, we observed a reduction of CD41[+] cells selectively at 144h, with no change in CD45[+] cells at the later timepoint of 216h (**Fig. 2S2B-C**). This suggests that the cells present at the later timepoint may not be the progeny of the former, thus supporting recapitulation of 2 independent hematopoietic waves, the latter putatively AGM. Additionally, the vulnerability of 144h CD41[+] to EZH2 inhibition suggests that they include YS-like EMP and do not predominantly reflect EZH2-independent primitive YS hematopoiesis.

HaemGx progenitors capture time-dependent signatures of YS-like and AGM-like hematopoiesis

To better characterize the extent and progression of developmental hematopoiesis in haemGx, we performed single-cell RNA-sequencing (scRNA-seq) time-course analysis of haemGx cell specification. We sorted cells from 2 independent haemGx cultures at 120, 144, 168, 192 and 216h, and profiled a total of 846 cells using the Smart-Seq2 protocol (Picelli *et al.,* 2014) (**Fig. 3S1A**). We analyzed unfractionated live single haemGx cells at 120-192h and enriched the dataset with sorted CD41[+] cells at their peak time of 144h, and sorted CD45[+] cells at 192 and 216h. At timepoints of significant enrichment in CD41[+] (144h) and CD45[+] cells (192h) (**Fig. 1D**), we also sorted C-Kit[+] cells. Library preparation and sequencing generated an average of 120000 reads/cell, which were mapped to an average of 4000 genes/cell, with minimal signs of cell stress / death as measured by mitochondrial DNA fraction (**Fig. 3S1B**).

Read and gene counts were similar between biological replicates, cell types and at
different timepoints (**Fig. 3S1C**). The exception was 120h unfractionated haemGx
cells, which despite similar sequencing depth (average read counts), were mapped to
twice the number of genes (**Fig. 3S1C**), possibly reflecting multi-gene program priming
at the onset of hemogenic specification. We selected highly varying genes (HVG)
before principal component analysis (PCA) dimensionality reduction and retained the
most relevant dimensions. In the PCA reduced space, we constructed a KNN graph
and used uniform manifold approximation and projection (UMAP) to visualize the data
on 2 dimensions, looking for cell communities using Leiden clustering (**Fig. 3S1D**). We
identified 12 cell clusters of which 2 (clusters 0 and 5) almost exactly mapped to C-
Kit$^+$ cells (also positive for Sca-1), 2 (clusters 1 and 8) contained CD45$^+$ sorted cells,
and cluster 4 uniquely captured the CD41$^+$ cells observed at 144h (**Fig. 3S1D**).
Although some unfractionated cells overlapped with the hemogenic cell clusters (**Fig.**
3S1D, middle panel), most occupied different transcriptional spaces, reflecting the
relative frequency of hematopoietic cells and suggesting the presence of potential
hemogenic niches.

To explore tissue and lineage affiliation of the different clusters, we performed
differential gene expression against all other cells using Wilcoxon ranking test, and
established cluster classifier gene lists (**Supplemental File S1**). Top classifier genes
for clusters 0 and 5 had endothelial affiliation (**Supplemental File S1-S2**), and the
cellular composition of the clusters broadly reflected their time signature (**Fig. 3S1D**):
cluster 5 comprised 144h cells and cluster 0 included most C-Kit$^+$ cells at 192h. Cluster
5 occupied the vicinity of 144h CD41$^+$ cells (**Fig. 3S1D**), which carried an erythroid-
biased signature (**Supplemental File S2**). On the other hand, 192/216h-CD45$^+$ cells
in cluster 8 expressed myeloid and lymphoid genes (**Supplemental File S1-S2**). The
remaining clusters largely corresponded to time-restricted unfractionated cells (**Fig.**
3S1D, bottom panel), with clusters of cells present at later timepoints matching
mesenchymal stromal cell and autonomic neuron signatures (**Fig. 3S1E**), potentially
configuring incipient formation of niches relevant to production of HSC (Kapeni *et al.,*
2022; Fitch *et al.,* 2012).

To specifically explore the extent to which haemGx capture putative YS and AGM-like
hematopoiesis, we re-clustered cells sorted on hemato-endothelial markers CD41, C-
Kit, and CD45 at the 144h, 192h and 216h timepoints (**Fig. 3A**). We mapped the

expression pattern of endothelial (*Pecam* and *Cd34*), HE (*Mecom* and *Cd44*) and

hematopoietic (*Runx1*, *Mllt3*, *Hoxa9*, and *Hlf*) markers across the time-dependent

clusters (**Fig. 3B**). We observed a coincidence of *Pecam*, *Cd34*, and *Mecom*

expression with clusters 0 and 3, which capture C-Kit$^+$ cells at 144 and 192h,

respectively. Some positive cells were also seen within CD41$^+$ and CD45$^+$-enriched

clusters 2, and 1 and 4, respectively, supporting the presence of phenotypic

progenitors, as suggested by flow cytometry analysis (see **Fig. 2**). CD41$^+$ and CD45$^+$-

sorted clusters, on the other hand, show expression of *Runx1*, confirming their

hematopoietic affiliation. *Cd44*, which is upregulated at the endothelial-to-

hematopoietic transition (EHT) and in association with the AGM region (Oatley *et al.*,

2020; Fadlullah *et al.*, 2022), is detected predominantly in C-Kit$^+$-enriched cluster 3

and CD45$^+$-sorted clusters 1 and 4, corresponding to the later timepoints of 192 and

216h. This supports the notion that the later timepoints may capture AGM-like EHT,

which is also suggested by significant increases in expression of *VE-cadherin* (*Cdh5*),

Sox17, as well as *Cd34*, *Cd44* and *Procr* (Fadlullah *et al.*, 2022; Jung *et al.*, 2019;

Oatley *et al.*, 2020) in 192h, endothelial-enriched, C-Kit$^+$ cells (**Fig. 3C-D**). Important

in this context, these cells express arterial endothelial markers, including *Flt1*, *Efnb2*

and *Dll4* steadily at both timepoints (**Fig. 3C and 3S2B**). *Hoxa9* expression also

associates with arterial specification and definitive AGM-like hematopoiesis: albeit at

low frequency, it is detected exclusively in C-Kit and CD45-sorted cells in the 192 and

216h-timepoints of haemGx differentiation (**Fig. 3B, E**). These cells also express *Mllt3*

and, rarely, *Hlf*, compatible with incipient specification of an AGM-like definitive blood

program (**Fig. 3B, E**). *Mllt3* is additionally expressed in CD41$^+$-sorted cells of cluster

2, in line with its role in early erythroid and megakaryocytic lineage identity (Pina *et al.*,

2008); *Myb* is also more frequently expressed in this population (**Fig. 3E**). In contrast,

Ikzf1, which regulates lymphoid specification (Huang *et al.*, 2019), gradually increases

its expression, and indeed frequency, in 216h-CD45$^+$ sorted cells. A handover from

erythroid/megakaryocytic to myelo-lymphoid lineage affiliation is apparent between

144h and 216h haemGx blood cells (**Fig. 3S2C**). Of note, a small number of erythroid-

affiliated cells at 216h express adult globins, suggesting a definitive origin (**Fig. 3S2C**).

Altogether, scRNA-seq analysis support the suggestion from phenotypic and *in vitro*

functional data, that the haemGx model captures two waves of hematopoietic

specification from HE-like cells and may configure pre-definitive and definitive
embryonic blood formation.

308
309

310
311
312
313

Late-stage haemGx cells transcriptionally resemble mouse AGM progenitors
and have short-term engraftment potential

To investigate the putative YS-like pre-definitve and AGM-like definitive nature of
haemGx-generated hematopoietic progenitors, we conducted projections of haemGx
sc-RNAseq data onto *bona fide* embryonic populations. We considered 3 distinct
scRNA-seq datasets, which (1) capture arterial and haemogenic specification in the
para-splanchnopleura (pSP) and AGM region between E8.0 and E11 (Hou *et al.* 2020)
(**Fig. 4S1A**); (2) uniquely capture YS, AGM and FL progenitors and the AGM EHT (Zhu
et al., 2020) (**Fig. 4S1B**); and (3) interrogate the mouse AGM at the point of HSC
emergence from the dorsal aorta HE (Thambyrajah *et al.,* 2024) (**Fig. 4S1C**). We
extracted highly-varying genes from each of the 3 datasets to construct dimensionality
reduction maps of the data (**Fig. 4S1A-C**), and conducted k-Nearest Neighbor (KNN)
analysis to classify and regress our haemGx scRNA-seq data onto those maps
considering 3 neighbors (**Fig. 4A**). HaemGx cells with correlation metric >0.7 were
projected onto the respective embryonic datasets (**Supplemental File S3)**.
Projections onto Hou *et al.* (2020) positioned clusters 0 and 3, corresponding to
endothelium, including arterial endothelium and HE-like cells, between days E8.5 and
E10, putatively spanning YS-like and AGM-like blood formation. A subset of CD45$^+$-
sorted cells, as well as a small numbers of CD41$^+$ cells from cluster 2 projected onto
E10 HE and intra-aortic cluster (IAC) cells, suggesting the presence of definitive HSPC
or their precursors in haemGx (**Fig. 4A**, **Supplemental File S3**). We sought to further
check the transcriptional affiliation of haemGx with intra-embryonic, definitive
hematopoiesis, by considering the dataset by Zhu *et al.* (2020), which uniquely
captures different hematopoietic locations in the same study (**Fig. 4A**). We matched

the identity of a large proportion of CD41$^+$-sorted cells to YS EMP, with some of the same cluster 2 cells mapping to erythroid progenitors, and some later CD45$^+$-sorted cells resembling myeloid progenitors (**Fig. 4A**, **Supplemental File S3**). On the other hand, a subset of CD45$^+$-sorted cells mapped to IAC and lymphoid-myeloid primed progenitors (LMPP) in the FL (**Fig. 4A**, **Supplemental File S3**), compatible with the presence of AGM-like definitive HSPC in haemGx. Interestingly, some CD41$^+$-sorted cells also map to IAC, and indeed to HE, suggesting that this earlier population may contain HSPC precursors, in addition to YS-like progenitors. IAC and HE-like cells were also found within C-Kit$^+$ cells, mostly from the 192h timepoint, while the majority of C-Kit$^+$ cells captured different endothelial identities, as expected from their transcriptional profiles. Projection onto Thambyrajah *et al.* (2024) (**Fig. 4A**, **Supplemental File S3**), conflates YS-like and AGM-like progenitors onto the candidate HSC-enriched embryo AGM cluster 2, highlighting the similarity between transcriptional programs at different sites of embryonic blood emergence. CD45$^+$ cells are more distally-projected onto the same cluster, as shown for E11.5 cells in the Thambyrajah *et al.* (2024) study, some of which correspond to the candidate FL-LMPP in Zhu *et al.* (2020) (**Supplemental File S3**). HaemGx cell similarities to Thambyrajah *et al.* (2024) HSC-enriched cluster 2 also include HE-like and IAC-like cells (as per Zhu *et al.* 2020 projections), mostly E9.5-like as per Hou *et al.* (2020), overall suggesting putative early specification of the AGM region within haemGx.

Late-stage haemGx cells can engraft hematopoietic tissues upon maturation

The presence of putative AGM-like hematopoietic cells within haemGx led us to investigate whether haemGx-derived cells had the ability to engraft the hematopoietic system of immunodeficient animals. In early experiments, we had transplanted dissociated haemGx cells obtained at 192 and 216h into irradiated immunocompetent C57Bl/6 recipients, sub-lethally irradiated immunodeficient NSG mice, or non-conditioned NSGW41 hosts, but failed to detect signs of engraftment of peripheral blood, spleen (Spl) or bone marrow (BM) at 12 days (including absence of spleen colony formation), at 4-6 weeks (short-term engraftment), or beyond 8-10 weeks (long-term engraftment) by either flow cytometry detection of CD45.2 or by genomic DNA (gDNA) PCR analysis of the *Flk1-GFP* locus.

Failure to engraft was perhaps not surprising given that transcriptional programs
position haemGx no later than E9.5-E10, which precedes HSC specification in the
AGM. Additionally, specification of HSC-supporting stroma within haemGx, namely
mesenchymal cells (Kapeni *et al.*, 2022) and elements of the sympathetic nervous
system (Fitch *et al.*, 2012), is only detectable at the 192h timepoint, suggesting that
further 3D maturation *in vivo* may be required. We thus tested implantation of
undissociated haemGx in the adrenal gland of *Nude* mice (**Fig. 4B**), a topography
more commonly used to support tumor development, but that was recently shown to
be able to support HSC development, possibly with contributions from the sympathetic
adrenal medullary niche (Schyrr *et al.,* 2023). In order to facilitate the analysis of
engrafted animals, and given the coincidence of CD45 isoform between *Flk1-GFP*
mouse ES cells and the *Nude* mice in which the experiments were performed, we
engineered constitutive expression of BFP from the *Rosa26* locus in the *Flk1-GFP*
mES cell line (*Rosa26-BFP::Flk1-GFP*) (**Fig. 4S2A**). We implanted 3 haemGx each
unilaterally in the adrenal gland of unconditioned recipient mice (**Fig. 4B**). Control mice
were injected with an equivalent volume of PBS. Animals were sacrificed at 4 or 8
weeks after implantation, with 4-5 experimental animals and 1 control collected at each
timepoint. We checked Spl and BM BFP engraftment, by flow cytometry and by PCR
analysis of DNA. Sensitivity of PCR analysis was determined by serial dilution of
Rosa26-BFP::Flk1-GFP mouse ES cells in the human K562 cell line, with detection of
as low as 10 pg of mouse input in a total of 100 ng gDNA (**Fig. 4S2B**), corresponding
to approximately 3 cells (in 300,000). PCR analysis of BFP engraftment of the Spl and
BM 4 weeks after adrenal implantation of 3 haemGx / animal detected *BFP* amplicon
in both tissues in 3 of the 5 samples (#11, 12 and 15) (**Fig. 4C and 4S2C**). We sampled
the gDNA of each recipient 5 times and detected the presence of the amplicon in 3-5
of the replicate samples of the positive animals; replicate sampling of the control was
consistently negative (**Fig. 4C and 4S2C**). The same 3 animals showed BFP detection
by flow cytometry above control, but no higher than 0.1% (**Fig. 4D**). PCR engraftment
was nevertheless detectable in both erythroid Ter119[+] and lympho-myeloid CD45[+]
fractions of the recipient animals showed PCR engraftment in both lineage fractions
(**Fig. 4S2D-E**). Analysis of engraftment 8 weeks after implantation showed low-level
detection (<0.1%) of BFP[+] cells with positivity for CD11b[+] myeloid and B220[+] lymphoid
compartments in BM of 1 recipient (**Fig. 4D and 4S2F-G**); a 4[th] recipient had to be
sacrificed early due to infection.

Overall, implantation in the adrenal niche and/or preservation of the 3D architecture
enabled low-level *in vivo* hematopoietic activity, although the mechanism remains
unclear. In some recipients, hematopoietic engraftment was accompanied by *in situ*
growth and multi-tissue differentiation of the implanted haemGx (**Fig. 4S2H**). However,
adrenal enlargement was not observed in all animals with hematopoietic engraftment
(e.g. 4 weeks' #11 had a normally sized adrenal, as did all animals analysed at 8
weeks), making it unlikely that hematopoietic activity results from teratoma formation
by residual pluripotent cells. Late-stage haemGx (216h) may thus contain
hematopoietic progenitors with some short-term engraftment potential but incomplete
functional maturation, which are present at low frequency. Engraftment is erythro-
myeloid at 4 weeks and lympho-myeloid at 8 weeks, reflecting different classes of
progenitors, putatively of YS-like and AGM-like affiliation.

HaemGx resolve the early hemogenic ontogeny targeted by infant AML gene
MNX1

The ability of haemGx to reconstitute putative YS and AGM-like HE and hemopoietic
progenitors positions it as an attractive model to probe cellular origins of
developmental leukemia. In particular, forms of AML unique to the first 2 years of life
– infant (inf)AML – have been shown to transform fetal liver (FL) but not adult
hematopoietic cells (Waraky *et al.,* 2024; Mercher *et al.,* 2009; Chen, W. *et al.,* 2011;
Ragusa *et al.,* 2023), compatible with targeting of a transient embryonic cell. In the
case of t(7;12) AML (herein MNX1-*r*) (Ragusa *et al.,* 2023), in which the translocation
results in transformation-driving ectopic expression of *MNX1* (Waraky *et al.,* 2024),
analysis of patient transcriptional signatures (Ragusa *et al.,* 2022) for over-
representation of cell atlas-defined affiliations, indicates enrichment in
cardiomyogenic, endothelial, HSC/progenitor and mast cells (**Fig. 5A**). These
signature enrichments are compatible with an early hemogenic cell derivative,
capturable in the haemGx model. Other cell type-enrichments reflect MNX1 functions
in pancreatic and neuronal development (Ragusa *et al.,* 2022) (**Fig. 5S1A**).
We overexpressed *MNX1* (MNX1-OE) in *Flk1-GFP* mouse ES cells by lentiviral
transduction (**Fig. 5B**) and analyzed progression of haemGx generation (**Fig. 5C**). We
used human *MNX1* cDNA to distinguish from the endogenous gene, but the degree of

homology is nevertheless high (84%), supporting functional equivalence. We
confirmed that *MNX1* overexpression in haemGx is maintained to endpoint (**Fig.
5S1B**). MNX1-OE haemGx activated polarized *Flk1-GFP* expression and elongated
with similar kinetics to empty-vector (EV) control (**Fig. 5C**), albeit with formation of
larger haemGx (**Fig. 5D**) denoting increased cellularity (**Fig. 5S1C**). From 192h
onwards, MNX1-OE haemGx had a higher frequency of spontaneously contractile
structures (**Fig. 5S1D**), compatible with the cardiogenic cell association of patients'
transcriptomes (**Fig. 5A**).

We interrogated the time-dependent hemogenic cell composition of haemGx by flow
cytometry, quantifying C-Kit, CD41 and CD45 (**Fig. 5E-G**, **5S1E-F**), through which we
had captured endothelial/HE-like cells, YS-EMP-like, and later putative AGM-like
progenitors, respectively. We observed a small but significant expansion of the C-Kit$^+$
compartment specifically at 144h (**Fig. 5E**, **5S1E**). All cell markers were relatively
unchanged at later timepoints, although absolute cell numbers were higher in MNX1-
OE haemGx (**Fig. 5S1C**).

To better understand the consequences of MNX1-OE on hemogenic development, we
performed RNA-sequencing (RNA-seq) of haemGx at 144h and 216h, capturing
putative YS and AGM-like hemopoietic waves, respectively (**Fig. 5H**). We confirmed
expression of the human *MNX1* transgene in the sequenced reads (**Fig. 5S2A**); in
contrast, we could not detect endogenous mouse *Mnx1* in any of the samples (**Fig. 5
S2A**), confirming that it does not normally play a role in hemogenic development.
DEGs between MNX1-OE and EV (**Supplemental File S4**) configured distinct MNX1-
driven programs at 144h and 216h, with >80% genes timepoint-specific (**Fig. 5H**). Both
programs include up-regulation of hematopoietic-associated genes, capturing EHT
and progenitor regulators (e.g. *Runx1*, and *Epor* and *Zfpm1*, respectively) at the earlier
timepoint, and more differentiated effectors, e.g. *Hba-a1*, *Hbb-bh1* and *Il3ra* at 216h,
compatible with selective targeting and/or expansion of YS EMP-like cells and their
later differentiated progeny (**Fig. 5H**). An overview of transcriptional regulatory
programs at both timepoints denotes a clearer hemogenic enrichment at 144h, with
over-representation of GATA2, GATA1 and RUNX1 binding targets (**Fig. 5I**), putatively
capturing an expansion of a HE-to-EMP-like transition, also represented in enriched
GO categories related to VEGF signalling and epithelial-to-mesenchymal transition
(EMT) (**Fig. 5J**). In contrast, no enrichment of hemogenic transcription factor (TF)-
ChIP targets or related gene ontologies (GO) categories was observed in MNX1-OE

haemGx at 216h (**Fig. 5I-J**), suggesting specific targeting of YS-like hemogenic programs. Instead, there was an enrichment in cardiogenic GO at 216h (**Fig. 5J**), reflecting the increase in contractile foci in MNX1-OE late haemGx.

We attempted to further deconvolve the cellular heterogeneity underlying bulk RNA-seq DEGs by interrogating the cluster signatures obtained from the single-cell analysis of haemGx development (**Fig. 5S2B-C**; refer to **Fig. 3S1D**). In support of an association with hemato-endothelial specification, MNX1-OE transcriptomes showed enrichment of haemGx HE cluster 5 and 0 signatures; the YS-EMP-like cluster 4 was also enriched (**Fig. 5S2B**). MNX1-OE DEGs at both timepoints also enriched for the relatively less differentiated cluster 10, which predominantly captures unfractionated haemGx cells at 120h, supporting an early ontogenic affiliation. Significantly, clusters 5 and 0 HE signatures were also enriched in *MNX1*-r infAML patient signatures (Ragusa *et al.*, 2022) (**Fig. 5S2B-C**), putatively aligning MNX1-OE effects with HE and the hemogenic transition. In contrast, MNX1-OE DEGs specific to 216h showed enrichment in myeloid progenitor / MLP-enriched cluster 8 which captures 192/216h haemGx cells, but diverges from *MNX1*-r infAML signatures (**Fig. 5S2B-C**), instead approximating the distinct myelo-monocytic and/or mixed-lineage affiliation of *KMT2A*-rearranged AML, which is also common in older pediatric and adult patients.

Serial replating of MNX1-OE haemGx cells identifies a candidate C-Kit$^+$ leukemia-propagating cell more susceptible to transformation in the YS period

We further investigated the potential of MNX1-OE haemGx to reflect *MNX1*-r AML biology by performing serial replating of CFC assays, a classical *in vitro* assay of leukemia transformation. We compared 144h and 216h haemGx to match functional and transcriptome data, and plated unfractionated haemGx, given the transient nature of C-Kit$^+$ enrichment. Both 144h (**Fig. 6A-B**) and 216h (**Fig. 6C-D**) MNX1-OE haemGx replated for at least 5 platings, significantly different to control (EV) (**Fig. 6B,D**). EV-haemGx replating could not be sustained beyond plate 3 in most experiments started from 216h (**Fig. 6C**); 144h EV-haemGx exhibited some low-level replating until plate 5 (**Fig. 6A**), but the colonies were formed by few scattered cells clearly distinct from the well-demarcated colonies obtained from MNX1-OE haemGx. We analyzed early-stage (plate 1) and late-stage (plate 5) colonies by flow cytometry to identify the nature

of the replating cells. Cells selected through replating were C-Kit$^+$ (**Fig. 6E**), matching

the cell phenotype transiently enriched at 144h. MNX1-OE C-Kit$^+$ cells co-expressed

CD31 (**Fig. 6S1A-B**), underpinning the endothelial signatures of *MNX1-r* leukemia.

Early replating of MNX1-OE haemGx also enriched for erythroid-affiliated Ter119$^+$

cells, particularly when initiated from the 144h timepoint (**Fig. 6S1C**), but this was not

sustained through replating. Myelo-monocytic-affiliated markers, including CD11b or

CD45, were not present (**Fig. 6E**, **6S1C**).

514
515

We performed RNA-seq analysis of MNX1-OE haemGx CFC cells obtained through

re-plating and integrated the data with the 144h and 216h haemGx timepoints to

identify genes enriched through the process of transformation. Hs*MNX1* reads were

increased in MNX1-OE CFC samples, supporting selective expansion of transduced

cells (**Fig. 7A**). We performed hierarchical clustering of all DEGs in pairwise

comparisons (**Fig. 7B**), and used Gene Set Enrichment Analysis (GSEA) of *MNX1*-r

patient signatures up-regulated in comparison to other AML in the infant (0-2) age

group, to interrogate the different clusters. Cluster k14 had significant enrichment of

MNX1-*r* signatures against all other AML subtypes (**Fig. 7C**), with considerable overlap

between leading-edge (i.e. significantly-enriched) genes (**Fig. 7S1A**). Importantly, k14

captures the subgroup of DEGs up-regulated upon CFC replating which are also up-

regulated in MNX1-OE haemGx at 144h (**Fig. 7B**), matching the time-dependent

enrichment of the putative MNX1-OE propagating population. Interrogation of cell-

type-specific signatures within the k14 MNX1-*r* leading edge genes against the single-

cell PanglaoDB atlas captured similarities to hemogenic and non-hemogenic cell

types, including interneuron, and other neural signatures (**Fig. 7D**), which reflect

MNX1 biological functions (Harrison *et al.,* 1999; Thaler *et al.,* 1999; Ragusa *et al.,*

2022). The most highly significant signatures capture early developmental hemogenic

components, such as 'endothelial cells' and 'erythroid cell precursors', reflecting the

144h, putative YS affiliation of MNX1-OE haemGx. The 'mast cells' affiliation of *MNX1-*

r AML signatures within the haemGx k14 cluster is also likely to reflect a YS EMP-like

affiliation (Chia *et al.,* 2023), and is accordingly aligned with MNX1-OE haemGx

isolated at 144h (**Fig. 7S1B**). Giemsa-Wright-stained cytospins of serially-replated

CFC assays initiated with 144h MNX1-OE cells showed a shift from blast-like at plate

3 to precursor cell morphology with some differentiated mast cells at plate 5 (**Fig. 7E,**

S1C). We also observed mast cells in 216h-initiated colonies (**Fig.7E**)**,** but these were

differentiated at an earlier plate, suggesting that MNX1-OE-responsive C-Kit$^+$

progenitors experience maturation between 144h and 216h.

Taken together, the data positioned the initiating cellular targets of MNX1-*r* AML at the

YS endothelial-to-EMP transition. More broadly, the analysis supports the potential of

haemGx to dissect the ontogeny of specific subtypes of hematological cancers

initiated *in utero*.

DISCUSSION

In this study, we developed and characterized a hemogenic gastruloid (haemGx) model from mouse ES cells, which recapitulates YS-like and AGM-like waves of definitive blood progenitor specification, and has incipient ability to engraft adult hematopoietic tissues short-term upon *in vivo* maturation. By exploring the haemGx model through enforced expression of the *MNX1* oncogene, which is a hallmark of infant AML harbouring the t(7;12) translocation, we identified the YS endothelial-to-hematopoietic transition as the transformation target of *MNX1* gene rearrangement, shedding light on to its exclusive association with the infant age group.

Our data positions the haemGx model as an advanced platform to probe physiological and pathological development of the hematopoietic system, that benefits from the time-dependent and topographic organization associated with gastruloid models. We constructed the haemGx model by introducing a pulse of Activin A and extending the protocol under hematopoiesis-promoting cytokines, to reach the equivalent of mouse embryonic day 10 (E10) and achieve sequential acquisition of YS-like and AGM-like blood production from HE, benchmarked by surface markers CD41 and CD45, respectively. In agreement with recent observations from Dias *et al.* (2025), which highlight a critical balance between WNT-driven posterior and Nodal / TGFβ anterior signals in patterning the midbody axis and endo-mesodermal derivatives, we observed that the combination of Activin A and CHIR99021 was critical for robust *Flk1/Kdr* activation and the efficient initiation of CD41 and CD45-marked hematopoietic specification. This is in contrast with the hematopoietic output of cardiogenic gastruloids reported by Rossi *et al.* (2022), which did not require Activin A. Although our strict dependence on Activin A to achieve initial activation of *Flk1* programs and specification of a vascular network and hematopoietic programs precluded a side-by-side comparison with the Rossi protocol (Rossi *et al.,* 2022), we note their apparent lower efficiency in blood formation, at least at the comparable timepoints of 144 and 168h, and their bias towards erythroid formation, reflecting a putative YS, and potentially primitive, nature. This bias may also be captured in the reported spleen colony formation; however, given the limited characterisation of the colonies in the manuscript, which does not appraise origin, morphology or cellularity, the nature of the spleen engraftment remains speculative. Importantly, standard gastruloid protocols

with only CHIR99021 pulse and no additional cytokines (Beccari *et al.,* 2018) can
capture a transcriptional erythrogenic wave.

Despite low frequency engraftment of adult hematopoietic tissues by haemGx cells,
we demonstrated donor origin of the engrafted cells by flow cytometry and PCR
detection of a BFP transgene up to 8 weeks. Engraftment progressed from erythroid-
biased at 4 weeks to myelo-lymphoid at 8 weeks, as observed in progenitor
engraftment from adult cells (Chun *et al.*, 2024). Critically, successful engraftment
required implantation of undissociated haemGx for further maturation in a supportive
microenvironment, in this case provided by the adrenal gland (Schyrr *et al.,* 2023). In
this study, we did not dissect the relative contribution of the continued maturation in
3D *vs* the influence of the niche for implantation, as we opted for performing the
transplantation as to maximise both. Future experiments will be necessary to
understand possible extrinsic niche contributions, including the sympathetic nervous
system we sought to include (Kapeni *et al.,* 2022; Fitch *et al.,* 2012), and how they can
be incorporated in the haemGx culture by extension and/or further modification of the
protocol. Although we observed continued growth and maturation of the haemGx *in*
situ after implantation, this was not the case in all animals with engraftment. On the
one hand, this dissociation excludes the likelihood of the engraftment resulting from
teratoma formation from the gastruloid itself. On the other hand, it raises the possibility
that the inefficiency of the engraftment is the consequence of a low frequency of cells
within each haemGx with HSPC properties, which we can seek to resolve by upscaling
of the haemGx culture or further expansion of dissociated HSPC (Ng *et al.*, 2024).

Although the system is shy of achieving HSC maturation *in situ*, it can mature, to
produce low-level engrafting multilineage progenitors, and potentially HSCs. This
suggests that access to the relevant hematopoietic molecular programs is preserved.
Incipient specification of non-hematopoietic tissues shown to support HSC
emergence, including PDGFRA[+] mesenchymal (Chandrakanthan *et al.,* 2022) and
neuro-mesenchymal (Miladinovic *et al.,* 2024) cells, and neural-crest derivatives
(Kapeni *et al.,* 2022; Fitch *et al.,* 2012), occurs in sync with development of AGM-like
hematopoiesis, to indicate the presence of cues that with further maturation may
facilitate HSPC specification at the appropriate developmental time. Interestingly,
through extending the haemGx culture, and in addition to AGM-like hematopoiesis, we

also captured the potential to form renal tubular structures upon *in vivo* maturation,

suggesting induction of lateral plate and intermediate mesoderm from which the AGM

region derives, and putative reconstitution of the intra-embryonic hematopoietic niche.

Establishment of a tractable *in vitro* model of hematopoietic development which

faithfully recapitulates time-dependent emergence of hematopoietic progenitors in

their physiological niche would propel translational applications by enabling high-

throughput studies of HSC emergence. Most ESC-based models of blood specification

do not achieve concomitant recapitulation of the surrounding microenvironment.

Protocols typically remove ESC-derived hemogenic cells from an early embryonic-like

niche to direct differentiation through genetic manipulation (Sugimura *et al.,* 2017)

and/or empirically-determined growth factor addition (Lis *et al.,* 2017; Vo and Daley,

2015; Fowler *et al.,* 2024; Nafria *et al.,* 2020; Sroczynska *et al.,* 2009; Piau *et al.,*

2023), with minimal probing of intermediate steps. When hematopoietic waves have

been specifically inspected, emergence of YS and AGM-like stages were conflated in

time (Pearson *et al.,* 2015; Murry and Keller, 2008), suggesting a bias towards intrinsic

hematopoietic cell potential, at the expense of cell fate modulation by time-coherent

extrinsic physiological cues, affecting the nature of cellular output. The present

haemGx system, in contrast, successfully deconvolutes YS-like and AGM-like stages

of blood production, including capturing differences in the HE generation, suggesting

a level of organisation amenable to further development into an integrated hemogenic

niche. Indeed, projections to scRNA-seq datasets of *in vivo* murine hematopoiesis

place our endothelial and hematopoietic outputs at coherent temporal windows.

The haemGx protocol could be further developed to include more supportive

structures of tissue organisation, as non-hematopoietic gastruloid models have

benefitted from mechanical, or chemo-mechanical, support of artificial ECM

embedding (Veenvliet *et al.,* 2020; van den Brink, Susanne C *et al.,* 2020; Dijkhuis *et*

al., 2023), to match temporal coherence and tissue polarisation, with detailed tissue

architecture illustrated by somite formation. Prior to HSC emergence in the AGM,

critical signals such as SCF and Shh, which we have delivered to the cultures, are

secreted by different cells on opposite dorsal-ventral locations (Souilhol *et al.,* 2016),

indicating the potential relevance of tissue organisation to improve and potentially

extend hematopoietic formation. Moreover, HSPC formation in the AGM is facilitated

by circulatory shear stress (North *et al.,* 2009; Diaz *et al.,* 2015; Lundin *et al.,* 2020),

suggesting that introduction of flow may improve hemato-endothelial organization and

boost blood production.

We attested the utility of haemGx's to interrogate the biology of leukemias with

developmental origins, thanks to a nearly full recapitulation of embryonic hemogenic

specification, including critical early stages of HE production and EHT to EMP and

MLP/MPPs. We modelled *MNX1*-r AML, a poorly characterized form of leukemia

exclusive to the infant age group (Ragusa *et al.,* 2023). Overexpression of *MNX1* in

the haemGx model propagated cells bearing transcriptional resemblance to patient

MNX1-r AML, overall positioning the haemGx as an attractive tractable model in which

to study developmental leukemias. We showed that MNX1-OE transiently expanded

a C-Kit⁺ cell with hemato-endothelial characteristics at the YS-like stage, and could

selectively sustain the phenotype through serial replating of CFC assays. Replating

CFCs could be initiated beyond the YS-like stage, albeit at the relative expense of an

early differentiation state and of the similarity to patient transcriptome, putatively

defining a window of susceptibility to MNX1-driven transformation.

Importantly, the haemGx model was able to resolve the cellularity and transcriptional

features inferred from patient transcriptomes, with the advantage of a higher resolution

in dissecting the contribution of cell-of-origin in the appropriate developmental context.

A recent paper using MNX1-OE has shown that MNX1 can initiate a transplantable

leukemia in immunodeficient mice by targeting FL, but not BM, cells (Waraky *et al.,*

2024). This suggests that ectopic expression of *MNX1* transforms an embryonic cell,

rather than depends on an embryonic niche. It also suggests that MNX1-OE can cause

leukemia as a single genetic hit, although it cannot be excluded that other mutations

are required for, and may indeed have contributed to, full transformation *in vivo*, as

observed in a recent report confirming the *in utero* origin of the leukemia (Bousquets-

Muñoz *et al.,* 2024). However, the mouse model does not address the nature of the

embryonic cell targeted by MNX1-OE. Like the MNX1-OE haemGx cells selected

through serial replating, MNX1-OE mouse leukemia are also C-Kit⁺ (Waraky *et al.,*

2024). Interestingly, clusters of genes up-regulated in the mouse MNX1-OE leukemia

(**Fig. 7S1D** – k1, 2, 5, 6, and 9) do not robustly separate *MNX1*-r from other infAML by

GSEA analysis (**Fig. 7S1E**), and have diverged more from the patient data in their cell-

type signature analysis than the original MNX1-OE cells in the FL (**Fig. 7S1F**),
putatively selecting a more generic program of leukemia transformation. With the
resolution provided by the haemGx model, we propose that the divergence from
MNX1-r infAML signatures increases from MNX1-OE haemGx 144h , to 216h to FL,
positioning the origin of the AML at the YS stage, likely at the HE-to-EMP transition.
This cell may persist in the haemGx culture system to the 216h timepoint, and migrate
to the FL in the embryo, eventually extinguishing through differentiation, as suggested
by the MNX1-OE haemGx CFC assays. Persistence of the original MNX1-OE target
cell will determine the sensitivity to MNX1-OE, and can be addressed in the haemGx
system through lineage tracing. Whether additional mutations, namely a reported
association between *MNX1*-r and tri(19) (Espersen *et al.,* 2018; Ragusa *et al.,* 2023),
contribute to prolonging the undifferentiated state of the candidate YS target cell with
increased probability of transformation, or otherwise expand the targeted cell in a more
classical second-hit fashion, can potentially also be screened in the haemGx system,
in combination with transplantation experiments.

Over the last decade, the effort to achieve robust hematopoietic production from PSC,
mouse and human, has been centred on the generation of HSC, a critical translational
need (Chang *et al.,* 2023; Dijkhuis *et al.,* 2023; Fidanza and Forrester, 2021). There
are good indications that this is achievable, both with (Vo and Daley, 2015; Dang *et*
al., 2002) and without (Piau *et al.,* 2023) genetic manipulation. However, success
remains episodic, mainly because mechanistic understanding of processes
underpinning blood generation during development is limiting, exacerbated by limiting
access to human embryonic and fetal material. On-going optimization of gastruloid
models and the success at integration of organoid cultures in microfluidic systems
(Saorin, Caligiuri and Rizzolio, 2023; Quintard *et al.,* 2024) will allow us to further
optimize haemGx culture. In conclusion, the haemGx represents a faithful *in vitro*
reference to replace and complement embryo studies, holding promise to understand
the biology of healthy and malignant developmental hematopoiesis and advance blood
production for clinical use.

EXPERIMENTAL PROCEDURES

MATERIALS

	SOURCE	IDENTIFIER
Antibodies		
Anti-mouse c-Kit (CD117) clone 2B8 - APC-Cy7	Biolegend	Cat. #105825
Anti-mouse CD41 clone MWReg30 - PE-Dazzle594	Biolegend	Cat. #133935
Anti-mouse CD45 clone 30F11 - APC	Biolegend	Cat. #103111
Anti-mouse CD144 (VE-cadherin) clone BV13 - APC	Biolegend	Cat. #138011
Anti-mouse CD43 clone S11 - PE	Biolegend	Cat. #143205
Anti-mouse Ter119 - PC7	Biolegend	Cat. #116223
Anti-mouse CD31 - biotin	BDPharmingen	Cat. #553371
Anti-mouse CD34 - APC	Biolegend	Cat. #128611
CD11b anti-mouse/human clone M1/70 - biotin	Biolegend	Cat. #101203
Mouse Anti-Mouse CD45.2 clone 104 - PE	BD Biosciences	Cat. #560695
Rat Anti-Mouse CD31 clone MEC13.3 - biotin	BD Biosciences	Cat. #553371
Goat Anti-Mouse c-Kit/CD117	R&D Systems	Cat. #AF1356SP
Bacterial and virus strains		
E. coli: NEB 5-alpha Competent *E. coli*	New England Biolabs (NEB)	Cat. #C2987H
Chemicals, peptides, and recombinant proteins		
Murine LIF	Peprotech	Cat. #250-02
StemMACS PD0325901	Miltenyi Biotec	Cat. #130-106-5411
Chiron (CHIR99021)	Biogems	Cat. #2520691
Activin A PLUS	QKine	Cat. #Qk005
Murine VEGF 165	Peprotech	Cat. #450-32
Murine FGF-basic	Peprotech	Cat. #450-33
Murine Sonic Hedgehog (Shh)	Peprotech	Cat. #315-22
Murine TPO	Peprotech	Cat. #315-14
Murine Flt3-Ligand	Peprotech	Cat. #250-31L
Murine SCF	Peprotech	Cat. #250-03
N2B27 medium (NDiff 227)	Takara Bio	Cat. #Y40002
DMEM/F-12	Gibco	Cat. #11320033
Neurobasal medium	Gibco	Cat. #21103049
N-2 Supplement	Gibco	Cat. #17502048
B-27 Supplement	Gibco	Cat. #17504044
Critical commercial assays		
Mouse Methylcellulose Complete Media	R&D Systems	Cat. #HSC007
Deposited data		
Single cell RNA-seq of hemogenic gastruloids at timepoints	This paper	ArrayExpress: E-MTAB-12148
Bulk RNA-seq of hemogenic gastruloids with MNX1 overexpression at 144h and 216 h	This paper	ArrayExpress: E-MTAB-12173
Experimental models: Cell lines		
Mus musculus: Flk-1-GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	Jakobsson *et al.*, 2010
Mus musculus: Sox17-GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	Niakan *et al.*, 2010
Mus musculus: T/Bra::GFP mouse embryonic stem cell line	Alfonso Martinez Arias Lab	Fehling *et al.*, 2003
Mus musculus: E14Tg2a mouse embryonic stem cell line	MMRRC, University of California Davis, US	Hooper *et al.*, 1987

Mus musculus: Flk-1-GFP::Rosa26-BFP mouse embryonic stem cell line	This paper	n/a
Homo sapiens: HEK293T	ATCC	CRL-3216
Oligonucleotides		
MNX1 (forward 5-GTTCAAGCTCAACAAGTACC-3; reverse 5- GGTTCTGGAACCAAATCTTC-3)	Gulino *et al.*, 2021	n/a
Ppia (forward 5- TTACCCATCAAACCATTCCTTCTG-3; reverse 5- AACCCAAAGAACTTCAGTGAGAGC-3)	Moris *et al.*, 2018	n/a
BFP (forward 5-GCACCGTGGACAACCATCACTT-3; reverse 5-CAGTTTGCTAGGGAGGTCGC-3)	This paper	n/a
Recombinant DNA		
pWPT-LSSmOrange-PQR	Ghevaert Lab, WT-MRC Cambridge Stem Cell Institute, UK	Dalby *et al.*, 2018
pWPT-LSSmOrange-PQR-MNX1	Cloned by Biomatik Corporation (Kitchener, Canada)	n/a
pcdna3-rosa26-rosa26-bfp	This paper	n/a
Software and algorithms		
GSEA software v4.2.3	Subramanian *et al.*, 2005	https://www.gsea-msigdb.org/gsea/index.jsp
ImageJ	Schneider *et al.*, 2012	https://imagej.nih.gov/ij/
Bowtie2	Langmead and Salzberg, 2012	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
Samtools	Li *et al.*, 2009	http://samtools.sourceforge.net/
Tophat2	Kim *et al.*, 2013	https://ccb.jhu.edu/software/tophat/index.shtml
HTseq	Anders *et al.*, 2015	https://htseq.readthedocs.io/en/master/
DEseq2	Love *et al.*, 2014	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
ExpressAnalyst	n/a	https://www.expressanalyst.ca/
PANTHER	Thomas *et al.*, 2022	http://www.pantherdb.org/
EnrichR	Chen *et al.*, 2013; Kuleshov *et al.*, 2016; Xie *et al.*, 2021	https://maayanlab.cloud/Enrichr/
Panglao BP	Franzen *et al.*, 2019	https://panglaodb.se
TARGET (Therapeutically Applicable Research to Generate Effective Treatments) Database	https://ocg.cancer.gov/programs/target https://xenabrowser.net/	GDC TARGET-AML
Single-cell RNA sequencing analysis scripts	DockerHub: dsblab/single_cell_analysis:0.5	https://github.com/dsb-lab/blood_gastruloids
Other		
CELLSTAR cell-repellent cell culture plate, 96 well, U-bottom	Greiner (Bio-One)	Cat. #650970

METHODS

Cell culture

Flk-1-GFP (Jakobsson *et al.,* 2010), Flk-1-GFP::Rosa26-BFP, Sox17-GFP (Niakan *et al.,* 2010), T/Bra::GFP (Fehling *et al.,* 2003), and E14Tg2A (Hooper *et al.,* 1987) mouse embryonic stem cells (mES) lines were cultured in ES+LIF medium in gelatinized flasks (0.1% gelatin) with daily medium change, as previously described (Turner *et al.,* 2017). The ES+LIF medium contained 500 ml Glasgow MEM BHK-21 (Gibco), 50 ml of fetal bovine serum (FBS, Embryonic Stem Cells tested, biosera, Nuaillé, France), 5 ml GlutaMAX supplement (Gibco), 5 ml MEM Non-Essential Amino Acids (Gibco), 5 ml Sodium pyruvate solution (100 mM, Gibco), and 1 ml 2-Mercaptoethanol (50mM, Gibco). Murine LIF (Peprotech) was added at 1000 U/ml. HEK293T cells were grown in Dulbecco's modified Eagle medium (Gibco) supplemented with 10% FBS (Gibco) and 1% penicillin/streptomycin (Gibco). All cultures were kept at 37°C and 5% CO_2.

Hemogenic gastruloids assembly, culture, and dissociation

mES were maintained in ES+LIF medium and transferred to 2i+LIF (containing Chiron and MEK inhibitor PD03) for 24 hours prior to the assembly into gastruloids. 250 untransduced mES cells (or 400 for GX-MNX1 and GX-EV) were seeded in each well of a U-bottom, cell-repellent 96-well plate (Greiner Bio-One, Stonehouse, UK) in 40 µl of N2B27 medium [Takara Bio or home-made as Cold Spring Harbour Protocols 'N2B27 Medium' (2017)]. The plate was centrifuged at 750 rpm for 2 minutes to promote deposition and aggregation of the cells and was then incubated at 37°C, 5% CO2 for 48 hours. After 48 hours, 150 µl of N2B27 medium supplemented with 100 ng/ml Activin A (QKine, Cambridge, UK) and 3 µM chiron (Peprotech) was added to each well. At 72 hours, 150 µl of medium were removed, without disrupting the gastruloids in the wells. 100 µl of N2B27 with 5 ng/ml of Vegf and Fgf2 each (Peprotech) were added to each well. From 72 h to 144 h, each day 100 µl of medium were removed and replaced with N2B27 + Vegf + Fgf2. At 144 h, the medium was further supplemented with Shh at 20 ng/ml. From 168 h to 216 h, the medium was

N2B27 + 5 ng/ml Vegf, plus 20 ng/ml mTpo, 100 ng/ml mFlt3l, and 100 ng/ml mScf

(Peprotech). To dissociate cells from the gastruloid structures, medium was removed

and individual gastruloids were collected using a pipette and precipitated at the bottom

of a microcentrifuge tube. The remaining medium was aspirated and the bulk of

gastruloids was washed in PBS. 50 µl of TrypLE express was added to pelleted

gastruloids to be incubated at 37°C for 2 minutes to dissociate cells.

Animals and implantation

Female athymic nude-Foxn1^{nu} (*nu/nu*) mice (Envigo, Bresso, Italy) were housed under

pathogen-free conditions. In accordance with the "3Rs policy", experiments were

reviewed and approved by the Animal Welfare Body (OPBA) of IRCCS Ospedale

Policlinico San Martino and by the Italian Ministry of Health (n. 883/2020-PR).

Intact haemGx (3 per mouse in 10 µL of PBS) were inoculated in the adrenal gland of

five-week-old mice. Briefly, mice were anesthetized with a mixture of xylazine

(ROMPUN, Bayer) and ketamine (LOBOTOR, Acme S.r.l.) and injected with haemGx,

after laparotomy, in the capsule of the left adrenal gland, as previously described

(Brignole *et al.,* 2023; Pastorino *et al.,* 2003). All mice survived surgery. Control mice

(CTR) received vehicle only. Mice body weight and general physical conditions were

recorded daily. One month after haemGx inoculation, mice were sacrificed and spleen

and bone marrow (BM) from femurs collected. BM cells and spleens, mechanically

reduced to cell suspension, were stored at -80°C until use. Adrenal glands were

paraffin-embedded for subsequent IHC studies.

Methylcellulose colony forming assays (CFC)

Disassembled gastruloids cells were plated on Mouse Methylcellulose Complete

Media (R&D Systems). Cells were first suspended in Iscove's Modified Dulbecco's

Medium (IMDM, Gibco) with 20% FBS (Gibco) before addition to the methylcellulose

medium. Cells were plated in duplicate 35 mm dishes with 2×10^5 cells/plate. Plates

were incubated at 37°C and 5% CO2 for 10 days, when colonies were scored. For

serial replating experiments, cells in methylcellulose were collected and washed in

phosphate buffer saline (PBS) to achieve single-cell suspensions and replated as

described above. Dissociated gastruloid cells from CFC were collected as cell
suspensions in PBS and centrifuged onto slides (Shandon Cytospin 2 Cytocentrifuge)
at 350rpm for 5-7min, with a PBS-only pre-spin at 1000rpm, 2min. Slides were stained
in Giemsa-Wright stain for 30 seconds followed by addition of phosphate buffer pH 6.5
for 5 min.

Immunofluorescence staining

Immunostaining of whole gastruloids was performed as described before (Baillie-
Johnson *et al.*, 2015). Briefly, gastruloids were fixed in 4% paraformaldehyde (PFA)
dissolved in PBS for 4 hours at 4°C on orbital shaking and permeabilised in PBSFT
(10% FBS and 0.2% Triton X-100), followed by one hour blocking in PBSFT at 4°C on
orbital shaking. Antibody dilutions were made in PBSFT at 1:200 for primary and 1:500
for secondary antibodies. Antibody incubations were performed overnight at 4°C on
orbital shaking, and subjected to optical clearing overnight in ScaleS4 clearing
solution. Individual gastruloids were then mounted on glass coverslips by pipetting as
droplets in ScaleS4 and DAPI nuclear stain.

Histology and H&E staining

Samples were fixed with 10% formalin at room temperature for 48 hours followed by
an ethanol series of ethanol and paraffin; ethanol 70% (90 minutes), ethanol 80% (120
minutes), ethanol 95% (180 minutes), 3 times ethanol 100% of 120, 120 and 180
minutes, 50% ethanol/paraffin (3x120 minutes) and 3 times paraffin (60, 90 and 120
minutes) before embedding in histological cassettes. Blocks were sectioned with a
HistoCore BIOCUT Microtome (Leica) at 10 mm and slides were stained with
Haematoxylin/Eosin.

Imaging

Images of cultured gastruloids and CFC plates were captured using the Cytation 5 Cell
Imaging Multi-Mode Reader (Biotek) plate reader using bright field and FITC channels.
ImageJ (Schneider *et al.*, 2012) was used for gastruloid size quantification. Confocal

microscopy was performed on LSM700 on a Zeiss Axiovert 200 M with Zeiss EC Plan-

Neofluar 10x/0.30 M27 and Zeiss LD Plan-Neofluar 20x/0.4 M27 objective lens. Tissue

sections were imaged with a THUNDER Imager 3D microscope system (Leica).

821   Lentiviral vector packaging and transduction

The lentiviral overexpression vector pWPT-LSSmOrange-PQR was used to clone the

MNX1 gene cDNA. The viral packaging vectors pCMV and pMD2.G, described in Pina

et al. (2008), were used to assemble lentiviral particles using HEK293T cells via

transfection using TurboFect Reagent (Invitrogen). Transduction of mES cells was

performed overnight by addition of lentivirus to culture medium and washed the

following day (Moris *et al.,* 2018). Transduced cells were sorted for positivity to

LSSmOrange.

830   Flow cytometry

Surface cell marker analysis was performed by staining using the antibodies listed in

Key Resources Table. Disassembled gastruloids cells were resuspended in PBS, 2%

FBS and 0.5 mM EDTA and stained at a dilution of 1:100 for primary antibodies for 20

minutes at 4°C. When indicated, streptavidin was added at a dilution of 1:200. Analysis

was performed on ACEA Novocyte (Agilent) or AttuneNxT (Thermo) analyzers, using

the respective software packages. Cell sorting was performed using a CS&T calibrated

BDFACS Aria III system (488nm 40 mW, 633nm 20 mW, 405nm 30 mW, and 561nm

50 mW), set with the 100μm nozzle at 20PSI and a 4-way purity mask or, in the case

of haemGx adrenal implants, a Beckman Coulter CytoFlex SRT (488nm 50 mW,

633nm 100 mW, 405nm 90 mW, and 561nm 30 mW), set with 100μm nozzle at 15PSI,

also using 4-way purity. Single-cell deposition in 96-well plates was performed using

single-cell sorting mode. Intact cells were gated on FSC-A vs SSC-A plot, followed by

doublet exclusion on FSC-A vs FSC-H and SSC-A vs SSC-H, prior to gating on

fluorescent parameters for the markers described in the results.

846   Single cell RNA sequencing of time-resolved hemogenic gastruloids

Gastruloids were collected at different timepoints of the protocol, disassembled and
FACS deposited into 96-well plates, either as unsorted (global) or sorted by CD45,
CD41 and C-Kit/ScaI markers (sorted). RNA from the cells were extracted from single
cells using Smart-seq2 technology at 500Kb and depth of 151 bases. Sequencing
reads were quality-checked using FastQC (v0.11.9). We trimmed the samples using
trimGalore! (0.6.6) with a cutoff of 30, clipping 15 base pairs and retaining reads of
more than 100 bases. Alignment was performed on STAR (2.7.8a) and Mus musculus
annotations from GENCODE vM26. Aligned BAM files were annotated using
featureCounts (v2.0.1) and count matrices were computed in python by directly
accessing the BAM files with gtfparse packages and collapsing lanes.

For quality control, based on the histogram of counts and multimodality distributions,
we set a minimum count threshold of 200000 counts, a minimum threshold of 1000
expressed genes, and a maximum threshold of 20% mitochondrial fraction per cell.
We performed the same procedure of the second dataset with a more restrictive
threshold of 400000 counts and similar expressed genes and mitochondrial
thresholds. Cells that did not pass the quality control metrics were omitted from
analysis. We normalized the cells to the mean count number per dataset and applied
a plus-one-log transformation of the data before proceeding to the downstream
analysis.

Dimensionality reduction was performed on feature selection of the gene space using
the function scanpy.highly_varying_genes with default parameters. Selection of
principal components in principal component analysis (PCA) was performed by
heuristic elbow method. Nearest neighbor analysis was constructed by KNN graphs
using a correlation metric and 10 nearest neighbors. Data was projected for low
dimensional visualization using the UMAP algorithm with default parameters as
implemented in scanpy.tl.umap. We used the leiden algorithm as implemented in
scanpy.tl.leiden to partition the data into clusters. In order to assess the election of the
resolution parameter we used Newman-Girvan modularity as a metric of clustering
quality. Differential expressed genes were computed comparing each cluster against
the rest using the Wilcoxon test with Benjamini-Hochberg correction.

Annotation and projection to additional datasets

Raw counts matrices were downloaded and processed following the same pipeline as
described above for scRNA-seq of gastruloids to generate UMAP and clustering. We
implemented the scmap algorithm to compare our scRNA sequencing of gastruloids
with the available datasets. We reduced the dimensionality of the space by selecting
highly varying genes from the annotated dataset. Then, we constructed a KNN
classifier with correlation metric and computed the nearest neighbors of the target
data. If neighbors with correlation metrics below 0.7 default standards, the projected
cells were not projected onto any cell from the annotated dataset. To visualize the cells
over the UMAP plots of the other datasets, we constructed a KNN regressor with three
neighbors and a correlation metric. Confusion matrices were used to visualize the
overlap between gastruloid clusters and the embryo dataset assigned clusters using
scmap.

Bulk RNA sequencing

Total RNA was extracted from disassembled gastruloid cells at 216 hours. Sequencing
was performed on NovaSeq PE150 platform, at 20M paired-end reads per sample.
Tophat2 with Bowtie2 were used to map paired-end reads to the reference Mus
musculus genome build GRCm39. GENCODE Release M30 (Frankish *et al.,* 2019)
was used as the reference mouse genome annotation. Aligned reads were filtered by
quality using samtools (Li, H. *et al.,* 2009) with a minimum threshold set at 30 (q30).
Transcript assembly and quantification was achieved using htseq (Putri *et al.*, 2022).
Differential expression between sample and control was performed by collapsing
technical replicates for each condition using Deseq2 (Love, Huber and Anders, 2014)
in R environment (Deseq2 library v 1.32.0). The differential expression was expressed
in the form of log2 fold change and filtered by false discovery rate (FDR) of 0.05.

Gene list enrichment analyses

Gene ontology (GO) analysis was performed in ExpressAnalyst (available at
www.expressanalyst.ca) using the PANTHER or GO Biological Process (BP)
repository. GO terms and pathways were filtered by false discovery rate (FDR) with a
cut-off of ≤ 0.1 for meaningful association. EnrichR (Xie *et al.,* 2021; Kuleshov *et al.,*
2016; Chen, E. Y. *et al.,* 2013) was used for cell type analysis using the Panglao DB

(Franzén, Gan and Björkegren, 2019) databases using a FDR threshold of ≤ 0.1 or p

value ≤ 0.01, where specified, as well as transcription factor (TF) binding site

enrichment using the ENCODE and ChEA Consensus TFs from ChIP database.

Gene Set Enrichment Analysis (GSEA)

Custom gene signatures were used as gene sets for GSEA analysis (Subramanian *et*

al., 2005) on the GSEA software v4.2.3 on RNA sequencing expression values in

counts units. GSEA was ran in 10000 permutations on gene set using the weighted

Signal2Noise metric. Enrichment metrics are shown as normalized enrichment score

(NES) and filtered by FDR ≤ 0.05. Leading edge genes (LEGs) are genes with a "Yes"

values for core enrichment. For AML patient analysis, clinical phenotype and

expression data (in counts units) were extracted from the GDC TARGET-AML cohorts

in the Therapeutically Applicable Research to Generate Effective Treatments project

(TARGET, https://ocg.cancer.gov/programs/target), downloaded from the University of

California Santa Cruz (UCSC) Xena public repository (last accessed 31st August

2022). Patient samples were selected according to the reported karyotype to include

t(7;12), inv(16), MLL, normal karyotype, and t(8;21). GSEA was performed comparing

RNA sequencing counts of t(7;12) samples against pooled AML subtypes (inv(16),

MLL, normal karyotype and t(8;21)) as "other AML".

Real-time polymerase chain reaction (qPCR)

Extracted RNA was reverse-transcribed into complementary DNA (cDNA) using High-

Capacity RNA-to-cDNA Kit (Applied Biosystems). QPCR was performed using

FastGene 2x IC Green Universal qPCR Mix (Nippon Genetics, Duren, Germany) using

primers for human *MNX1* (forward 5-GTTCAAGCTCAACAAGTACC-3; reverse 5-

GGTTCTGGAACCAAATCTTC-3) (Gulino *et al.*, 2021) and *Ppia* for endogenous

control   (forward   5-TTACCCATCAAACCATTCCTTCTG-3;   reverse   5-

AACCCAAAGAACTTCAGTGAGAGC-3) (Moris *et al.,* 2018),   or Taqman gene

expression   assays   (Mm00501741_m1   *Myb*;   Mm00439364_m1   *Hoxa9*;

Mm00723157_m1   *Hlf*;   Mm01213404_m1   *Runx1*;   Mm00444619_m1   *Dll4*;

Mm00491303_m1 *Mecom*; Mm02342430_g1 *Ppia*) using Taqman Gene Expression

Mastermix (Applied Biosystems). Differential gene expression was calculated using
the delta delta Ct (ΔΔCt) method.

Polymerase chain reaction (PCR)

Genomic DNA was extracted using Monarch Genomic DNA Purification Kit (New
England Biolabs, Hitchin, UK) using manufacturer's instructions. PCR on genomic
DNA was performed using Phire PCR Master Mix (Thermo Scientific) to amplify the
BFP locus using forward primer 5'-GCACCGTGGACAACCATCACTT-3' and reverse
primer 5'-CAGTTTGCTAGGGAGGTCGC-3'.

Statistical analysis

Experiments were performed at least in triplicates, unless specified otherwise. Data
are plotted to include standard deviation (+/-SD) between replicates. Statistical
significance was set at a threshold of p value < 0.05. Statistical analysis was performed
in R environment (version 4.1.3) or using GraphPad Prism 8.0 software and is detailed
in respective figure legends.

Data availability

Raw data as well as processed count matrices and post-processed files from single-
cell RNA-seq for the time-resolved data is available at E-MTAB-12148. Bulk RNA-seq
of MNX1 overexpressing gastruloids is available at Array Express with accession code
E-MTAB-12173. The post-processing was performed in Python on DockerHub:
dsblab/single_cell_analysis:0.5. Scripts are available in https://github.com/dsb-
lab/blood_gastruloids and Zenodo (https://doi.org/10.5281/zenodo.7053423). The
results published here are partly based upon data generated by the Therapeutically
Applicable Research to Generate Effective Treatments (TARGET)
(https://ocg.cancer.gov/programs/target) initiative, of the Acute Myeloid Leukemia
(AML) cohort GDC TARGET-AML. The data used for this analysis are available at
https://portal.gdc.cancer.gov/projects and https://xenabrowser.net/.

ACKNOWLEDGEMENTS

This project was funded by a start-up grant and a BRIEF award from Brunel University London, a Little Princess Trust (LPT) Project Grant (CCLGA 2023 22 Pina) and a National Centre for the Replacement, Refinement and Reduction of Animals in Research (NC3Rs) project grant (NC/Z500677/1) to CP, by an Agencia Estatal de Investigación grant (PLEC2021-007518) to AB and AMA, and by an ERC Advanced Grant (MiniEmbryoBlueprint 834580) to AMA. LPT research is funded in partnership with CCLG through the CCLG Charity Research Network. DR received funding from the Royal Society of Biology (MRSB Travel Grant). GTC was funded by grant FPU18/05091 from the Spanish Ministry of Universities. AJ is funded by a Lady Tata Memorial Trust International Scholarship (2022). SvdB was funded by an EMBO Postdoctoral Fellowship (ALTF 195-2021) and is in receipt of a HFSP Postdoctoral Fellowship (LT0047/2022-L). Work in the CP lab was also funded by a KKLF Intermediate Fellowship (KL888), a Leuka John Goldman Fellowship for Future Science (2017-2019), and a Wellcome Trust / ISSF Bridge Funding award at the University of Cambridge (2019). JGO acknowledges financial support from the Spanish Ministry of Science and Innovation and FEDER (grant PGC2018-101251-B-I00), by the Maria de Maeztu Programme for Units of Excellence in R&D (grant CEX2018-000792-M), and by the Generalitat de Catalunya (ICREA Academia programme). MP acknowledges funding from the Italian Ministry of Health (Ricerca Finalizzata 5 per mille and Ricerca Corrente to MP). Library preparation and next-generation sequencing for single-cell RNA-seq analysis were performed by the Single Cell Genomics Group at the National Centre for Genomic Analysis – Centre for Genomic Regulation (CNAG-CRG), Barcelona. The Authors wish to acknowledge Tina Balayo, Ana Filipa Domingues, Shikha Gupta, Oliver Davies, Kristen Place and Remisha Gurung's technical support at different stages of the project.

AUTHOR CONTRIBUTIONS

Conceptualization: CP, AMA; Methodology: CWS, DR, GTC, FP, SvdB, MC,MP, KRK,

VH-H, AB, JGO, AMA, CP; Software: GTC, JGO; Validation: CWS, DR, AJ, YC, SvdB,

CP; Investigation: CWS, DR, GTC, FP, AJ, YC, LD, CB, G-AI, CP; Formal Analysis:

GTC, DR, JGO, CP; Resources: GTC, JC, MP, JGO, AMA; Data curation: DR, GTC,

JGO; Writing – Original Draft: CP, DR, GTC; Writing – Review and editing: CP, DR,

GTC, AB, JGO, AMA; Visualisation: DR, GTC, CWS, CP; Supervision: CP; Project

administration: CP, AMA, JGO; Funding acquisition: CP, AMA, AB.

1010

1011

1012   **DECLARATION OF INTERESTS**

1013   Authors declare no competing interests.

SUPPLEMENTAL INFORMATION

Supplemental Figures S1-S7

Supplemental Files S1-S3 (Excel files):

Supplemental File S1: Cluster classifier gene lists for each cluster by differential gene expression analysis of scRNA-seq of haemGx (all conditions), obtained by Wilcoxon rank test of each cluster against all other clusters.

Supplemental File S2: Cell type enrichment analysis of cluster classifier genes, showing the inferred identity of clusters of scRNA-seq of haemGx (all conditions). Classifier genes (in Supplemental File S1) were obtained by differential expression in comparison to all other clusters and subjected to enrichment analysis using the PanglaoDB repository (Franzén, Gan and Björkegren, 2019).

Supplemental File S3: Summary of projections of scRNA-seq of haemGx cells onto available datasets by Hou et al. (2020), Zhu et al. (2020), and Thambyrajah et al. (2024), showing the identity of each haemGx cell projecting to annotated clusters of the respective studies.

Supplemental File S4: List of differentially expressed genes between haemGx-MNX1 and haemGx-MNX1 at 144h and 216h compared to empty vector (EV) control.

  FIGURE LEGENDS

  Figure 1 – Hemogenic gastruloids (haemGx) produced from mES cells promote hemato-
  endothelial specification with spatio-temporally accurate ontogeny. (A) Timeline of mES
  cells assembly and culture into gastruloids over a 216h period with the addition of specified
  factors for the promotion of hemato-endothelial specification. The 24-hour pre-treatment in
  2i+LIF was omitted when mES cultures showed no signs of differentiation. **(B)** Imaging of
  haemGx over time from 72h to 216h at 10x magnification, showing the assembly and growth
  of the 3D structures and the polarization of the Flk-1-GFP marker from 96h; scale bar: 100mm.
  (C) Flow cytometry timecourse analysis of haemGx for the presence of CD144 (VE-cadherin),
  C-Kit, CD41, and CD45 markers; representative plots. **(D)** Quantification of analysis in (C).
  Violin plots with median and interquartile range of up to 26 replicates; Kruskal-Wallis CD144
  (p=0.0755), C-Kit (p<0.0001), CD41 (p<0.0342), CD45 (p<0.0001). Shown are significant
  Dunn's multiple comparisons tests; adjusted q-value. **(E)** Immunostaining of whole individual
  haemGx at 216h showing the localized expression of Flk1-GFP (green), C-Kit (yellow), and
  CD45 (red) markers; nuclear staining is in DAPI (blue); scale bar: 100μm.

  Figure 1 Supplement 1 – Optimization and reproducibility of the haemGx protocol. (A)
  Imaging of a 120h-haemGx with and without the addition of Activin A in conjunction with Chiron
  99021 (CHIR) to achieve polarized expression of *Flk-1-GFP*; scale bar: 100mm. **(B)** Titration
  of Activin A dosage (from 25 ng/ml to 150 ng/ml) required for *Flk-1-GFP* expression activation
  (green). **(C)** Representative flow cytometry plots of %CD45$^+$ cells at 216h without and without
  addition of SCF, Flt3L and TPO (SFT) from 168h; representative plots. **(D)** Quantification of
  %CD45$^+$ cells by flow cytometry analysis of haemGx cultured with or without SFT, between
  192-216h and 168-216h. Mean ± SD of n=7-8 independent experiments; Tukey's multiple
  testing with significant q-value<0.05. **(E)** Quantification of %CD45$^+$ cells at 216h by flow
  cytometry analysis of haemGx cultured in VEGF+FGF or VEGF-only during the 168-216h
  period. Mean ± SD of n=7 independent experiments; unpaired t-test; ns, not significant. **(F)**
  Flow cytometry monitoring of C-Kit, CD41, and CD45 markers in haemGx established from
  E14 and *Flk-1-GFP* mES lines; n=3-4 replicate experiments (n=1, CD45$^+$ 120h). Mixed effects
  model with Sidak's multiple comparison test; significant adjusted p-value <0.05. **(G)**
  Representative flow cytometry plots of CD41$^+$ and C-Kit$^+$ cells in E14 and *Flk-1-GFP* mES cell-
  initiated haemGx. **(H)** Representative flow cytometry plots of CD45$^+$ cells in E14 and *Flk-1-*
  GFP mES cell-initiated haemGx. **(I)** Quantification of %CD45$^+$ cells in *Flk-1-GFP*, *Sox17-GFP*,
  and *TBra-GFP* mES cell- initiated haemGx; n=5, mean ± SD. Welch's t-test Sox17-GFP vs.
  Flk1-GFP, p=0.5254; Tbra-GFP vs. Flk1-GFP, p=0.3782.

1069

Figure 2 – HaemGx produce hematopoietic output following the emergence of specialised endothelium. (A) Immunostaining of whole individual haemGx at 144h showing the establishment of a vascular network by co-expression of CD31 and Flk1-GFP; nuclear staining is in DAPI (blue); scale bar: 100μm. **(B)** Flow cytometry analysis of haemGx cells at 144h and 216h stained for CD41, CD45, CD31, and CD34. **(C)** Quantification of flow cytometry analysis in (B) of CD41+ and CD45+ fractions co-expressing CD34 and CD31. Mixed effect analysis with a Ŝidàk test for multiple comparisons. **(D)** Colony-forming cell (CFC) assay of 216h-haemGx in multipotential methylcellulose-based medium. GEM: granulocyte-erythroid-monocyte; GM: granulocyte-monocyte; M: monocyte; E: erythroid. Please note that the medium does not include TPO and does not assess the presence of megakaryocytic progenitors. CFC frequency in 3 haemGx, n=3, mean±SD. **(E)** Representative photographs of colonies quantified in C; scale bar=2mm. **(F)** Real-time quantitative (qPCR) analysis of expression of hemato-endothelial genes by timepoint. Relative gene expression fold change calculated by normalization to Ppia. Bars represent mean of 3 replicates and show individual data points; p values by ANOVA across datapoints; post-hoc statistical significance between specific variables by Tukey's test and shown by brackets.

Figure 2 Supplement 1 – Multi-color flow cytometry detection of surface markers in haemGx. (A) Representative flow cytometry plots of CD31 and CD34 staining of haemGx at 144h and 216h gated on Flk-1-GFP$^+$. **(B)** Quantification of Flk-1-GFP$^+$ haemGx populations at 144h and 216h expressing CD34 and CD31. Crossbar shows mean of 4 replicate experiments with individual data points shown; Welch's t-test; p value significant < 0.05. **(C-E)** Unstained controls (C-D) and fluorescence minus one (FMO) controls (E) relative to flow cytometry detection of multi-color of surface markers in Figure 2. **(F)** Representative flow cytometry plot of haemGx of CD45$^+$CD41lo fractions co-expressing C-Kit and Cd144 (Ve-Cadherin).

Figure 2 Supplement 2 – Characterization of hematopoietic output from haemGx. (A) Representative image of cytospins of dissociated haemGx at 216h stained with Giemsa-Wright's stain. Annotated are cells in the monocytic (dashed open arrow), granulocytic (solid open arrow), megakaryocytic (solid arrow) and erythroid (asterisk) lineages; arrowheads indicate cells with a non-specific blast-like morphology. 40x magnification, scale bar = 100 μm. **(B)** Representative flow cytometry plots of CD41 and CD45 expressing populations in haemGx treated with 0.5 μM EZH2 inhibitor GSK126 or with 0.05% DMSO (control) at 144h and 216h. **(C)** Quantification of flow cytometry analysis in (B) showing the proportion of CD41$^+$ and CD45$^+$CD41^{lo} populations in GSK126-treated haemGx. Welch's t-test; p value significant < 0.05.

1107

Figure 3 – Time-resolved analysis of haemGx by scRNA-seq captures successive waves of hematopoietic specification. (A) UMAP clustering of subsets of sequenced cells expressing CD41, C-Kit, and CD45 at 144h, 192h and 216h, annotated by timepoint (left panel), sorting condition (middle panel), and Leiden clustering (right panel). **(B)** UMAP panels highlighting the expression of specific hemato-endothelial markers in the transcriptional spaces defined in (A). Color scale indicates expression levels in counts. **(C)** Heatmaps of differentiation of an arterial endothelial programme in haemGx from scRNA-seq data of sorted on expression of C-Kit at 144h and 192h. Color scale indicates expression in counts. **(D)** Violin plots quantifying the expression of hemato-endothelial markers in C-Kit+ fraction from clusters in (A) comparing 144h and 192h timepoints. Wilcoxon test; significant p value<0.05. **(E)** Expression of definitive hematopoietic genes in in C-Kit$^+$, CD41$^+$ and CD45$^+$ cells at 144h and 192/216h timepoints; expression levels as normalised counts per million reads. Mann-Whitney or Kruskal-Wallis with Dunn's multiple comparisons; significant p or q-value<0.05.

Figure 3 Supplement 1 – Single-cell-RNAseq (scRNAseq) analysis of haemGx identifies time-dependent signatures of endothelial, hemogenic and stromal cells. **(A)** Summary of plating strategy for scRNA-seq analysis of gastruloids at 120h, 144h, 168h, 192h, and 216h without selection of surface markers ('all') or sorted as C-Kit/Sca1+ (green shading), CD41+ (blue), or CD45+ (orange) cells; P = plate. **(B)** ScRNA-seq quality control measures (quantification of mapped reads, individual genes and mapping of reads on to mitochondrial (mt)DNA. **(C)** Mapped read and gene counts for individual Smart-seq2 libraries (plate, P), biological replicate (repl) populations of single cells, and read and map distributions across unsorted and cell surface phenotype-sorted cells. **(D)** UMAP projection of all sequenced cells colored by annotated clusters (left), by sorting markers C-Kit/Sca1, CD41, and CD45, or unfractionated cells ('whole' corresponding to 'all' in panel A) (center), and by haemGx culture time (right). **(E)** Cell type enrichment analysis of cluster classifier genes, using the PanglaoDB repository (Franzén, Gan and Björkegren, 2019); classifier genes are obtained by differential expression in comparison to all other clusters. Represented are clusters 1, 3 and 9, which are characteristic of timepoints 168-216h, and capture stromal cells putatively relevant for hemogenic support, including autonomic neurons (cluster 9). The statistical power of representation of individual cell types is expressed as the EnrichR combined score with a p-value threshold of < 0.05.

Figure 3 Supplement 2 – Single-cell-RNAseq (scRNAseq) analysis of haemGx. (A) Heatmap representation of the expression of endothelial and hematopoietic marker genes in

individual haemGx CD41$^+$ and CD45$^+$ cells sorted at 192 and 216h. Cells and genes ordered by unsupervised hierarchical clustering. **(B)** Violin plots quantifying the expression of arterial markers in C-Kit+ fraction from clusters in Fig. 3A comparing 144h and 192h timepoints. Wilcoxon test; significant p value<0.05. **(C)** Summary representation of the proportion of hGx cells expressing endothelial (endo), erythroid/myeloid (Ery+My), myeloid/lymphoid (My+Ly) and HSPC gene signatures. Endo: *Kdr*; Ery+My: *Epor±Gata1±Klf1±Hbb±Hba±Hbg + Spi1±Mpo±Anpep±Csf1/2/3r*); My+Ly: *Spi1±Csf1/2a/3r + Ikzf1±Ighm/d±Igk*); HSPC: *Ptprc + Myb*).

Figure 4 – Late-stage haemGx contain hematopoietic output with transcriptional alignment to mouse embryonic populations and in vivo engraftment potential. (A) Projection of clustered haemGx cells (CD41, C-Kit, and CD45 at 144h, 192h and 216h; Fig. 3A) onto mouse single-cell RNA-seq datasets capturing: arterial and haemogenic specification in the para-splanchnopleura (pSP) and AGM region between E8.0 and E11 (Hou et al. 2020); YS, AGM and FL progenitors and the AGM EHT (Zhu et al., 2020); HSC emergence from the dorsal aorta HE (Thambyrajah et al., 2024). Gastruloid cell projection identified as per their cluster of origin (Fig. 3A). **(B)** Schematic representation of the experimental workflow for execution and analysis of haemGx implantation in the adrenal gland of immunodeficient mice. **(C)** PCR detection of haemGx genomic (g)DNA in the bone marrow (BM) and spleen (Spl) of immunodeficient mice 4 weeks after unilateral adrenal implantation of 3 gastruloids/gland. Analysis of 5 replicates of 100ng gDNA/recipient tissue; control animal was injected unilaterally with PBS in an adrenal gland in parallel with experimental implantation. Reaction positive control used 100ng of gDNA from *Rosa26-BFP*::*Flkj1-GFP* mES cells (mES) used to generate haemGxs. NTC: no template control. **(D)** Flow cytometry plots of engraftment detection by BFP expression in bone marrow (BM) of recipient mice 4 weeks following implantation of 216h haemGx. **(E)** Flow cytometry plots of lineage affiliation of BFP$^+$ engraftment in bone marrow (BM) of recipient mice 8 weeks following implantation of 216h haemGx.

Figure 4 Supplement 1 – Analysis of mouse embryonic scRNA-seq datasets for transcriptional similarities of haemGx outputs. (A-C) UMAPs of mouse single-cell RNA-seq datasets annotated as in respective publications: (A) arterial and haemogenic specification in the para-splanchnopleura (pSP) and AGM region between E8.0 and E11 (Hou et al. 2020); (B) YS, AGM and FL progenitors and the AGM EHT (Zhu et al., 2020); (C) HSC emergence from the dorsal aorta HE (Thambyrajah et al., 2024).

Figure 4 Supplement 2 – Detection of engraftment potential of late-stage haemGx in adrenal glands of Nude mice. (A) Map of the *Rosa26-BFP* construct used to target *Flk1-GFP* mES cells employed in implant assays. Primers used for gDNA PCR analysis are indicated. **(B)** PCR detection limit of the BFP amplicon using the primers in (A), in a serial dilution of *Rosa26-BFP::Flk1-GFP* mES gDNA into human leukemia cell line K562 gDNA; total 100ng DNA per lane. **(C)** Uncropped image of agarose gel electrophoresis of BFP detection shown in Fig. 4C. **(D)** Representative flow cytometry plot of the spleen of immunodeficient mouse recipients, showing myelo-lymphoid (CD45$^+$Ter119$^-$) and erythroid (Ter119$^+$CD45$^{+/-}$) cell fractions at 4 weeks post implantation. **(E)** PCR analysis of *Rosa26-BFP::Flk1-GFP* gDNA in the cell fractions in (D) of implanted animal #15 at 4 weeks post-implantation. PCR run in triplicate material from each sample; positive and NTC controls as per Fig. 4C. **(F-G)** Representative flow cytometry interrogation of BFP detection within lineage compartments in the bone marrow (BM) of control and implanted recipient animal #2 (F) and animal #1 (G) at 8 weeks after hGx implantation. **(H)** Histology of implanted and control adrenal glands; paraffin sections stained with hematoxylin & eosin (H&E). Inserts in implanted animal highlight neural (orange) and hematopoietic and renal tubular epithelium (green); inserts in control animal highlight normal cortical (blue) and medullar (red) adrenal architecture, absent in implanted adrenal glands.

Figure 5 – HaemGx with *MNX1* overexpression have increased cellularity and enhanced hemogenic endothelial potential. (A) Cell type enrichment analysis in MNX1-r AML transcriptomes, compared to normal paediatric bone marrow (BM) or other paediatric AML from the TARGET database. GSEA used representative cell type gene sets from the 2021 DB database; bubble plot shows NES scores (color gradient) and statistical significance (–log10(FDR), bubble size). **(B)** Schematic representation of generation of *MNX1-*overexpressing and empty vector (EV) mES cells by lentiviral transduction, with subsequent assembly into haemGxs. **(C)** Imaging of haemGxs with *MNX1* overexpression and EV controls at 10x magnification, showing appropriate assembly and polarization of the Flk1-GFP marker. scale bar: 300mm. **(D)** Size of haemGx at each timepoint, determined by the distance between the furthest extreme points in μm. Mean ± SD of 3 replicate experiments; 2-tailed t-test, p< 0.05 (*), 0.001 (**), 0.0001 (***), and 0.00001 (****). **(E-G)** Flow cytometry timecourse analysis of **(E)** C-Kit$^+$, **(F)** CD41$^+$, and **(G)** CD45$^+$ cell abundance in MNX1 and EV haemGxs (120-216h). Mean ± standard deviation of 3-7 independent experiments; 2-way ANOVA and Sidak's multiple comparison test significant at p<0.05 for C-Kit$^+$ cells only (construct contribution to variance p=0.0191; 144h comparison *p=0.0190). **(H)** Volcano plots of differentially expressed genes (DEGs) by bulk RNA-seq between MNX1 and EV haemGxs at 144h and 216h. Intersection of statistically significant DEGs between 144 h and 216 h (shown in Venn diagram

on top right) identified unique DEGs for each condition (labelled in red for 144 h and blue for
216 h). **(J)** Gene Ontology (GO) term enrichment for differentially upregulated genes between
MNX1 and EV haemGx at 144h (red) and 216h (blue), computed in EnrichR using the GO
Biological Process repository. **(I)** Transcription factor (TF) binding site enrichment on
differentially upregulated genes between MNX1 and EV haemGx at 144h (red) and 216 hr
(blue) using the ENCODE and ChEA Consensus TFs from ChIP database on EnrichR.

1222

Figure 5 Supplement 1 – MNX1 overexpression promotes hemogenic specification in
haemGx. **(A)** Cell type enrichment analysis in MNX1-r AML transcriptomes, compared to
normal paediatric bone marrow (BM) or other paediatric AML from the TARGET database.
GSEA used representative cell type gene sets from the 2021 DB database; bubble plot shows
NES scores (color gradient) and statistical significance (–log10(FDR), bubble size). **(B)**
Quantitative (q)RT-PCR analysis of *MNX1* overexpression in 216h-haemGx. Gene expression
fold change calculated by normalization to HPRT1. Mean ± SD of 3 replicates; 2-tailed t-test,
p< 0.001 (**). **(C)** Cell counts of disassembled haemGx at 216h. Mean ± SD of 3 replicate
experiments; 2-tailed t-test, p<0.001 (**). **(D)** Proportion of EV and MNX1 haemGx exhibiting
spontaneous unifocal contractility at 192h, observed in commercial N2B27 medium (see
Experimental Procedures); mean ± SD of 3 replicate experiments; 2-tailed t-test, p<0.001 (**).
(E-F) Flow cytometry analysis of specification of **(E)** hemato-endothelial – C-Kit[+] and VE-
Cadherin[+] – and **(F)** hematopoietic progenitor – CD41[+] and CD45[+] – cells over a 120-216h
timecourse of EV and MNX1 haemGx cultures; representative plots.

1237

Figure 5 Supplement 2 – Transcriptional analysis of haemGx with MNX1
overexpression. (A) Expression of human *MNX1* and murine *Mnx1* genes in FPKM units
from RNA-seq of MNX1 and EV haemGx. **(B)** UMAP of time-resolved global clustering of
scRNA-seq of haemGx showing the enriched clusters in GX-MNX1 at 144 hr (orange boxes)
and 216 hr (blue boxes), and MNX1-*r* AML (black boxes) or MLL-AML (green boxes)
determined by GSEA. **(C)** Bubble plot of GSEA NES values and statistical significance by –
log10(FDR) of enrichments in specific time clusters and corresponding global clusters in GX-
MNX1 at 144 hr and 216 hr compared to respective GX-EV, and MNX1-*r* or MLL AML samples
compared to other pediatric AML.

1247

1248   **Figure 6 – MNX1 selects C-Kit[+] clonogenic cells and selectively transforms end-stage**
1249   **haemGx. (A)** Representative photographs of serial replating of colony-forming cell assays
1250   initiated from EV or MNX1 cells obtained at 144h of the haemGx protocol. **(B)** Quantification
1251   of colony-replating efficiency of EV and MNX1 144h-haemGx (GX) cells. Mean+SD of n>3
1252   replicates; Kruskal-Wallis with Dunn's multiple comparison testing at significant q<0.05. **(C)**

Representative photographs of serial replating of colony-forming cell assays initiated from EV or MNX1 cells obtained at 216h of the haemGx protocol. **(D)** Quantification of colony-replating efficiency of EV and MNX1 216h-haemGx cells. Mean+SD of n=5-8 replicates; Kruskal-Wallis with Dunn's multiple comparison testing at significant q<0.05. **(E)** Flow cytometry analysis of hemogenic progenitor C-Kit+ cells and myeloid-affiliated CD11b+ cells at first round of plating of 144h and 216h-haemGx initiated from EV and MNX1 cells; representative plots.

Figure 6 Supplement 1 – Characterization of MNX1-overexpressing replating cells from haemGx. (A) Representative flow cytometry plot of CD31 and C-Kit staining of CFC plating of GX-MNX1 (left) and quantification of sustained expression of CD31+CKit+ cells through replatings. **(B)** Representative flow cytometry plots of serially re-plating MNX1 cells from 144h and 216h-gastruloids (plate 5 cells) stained with Ter119 and CD45.

Figure 7 – HaemGx with *MNX1* overexpression recapitulate MNX1-Acute Myeloid Leukemia patient signatures. (A) Quantification of human *MNX1* transcripts in FPKM units from RNA-seq of GX-EV, GX-MNX1, and CFC at 144 hr and 216 hr. Bars represent mean of replicates. **(B)** Heatmap comparing the expression of all differentially expressed genes (DEGs) between all conditions (GX-MNX1 144 hr vs GX-EV 144 hr; GX-MNX1 216 hr vs GX-EV 216 hr; CFC MNX1 144 hr vs GX MNX1 144 hr; CFC MNX1 216 hr vs GX MNX1 216 hr) as Z score. Hierarchical clustering by Ward D method on Euclidean distances identifies 14 clusters (k). **(C)** GSEA plots for gene set extracted from k14 from **(B)** against MNX1-r patients RNA-seq counts vs MLL, core-binding factors (CBF), or other pediatric AML. **(D)** Cell type analysis of the intersect of leading edge genes LEGs (n=476) from Fig. S5D in MNX1-r patients (k14 LEGs) **(B)** using the Panglao DB 2021 database. **(E)** Representative Giemsa-Wright stained dissociated CFC replating (4th plating) MNX1 haemGx cells from 144h and 216h. Solid black arrows, mast cell precursors; dashed black arrows, mast cells.

Figure 7 Supplement 1 – Transcriptional analyses of haemGx with *MNX1* overexpression compared to patient signatures and current *in vivo* model. (A) Venn diagram showing the intersection of leading-edge genes enriched in *MNX1*-r patient vs MLL, core-binding factors (CBF), or other pediatric AML from GSEA analysis using the K14 signature. **(B)** Cell type enrichment analysis in *MNX1*-r AML transcriptomes, compared to normal paediatric bone marrow (BM) or other paediatric AML from the TARGET database, with comparison with transcriptomes of haemGx and CFCs at from MNX1 vs EV conditions at 144h and 216h. GSEA used representative cell type gene sets from the Panglao DB 2021database; bubble plot shows NES scores (color gradient) and statistical significance (–log10(FDR), bubble size). **(C)** Representative Giemsa-Wright stained dissociated CFC replating MNX1

haemGx cells from 144h; purple arrows, undifferentiated blasts; solid black arrow, mast cell

precursors; dashed black arrow, mast cells. **(D)** Transcriptional re-analysis of Waraky et al

(2024) in vivo model of MNX1-OE leukemia from transplanted fetal liver (FL) cells. Heatmap

comparing the expression levels of all DEG between all conditions, i.e. FL control (FL Ctrl) vs

FL with MNX1-OE (FL MNX1) in liquid culture, and bone marrow (BM) from mice transplanted

with FL Ctrl (BM TX Ctrl) vs leukemic animals transplanted with FL MNX1 cells (BM TX MNX1).

Hierarchical clustering by Ward D method on Euclidean distances identifies 12 clusters (k).

Clusters k1, k2, k5, k6 and k9 are specifically up-regulated in BM TX MNX1 and putatively

represent MNX1-OE leukemia programs. Analysis matches Fig. 7B for MNX1-OE haemGx.

(E) GSEA plots of significant results (by p value < 0.05) for gene set extracted from k1, k2, k5,

k6 and k9, individually or combined, from (D) against MNX1-r patients RNA-seq counts vs

MLL, core-binding factors (CBF), or other pediatric AML. FDR values are indicated for each

enrichment point. Analysis matches Fig. 7C for MNX1-OE haemGx. **(F)** Cell type enrichment

analysis in MNX1-r AML transcriptomes, compared to normal pediatric bone marrow (BM) or

other pediatric AML from the TARGET database. GSEA used representative cell type gene

sets from the Panglao DB 2021 database; bubble plot shows NES scores (color gradient) and

statistical significance (–log10(FDR), bubble size). Analysis matches Fig. S6E for MNX1-OE

haemGx.

1308

1309

1310

REFERENCES

Alexander, T.B. and Mullighan, C.G. (2021) 'Molecular biology of childhood leukemia', *Annual Review of Cancer Biology,* 5, pp. 95-117.

Ballabio, E., Cantarella, C.D., Federico, C., Di Mare, P., Hall, G., Harbott, J., Hughes, J., Saccone, S. and Tosi, S. (2009) 'Ectopic expression of the HLXB9 gene is associated with an altered nuclear position in t (7; 12) leukaemias', *Leukemia,* 23(6), pp. 1179-1183.

Beccari, L., Moris, N., Girgin, M., Turner, D.A., Baillie-Johnson, P., Cossy, A., Lutolf, M.P., Duboule, D. and Arias, A.M. (2018) 'Multi-axial self-organization properties of mouse embryonic stem cells into gastruloids', *Nature,* 562(7726), pp. 272-276.

Boisset, J., van Cappellen, W., Andrieu-Soler, C., Galjart, N., Dzierzak, E. and Robin, C. (2010) 'In vivo imaging of haematopoietic cells emerging from the mouse aortic endothelium', *Nature,* 464(7285), pp. 116-120.

Bolouri, H., Farrar, J.E., Triche Jr, T., Ries, R.E., Lim, E.L., Alonzo, T.A., Ma, Y., Moore, R., Mungall, A.J. and Marra, M.A. (2018) 'The molecular landscape of pediatric acute myeloid leukemia reveals recurrent structural alterations and age-specific mutational interactions', *Nature medicine,* 24(1), pp. 103.

Bousquets-Muñoz, P., Molina, O., Varela, I., Álvarez-Eguiluz, Á, Fernández-Mateos, J., Gómez, A., Sánchez, E.G., Balbín, M., Ruano, D. and Ramírez-Orellana, M. (2024) 'Backtracking NOM1:: ETV6 fusion to neonatal pathogenesis of t (7; 12)(q36; p13) infant AML', *Leukemia,* , pp. 1-5.

Bowie, M.B., McKnight, K.D., Kent, D.G., McCaffrey, L., Hoodless, P.A. and Eaves, C.J. (2006) 'Hematopoietic stem cells proliferate until after birth and show a reversible phase-specific engraftment defect', *The Journal of clinical investigation,* 116(10), pp. 2808-2816.

Brignole, C., Calarco, E., Bensa, V., Giusto, E., Perri, P., Ciampi, E., Corrias, M.V., Astigiano, S., Cilli, M. and Loo, D. (2023) 'Antitumor activity of the investigational B7-H3 antibody-drug conjugate, vobramitamab duocarmazine, in preclinical models of neuroblastoma', *Journal for Immunotherapy of Cancer,* 11(9).

Calvanese, V., Nguyen, A.T., Bolan, T.J., Vavilina, A., Su, T., Lee, L.K., Wang, Y., Lay, F.D., Magnusson, M. and Crooks, G.M. (2019) 'MLLT3 governs human haematopoietic stem-cell self-renewal and engraftment', *Nature,* 576(7786), pp. 281-286.

Cazzola, A., Cazzaniga, G., Biondi, A., Meneveri, R., Brunelli, S. and Azzoni, E. (2021) 'Prenatal origin of pediatric leukemia: lessons from hematopoietic development', *Frontiers in cell and developmental biology,* 8, pp. 618164.

Chandrakanthan, V., Rorimpandey, P., Zanini, F., Chacon, D., Olivier, J., Joshi, S., Kang, Y.C., Knezevic, K., Huang, Y. and Qiao, Q. (2022) 'Mesoderm-derived PDGFRA cells regulate the emergence of hematopoietic stem cells in the dorsal aorta', *Nature cell biology,* 24(8), pp. 1211-1225.

Chang, Y., Hummel, S.N., Jung, J., Jin, G., Deng, Q. and Bao, X. (2023) 'Engineered
hematopoietic and immune cells derived from human pluripotent stem cells',
Experimental hematology, .

Chaudhury, S., O'Connor, C., Cañete, A., Bittencourt-Silvestre, J., Sarrou, E., Prendergast,
Á, Choi, J., Johnston, P., Wells, C.A. and Gibson, B. (2018) 'Age-specific biological
and molecular profiling distinguishes paediatric from adult acute myeloid leukaemias',
Nature communications, 9(1), pp. 5280.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V. and Clark, N.R. (2013)
'A. 1126 Ma'ayan, Enrichr: Interactive and collaborative HTML5 gene list enrichment
analysis tool. BMC 1127', *Bioinformatics,* .

Chen, W., O'Sullivan, M.G., Hudson, W. and Kersey, J. (2011) 'Modeling human infant MLL
leukemia in mice: leukemia from fetal liver differs from that originating in postnatal
marrow', *Blood,* 117(12), pp. 3474.

Chia, S.L., Kapoor, S., Carvalho, C., Bajénoff, M. and Gentek, R. (2023) 'Mast cell ontogeny:
From fetal development to life‐long health and disease', *Immunological reviews,*
315(1), pp. 31-53.

Chun, I.Y., Maser, R., Marches, F., Banchereau, J. and Palucka, K. (2024) 'Engraftment of
adult hematopoietic stem and progenitor cells in a novel model of humanized
mice' *Iscience*, 27(3).

Ciau-Uitz, A., Pinheiro, P., Gupta, R., Enver, T. and Patient, R. (2010) 'Tel1/ETV6 specifies
blood stem cells through the agency of VEGF signaling', *Developmental cell,* 18(4),
pp. 569-578.

Ciriza, J., Thompson, H., Petrosian, R., Manilay, J.O. and García-Ojeda, M.E. (2013) 'The
migration of hematopoietic progenitors from the fetal liver to the fetal bone marrow:
lessons learned and possible clinical applications', *Experimental hematology,* 41(5),
pp. 411-423.

Costa, G., Kouskoff, V. and Lacaud, G. (2012) 'Origin of blood cells and HSC production in
the embryo', *Trends in immunology,* 33(5), pp. 215-223.

Dang, S.M., Kyba, M., Perlingeiro, R., Daley, G.Q. and Zandstra, P.W. (2002) 'Efficiency of
embryoid body formation and hematopoietic development from embryonic stem cells
in different culture systems', *Biotechnology and bioengineering,* 78(4), pp. 442-453.

Deguchi, Y. and Kehrl, J.H. (1991) 'Selective expression of two homeobox genes in CD34-
positive cells from human bone marrow', *Blood,* 78(2), pp. 323-328.

Dias, A., Pascual-Mas, P., Torregrosa-Cortés, G., McNamara, H.M., Wehmeyer, A.E., Arnold,
S.J., Martinez Arias A. (2025) 'A temporal coordination between Nodal and Wnt
signalling governs the emergence of the mammalian body plan' *bioRxiv*
2025.01.11.632562; doi: https://doi.org/10.1101/2025.01.11.632562

Diaz, M.F., Li, N., Lee, H.J., Adamo, L., Evans, S.M., Willey, H.E., Arora, N., Torisawa, Y.,
Vickers, D.A. and Morris, S.A. (2015) 'Biomechanical forces promote blood
development through prostaglandin E2 and the cAMP–PKA signaling axis', *Journal of
Experimental Medicine,* 212(5), pp. 665-680.

Dijkhuis, L., Johns, A., Ragusa, D., van den Brink, Susanne C and Pina, C. (2023)
'Haematopoietic development and HSC formation in vitro: promise and limitations of
gastruloid models', *Emerging Topics in Life Sciences,* 7(4), pp. 439-454.

Dong, Y., Shi, O., Zeng, Q., Lu, X., Wang, W., Li, Y. and Wang, Q. (2020) 'Leukemia incidence
trends at the global, regional, and national level between 1990 and 2017', *Experimental*
hematology & oncology, 9, pp. 1-11.

Dzierzak, E. and Bigas, A. (2018) 'Blood development: hematopoietic stem cell dependence
and independence', *Cell stem cell,* 22(5), pp. 639-651.

Espersen, A.D.L., Noren‐Nyström, U., Abrahamsson, J., Ha, S., Pronk, C.J., Jahnukainen,
K., Jónsson, ÓG., Lausen, B., Palle, J. and Zeller, B. (2018) 'Acute myeloid leukemia
(AML) with t (7; 12)(q36; p13) is associated with infancy and trisomy 19: Data from
Nordic Society for Pediatric Hematology and Oncology (NOPHO‐AML) and review of
the literature', *Genes, Chromosomes and Cancer,* 57(7), pp. 359-365.

Fadlullah, M.Z.H., Neo, W.H., Lie-A-Ling, M., Thambyrajah, R., Patel, R., Mevel, R., Aksoy, I.,
Do Khoa, N., Savatier, P., Fontenille, L. and Baker, S.M. (2022) 'Murine AGM single-
cell profiling identifies a continuum of hemogenic endothelium differentiation marked
by ACE' *Blood, 139*(3), pp.343-356.

Fehling, H.J., Lacaud, G., Kubo, A., Kennedy, M., Robertson, S., Keller, G. and Kouskoff, V.
(2003) 'Tracking mesoderm induction and its specification to the hemangioblast during
embryonic stem cell differentiation', .

Fidanza, A. and Forrester, L.M. (2021) 'Progress in the production of haematopoietic stem
and progenitor cells from human pluripotent stem cells', *Journal of Immunology and*
Regenerative Medicine, 13, pp. 100050.

Fitch, S.R., Kimber, G.M., Wilson, N.K., Parker, A., Mirshekar-Syahkal, B., Göttgens, B.,
Medvinsky, A., Dzierzak, E. and Ottersbach, K. (2012) 'Signaling from the sympathetic
nervous system regulates hematopoietic stem cell emergence during embryogenesis',
Cell stem cell, 11(4), pp. 554-566.

Fowler, J.L., Zheng, S.L., Nguyen, A., Chen, A., Xiong, X., Chai, T., Chen, J.Y., Karigane, D.,
Banuelos, A.M. and Niizuma, K. (2024) 'Lineage-tracing hematopoietic stem cell
origins in vivo to efficiently make human HLF HOXA hematopoietic progenitors from
pluripotent stem cells', *Developmental Cell,* 59(9), pp. 1110-1131. e22.

Frankish, A., Diekhans, M., Ferreira, A., Johnson, R., Jungreis, I., Loveland, J., Mudge, J.M.,
Sisu, C., Wright, J. and Armstrong, J. (2019) 'GENCODE reference annotation for the
human and mouse genomes', *Nucleic acids research,* 47(D1), pp. D766-D773.

Franzén, O., Gan, L. and Björkegren, J.L. (2019) 'PanglaoDB: a web server for exploration of
mouse and human single-cell RNA sequencing data', *Database,* 2019, pp. baz046.

Fujimoto, T., Ogawa, M., Minegishi, N., Yoshida, H., Yokomizo, T., Yamamoto, M. and
Nishikawa, S. (2001) 'Step‐wise divergence of primitive and definitive haematopoietic
and endothelial cell lineages during embryonic stem cell differentiation', *Genes to*
Cells, 6(12), pp. 1113-1127.

Ganuza, M., Hall, T., Myers, J., Nevitt, C., Sánchez-Lanzas, R., Chabot, A., Ding, J., Kooienga, E., Caprio, C. and Finkelstein, D. (2022) 'Murine foetal liver supports limited detectable expansion of life-long haematopoietic progenitors', *Nature cell biology,* 24(10), pp. 1475-1486.

Ghiaur, G., Ferkowicz, M.J., Milsom, M.D., Bailey, J., Witte, D., Cancelas, J.A., Yoder, M.C. and Williams, D.A. (2008) 'Rac1 is essential for intraembryonic hematopoiesis and for the initial seeding of fetal liver with definitive hematopoietic progenitor cells', *Blood, The Journal of the American Society of Hematology,* 111(7), pp. 3313-3321.

Harrison, K.A., Thaler, J., Pfaff, S.L., Gu, H. and Kehrl, J.H. (1999) 'Pancreas dorsal lobe agenesis and abnormal islets of Langerhans in Hlxb9-deficient mice', *Nature genetics,* 23(1), pp. 71-75.

Hooper, M., Hardy, K., Handyside, A., Hunter, S. and Monk, M. (1987) 'HPRT-deficient (Lesch–Nyhan) mouse embryos derived from germline colonization by cultured cells', *Nature,* 326(6110), pp. 292-295.

Hou, S., Li, Z., Zheng, X., Gao, Y., Dong, J., Ni, Y., Wang, X., Li, Y., Ding, X., Chang, Z. and Li, S. (2020) 'Embryonic endothelial evolution towards first hematopoietic stem cells revealed by single-cell transcriptomic and functional analyses' *Cell research*, 30(5), pp.376-392.

Huang, Y., Lu, Y., He, Y., Feng, Z., Zhan, Y., Huang, X., Liu, Q., Zhang, J., Li, H., Huang, H. and Ma, M. (2019) 'Ikzf1 regulates embryonic T lymphopoiesis via Ccr9 and Irf4 in zebrafish' *Journal of Biological Chemistry*, *294*(44), pp.16152-16163.

Ivanovs, A., Rybtsov, S., Welch, L., Anderson, R.A., Turner, M.L. and Medvinsky, A. (2011) 'Highly potent human hematopoietic stem cells first emerge in the intraembryonic aorta-gonad-mesonephros region', *Journal of Experimental Medicine,* 208(12), pp. 2417-2427.

Jakobsson, L., Franco, C.A., Bentley, K., Collins, R.T., Ponsioen, B., Aspalter, I.M., Rosewell, I., Busse, M., Thurston, G. and Medvinsky, A. (2010) 'Endothelial cells dynamically compete for the tip cell position during angiogenic sprouting', *Nature cell biology,* 12(10), pp. 943-953.

Jung, H.S., Uenishi, G., Park, M.A., Liu, P., Suknuntha, K., Raymond, M., Choi, Y.J., Thomson, J.A., Ong, I.M. and Slukvin, I.I. (2021) 'SOX17 integrates HOXA and arterial programs in hemogenic endothelium to drive definitive lympho-myeloid hematopoiesis' *Cell reports*, *34*(7).

Kapeni, C., Nitsche, L., Kilpatrick, A.M., Wilson, N.K., Xia, K., Mirshekar-Syahkal, B., Chandrakanthan, V., Malouf, C., Pimanda, J.E. and Göttgens, B. (2022) 'p57Kip2 regulates embryonic blood stem cells by controlling sympathoadrenal progenitor expansion', *Blood, The Journal of the American Society of Hematology,* 140(5), pp. 464-477.

Kikuchi, K. and Kondo, M. (2006) 'Developmental switch of mouse hematopoietic stem cells from fetal to adult type occurs in bone marrow after birth', *Proceedings of the National Academy of Sciences,* 103(47), pp. 17852-17857.

Kiselev, V.Y., Yiu, A. and Hemberg, M. (2018) 'scmap: projection of single-cell RNA-seq data across data sets', *Nature methods,* 15(5), pp. 359-362.

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S.L., Jagodnik, K.M. and Lachmann, A. (2016) 'Enrichr: a comprehensive gene set enrichment analysis web server 2016 update', *Nucleic acids research,* 44(W1), pp. W90-W97.

Lacaud, G. and Kouskoff, V. (2017) 'Hemangioblast, hemogenic endothelium, and primitive versus definitive hematopoiesis', *Experimental hematology,* 49, pp. 19-24.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R. and 1000 Genome Project Data Processing Subgroup (2009) 'The sequence alignment/map format and SAMtools', *bioinformatics,* 25(16), pp. 2078-2079.

Li, W., Ferkowicz, M.J., Johnson, S.A., Shelley, W.C. and Yoder, M.C. (2005) 'Endothelial cells in the early murine yolk sac give rise to CD41-expressing hematopoietic cells', *Stem cells and development,* 14(1), pp. 44-54.

Lis, R., Karrasch, C.C., Poulos, M.G., Kunar, B., Redmond, D., Duran, J.G.B., Badwe, C.R., Schachterle, W., Ginsberg, M. and Xiang, J. (2017) 'Conversion of adult endothelium to immunocompetent haematopoietic stem cells', *Nature,* 545(7655), pp. 439-445.

Love, M., Huber, W. and Anders, S. (2014) 'No title', *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2 Genome Biology 15 (12), 550,* .

Lundin, V., Sugden, W.W., Theodore, L.N., Sousa, P.M., Han, A., Chou, S., Wrighton, P.J., Cox, A.G., Ingber, D.E. and Goessling, W. (2020) 'YAP regulates hematopoietic stem cell formation in response to the biomechanical forces of blood flow', *Developmental cell,* 52(4), pp. 446-460. e5.

Masetti, R., Vendemini, F., Zama, D., Biagi, C., Pession, A. and Locatelli, F. (2015) 'Acute myeloid leukemia in infants: biology and treatment', *Frontiers in pediatrics,* 3, pp. 37.

McGrath, K.E. and Palis, J. (2005) 'Hematopoiesis in the yolk sac: more than meets the eye', *Experimental hematology,* 33(9), pp. 1021-1028.

McGrath, K.E., Frame, J.M., Fegan, K.H., Bowen, J.R., Conway, S.J., Catherman, S.C., Kingsley, P.D., Koniski, A.D. and Palis, J. (2015) 'Distinct sources of hematopoietic progenitors emerge before HSCs and provide functional blood cells in the mammalian embryo', *Cell reports,* 11(12), pp. 1892-1904.

Medvinsky, A. and Dzierzak, E. (1996) 'Definitive hematopoiesis is autonomously initiated by the AGM region', *Cell,* 86(6), pp. 897-906.

Medvinsky, A., Rybtsov, S. and Taoudi, S. (2011) 'Embryonic origin of the adult hematopoietic system: advances and questions', *Development,* 138(6), pp. 1017-1031.

Mercher, T., Raffel, G.D., Moore, S.A., Cornejo, M.G., Baudry-Bluteau, D., Cagnard, N., Jesneck, J.L., Pikman, Y., Cullen, D. and Williams, I.R. (2009) 'The OTT-MAL fusion oncogene activates RBPJ-mediated transcription and induces acute megakaryoblastic leukemia in a knockin mouse model', *The Journal of clinical investigation,* 119(4), pp. 852-864.

Miladinovic, O., Canto, P., Pouget, C., Piau, O., Radic, N., Freschu, P., Megherbi, A., Brujas Prats, C., Jacques, S. and Hirsinger, E. (2024) 'A multistep computational approach reveals a neuro-mesenchymal cell population in the embryonic hematopoietic stem cell niche', *Development,* 151(7), pp. dev202614.

Moore, M.A. and Metcalf, D. (1970) 'Ontogeny of the haemopoietic system: yolk sac origin of in vivo and in vitro colony forming cells in the developing mouse embryo', *British journal of haematology,* 18(3), pp. 279-296.

Moris, N., Edri, S., Seyres, D., Kulkarni, R., Domingues, A.F., Balayo, T., Frontini, M. and Pina, C. (2018) 'Histone acetyltransferase KAT2A stabilizes pluripotency with control of transcriptional heterogeneity', *Stem cells,* 36(12), pp. 1828-1838.

Murry, C.E. and Keller, G. (2008) 'Differentiation of embryonic stem cells to clinically relevant populations: lessons from embryonic development', *Cell,* 132(4), pp. 661-680.

Nafria, M., Bonifer, C., Stanley, E.G., Ng, E.S. and Elefanty, A.G. (2020) 'Protocol for the generation of definitive hematopoietic progenitors from human pluripotent stem cells', *STAR protocols,* 1(3), pp. 100130.

Nagel, S., Scherr, M., Quentmeier, H., Kaufmann, M., Zaborski, M., Drexler, H.G. and MacLeod, R. (2005) 'HLXB9 activates IL6 in Hodgkin lymphoma cell lines and is regulated by PI3K signalling involving E2F3', *Leukemia,* 19(5), pp. 841.

Neo, W.H., Booth, C.A., Azzoni, E., Chi, L., Delgado-Olguín, P., de Bruijn, M.F., Jacobsen, S.E.W. and Mead, A.J. (2018) 'Cell-extrinsic hematopoietic impact of Ezh2 inactivation in fetal liver endothelial cells' *Blood*, 131(20), pp.2223-2234.

Neo, W.H., Meng, Y., Rodriguez-Meira, A., Fadlullah, M.Z., Booth, C.A., Azzoni, E., Thongjuea, S., de Bruijn, M.F., Jacobsen, S.E.W., Mead, A.J. and Lacaud, G. (2021) 'Ezh2 is essential for the generation of functional yolk sac derived erythro-myeloid progenitors' *Nature Communications*, *12*(1), p.7019.

Ng, E.S., Sarila, G., Li, J.Y., Edirisinghe, H.S., Saxena, R., Sun, S., Bruveris, F.F., Labonne, T., Sleebs, N., Maytum, A. and Yow, R.Y. (2024) 'Long-term engrafting multilineage hematopoietic cells differentiated from human induced pluripotent stem cells' *Nature biotechnology*, pp.1-14.

Niakan, K.K., Ji, H., Maehr, R., Vokes, S.A., Rodolfa, K.T., Sherwood, R.I., Yamaki, M., Dimos, J.T., Chen, A.E. and Melton, D.A. (2010) 'Sox17 promotes differentiation in mouse embryonic stem cells by directly regulating extraembryonic gene expression and indirectly antagonizing self-renewal', *Genes & development,* 24(3), pp. 312-326.

North, T.E., Goessling, W., Peeters, M., Li, P., Ceol, C., Lord, A.M., Weber, G.J., Harris, J., Cutting, C.C. and Huang, P. (2009) 'Hematopoietic stem cell development is dependent on blood flow', *Cell,* 137(4), pp. 736-748.

Oatley, M., Bölükbası, Ö.V., Svensson, V., Shvartsman, M., Ganter, K., Zirngibl, K., Pavlovich, P.V., Milchevskaya, V., Foteva, V., Natarajan, K.N. and Baying, B. (2020) 'Single-cell transcriptomics identifies CD44 as a marker and regulator of endothelial to haematopoietic transition' *Nature communications*, *11*(1), p.586.

Palis, J., Robertson, S., Kennedy, M., Wall, C. and Keller, G. (1999) 'Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse', *Development,* 126(22), pp. 5073-5084.

Pastorino, F., Brignole, C., Marimpietri, D., Cilli, M., Gambini, C., Ribatti, D., Longhi, R., Allen, T.M., Corti, A. and Ponzoni, M. (2003) 'Vascular damage and anti-angiogenic effects of tumor vessel-targeted liposomal chemotherapy', *Cancer research,* 63(21), pp. 7400-7409.

Pearson, S., Cuvertino, S., Fleury, M., Lacaud, G. and Kouskoff, V. (2015) 'In vivo repopulating activity emerges at the onset of hematopoietic specification during embryonic stem cell differentiation', *Stem Cell Reports,* 4(3), pp. 431-444.

Piau, O., Brunet-Manquat, M., L'homme, B., Petit, L., Birebent, B., Linard, C., Moeckes, L., Zuliani, T., Lapillonne, H. and Benderitter, M. (2023) 'Generation of transgene-free hematopoietic stem cells from human induced pluripotent stem cells', *Cell Stem Cell,* 30(12), pp. 1610-1623. e7.

Picelli, S., Faridani, O.R., Björklund, ÅK., Winberg, G., Sagasser, S. and Sandberg, R. (2014) 'Full-length RNA-seq from single cells using Smart-seq2', *Nature protocols,* 9(1), pp. 171-181.

Pina, C., May, G., Soneji, S., Hong, D. and Enver, T. (2008) 'MLLT3 regulates early human erythroid and megakaryocytic cell fate', *Cell stem cell,* 2(3), pp. 264-273.

Quintard, C., Tubbs, E., Jonsson, G., Jiao, J., Wang, J., Werschler, N., Laporte, C., Pitaval, A., Bah, T. and Pomeranz, G. (2024) 'A microfluidic platform integrating functional vascularized organoids-on-chip', *Nature Communications,* 15(1), pp. 1452.

Ragusa, D., Ciciro, Y., Federico, C., Saccone, S., Bruno, F., Saeedi, R., Sisu, C., Pina, C., Sala, A. and Tosi, S. (2022) 'Engineered model of t (7; 12)(q36; p13) AML recapitulates patient-specific features and gene expression profiles', *Oncogenesis,* 11(1), pp. 50.

Ragusa, D., Dijkhuis, L., Pina, C. and Tosi, S. (2023) 'Mechanisms associated with t (7; 12) acute myeloid leukaemia: from genetics to potential treatment targets', *Bioscience reports,* 43(1), pp. BSR20220489.

Rossi, G., Broguiere, N., Miyamoto, M., Boni, A., Guiet, R., Girgin, M., Kelly, R.G., Kwon, C. and Lutolf, M.P. (2021) 'Capturing cardiogenesis in gastruloids', *Cell stem cell,* 28(2), pp. 230-240. e6.

Rossi, G., Giger, S., Hübscher, T. and Lutolf, M.P. (2022) 'Gastruloids as in vitro models of embryonic blood development with spatial and temporal resolution', *Scientific Reports,* 12(1), pp. 13380.

Rybtsov, S., Batsivari, A., Bilotkach, K., Paruzina, D., Senserrich, J., Nerushev, O. and Medvinsky, A. (2014) 'Tracing the origin of the HSC hierarchy reveals an SCF-dependent, IL-3-independent CD43− embryonic precursor', *Stem cell reports,* 3(3), pp. 489-501.

Saorin, G., Caligiuri, I. and Rizzolio, F. (2023) *Microfluidic organoids-on-a-chip: The future of human models.* Elsevier, pp. 41.

Schyrr, F., Alonso-Calleja, A., Vijaykumar, A., Gebhard, S., Sarkis, R., Lopes, S.F., Oggier, A., De Leval, L., Nombela-Arrieta, C. and Naveiras, O. (2023) 'Adrenal extramedullary hematopoiesis as an inducible model of the adult hematopoietic niche', *bioRxiv, ,* pp. 2023.03. 15.531679.

Souilhol, C., Gonneau, C., Lendinez, J.G., Batsivari, A., Rybtsov, S., Wilson, H., Morgado-Palacin, L., Hills, D., Taoudi, S. and Antonchuk, J. (2016) 'Inductive interactions mediated by interplay of asymmetric signalling underlie development of adult haematopoietic stem cells', *Nature communications,* 7(1), pp. 10784.

Sroczynska, P., Lancrin, C., Pearson, S., Kouskoff, V. and Lacaud, G. (2009) 'In vitro differentiation of embryonic stem cells as a model of early hematopoietic development', *Leukemia: Methods and Protocols,* , pp. 317-334.

Steliarova-Foucher, E., Colombet, M., Ries, L.A., Moreno, F., Dolya, A., Bray, F., Hesseling, P., Shin, H.Y. and Stiller, C.A. (2017) 'International incidence of childhood cancer, 2001–10: a population-based registry study', *The Lancet Oncology,* 18(6), pp. 719-731.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R. and Lander, E.S. (2005) 'Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles', *Proceedings of the National Academy of Sciences,* 102(43), pp. 15545-15550.

Sugimura, R., Jha, D.K., Han, A., Soria-Valles, C., Da Rocha, E.L., Lu, Y., Goettel, J.A., Serrao, E., Rowe, R.G. and Malleshaiah, M. (2017) 'Haematopoietic stem and progenitor cells from human pluripotent stem cells', *Nature,* 545(7655), pp. 432-438.

Taketani, T., Taki, T., Sako, M., Ishii, T., Yamaguchi, S. and Hayashi, Y. (2008) 'MNX1–ETV6 fusion gene in an acute megakaryoblastic leukemia and expression of the MNX1 gene in leukemia and normal B cell lines', *Cancer genetics and cytogenetics,* 186(2), pp. 115-119.

Thaler, J., Harrison, K., Sharma, K., Lettieri, K., Kehrl, J. and Pfaff, S.L. (1999) 'Active suppression of interneuron programs within developing motor neurons revealed by analysis of homeodomain factor HB9', *Neuron,* 23(4), pp. 675-687.

Thambyrajah, R., Maqueda, M., Neo, W.H., Imbach, K., Guillén, Y., Grases, D., Fadlullah, Z., Gambera, S., Matteini, F. and Wang, X. (2024) 'Cis inhibition of NOTCH1 through JAGGED1 sustains embryonic hematopoietic stem cell fate', *Nature Communications,* 15(1), pp. 1604.

Tosi, S., Kamel, Y.M., Owoka, T., Federico, C., Truong, T.H. and Saccone, S. (2015) 'Paediatric acute myeloid leukaemia with the t (7; 12)(q36; p13) rearrangement: a review of the biological and clinical management aspects', *Biomarker research,* 3(1), pp. 1.

Turner, D.A., Girgin, M., Alonso-Crisostomo, L., Trivedi, V., Baillie-Johnson, P., Glodowski, C.R., Hayward, P.C., Collignon, J., Gustavsen, C. and Serup, P. (2017) 'Anteroposterior polarity and elongation in the absence of extra-embryonic tissues and of spatially localised signalling in gastruloids: mammalian embryonic organoids', *Development,* 144(21), pp. 3894-3906.

van den Brink, Susanne C and van Oudenaarden, A. (2021) '3D gastruloids: A novel frontier in stem cell-based in vitro modeling of mammalian gastrulation', *Trends in cell biology,* 31(9), pp. 747-759.

van den Brink, Susanne C, Alemany, A., van Batenburg, V., Moris, N., Blotenburg, M., Vivie, J., Baillie-Johnson, P., Nichols, J., Sonnen, K.F. and Martinez Arias, A. (2020) 'Single-cell and spatial transcriptomics reveal somitogenesis in gastruloids', *Nature,* 582(7812), pp. 405-409.

Van den Brink, Susanne C, Baillie-Johnson, P., Balayo, T., Hadjantonakis, A., Nowotschin, S., Turner, D.A. and Martinez Arias, A. (2014) 'Symmetry breaking, germ layer specification and axial organisation in aggregates of mouse embryonic stem cells', *Development,* 141(22), pp. 4231-4242.

Veenvliet, J.V., Bolondi, A., Kretzmer, H., Haut, L., Scholze-Wittler, M., Schifferl, D., Koch, F., Guignard, L., Kumar, A.S. and Pustet, M. (2020) 'Mouse embryonic stem cells self-organize into trunk-like structures with neural tube and somites', *Science,* 370(6522), pp. eaba4937.

Vo, L.T. and Daley, G.Q. (2015) 'De novo generation of HSCs from somatic and pluripotent stem cell sources', *Blood, The Journal of the American Society of Hematology,* 125(17), pp. 2641-2648.

Waraky, A., Östlund, A., Nilsson, T., Weichenhan, D., Lutsik, P., Bähr, M., Hey, J., Tunali, G., Adamsson, J. and Jacobsson, S. (2024) 'Aberrant MNX1 expression associated with t (7; 12)(q36; p13) pediatric acute myeloid leukemia induces the disease through altering histone methylation', *Haematologica,* 109(3), pp. 725.

Weichenhan, D., Riedel, A., Sollier, E., Toprak, U.H., Hey, J., Breuer, K., Wierzbinska, J.A., Touzart, A., Lutsik, P. and Bähr, M. (2023) 'Altered enhancer-promoter interaction leads to MNX1 expression in pediatric acute myeloid leukemia with t (7; 12)(q36; p13)', *bioRxiv,* , pp. 2023.09. 13.557546.

Wildenhain, S., Ingenhag, D., Ruckert, C., Degistirici, O., Dugas, M., Meisel, R., Hauer, J. and Borkhardt, A. (2012) 'Homeobox protein HB9 binds to the prostaglandin E receptor 2 promoter and inhibits intracellular cAMP mobilization in leukemic cells', *The Journal of biological chemistry,* 287(48), pp. 40703-40712 Available at: 10.1074/jbc.M111.308965 [doi].

Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J., Evangelista, J.E., Jenkins, S.L., Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E. and Jagodnik, K.M. (2021) 'Gene set knowledge discovery with Enrichr', *Current protocols,* 1(3), pp. e90.

Yokomizo, T., Ideue, T., Morino-Koga, S., Tham, C.Y., Sato, T., Takeda, N., Kubota, Y., Kurokawa, M., Komatsu, N. and Ogawa, M. (2022) 'Independent origins of fetal liver haematopoietic stem and progenitor cells', *Nature,* 609(7928), pp. 779-784.

Yoshimoto, M., Kosters, A., Cornelius, S., Valiente, N., Cheng, H., Latorre, A., Nishida, C., Ghosn, E.E. and Kobayashi, M. (2022) 'Mast cell repopulating ability is lost during the transition from pre-HSC to FL HSC', *Frontiers in Immunology,* 13, pp. 896396.

Zhu, Q., Gao, P., Tober, J., Bennett, L., Chen, C., Uzun, Y., Li, Y., Howell, E.D., Mumau, M., Yu, W. and He, B. (2020) 'Developmental trajectory of prehematopoietic stem cell formation from endothelium' *Blood*, 136(7), pp.845-856.

1679   Zovein, A.C., Hofmann, J.J., Lynch, M., French, W.J., Turlo, K.A., Yang, Y., Becker, M.S.,
1680   Zanetta, L., Dejana, E. and Gasson, J.C. (2008) 'Fate tracing reveals the endothelial
1681   origin of hematopoietic stem cells', *Cell stem cell,* 3(6), pp. 625-636.

1682