Title:

The deep-sea coral, *Callogorgia delta,* associates with bacteria belonging to a novel

marine branch of the Mollicutes.

Authors:

Samuel A. Vohsen, Harald R. Gruber-Vodicka, Martina Meyer, Miriam Sadowski,

Nicole Dubilier, Charles R. Fisher, Iliana B. Baums.

Abstract:

In addition to abundant animal communities, corals from all ocean depths support

diverse microbial associates that are important to coral health. While some of these

microbes have been classified taxonomically*,* understanding the metabolic potential of

coral-associated bacteria and how they interact with their coral hosts is limited by a lack

of genomic data. One example is *Mycoplasma* and other members of the class Mollicutes

which are widespread coral associates. Here we investigated the association between two

novel members of the class Mollicutes and the deep-sea octocoral *Callogorgia delta.* We

screened *C. delta,* a closely related species *C. americana,* sediment, and water for

mollicutes using 16S metabarcoding. One ASV was found in most colonies screened

(99/108) and often dominated the microbiome (up to 99%). Another ASV was detected at

lower abundance and prevalence in these corals. Both were absent in all water and were

absent or rare in the sediment. We sequenced metagenomes and metatranscriptomes to

assemble and annotate genomes and propose the names *Ca. Oceanoplasma callogorgiae*

and *Ca. Thalassoplasma callogorgiae.* The genomes were small, revealed a reliance on

the arginine dihydrolase pathway for ATP production, and contained CRISPR-Cas

systems with extensive arrays. CARD-FISH microscopy unveiled an abundant bacterium

in the mesoglea which is likely to be *Ca. O. callogorgiae*. These novel mollicutes cluster

with others from diverse invertebrate hosts. Altogether, this work describes the

association of these novel mollicutes in *C. delta,* provides insight into widespread coral

associates, and identifies a novel clade of marine mollicutes whose diversity remains

largely undiscovered.

Keywords:

Deep-sea corals, Mollicutes, symbiosis

Introduction:

Corals are important foundation species that support diverse animal communities

from shallow waters to the deep sea. In addition to abundant animal communities, corals

also associate with a diverse array of microbes (Bourne et al. 2016). Besides the well-

studied algal symbionts of the family Symbiodiniaceae, corals associate with a diverse

array of microbial taxa that have proven to be important to holobiont function by

performing a variety of roles including providing nitrogen through fixation (M. P. Lesser

et al. 2007; Michael P. Lesser et al. 2004; S. H. Yang et al. 2016) and causing disease

(Baker et al. 2022; J. G. Klinges et al. 2022; G. Klinges et al. 2020; J. G. Klinges et al.

2019). Some of the most widespread taxa commonly found in corals include corallicolid

apicomplexans (Vohsen et al. 2020; Kwong et al. 2019; Kirk et al. 2013a; Kirk et al.

2013b), *Endozoicomonas* (Pollock et al. 2018)*,* and *Mycoplasma*.

Mycoplasma and other members of the class Mollicutes are widespread coral

associates. While they are most commonly found in octocorals (Gray et al. 2011; Penn et

al. 2006; Weiler et al. 2018; Holm and Heidelberg 2016; Kellogg et al. 2016; Porporato et

al. 2013; Shirur et al. 2016; van de Water et al. 2018; van de Water et al. 2017; McCauley

et al. 2020), they have also been found in black corals (van de Water et al. 2020) and

stony corals (Neulinger et al. 2009; Kellogg et al. 2009). The corals that they inhabit are

found in a wide breadth of habitats from shallow-water reefs (Woo et al. 2017; Shirur et

al. 2016; McCauley et al. 2020), through the mesophotic (van de Water et al. 2020) and

the deep sea (Neulinger et al. 2009; Weiler et al. 2018; Penn et al. 2006) to abyssal depths

(Quintanilla et al. 2022). Beyond this, some studies have yielded further insight into the

association between mollicutes and corals. For instance, *Mycoplasma corallicola* resides

within its host coral's ectoderm in the deep-sea scleractinian, *Lophelia pertusa*

(Neulinger et al. 2009). Further, *Mycoplasma* spp. are abundant and dominant in multiple

coral species, comprising more than 50% of their associated microbial community

(Weiler et al. 2018; Woo et al. 2017; McCauley et al. 2020). Interestingly, the relative

abundance of *Mycoplasma* can show substantial variation across the ranges of their host

corals. In some coral species, colonies from only a subset of geographic locations are

dominated by *Mycoplasma* (Woo et al. 2017; van de Water et al. 2017). Thus, this

association may be influenced by environmental conditions or otherwise shaped by

geography such as through limited dispersal. *Mycoplasma* may have also coevolved with

their coral hosts because they exhibit varying degrees of host specificity. Closely related

octocoral species from the same locations differ in the abundance and identity of the

Mycoplasma phylotypes that they host (McCauley et al. 2020; van de Water et al. 2017;

Holm and Heidelberg 2016). Despite these insights, the interactions between

mycoplasmas and other mollicutes with their host corals remain unknown.

Almost all other members of the class Mollicutes are symbionts that associate

with a wide diversity of plant, fungal, and animal host species including humans (Ing-

Ming Lee et al. 2000; Tully et al. 1993). Many members are well-studied parasites

(Atkinson 2018), while some are unclear such as the ubiquitous endocellular symbionts

of arbuscular mycorrhizal fungi, an ancient association that dates back to the era when

plants first colonized the land (Bonfante and Desirò 2017; Desiro et al. 2015; Kuga et al.

2021). Yet others are mutualists with well-established benefits to their hosts such as some

Spiroplasma spp. that infect insects and confer protection against nematodes, parasitoid

wasps, and fungi through the production of unique toxins (Ballinger and Perlman 2019).

Recently, many novel and divergent members of this class have been discovered in

marine invertebrates from a variety of environments including a holothurian from a deep-

sea trench (He et al. 2018), a pelagic photosynthetic jellyfish (Viver et al. 2017; Cortés-

Lara et al. 2015), wood-boring deep-sea chitons (Duperron et al. 2013), ascidians from a

coastal lagoon (Tait et al. 2007), and crown-of-thorns sea stars from the Great Barrier

Reef (Høj et al. 2018). The genome of the holothurian associate has been sequenced and

it encodes an extensive CRISPR-Cas9 array leading to speculation that this associate

provides viral defense for its host (He et al. 2018). Similar insights about the interactions

between corals and their associated mollicutes may be gleaned by obtaining genomes of

their associates.

Here we discovered two novel bacteria of the class Mollicutes which are abundant

in the deep-sea octocoral *Callogorgia delta. C. delta* is a common coral along the

continental slope in the Gulf of Mexico between 400 m and 900 m depth, where it is

often the dominant habitat-forming coral (Quattrini et al. 2015; Quattrini et al. 2013). *C.*

delta colonies create habitat for numerous animal species including *Asteroschema*

ophiuroids and the chain catshark, *Scyliorhinus retifer,* which lay their eggs directly on

colonies (Etnoyer and Warrenchuk 2007). The closely related coral, *C. americana,* is also

a dominant habitat forming species. *C. delta* was originally classified as a subspecies of

C. americana until it was raised to the level of species in 2015 (Bayer et al. 2015). Both

species occur in the Gulf of Mexico but exhibit different ecological niches wherein *C.*

americana is found at shallower depths (300 - 400m) and *C. delta* is often found near

hydrocarbon seeps (Quattrini et al. 2013).

In order to characterize the association between *Callogorgia delta* and these

newly discovered mollicutes, we screened colonies from six sites in the Gulf of Mexico

across three years of sampling using 16S metabarcoding to determine the prevalence of

these novel mollicutes and quantify their relative abundances. We also screened the

closely related species *Callogorgia americana,* sediment, and water from the surrounding

environment for comparison. Further, we sequenced metagenomes and

metatranscriptomes of *C. delta* to assemble the genomes of these novel mollicutes, assess

their metabolic capabilities, and to determine their phylogenetic positions. Finally, we

used catalyzed reporter deposition fluorescence in situ hybridization (CARD-FISH)

microscopy to locate bacteria within coral tissue.

Methods:

Collections

Collections are a subset of those described in Vohsen et al. 2020. One hundred

and eight *Callogorgia delta* colonies were collected from six sites spanning over 350 km

in the northern Gulf of Mexico in 2015, 2016, and 2017 (Fig. 1). The sites are named

after the Bureau of Ocean Energy Management's designations for lease blocks in which

corals were sampled and include Mississippi Canyon (MC) 751 (n=19 colonies, 435-443

m, 28.193 N -89.800 W), Viosca Knoll (VK) 826 (n=3 colonies, 547-559 m, 29.159 N -

88.010 W), Green Canyon (GC) 234 (n=37 colonies, 505-536 m, 27.746 N -91.122 W),

MC885 (n=25 colonies, 624-642 m, 28.064 N -89.718 W), GC249 (n=14 colonies, 789-

795 m, 27.724 N -90.514 W), and GC290 (n=10 colonies, 852 m, 27.689 N -90.646 W).

To compare *Callogorgia delta* to a closely related coral species, we additionally

processed five *Callogorgia americana* colonies that were collected in 2010 from Garden

Banks (GB) 299 (n=3 colonies, 358-359m, 27.689 N -92.218 W) and VK862 (n=2

colonies, 352-357m, 29.109 N -88.387 W).

Colonies were sampled using specially designed coral cutters mounted on the

manipulator arm of remotely operated vehicles (ROVs). Coral fragments were removed,

placed in separate temperature insulated containers until recovery of the ROV, and were

maintained at 4 °C for up to 4 hours until preservation in ethanol or frozen in liquid

nitrogen.

Sediment samples were taken in close proximity to many of the *Callogorgia delta*

collections using push cores with a diameter of 6.3 cm. Upon recovery of the ROV, 1 mL

of sediment from the top 1 cm of each sediment core was frozen in liquid nitrogen for

later DNA extraction for microbiome analysis. Water was sampled in 2015 using a Large

Volume Water Transfer System (McLane Laboratories Inc., Falmouth, MA) which

filtered approximately 400 L of water through a 0.22 micron porosity filter with a

diameter of 142 mm. One quarter of each filter was preserved in ethanol and another

quarter was frozen in liquid nitrogen for these analyses. In 2016 and 2017, 2.5 L Niskin

bottles mounted on the ROV were fired above corals. Upon recovery of the ROV, this

water was filtered through 0.22 micron filters which were frozen in liquid nitrogen.

16S metabarcoding

140   The microbiomes of *Callogorgia delta, Callogorgia americana*, water, and

sediment were analyzed using 16S metabarcoding. DNA was extracted from coral tissue

and sediment samples using DNeasy PowerSoil kits (Qiagen, Hilden, Germany)

following manufacturer protocols using approximately 1 cm of coral branches and about

0.25 g of sediment. DNeasy PowerSoil kits were also used for water samples collected in

2015 using 1 cm^2 of filter. For all water samples from 2015, replicate extractions were

performed on quarters of the filter preserved in both ethanol and frozen in liquid nitrogen.

DNA was extracted from all other water samples using Qiagen DNeasy PowerWater kits.

148   The V1 and V2 regions of the 16S rRNA gene were amplified using universal

bacterial primers 27F and 355R (Rodriguez-Lanetty et al. 2013) with CS1 and CS2

adapters (Illumina, San Diego, CA). PCR was conducted with the following reaction

composition: 0.1 U/ µL Gotaq (Promega, Madison, WI), 1X Gotaq buffer, 0.25mM of

each dNTP, 2.5 mM MgCl$_2$, and 0.25 µM of each primer, and the following

thermocycler conditions: 95 °C for 5 min; 30 cycles of 95 °C for 30 sec, 51 °C for 1 min,

and 72 °C for 1 min; and finally 72 °C for 7 min. Libraries were prepared by the

University of Illinois Chicago DNA services facility and sequenced on two separate runs

on an Illumina MISEQ platform (Naqib et al. 2018). Raw sequence data are available on

the NCBI database under BioProject PRJNA565265 and BioSample IDs

SAMN12824733 – 4765, SAMN12824767 – 4875.

Amplicon sequence data were analyzed as described in (Vohsen et al. 2020) using

QIIME 2 (ver2017.11) (Bolyen et al. 2018). Reads were joined using vsearch and quality

filtered using q-score-joined. Deblur denoise was used to detect chimeras, construct sub-

operational taxonomic units (later referred to as ASVs), and trim assembled reads to

300bp. An ASV table was exported into R to create bar plots.

Metagenomes and Metatranscriptomes

Metagenomes and metatranscriptomes of *C. delta* were sequenced to assemble

genomes of novel mollicutes which were used to assess their phylogenetic positions and

metabolic potential. DNA was extracted from eight *Callogorgia delta* colonies as

described in Vohsen et al. 2020. In brief, DNA was extracted using DNA/RNA allprep

kits (Qiagen, Hilden, Germany) with β-mercaptoethanol added to the RLT+ lysis buffer

following the manufacturer's instructions. In addition to these samples, DNA was

extracted from an additional 18 samples corresponding to 16 additional colonies.

RNA was extracted from two of those *Callogorgia delta* colonies using an

RNeasy extraction kit (Qiagen, Hilden, Germany). The RNA extracts were enriched for

bacteria by depleting coral host rRNA using a Ribo-Zero Gold Yeast kit (Illumina, San

Diego, CA, USA) following the manufacturer's protocols.

The original eight DNA extracts were sent to the Max Planck Genome Center in

Cologne, Germany for library preparation and were sequenced on an Illumina

HISEQ2500 platform using 150 bp paired-end reads. The two *Callogorgia delta* libraries

with the highest coverage of mollicutes were sequenced a second time. Raw sequence

data are available on the NCBI database under BioProject ID PRJNA574146 and

BioSample IDs SAMN12856800 – SAMN12856807. Both the additional 18 DNA

extracts and RNA extracts were submitted to the Pennsylvania State University

Genomics Core facility for library preparation and were sequenced on a HISEQ with

100bp paired-end reads and a MISEQ platform with 100 bp single-end reads,

respectively. Raw sequence data are available on the NCBI database under BioProject ID

PRJNA565265 and SRA accession numbers SRR10174410 and SRR10174411.

All eight initial DNA sequence libraries were screened for bacterial SSU rRNA

using phyloFlash ver3.3 (Gruber-Vodicka et al. 2019). The two libraries with the highest

coverage of mollicute 16S rRNA were sequenced a second time. Of these, the library

with higher coverage was chosen for further processing. Reads were trimmed to a quality

score of 2 from both ends using BBduk and a kmer frequency analysis was performed

using BBnorm. The library was kfiltered to a depth of 1,000 or greater using BBnorm and

downsized to 10% using reformat with a single pass. The library was then assembled

using SPAdes (Bankevich et al. 2012) with a kmer progression of 21,33,55,77,99,127.

Bandage ver0.8.1 (Wick et al. 2015) was used to identify a single circular scaffold of

contigs containing the dominant mollicute 16S rRNA gene. The overlaps on the end were

manually trimmed off and the scaffold was recentered to the origin of replication

predicted by Ori-Finder (Gao and Zhang 2008). This constituted the draft genome for the

dominant mollicute and was annotated using classic RAST ver2.0 (Aziz et al. 2008;

Overbeek et al. 2005).

In order to generate a genome for the less abundant mollicute, the two libraries

with the highest coverage of its 16S rRNA gene were coassembled using megahit (D. H.

Li et al. 2015). All libraries were mapped to this coassembly using BBmap with a kfilter

of 22, subfilter of 15, and a maxindel of 80. Alignments were converted from SAM to

BAM formats and sorted using samtools (H. Li et al. 2009) and used for binning using

metabat (Kang et al. 2015) with default parameters. All scaffolds with contigs belonging

to bin 1 constituted a draft genome that was annotated using RAST and used for

phylogenomic analysis.

The presence of CRISPRs and associated genes and the number of spacers was

determined using RAST annotations and the tool CRISPRfinder (Symeonidi et al. 2021;

Grissa et al. 2007). OrthoANI (Lee et al. 2016) was used to estimate average nucleotide

identity between genomes. To determine which genes were expressed and to quantify

their expression, the two RNA sequence libraries were pseudoaligned to the gene

annotations of both genomes using kallisto ver0.44.0 (Bray et al. 2016) using 100

bootstrap replicates, an estimated fragment length of 180 with a standard deviation of 20,

and an index with a k-mer size of 31.

Phylogenetic Analyses

Phylogenetic trees were constructed to infer the phylogenetic positions of the

mollicutes that were detected in *Callogorgia.* First a tree of 16S sequences was generated

to place these novel mollicutes within the wide diversity of mollicutes with available 16S

sequences. Full length 16S rRNA genes were assembled from the metagenomes using

phyloFlash. These full length sequences were aligned with publicly available Mollicutes

sequences that were over 1,000 bp using MUSCLE ver3.8.425 (Edgar 2004). A

maximum likelihood tree was constructed using the IQ-TREE web server ver1.6.10

(Nguyen et al. 2014; Trifinopoulos et al. 2016). Within IQ-TREE, ModelFinder

(Kalyaanamoorthy et al. 2017) was used to choose a general time reversible model with

unequal rates and empirical base frequencies and a FreeRate model (Soubrier et al. 2012;

Z. Yang 1995) for rate heterogeneity across sites with 3 categories (GTR+F+R3) based

on the highest BIC score. UFBoot2 (Hoang et al. 2017) was used to obtain ultrafast

bootstrap support values (UFBoot) using 1000 replicates.

In order to get a more confident phylogenetic placement of these novel

mollicutes, a phylogenomic tree was constructed including genomes from both mollicutes

in *C. delta* plus an additional 66 publicly available genomes. One more genome was

assembled from a publicly available dataset because an assembled genome was not

available. This genome corresponded to the novel bacterium that associates with the

jellyfish, *Cotylorhiza tuberculata*, which those authors classified as *Mycoplasma.*

PhyloFlash (Gruber-Vodicka et al. 2019) was performed on all 4 available libraries to

assess bacterial compositions and the library with the highest coverage of *Mycoplasma*

(Library M3) was assembled using megahit (D. H. Li et al. 2015) with a kmax of 241. All

4 libraries were mapped to this assembly using bbmap with a kfilter of 22, subfilter of 15,

and maxindel of 80. The alignments were subsequently used to bin the assembly using

metabat (Kang et al. 2015) using default parameters. Bin 1 corresponded to the purported

Mycoplasma genome. This bin and all associated scaffolds were used to represent this

genome in phylogenomic analyses. A concatenated amino acid sequence alignment of 43

genes from all 69 genomes was generated with CheckM (ver1.1.3) (Parks et al. 2015).

This was used to generate a maximum likelihood phylogenetic tree using IQ-TREE

(ver1.6.12) (Hoang et al. 2017; Nguyen et al. 2014).

Both phylogenetic trees were annotated using the revised phylogeny of the class

Mollicutes established in (Radhey S. Gupta et al. 2018; Radhey S Gupta and Oren 2020;

Radhey S Gupta et al. 2019).

Microscopy

Catalyzed reporter deposition fluorescence in situ hybridization (CARD-FISH)

microscopy was employed to localize bacteria within *Callogorgia delta.* Colonies

collected in 2016 were preserved for microscopy. During subsampling, 0.5 cm long

branches were fixed in 2 mL of 4% formaldehyde with 1X phosphate-buffered saline

(PBS) at 4 °C overnight (4-8 hours). Samples were washed 3 times and then stored in

equal parts ethanol and 1X PBS for up to 21 months. Samples were decalcified in 0.45M

EDTA in 1X PBS for 7 days, refreshing the buffer after 4 days, then washed in 1X PBS.

Samples were post-fixed in 1X PBS with 1% paraformaldehyde for 1 hour then washed 3

times in 1X PBS and stored in 50:50 1X PBS:ethanol overnight. The next day, the

samples were embedded in TissueTek (Sakura Finetek, Maumee, OH) and left at 4 °C

overnight. The next day, the TissueTek was replaced and the samples were frozen.

Samples were sectioned to a thickness of 4 µm using a Leica (Wetzlar, Germany)

cryostat CM3050 S. Adjacent sections were placed on polysine slides and stored at 4 °C

before hybridization.

Hybridization began by removing TissueTek from tissue sections via soaking

slides in milliQ water for 3 minutes. Autofluorescence was then diminished by soaking

slides in ethanol with 0.1% Sudan black B for 10 minutes. Sections were washed three

times in 1X PBS and left in 1X PBS for a final 24 minutes. Endogenous peroxidases were

inactivated by soaking sections in 0.2M HCl for 12 minutes, then 20 mM Tris-HCl for 20

minutes, lysozyme solution (0.01 g/mL) for 30 minutes at 37 °C, 20 mM Tris-HCl again

for 20 minutes, and finally 3 minutes in milliQ water. Slides were then airdried and

sections were circled with a PAP pen. Two probes were used: EUB I-III which targets

eubacteria and a control probe (non-EUB) consisting of the reverse complement.

Hybridization mixes were created by combining 1 µL of probe (50 ng/µL) with 299 µL

hybridization buffer (30% formamide with 0.102 M NaCl, 0.02 M Tris HCl, 10%

Blocking Reagent, 0.01% sodium dodecyl sulfate, and 0.1 g/mL dextran sulfate). Each

section was covered with approximately 10-20 µL of one of the hybridization mixes.

Two adjacent sections were hybridized with the EUB I-III and non-EUB probes

separately to produce a negative control of the same tissue region. Each slide was then

placed in a humidity chamber consisting of a closed 50 mL tube with a kimwipe soaked

in 2 mL of 30% formamide in milliQ water. Hybridization was accomplished by

incubating slides at 46 °C for 2 hours within their humidity chambers.

After hybridization, slides were washed for 15 minutes at 48 °C in washing buffer

(0.102M NaCl, 0.02M Tris HCl, 0.005M EDTA, 0.01% SDS) followed by 15 minutes in

1X PBS at room temperature. A fresh amplification mix was created by first combining

299 µL 1X PBS and 1 µL H_2O_2, then adding 10 µL of that solution to 1 mL of

amplification buffer (1X PBS, 0.01% Blocking Reagent, 2M NaCl, 0.1 g/mL dextran

sulfate) and 1 µL of labelled tyramide (1mg/mL ALEXA 488-labelled tyramide in

dimethylformamide with 20 mg/mL p-iodophenylboronic acid). Slides were again placed

in 50 mL humidity chambers with a kimwipe soaked in 2 mL of milliQ water.

Amplification was accomplished by incubating slides at 37 °C for 30 minutes in these

humidity chambers.

After amplification, slides were washed by dipping in 1X PBS followed by 10

minutes in 1X PBS, and a final dip in milliQ water. Slides were air dried and sections

were stained with DAPI (1 µg/mL) for 10 minutes in the dark. Slides were washed a final

time in milliQ water for 3 minutes. Once airdried, slides were mounted using Vectashield

(Vector Laboratories, Burlingame, CA) and viewed on an Axioscope epifluorescence

microscope (Carl Zeiss AG, Oberkochen, Germany) with filters restricted to the emission

spectra of ALEXA 488 and DAPI. Images were taken with a Zeiss AxioCam camera and

visualized using AxioVision software. CARD-FISH signals were considered positive if

they appeared consistently within a structure in the coral, were absent in the same

structures in negative control sections, and overlapped DAPI signals.

Results:

Occurrence of novel Mollicutes in and around Callogorgia

An ASV (1) belonging to class Mollicutes was dominant in *Callogorgia delta*

colonies (Fig. 2). It was detected in 99 out of 108 colonies and was present in samples

from all three sampling years (2015-2017) and all six sampling locations (Fig. 2). It

averaged 78% of the microbial community among corals which harbored it and reached a

maximum of 99% of the community in some samples. It was also present in all five

Callogorgia americana colonies where it averaged 91% of the microbial community. An

additional ASV (2) was identified that differed from ASV 1 by a single base pair among

300. This closely related ASV was detected in 4 of 10 colonies from the deepest site,

GC290, and averaged 90% of the microbial community in those samples while ASV 1

was undetected (Fig. 2).

Another ASV (3) was identified that belonged to class Mollicutes. While it was

prevalent and abundant in some colonies, it was less so compared to ASV 1 (Fig. 2). This

ASV was present in 69 out of 108 *C. delta* colonies (up to 82% of the coral-associated

microbes) and three out of five *C. americana* colonies (up to 15% of the coral-associated

microbes). However, this ASV was absent in all *C. delta* colonies from sites GC249

(n=14) and GC290 (n=10).

Neither of these mollicute ASVs were detected in any water sample (n=30).

However, ASVs 1 and 3 were detected in 35 and 11 out of 47 total sediment samples with

maximum relative abundances of 1.1 and 0.16% of the sediment microbial community

respectively.

Phylogenetic positions of the mollicute in Callogorgia delta

Twenty-two full length 16S rRNA gene sequences that correspond to ASV 1 were

assembled from separate metagenomes and an additional three were assembled that

correspond to ASV 3. Phylogenetic analysis using these sequences revealed that they

were distinct from other known mollicutes and clustered with a diverse array of

mollicutes recently discovered in association with marine invertebrates (Fig. 3a,b).

Interestingly, the two novel mollicutes in *Callogorgia delta* are not each other's closest

relative and shared only 84% sequence identity which is below the level which has been

suggested to differentiate bacterial families (Yarza et al. 2014). Phylogenomic analysis of

shared amino acid sequences among mollicute genomes reiterated this pattern (Fig. 3b).

A clade of novel Mollicutes that associates with marine invertebrates was well-supported

in both 16S and phylogenomic trees (UFBoot value of 100%). Branch lengths between

novel marine mollicutes were long and the average nucleotide identity (ANI) of these

novel mollicutes with their closest known relatives were <67% (OrthoANI) suggesting

that the mollicutes described here are novel genera and that there is great diversity within

this group. We propose the names *Candidatus Oceanoplasma callogorgiae* gen. nov. sp.

nov. and *Ca. Thalassoplasma callogorgiae* gen. nov. sp. nov. for the dominant and less

abundant novel mollicutes respectively.

Metabolic capabilities of the novel mollicutes in Callogorgia delta

The genome assembled for the dominant mollicute, *Ca. Oceanoplasma*

callogorgiae, was very reduced. It was small (446,099 bp) and encoded only 405

predicted genes. Of these, 265 had functional annotations (65%) and 140 were classified

as only hypothetical genes. A single gene was identified that may be involved in

carbohydrate metabolism, glyceraldehyde-3-phosphate dehydrogenase. Instead, the

genome encoded the three genes that conduct the arginine dihydrolase pathway (*arcA* =

arginine deiminase, *arcB* = ornithine carbamoyltransferase, and *arcC* = carbamate

kinase). These genes along with *arcD* (arginine/ornithine antiporter) generate ATP

through the catabolism of arginine and were among the most highly expressed genes in

both RNA libraries (Fig. 4). Like other mollicutes, this genome encoded clustered

regularly interspaced short palindromic repeats (CRISPRs) and cas genes (*cas1*, *cas2,*

and *cas9*) comprising a Class 2 Type II-C system. It contained more spacer sequences, at

107, than most mollicutes with available genomes (Ipoutcha et al. 2019). It also contained

type I, II, and III restriction modification systems.

The genome of *Ca. Thalassoplasma callogorgiae* was very similar to that of *Ca.*

O. callogorgiae. It was reduced in size (445,456 bp) and metabolic capabilities with only

416 gene annotations (163 hypothetical). It did not encode pathways for carbohydrate

metabolism but did encode glyceraldehyde-3-phosphate dehydrogenase, ribose-5-

phosphate isomerase, ribose-phosphate pyrophosphokinase, and a ribose importer. Like

Ca. O. callogorgiae, it contained the genes involved in the arginine deiminase pathway, a

type I restriction modification system, and a Class 2 Type II-C CRISPR-Cas system.

Only 14 spacer sequences were identified however this spacer array spanned a gap

between scaffolds and is likely an underestimate.

Localization of bacteria in Callogorgia delta

372   Fluorescent signals were observed in the mesoglea of two separate *Callogorgia*

delta colonies and interpreted as abundant bacteria (Fig. 5 and 6)*.* These signals were

present in tissue sections hybridized with the EUB I-III probe but absent in adjacent

sections hybridized with the control probe (non-EUB). These signals were <1 µm in

diameter and overlapped DAPI fluorescence. These signals were observed in both the

mesoglea within polyps (Fig. 5c,e,g) and in mesoglea within the coenosarc surrounding

the proteinaceous central axis (Fig. 5a). They formed large aggregates potentially

comprised of hundreds or thousands of cells (Fig. 6). Another fluorescence signal was

observed in the base of the tentacles in one colony but was far less aggregated (Fig. 6c,d).

Discussion:

Prevalence of novel Mollicutes in Callogorgia delta

383   The association between *Ca. Oceanoplasma callogorgiae* and *Callogorgia delta*

was stable across geography and time. Including both ASV 1 and 2, over 95% of *C. delta*

colonies hosted *Ca. O. callogorgiae* including samples from every sampling year and

every site. *Ca. T. callogorgia* was less prevalent. While it was detected in samples from

every year, it was detected in fewer samples, those samples hosted a smaller relative

abundance, and they were completely absent from the deepest two sites. These microbes

may associate with corals in the genus *Callogorgia* more broadly since both *Ca. O.*

callogorgiae and *Ca. T. callogorgiae* were detected in the closely related species *C.*

americana but were absent or rare in the water and sediment from the same sites.

These results suggest that *Callogorgia* is the primary habitat for these mollicutes.

This is corroborated by our observation of abundant bacteria in the mesoglea of *C. delta.*

The bacteria living in the mesoglea are likely to be *Ca. Oceanoplasma callogorgiae*

given its abundance in the amplicon dataset and in the metagenomes where its genomic

coverage is as high as 100 times that of *C. delta*.

Metabolic capabilities and interaction with the coral host

The genomes of both *Ca. Oceanoplasma callogorgiae* and *Ca. Thalassoplasma*

callogorgia were reduced like many symbiotic bacteria and possessed almost identical

metabolic capabilities. Both genomes revealed a dependence on arginine to generate ATP

which they likely receive from their coral hosts. However, it has been suggested that the

arginine dihydrolase pathway alone is not efficient enough to support the ATP demands

of mollicutes (Romano et al. 1980). The energy demands of some non-fermentative

mollicutes are additionally supported by the generation of ammonium which generates a

proton gradient that in turn drives ATP-synthases (Romano et al. 1980; Smith et al.

1993). This likely occurs in these mollicutes since the arginine dihydrolase pathway

produces ammonium and ATP-synthases were among their most highly expressed genes.

The catabolism of arginine may be a key pathway at the center of the interaction

between *Callogorgia delta* and its dominant mollicute. Arginine is an important regulator

in the tripartite symbiosis between plants, their mycorrhizal associates, and the

endosymbiotic mollicutes which are found within the mycorrhizae. Arginine constitutes

over 90% of the amino acids in arbuscular mycorrhizal fungi and serves as the primary

intermediate in nitrogen exchange between fungal and plant partners (Govindarajulu et al.

2005; Tian et al. 2010). Abundant arginine may similarly be provided by *Callogorgia*

delta, however it is unclear how either coral or mollicute may process the resulting

ammonia and ornithine.

Another commonality between these novel mollicutes was the presence of

CRISPR-Cas systems in both of their genomes. *Ca. Oceanoplasma callogorgiae*

possessed substantially more spacer sequences than most other mollicutes (Ipoutcha et al.

2019). This suggests that they possess extensive viral defenses and/or often encounter

foreign DNA. Other metabolic processes may also underly the basic functioning of these

novel mollicutes since many genes were unannotated. In fact, some of the most highly

expressed genes had unknown functions. Further study is required to describe the

complete metabolic potential of these mollicutes.

While there is strong evidence that the dominant mollicute is a symbiont, it is still

unclear if the mollicute provides any benefit to its coral host or if it is commensal or a

parasite like many other members of the Mollicutes. The metabolism of arginine by the

mollicute may provide the coral with an alternate pathway to process nitrogenous waste

or may even function to recycle nitrogen which may be useful for corals which rely on a

nitrogen-poor diet of marine snow (Kriest 2002). However, these possibilities are

speculative without a complete understanding of the capabilities of both the mollicute and

the coral host. Another possible benefit that the mollicute may provide is protection from

viruses using their CRISPR-Cas systems as suggested for the symbionts of hadal sea

cucumbers (He et al. 2018) which are the closest relatives of *Ca. O. callogorgiae*.

additional experiments are required to fully understand the metabolic interactions

between these mollicutes and their coral host and any potential benefit this may provide

for the coral.

438
Phylogenetic position of the novel mollicutes in C. delta

Ca. Oceanoplasma callogorgiae and *Ca. Thalassoplamsa callogorgiae* were both

novel and highly divergent members of the mollicutes. Phylogenetic analysis grouped

them with other novel mollicutes that were isolated from other marine invertebrates with

high support. These other invertebrates were phylogenetically diverse, exhibited diverse

lifestyles, and originated from wide ranging habitats including holothurians from hadal

trenches, sea stars from the Great Barrier Reef, wood-boring chitons from the deep sea,

ascidians from a coastal lagoon, and a pelagic jellyfish with algal symbionts (Cortés-Lara

et al. 2015; Duperron et al. 2013; Tait et al. 2007; Viver et al. 2017; He et al. 2018). The

diversity of these related associates suggests that this novel clade associates with marine

invertebrates broadly and is not restricted to hosts with specific lifestyles or from specific

habitats. Further, the large phylogenetic distances between mollicutes in this clade

suggests that a substantial amount of diversity remains undiscovered. Interestingly, this

novel clade of marine mollicutes clusters alongside the endocellular symbionts of

mycorrhizal fungi so these two clades may be more closely related to each other than to

other mollicutes.

In addition to the novel mollicutes in *C. delta,* the metabolic capabilities of two

other members of this clade were able to be characterized: the associate of hadal sea

cucumbers named *Spiroplasma holothuricola* and the associate of a pelagic jellyfish

classified as *Mycoplasma*. These two genomes showed many similarities with the

genomes of the novel mollicutes in *C. delta.* All four genomes were very small ranging

between 380 kbp and 450 kbp placing them among the smallest genomes of any bacteria

and smaller than other members of the class Mollicutes (Faburay and McVey 2022). All

four genomes were very restricted in metabolic capabilities. All encoded the arginine

dihydrolase pathway and no more than four genes involved in carbohydrate metabolism.

The mollicute from the jellyfish encoded lactate dehydrogenase, ribulose-phosphate 3-

epimerase, and two genes in common with *Ca. Thalassoplasma callogorgiae*: ribose-5-

phosphate isomerase and ribose-phosphate pyrophosphokinase. *Spiroplasma*

holothuricola encoded only phosphate acetyltransferase and acetate kinase. All four

genomes encoded Class 2 Type II-C CRISPR-Cas systems. The two genomes with

scaffolds that spanned the entire CRISPR array contained 107 and 76 spacers. This is

more than any other member of the Mollicutes except for *Mycoplasma liparidae,* a gut

associate of a hadal snailfish, which encoded 118 (Lian et al. 2020). The extent of

CRISPR arrays in these deep sea mollicutes may reflect a common adaptation to deep-sea

environments.

Conclusion

Here we discovered two novel members of the Mollicutes that associate with the

coral *Callogorgia delta* and propose the names *Ca. Oceanoplasma callogorgiae* and *Ca.*

Thalassoplasma callogorgiae. We demonstrated their prevalence across sites, years, and

in a congener, generated genomes, characterized their metabolic potential, located

abundant bacteria within the mesoglea of the coral host, and uncovered their phylogenetic

position among a novel clade of marine mollicutes that has not yet been recognized. This

work advances our understanding of the widespread coral associates of the class

Mollicutes by providing genomic resources and metabolic inferences but also reveals that

a large diversity of these novel marine mollicutes remain undiscovered.

484
485

References:

Atkinson, T. P. (2018). Mollicutes: Mycoplasma pneumoniae. In G. Ragab, T. P.
Atkinson, & M. L. Stoll (Eds.), *The Microbiome in Rheumatic Diseases and*
Infection (pp. 103-111). Cham: Springer International Publishing.
Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., et al. (2008).
The RAST Server: rapid annotations using subsystems technology. *BMC*
Genomics, 9, 75, doi:10.1186/1471-2164-9-75.
Baker, L. J., Reich, H. G., Kitchen, S. A., Grace Klinges, J., Koch, H. R., Baums, I. B., et al.
(2022). The coral symbiont Candidatus Aquarickettsia is variably abundant
in threatened Caribbean acroporids and transmitted horizontally. *The ISME*
Journal, 16(2), 400-411, doi:10.1038/s41396-021-01077-8.
Ballinger, M. J., & Perlman, S. J. (2019). The defensive spiroplasma. *Current opinion in*
insect science, 32, 36-41.
Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., et al.
(2012). SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *Journal of computational biology, 19*(5), 455-477.
Bayer, F. M., Cairns, S. D., Cordeiro, R. T., & Pérez, C. D. (2015). New records of the
genus Callogorgia (Anthozoa: Octocorallia) in the western Atlantic, including
the description of a new species. *Journal of the Marine Biological Association*
of the United Kingdom, 95(5), 905-911.
Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A., et al.
(2018). QIIME 2: Reproducible, interactive, scalable, and extensible
microbiome data science. *PeerJ Preprints, 6*, e27295v27292,
doi:10.7287/peerj.preprints.27295v2.
Bonfante, P., & Desirò, A. (2017). Who lives in a fungus? The diversity, origins and
functions of fungal endobacteria living in Mucoromycota. [Mini Review]. *The*
ISME Journal, 11, 1727, doi:10.1038/ismej.2017.21.
Bourne, D. G., Morrow, K. M., & Webster, N. S. (2016). Insights into the Coral
Microbiome: Underpinning the Health and Resilience of Reef Ecosystems.
Annual Review of Microbiology, 70(1), 317-340, doi:10.1146/annurev-micro-
102215-095440.
Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. *Nature Biotechnology, 34*, 525, doi:10.1038/nbt.3519
https://www.nature.com/articles/nbt.3519#supplementary-information.
Cortés-Lara, S., Urdiain, M., Mora-Ruiz, M., Prieto, L., & Rosselló-Móra, R. (2015).
Prokaryotic microbiota in the digestive cavity of the jellyfish Cotylorhiza
tuberculata. *Systematic and Applied Microbiology, 38*(7), 494-500,
doi:https://doi.org/10.1016/j.syapm.2015.07.001.
Desiro, A., Faccio, A., Kaech, A., Bidartondo, M. I., & Bonfante, P. (2015). *Endogone*,
one of the oldest plant-associated fungi, host unique Mollicutes-related
endobacteria. *New Phytol, 205*(4), 1464-1472, doi:10.1111/nph.13136.
Duperron, S., Pottier, M.-A., Léger, N., Gaudron, S. M., Puillandre, N., Le Prieur, S., et
al. (2013). A tale of two chitons: is habitat specialisation linked to distinct
associated bacterial communities? *FEMS Microbiology Ecology, 83*(3), 552-
567, doi:10.1111/1574-6941.12014.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and
high throughput. *Nucleic Acids Res, 32*(5), 1792-1797,
doi:10.1093/nar/gkh340.
Etnoyer, P., & Warrenchuk, J. (2007). A catshark nursery in a deep gorgonian field in
the Mississippi Canyon, Gulf of Mexico. *Bulletin of Marine Science, 81*, 553-
559.
Faburay, B., & McVey, D. S. (2022). Mollicutes. *Veterinary Microbiology*, 364-376.
Gao, F., & Zhang, C.-T. (2008). Ori-Finder: a web-based system for finding oriC s in
unannotated bacterial genomes. *BMC Bioinformatics, 9*(1), 1-6.
Govindarajulu, M., Pfeffer, P. E., Jin, H., Abubaker, J., Douds, D. D., Allen, J. W., et al.
(2005). Nitrogen transfer in the arbuscular mycorrhizal symbiosis. *Nature,*
435(7043), 819-823, doi:10.1038/nature03610.
Gray, M. A., Stone, R. P., McLaughlin, M. R., & Kellogg, C. A. (2011). Microbial
consortia of gorgonian corals from the Aleutian islands. *FEMS Microbiology*
Ecology, 76(1), 109-120, doi:10.1111/j.1574-6941.2010.01033.x.
Grissa, I., Vergnaud, G., & Pourcel, C. (2007). CRISPRFinder: a web tool to identify
clustered regularly interspaced short palindromic repeats. *Nucleic Acids*
Research, 35(suppl_2), W52-W57.
Gruber-Vodicka, H. R., Seah, B. K. B., & Pruesse, E. (2019). phyloFlash – Rapid SSU
rRNA profiling and targeted assembly from metagenomes. *bioRxiv*, 521922,
doi:10.1101/521922.
Gupta, R. S., & Oren, A. (2020). Necessity and rationale for the proposed name
changes in the classification of Mollicutes species. Reply to:'Recommended
rejection of the names Malacoplasma gen. nov., Mesomycoplasma gen. nov.,
Metamycoplasma gen. nov., Metamycoplasmataceae fam. nov.,
Mycoplasmoidaceae fam. nov., Mycoplasmoidales ord. nov., Mycoplasmoides
gen. nov., Mycoplasmopsis gen. nov.[Gupta, Sawnani, Adeolu, Alnajar and
Oren 2018] and all proposed species comb. nov. placed therein', by M. Balish
et al.(Int J Syst Evol Microbiol, 2019; 69: 3650–3653). *International Journal of*
Systematic and Evolutionary Microbiology, 70(2), 1431-1438.
Gupta, R. S., Sawnani, S., Adeolu, M., Alnajar, S., & Oren, A. (2018). Phylogenetic
framework for the phylum Tenericutes based on genome sequence data:
proposal for the creation of a new order *Mycoplasmoidales* ord. nov.,
containing two new families *Mycoplasmoidaceae* fam. nov. and
Metamycoplasmataceae fam. nov. harbouring *Eperythrozoon*, *Ureaplasma* and
five novel genera. *Antonie van Leeuwenhoek, 111*(9), 1583-1630,
doi:10.1007/s10482-018-1047-3.
Gupta, R. S., Son, J., & Oren, A. (2019). A phylogenomic and molecular markers based
taxonomic framework for members of the order Entomoplasmatales:
proposal for an emended order Mycoplasmatales containing the family
Spiroplasmataceae and emended family Mycoplasmataceae comprised of six
genera. *Antonie van Leeuwenhoek, 112*(4), 561-588.
He, L.-S., Zhang, P.-W., Huang, J.-M., Zhu, F.-C., Danchin, A., & Wang, Y. (2018). The
Enigmatic Genome of an Obligate Ancient *Spiroplasma* Symbiont in a Hadal
Holothurian. *Applied and Environmental Microbiology, 84*(1), e01965-01917,
doi:10.1128/aem.01965-17.

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2017).
UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular*
Biology and Evolution, 35(2), 518-522, doi:10.1093/molbev/msx281.
Høj, L., Levy, N., Baillie, B. K., Clode, P. L., Strohmaier, R. C., Siboni, N., et al. (2018).
Crown-of-Thorns Sea Star, *Acanthaster* cf. *solaris*, Has Tissue-Characteristic
Microbiomes with Potential Roles in Health and Reproduction. *Applied and*
Environmental Microbiology, 84(13), e00181-00118,
doi:10.1128/aem.00181-18.
Holm, J. B., & Heidelberg, K. B. (2016). Microbiomes of Muricea californica and M.
fruticosa: Comparative Analyses of Two Co-occurring Eastern Pacific
Octocorals. [Original Research]. *Frontiers in Microbiology, 7*(917),
doi:10.3389/fmicb.2016.00917.
Ing-Ming Lee, Robert E. Davis, & Gundersen-Rindal, D. E. (2000). Phytoplasma:
Phytopathogenic Mollicutes. *Annual Review of Microbiology, 54*(1), 221-255,
doi:10.1146/annurev.micro.54.1.221.
Ipoutcha, T., Tsarmpopoulos, I., Talenton, V., Gaspin, C., Moisan, A., Walker, C. A., et
al. (2019). Multiple origins and specific evolution of CRISPR/Cas9 systems in
minimal bacteria (Mollicutes). *Frontiers in Microbiology, 10*, 2701.
Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., & Jermiin, L. S.
(2017). ModelFinder: fast model selection for accurate phylogenetic
estimates. *Nature Methods, 14*, 587, doi:10.1038/nmeth.4285
https://www.nature.com/articles/nmeth.4285#supplementary-information.
Kang, D. D., Froula, J., Egan, R., & Wang, Z. (2015). MetaBAT, an efficient tool for
accurately reconstructing single genomes from complex microbial
communities. *PeerJ, 3*, e1165, doi:10.7717/peerj.1165.
Kellogg, C. A., Lisle, J. T., & Galkiewicz, J. P. (2009). Culture-Independent
Characterization of Bacterial Communities Associated with the Cold-Water
Coral *Lophelia pertusa* in the Northeastern Gulf of Mexico. *Applied and*
Environmental Microbiology, 75(8), 2294-2303, doi:10.1128/aem.02357-08.
Kellogg, C. A., Ross, S. W., & Brooke, S. D. (2016). Bacterial community diversity of
the deep-sea octocoral *Paramuricea placomus. PeerJ, 4*, e2529,
doi:10.7717/peerj.2529.
Kirk, N. L., Ritson-Williams, R., Coffroth, M. A., Miller, M. W., Fogarty, N. D., & Santos,
S. R. (2013a). Tracking Transmission of Apicomplexan Symbionts in Diverse
Caribbean Corals. [Article]. *PLOS ONE, 8*(11), 10,
doi:10.1371/journal.pone.0080618.
Kirk, N. L., Thornhill, D. J., Kemp, D. W., Fitt, W. K., & Santos, S. R. (2013b). Ubiquitous
associations and a peak fall prevalence between apicomplexan symbionts
and reef corals in Florida and the Bahamas. [Article]. *Coral Reefs, 32*(3), 847-
858, doi:10.1007/s00338-013-1038-9.
Klinges, G., Maher, R. L., Vega Thurber, R. L., & Muller, E. M. (2020). Parasitic
'Candidatus Aquarickettsia rohweri' is a marker of disease susceptibility in
Acropora cervicornis but is lost during thermal stress.
[https://doi.org/10.1111/1462-2920.15245]. *Environmental Microbiology,*
22(12), 5341-5355, doi:https://doi.org/10.1111/1462-2920.15245.

Klinges, J. G., Patel, S. H., Duke, W. C., Muller, E. M., & Vega Thurber, R. L. (2022).
Phosphate enrichment induces increased dominance of the parasite
Aquarickettsia in the coral Acropora cervicornis. *FEMS Microbiology Ecology*,
fiac013, doi:10.1093/femsec/fiac013.
Klinges, J. G., Rosales, S. M., McMinds, R., Shaver, E. C., Shantz, A. A., Peters, E. C., et al.
(2019). Phylogenetic, genomic, and biogeographic characterization of a novel
and ubiquitous marine invertebrate-associated Rickettsiales parasite,
Candidatus Aquarickettsia rohweri, gen. nov., sp. nov. *The ISME Journal,*
13(12), 2938-2953, doi:10.1038/s41396-019-0482-0.
Kriest, I. (2002). Different parameterizations of marine snow in a 1D-model and
their influence on representation of marine snow, nitrogen budget and
sedimentation. *Deep Sea Research Part I: Oceanographic Research Papers,*
49(12), 2133-2162, doi:https://doi.org/10.1016/S0967-0637(02)00127-9.
Kuga, Y., Wu, T.-D., Sakamoto, N., Katsuyama, C., & Yurimoto, H. (2021). Allocation of
Carbon from an Arbuscular Mycorrhizal Fungus, Gigaspora margarita, to Its
Gram-Negative and Positive Endobacteria Revealed by High-Resolution
Secondary Ion Mass Spectrometry. *Microorganisms, 9*(12), 2597.
Kwong, W. K., del Campo, J., Mathur, V., Vermeij, M. J. A., & Keeling, P. J. (2019). A
widespread coral-infecting apicomplexan with chlorophyll biosynthesis
genes. [Article]. *Nature, 568*(7750), 103-107, doi:10.1038/s41586-019-
1072-z.
Lee, I., Kim, Y. O., Park, S.-C., & Chun, J. (2016). OrthoANI: an improved algorithm and
software for calculating average nucleotide identity. *International Journal of*
Systematic and Evolutionary Microbiology, 66(2), 1100-1103.
Lesser, M. P., Falcón, L. I., Rodrí-guez-Román, A., Enr-íquez, S., Hoegh-Guldberg, O., &
Iglesias-Prieto, R. (2007). Nitrogen fixation by symbiotic cyanobacteria
provides a source of nitrogen for the scleractinian coral Montastraea
cavernosa. *Marine Ecology Progress Series, 346*, 143-152.
Lesser, M. P., Mazel, C. H., Gorbunov, M. Y., & Falkowski, P. G. (2004). Discovery of
Symbiotic Nitrogen-Fixing Cyanobacteria in Corals. *Science, 305*(5686), 997-
1000, doi:10.1126/science.1099128.
Li, D. H., Liu, C. M., Luo, R. B., Sadakane, K., & Lam, T. W. (2015). MEGAHIT: an ultra-
fast single-node solution for large and complex metagenomics assembly via
succinct de Bruijn graph. [Article]. *Bioinformatics, 31*(10), 1674-1676,
doi:10.1093/bioinformatics/btv033.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The
sequence alignment/map format and SAMtools. *Bioinformatics, 25*(16), 2078-
2079.
Lian, C.-A., Yan, G.-Y., Huang, J.-M., Danchin, A., Wang, Y., & He, L.-S. (2020). Genomic
characterization of a novel gut symbiont from the hadal snailfish. *Frontiers in*
Microbiology, 10, 2978.
McCauley, M., Jackson, C. R., & Goulet, T. L. (2020). Microbiomes of Caribbean
octocorals vary over time but are resistant to environmental change.
Frontiers in Microbiology, 11, 1272.
Naqib, A., Poggi, S., Wang, W., Hyde, M., Kunstman, K., & Green, S. J. (2018). Making
and sequencing heavily multiplexed, high-throughput 16S ribosomal RNA

gene amplicon libraries using a flexible, two-stage PCR protocol. In *Gene*
expression analysis (pp. 149-169): Springer.
Neulinger, S. C., Gärtner, A., Järnegren, J., Ludvigsen, M., Lochte, K., & Dullo, W.-C.
(2009). Tissue-Associated "*Candidatus* Mycoplasma corallicola" and
Filamentous Bacteria on the Cold-Water Coral *Lophelia pertusa* (Scleractinia).
Applied and Environmental Microbiology, 75(5), 1437-1444,
doi:10.1128/aem.01781-08.
Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2014). IQ-TREE: A Fast
and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
Phylogenies. *Molecular Biology and Evolution, 32*(1), 268-274,
doi:10.1093/molbev/msu300.
Overbeek, R., Begley, T., Butler, R. M., Choudhuri, J. V., Chuang, H. Y., Cohoon, M., et al.
(2005). The subsystems approach to genome annotation and its use in the
project to annotate 1000 genomes. *Nucleic Acids Res, 33*(17), 5691-5702,
doi:10.1093/nar/gki866.
Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015).
CheckM: assessing the quality of microbial genomes recovered from isolates,
single cells, and metagenomes. *Genome research, 25*(7), 1043-1055.
Penn, K., Wu, D., Eisen, J. A., & Ward, N. (2006). Characterization of Bacterial
Communities Associated with Deep-Sea Corals on Gulf of Alaska Seamounts.
Applied and Environmental Microbiology, 72(2), 1680-1683,
doi:10.1128/aem.72.2.1680-1683.2006.
Pollock, F. J., McMinds, R., Smith, S., Bourne, D. G., Willis, B. L., Medina, M., et al.
(2018). Coral-associated bacteria demonstrate phylosymbiosis and
cophylogeny. *Nature Communications, 9*(1), 4921, doi:10.1038/s41467-018-
07275-x.
Porporato, E. M. D., Lo Giudice, A., Michaud, L., De Domenico, E., & Spanò, N. (2013).
Diversity and Antibacterial Activity of the Bacterial Communities Associated
with Two Mediterranean Sea Pens, *Pennatula phosphorea* and *Pteroeides*
spinosum (Anthozoa: Octocorallia). [journal article]. *Microbial Ecology, 66*(3),
701-714, doi:10.1007/s00248-013-0260-x.
Quattrini, A. M., Baums, I. B., Shank, T. M., Morrison, C. L., & Cordes, E. E. (2015).
Testing the depth-differentiation hypothesis in a deepwater octocoral.
Proceedings of the Royal Society B: Biological Sciences, 282(1807), 20150008,
doi:doi:10.1098/rspb.2015.0008.
Quattrini, A. M., Georgian, S. E., Byrnes, L., Stevens, A., Falco, R., & Cordes, E. E.
(2013). Niche divergence by deep-sea octocorals in the genus *Callogorgia*
across the continental slope of the Gulf of Mexico. *Molecular Ecology, 22*(15),
4123-4140, doi:10.1111/mec.12370.
Quintanilla, E., Rodrigues, C. F., Henriques, I., & Hilário, A. (2022). Microbial
Associations of Abyssal Gorgonians and Anemones (> 4,000 m Depth) at the
Clarion-Clipperton Fracture Zone. *Frontiers in Microbiology, 13*.
Rodriguez-Lanetty, M., Granados-Cifuentes, C., Barberan, A., Bellantuono, A. J., &
Bastidas, C. (2013). Ecological Inferences from a deep screening of the
Complex Bacterial Consortia associated with the coral, Porites astreoides.
Molecular Ecology, 22(16), 4349-4362, doi:10.1111/mec.12392.

Romano, N., Tolone, G., Ajello, F., & La Licata, R. (1980). Adenosine 5'-triphosphate
synthesis induced by urea hydrolysis in *Ureaplasma urealyticum*. *Journal of*
Bacteriology, 144(2), 830-832.
Shirur, K. P., Jackson, C. R., & Goulet, T. L. (2016). Lesion recovery and the bacterial
microbiome in two Caribbean gorgonian corals. [journal article]. *Marine*
Biology, 163(12), 238, doi:10.1007/s00227-016-3008-6.
Smith, D. G., Russell, W. C., Ingledew, W. J., & Thirkell, D. (1993). Hydrolysis of urea
by *Ureaplasma urealyticum* generates a transmembrane potential with
resultant ATP synthesis. *Journal of Bacteriology, 175*(11), 3253-3258,
doi:10.1128/jb.175.11.3253-3258.1993.
Soubrier, J., Steel, M., Lee, M. S. Y., Der Sarkissian, C., Guindon, S., Ho, S. Y. W., et al.
(2012). The Influence of Rate Heterogeneity among Sites on the Time
Dependence of Molecular Rates. *Molecular Biology and Evolution, 29*(11),
3345-3358, doi:10.1093/molbev/mss140.
Symeonidi, E., Regalado, J., Schwab, R., & Weigel, D. (2021). CRISPR-finder: A high
throughput and cost-effective method to identify successfully edited
Arabidopsis thaliana individuals. *Quantitative Plant Biology, 2*.
Tait, E., Carman, M., & Sievert, S. M. (2007). Phylogenetic diversity of bacteria
associated with ascidians in Eel Pond (Woods Hole, Massachusetts, USA).
Journal of Experimental Marine Biology and Ecology, 342(1), 138-146,
doi:https://doi.org/10.1016/j.jembe.2006.10.024.
Tian, C., Kasiborski, B., Koul, R., Lammers, P. J., Bücking, H., & Shachar-Hill, Y. (2010).
Regulation of the Nitrogen Transfer Pathway in the Arbuscular Mycorrhizal
Symbiosis: Gene Characterization and the Coordination of Expression with
Nitrogen Flux. *Plant Physiology, 153*(3), 1175-1187,
doi:10.1104/pp.110.156430.
Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., & Minh, B. Q. (2016). W-IQ-TREE: a
fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids*
Research, 44(W1), W232-W235.
Tully, J. G., Bové, J. M., Laigret, F., & Whitcomb, R. F. (1993). Notes: Revised
Taxonomy of the Class Mollicutes: Proposed Elevation of a Monophyletic
Cluster of Arthropod-Associated Mollicutes to Ordinal Rank
(Entomoplasmatales ord. nov.), with Provision for Familial Rank To Separate
Species with Nonhelical Morphology (Entomoplasmataceae fam. nov.) from
Helical Species (Spiroplasmataceae), and Emended Descriptions of the Order
Mycoplasmatales, Family Mycoplasmataceae. *International Journal of*
Systematic and Evolutionary Microbiology, 43(2), 378-385,
doi:https://doi.org/10.1099/00207713-43-2-378.
van de Water, J. A. J. M., Coppari, M., Enrichetti, F., Ferrier-Pagès, C., & Bo, M. (2020).
Local Conditions Influence the Prokaryotic Communities Associated With the
Mesophotic Black Coral Antipathella subpinnata. [Original Research].
Frontiers in Microbiology, 11(2423), doi:10.3389/fmicb.2020.537813.
van de Water, J. A. J. M., Melkonian, R., Voolstra, C. R., Junca, H., Beraud, E., Allemand,
D., et al. (2017). Comparative Assessment of Mediterranean Gorgonian-
Associated Microbial Communities Reveals Conserved Core and Locally

Variant Bacteria. [journal article]. *Microbial Ecology, 73*(2), 466-478,
doi:10.1007/s00248-016-0858-x.
van de Water, J. A. J. M., Voolstra, C. R., Rottier, C., Cocito, S., Peirano, A., Allemand, D.,
et al. (2018). Seasonal Stability in the Microbiomes of Temperate Gorgonians
and the Red Coral *Corallium rubrum* Across the Mediterranean Sea. [journal
article]. *Microbial Ecology, 75*(1), 274-288, doi:10.1007/s00248-017-1006-y.
Viver, T., Orellana, L. H., Hatt, J. K., Urdiain, M., Díaz, S., Richter, M., et al. (2017). The
low diverse gastric microbiome of the jellyfish *Cotylorhiza tuberculata* is
dominated by four novel taxa. *Environmental Microbiology, 19*(8), 3039-
3058, doi:10.1111/1462-2920.13763.
Vohsen, S. A., Anderson, K. E., Gade, A. M., Gruber-Vodicka, H. R., Dannenberg, R. P.,
Osman, E. O., et al. (2020). Deep-sea corals provide new insight into the
ecology, evolution, and the role of plastids in widespread apicomplexan
symbionts of anthozoans. *Microbiome, 8*(1), 34, doi:10.1186/s40168-020-
00798-w.
Weiler, B. A., Verhoeven, J. T. P., & Dufour, S. C. (2018). Bacterial Communities in
Tissues and Surficial Mucus of the Cold-Water Coral *Paragorgia arborea*.
[Original Research]. *Frontiers in Marine Science, 5*(378),
doi:10.3389/fmars.2018.00378.
Wick, R. R., Schultz, M. B., Zobel, J., & Holt, K. E. (2015). Bandage: interactive
visualization of de novo genome assemblies. *Bioinformatics, 31*(20), 3350-
3352, doi:10.1093/bioinformatics/btv383.
Woo, S., Yang, S.-H., Chen, H.-J., Tseng, Y.-F., Hwang, S.-J., De Palmas, S., et al. (2017).
Geographical variations in bacterial communities associated with soft coral
Scleronephthya gracillimum. *PLOS ONE, 12*(8), e0183663,
doi:10.1371/journal.pone.0183663.
Yang, S. H., Lee, S. T., Huang, C. R., Tseng, C. H., Chiang, P. W., Chen, C. P., et al. (2016).
Prevalence of potential nitrogen‐fixing, green sulfur bacteria in the skeleton
of reef‐building coral I sopora palifera. *Limnology and Oceanography, 61*(3),
1078-1086.
Yang, Z. (1995). A space-time process model for the evolution of DNA sequences.
Genetics, 139(2), 993-1005.
Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F. O., Ludwig, W., Schleifer, K.-H., et al.
(2014). Uniting the classification of cultured and uncultured bacteria and
archaea using 16S rRNA gene sequences. *Nature Reviews Microbiology, 12*(9),
635-645.
795

Figure 1. Map of sampling locations. Black circles mark locations where *Callogorgia*

delta was sampled and gray diamonds for *C. americana*. Below site names, the depth

range of collected samples is shown followed by the number of colonies, sediment, and

niskin water samples in parentheses. If no water or sediment were collected, only the

number of colonies is shown. A star signifies whether a 400L water sample was taken in

addition to water collected with niskins. Bathymetric map obtained from BOEM (credit:

803   William Shedd, Kody Kramer). Image of *C. americana* and electron micrographs taken

804   from (Quattrini et al. 2013).

806   Figure 2. The relative abundances of three ASVs corresponding to novel mollicutes in

807   *Callogorgia delta* and *C. americana.* Each bar represents the microbial composition of a

808   single colony and are organized by species, site, and sampling year.

Figure 3. Maximum likelihood phylogenetic trees of class Mollicutes using the 16S

rRNA gene (a) and amino acid sequences of 43 genes (b). UFBoot support values are

reported at each node. Numbers in parentheses designate the number of sequences within

collapsed nodes. *Ca. Oceanoplasma callogorgiae* and *Ca. Thalassoplasma callogorgiae*

are presented in bold text and underlined.

Figure 4. Metabolic map of the arginine dihydrolase pathway encoded in the genomes of

both *Ca. Oceanoplasma callogorgiae* and *Ca. Thalassoplasma callogorgiae.* Genes are

in bold and underlined text. arcA = arginine deiminase, arcB = ornithine

carbamoyltransferase, arcC = carbamate kinase, arcD = arginine/ornithine antiporter.

Figure 5. Bacterial signals in the mesoglea of *Callogorgia delta* using CARD-FISH with

a general eubacterial probe (EUB I-III) and DAPI staining of DNA. Fluorescence in the

emission range of DAPI is shown in blue while green shows emission in the range of the

fluorophore ALEXA 488 used by both EUB I-III and control probes. Paired images

compare adjacent sections where left images display sections hybridized with EUB I-III

and show bacterial signals (a,c,e) while right images display sections hybridized with a

negative control probe and only show autofluorescence (b,d,f). Signals were observed in

transverse sections through the central axis (a,b) and transverse sections through a single

polyp (c,d,e,f,g). Arrows in (a) and (c) show examples of EUB I-III fluorescence. A =

central axis, G = gastrovascular cavity, M = gastric mesentery, L = sclerite lacuna.

Figure 6. Higher magnification of bacterial signals in the mesoglea and tentacles of

Callogorgia delta using CARD-FISH with a general eubacterial probe (EUB I-III) and

DAPI staining of DNA. Fluorescence in the emission range of DAPI is shown in blue

while green shows emission in the range of the fluorophore ALEXA 488 used by both

EUB I-III and control probes. Paired images compare adjacent sections where left images

display sections hybridized with EUB I-III and show bacterial signals (a,c,e) while right

images display sections hybridized with a negative control probe and only show

autofluorescence (b,d,f). Transverse sections through the central axis (a,b) exhibit EUB I-

III fluorescence (a, arrows) in the surrounding mesoglea and an overlap of DAPI (c,

arrow). (a, arrow). Transverse sections through a polyp (c,d) show EUB I-III fluorescence

(c, arrows) in the base of a tentacle. Transverse sections through the body wall of a polyp

show granular (<1µm) EUB I-III fluorescence in the mesoglea (e,f,g). A = central axis, T

= tentacle.