Title:

Discovery of deep-sea coral symbionts from a novel family of marine bacteria,

Oceanoplasmataceae, with severely reduced genomes

Authors:

Samuel A. Vohsen, Harald R. Gruber-Vodicka, Nicole Dubilier, Charles R. Fisher, Iliana B.

Baums*

Abstract:

Microbes perform critical functions in corals yet most knowledge is derived from the

photic zone. Here, we discovered two mollicutes that dominate the microbiome of the deep-sea

octocoral, *Callogorgia delta,* and reside in the mesoglea. These symbionts were abundant across

the host's range, absent in the water, and rare in sediments. The symbionts lack all known

fermentative capabilities including glycolysis and can only generate energy from arginine

provided by the coral host. Their genomes feature extensive mechanisms to interact with foreign

DNA which may be indicative of their role in symbiosis. We erect the novel family

Oceanoplasmataceae which includes these symbionts and others associated with four marine

invertebrate phyla. Its exceptionally broad host range suggests that the diversity of this enigmatic

family remains largely undiscovered. Oceanoplasmataceae genomes are the most highly reduced

among mollicutes providing new insight into their reductive evolution and the roles of coral

symbionts.

Keywords:

Deep-sea corals, Mollicutes, symbiosis, microbiome, mesoglea

Main:

Introduction:

Corals are foundation species that support diverse animal communities from shallow waters to the deep sea. Corals also associate with a diversity of microbes[1] including the well-studied algal symbionts of the family Symbiodiniaceae and other microbial taxa that perform a variety of roles from providing nitrogen through fixation[2] to causing disease[3]. Most of these microbes were identified in scleractinian corals from the photic zone, while studies investigating the roles of microbes in octocorals and deep-sea corals are rarer because of the limitations that great depths impose on sampling and experimentation. Such work requires the use of remotely operated vehicles or submersibles launched from ships. Still, studies have demonstrated that octocorals and deep-sea corals host some of the same associates as shallow-water scleractinians such as corallicolid apicomplexans[4-7] and *Endozoicomonas*[8-11]. Deep-sea octocorals also host associates that are rare or absent in shallow-water and/or scleractinian corals. This includes bacteria from the SUP05 cluster whose role is linked to cold seeps in the deep sea[12] and commonly members of the class Mollicutes.

Members of the class Mollicutes associate with many coral species and with a wide diversity of plant, fungal, and animal hosts including humans[13,14]. Several members are well-studied parasites[15], while the impact of others on their hosts are unclear such as the ubiquitous intracellular symbionts of arbuscular mycorrhizal fungi[16-18]. Yet others are mutualists such as *Spiroplasma* spp. that infect insects and confer protection against nematodes, parasitoid wasps, and fungi through the production of unique toxins[19]. Reflecting their lifestyles as symbionts, their evolutionary history is dominated by genome reduction with some species considered to have the most reduced genomes capable of supporting cellular life[20,21]. Recently, novel and

46   divergent mollicutes were discovered in diverse marine invertebrates including a deposit-feeding

47   holothurian from a deep-sea trench[22]; pelagic, photosynthetic jellyfish[23,24]; wood-boring, deep-

48   sea chitons[25]; ascidians from a coastal lagoon[26], and crown-of-thorns sea stars from the Great

49   Barrier Reef[27]. However, the impacts these symbionts have on their hosts remain unclear.

50   Among coral species, mollicutes are most commonly found in octocorals[28-37], but have

also been reported in black corals[38] and stony corals[39,40]. These corals occupy a wide breadth of

habitats from shallow-water reefs[34,37,41], through the mesophotic zone[38] and the deep sea[29,30,39] to

abyssal depths[42]. In many of these coral species, *Mycoplasma* spp. are the dominant microbes,

comprising more than 50% of their associated microbial communities[30,37,41]. Interestingly, the

relative abundances of *Mycoplasma* vary substantially across the ranges of individual coral

species[36,41]. Thus, this association may be influenced by environmental conditions or otherwise

shaped by geography such as through limited dispersal. In addition, *Mycoplasma* spp. exhibit

varying degrees of host specificity[31,36,37]. In cases with high specificity, *Mycoplasma* spp. may

have coevolved with their coral hosts and closely interact with them as symbionts. Despite these

insights, the interactions between mollicutes and their host corals remain unknown.

61   Here we discovered two novel bacteria of the class Mollicutes which are abundant in the

62   deep-sea octocoral *Callogorgia delta. C. delta* is common along the continental slope in the Gulf

63   of Mexico between 400 m and 900 m depth, where it is often the dominant habitat-forming coral

64   species[43,44]. Like many deep-sea coral species, *C. delta* colonies create habitat for numerous

65   animal species including *Asteroschema* ophiuroids and the chain catshark, *Scyliorhinus retifer,*

66   which lay their eggs directly on *C. delta* colonies[45].

67   To characterize the association between *Callogorgia delta* and these newly discovered

mollicutes, we screened colonies from six sites in the Gulf of Mexico across three years of

sampling (Fig 1a) using 16S metabarcoding to determine the prevalence of these novel

mollicutes and quantify their relative abundances. We also screened the closely related species

Callogorgia americana to assess the phylogenetic breadth of the association. *C. americana,* is

also a dominant habitat-forming species in the northern Gulf of Mexico and provides a good

comparison species. *C. americana* occurs at shallower depths (300 – 400m) while *C. delta* is

often found near hydrocarbon seeps[44].

We determined the specificity of the association to corals by screening sediment and

water in addition to *Callogorgia*. Further, we assembled the genomes of these novel mollicutes

to assess their metabolic capabilities, determine their phylogenetic positions, and compare them

to other mollicutes. Finally, we located these mollicutes within coral tissue using catalyzed

reporter deposition fluorescence in situ hybridization (CARD-FISH) microscopy.

Results:

Occurrence of novel mollicutes in and around Callogorgia

We investigated symbiont diversity using V1-V2 16S rRNA-based metabarcoding on an

Illumina MiSeq. Three abundant amplicon sequence variants (ASVs) were identified among 108

Callogorgia delta colonies that belong to the class Mollicutes (Molli-1,2,3). Molli-1 was the

most prevalent ASV (Fig. 1b). It was detected in 99 out of 108 colonies and was present in

samples from all three sampling years (2015-2017) and all six sampling locations (Fig. 1b). It

had an average relative abundance of 78% among corals which harbored it and reached a

maximum of 99% of the microbial community in some samples. It was also present in all five

Callogorgia americana colonies analyzed where it averaged 91% of the microbial community.

Molli-2 differed from Molli-1 by a single base pair among 300. This closely related ASV was

detected in 4 of 10 colonies from the deepest site, GC290. Among those four colonies, Molli-2

averaged 90% of the microbial community while Molli-1 was undetected (Fig. 1b).

Molli-3, which was 72% identical to Molli-1, was prevalent and abundant in some

colonies, but less so compared to Molli-1 (Fig. 1b). Molli-3 was present in 69 out of 108 *C. delta*

colonies (up to 82% of the community) and three out of five *C. americana* colonies (up to 15%).

However, Molli-3 was absent in all *C. delta* colonies from the deepest sites GC249 (n=14) and

GC290 (n=10).

None of these ASVs were detected in any water sample (n=30). However, Molli-1 and

Molli-3 were detected in 35 and 11 out of 47 total sediment samples with maximum relative

abundances of 1.1 and 0.16% of the sediment microbial community respectively.

Phylogenetic positions of the mollicute in Callogorgia delta

Twenty-two full length 16S rRNA gene sequences that correspond to Molli-1 were

assembled from separate *C. delta* metagenomes and three additional sequences were assembled

that correspond to Molli-3. These sequences were distinct from those of other known mollicutes

and clustered with sequences from recently discovered associates of marine invertebrates

(UFBoot 100%, Fig. 2a). Interestingly, the two novel mollicutes in *Callogorgia delta* are not

each other's closest relative and shared only 84% sequence identity across the entire 16S rRNA

gene which is below the cutoff which has been suggested to differentiate bacterial families[46].

A circular and closed genome corresponding to Molli-1 was assembled from a deeply

sequenced metagenome of *Callogorgia delta.* The metagenome was filtered to kmers with a

depth of 1,000 or greater and the corresponding reads were subsampled at a rate of 10% and

assembled with SPAdes[47] to produce a single circular scaffold. Additionally, a genome

corresponding to Molli-3 was obtained by co-assembling the two metagenomes with the highest

16S coverage and binning contigs using coverages from all 26 metagenomes. Phylogenomic

analysis of shared amino acid sequences among mollicute genomes reiterated phylogenetic

inferences based on 16S rRNA (Fig. 2b). A novel clade of Mollicutes that associates with marine

invertebrates was well-supported (UFBoot 100%) and this clade clustered alongside the

intracellular symbionts of mycorrhizal fungi with high confidence (UFBoot ≥98%). Branch

lengths between members of the novel clade of marine mollicutes were long and the average

nucleotide identities (ANI) between the four representative genomes ranged from 63 to 65%

(OrthoANI[48]) suggesting that both mollicutes described here are novel genera. We propose the

names *Candidatus Oceanoplasma callogorgiae* gen. nov. sp. nov. and *Ca. Thalassoplasma*

callogorgiae gen. nov. sp. nov. corresponding to Molli-1 and Molli-3 respectively. Further, we

erect the novel family Oceanoplasmataceae with *Ca. Oceanoplasma callogorgieae* as its type

species and which includes other associates of marine invertebrates.

Metabolic capabilities of the novel mollicutes in Callogorgia delta

The genome of *Ca. Oceanoplasma callogorgiae* was highly reduced in size (446,099 bp,

Table 1, Fig. 2c) and function with only 359 predicted protein-encoding genes (241 with KEGG

orthology annotations). The genome lacked any pathway to generate biomass or energy from

carbohydrates and instead contained the genes comprising the arginine dihydrolase pathway

(*arcABCD*) and ATP synthase. These genes were among the most highly expressed genes in

both RNA libraries (Fig. 3ab) and constitute the sole pathway in this genome that is capable of

producing ATP. This pathway generates ATP from the equimolar catabolism of arginine. In

other non-fermentative mollicutes, additional ATP is generated as the resulting ammonia

establishes a proton gradient which drives ATP synthase[49,50]. Besides arginine catabolism, a few

genes could potentially be involved in ATP production: glyceraldehyde-3-phosphate

dehydrogenase (*gapA*), phosphoglucomutase (*pgm*), ribose-phosphate pyrophosphokinase

(*prsA*), and acetate kinase (*ackA*).

The genome featured a very limited ability to synthesize essential compounds. It encoded

only one enzyme involved in amino acid synthesis (threonine dehydrogenase) but encoded

several protein-degrading enzymes and amino acid importers. Similarly, it encoded no enzymes

that synthesize vitamins, cofactors, or coenzymes besides NAD^+ as well as FMN and FAD from

imported riboflavin (Fig. 3a).

The genome contained extensive and active mechanisms to interact with foreign DNA

including type I, II, and III restriction modification systems and a Class 2 Type II-C CRISPR-

Cas system. One of the five most highly expressed genes in both libraries was the specificity

subunit S of its type I restriction modification system (Fig. 3b). Further, it contained the second

highest number of spacer sequences, at 107, of all mollicutes with available genomes[51,52].

Four predicted genes with no functional annotations were among the ten most highly

expressed (Fig. 3b). The highest of these (named U1) was predicted to contain a transmembrane

segment and a signal peptide (Hhblits[53]). It also contained a region that matches the EAL domain

of various proteins. The EAL domain is a diguanylate phosphodiesterase which degrades the

second messenger cyclic di-GMP[54-56]. Another highly expressed, unannotated gene (U2) was

predicted to contain a GIY-YIG endonuclease domain with high confidence (>95%, Hhblits).

The fourth most highly expressed, unannotated gene (U4) belongs to the YneF family whose

function is unknown but is conserved across Mollicutes and most Bacilli where it is highly

expressed and considered an essential protein[57].

The genome of *Ca. Thalassoplasma callogorgiae* was very similar to that of *Ca. O.*

callogorgiae. It was reduced in size (445,456 bp) and metabolic capabilities with only 385

predicted protein-coding genes (237 with KEGG orthology annotations). It possessed all the

pathways and genes mentioned above for *Ca. O. callogorgiae* except it lacked a type II

restriction modification system and homologs of U1 and U2. In contrast, it possessed ribose-5-

phosphate isomerase (*rpiB*) and alternative genes to synthesize NAD^+ (Fig. 3a).

Comparison to other mollicutes

The genomes of *Ca. O. callogorgiae* and *Ca. T. callogorgiae* were similar to others

within Oceanoplasmataceae. The genome of "*Ca. Spiroplasma holothuricola*" is publicly

available and we asssembled a fourth genome corresponding to the closely related associate of

the jellyfish *Cotylorhiza tuberculata*[23]. We assembled the publicly available metagenome with

the highest 16S coverage and used all metagenomes from the study for binning. These four

genomes shared 211 clusters of orthologous genes (COGs) which comprised 60-66% of the

COGs in each genome. Besides housekeeping genes, all four genomes encode the arginine

dihydrolase pathway and CRISPR-Cas systems. The genome of the jellyfish associate was the

most divergent and may have further but limited means to generate ATP since it encoded lactate

dehydrogenase, glycerate kinase, ribulose-phosphate 3-epimerase, and an agmatine/putrescine

antiporter.

Overall, the genomes of Oceanoplasmataceae were distinct from other mollicutes and

were the most reduced (Fig. 4). They formed a distinct cluster based on the presence/absence of

COGs (Fig. 4a), were the smallest (389 – 446 kbp versus >564 kbp[58]; Fig. 3c), and had the

fewest protein-encoding genes (pegs) (336 – 385 versus >515; Fig. 4b). Further, they were the

only mollicutes missing the genes necessary to conduct glycolysis besides *Moeniiplasma* spp.

with which they shared other physiological similarities. Compared to other mollicutes, they had

the fewest COGs with KEGG orthology terms and the fewest genes involved in carbohydrate

metabolism. Oceanoplasmataceae genomes were distinguishable from those of *Moeniiplasma*

spp. by the presence of genes encoding ribose-phosphate pyrophosphokinase, glyceraldehyde 3-

phosphate dehydrogenase, and ATP synthase.

All Oceanoplasmataceae genomes contained twenty-three COGs that were absent in all

other mollicutes. Notably, these included a Ca^{2+}/Na^+ antiporter, several other transporters, and a

thymidine kinase. Thymidine kinases were enriched in the genomes of Oceanoplasmataceae

which each contained four copies while all other genomes had one or zero except two

Moeniiplasma spp. which had two copies.

Localization of bacteria in Callogorgia delta

To locate the symbiotic bacteria, we used probes that bind to ribosomal RNA permitting

deposition of a fluorophore through catalyzed reporter deposition fluorescence in situ

hybridization (CARD-FISH). Fluorescence was observed in the mesoglea of two separate

Callogorgia delta colonies representing abundant bacteria (Fig. 5 and 6). These signals were

present in tissue sections hybridized with the EUB338 I-III probe mix but absent in adjacent

sections hybridized with the negative control probe (non-EUB). These signals were <1 µm in

diameter and overlapped DAPI fluorescence. These signals of bacterial symbionts were observed

in both the mesoglea within polyps (Fig. 5c,e,g) and in mesoglea within the coenosarc

surrounding the proteinaceous central axis (Fig. 5a). They formed large aggregates potentially

comprised of hundreds or thousands of cells (Fig. 6). Another fluorescence signal was observed

in the base of the tentacles in one colony but was far less aggregated (Fig. 6c,d).

Discussion:

Novel mollicutes in Callogorgia delta are symbionts

The data presented here suggests that *Ca. Oceanoplasma callogorgiae* and *Ca.*

Thallasoplasma callogorgiae are symbionts of *Callogorgia delta sensu*[59]. First, their association

was stable across geography and time. Additionally, these microbes seem to associate

specifically with corals in the genus *Callogorgia* since both were absent or rare in the water and

sediment but were also detected in the closely related species *C. americana*. Further, their

genomes were very reduced and demonstrated a reliance on compounds like arginine and amino

acids that are probably obtained from *C. delta.* Finally, *Ca. Oceanoplasma callogorgiae* likely

resides within the coral. Abundant bacteria were observed in the mesoglea of *C. delta* and *Ca. O.*

callogorgiae was the most abundant bacterium by far in both the amplicon dataset and the

metagenomes where the genomic coverage of the symbiont was up to 50 times as high as that of

the host.

Metabolic processes and other fundamental activities of the symbionts

Both *Ca. Oceanoplasma callogorgiae* and *Ca. Thalassoplasma callogorgiae* lacked

genes encoding complete fermentative pathways and depended on arginine to generate ATP.

They likely receive this arginine from their coral hosts as well as essential compounds such as

amino acids, riboflavin, and biotin. Conversely, they export ornithine and possibly short peptides

to their coral host.

Ca. Oceanoplasma callogorgiae possessed extensive mechanisms to defend against

viruses or other forms of foreign DNA. Its genome contained an extensive CRISPR array and

high expression of restriction modification systems and endonucleases. Further, it shared

additional genomic features with *Ca. Thalassoplasma callogorgiae* including *comEC* to import

foreign DNA and enrichment of thymidine kinase genes to salvage thymidine from DNA

degradation. The presence of these anti-viral characteristics in such highly reduced bacterial

genomes suggests that they play a central role in the lifestyles of these bacteria.

Other, not yet well-understood processes must underly the basic functioning of these

novel mollicutes since some of the most highly expressed genes had unknown functions. This

includes U1 which is likely membrane-bound and potentially degrades cyclic di-GMP. Cyclic di-

GMP is a second messenger involved in morphogenesis, motility, and virulence[60]. U1 may

respond to external, or host-derived, stimuli as part of signaling pathways that could be linked to

host innate immunity and symbiont recognition.

Residence in the Mesoglea

Ca. O. callogorgiae likely resides in the mesoglea of *C. delta.* The mesoglea of

cnidarians is composed of a largely acellular, mucoid matrix of collagen containing some cells

such as amoebocytes[61]. The mesoglea provides rigidity upon which the muscles of the polyp can

act but also facilitates the transport of nutrients[62,63]. Some compounds diffuse through the

mesoglea like glucose and fatty acids[62,63] while amoebocytes can also shuttle compounds such as

amino acids[62]. *Ca. O. callogorgiae* probably acquires compounds including arginine, other

amino acids, and the precursors of coenzymes such as riboflavin from the mesoglea because it

lacks the genes necessary to synthesize these.

The mesoglea in cnidarians is also a site of immune defense[64]. Bacteria that reside in the

mesoglea must evade the coral's immune defenses such as phagocytic amoebocytes[64]. A few

bacteria have been localized within the mesoglea of cnidarians including *Pseudomonas* and

spirochetes in *Hydra* spp.[65,66] and pathogenic cyanobacteria in the coral *Orbicella annularis*

suffering from black band disease[67]. In contrast, the only mollicute previously localized in corals

was *Mycoplasma corallicola* which was found on the surface of the tentacle ectoderm of

Lophelia pertusa[39]. Unlike *M. corallicola*, *Ca. O. callogorgiae* likely resides within its host coral

and therefore must somehow evade phagocytic amoebocytes in the mesoglea and potentially

other immune defenses.

Consequences for the coral host

It is still unclear if *Ca. Oceanoplasma callogorgiae* incurs a cost or provides any benefit

to its coral host. These novel mollicutes may simply be commensals or parasites with a minimal

impact on their host corals since they did not display any obvious pathology and nearly every

colony was infected. Conversely, the metabolism of arginine may provide the coral with an

alternative pathway to process nitrogenous waste or could function to recycle nitrogen which

would be useful for deep-sea corals because they rely on a nitrogen-poor diet of marine snow[68].

The symbiont might provide protection from viruses using its CRISPR-Cas and restriction

modification systems similar to what has been proposed for the symbiont of hadal sea cucumbers

which was the closest relative to *Ca. O. callogorgiae*[22]. It is also possible that some of the

unannotated genes are peptides that confer protection from pathogens or parasites like those that

Spiroplasma spp. produce in insects[19].

268
The novel family Oceanoplasmataceae

We erect the family Oceanoplasmataceae which includes *Ca. Oceanoplasma*

callogorgiae, *Ca. Thalassoplasma callogorgiae*, and other associates of marine invertebrates.

These hosts are phylogenetically diverse comprising four animal phyla, exhibit diverse lifestyles

from being photosynthetic to wood-borers, and originate from a wide range of habitats from the

epipelagic zone to hadal trenches[22-26]. In addition to this, the large phylogenetic distances

between members of the Oceanoplasmataceae suggests that a substantial amount of diversity

remains undiscovered. Interestingly, *Ca. Moeniiplasma* spp. appear to be the closest relatives of

Oceanoplasmataceae. They clustered together in the phylogenetic analyses and shared genomic

similarities such as a lack of glycolysis.

Compared to other mollicutes, Oceanoplasmataceae exhibit extensive viral defense

systems. All were enriched in thymidine kinase genes and the two genomes with unfragmented

CRISPR arrays had more spacers than other mollicutes. Only the associate of a hadal snailfish,

"*Mycoplasma*" *liparidae,* had more spacers[52]. This may reflect a general evolutionary trend

among animal-microbial symbioses in marine environments where exposure to viruses is high.

Similarly, the symbionts of marine sponges are enriched in CRISPR-Cas systems compared to

surrounding seawater [69,70].

Oceanoplasmataceae possess the most reduced genomes among all mollicutes. They

possess the smallest genomes which contain the fewest protein-encoding genes and lack any

complete fermentative pathway. Extensive genome reduction dominates the evolutionary history

of Mollicutes. Mollicute genomes are used to identify the minimum set of genes essential for

cellular life[21] and are utilized as precursors in efforts to design minimal genomes[71,72]. These

synthetic genomes retain glycolysis. However, the lack of glycolysis in Oceanoplasmataceae and

Moeniiplasma spp. suggests that these synthetic genomes could be reduced further. Our

comparative genomics analysis also revealed that *Moeniiplasma* spp. lack ATP-synthase

suggesting alternative gene sets may permit further reduction.

Strangely, no mollicutes except *Acholeplasma laidlawii* are known to use cyclic di-GMP

and all mollicutes instead use cyclic di-AMP[73]. The presence of an EAL domain in gene U1 of

Ca. O. callogorgiae implies that the use of cyclic di-GMP was not lost in all other mollicutes or

was regained through horizontal gene transfer in some lineages.

Conclusion

Here we describe two novel members of the Mollicutes that associate with the deep-sea

coral, *Callogorgia delta,* and propose the names *Ca. Oceanoplasma callogorgiae* and *Ca.*

Thalassoplasma callogorgiae. We characterize their association with *C. delta,* generate genomic

resources, and produce phylogenetic and metabolic inferences advancing our understanding of

coral-associated mollicutes and symbiosis in the deep sea. Further, we erect a new family,

Oceanoplasmataceae, which has not yet been recognized and whose diversity remains largely

uncharacterized. We show that its genome reduction exceeds what was known among Mollicutes

informing their evolutionary history, the consequences of symbiosis, and the concept of minimal

bacteria.

Methods:

Collections

Collections are a subset of those described in Vohsen et al. 2020. One hundred and eight

Callogorgia delta colonies were collected from six sites spanning over 350 km in the northern

Gulf of Mexico in 2015, 2016, and 2017 (Fig. 1). The sites are named after the Bureau of Ocean

Energy Management's designations for lease blocks in which corals were sampled and include

Mississippi Canyon (MC) 751 (n=19 colonies, 435-443 m, 28.193 N -89.800 W), Viosca Knoll

(VK) 826 (n=3 colonies, 547-559 m, 29.159 N -88.010 W), Green Canyon (GC) 234 (n=37

colonies, 505-536 m, 27.746 N -91.122 W), MC885 (n=25 colonies, 624-642 m, 28.064 N -

89.718 W), GC249 (n=14 colonies, 789-795 m, 27.724 N -90.514 W), and GC290 (n=10

colonies, 852 m, 27.689 N -90.646 W). To compare *Callogorgia delta* to a closely related coral

species, we additionally processed five *Callogorgia americana* colonies that were collected in

2010 from Garden Banks (GB) 299 (n=3 colonies, 358-359m, 27.689 N -92.218 W) and VK862

(n=2 colonies, 352-357m, 29.109 N -88.387 W).

Colonies were sampled using specially designed coral cutters mounted on the

manipulator arm of remotely operated vehicles (ROVs). Coral fragments were removed, placed

in separate temperature insulated containers until recovery of the ROV, and were maintained at 4

°C for up to 4 hours until preservation in ethanol or freezing in liquid nitrogen.

Sediment samples were taken in close proximity to many of the *Callogorgia delta*

collections using push cores with a diameter of 6.3 cm. Upon recovery of the ROV, 1 mL of

sediment from the top 1 cm of each sediment core was frozen in liquid nitrogen for later DNA

extraction for microbiome analysis. Water was sampled in 2015 using a Large Volume Water

Transfer System (McLane Laboratories Inc., Falmouth, MA) which filtered approximately 400 L

of water through a 0.22 micron porosity filter with a diameter of 142 mm. One quarter of each

filter was preserved in ethanol and another quarter was frozen in liquid nitrogen for these

analyses. In 2016 and 2017, 2.5 L Niskin bottles mounted on the ROV captured water above

corals. Upon recovery of the ROV, this water was filtered through 0.22 micron filters which

were frozen in liquid nitrogen.

16S rRNA metabarcoding

The microbiomes of *Callogorgia delta, Callogorgia americana*, water, and sediment

were analyzed using 16S metabarcoding. DNA was extracted from coral tissue and sediment

samples using DNeasy PowerSoil kits (Qiagen, Hilden, Germany) following manufacturer

protocols using approximately 1 cm of coral branches and about 0.25 g of sediment. DNeasy

PowerSoil kits were also used for water samples collected in 2015 using 1 cm^2 of filter. For all

water samples from 2015, replicate extractions were performed on quarters of the filter preserved

in both ethanol and frozen in liquid nitrogen. DNA was extracted from all other water samples

using Qiagen DNeasy PowerWater kits.

The V1 and V2 regions of the 16S rRNA gene were amplified using universal bacterial

primers 27F and 355R[74] with CS1 and CS2 adapters (Illumina, San Diego, CA). PCR was

conducted with the following reaction composition: 0.1 U/ µL Gotaq (Promega, Madison, WI),

1X Gotaq buffer, 0.25mM of each dNTP, 2.5 mM $MgCl_2$, and 0.25 µM of each primer, and the

following thermocycler conditions: 95 °C for 5 min; 30 cycles of 95 °C for 30 sec, 51 °C for 1

min, and 72 °C for 1 min; and finally 72 °C for 7 min. Libraries were prepared by the University

of Illinois Chicago DNA services facility and sequenced on two separate runs on an Illumina

MISEQ platform[75]. Raw sequence data are available on the NCBI database under BioProject

PRJNA565265 and BioSample IDs SAMN12824733 – 4765, SAMN12824767 – 4875.

Amplicon sequence data were analyzed as described in[4] using QIIME 2 (ver2017.11)[76].

Reads were joined using vsearch and quality filtered using q-score-joined. Deblur denoise was

used to detect chimeras, construct sub-operational taxonomic units (referred to as ASVs), and

trim assembled reads to 300bp.

Metagenomes and Metatranscriptomes

Metagenomes and metatranscriptomes of *C. delta* were sequenced to assemble genomes

of novel mollicutes which were used to assess their phylogenetic positions and metabolic

potential. DNA was extracted from eight *Callogorgia delta* colonies as described in Vohsen et al.

2020. In brief, DNA was extracted using DNA/RNA allprep kits (Qiagen, Hilden, Germany)

with β-mercaptoethanol added to the RLT+ lysis buffer following the manufacturer's

instructions. These DNA extracts were sequenced on an Illumina HISEQ2500 platform using

150 bp paired-end reads. The two libraries with the highest coverage of mollicutes were

sequenced a second time. In addition to these, DNA was extracted from an additional 18 samples

corresponding to 16 additional colonies. These were sequenced on an Illumina HISEQ platform

with 100bp paired-end.

RNA was extracted from two *Callogorgia delta* colonies using an RNeasy extraction kit

(Qiagen, Hilden, Germany). The RNA extracts were enriched for bacteria by depleting coral

host rRNA using a Ribo-Zero Gold Yeast kit (Illumina, San Diego, CA, USA) following the

manufacturer's protocols. The RNA extracts were sequenced on a MISEQ platform with 150 bp

single-end reads.

All DNA sequence libraries were screened for bacterial SSU rRNA using phyloFlash

ver3.3[77]. The library with highest coverage of *Ca. Oceanoplasma callogorgiae* was used to

assemble its genome. Reads were trimmed to a quality score of 2 from both ends using BBduk

and a kmer frequency analysis was performed using BBnorm. The library was kfiltered to a

depth of 1,000 or greater using BBnorm and downsized to 10% using reformat with a single

pass. The library was then assembled using SPAdes[47] with a kmer progression of

21,33,55,77,99,127. Bandage ver0.8.1[78] was used to identify a single circular scaffold of contigs

containing the 16S rRNA gene of *Ca. Oceanoplasma callogorgiae*. The overlaps on the end were

manually trimmed off and the scaffold was recentered to the origin of replication predicted by

Ori-Finder[79]. This genome was then annotated using classic RAST ver2.0[80,81].

In order to generate a genome for *Ca. Thalassoplasma callogorgiae*, the two libraries

with the highest coverage of its 16S rRNA gene were coassembled using megahit[82]. All libraries

were mapped to this coassembly using BBmap with a kfilter of 22 bp, subfilter of 15 bp, and a

maxindel of 80 bp. Alignments were converted from SAM to BAM formats, sorted using

samtools[83], and used for binning using metabat[84] with default parameters. All scaffolds with

contigs belonging to bin one constituted a draft genome that was annotated using RAST and used

for phylogenomic analysis.

The presence of CRISPRs and associated genes along with the number of spacers were

determined using RAST annotations and CRISPRfinder[85,86]. OrthoANI[48] was used to estimate

average nucleotide identity between genomes. To determine which genes were expressed and to

quantify their expression, the two RNA sequence libraries were pseudoaligned to the DRAM

gene annotations described below of both genomes with kallisto ver0.44.0[87] using 100 bootstrap

replicates, an estimated fragment length of 180 bp with a standard deviation of 20 bp, and an

index with a k-mer size of 31 bp. Coding regions with high expression levels but no functional

annotations from RAST were further investigated using blastp and HHBlits[53] to inform potential

function.

Phylogenetic Analyses

Phylogenetic trees were constructed to infer the phylogenetic positions of the mollicutes

that were detected in *Callogorgia delta.* First a tree of 16S sequences was generated to place

these novel mollicutes within the wide diversity of mollicutes with available 16S sequences. Full

length 16S rRNA genes were assembled from the metagenomes using phyloFlash ver 3.3[77].

These full-length sequences were aligned with publicly available Mollicutes sequences that were

over 1,000 bp using MUSCLE ver3.8.425[88]. A maximum likelihood tree was constructed using

the IQ-TREE web server ver1.6.10[89,90]. Within IQ-TREE, ModelFinder[91] was used to choose a

general time reversible model with unequal rates and empirical base frequencies as well as a

FreeRate model[92,93] for rate heterogeneity across sites with 3 categories (GTR+F+R3) based on

the highest Bayesian information criterion (BIC) score. UFBoot2[94] was used to obtain ultrafast

bootstrap support values (UFBoot) using 1000 replicates.

To obtain a more confident phylogenetic placement of the coral-associated mollicutes, a

phylogenomic tree was constructed including genomes from the two mollicutes in *C. delta* and

66 mollicute genomes accessed through GenBank and the European Nucleotide Archive (ENA)

(Supplementary File 1g). One more mollicute genome was assembled from a dataset available on

ENA because an assembled genome was not available. This genome corresponded to the novel

bacterium that associates with the jellyfish, *Cotylorhiza tuberculata*, which was tentatively

identified as *Mycoplasma*[23]. Among the four libraries, M3 was chosen for assembly since it had

the highest coverage of *Mycoplasma* based on the coverages of 16S rRNA assembled using

phyloFlash[77]. This library was then assembled using megahit[82] with a kmax of 241. All four

libraries were mapped to this assembly using bbmap with a kfilter of 22 bp, subfilter of 15 bp,

and maxindel of 80 bp. The alignments were subsequently used to bin the assembly using

metabat[84] using default parameters. Bin one corresponded to the purported *Mycoplasma* genome.

This bin and all associated scaffolds were used to represent this genome in phylogenomic

analyses and was also annotated using RAST to compare to other genomes. A concatenated

amino acid sequence alignment of 43 genes from all 69 genomes was generated with CheckM

(ver1.1.3)[95]. A maximum likelihood phylogenomic tree was then constructed using IQ-TREE

(ver1.6.12)[89,94]. Both phylogenetic trees were annotated using the recently revised phylogeny of

the class Mollicutes[96-98].

Comparative Genomics Analyses

All mollicute genomes included in phylogenetic analyses that use translation table 4 (all

except Acholeplasma-Anaeroplasma-Phytoplasma group) were annotated using DRAM[99].

Clusters of orthologous genes (COGs) were identified among all genomes using orthoMCL[100] on

KBase[101] with the "BuildPangenome with OrthoMCL" app (version 2.0) using amino acid

translations from all protein-encoding genes (pegs) and default parameters. To visualize

differences in genomic composition, non-metric multidimensional scaling was performed using

Jaccard distances based on the presence/absence of COGs. Only COGs that were present in more

than two genomes were included.

Microscopy

CARD-FISH was employed to localize bacteria within *Callogorgia delta.* Colonies

collected in 2016 were preserved for microscopy. During subsampling, 0.5 cm long branches

were fixed in 2 mL of 4% formaldehyde with 1X phosphate-buffered saline (PBS) at 4 °C

overnight (4-8 hours). Samples were washed 3 times and then stored in equal parts ethanol and

1X PBS for up to 21 months. Samples were decalcified in 0.45M EDTA in 1X PBS for 7 days,

refreshing the buffer after 4 days, then washed in 1X PBS. Samples were post-fixed in 1X PBS

with 1% paraformaldehyde for 1 hour then washed 3 times in 1X PBS and stored in 50:50 1X

PBS:ethanol overnight. The next day, the samples were embedded in TissueTek (Sakura Finetek,

Maumee, OH) and left at 4 °C overnight. The next day, the TissueTek was replaced and the

samples were frozen. Samples were sectioned to a thickness of 4 µm using a Leica (Wetzlar,

Germany) cryostat CM3050 S. Adjacent sections were placed on polysine slides and stored at 4

°C before hybridization.

Hybridization began by removing TissueTek from tissue sections via soaking slides in

milliQ water for 3 minutes. Autofluorescence was then diminished by soaking slides in ethanol

with 0.1% Sudan black B for 10 minutes. Sections were washed three times in 1X PBS and left

in 1X PBS for a final 24 minutes. Endogenous peroxidases were inactivated by soaking sections

in 0.2M HCl for 12 minutes, then 20 mM Tris-HCl for 20 minutes, lysozyme solution (0.01

g/mL) for 30 minutes at 37 °C, 20 mM Tris-HCl again for 20 minutes, and finally 3 minutes in

milliQ water. Slides were then airdried and sections were circled with a PAP pen. Two probes

were used: a mix of EUB338 I-III which target eubacteria[102,103] and a negative control probe

(non-EUB) consisting of the reverse complements. Hybridization mixes were created by

combining 1 µL of probe (50 ng/µL) with 299 µL hybridization buffer (30% formamide with

0.102 M NaCl, 0.02 M Tris HCl, 10% Blocking Reagent, 0.01% sodium dodecyl sulfate, and 0.1

g/mL dextran sulfate). Each section was covered with approximately 10-20 µL of one of the

hybridization mixes. Two adjacent sections were hybridized with the EUB338 I-III probe mix

and non-EUB probes separately to produce a negative control of the same tissue region. Each

slide was then placed in a humidity chamber consisting of a closed 50 mL tube with a kimwipe

soaked in 2 mL of 30% formamide in milliQ water. Hybridization was accomplished by

incubating slides at 46 °C for 2 hours within their humidity chambers.

471   After hybridization, slides were washed for 15 minutes at 48 °C in washing buffer

(0.102M NaCl, 0.02M Tris HCl, 0.005M EDTA, 0.01% SDS) followed by 15 minutes in 1X

PBS at room temperature. A fresh amplification mix was created by first combining 299 µL 1X

PBS and 1 µL H_2O_2, then adding 10 µL of that solution to 1 mL of amplification buffer (1X

PBS, 0.01% Blocking Reagent, 2M NaCl, 0.1 g/mL dextran sulfate) and 1 µL of labelled

tyramide (1mg/mL ALEXA 488-labelled tyramide in dimethylformamide with 20 mg/mL p-

iodophenylboronic acid). Slides were again placed in 50 mL humidity chambers with a kimwipe

soaked in 2 mL of milliQ water. Amplification was accomplished by incubating slides at 37 °C

for 30 minutes in these humidity chambers.

480   After amplification, slides were washed by dipping in 1X PBS followed by 10 minutes in

1X PBS, and a final dip in milliQ water. Slides were air dried and sections were stained with

DAPI (1 µg/mL) for 10 minutes in the dark. Slides were washed a final time in milliQ water for

3 minutes. Once airdried, slides were mounted using Vectashield (Vector Laboratories,

Burlingame, CA) and viewed on an Axioscope epifluorescence microscope (Carl Zeiss AG,

Oberkochen, Germany) with filters restricted to the emission spectra of ALEXA 488 and DAPI.

Images were taken with a Zeiss AxioCam camera and visualized using AxioVision software.

CARD-FISH signals were considered positive if they appeared consistently within a structure in

the coral, were absent in the same structures in negative control sections, and overlapped DAPI

signals.

Data availability:

Raw sequence data are available on the NCBI Sequence Read Archive under BioProjects

PRJNA574146 (BioSample IDs SAMN12856800 – SAMN12856807) and PRJNA565265 (SRA

accession numbers SRR10174410 and SRR10174411). Assembled 16S rRNA sequences and

genomes are available under accession numbers OR679038 – 62, CP125803, and

JARVCM000000000. Raw microscopy images are available upon request. Data used to

construct figures are provided in Supplementary File 1.

Code availability:

No custom scripts were developed for this manuscript that are central to the main findings. Code

used in this manuscript followed manual guidelines with details described in the methods. Scripts

are available upon request from the corresponding author.

References:

1   Bourne, D. G., Morrow, K. M. & Webster, N. S. Insights into the coral microbiome: Underpinning the health and resilience of reef ecosystems. *Annual Review of Microbiology* **70**, 317-340 (2016).

2   Lesser, M. P. *et al.* Nitrogen fixation by symbiotic cyanobacteria provides a source of nitrogen for the scleractinian coral *Montastraea cavernosa*. *Marine Ecology Progress Series* **346**, 143-152 (2007).

3   Ben-Haim, Y. *et al. Vibrio coralliilyticus* sp. nov., a temperature-dependent pathogen of the coral Pocillopora damicornis. *International journal of systematic and evolutionary microbiology* **53**, 309-315 (2003).

4   Vohsen, S. A. *et al.* Deep-sea corals provide new insight into the ecology, evolution, and the role of plastids in widespread apicomplexan symbionts of anthozoans. *Microbiome* **8**, 34 (2020).

5   Kwong, W. K., del Campo, J., Mathur, V., Vermeij, M. J. A. & Keeling, P. J. A widespread coral-infecting apicomplexan with chlorophyll biosynthesis genes. *Nature* **568**, 103-107 (2019).

6   Kirk, N. L. *et al.* Tracking transmission of apicomplexan symbionts in diverse caribbean corals. *Plos One* **8**, 10 (2013).

7   Kirk, N. L., Thornhill, D. J., Kemp, D. W., Fitt, W. K. & Santos, S. R. Ubiquitous associations and a peak fall prevalence between apicomplexan symbionts and reef corals in Florida and the Bahamas. *Coral Reefs* **32**, 847-858 (2013).

8   Pollock, F. J. *et al.* Coral-associated bacteria demonstrate phylosymbiosis and cophylogeny. *Nature Communications* **9**, 4921 (2018).

9   Kellogg, C. A. & Pratte, Z. A. Unexpected diversity of *Endozoicomonas* in deep-sea corals. *Marine Ecology Progress Series* **673**, 1-15 (2021).

10   Dannenberg, R. P. Characterization and oil response of the deep sea coral-associated microbiome. (2015).

11   Galand, P. E., Chapron, L., Meistertzheim, A.-L., Peru, E. & Lartaud, F. The effect of captivity on the dynamics of active bacterial communities differs between two deep-sea coral species. *Frontiers in Microbiology* **9**, 2565 (2018).

12   Vohsen, S. A. *et al.* Deep-sea corals near cold seeps associate with chemoautotrophic bacteria that are related to the symbionts of cold seep and hydrothermal vent mussels. *bioRxiv*, 2020.2002.2027.968453 (2020).

13   Ing-Ming Lee, Robert E. Davis & Gundersen-Rindal, D. E. Phytoplasma: Phytopathogenic mollicutes. *Annual Review of Microbiology* **54**, 221-255 (2000).

14   Tully, J. G., Bové, J. M., Laigret, F. & Whitcomb, R. F. Notes: Revised taxonomy of the class Mollicutes: Proposed elevation of a monophyletic cluster of arthropod-associated mollicutes to ordinal rank (Entomoplasmatales ord. nov.), with provision for familial rank to separate species with nonhelical morphology (Entomoplasmataceae fam. nov.) from helical species (Spiroplasmataceae), and emended descriptions of the order Mycoplasmatales, family Mycoplasmataceae. *International Journal of Systematic and Evolutionary Microbiology* **43**, 378-385 (1993).

15   Atkinson, T. P. in *The Microbiome in Rheumatic Diseases and Infection* Mollicutes: *Mycoplasma pneumoniae*. (eds Ragab G., Atkinson T. P., & Stoll M. L.) 103-111 (Springer International Publishing, 2018).

16 Bonfante, P. & Desirò, A. Who lives in a fungus? The diversity, origins and functions of
fungal endobacteria living in Mucoromycota. *The Isme Journal* **11**, 1727 (2017).
17 Desiro, A., Faccio, A., Kaech, A., Bidartondo, M. I. & Bonfante, P. *Endogone*, one of the
oldest plant-associated fungi, host unique mollicutes-related endobacteria. *The New*
phytologist **205**, 1464-1472 (2015).
18 Kuga, Y., Wu, T.-D., Sakamoto, N., Katsuyama, C. & Yurimoto, H. Allocation of carbon
from an arbuscular mycorrhizal fungus, *Gigaspora margarita*, to its gram-negative and
positive endobacteria revealed by high-resolution secondary ion mass spectrometry.
Microorganisms **9**, 2597 (2021).
19 Ballinger, M. J. & Perlman, S. J. The defensive *Spiroplasma*. *Current opinion in insect*
science **32**, 36-41 (2019).
20 Sirand-Pugnet, P., Citti, C., Barré, A. & Blanchard, A. Evolution of Mollicutes: Down a
bumpy road with twists and turns. *Research in microbiology* **158**, 754-766 (2007).
21 Peterson, S. N. & Fraser, C. M. The complexity of simplicity. *Genome biology* **2**, 1-7
(2001).
22 He, L.-S. *et al.* The enigmatic genome of an obligate ancient *Spiroplasma* symbiont in a
hadal holothurian. *Applied and Environmental Microbiology* **84**, e01965-01917 (2018).
23 Viver, T. *et al.* The low diverse gastric microbiome of the jellyfish *Cotylorhiza*
tuberculata is dominated by four novel taxa. *Environmental Microbiology* **19**, 3039-3058
(2017).
24 Cortés-Lara, S., Urdiain, M., Mora-Ruiz, M., Prieto, L. & Rosselló-Móra, R. Prokaryotic
microbiota in the digestive cavity of the jellyfish *Cotylorhiza tuberculata*. *Systematic and*
Applied Microbiology **38**, 494-500 (2015).
25 Duperron, S. *et al.* A tale of two chitons: Is habitat specialisation linked to distinct
associated bacterial communities? *FEMS Microbiology Ecology* **83**, 552-567 (2013).
26 Tait, E., Carman, M. & Sievert, S. M. Phylogenetic diversity of bacteria associated with
ascidians in eel pond (woods hole, massachusetts, USA). *Journal of Experimental Marine*
Biology and Ecology **342**, 138-146 (2007).
27 Høj, L. *et al.* Crown-of-thorns sea star, *Acanthaster cf. solaris*, has tissue-characteristic
microbiomes with potential roles in health and reproduction. *Applied and Environmental*
Microbiology **84**, e00181-00118 (2018).
28 Gray, M. A., Stone, R. P., McLaughlin, M. R. & Kellogg, C. A. Microbial consortia of
gorgonian corals from the Aleutian islands. *FEMS Microbiology Ecology* **76**, 109-120
(2011).
29 Penn, K., Wu, D., Eisen, J. A. & Ward, N. Characterization of bacterial communities
associated with deep-sea corals on Gulf of Alaska seamounts. *Applied and Environmental*
Microbiology **72**, 1680-1683 (2006).
30 Weiler, B. A., Verhoeven, J. T. P. & Dufour, S. C. Bacterial communities in tissues and
surficial mucus of the cold-water coral *Paragorgia arborea*. *Frontiers in Marine Science*
5 (2018).
31 Holm, J. B. & Heidelberg, K. B. Microbiomes of *Muricea californica* and *M. ruticosa*:
Comparative analyses of two co-occurring eastern pacific octocorals. *Frontiers in*
Microbiology **7** (2016).
32 Kellogg, C. A., Ross, S. W. & Brooke, S. D. Bacterial community diversity of the deep-
sea octocoral *Paramuricea placomus*. *PeerJ* **4**, e2529 (2016).

33   Porporato, E. M. D., Lo Giudice, A., Michaud, L., De Domenico, E. & Spanò, N.
Diversity and antibacterial activity of the bacterial communities associated with two
mediterranean sea pens, *Pennatula phosphorea* and *Pteroeides spinosum* (anthozoa:
Octocorallia). *Microbial Ecology* **66**, 701-714 (2013).
34   Shirur, K. P., Jackson, C. R. & Goulet, T. L. Lesion recovery and the bacterial
microbiome in two caribbean gorgonian corals. *Marine Biology* **163**, 238 (2016).
35   van de Water, J. A. J. M. *et al.* Seasonal stability in the microbiomes of temperate
gorgonians and the red coral *Corallium rubrum* across the mediterranean sea. *Microbial*
Ecology **75**, 274-288 (2018).
36   van de Water, J. A. J. M. *et al.* Comparative assessment of mediterranean gorgonian-
associated microbial communities reveals conserved core and locally variant bacteria.
Microbial Ecology **73**, 466-478 (2017).
37   McCauley, M., Jackson, C. R. & Goulet, T. L. Microbiomes of caribbean octocorals vary
over time but are resistant to environmental change. *Frontiers in microbiology* **11**, 1272
(2020).
38   van de Water, J. A. J. M., Coppari, M., Enrichetti, F., Ferrier-Pagès, C. & Bo, M. Local
conditions influence the prokaryotic communities associated with the mesophotic black
coral *Antipathella subpinnata*. *Frontiers in Microbiology* **11** (2020).
39   Neulinger, S. C. *et al.* Tissue-associated "*Candidatus Mycoplasma corallicola*" and
filamentous bacteria on the cold-water coral *Lophelia pertusa* (scleractinia). *Applied and*
Environmental Microbiology **75**, 1437-1444 (2009).
40   Kellogg, C. A., Lisle, J. T. & Galkiewicz, J. P. Culture-independent characterization of
bacterial communities associated with the cold-water coral *Lophelia pertusa* in the
northeastern Gulf of Mexico. *Applied and Environmental Microbiology* **75**, 2294-2303
(2009).
41   Woo, S. *et al.* Geographical variations in bacterial communities associated with soft coral
Scleronephthya gracillimum. *PLOS ONE* **12**, e0183663 (2017).
42   Quintanilla, E., Rodrigues, C. F., Henriques, I. & Hilário, A. Microbial associations of
abyssal gorgonians and anemones (> 4,000 m depth) at the clarion-clipperton fracture
zone. *Frontiers in Microbiology* **13** (2022).
43   Quattrini, A. M., Baums, I. B., Shank, T. M., Morrison, C. L. & Cordes, E. E. Testing the
depth-differentiation hypothesis in a deepwater octocoral. *Proceedings of the Royal*
Society B: Biological Sciences **282**, 20150008 (2015).
44   Quattrini, A. M. *et al.* Niche divergence by deep-sea octocorals in the genus *Callogorgia*
across the continental slope of the Gulf of Mexico. *Molecular Ecology* **22**, 4123-4140
(2013).
45   Etnoyer, P. & Warrenchuk, J. A catshark nursery in a deep gorgonian field in the
Mississippi Canyon, Gulf of Mexico. *Bulletin of Marine Science* **81**, 553-559 (2007).
46   Yarza, P. *et al.* Uniting the classification of cultured and uncultured bacteria and archaea
using 16S rRNA gene sequences. *Nature Reviews Microbiology* **12**, 635-645 (2014).
47   Bankevich, A. *et al.* Spades: A new genome assembly algorithm and its applications to
single-cell sequencing. *Journal of computational biology* **19**, 455-477 (2012).
48   Lee, I., Kim, Y. O., Park, S.-C. & Chun, J. OrthoANI: An improved algorithm and
software for calculating average nucleotide identity. *International journal of systematic*
and evolutionary microbiology **66**, 1100-1103 (2016).

49   Romano, N., Tolone, G., Ajello, F. & La Licata, R. Adenosine 5'-triphosphate synthesis
639   induced by urea hydrolysis in *Ureaplasma urealyticum*. *Journal of Bacteriology* **144**,
640   830-832 (1980).
641 50   Smith, D. G., Russell, W. C., Ingledew, W. J. & Thirkell, D. Hydrolysis of urea by
642   *Ureaplasma urealyticum* generates a transmembrane potential with resultant ATP
643   synthesis. *Journal of Bacteriology* **175**, 3253-3258 (1993).
644 51   Ipoutcha, T. *et al.* Multiple origins and specific evolution of CRISPR/cas9 systems in
645   minimal bacteria (Mollicutes). *Frontiers in microbiology* **10**, 2701 (2019).
646 52   Lian, C.-A. *et al.* Genomic characterization of a novel gut symbiont from the hadal
647   snailfish. *Frontiers in microbiology* **10**, 2978 (2020).
648 53   Remmert, M., Biegert, A., Hauser, A. & Söding, J. Hhblits: Lightning-fast iterative
649   protein sequence searching by hmm-hmm alignment. *Nature methods* **9**, 173-175 (2012).
650 54   Schmidt, A. J., Ryjenkov, D. A. & Gomelsky, M. The ubiquitous protein domain EAL is
651   a cyclic diguanylate-specific phosphodiesterase: Enzymatically active and inactive EAL
652   domains. *Journal of bacteriology* **187**, 4774-4781 (2005).
653 55   Minasov, G. *et al.* Crystal structures of YkuI and its complex with second messenger
654   cyclic di-gmp suggest catalytic mechanism of phosphodiester bond cleavage by EAL
655   domains. *J. Biol. Chem.* **284**, 13174-13184 (2009).
656 56   Tchigvintsev, A. *et al.* Structural insight into the mechanism of c-di-GMP hydrolysis by
657   eal domain phosphodiesterases. *Journal of molecular biology* **402**, 524-538 (2010).
658 57   Wicke, D., Meißner, J., Warneke, R., Elfmann, C. & Stülke, J. Understudied proteins and
659   understudied functions in the model bacterium *Bacillus subtilis*—a major challenge in
660   current research. *Molecular Microbiology* (2023).
661 58   Faburay, B. & McVey, D. S. Mollicutes. *Veterinary Microbiology*, 364-376 (2022).
662 59   Goff, L. J. Symbiosis and parasitism: Another viewpoint. *BioScience* **32**, 255-256 (1982).
663 60   Jenal, U., Reinders, A. & Lori, C. Cyclic di-GMP: Second messenger extraordinaire.
664   *Nature Reviews Microbiology* **15**, 271-284 (2017).
665 61   Berzins, I. K., Yanong, R. P., LaDouceur, E. E. & Peters, E. C. Cnidaria. *Invertebrate*
666   *Histology*, 55-86 (2021).
667 62   Van Praet, M. Absorption of substances dissolved in the environment, particles and
668   products of extracellular digestion in *Actinia equina* (Cnidaria, Actiniaria). *Reproduction,*
669   *Nutrition, Developpement* **20**, 1393-1399 (1980).
670 63   Lebouvier, M., Miramón-Puértolas, P. & Steinmetz, P. R. Evolutionarily conserved
671   aspects of animal nutrient uptake and transport in sea anemone vitellogenesis. *Current*
672   *Biology* **32**, 4620-4630. e4625 (2022).
673 64   Menzel, L., Tondo, C., Stein, B. & Bigger, C. Histology and ultrastructure of the
674   coenenchyme of the octocoral *Swiftia exserta*, a model organism for innate
675   immunity/graft rejection. *Zoology* **118**, 115-124 (2015).
676 65   Hufnagel, L. A. & Myhal, M. L. Observations on a spirochaete symbiotic in *Hydra*.
677   *Transactions of the American Microscopical Society*, 406-411 (1977).
678 66   Rathje, K. *et al.* Dynamic interactions within the host-associated microbiota cause tumor
679   formation in the basal metazoan *Hydra*. *PLoS Pathogens* **16**, e1008375 (2020).
680 67   Miller, A. W., Blackwelder, P., Al-Sayegh, H. & Richardson, L. L. Fine-structural
681   analysis of black band disease-infected coral reveals boring cyanobacteria and novel
682   bacteria. *Diseases of aquatic organisms* **93**, 179-190 (2011).

68   Kriest, I. Different parameterizations of marine snow in a 1d-model and their influence
on representation of marine snow, nitrogen budget and sedimentation. *Deep Sea*
Research Part I: Oceanographic Research Papers **49**, 2133-2162 (2002).
69   Horn, H. *et al.* An enrichment of CRISPR and other defense-related features in marine
sponge-associated microbial metagenomes. *Frontiers in microbiology* **7**, 1751 (2016).
70   Fan, L. *et al.* Functional equivalence and evolutionary convergence in complex
communities of microbial sponge symbionts. *Proceedings of the National Academy of*
Sciences **109**, E1878-E1887 (2012).
71   Hutchison III, C. A. *et al.* Design and synthesis of a minimal bacterial genome. *Science*
351, aad6253 (2016).
72   Breuer, M. *et al.* Essential metabolism for a minimal cell. *Elife* **8**, e36842 (2019).
73   Galperin, M. Y. Genomes of model organisms: Know thy tools. *Environmental*
microbiology **10**, 1383 (2008).
74   Rodriguez-Lanetty, M., Granados-Cifuentes, C., Barberan, A., Bellantuono, A. J. &
Bastidas, C. Ecological inferences from a deep screening of the complex bacterial
consortia associated with the coral, *Porites astreoides*. *Molecular Ecology* **22**, 4349-4362
(2013).
75   Naqib, A. *et al.* In *Gene Expression Analysis. Methods in Molecular Biology* Vol. 1783
Making and sequencing heavily multiplexed, high-throughput 16S ribosomal RNA gene
amplicon libraries using a flexible, two-stage PCR protocol. 149-169 (Springer, 2018).
76   Bolyen, E. *et al.* Qiime 2: Reproducible, interactive, scalable, and extensible microbiome
data science. *PeerJ Preprints* **6**, e27295v27292 (2018).
77   Gruber-Vodicka, H. R., Seah, B. K. B. & Pruesse, E. Phyloflash – rapid SSU rRNA
profiling and targeted assembly from metagenomes. *bioRxiv*, 521922 (2019).
78   Wick, R. R., Schultz, M. B., Zobel, J. & Holt, K. E. Bandage: Interactive visualization of
de novo genome assemblies. *Bioinformatics* **31**, 3350-3352 (2015).
79   Gao, F. & Zhang, C.-T. Ori-finder: A web-based system for finding oric s in unannotated
bacterial genomes. *BMC bioinformatics* **9**, 1-6 (2008).
80   Aziz, R. K. *et al.* The rast server: Rapid annotations using subsystems technology. *BMC*
genomics **9**, 75 (2008).
81   Overbeek, R. *et al.* The subsystems approach to genome annotation and its use in the
project to annotate 1000 genomes. *Nucleic acids research* **33**, 5691-5702 (2005).
82   Li, D. H., Liu, C. M., Luo, R. B., Sadakane, K. & Lam, T. W. Megahit: An ultra-fast
single-node solution for large and complex metagenomics assembly via succinct de
bruijn graph. *Bioinformatics* **31**, 1674-1676 (2015).
83   Li, H. *et al.* The sequence alignment/map format and samtools. *Bioinformatics* **25**, 2078-
2079 (2009).
84   Kang, D. D., Froula, J., Egan, R. & Wang, Z. Metabat, an efficient tool for accurately
reconstructing single genomes from complex microbial communities. *PeerJ* **3**, e1165
(2015).
85   Symeonidi, E., Regalado, J., Schwab, R. & Weigel, D. Crispr-finder: A high throughput
and cost-effective method to identify successfully edited arabidopsis thaliana individuals.
Quantitative Plant Biology **2** (2021).
86   Grissa, I., Vergnaud, G. & Pourcel, C. Crisprfinder: A web tool to identify clustered
regularly interspaced short palindromic repeats. *Nucleic acids research* **35**, W52-W57
(2007).

87   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nature Biotechnology* **34**, 525 (2016).

88   Edgar, R. C. Muscle: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic acids research* **32**, 1792-1797 (2004).

89   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies.
Molecular Biology and Evolution **32**, 268-274 (2014).

90   Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A. & Minh, B. Q. W-IQ-TREE: A fast
online phylogenetic tool for maximum likelihood analysis. *Nucleic acids research* **44**,
W232-W235 (2016).

91   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
Modelfinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods*
14, 587 (2017).

92   Soubrier, J. *et al.* The influence of rate heterogeneity among sites on the time dependence
of molecular rates. *Molecular Biology and Evolution* **29**, 3345-3358 (2012).

93   Yang, Z. A space-time process model for the evolution of DNA sequences. *Genetics* **139**,
993-1005 (1995).

94   Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:
Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**,
518-522 (2017).

95   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
Assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome research* **25**, 1043-1055 (2015).

96   Gupta, R. S., Sawnani, S., Adeolu, M., Alnajar, S. & Oren, A. Phylogenetic framework
for the phylum Tenericutes based on genome sequence data: Proposal for the creation of
a new order *Mycoplasmoidales* ord. nov., containing two new families
Mycoplasmoidaceae fam. nov. and *Metamycoplasmataceae* fam. nov. harbouring
Eperythrozoon, *Ureaplasma* and five novel genera. *Antonie van Leeuwenhoek* **111**, 1583-
1630 (2018).

97   Gupta, R. S. & Oren, A. Necessity and rationale for the proposed name changes in the
classification of Mollicutes species. Reply to:'Recommended rejection of the names
Malacoplasma gen. nov., *Mesomycoplasma* gen. nov., *Metamycoplasma* gen. nov.,
Metamycoplasmataceae fam. nov., *Mycoplasmoidaceae* fam. nov., Mycoplasmoidales
ord. nov., Mycoplasmoides gen. nov., *Mycoplasmopsis* gen. nov.[gupta, sawnani, adeolu,
alnajar and oren 2018] and all proposed species comb. Nov. Placed therein', by m. Balish
et al.(int j syst evol microbiol, 2019; 69: 3650–3653). *International Journal of Systematic*
and Evolutionary Microbiology **70**, 1431-1438 (2020).

98   Gupta, R. S., Son, J. & Oren, A. A phylogenomic and molecular markers based
taxonomic framework for members of the order Entomoplasmatales: Proposal for an
emended order Mycoplasmatales containing the family Spiroplasmataceae and emended
family Mycoplasmataceae comprised of six genera. *Antonie Van Leeuwenhoek* **112**, 561-
588 (2019).

99   Shaffer, M. *et al.* DRAM for distilling microbial metabolism to automate the curation of
microbiome function. *Nucleic acids research* **48**, 8883-8900 (2020).

100   Li, L., Stoeckert, C. J. & Roos, D. S. OrthoMCL: Identification of ortholog groups for
774   eukaryotic genomes. *Genome research* **13**, 2178-2189 (2003).

101 Arkin, A. P. *et al.* Kbase: The united states department of energy systems biology
knowledgebase. *Nature biotechnology* **36**, 566-569 (2018).
102 Amann, R. I. *et al.* Combination of 16S rRNA-targeted oligonucleotide probes with flow
cytometry for analyzing mixed microbial populations. *Applied and environmental*
microbiology **56**, 1919-1925 (1990).
103 Daims, H., Brühl, A., Amann, R., Schleifer, K.-H. & Wagner, M. The domain-specific
probe EUB338 is insufficient for the detection of all bacteria: Development and
evaluation of a more comprehensive probe set. *Systematic and applied microbiology* **22**,
434-444 (1999).
784

Acknowledgements:

We would like to thank Martina Meyer, Miriam Sadowski, Silke Wetzel, and Meghann

Devlin-Durante for their help with microscopy and DNA extractions. We would also like to

thank the crews of all research vessels and ROV teams involved in collecting samples. This

research was made possible by a grant from The Gulf of Mexico Research Initiative which was

awarded to the Ecosystem Impacts of Oil and Gas Inputs to the Gulf (ECOGIG) consortium.

Additional funding was provided from the Max Planck Society. The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Information:

Authors and Affiliations

Department of Biology, The Pennsylvania State University, State College, PA, USA

Samuel A. Vohsen, Charles R. Fisher, Iliana B. Baums

Department of Symbiosis, Max Planck Institute for Marine Microbiology, Bremen, Germany

Harald R. Gruber-Vodicka, Nicole Dubilier

Contributions

SAV conducted sampling, processing, and analyses. IBB and CRF acquired funding, led research

cruises, and supervised work. ND contributed additional funding and supervised work. HGV

advised analyses. SAV wrote the manuscript. All authors reviewed and edited the manuscript.

Corresponding author

Correspondence to Iliana B. Baums

Ethics Declaration:
806   Competing Interests

807   The authors declare no competing interests

808   **Tables and Figures:**

810   **Table 1**. Summary of the genome statistics for the novel mollicutes in *Callogorgia delta* and
811   their closest known relatives. *LSU rRNA fragments. Pegs (w/KO) = CDS = coding sequences

Organism name	Genome Size (bp)	Contigs	GC (%)	DRAM	Classic RAST			checkm			
				pegs (w/ KO)	CDS	tRNAs	rRNAs	Completeness (%)	Contamination (%)	Predicted Genes	Coding Density (%)
Ca. Oceanoplasma callogorgiae	446,099	3	31.0	359 (241)	405	34	2 + 7*	70.4	2.6	392	88.2
Ca. Thalassoplasma callogorgiae	445,436	21	24.9	385 (237)	416	30	2	72.4	2.6	396	87.6
Ca. Spiroplasma holothuricola[1]	424,539[1]	2[1]	29.5[1]	336 (224)	399 (347)	32	2	67.4	2.3	367	89.9
Mycoplasma- like from jellyfish[2]	389,210 (409,158)[2]	6 (11)[2]	33.1 (32.9)[2]	342 (225)	358	32	2	73.4 (74.1)	2.3 (2.3)	356	92.5

References: [1]He et al. 2018, [2]Viver et al 2017

Fig. 1: Sampling locations and prevalence of novel mollicutes in *Callogorgia delta* and *C.*

americana

a) Map of locations where *Callogorgia delta* (black circles) and *C. americana* (gray

diamonds) were sampled. The depth range of samples collected from each site is denoted below

the site name followed by the number of colonies, sediment, and water samples in parentheses. If

no water or sediment samples were collected, only the number of colonies is shown. A star

signifies whether a water sample was collected using a McLane pump (~400L) in addition to

niskin bottles (~2.5L). Bathymetric map was obtained from BOEM (credit: William Shedd,

Kody Kramer). b) Image of *Callogorgia delta.* c) The relative abundances of novel mollicute

ASVs. Each column represents the microbial composition based on 16S rRNA libraries obtained

from a single colony. Colonies are organized by species, site, and sampling year. *C. delta* sites

are ordered left to right by increasing depth.

Fig. 2: Phylogenetic analyses of the novel mollicutes in *C. delta*

Maximum likelihood phylogenetic trees of class Mollicutes using the a) 16S rRNA gene

and b) amino acid sequences of 43 genes. UFBoot support values are reported at each node.

Numbers in parentheses designate the number of sequences within collapsed nodes. *Ca.*

Oceanoplasma callogorgiae and *Ca. Thalassoplasma callogorgiae* are presented in bold text and

underlined. Genome sizes (in Mbp) are depicted in c).

Fig. 3: Simplified metabolic model and transcriptomic profile of novel mollicutes

838   a) Simplified metabolic model of *Ca. Oceanoplasma callogorgiae* and *Ca.*

Thalassoplasma callogorgiae. Gene names are denoted in italics and unidentified proteins in red

text. ATP is generated by the arginine dihydrolase pathway. Additional ATP is likely generated

by ATP-synthase utilizing the proton gradient resulting from the consumption of H^+ and

production of NH_3. Nicotinamide (NAM), nicotinamide riboside (NR), nicotinamide

mononucleotide (NMN), 5-phosphoribosyl diphosphate (PRPP) b) Expression level of *Ca. O.*

callogorgiae genes from two libraries sorted by functional annotations. Only genes with a

maximum of 1.5 transcripts per thousand among both libraries are displayed.

Fig. 4: Comparative Genomics of related mollicutes

a) Non-metric multidimensional scaling (NMDS) plot of cluster of orthologous genes

(COG) using Jaccard distances. Only COGs present in more than two genomes were included. b)

the number of predicted protein-encoding genes for each genome. Oceanoplasmataceae had the

fewest predicted protein-encoding genes. c) the presence of COGs involved in carbohydrate

metabolism for each genome. COGs are labeled using their KEGG orthology identifier

annotations. d) metabolic map of glycolysis and other reactions associated with the COGs in c).

The map was drawn using reactions listed on the KEGG database for each KO identifier.

Phosphoribosyl diphosphate

Fig. 5: Bacterial signals in the mesoglea of *Callogorgia delta*

Tissue sections were hybridized using CARD-FISH with the general eubacterial probe

mix (EUB338 I-III) and DNA was stained with DAPI. Fluorescence in the emission range of

DAPI is shown in blue while green shows emission in the range of the fluorophore ALEXA 488

used by both EUB338 I-III and control probes. Paired images compare adjacent sections where

sections on the left were hybridized with EUB338 I-III and show bacterial signals (a,c,e) while

sections on the right were hybridized with a negative control probe and only show

autofluorescence (b,d,f). Signals were observed in transverse sections through the central axis

(a,b) and transverse sections through a single polyp (c,d,e,f,g). Arrows in (a) and (c) show

examples of EUB338 I-III fluorescence. A = central axis, G = gastrovascular cavity, M = gastric

mesentery, L = sclerite lacuna. Scale bars = 100 µm (a,b,g), 200 µm (c,d,e,f)

Fig. 6: Higher magnification of bacterial signals in the mesoglea and tentacles of *Callogorgia*

delta

Tissue sections were hybridized using CARD-FISH with the general eubacterial probe

mix (EUB338 I-III) and DNA was stained with DAPI. Fluorescence in the emission range of

DAPI is shown in blue while green shows emission in the range of the fluorophore ALEXA 488

used by both EUB338 I-III and control probes. Paired images compare adjacent sections where

sections on the left were hybridized with EUB338 I-III and show bacterial signals (a,c,e) while

sections on the right were hybridized with a negative control probe and only show

autofluorescence (b,d,f). Transverse sections through the central axis (a,b) exhibit EUB338 I-III

fluorescence (a, arrows) in the surrounding mesoglea and an overlap of DAPI (c, arrow). (a,

arrow). Transverse sections through a polyp (c,d) show EUB338 I-III fluorescence (c, arrows) in

the base of a tentacle. Transverse sections through the body wall of a polyp show granular

(<1µm) EUB338 I-III fluorescence in the mesoglea (e,f,g). A = central axis, T = tentacle. Scale

bars = 10 µm (a,c,d,e,f), 20 µm (g)