1 **Fast prediction in marmoset reach-to-grasp movements for dynamic prey**

2 Luke Shaw[1], Kuan Hong Wang[1,3], Jude Mitchell[2,3]

3 [1] Department of Neuroscience, University of Rochester Medical Center, Rochester, NY
4 [2] Department of Brain and Cognitive Sciences, University of Rochester, Rochester, NY
5 [3] Lead Contact
6

Summary

Primates have evolved sophisticated visually guided reaching behaviors for interacting with
dynamic objects, such as insects during foraging(P. S. Archambault, Ferrari-Toniolo, & Battaglia-
Mayer, 2011; Bicca-Marques, 1999; Ngo et al., 2022; Smith & Smith, 2013; Sustaita et al., 2013).
Reaching control in dynamic natural conditions requires active prediction of the target's future
position to compensate for visuo-motor processing delays and enhance online movement
adjustments(Catania, 2009; Desmurget & Grafton, 2000; Fujioka, Aihara, Sumiya, Aihara, &
Hiryu, 2016; Merchant & Georgopoulos, 2006; Mischiati et al., 2015; R. Shadmehr, Smith, &
Krakauer, 2010; Wolpert & Kawato, 1998). Past reaching research in non-human primates mainly
focused on seated subjects engaged in repeated ballistic arm movements to either stationary
targets, or targets that instantaneously change position during the movement(Philippe S.
Archambault, Caminiti, & Battaglia-Mayer, 2009; Battaglia-Mayer et al., 2013; Dickey, Amit, &
Hatsopoulos, 2013; Georgopoulos, Kalaska, Caminiti, & Massey, 1983; Georgopoulos, Kalaska,
& Massey, 1981). However, those approaches impose task constraints that limit the natural
dynamics of reaching. A recent field study in marmoset monkeys highlights predictive aspects of
visually-guided reaching during insect prey capture among wild marmoset monkeys(Ngo et al.,
2022). To examine the complementary dynamics of similar natural behavior within a laboratory
context we developed an ecologically motivated unrestrained reach-to-grasp task involving live
crickets. We used multiple high-speed video cameras to capture the movements of marmosets
and crickets stereoscopically and applied machine vision algorithms for marker-free object and
hand tracking. Contrary to estimates under traditional constrained reaching paradigms, we find
that reaching for dynamic targets can operate at incredibly short visuo-motor delays around 80
milliseconds, rivaling the speeds that are typical of the oculomotor systems during closed-loop
visual pursuit(Cloherty, Yates, Graf, DeAngelis, & Mitchell, 2020). Multivariate linear regression
modeling of the kinematic relationships between the hand and cricket velocity revealed that
predictions of the expected future location can compensate for visuo-motor delays during fast
reaching. These results suggest a critical role of visual prediction facilitating online movement
adjustments for dynamic prey.

Keywords

Measurement of visuomotor delay in marmosets, evidence for fast predictive reaching,
naturalistic paradigm, crickets, markerless video tracking

Results

Tracking of marmoset reaches

Primates exhibit a sophisticated repertoire of visually-guided reaching behaviors that enable them
to interact with objects and navigate arboreal environments. While the motor aspects of these
reaching movements are largely conserved across mammalian species, primates are unique in

their integration of visual information to guide reaching movements(De Bruin, Sacrey, Brown,
Doan, & Whishaw, 2008; J. H. Kaas, Gharbawie, & Stepniewska, 2011; Jon H. Kaas &
Stepniewska, 2016; Sacrey & Whishaw, 2012; Wang et al., 2017). A recent study used video in
the field to examine reach-to-grasp movements for flying insects among wild marmoset
monkeys(Ngo et al., 2022). They found evidence to support an online active vision strategy during
prey capture in which the head angle of marmosets actively tracked insects during reaching.
However, the video resolution in the wild does not afford enough precision to measure visuo-
motor delay nor test if the motor control strategy exploits predictions of future target location to
compensate for processing delays. Here we establish an ecologically motivated reaching task
within a laboratory setting using high-speed video to test the kinematics of reach-to-grasp
movements for dynamic prey.

Figure 1. High-speed, multi-camera video reconstructs reaching for dynamic prey. A: A wheeled video platform allowed tests of individual marmosets from inside their housing in a colony. **B:** Three GoPro cameras were calibrated around a semi-circular reaching arena where a live cricket was placed. Marmosets reached through a rectangular aperture to acquire crickets. **C:** Feature points were manually labeled in a subset of video frames to train DeepLabCut. Hand and cricket markers (*) were used for reaching analyses and knuckle markers (^) for grasping analyses. **D:** Arena markers were used to validate spatial coordinates and correct for any post-calibration camera movement. **E-G:** DeepLabCut labeling of synchronized frames of video from three camera angles. DeepLabCut tracked labeled points with a RMSE of ~3 and ~7 pixels for the cricket and hand, respectively, compared to ~37 pixels for the size of the marmoset wrist.

We used high-speed, multi-camera video and machine learning methods to track marmoset
reaches for live, moving crickets. Reaching experiments were conducted by docking a mobile
video recording platform to marmoset family cages within the marmoset colony and recording
reaches using 3 GoPro cameras **(Figure 1A-B)**. Hands, fingers, crickets, and the experimental
apparatus were tracked from three cameras using DeepLabCut(Mathis et al., 2018) and 3D
position triangulated using Anipose(Karashchuk et al., 2021) **(Figure 1C-G)**. DeepLabCut
performance using the labels illustrated in **Figure 1C-D** is quantified in **Supplemental Figure 1**.
Although we were able to reconstruct 3D trajectories of reaching, most subsequent analyses
focused on 2D tracking from the top view camera angle depicted in **Figure 1F**, because crickets
mainly limited their motion to the horizontal plane during the reaching tasks.

Marmoset reach-to-grasp dynamics

Earlier work in laboratory settings have shown that marmosets are equally likely to prefer using
their left or right hands in reaching(Hook & Rogers, 2008), and typically employ a power grasp
strategy to reach items in which all fingers open in unison and close around the target to press it
within the palm of the hand, which has been shown in marmoset (Fox, Mundinano, & Bourne,
2019) and macaque(Castiello & Dadda, 2019; Sartori, Camperio Ciani, Bulgheroni, & Castiello,
2013). A key difference from macaques is that marmosets do not use a precision grip. Old World
primates have opposable thumbs that allow them to grasp small objects between the thumb and
fingers (Marzke, 1997), while most New World primates, including the marmoset, use a power
grasp(Bishop, 1962). We found that marmosets in our experiments predominantly employed a
left or right single-handed (LH or RH) reaching strategy akin to a pause and lunge approach
utilized by marmosets in the wild(Ngo et al., 2022; Schiel & Souto, 2017). This strategy may have
been favored due to the reaching aperture utilized in our setup, which prevented marmosets from
using their mouths or a two-handed pounce (P) strategy. We recorded a total of 271 reaches for
live crickets (137 LH, 134 RH, 10 P) from 10 marmosets. Gross quantification of these reaches
through a left-handedness index (LH reaches/total reaches) indicates an evenly split mixed hand
preference over the population of reaches. Assessing this quantification at the marmoset
participant level shows two marmosets with a strong right-handed preference (LHI=.31 and .04)
and three marmosets with a strong left-handed preference (.83, .87, and .92), while the other 5
marmosets had little preference (.60, .68, .64, .66, .44, .40). Of the total recorded set of 271
reaches for crickets, 78 reaches contained crickets that moved during or prior to the reach, which
were selected for subsequent analyses (see **Star Methods** for inclusion criteria).

Marmosets performed reach-to-grasp movements for moving crickets using a power grasp to
enclose crickets within their hand at the end of the reach (**Figure 2**). In an individual reaching
trial, the opening of grasp aperture begins as hand velocity drops with increasing proximity to
target (**Figure 2A-C**), which is consistent with findings in previous NHP studies from both
marmosets and macaques (Castiello & Dadda, 2019; Fox et al., 2019; Sartori et al., 2013). Finger
separation distance was calculated as the average of the distances between the index finger and
middle finger, and the middle finger and ring finger, using knuckle markers (**Figure 2B, dark blue
points**). For this reach, there is a clear temporal order with the peak in hand speed followed by
an increase in finger spread as the hand speed decreases (**Figure 2C**). This pattern was typical
of the population of reaches. A frontal view of a typical reach (**Supplemental Figure 2**) shows
the fingers wrapping around the cricket.

We used dimensionality reduction methods to further characterize the grasping strategies
employed in the population of reaches. For each reach, a time series of finger spread distance

Figure 2. Hand aperture opens earlier with increasing hand speed in a power grasp. A: Cropped video frames show a single reaching trial at two time points indicated by light and dark gray points. **B:** The hand position for the corresponding time points in A is shown from the top camera view. The underlying light blue trace shows the trajectory of the reach from the central hand marker and the red traces indicates cricket motion. **C:** Within the example reach the speed of the hand (shown in black, left vertical scale) peaked and began to fall before the finger spread increased, depicted in gray (right vertical scale). **D:** The finger spread over time was submitted to a cluster analysis (see methods) which revealed a continuum along the principal component axes that we split into three clusters (green, purple, and orange) for visualization. The two outliers present in Cluster 3 (yellow, at the bottom) were due to finger contact against the retaining wall at the end of the reach. **E:** Finger separation peaked at longer delays in the reach for the three identified clusters when plotted over normalized time from reach start to cricket occlusion. **F:** The corresponding hand velocity traces for the same clusters reveal slower peak velocities correlating with later grasp opening. **G-I:** Three frames from an example reach to grasp for each of the clusters reveals a power grasp enclosing the cricket; on the right, plots of the three knuckle markers used to measure finger spread at corresponding times.

was calculated based on the knuckle markers. The distance values were interpolated over a fixed
number of timesteps over the total reach duration, with the reach start defined from the moment
the hand first entered the arena to the end being when the fingers first began to close around the

cricket. Principal component analysis of these reach trials captured 69.1% of the variance within
the first two components. We found a continuum of points, where each point corresponds to one
time-normalized reach, that extended along the first principal component as opposed to discrete
clusters (**Figure 2D**). To visualize the variations in finger separation across the population of
reaches we performed a k-means clustering and split the space into the best fitting three clusters.

The power grasp strategy shows a temporal pattern of earlier grasp opening for faster
movements. Cluster 1 shows an early finger spread after reach initiation, which continues
throughout the reach (**Figure 2E1**). By comparison in Cluster 2, the finger spread initiates later
during hand transport (**Figure 2E2**), and in Cluster 3 finger spread initiates only at the very end
of the reach **(Figure 2E3)**. Plotting hand velocity time series for the same clusters we find higher
and earlier peak velocities for Cluster 1 compared to Cluster 2, and likewise for Cluster 2 to 3
(**Figure 2F1-3**). Thus, the timing of grasp opening varies with reach speed. Video frames taken
from an example reach in each of the three clusters illustrate the power grasp, with variation in
the timing of grasp opening across the examples **(Figure 2G-I)**. In the third case with the slowest
speed the fingers have still not fully separated by movement end (**Figure 2I**). In contrast, rodents
often use an arpeggio grasp to first contact the object with fingers extended flatly, and then based
on somatosensory feedback, close around it(Whishaw & Gorny, 1994). Here even the slowest
reaches appear to involve opening of the grasp near contact (**Fig. 2E3**), and thus are more
consistent with a vision-guided power grasp strategy typical of macaque reaching.

Hand velocity converges to match lateral cricket velocity during reaching

Movement dynamics during prey capture have been found to follow control strategies for target
interception such as pure pursuit and proportional navigation across a range of species including
dragonflies(Mischiati et al., 2015), hawks(Brighton & Taylor, 2019), and humans(Lenoir, Musch,
Janssens, Thiery, & Uyttenhove, 1999). In pure pursuit the rotational acceleration, or steering, of
the body (or hand) is dictated by the instantaneous offset from the target location, called the range
vector. Pure pursuit seeks to close the distance of the range vector but ignores motion of the
target and is not anticipatory of its future location. By contrast, proportional navigation aims to
maintain a constant bearing angle of the range vector as the pursuer closes the distance. One
way to visualize the distinction between these strategies is to decompose velocity into a
component that lies along the range vector at each movement, the direct axis, and another
component that lies on its orthogonal projection, the lateral axis. Both strategies predict changes
in velocity related to closing distance on the direct axis, but in addition proportional navigation
predicts a matching of the velocity of the body (or hand) with the lateral velocity of the target to
maintain a constant bearing angle. Another feature characteristic of proportional navigation is that
the derivative of the range vector over time will be negatively correlated at -1 with the range vector
itself(Mischiati et al., 2015).

We examined the direct and lateral components of hand velocity as marmosets performed
reaches to moving crickets in the arena (**Figure 3)**. Although we are able to reconstruct the full
3D kinematics of reaching movements **(Figure 3A)**, we focus our subsequent analyses on the
horizontal plane seen from the top camera view which captured most of the cricket motion. To
examine the influence of cricket velocity on hand velocity, we decomposed velocity of the hand
(H) and cricket (C) into direct and lateral components relative to the range vector (RV) connecting
the instantaneous position of the two **(Figure 3B)**. In the set of reaches, most of the hand motion
falls along the direct axis, *Dh*, which is necessary to traverse the arena to close on the cricket and
exhibits a peaked shape typical of ballistic reaching movements (**Figure 3C**, dashed lines**)**.

Conversely, most of cricket motion falls along the lateral axis, *Lc*, due to the tendency of crickets
to crawl along the perimeter of the reaching arena. The lateral component of absolute hand
velocity, *Lh*, was closely matched in magnitude to cricket lateral velocity, *Lc* (**Figure 3C**, solid
lines). We thus sought to determine to what extent the lateral hand velocity might match cricket
velocity, both in sign and magnitude over the reach, as predicted during proportional navigation.

A typical reaching trial demonstrates that hand velocity tracks the lateral motion of the cricket
target in a manner more consistent with a proportional navigation strategy for interception **(Figure
3D-F)**. In this trial, the cricket moves in the horizontal plane along the boundary of the arena and
the hand adopts a curved trajectory in pursuit (**Figure 3D**). We applied computational methods
from a previous study of hawk prey pursuit(Brighton & Taylor, 2019) to simulate the hand's
trajectory using either a pure pursuit (PP) or proportional navigation (PN) steering model. For this
trial the PN model provides a closer match to the actual hand trajectory (R^2=.99) compared to the
PP model (R^2=.94) (**Figure 3E**). In line with the better fit of the PN model, the correlation of the
range vectors to their derivatives over time for this example trial takes on values near -1,
especially near the end of the reach as the hand converges upon the target. In pure pursuit, the
lateral velocity is not necessarily matched, leading to overall weaker negative correlations. We
find that the correlation increasingly converges towards -1 up to interception in this reach (**Figure
3F**).

When steering strategies are examined over the set of reaches for moving crickets, the PN model
better explains the hand kinematics than the PP model. We applied PN and PP model simulations
to the population of reaches and compared the range vector correlations of the models against
the actual movements. We find that the mean correlation coefficient converges towards -1 over
the duration of reaches, as in the previous example trial, and that the correlation trace of the PN
steering model better matches that of the hand than the PP model, especially as the hand nears
interception (**Figure 3G**). To examine if the lateral velocity of the cricket was matched by hand
velocity, we examined the mean of the lateral velocity of the hand multiplied by the sign of
instantaneous cricket velocity, such that correspondence would give similar mean velocity and no
correlation in direction would converge towards zero. This process was repeated for the lateral
velocity components of PP and PN simulations. The mean lateral velocity of the hand, *Lh,*
increasingly matches that of the cricket, *Lc*, over the time-course of reaches, which is very closely
approximated by the PN model, but not the PP model (**Figure 3H**). Together these results provide
evidence to support a proportional navigation strategy of hand guidance. However, a weakness
of any of these steering models stems from their ability to compensate for visual processing
delays, which represent a critical feature of any real biological or engineered system. Indeed,
when we incorporate a visual delay of cricket position into the PP and PN models we find the
goodness of fit for simulated kinematics break down progressively with visuomotor delay **(Figure
3I)**. A previous study found that a mixed pursuit strategy, including an additive combination of PP
and PN guidance commands, was more robust than single strategy models(Brighton & Taylor,
2019). However, even implementing a mixed PP+PN strategy here, we find that performance
matching the hand kinematics breaks down progressively with delay. Therefore, we sought to
determine visuo-motor delay in marmoset reaching for this task, and furthermore to test if an
explicitly predictive control strategy would be able to better explain observed reach kinematics
while using realistic delay values.

Figure 3. Hand velocity during reaching is consistent with proportional navigation. A: Representative 3D reconstructions of the population of reaches for moving crickets. **B:** At any moment, the hand and cricket velocity can be broken down into two components, the direct and lateral velocity. Direct velocity is the component projected along the range vector while lateral is the orthogonal projection. **C:** The average absolute cricket and hand velocity for the direct and lateral components as illustrated in **B** over interpolated time for the population of reaches. The dashed box indicates the period where the cricket was visible. **D:** Path of the hand (blue) and cricket (red) in an individual example reaching trial. Lighter points indicate the beginning of the reach. Gray lines indicate line of sight, or range vectors, for corresponding hand and cricket points, 1:4 down-sampled. **E:** A pure pursuit (PP) steering strategy aligns hand velocity with the range vector. Proportional navigation (PN) rotates hand velocity to mirror range vector rotations. Overlaid simulations of PP and PN show that PN best approximates this reach. **F:** PN strategies predict matching the lateral component of the hand and cricket velocity while the range to the target is close, and if followed, the derivative of the range vector during pursuit should be in opposite direction of the range vector. For the example reach, the range vectors are shown in gray (matching gray lines in **D**) and their derivatives in black, with the correlation between vectors over time quantified below (black). The trace is aligned on the right to the moment of interception. **G:** The correlations between the range vectors and their derivatives converge towards -1 on average across the population of 78 reaches (individual reaches shown as gray lines, average in black, +/- 1 SEM in gray shade). This analysis was also performed for PP and PN simulations, showing that PN is a better match to actual data. **H**: Lateral velocities from the hand and pursuit simulations were made positive when traveling in the same direction as cricket velocity. Hand lateral velocity increasingly approaches cricket lateral velocity throughout the reach. This is better matched by a PN simulation than a PP simulation. **I:** Pursuit strategy simulations perform worse when a delay is introduced. We include a mixed model simulation, which uses a combination of PN and PP to steer hand velocity and is moderately more robust than proportional navigation alone. Goodness of fit quantified by R^2 values that compare model simulations to actual hand data. All error fields are +/- 1 SEM.

Hand velocity incorporates target predictions to compensate for visuo-motor delays

A key problem in guiding the hand during pursuit is that positional and velocity information about the target arrives at a processing delay in the visual system before it can influence actions. Motor control theories often assume predictive models that estimate future body and target positions to compensate for such delays in planning movements(Reza Shadmehr & Krakauer, 2008; Wolpert & Kawato, 1998). In human experiments employing double-step reaching tasks, where an initial target turns off and a new illuminated target turns on mid-reach, there are visuo-motor delays ranging from 125-225 ms(Brenner & Smeets, 1997; Day & Lyon, 2000; De Brouwer & Spering, 2021; Prablanc & Martin, 1992). By contrast, studies with head-restrained macaques performing double-step tasks have found substantially longer motor delays around 200-300ms(Philippe S. Archambault et al., 2009; P. S. Archambault et al., 2011; Battaglia-Mayer et al., 2013; Dickey et al., 2013; Georgopoulos et al., 1981). This slower timing may reflect constraints due to head-restraint and the artificial nature of the tasks. Thus, it was critical to first asses the visuo-motor delays in the current task.

Visuo-motor delays in dynamic reaching for cricket prey were much shorter than those found in previous non-human primate studies, ranging between 80-100 ms. To estimate visuo-motor delay, we identified discrete cricket acceleration events across task trials in which the cricket increased its motion while the reach was ongoing. The lateral speed of the cricket was filtered to flag the top 5% of moments where lateral velocity increased. We observed that immediately following an increase in lateral cricket velocity, there is a corresponding increase in lateral hand velocity in the same direction that begins around 80 ms (**Figure 4A**). This analysis, however, relies on discrete events in the total data set. To determine if it was consistent throughout the entire data set, we applied a simple linear regression on all the velocity data to predict lateral hand-velocity based on the recent history of cricket lateral velocity (see **Star Methods, Equation 1**). This enabled us to fit a linear kernel for hand velocity based on cricket velocity (i.e., the impulse response function corrected for autocorrelations in cricket velocity). Consistent with the visuo-motor delay observed due to sudden jumps, the linear prediction found a significant prediction coefficient near 80ms (p<.0001 at 87.5ms and p=.0559 at 100ms), but not at other delays (**Figure 4B**). This suggests that changes in hand velocity track the cricket velocity at relatively short delays around 80-100ms, a value much faster than the 200-300ms reported in constrained tasks (Philippe S. Archambault et al., 2009; Battaglia-Mayer et al., 2013; Dickey et al., 2013; Georgopoulos et al., 1981). However, this value is within range of the delay seen in ocular following responses in marmosets, which is close to 75ms(Cloherty et al., 2020). Of interest, like the reaches for live crickets, studies of oculomotor pursuit employ fixation targets that are moving continuously whereas the double-step reaching studies in macaques used instantaneous jumps in target positions.

While the visuo-motor delays are briefer in the current task than previous studies, there may still be significant advantages for taking them into account by planning predictive movements that incorporate the cricket's velocity. To test the role of prediction in guiding the hand, we fit an auto-regressive model of hand velocity that included terms for the visuo-motor delay, τ_d, as well as a prediction time constant, τ_p, that dictated how far into the future the cricket's position was projected based on its velocity. We followed a generative model formulation from a previous study in macaques(Yoo, Tu, Piantadosi, & Hayden, 2020), which described the hand velocity in terms of

Figure 4. Cricket velocity predictively guides hand velocity to compensate for visuo-motor delay. A: Increases in the lateral cricket velocity are followed by increases in lateral hand velocity at a latency of roughly 80ms. A Difference of Gaussian filter was applied to the absolute velocity of cricket motion to identify discrete acceleration events (N=68). The average cricket velocity (in red) time-locked to those events reflects increases in speed at time zero, as expected, while hand velocity follows with a comparable acceleration at a delay (blue), shown as +/- 1 SEM. **B.** A linear regression of lateral hand velocity with the offset lateral cricket velocity from delays ranging from -25 to 150 ms reveals a significant regression coefficient between 80-100 ms. Error shown in 1 SEM estimated from a jackknife procedure. **C:** The multivariate linear model equation used to test the influence of visuomotor delay τ_d and prediction time constant τ_p on the fit of hand velocity. **D:** The model estimates the hand velocity at the next time step (t+1) based on the range vector (R) at time t- τ_d and momentum from the previous hand velocity. A predictive term based on cricket velocity (C) at time t- τ_d (teal dashed lines) is multiplied by the prediction term τ_p to project the range vector into its estimated future location (teal dashed lines). The gray field represents the range of tested prediction τ_p values. **E:** The goodness of fit as a function of τ_d and τ_p is shown in color scale. Upon setting τ_d to 80ms, the best fit of this model is achieved using a τ_p prediction = 115ms (indicated by intersection of black dashed lines). Color scale indicates the goodness-of-fit surface measured by a difference of AIC values between the full predictive model (equation as in **C**) and a model that only uses hand velocity (momentum term) as a predictor. (Left panel) The goodness-of-fit with visuo-motor delay τ_d fixed at 80ms and varying the prediction time constant τ_p is shown with 1 SEM estimated from a jackknife procedure. **F:** When this predictive pursuit model is applied in an approach to simulate hand trajectories (as in Figure 3) it provides a better fit (teal curve) that degrades more gradually than the best mixed PP+PN steering models (orange curve, left vertical axis). Colored traces indicate mean R^2 of simulation fits with 1 SEM fields. The model improvement for the predictive vs. best steering model (black curve, right vertical axis) at 80ms delay and beyond is significantly better for the predictive models (Sign Rank test p=.0105).

a momentum term plus a scaled version of a range vector, $R(t-\tau_d)$, at some visuo-motor delay
(**see equation, Figure 4C**). To compensate for delays in the range vector (**cyan dashed line,**
Figure 4D) the model includes a term to estimate the cricket's future position, $\tau_p\, C(t-\tau_d)$, where τ_p
is a linear temporal weight that determines how far in the future to adjust for the cricket's future
position (**dark gray dashed line, Figure 4D**). We pooled data from 78 reaches for moving crickets
and tested the predictive model fit for visuo-motor delays τ_d over a range from 0ms to 145ms in
the past and prediction time constant τ_p over a range from 250ms in the past up to 250ms in the
future. We assessed model fit by the difference of Akaike Information Criterion (AIC)(Akaike, 1973
) values for the full model (including cricket velocity prediction) against a baseline model that only
contains information about hand velocity (momentum term).

Auto-regressive modeling indicates that hand velocity is best explained by a model where visuo-
motor delay is compensated for by using predictions based on cricket velocity. The AIC difference,
or goodness-of-fit, between the full predictive model and the hand velocity model based on
momentum showed the best fit with a positively correlated diagonal structure reflecting that
positive temporal prediction constants were matched to compensate for visuomotor delays
(**yellow band in Figure 4E**). When we set delay to 80ms as determined above (**Figure 4A, B**),
the best fit is achieved using a prediction constant of 115 ms into the future, (**vertical dashed**
line, Fig. 4E). To visualize standard error in these surface estimates, the cross-section at that
80ms visuo-motor delay was examined as a function of the temporal prediction parameter, with
its confidence intervals estimated using a jackknife procedure (**left panel, Fig. 4E**). The prediction
model demonstrated a significant improvement in model performance for temporal constants
weighted into the future. To obtain a more direct comparison of this model against the PP, PN,
and PP+PN mixed steering models **(Figure 3I)**, we tested this prediction model using a simulation
approach directly comparable to earlier analyses fitting hand kinematics, and tested a range of
visuo-motor delays. When compared against the best steering model, the PP+PN mixed strategy,
the predictive model outperformed it at each delay, showing a more gradual breakdown in
performance for longer visuo-motor delays **(Figure 4F, cyan versus orange curves)**. Taking the
difference in performance between models, the full predictive pursuit model significantly
outperformed the mixed PP+PN steering model at 80 ms (Sign Rank test, p = 0.0105) and longer
delays **(Figure 4F, black trace)**. These findings support that a predictive term compensating for
visuo-motor delay provides a better description of reaching kinematics at realistic delays.
Furthermore, while steering models such as proportional navigation successfully describe
marmoset reaching for moving crickets when they are formulated with no processing delays
(Figure 3), they lack the ability to do so given systematic delays characteristic of the real system.

Discussion

The present findings demonstrate that during an ecologically motivated reaching paradigm for
moving crickets, marmoset monkeys utilize a power grasp that opens earlier in the reach for faster
velocity reaches. Furthermore, marmosets guide their hand movements at very brief visuo-motor
delays, comparable to oculomotor pursuit. In an examination of hand guidance steering models,
we find that proportional navigation explains hand motion reasonably well and provides better fits
than a pure pursuit model (Figure 3). However, these models perform worse in the presence of
realistic visuo-motor delays. To explain the reach kinematics given a realistic visuo-motor delay,
which we estimate to be around 80 ms, we find that it is necessary to use a predictive pursuit
model that incorporates a term to extrapolate the future location of the target based on cricket

velocity. This predictive pursuit model outperforms the proportional navigation or mixed strategy
model(Brighton & Taylor, 2019) at the estimated delay. We find that the prediction constant
closely matches the scale of the visuo-motor delay, extending slightly beyond it about 35 ms into
the future. Such a prediction constant, that remains close to or slightly beyond the visuo-motor
delay, is often close to optimal for maintaining stability and achieving optimal solutions in Bayesian
integration(Körding & Wolpert, 2004) or Smith prediction(Read et al., 2022).

The current finding emphasizes that moving towards realistic dynamics in experimental designs
c(Parker, Abe, Leonard, Martins, & Niell, 2022)an better inform our understanding of reaching
behavior, and provide insights into the underlying neural correlates and the evolutionary contexts
that have driven their development. A previous study also found that primates use prediction in a
pursuit task involving a simulated evading virtual target that was captured by a screen cursor
under joystick control(Yoo et al., 2020). In our study we have adapted their approach to modeling
prediction, but have done so within a task where reaching movements unfold at natural
timescales. The former study identified predictive terms beyond 500 ms into the future, which
match the delays imposed by using a joystick for control in their task. Here we find that similar
predictive strategies are involved to compensate delays when reaching for live prey, but that they
occur at much shorter latencies where visuo-motor delays are comparable to oculomotor pursuit
and under 100 ms. While the present findings were sufficient to characterize the visuomotor
strategies in the current reaching task, future studies could leverage recent advances in head
mounted eye tracking(Meyer, Poort, O'Keefe, Sahani, & Linden, 2018; Parker et al., 2022) to
characterize coordination between head and eye gaze in guiding predictive reaching.

Visually guided insect predation and fruit/leaf eating in an arboreal environment are hypothesized
to play a role in the evolution of primates(Cartmill, 1974; Sussman, Tab Rasmussen, & Raven,
2013). Although not ubiquitous, insect hunting is not only observed in New World monkeys, like
marmosets, but also in Old World monkeys and hominoids(McGrew, 2014; Raubenheimer &
Rothman, 2013). Fronto-parietal neural circuits for motor control in primates are thought to have
evolved in primates to optimize visuo-motor strategies for coordination and speed of reach-to-
grasp movements(Karl & Whishaw, 2013). Much evidence suggests that motor commands are
sent via feedback pathways from premotor cortex to higher order sensory areas in parietal cortex,
and could mediate the forward models necessary to implement predictive on-line control(Reza
Shadmehr & Krakauer, 2008; Wagner & Smith, 2008). The current experimental paradigm shows
that predictive strategies in reaching emerge naturally even within a laboratory context for
dynamic targets, such as crickets, affording new opportunities to study the underlying neural
mechanisms within an ethologically motivated paradigm.

Acknowledgments

315 We would like to thank Dina Graf and members of the Mitchell lab for help with marmoset care
316 and handling. We also thank Cory Miller, Alex Huk, and Cris Niell for comments on this
317 manuscript. This work was supported by NIH grant UF1NS116377 (JFM and LS) and a grant from
318 the Schmitt Program on Integrative Neuroscience (KHW and JFM) and the Del Monte Institute for
319 Neuroscience at the University of Rochester (KHW and LS).

Author Contributions

321 Conceptualization, all authors; formal analysis, all authors; investigation, all authors;
322 methodology, all authors; data acquisition, LS; writing – review & editing, all authors.

Declarations of interests

The authors declare no competing interests.

Inclusion and Diversity

None of the authors of this paper self-identifies as an underrepresented ethnic minority in their field of research or within their geographical location. We support inclusive, diverse, and equitable conduct of research.

STAR METHODS

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contacts Jude Mitchell (jmitch27@ur.rochester.edu) and Kuan Hong Wang (kuanhong_wang@urmc.rochester.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

All data reported in this paper will be shared by the lead contact upon request.

All original code in this paper is available in Supplemental Materials.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All experimental procedures described here are approved by the University Committee on Animal Resources at University of Rochester. 10 marmoset monkeys (5 females, 5 males) were recorded by multiple cameras while reaching for live crickets and cheerios. Their ages ranged from 1.2 years old to 7.1 years old. The crickets used in experiments served as additional enrichment to their daily diet.

METHOD DETAILS

Reaching Task and Video Recording

Reaching experiments were conducted by docking a mobile video recording platform to marmoset family cages within the marmoset colony and recording reaches using 3 GoPro Hero 7 Black cameras **(Figure 1B)**. Marmosets were acclimated to the experimental apparatus and allowed to become comfortable reaching for cheerios as food treats, and then also for crickets. Acclimation was typically less than 30 minutes and was necessary only on the first day of experiments. To calibrate cameras prior to reaching experiments, we recorded a brief video of an 8x6 checkerboard array that was moved and rotated continuously within the reach arena **(Figure 1B,D)**. Frames of this video were extracted and fed into triangulation algorithms provided in the Anipose toolkit(Karashchuk et al., 2021). A manually triggered photography flash aimed at the

cameras was used to deliver a synchronization signal to all three cameras. This created 1-2
frames of video that were significantly over-exposed and easily identified using custom Matlab
(Mathworks) and ffmpeg (Ffmpeg Developers) scripts.

Up to 15 cricket reaches were recorded from an individual marmoset on a given recording day. A
sliding plate at the entry of the reaching enclosure was closed to isolate individual marmosets
briefly from their family groups on the test platform. Reaching trials began by sliding an opaque
acrylic plate upwards that occluded the view of the reaching platform, thus allowing the marmoset
to view the target and begin hunting. The marmoset was able to reach through a large rectangular
aperture (9cm x 13.5cm) that contained two laterally traversing 2mm stainless-steel bars at 3cm
and 6cm height and was built into an acrylic plate. This aperture allowed for a wide range of reach
start points while also preventing the animal from getting outside the test chamber **(Figure 1B)**.
Before lifting the clear acrylic sheet to allow reaching to begin, the crickets were dropped onto the
platform and allowed to move freely. Crickets were kept from crawling off the platform with a 4cm
tall acrylic wall molded into an arc and cemented to the platform with a center maximum y-
dimension of 9cm and base maximum x-dimension of 20cm. The platform was sanded to
eliminate reflections and increase hand and target contrast. Reflection reduction was useful due
to the competition of the features of a tracked subject's reflection against the features of the
subject itself.

Our video recordings were optimized to capture high velocity ballistic reaches. The GoPro Hero
7 Black was specifically chosen for its high-speed capture capability and manual control of
recording settings. High frame rate (1/240s) and high shutter speeds (1/960s) were sufficient to
render all hand features in full detail without observable motion blur. To attain proper exposure at
this high shutter speed and to avoid shining a bright light directly into the eyes of our subjects, a
LED light source (Nanlite Forza 60), was positioned to project light at a 65-degree angle 45cm
from the reaching platform **(Figure 1B)**. Light intensity at the platform was measured to be 4.6E5
Lux, comparable to direct sun on a bright day. The cameras were arranged around the reaching
arena to provide a view from above and a downward angled view from the right and left **(Figure
1E-G)**. The cameras were fixed in position using articulating camera brackets mounted to an
aluminum breadboard.

Video Preparation and DeepLabCut Pose Estimation

Videos were preprocessed by the experimenter to indicate which video frames contained reaches
for crickets and to provide metadata for each reach. Reach success, handedness, and whether
reaches were composed of a single volley or multiple volleys were documented for each reach in
the data workflow. We defined volley multiplicity by whether the hand returned fully to the reach
aperture opening on the platform before extending again. Selected frames spanned from ~200ms
prior to the hand moving through the reaching aperture to full retraction of the arm, which was
typically accompanied by delivery of the target to the mouth. These data were used to generate
standalone video files for each reach that were synchronized for all camera views. Adobe
Premiere Pro was used for video viewing and scoring. Videos were cropped to 760x720 pixels to
remove pixel areas with irrelevant information. DeepLabCut 2.1.8.2 performed feature tracking
from the reaching videos.

DeepLabCut labeling utilized a marker set that was chosen to track the position of fingers, hands,
cricket, and features of the reaching platform. We trained the pre-trained ResNet50 network
supplied by DeepLabCut using 574 manually labeled, manually selected frames of video. The set

of frames was selected to fully characterize the diversity of marmosets, camera angles, hand
positions, and targets of our reaching video data. The final model was trained for 800k iterations
and used the imgaug augmentation pack. Deeplabcut was deployed through Anipose, which
generated unfiltered 2D tracking files using DeepLabCut algorithms. In some later recordings, an
unexpected minor shift in the top-view camera position **(Figure 1F)** offset the accuracy of the
original triangulation calibration. To correct for this, we re-aligned the 2D DeepLabCut tracking
data from each reach video segment using a linear transformation matrix generated by comparing
the arena feature markers **(Figure 1D)** taken from the original calibration videos to the arena
feature markers of each reaching video. We measured the distance each point was moved to
confirm that transformation did not introduce unintended scaling artifacts. These steps were
performed using custom Matlab scripts after converting tracking data files to .csv files. All tracked
points were transformed accordingly, re-packaged into .hd5 files using custom Python scripts,
and then triangulated into 3D space using Anipose.

Reaching Kinematic Data Preparation

Manual reach scoring provided metadata for each reach, which guided the identification of video
segments with hand extension and target motion position data. Position estimates with less than
a .7 estimation likelihood were excluded from analyses, which is a metric provided by DeepLabCut
indicating label accuracy probability. Hand and cricket tracking points were filtered using a 5
sample Gaussian smoothing window. We applied a 180-degree rotation of right-handed reaches
and the cricket position about the reach platform midline to reduce any potential effects of any
handedness-related directionality biases.

We defined a reach as the first tracked appearance of the central hand point beyond the reach
aperture plane to the last tracked point of the cricket. Cricket tracking at the very end of reaches,
during deacceleration and grasp, was typically lost due to occlusion by the fingers or hand.
Windows of reaching data were defined by viewing videos with overlayed tracked points and
marking the beginning and end frame number. These frame number values were used to refer
directly to the .csv tracking data files in custom Matlab analysis scripts. Reaches for crickets that
moved greater than a 0.5 cm distance during the reach and moved continuously were included in
analyses. Crickets that moved this distance by a jump or that ceased motion during the reach
were not included in the data set, nor were crickets that moved less than 0.5cm in total.

Grasp Cluster Analyses

Reaches were selected by visualizing all the tracked points of each of the 78 selected reaches
with moving crickets and excluding 14 reaches with poor knuckle and finger tracking. Each reach
was interpolated to a 51-sample vector where each point is equal to the mean distance between
the pointer and middle knuckle and middle and ring knuckle markers. Knuckle markers were
chosen because they were better tracked and less subject to perturbation by the cricket retaining
wall. Each finger separation vector was z scored and the population was subjected to a kmeans
cluster analysis. The maximum number of clusters was set manually based on testing 2-6
clusters. We repeated clustering using different initial centroids 10 times. The results of this
process were nearly identical to a hierarchical wavelet-based clustering process performed using
the mdwtcluster function provided by Matlab.

Pursuit Models and Velocity Decomposition

To investigate the degree to which marmoset reaching adheres to steering guidance laws such
as pure pursuit and proportional navigation, we adopted Matlab analysis scripts used to
investigate hawk pursuit of moving targets(Brighton & Taylor, 2019). These scripts provided the
framework to simulate marmoset reaches by applying the sample-by-sample steering commands
while matching the magnitude of each step to the measured actual hand speed. Simulation fits to
tracked hand data were measured using the R-squared between the simulated and actual hand
velocity across each reach, then pooled across all reaches. To set initial conditions for simulated
reaches, we matched simulated hand velocity in the first 75ms of the reach to the mean of the
actual hand velocity averaged over the same period. Following Brighton and Taylor(Brighton &
Taylor, 2019), the proportional navigation guidance law utilized by these scripts is defined as

Eq 1 $\dot{\gamma}(t)_{pn} = N\dot{\lambda}(t-\tau)$

Where $\dot{\gamma}(t)_{pn}$ is turning rate, N is an experimentally derived gain constant, $\dot{\lambda}(t)$ is the line of sight
rate, t is time, and τ is a delay parameter. Proportional pursuit is defined as

Eq 2 $\dot{\gamma}(t)_{pp} = -K\delta(t-\tau)$

Where $\dot{\gamma}(t)_{pp}$ is turning rate, K is an experimentally derived gain constant, δ is the range vector
angle with respect to the hand's velocity, t is time, and τ is a delay parameter. A mixed strategy is
defined as the sum of the two above terms as follows:

Eq 3 $\dot{\gamma}(t)_{pnp} = \dot{\gamma}(t)_{pn} + \dot{\gamma}(t)_{pp}$

Model parameters for gain were set by maximizing the total R-squared for hand velocity against
model hand velocity. All reach trials (N=78) were included to fit a gain parameter that was fixed
across trials. Confidence intervals on the R-squared performance were computed from the
standard deviation of trial-by-trial R-squared performance.

Proportional navigation strategies yield parallel pursuit, which can be described as matching
velocity orthogonal to the line of sight, or range vector. To investigate the behavior of range-
vector-relative decomposed velocities in the above steering model frameworks, target and hand
velocity vectors were decomposed into orthogonal velocity components relative to a range vector,
which connects hand position to cricket position. We refer to the velocity component parallel to
the range vector as *direct* velocity. We refer to the velocity component orthogonal to the range
vector as *lateral* velocity. This distinction is important due to the typical movement of crickets
along the perimeter of the reaching platform. When cricket velocity is decomposed in this way,
the majority of cricket velocity is contained within the *lateral* velocity component. A summary of
this decomposition is provided in **Figure 3D**.

A proportional navigation analysis was conducted according to methods used in dragonfly
pursuit(Mischiati et al., 2015). Range vectors were calculated as a subtraction of hand position
from cricket position. The time derivative of the series of range vectors was calculated and
correlated with the range vector itself using cosine vector correlation. Proportional navigation
predicts a negative correlation of -1 between the range vector and its derivative.

Predictive Modeling and Reaching Analyses

We examined if sudden increases in cricket velocity during reaching led to corresponding
increases in hand velocity. To identify increases in cricket velocity we filtered the lateral speed of
the cricket (absolute value of lateral velocity) using a difference filter that was defined by a

Gaussian kernel (σ = 20ms) with negative amplitude at negative lags and positive amplitude as
positive lags. This filter produces the largest values for epochs where low speed is directly
followed by an increase in speed, regardless of whether speed increase is due to the cricket
moving to the left or right. We used a threshold to flag the top 5% of the filtered values and identify
peaks, ensuring that no two peaks occurred within 100 ms of each other. From 78 total reach
trials we flagged 68 events. We averaged the lateral cricket velocity and hand velocity traces from
-100 to 150 ms time-locked around these events (**Figure 4A**). Results were similar if we relaxed
our acceleration threshold to include the top 10% or 20% of events (not shown).

Decomposed hand and cricket velocity components were used to investigate the predictive
influence of cricket velocity on hand velocity. We first tested a linear prediction in which the lateral
hand velocity was predicted by a weighted linear sum of the lateral cricket velocity history. The
linear predictor equations was defined as

Eq 4 $$H_{(t)} = \Sigma^{\tau}\ \kappa_{\tau}C_{(t-\tau)}+\varepsilon_{(t)}$$

where $H_{(t)}$ represents lateral hand velocity at time t, C is the lateral cricket velocity at time t minus
time delay τ and κ_{τ} is a linear prediction coefficient at this time delay. To reduce the number of
parameters we down-sampled the velocity traces from 240 to 80 hertz, giving 12.5ms steps. We
tested a range of τ values from -25 to 150ms. We defined the range vector used to separate direct
and lateral velocities as the vector from the hand to cricket at the start of the reach (when the
hand first entered the arena). We considered the case of updating the range vector at each delay
during the reach as in the full auto-regressive model discussed below and settled upon this
approach for the sake of simplicity.

A recent study examined a pursuit strategy embedded in a 2D coordinate system with a term
incorporating prey velocity to test for the role of prediction (Yoo et al., 2020). To apply the same
model, we employed an auto-regressive linear regression to fit parameters in Eq 5 shown below
which includes a momentum term for hand velocity, and a term based on the predicted range
vector. The parameters for visuo-motor delay, $\tau_{d,}$ and a prediction time constant, $\tau_{p,}$ were fit to
optimize the match to actual hand velocity. This analysis was evaluated using the Matlab
mvregress function. Concatenated velocity and range vector magnitude values from each reach
were supplied to the regression function. Time is defined by frames of video, each of which
captures 1/240 second, or 4.2ms of behavior.

Eq 5 $$H_{(t+1)} = \alpha H_{(t)} + \beta(R_{(t-\tau d)}+\tau_{p}C_{(t-\tau d)}) + \varepsilon_{(t)}$$

In the above Equation 5, $H_{(t+1)}$ represents hand velocity one frame of video into the future (t+1)
relative to the current $H_{(t)}$ velocity at time t. The velocity of the cricket C at a delayed time, t-τ_{d}, is
multiplied by the prediction time constant, $\tau_{p,}$ to project cricket position into the future or past. We
also include a variable representing the range vector from hand to the cricket position as R. This
vector connects hand position at time t and cricket position at t-τ_{d}. The sequence of operations

to generate these variables was to first temporally offset tracked cricket position values according
to t- τ_d. Velocities and range vectors were calculated. Finally cricket velocities were multiplied by
τ_p. The predictor variables are weighted by a set of coefficients α and β, respectively. All terms in
this equation are composed of cartesian x and y constituents and thus are bivariate. These data
were submitted to a least-square regression where $\varepsilon_{(t)}$ represents the residual error.

We varied the values of τ_p and τ_d to determine the effects of visuomotor delay and prediction on
model fit. The parameter τ_d was varied from 0 to 35 frames of video (equivalent to 0ms to 146ms)
such that time shifted cricket position values were aligned to present hand values. Velocities and
range vectors were calculated after this offset was applied. Following this we multiplied cricket
velocity by a range of τ_p values ranging from -60 to 60 frames of video (-250ms to 250ms). To
quantitatively assess the influence of cricket velocity on model fit and to subsequently find offsets
with significant cricket velocity influence, we compared the Akaike Information Criterion (AIC)
(Akaike, 1973) of the model described in Eq5 to analogous models that exclude the range vector
and cricket velocity predictor terms R and C. This was performed for each τ_p and τ_d in the above
range. When the AIC value of one model is more negative than another, it can be interpreted to
have a better fit. More negative AIC values in the cricket inclusion models would suggest that
prediction based on cricket velocity is a significant contributing factor to reach kinematics. We
plotted the AIC difference values and used a jackknife approach (leaving ~8 of 78 reaches out for
10 jackknife iterations) to estimate jackknife AIC mean and calculate standard error.

To compare the predictive pursuit model against steering models we applied the simulation
approach discussed above in Eq. 1-3 and compared its R squared performance for matching
hand velocity against those models. All features of the simulated reaches were applied as before
for the steering models with the guidance law described by Eq. 6 as

Eq 6 $$A_{pred} = \beta(\ R(t-\tau_d) + \tau_p C(t-\tau_d)\ )$$

where A_{pred} is acceleration applied to hand velocity, β is an experimentally derived gain constant,
$R(t-\tau_d)$ is the range vector and $C(t-\tau_d)$ is the cricket velocity, both at delay τ_d, and τ_p is the
prediction time constant. All 78 trials were included to maximize the total R-squared fit across
trials in fitting β and τ_p parameters for each tested visuo-motor delay τ_d. Confidence intervals on
the R-squared performance were computed from the standard deviation of trial-by-trial R-squared
values with statistical tests comparing trial R-squared between models (Sign Rank tests).

Supplemental Discussion

The benefits of utilizing naturalistic behavior for neuroscience research are tempered by
experimental variability. However, modeling approaches provide a tractable alternative to
traditional trial averaging for addressing this variability issue. Naturalistic behavior can tap into
more dynamic aspects of visually-guided reaching including prediction. Furthermore, this
approach presents animals with tasks that they are already accustomed to performing. As a clear
example, our research was conducted with none of the extensive animal training effort that is
required for traditional vision task-related experiments with non-human primates. However, both
the subject and the experimental environment (e.g., crickets) have larger intrinsic variability, which
increases the difficulty of collecting trials that meet inclusion criteria and performing traditional
averaging over repeated trials. We circumvented the lack of identical repeats by performing model
fitting, which made it possible to determine the influence of visuomotor delay prediction on

behavior. Future research could benefit from approaches that leverage the variability of
movements to examine the sensitivity of individual reach trajectories to external or internal
disturbances such as unexpected cricket motion or gaze shifts. Finally, while the reaching task
utilized here is less constrained than traditional reaching tasks, it still imposes some physical
constraints on reaching. This may have precluded more diverse strategies including direct capture
using the mouth, which has been found in the wild(Ngo et al., 2022; Schiel, Souto, Huber, &
Bezerra, 2010). Nonetheless, the experiments and analyses described here represent an
intermediate step towards growing efforts to achieve more naturalistic behavior in laboratory
settings (Datta, Anderson, Branson, Perona, & Leifer, 2019).

Previous model-based approaches have shown that primates utilize simple physical parameters
to predictively pursue an evading virtual target with a joystick (Yoo et al., 2020). A linear prediction
term in the model employed by Yoo and colleagues allowed them to estimate a visuo-motor delay
in their task. Similar to our result, they find evidence for a predictive strategy, although their
estimate of visuo-motor delay differs substantially from the value reported here. This can likely be
attributed to differences in their task design because movements of the cursor on the screen were
limited by the speed allowed under joystick control and momentum terms therein, with overall
much slower timescales for target and cursor speeds on the order of seconds. By contrast the
natural dynamics of marmoset reaches unfold typically in under 200ms. The approach taken in
our study emphasizes the dynamics of natural behavior and motor control, which enables
interrogation of features of the visuomotor system within the natural timescales of visually-guided
reaching and target pursuit.

The derived estimate of marmoset visuomotor delay is consistent with experimentally derived
estimates in humans given the quicker processing and neural transduction times of smaller
primates. A widely studied paradigm that interrogates the on-line visuomotor response delay is
the double step task. In this task, subjects reach for an initial target, which is extinguished and
replaced by a second target during the flight of the hand. The limited number of studies with
macaques that have used double-step reaching tasks to probe online correction have found
substantially longer motor delays than our estimation, around 200-300ms(Philippe S.
Archambault et al., 2009; Battaglia-Mayer et al., 2013; Dickey et al., 2013; Georgopoulos et al.,
1981). Of note, the redirected reaches are typically preceded by a saccadic eye movement at
100-200ms latency to the new jumped location followed by changes in the hand trajectory at 50-
100ms (Archambault et al, 2015). Further, the jumps of targets used in non-human primate
studies have often been much larger than those in human studies, which could impose additional
delays in redirecting attention before movement adjustment. On-line corrective reaction times,
which are analogous to an on-line visuomotor response delay, have been reported in humans
from 125ms to > 200ms (Brenner & Smeets, 1997; Brouwer, Brenner, & Smeets, 2002; Day &
Lyon, 2000; De Brouwer & Spering, 2021; Prablanc & Martin, 1992; Reichenbach, Thielscher,
Peer, Bulthoff, & Bresciani, 2009). This range is likely due to differences in task parameters (eg.
pointing with a finger, holding a rod while reaching, using a pantograph, etc.), measurement
methods (eg. touch plates, receiver operating characteristic analyses, scored deviation from
control, etc.), and the influence of separate motor control pathways with different latencies (eg.
tectospinal versus corticospinal). If we take the bottom line from one of these human experiments
that derives delay from a deviation in hand kinematics (Day & Lyon, 2000), which is 125ms, and
compare that to our estimate of marmoset reaching delay, which is 80ms, we are within range of
a human/marmoset difference seen in a separate, but related comparison of human and
marmoset kinematics. The ocular following response measures the automatic movement of the

eye when presented with coherently moving dot patches. In a study of both humans and
marmosets, humans demonstrate a 125ms delay in ocular following response while marmosets
demonstrate a 75ms delay, which aligns well with our findings (Cloherty et al., 2020). While the
motion of the crickets in this reaching paradigm is not controlled in the same way as in double
step psychophysics experiments, it does demand adjustments in reaching to account for its
motion and thus provides sufficient variation to estimate visuomotor delay through model fitting.

While insect hunting is effective for tests of marmoset reaching and has been quantified in natural
behavior recently in wild marmosets(Ngo et al., 2022), we do find some limitations using this
approach that could be easily addressed in future laboratory studies. In particular, for our study
the number of reaching movements in which the cricket moved during the reach were relatively
small in proportion (~30% of reaches). Further, the number of reaching movements that we could
obtain in any single behavioral session was limited to roughly 15 crickets in a day, more than
enough to satiate a marmoset. Given that only a small fraction of those movements involves
cricket motion during the reach, we obtained relatively few dynamic reaches per day in any given
animal. This limitation of data quantity motivated our pooled approach across 10 marmosets to
analyses. By pooling movements across animals and days we were able to estimate the
visuomotor delay, but it would be difficult to employ this same paradigm to test causal
manipulations of reaching behavior measurable within a single behavioral session. Future work
may seek an intermediate solution without live crickets, such as reaching to small food treats that
are moved on a robotic arm, in which the movement of the target can be controlled under online
feedback during the reach.

We found that DeepLabCut-powered tracking provided reliable descriptions of the trajectory of
the hand supporting the reliability reported in a previous study(Moore, Walker, MacLean, &
Hatsopoulos, 2022). DeepLabCut affords a key advantage for studying natural movements over
other methods that affix optical tracking markers to hands and arms, which may impose
constraints on animal reaching and require additional time and care to install. We found that
GoPro cameras were well suited for data collection in these reaching tasks due to their ability to
capture high frame rate video with user-set shutter speeds that reduce motion blur. While the
arrangement of cameras was optimized for a 3-D rendering of the top-down view of the hand,
occlusion of the fingers was common due to the wide variability of reach trajectories, the
pronation of the hand, and the tendency of the fingers to make contact with the cricket retaining
wall. Regardless, it is impressive that DLC could provide tracking of features with such small
degrees of error and could discriminate between the right and left hands without confusing their
features. One limitation of our current tracking approach is that it is difficult to track body parts
that are occluded by featureless fur such as the shoulder or elbow. Future studies could
characterize the detailed interaction between shoulder and elbow movement in positioning the
final hand position during motor control, as well as the role of gaze in guiding reaching
movements. Finally, we found it difficult to achieve low-noise 3D depictions of reaching and
grasping and thus used 2D tracking for our analyses. We feel this problem is likely solved by
additional camera angles to both generate more camera coverage overlap and to better cover
areas that were poorly captured by the set of 3 cameras used in this study.

  Supplemental Figures

Supplemental Figure 1. Tracking accuracy in DeepLabCut video analysis: Three DeepLabCut networks were trained to show tracking errors from the testing and training sets in RMSE quantified in pixels and converted to mm using the pixel to mm conversion respective to the above-view camera (Figure 1F). The label 'stage' on the x-axis refers to the reach platform stage and corresponding arena points (Figure 1D). Fingers refers to the average of both the fingertip and knuckle points, which equals 10 points for each hand. The average RMSE values depict the stability of the training approach over 3 different training iterations. Each point depicts the labeling error of a model trained with a different "shuffle" of the test and training data. The model used for all tracking analyses performed with a testing RMSE of 2.79 pixels for the stage and arena markers, 3.28px for the cricket marker, 8.21px for the left hand, 5.1px for the left fingertips, 5.9px for the left knuckles, 6.2px for the right hand, 5.8px for the right fingertips, and 5.0px for the right knuckles. For comparison, the width of the wrist is typically around ~1.5cm, which equates to 37px. The length of the cricket ranged from ~1.25cm to ~3cm. We utilize the hand marker for reach modeling and correlation analyses because of its tracking reliability across all three camera views and its reduced susceptibility to occlusion-related tracking drops, which were more common in the finger markers.

Supplemental Figure 2. Further grasp characterization in relation to hand speed and distance. A: Grasp visualization from a low camera angle to show parallel hand orientation and simultaneous finger closure. **B-D:** Scatter plots illustrate the relationships between distance to target, hand speed, and finger separation across reaching movements. Points show a 1/4 subsample of all tracked points from the population of reaches. Each plot is fit with a line (B,C) or 2nd order polynomial (D), which is shown with a 95% confidence interval. **B:** Across many reaches the finger separation increases with proximity to target. **C:** Finger separation increases with decreased hand speed. **D:** Hand speeds are lower when at greater distances from target and when in close proximity to the target.

REFERENCES

Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle In B. Petrov & F. Csaki (Eds.), *2nd International Symposium on Information Theory* (pp. 267-281). Budapest, Hungary Akadémiai Kiadó

Archambault, P. S., Caminiti, R., & Battaglia-Mayer, A. (2009). Cortical Mechanisms for Online Control of Hand Movement Trajectory: The Role of the Posterior Parietal Cortex. *Cerebral Cortex, 19*(12), 2848-2864. doi:10.1093/cercor/bhp058

Archambault, P. S., Ferrari-Toniolo, S., & Battaglia-Mayer, A. (2011). Online control of hand trajectory and evolution of motor intention in the parietofrontal system. *J Neurosci, 31*(2), 742-752. doi:10.1523/JNEUROSCI.2623-10.2011

Battaglia-Mayer, A., Ferrari-Toniolo, S., Visco-Comandini, F., Archambault, P. S., Saberi-Moghadam, S., & Caminiti, R. (2013). Impairment of Online Control of Hand and Eye Movements in a Monkey Model of Optic Ataxia. In *Cerebral Cortex* (Vol. 23, pp. 2644-2656): Oxford Academic.

Bicca-Marques, J. C. (1999). Hand specialization, sympatry, and mixed-species associations in
callitrichines. *Journal of Human Evolution, 36*(4), 349-378.
doi:https://doi.org/10.1006/jhev.1998.0272

Bishop, A. (1962). CONTROL OF THE HAND IN LOWER PRIMATES. *Annals of the New York Academy of
Sciences, 102*(2), 316-337. doi:10.1111/j.1749-6632.1962.tb13649.x

Brenner, E., & Smeets, J. B. (1997). Fast Responses of the Human Hand to Changes in Target Position. *J
Mot Behav, 29*(4), 297-310. doi:10.1080/00222899709600017

Brighton, C. H., & Taylor, G. K. (2019). Hawks steer attacks using a guidance system tuned for close pursuit
of erratically manoeuvring targets. *Nat Commun, 10*(1), 2462. doi:10.1038/s41467-019-10454-z

Brouwer, A.-M., Brenner, E., & Smeets, J. B. J. (2002). Hitting moving objects: is target speed used in
guiding the hand? *Experimental Brain Research, 143*(2), 198-211. doi:10.1007/s00221-001-0980-
x

Cartmill, M. (1974). Rethinking primate origins. *Science, 184*(4135), 436-443.
doi:10.1126/science.184.4135.436

Castiello, U., & Dadda, M. (2019). A review and consideration on the kinematics of reach-to-grasp
movements in macaque monkeys. *J Neurophysiol, 121*(1), 188-204. doi:10.1152/jn.00598.2018

Catania, K. C. (2009). Tentacled snakes turn C-starts to their advantage and predict future prey behavior.
Proceedings of the National Academy of Sciences, 106(27), 11183 - 11187. Retrieved from
https://www.pnas.org/doi/epdf/10.1073/pnas.0905183106

Cloherty, S. L., Yates, J. L., Graf, D., DeAngelis, G. C., & Mitchell, J. F. (2020). Motion Perception in the
Common Marmoset. *Cereb Cortex, 30*(4), 2658-2672. doi:10.1093/cercor/bhz267

Datta, S. R., Anderson, D. J., Branson, K., Perona, P., & Leifer, A. (2019). Computational Neuroethology: A
Call to Action. In *Neuron* (Vol. 104, pp. 11-24): Cell Press.

Day, B. L., & Lyon, I. N. (2000). Voluntary modification of automatic arm movements evoked by motion of
a visual target. *Experimental Brain Research, 130*(2), 159-168. doi:10.1007/s002219900218

De Brouwer, A. J., & Spering, M. (2021). *Eye-hand coordination during online reach corrections is task-
dependent*. Cold Spring Harbor Laboratory. Retrieved from
https://dx.doi.org/10.1101/2021.06.13.448238

De Bruin, N., Sacrey, L. A. R., Brown, L. A., Doan, J., & Whishaw, I. Q. (2008). Visual guidance for hand
advance but not hand withdrawal in a reach-to-eat task in adult humans: Reaching is a composite
movement. In *Journal of Motor Behavior* (Vol. 40, pp. 337-346).

Desmurget, M., & Grafton, S. (2000). Forward modeling allows feedback control for fast reaching
movements. *Trends in Cognitive Sciences, 4*(11), 423-431. doi:10.1016/s1364-6613(00)01537-0

Dickey, A. S., Amit, Y., & Hatsopoulos, N. G. (2013). Heterogeneous neural coding of corrective movements
in motor cortex. *Front Neural Circuits, 7*, 51. doi:10.3389/fncir.2013.00051

Fox, D. M., Mundinano, I. C., & Bourne, J. A. (2019). Prehensile kinematics of the marmoset monkey:
Implications for the evolution of visually-guided behaviors. In *Journal of Comparative Neurology*
(pp. 1495-1507).

Fujioka, E., Aihara, I., Sumiya, M., Aihara, K., & Hiryu, S. (2016). Echolocating bats use future-target
information for optimal foraging. *Proc Natl Acad Sci U S A, 113*(17), 4848-4852.
doi:10.1073/pnas.1515091113

Georgopoulos, A. P., Kalaska, J. F., Caminiti, R., & Massey, J. T. (1983). Interruption of motor cortical
discharge subserving aimed arm movements. *Experimental Brain Research, 49*(3), 327-340.
doi:10.1007/BF00238775

Georgopoulos, A. P., Kalaska, J. F., & Massey, J. T. (1981). Spatial trajectories and reaction times of aimed
movements: effects of practice, uncertainty, and change in target location. *Journal of
Neurophysiology, 46*(4), 725-743. doi:10.1152/jn.1981.46.4.725

Hook, M. A., & Rogers, L. J. (2008). Visuospatial reaching preferences of common marmosets (Callithrix jacchus): an assessment of individual biases across a variety of tasks. *J Comp Psychol, 122*(1), 41-51. doi:10.1037/0735-7036.122.1.41

Kaas, J. H., Gharbawie, O. A., & Stepniewska, I. (2011). The organization and evolution of dorsal stream multisensory motor pathways in primates. *Front Neuroanat, 5*, 34. doi:10.3389/fnana.2011.00034

Kaas, J. H., & Stepniewska, I. (2016). Evolution of posterior parietal cortex and parietal-frontal networks for specific actions in primates. *Journal of Comparative Neurology, 524*(3), 595-608. doi:10.1002/cne.23838

Karashchuk, P., Rupp, K. L., Dickinson, E. S., Walling-Bell, S., Sanders, E., Azim, E., . . . Tuthill, J. C. (2021). Anipose: A toolkit for robust markerless 3D pose estimation. *Cell Reports, 36*(13), 109730. doi:10.1016/j.celrep.2021.109730

Karl, J. M., & Whishaw, I. Q. (2013). Different evolutionary origins for the reach and the grasp: an explanation for dual visuomotor channels in primate parietofrontal cortex. *Front Neurol, 4*, 208. doi:10.3389/fneur.2013.00208

Körding, K. P., & Wolpert, D. M. (2004). Bayesian integration in sensorimotor learning. *Nature, 427*(6971), 244-247. doi:10.1038/nature02169

Lenoir, M., Musch, E., Janssens, M., Thiery, E., & Uyttenhove, J. (1999). Intercepting Moving Objects During Self-Motion. *J Mot Behav, 31*(1), 55-67. doi:10.1080/00222899909601891

Marzke, M. W. (1997). Precision grips, hand morphology, and tools. *Am J Phys Anthropol, 102*(1), 91-110. doi:10.1002/(sici)1096-8644(199701)102:1<91::Aid-ajpa8>3.0.Co;2-g

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience, 21*(9), 1281-1289. doi:10.1038/s41593-018-0209-y

McGrew, W. C. (2014). The 'other faunivory' revisited: Insectivory in human and non-human primates and the evolution of human diet. *Journal of human evolution, 71*, 4-11. doi:10.1016/j.jhevol.2013.07.016

Merchant, H., & Georgopoulos, A. P. (2006). Neurophysiology of perceptual and motor aspects of interception. *J Neurophysiol, 95*(1), 1-13. doi:10.1152/jn.00422.2005

Meyer, A. F., Poort, J., O'Keefe, J., Sahani, M., & Linden, J. F. (2018). A Head-Mounted Camera System Integrates Detailed Behavioral Monitoring with Multichannel Electrophysiology in Freely Moving Mice. *Neuron, 100*(1), 46-60.e47. doi:10.1016/j.neuron.2018.09.020

Mischiati, M., Lin, H. T., Herold, P., Imler, E., Olberg, R., & Leonardo, A. (2015). Internal models direct dragonfly interception steering. *Nature, 517*(7534), 333-338. doi:10.1038/nature14045

Moore, D. D., Walker, J. D., MacLean, J. N., & Hatsopoulos, N. G. (2022). Validating markerless pose estimation with 3D X-ray radiography. *J Exp Biol, 225*(9). doi:10.1242/jeb.243998

Ngo, V., Gorman, J. C., De la Fuente, M. F., Souto, A., Schiel, N., & Miller, C. T. (2022). Active vision during prey capture in wild marmoset monkeys. *Curr Biol, 32*(15), 3423-3428 e3423. doi:10.1016/j.cub.2022.06.028

Parker, P. R. L., Abe, E. T. T., Leonard, E. S. P., Martins, D. M., & Niell, C. M. (2022). Joint coding of visual input and eye/head position in V1 of freely moving mice. *Neuron, 110*(23), 3897-3906.e3895. doi:10.1016/j.neuron.2022.08.029

Prablanc, C., & Martin, O. (1992). Automatic Control During Hand Reaching at Undetected Two-Dimensional Target Displacements. *Journal of Neurophsyiology, 67*(2), 455-469.

Raubenheimer, D., & Rothman, J. M. (2013). Nutritional ecology of entomophagy in humans and other primates. *Annu Rev Entomol, 58*, 141-160. doi:10.1146/annurev-ento-120710-100713

Read, J. C. A., Kaspiris-Rousellis, C., Wood, T. S., Wu, B., Vlaskamp, B. N. S., & Schor, C. M. (2022). Seeing the future: Predictive control in neural models of ocular accommodation. *J Vis, 22*(9), 4. doi:10.1167/jov.22.9.4

Reichenbach, A., Thielscher, A., Peer, A., Bulthoff, H. H., & Bresciani, J. P. (2009). Seeing the hand while
reaching speeds up on-line responses to a sudden change in target position. *J Physiol, 587*(Pt 19),
4605-4616. doi:10.1113/jphysiol.2009.176362
Sacrey, L. A. R., & Whishaw, I. Q. (2012). Subsystems of sensory attention for skilled reaching: Vision for
transport and pre-shaping and somatosensation for grasping, withdrawal and release. In
Behavioural Brain Research (Vol. 231, pp. 356-365): Elsevier B.V.
Sartori, L., Camperio Ciani, A., Bulgheroni, M., & Castiello, U. (2013). Reaching and grasping behavior in
Macaca fascicularis: a kinematic study. *Experimental Brain Research, 224*(1), 119-124.
doi:10.1007/s00221-012-3294-2
Schiel, N., & Souto, A. (2017). The common marmoset: An overview of its natural history, ecology and
behavior. In *Developmental Neurobiology* (Vol. 77, pp. 244-262).
Schiel, N., Souto, A., Huber, L., & Bezerra, B. M. (2010). Hunting strategies in wild common marmosets are
prey and age dependent. In *American Journal of Primatology* (Vol. 72, pp. 1039-1046).
Shadmehr, R., & Krakauer, J. W. (2008). A computational neuroanatomy for motor control. *Experimental
Brain Research, 185*(3), 359-381. doi:10.1007/s00221-008-1280-5
Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and adaptation
in motor control. *Annu Rev Neurosci, 33*, 89-108. doi:10.1146/annurev-neuro-060909-153135
Smith, J. M., & Smith, A. C. (2013). An investigation of ecological correlates with hand and foot morphology
in callitrichid primates. *Am J Phys Anthropol, 152*(4), 447-458. doi:10.1002/ajpa.22378
Sussman, R. W., Tab Rasmussen, D., & Raven, P. H. (2013). Rethinking primate origins again. *Am J Primatol,
75*(2), 95-106. doi:10.1002/ajp.22096
Sustaita, D., Pouydebat, E., Manzano, A., Abdala, V., Hertel, F., & Herrel, A. (2013). Getting a grip on
tetrapod grasping: form, function, and evolution. *Biol Rev Camb Philos Soc, 88*(2), 380-405.
doi:10.1111/brv.12010
Wagner, M. J., & Smith, M. A. (2008). Shared Internal Models for Feedforward and Feedback Control. In
Journal of Neuroscience (Vol. 28, pp. 10663-10673): Society for Neuroscience.
Wang, X., Liu, Y., Li, X., Zhang, Z., Yang, H., Zhang, Y., . . . He, Z. (2017). Deconstruction of Corticospinal
Circuits for Goal-Directed Motor Skills. *Cell, 171*(2), 440-455 e414. doi:10.1016/j.cell.2017.08.014
Whishaw, I. Q., & Gorny, B. (1994). Arpeggio and fractionated digit movements used in prehension by
rats. *Behavioural Brain Research, 60*, 15-24.
Wolpert, D. M., & Kawato, M. (1998). Multiple paired forward and inverse models for motor control.
Neural Networks, 11(7-8), 1317-1329. doi:10.1016/s0893-6080(98)00066-5
Yoo, S. B. M., Tu, J. C., Piantadosi, S. T., & Hayden, B. Y. (2020). The neural basis of predictive pursuit. In
Nature Neuroscience (Vol. 23, pp. 252-259): Nat Neurosci.

805