Tumor Spheroids Layered in an Imageable Cancer Environment (T-SLICE): a novel *in*

***vitro* platform to study tumor biology**

Authors/Affiliations:

Morgan Pugh-Toole[*1,2], Nicholas Dawe[*2,3], Jeanette E. Boudreau[1,2,4], Brendan M. Leung[1,2,3,5]

[1]Department of Pathology, Dalhousie University

[2]Beatrice Hunter Cancer Research Institute

[3]School of Biomedical Engineering, Dalhousie University

[4]Department of Microbiology & Immunology, Dalhousie University

[5]Department of Applied Oral Sciences, Dalhousie University

Correspondance: Dr. Brendan Leung, bleung@dal.ca

Mailing Address: 5891 University Avenue, Dentistry Building, Room 5193, Halifax, NS, B3H

1W2

*These authors contributed equally to the manuscript.

Keywords: Tumor Microenvironment, 3-D culture system, hypoxia, microscopy

19
20

Summary

Cancer treatment is shifting away from traditional to precise therapeutic approaches informed by patient and tumor profiles. To achieve better outcomes for cancer patients, it will be necessary to design and test therapies that target the heterogeneity of tumors that exists within and between patients and even within the same tumor. Existing *in vitro* model systems do not fully recapitulate the physio-chemical gradients of tumors and may fall short of representing a living organism. Mouse models remain the gold standard for preclinical testing but lack the throughput needed to quickly understand mechanisms and iterate quickly to maximize therapeutic efficacy. Microfluidic systems can fill this gap but remain out of reach for most biologic laboratories due to their complexity in assembly and use. We created T-SLICE (Tumor Spheroids Layered in an Imageable Cancer Environment) to enable high-throughput imageable studies within a complex simulated tumor microenvironment (TME). Tumor spheroids are seeded on a monolayer of fibroblasts and cultured between two parallel, closely spaced glass coverslips (300-700 μm). The restrictive gap distance limits permeation of nutrients and oxygen to create gradients that recapitulate those observed in the *in vivo* TME. In T-SLICE, physio-chemical gradients are defined by cellular metabolism, akin to an actual tumor. We demonstrate that this is associated with changes in cellular viability, mitochondrial trans-membrane potential, and proliferation. T-SLICE provides a live-cell imaging compatible, high throughput platform that allows real-time monitoring of cellular phenotypes, function, and sociology within a cell-driven TME physio-chemistry amenable to therapeutic optimization, screening, and patient avatar modelling.

Introduction

Tumors are dynamic communities of cells that together form a microenvironment with gradients of nutrient and oxygen availability[1]. Within the tumor microenvironment (TME), the relative frequencies of stromal and tumor cells can vary, impacting tumor progression and prognosis[2]. Research has begun to uncover the dynamic nature of the TME and physio-chemical factors that allow cancer cells to persist and invade healthy tissues, but model systems have not allowed faithful recapitulation of these complex TMEs[1,3].

The hypoxic cellular response is of particular significance as it is implicated in both DNA damage and repair and apoptosis of malignant cells[4]. Acute hypoxic responses lead to the accumulation of reactive oxygen species (ROS), leading to DNA damage, and chronic hypoxia leads to impairments in DNA damage repair. Over time, such responses in cancer cells can lead to the accumulation of irreparable DNA mutations and give rise to advanced malignancies[5]. Cell metabolism within the TME generates physio-chemical gradients that increase with tumor size and availability of blood supply, compounded by relatively poor vascularization and reduced ability of oxygen and nutrients to diffuse through the tumor[6]. A nutrient-poor, hypoxic core that favors anaerobic metabolism is established as a result. Tumor cells persisting in these conditions must adapt, which often manifests as decreased proliferation, increased mutation rates, and resistance to apoptosis[6,7].

The heterogeneous TME drives diversification of cancer cell phenotypes, making it unlikely that a single intervention will completely eradicate the tumor. Using mainstream drug development pipelines, oncology drug candidates promising in preclinical trials have a low probability of success by the time they reach phase III clinical trials, with a cumulative failure rate of approximately 97%[8]. Immunotherapies are showing promise for treating solid tumors, but

65   most patients still fail to achieve long-term durable responses[9]. Likely, these studies reflect that

66   most tumors are only partially sensitive to therapeutic interventions because their diversity is

67   underestimated in existing preclinical models. Strategies to experimentally recapitulate the

68   complex TME are urgently required.

69   To enable high-throughput and mechanistic studies that recapitulate the TME, we created

Tumor Spheroids Layered in an Imageable Cancer Environment (T-SLICE). The T-SLICE

culture model is designed to be compatible with standard life science laboratory techniques. T-

SLICE incorporates tunable physio-chemical parameters to recreate specific features of the TME

of different malignancies for experimental study. In T-SLICE, a fibroblast monolayer is seeded

with tumor spheroids between two glass coverslips with a narrow vertical gap (300-700 μm).

The small gap space negates most convective transport caused by bulk fluid flow, creating a

diffusional resistance that is inversely proportional to its height. As a result of metabolism, cells

closer to the edge of the parallel glass coverslips preferentially receive oxygen; thus, a hypoxic

gradient is formed from the periphery toward the middle of the T-SLICE. The monolayer of

fibroblasts seeded within the gap creates a gradient profile that simulates a TME of a tumor the

same size as the coverslip. By adding small spheroids within the gap space, we address the

shortcomings of conventional spheroid culture because we uncouple tissue architecture from

metabolically-generated physio-chemical gradients. T-SLICE enables control of physio-chemical

gradients independently from tumor tissue density, allowing live imaging of the hypoxic region

of the TME, and therefore empowers laboratories to better test interventions in a faithful *in vitro*

system.

Methods

3-D mold design and printing

T-SLICE design and construction are depicted in Figure 1A-B. T-SLICE chips were designed using Autodesk® Fusion 360™ Computer-Aided Design (CAD) software as '.stl' files, then converted to '.ctb' files using Chitubox® 3-D printing preprocessing software. Designs were 3-D printed using an ELEGOO Mars 2 Pro MSLA 3-D Printer (ELEGOO, Shenzhen. China). Molds were created from Standard LCD UV-Curing Photopolymer Rapid Resin (ELEGOO, Shenzhen. China) in 0.05mm layers, each cured by 6 seconds of UV light exposure (405nm). Resin molds were washed in isopropanol (20 min) and UV-cured (405 nm, 30 min) (ELEGOO Mercury Plus Washing and Curing Machine, Shenzhen. China). Uncured resin monomers were extracted from molds by soaking in 95% ethanol (3x24h, room temperature, with agitation).

T-SLICE chips were prepared by pouring Polydimethylsiloxane (PDMS) (10:1 elastomer to curing agent, Sylgard 184™ Silicone Elastomer Kit, Corning) into the resin molds, degassed in a vacuum chamber for 15 min, then cured (60° for 24 h) and removed using forceps. A No. 1 18mm coverslip was covalently bonded to the underside of the chip by plasma oxidation (1-2 min.). Before use in cell culture, T-SLICE chips were sterilized in 70% ethanol and exposed to UV light for 30 min.

Cell Culture

The NIH/3T3 (fibroblast), MCF-7 (breast cancer), and HCC-1806 (breast cancer) cell lines were obtained from the American Type Culture Collection (ATCC), and the ID8 $Trp53^{-/-}$ (ovarian cancer) cell line was generously gifted by Dr. Madhuri Koti (Queen's University). Cells were maintained in a humidified incubator at 37°C and 5% CO_2. MCF-7 and NIH/3T3 cells were

maintained Dulbecco's Modified Eagle Media (DMEM; Gibco) supplemented with 10% fetal

bovine serum (FBS; Gibco) and 1% antibiotic-antimycotic (AA;100 IU/mL penicillin, 0.1

mg/mL streptomycin and 0.25 µg/mL amphotericin-B, Gibco,). HCC-1806 cells were maintained

in RPMI-1640 media (Gibco) supplemented with 10% FBS and 1% AA. ID8-*Trp53*[-/-] cells[10]

were maintained in DMEM supplemented with 10% FBS, 1% penicillin-streptomycin (PS;

Wisent), and 1% Insulin Transferrin Selenite (ITS; Wisent)[10].

Spheroid culture in 384-hanging drop plates and T-SLICE assembly

Tumor spheroids were created using Perfecta3-D^{TM} 384-well hanging drop culture plates (3-D

Biomatrix Inc. Ann Arbor, MI). Before seeding, plates were pretreated in 0.1% w/v Pluronic® F-

108 solution (Sigma-Aldrich) in distilled water, then rinsed with distilled water and UV-

sterilized for 30 min. MCF-7 (50 cells) or HCC1806 and ID8-*Trp53-/-* (100 cells) were seeded in

25 µL droplets containing culture media supplemented with 2.4 mg/mL MethoCel® A4M

(Sigma-Aldrich) to create spheroids with diameters of approximately 150 µm. To prevent

spheroid merging, droplets were seeded in a lattice pattern, yielding 192 spheroids for each 384-

well hanging drop plate. Spheroids were maintained under standard cell culture conditions

(37°C, 5% CO_2). Following spheroid formation (24h for MCF-7; 48h for HCC1806 and ID8-

Trp53-/-), cells were pooled and seeded onto fibroblast monolayers in T-SLICE.

T-SLICE culture is assembled within a standard six-well cell culture-coated plate.

NIH/3T3 fibroblasts are cultured on autoclaved No. 1 18mm round coverslips (VWR) pretreated

with fibronectin (10 µg/mL in PBS; Sigma-Aldrich) for 45 min. Coverslips are with NIH/3T3

fibroblasts overnight at 200,000 cells per coverslip (400,000 cells/mL), incubated overnight to

reach confluency, and then seeded with 192 spheroids/coverslip. Spheroids were collected,

centrifuged at 200 x *g*, and resuspended in 500 μL of complete cell culture media, which was

then used to replace the culture media atop the fibroblasts. Fibroblasts and spheroids were co-

cultured overnight, then the T-SLICE chip was placed, and 2mL of cell culture media was added

to the sides until the area under the chip was filled by capillary action. Media was changed in

500μL increments every 24h. Control wells without T-SLICE were cultured in parallel. Zones of

T-SLICE were defined as middle, intermediate and peripheral zones, representing concentric

rings of 0-3mm, 3-6mm or 6-9mm diameter, respectively.

EF5 for characterization of hypoxic responses within T-SLICE of MCF-7, HCC1806 and ID8

Trp53-/- cells

To characterize hypoxic cellular responses, we cultured T-SLICE in the presence of 100 μM of

the EF5 Hypoxia Detection Kit (Sigma-Aldrich). Cell culture media was supplemented with 100

μM EF5 incorporated into the media for 6, 12, 24, or 48 h. EF5 (2-(2-Nitro-1H-imidazol-1-yl)-

N-(2,2,3,3,3-pentafluoropropyl) acetamide), is a 2-nitroimidazole compound that is reduced in

hypoxic cells to form adducts at the cell membrane that can be fluorescently labelled with a

fluorophore-conjugated anti-EF5 antibody[11]. After the T-SLICE chip was removed, cells were

fixed in 4% paraformaldehyde in PBS overnight at 4°C. Cells were then washed and prepared for

imaging according to the manufacturer's instructions, using the anti-EF5, clone ELK3-51

Cyanine 3 conjugate mouse monoclonal antibody (5μg/mL in 1% BSA/PBS; Sigma-Aldrich).

Imaging was done with the EVOS™ Light Cube, RFP 2.0 (Ex: 542 nm; Em: 593nm).

Measuring cellular proliferation using the EdU Cell Proliferation Kit

Cell proliferation in T-SLICE zones was analyzed using an EdU (5-ethynyl-2'-deoxyuridine)

Cell Proliferation Kit (Click-iT™ EdU Cell Proliferation Kit for Imaging, Alexa Fluor™ 488

dye: Invitrogen). Cell culture media was supplemented throughout with 10 uM EdU. After 72h,

T-SLICE chips were removed, and the cells were fixed with 4% paraformaldehyde (PFA)

solution in PBS for 15 min at RT, then stained according to the manufacturer's instructions.

Imaging was performed using the EVOS™ Light Cube, GFP 2.0.

Assessment of cell viability using Calcein-AM and ethidium homodimer-1

Imaging was performed on the EVOS™ FL Auto 2 Cell Imaging System, all assessed with

ImageJ. Viability of cells cultured within T-SLICE was assessed using Calcein-AM

(ThermoFisher Scientific) to label live cells and ethidium homodimer-1 (EthD; Invitrogen) to

label dead cells. At specified timepoints, T-SLICE chips were removed, cells washed once with

PBS, and stained with a 1μM Calcein-AM, 1 μM ethidium homodimer-1 solution in PBS for 30

min at room temperature (RT) in the dark. Cells were washed twice with 1 mL PBS and then

immersed in PBS for imaging using the EVOS™ Light Cube, GFP 2.0 (Ex: 482 nm, Em: 524

nm) and EVOSTM Light Cube, RFP 2.0 (Ex: 542 nm, Em: 593 nm).

Measuring Mitochondrial Membrane Potential using MitoView 633

MitoView™ 633 (Biotium) was used to measure mitochondrial membrane potential (MMP)

according to the manufacturer's recommendation. Cell culture media was supplemented with 100

nM MitoView 633 (Biotium) for the duration of T-SLICE culture. Live T-SLICE cultures were

imaged every 24h using the EVOS™ Light Cube, Cy5 2.0 (Ex: 635 nm, Em: 692 nm).

Statistical analysis

Statistical analysis of data was performed using GraphPad Prism 9 (GraphPad Software). Two-

way analysis of variance (ANOVA) with Tukey's multiple comparisons test was performed to

assess multi-group comparisons. *P*-values are represented as follows: *$p<0.05$; **$p<0.01$;*

****$p<0.001$; ****$p<0.0001$.*

Results

Design and optimization of T-SLICE

We sought to create a synthetic TME that recapitulates the physio-chemical properties of a solid

tumor, focusing on oxygen tension and mitochondrial stress. We iterated on design, considering

both ease of device fabrication and experimental use. Chip geometry, cell growth surface area,

and distance between the two parallel-spaced glass coverslips are all factors that can influence

the biochemical gradients within T-SLICE and were therefore considered together during

conceptualization and design. Circle and square geometries were evaluated using prototypes with

gap heights ranging from 300-700 μm. In comparison to squares, circular geometry has a

symmetrical center-to-edge distance for easier measurement of cell distances. To recapitulate

tumor microenvironments with different oxygen tensions, we created T-SLICE chips with a

range of gap heights **(Figure 1A, B).** For the current study, we chose a circular design using an

18mm glass coverslip bonded on the underside of a PDMS chip of 24 mm diameter and 4 mm

total height. Three different T-SLICE chip gap heights of 300 μm, 500 μm, or 700 μm were

created and placed on top of a bottom coverslip, between which cells are cultured.

Fibroblasts are a prominent cell type found in the stromal region of tumors and play an

essential role within the TME. To capture aspects of the TME within T-SLICE, we used

NIH/3T3 embryonic murine fibroblasts as the gradient-driving monolayer. On an 18 mm

coverslip within a 6-well plate, we grew NIH/3T3 cells to confluency and seeded tumor

spheroids. We tested a range of cell seeding densities and found that 2×10^5 NIH/3T3 cells best

established a confluent monolayer in 24h, onto which tumor spheroids could be supported

without direct contact with the coverslip and any associated deformation due to spheroid

adhesion spreading **(Figure 1C)**. In parallel, we optimized cancer cell seeding densities in

hanging drop culture to create spheroids with a diameter <150 µm. We chose this limit to

exclude the possibility of generating internal physio-chemical gradients and to ensure that the

hypoxia level of a given spheroid is dependent only on its position under the T-SLICE chip. The

small size of the spheroid further allows imaging and microscopic detection and quantification of

changes within it, as confirmed using brightfield microscopy (**Figure 1C**).

T-SLICE culture generates sustained hypoxic gradients

Hypoxia is a hallmark biochemical pressure within heterogeneous TME of solid tumors that

results from cellular oxygen consumption and limited availability, especially in regions distal to

the blood supply[12]. We expected that hypoxia would occur most prominently at the center of the

T-SLICE. To assess this, we identified three concentric zones for analysis: middle (M, 0– 3 mm

from centre), intermediate (I, 3– 6 mm from centre), and periphery (P, 6– 9 mm from centre).

Using EF5 staining, we measured hypoxia within T-SLICE after 6, 12, 24 and 48h of culture. At

all three gap heights tested, increasing time resulted in greater hypoxia among MCF7 cells

($p<0.0001$, **Figure 2**); this result was confirmed in a second cell line, HCC1806

(**Supplementary Figure 1**). For all cell lines and gap heights tested, EF5 fluorescence intensity

was highest in the middle zone of T-SLICE and progressively decreased in the intermediate and

peripheral zones. EF5 staining was observed as early as 6 h and increased during the 48h during

which we studied cells in T-SLICE for this experiment ($p<0.00-5$ for all gap heights). Hypoxic

reactions were most pronounced with the smallest gap height (300 µm) compared to the larger

gap heights, consistent with our prediction of decreasing oxygen availability with restricted

diffusion as gap height decreases ($p<0.0001$ at all timepoints). Together, our results indicate that

T-SLICE functions to establish cell-driven hypoxic gradients that can be tuned by altering gap

heights.

Cellular proliferation is diminished in the hypoxic regions of T-SLICE

A direct impact of hypoxia in cancer cells is the reduction of proliferation to maintain viability[12].

We next used the EdU assay to assess whether cellular proliferation is diminished in the hypoxic

regions of T-SLICE. After 72h of culture, proliferation of tumor cells within spheroids and

fibroblasts was most present in the peripheral zone of T-SLICE ($p < 0.0001$, **Figure 3 and**

Supplementary Figure 2). Newly divided MCF-7 or HCC1806 cells were not detected in the

middle zone, at any gap height. In MCF7 cells, cell proliferation increased with higher gap

heights ($p < 0.001$), confirming that cellular proliferation and hypoxia are inversely related in T-

SLICE.

Cell viability is reduced in the hypoxic regions of T-SLICE

Concurrent with reduced proliferation, loss of cell viability is commonly observed in hypoxic

regions due to deprivation of essential nutrients and oxygen[12]. To examine cellular viability

within T-SLICE zones, we cultured cells for 72 h and used Calcein-AM and ethidium

homodimer-1 to stain in tandem to identify live (green) and dead (red) cells, respectively **(Figure**

4). Cell viability was near 100% in the peripheral zone of T-SLICE at all gap heights tested. In

MCF-7 spheroids cultured with 300 µm gap height, cell viability dropped to ~50% and ~0% in

the intermediate and middle zones, respectively, but this diminution was nearly completely

abrogated when gap heights were raised to 500 µm and 700 µm. We observed similar cell

viability trends when we instead used ID8 *Trp53-/-* cells in T-SLICE (Supplemental Figure 3).

Hypoxia, slowed proliferation and cell death in T-SLICE is associated with loss of mitochondrial

membrane potential

Loss of mitochondrial trans-membrane potential MMP is a characteristic early feature of

apoptosis and can be brought about by hypoxic stress[13]. MitoView 633 was used to analyze the

mitochondrial membrane potential of cancer cells cultured in T-SLICE for up to 48h **(Figure 5**

and Supplementary Figure 4). In both MCF-7 and ID8 *Trp53-/-* cells, MMP was significantly

diminished in the intermediate and middle zones of T-SLICE compared with the periphery

($p < 0.001$). This impact was significantly greater after 48h ($p < 0.001$). Altogether, our results

indicate that T-SLICE greats physio-chemical gradients of hypoxia that correspond to loss of

cellular transmembrane potential, impacting cellular viability and proliferation.

Discussion

Hypoxia is a characteristic of tumors, whose rapid growth often outpaces by blood vessel

formation[14]. The limitation in oxygen, nutrients, and waste removal force tumor cells to adapt

their metabolic features or undergo apoptosis. These adaptations are associated with increased

metastasis, genomic instability, treatment resistance and immune escape[4,7,12,15]. T-SLICE

represents a simple and effective *in vitro* 3-D system that mimics key features of the TME. By

growing a fibroblast monolayer and cancer cell spheroids between two glass coverslips, T-

SLICE creates physio-chemical gradients in which we observe hypoxic response with

diminutions in cellular proliferation and viability, as well as corresponding loss of MMP moving

from the periphery towards the middle of T-SLICE.

274  In our studies, we worked with three tumor cell lines representing breast and ovarian

cancer. Although the degree of the physio-chemical gradients varied for each, they each

demonstrated hypoxic responses, with spatially correlated cellular proliferation, apoptosis (loss

of mitochondrial transmembrane potential), and viability. Compered to cancer models that rely

on extrinsically imposed oxygen gradients[16,17], a cell-driven gradient generated by metabolic

demand has a better chance of faithfully reflecting the TME of different tumour types and

geometries. Here we did not measure oxygen tension within the T-SLICE to evidence hypoxia,

instead we relied on EF5, cell viability and proliferation markers to detect the cellular hypoxic

response. Trends described, including increasing hypoxic response as well as reduced viability

and cell proliferation moving towards the center of T-SLICE, have been observed in other cell-

driven hypoxic cancer models[18,19].

285  We confirmed that the strength and area of hypoxic zone within T-SLICE can be

increased by reducing the device gap height from 700 to 300 μm. A minimum gap height must

be maintained to accommodate the spheroids (<150 μm diameter), but by increasing the diameter

of T-SLICE this increases the diffusion distance from periphery to middle thereby decreasing the

hypoxic area. This effectively recapitulates TMEs and hypoxia as it would occur in different-

sized tumors or at varying distances from blood vessels. By comparison, in spheroid culture, an

increase in hypoxic effect can only be achieved with more cells and larger spheroid diameters

which will also directly impair the ability to observe cell behaviors within the spheroids. Using

T-SLICE, physio-chemical gradients can be uncoupled from tissue architecture, therefore

making it possible to observe cellular events within a comparable TME from a much larger

spheroid, or a centimeter sized human tumour.

With increasing focus on personalized and precision medicine approaches to cancer

therapy, there is an urgent need to develop faithful and high-throughput systems that can be

efficiently adopted by life science research laboratories around the world. Conventional 2-D

culture and model organisms have been informative for understanding mechanism and disease

processes; however, 2-D systems lack a tumor's structural and physio-chemical characteristics.

Model organisms, including xenografted or humanized mice, are excellent tools to study cancer

in a complex microenvironment, but lack a human immune system, require extensive workup

and monitoring[20], and can be quite heterogeneous. Microfluidic platforms in combination with

3-D culture systems allow a small number of explanted tumor, immune and accessory cells to be

grown and studied, but their complex set-ups make them out of reach for most life science

laboratories[21]. Moreover, endpoint sample recovery can be challenging in sealed microfluidic

devices, thus further impairing their usability as a standard model to study cancer biology. In

contrast, T-SLICE can be easily assembled using components that can be readily made to fit

inside standard cell culture vessels (e.g. 6-well plate), allows high-throughput tumor studies with

continuous monitoring capability in a broadly affordable and feasible way. The transparency of T-SLICE combined with thin cell layers enable brightfield and fluorescent live-cell imaging. By simply lifting the T-SLICE with forceps, selected portions of cells or the entire culture can be harvested for endpoint histologic analysis using standard staining protocol, all without disrupting the overall spatial arrangement of cells.

An advantage that T-SLICE shares with microfluidic culture platforms is the low cell number needed to establish culture and strong paracrine signalling gradients due to the small culture volume. With T-SLICE, fewer than a million cells harvested from patients can be cultured within an environment that mimics the TME of a centimeter sized tumour. Such models using conventional spheroid culture would require several magnitudes more cells and will not be imageable. With a small number of cells, features and interventions can be studied with iteration in adjacent wells of the same plate. Limitations in harvesting sufficient cells to study interventions are circumvented in T-SLICE, making it feasible to routinely incorporate precious patient biopsy samples for therapy screening. Patient avatars could be created to test various strategies and combinations[22,23]. We show that simple adjustments, like changing the gap height, have predictable impacts on the hypoxic gradient; this is relevant because oxygen availability varies even within the same tumor[22,23]. Presently, the hypoxia in T-SLICE is driven primarily by the fibroblast monolayer; however, relative proportions of the tumor, stromal and other cells should be carefully considered and included to reflect specific applications of T-SLICE. Oxygen, nutrients, small molecules, and immune cells will be subject to the challenges created in T-SLICE, just as they are in a tumor. Thus, by designing studies which accurately simulate TMEs within T-SLICE, we can more faithfully capture *in vivo* like drug responses in pre-clinical drug screening, promoting success in the later stages of clinical trials.

Acknowledgements

The authors thank Ms. Sarah Nersesian and Dr. Naeimeh Jafari for technical assistance and

discussions on T-SLICE development. MPT and ND are trainees in the Cancer Research

Training Program of the Beatrice Hunter Cancer Research Institute, with funds for MPT

provided by the Bruce and Dorothy Rosetti Endowment through the Dalhousie Medical Research

Foundation and funds for ND provided by the Helyer/Williams Studentship. MPT is funded by a

CIHR Canadian Graduate Scholarship - Masters and a Nova Scotia Graduate Scholarship. ND is

supported by a Canadian Graduate Scholarship (Masters; NSERC) and a Genomics in Medicine

Scholarship through the Dalhousie Medical Research Foundation. JEB is the Dalhousie Medical

Research Foundation Cameron Cancer Scientist. We acknowledge the support of the

Government of Canada's New Frontiers in Research Fund (NFRF), [NFRFE-2019-00415] and

the OvCAN models consortium for funding this research. We thank Dr. Madhuri Koti (Queen's

University) for the ID8-Trp53$^{-/-}$ cell line.

Author Contributions

T-SLICE model and study was conceptualized by MPT, ND, JEB, BML; experiments were

designed, conducted, and validated by MPT and ND. Data analysis was conducted by MPT, ND,

JEB, BML. The manuscript was written by MPT, ND, JEB, and BML, reviewed, edited, and

approved by all authors.

Declaration of interests

The authors declare that no competing interests exist.

Inclusion and Diversity statement

Dalhousie University is located on the ancestral and unceded territory of the Mi'kmaq peoples.

We strive toward understanding, inclusion and awareness.

References

1 Emon, B., Bauer, J., Jain, Y., Jung, B. & Saif, T. Biophysics of Tumor Microenvironment
and Cancer Metastasis - A Mini Review. *Computational and Structural Biotechnology*
Journal **16**, 279-287, doi:10.1016/j.csbj.2018.07.003 (2018).

2 Balkwill, F. R., Capasso, M. & Hagemann, T. The tumor microenvironment at a glance.
Journal of Cell Science **125**, 5591-5596, doi:10.1242/jcs.116392 (2012).

3 Imparato, G., Urciuolo, F. & Netti, P. A. In vitro three-dimensional models in cancer
research: a review. *International Materials Reviews* **60**, 297-311,
doi:10.1179/1743280415y.0000000003 (2015).

4 Petrova, V., Annicchiarico-Petruzzelli, M., Melino, G. & Amelio, I. The hypoxic tumour
microenvironment. *Oncogenesis* **7**, doi:10.1038/s41389-017-0011-9 (2018).

5 Jing, X. *et al.* Role of hypoxia in cancer therapy by regulating the tumor
microenvironment. *Molecular Cancer* **18**, doi:10.1186/s12943-019-1089-9 (2019).

6 Carmona-Fontaine, C. *et al.* Metabolic origins of spatial organization in the tumor
microenvironment. *Proceedings of the National Academy of Sciences* **114**, 2934-2939,
doi:10.1073/pnas.1700600114 (2017).

7 Abou Khouzam, R. *et al.* Tumor Hypoxia Regulates Immune Escape/Invasion: Influence
on Angiogenesis and Potential Impact of Hypoxic Biomarkers on Cancer Therapies.
Frontiers in Immunology **11**, doi:10.3389/fimmu.2020.613114 (2021).

8 Wong, C. H., Siah, K. W. & Lo, A. W. Estimation of clinical trial success rates and
related parameters. *Biostatistics* **20**, 273-286, doi:10.1093/biostatistics/kxx069 (2019).

9 Spagnolo, F. *et al.* Treatment beyond progression with anti-PD-1/PD-L1 based regimens
in advanced solid tumors: a systematic review. *BMC Cancer* **21**, doi:10.1186/s12885-
021-08165-0 (2021).

10 Walton, J. *et al.* CRISPR/Cas9-Mediated Trp53 and Brca2 Knockout to Generate
Improved Murine Models of Ovarian High-Grade Serous Carcinoma. *Cancer Research*
76, 6118-6129, doi:10.1158/0008-5472.Can-16-1272 (2016).

11 Koch, C. J. in *Redox Cell Biology and Genetics Part A Methods in Enzymology* 3-31
(2002).

12 Bhandari, V. *et al.* Molecular landmarks of tumor hypoxia across cancer types. *Nature*
Genetics **51**, 308-318, doi:10.1038/s41588-018-0318-2 (2019).

13 Zhao, X. *et al.* Hypoxia-Inducible Factor 1-α (HIF-1α) Induces Apoptosis of Human
Uterosacral Ligament Fibroblasts Through the Death Receptor and Mitochondrial
Pathways. *Medical Science Monitor* **24**, 8722-8733, doi:10.12659/msm.913384 (2018).

14 Hanahan, D. Hallmarks of Cancer: New Dimensions. *Cancer Discovery* **12**, 31-46,
doi:10.1158/2159-8290.Cd-21-1059 (2022).

15 Busk, M., Overgaard, J. & Horsman, M. R. Imaging of Tumor Hypoxia for Radiotherapy:
Current Status and Future Directions. *Seminars in Nuclear Medicine* **50**, 562-583,
doi:10.1053/j.semnuclmed.2020.05.003 (2020).

16 Orcheston-Findlay, L., Hashemi, A., Garrill, A. & Nock, V. A microfluidic gradient
generator to simulate the oxygen microenvironment in cancer cell culture.
Microelectronic Engineering **195**, 107-113, doi:10.1016/j.mee.2018.04.011 (2018).

17 Nam, H., Funamoto, K. & Jeon, J. S. Cancer cell migration and cancer drug screening in
oxygen tension gradient chip. *Biomicrofluidics* **14**, doi:10.1063/5.0011216 (2020).

18 Mosadegh, B. *et al.* A paper-based invasion assay: Assessing chemotaxis of cancer cells
in gradients of oxygen. *Biomaterials* **52**, 262-271,
doi:10.1016/j.biomaterials.2015.02.012 (2015).

19 Rodenhizer, D., Dean, T., Xu, B., Cojocari, D. & McGuigan, A. P. A three-dimensional
engineered heterogeneous tumor model for assessing cellular environment and response.
Nature Protocols **13**, 1917-1957, doi:10.1038/s41596-018-0022-9 (2018).

20 Walsh, N. C. *et al.* Humanized Mouse Models of Clinical Disease. *Annual Review of*
Pathology: Mechanisms of Disease **12**, 187-215, doi:10.1146/annurev-pathol-052016-
100332 (2017).

21 Hammel, J. H., Cook, S. R., Belanger, M. C., Munson, J. M. & Pompano, R. R. Modeling
Immunity In Vitro: Slices, Chips, and Engineered Tissues. *Annual Review of Biomedical*
Engineering **23**, 461-491, doi:10.1146/annurev-bioeng-082420-124920 (2021).

22 Yoshida, G. J. Applications of patient-derived tumor xenograft models and tumor
organoids. *Journal of Hematology & Oncology* **13**, doi:10.1186/s13045-019-0829-z
(2020).

23 Voabil, P. *et al.* An ex vivo tumor fragment platform to dissect response to PD-1
blockade in cancer. *Nature Medicine* **27**, 1250-1261, doi:10.1038/s41591-021-01398-3
(2021).
423
424

Figures and Captions

Figure 1: T-SLICE Schematic, Design & Assembly. (A) Top, bottom and side-view

schematic of T-SLICE demonstrating the placement of chips and resulting physio-chemical

gradients. (B) photograph of (i) two-part T-SLICE mold (shown both assembled, top right, and

in component ring and insert pieces (bottom and top left); (ii) top-down and (iii) side view of

assembled T-SLICE. (C) photographs depicting tumor cell line spheroids atop fibroblast

monolayers imaged through a T-SLICE chip: (i) MCF7; (ii) HCC1806; (iii) ID8 *Trp53*[-/-].

Figure 2: T-SLICE generates hypoxic gradients from the periphery to the center. MCF7

spheroids were seeded atop a fibroblast monolayer for 6, 12, 24 or 48h. Thereafter, cells were

fixed and stained with EF5 to ascertain hypoxic responses (A-C). (i) photographs; (ii)

fluorescence quantifications. (D and E) control wells either did not have a T-SLICE installed (D)

or did not include dye (E). P, periphery (blue); M, middle (grey); I, intermediate (red); a.u.,

Arbitrary units. Scale bars represent 275 μm. Data are representative of three technical replicates

with measurements of fluorescence intensity in three fields of view for each zone (p<0.0001,

error bars represent standard deviation (SD)).

Figure 3. Cellular proliferation is halted in the middle region of T-SLICE. MCF-7 cells were

seeded on a fibroblast monolayer in T-SLICE in the presence of EDU. After 72h incubation,

cellular proliferation (EDU incorporation) was ascertained by fluorescence microscopy. (A)

peripheral zone; (B) Middle zone; C) No T-SLICE control; D) unstained control; (E)

quantification of fluorescence. a.u., Arbitrary units. Blue bars represent quantification from the

peripheral zone; red bars represent quantification from the middle zone. Scale bars represent

275μm. Data are representative of three technical replicates with measurements of fluorescence

intensity in three fields of view for each zone (p<0.0001, error bars represent SD).

Figure 4: Cell viability is lost in the most hypoxic regions of T-SLICE. MCF7 spheroids were

cultured in T-SLICE on a fibroblast monolayer for 72h, then stained with Calcein-AM and

ethidium homodimer to visualize cell viability. Images are representative of three technical

replicates and three fields of view for each T-SLICE zone. P, periphery; I, intermediate; M,

middle.

Figure 5. Mitochondrial trans-membrane potential is reduced in T-SLICE. MMP was

measured using MitoView 633 staining after 24 or 48h in MCF7 cells. Significant diminishment

was observed in the intermediate and middle zones compared to the periphery (p<0.001) with

greater impact after 48h (p<0.001). (A-C) each of 300, 500 and 700 μm gap heights were tested.

(i) representative images; and (ii) quantification are shown. (D) controls included unstained cells

in T-SLICE, and stained cells with no T-SLICE. P, periphery (blue); M, middle (grey); I,

intermediate (red); a.u., Arbitrary units. Scale bars represent 275μm. Data are representative of

three technical replicates with measurements of fluorescence intensity in three fields of view for

each zone (p<0.001 after 24h, p<0.0001 after 48h, error bars represent SD).

      Supplemental Information:

Supplementary Figure 1: T-SLICE hypoxic gradients from the periphery to the center in a

second cell line (related to Figure 2). HCC1806 spheroids were seeded atop a fibroblast

monolayer for 6, 12, 24 or 48h. Thereafter, cells were fixed and stained with EF5 to ascertain

hypoxic responses (A-C). (i) photographs; (ii) fluorescence quantifications. (D and E) control

wells either did not contain dye (D) or did not have a T-SLICE chip installed I. P, periphery; M,

middle; I, intermediate. Scale bars represent 275µm. Data are representative of three technical

replicates with measurements of fluorescence intensity in three fields of view for each zone

(p<0.0001, error bars represent SD).

Supplementary Figure 2: Cellular proliferation is slowed in a second cell line model in T-SLICE. HCC1806 cells were seeded on a fibroblast monolayer in T-SLICE in the presence of EDU. After 72h incubation, cellular proliferation (EDU incorporation) was ascertained by fluorescence microscopy. (A) peripheral zone; (B) Middle zone; C) No T-SLICE control; D) unstained control; (D) quantification of fluorescence. a.u., Arbitrary units. Scale bars represent 275μm. Data are representative of three technical replicates with measurements of fluorescence intensity in three fields of view for each zone (p<0.0001, error bars represent SD).

495 **Supplementary Figure 3: Cellular viability is lost in the most hypoxic regions of T-SLICE**

496 **in a second cell line.** ID8-Trp53$^{-/-}$ spheroids were cultured in T-SLICE on a fibroblast

497 monolayer for 72h, then stained with Calcein-AM and ethidium homodimer to visualize cell

498 viability. Images are representative of three technical replicates and three fields of view for each

499 T-SLICE zone. P, periphery; I, intermediate; M, middle.

Supplementary Figure 4. MMP is reduced in T-SLICE in a second cell line. MMP was

measured using MitoView 633 staining after 24 or 48h in ID8-Trp53-/- cells. (A-C) each of 300,

500 and 700 μm gap heights were tested. (i) representative images; and (ii) quantification are

shown. (D) controls included unstained cells in T-SLICE, and stained cells with no T-SLICE. P,

periphery (blue); M, middle (grey); I, intermediate (red); a.u., Arbitrary units. Scale bars

represent 275 μm. Data are representative of three technical replicates with measurements of

fluorescence intensity in three fields of view for each zone (p<0.001 after 24h, p<0.0001 after

48h, error bars represent SD).

510
511