Molecular recognition of two endogenous hormones by the human parathyroid hormone receptor-1

Li-Hua Zhao[1,2]*†, Qingning Yuan[1]*, Antao Dai[1,3]*, Chuan-Wei Chen[4], Xinheng He[1,2], Chao Zhang[5], Youwei Xu[1], Yan Zhou[1,3], Ming-Wei Wang[4,6,7], Dehua Yang[1,2,3]†, H. Eric Xu[1,2]†

[1]The CAS Key Laboratory of Receptor Research, Center for Structure and Function of Drug Targets, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China

[2]University of Chinese Academy of Sciences, Beijing 100049, China

[3]The National Center for Drug Screening, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China.

[4]Research Center for Deepsea Bioresources, Sanya, Hainan 572025, China.

[5]School of Pharmacy, Fudan University, Shanghai 201203, China.

[6]Department of Pharmacology, School of Basic Medical Sciences, Fudan University, Shanghai 200032, China

[7]Department of Chemistry, School of Science, The University of Tokyo, Tokyo 113-0033, Japan.

*These authors contributed equally: Li-Hua Zhao, Qingning Yuan and Antao Dai

†Correspondence: zhaolihuawendy@simm.ac.cn (Li-Hua Zhao); dhyang@simm.ac.cn (Dehua Yang); eric.xu@simm.ac.cn (H. Eric Xu)

ABSTRACT

Parathyroid hormone (PTH) and PTH-related peptide (PTHrP) are two endogenous hormones recognized by PTH receptor -1 (PTH1R), a member of class B G protein couple receptors (GPCRs). Both PTH and PTHrP analogs including teriparatide and abaloparatide, are approved drugs for osteoporosis, but they exhibit distinct pharmacology. Here we report two cryo-EM structures of human PTH1R bound to PTH and PTHrP in the G protein-bound state at resolutions of 2.6 and 3.2 Å, respectively. Detailed analysis of these structures uncovers both common and unique features for the agonism of PTH and PTHrP. Molecular dynamics (MD) simulation and together with site-directed mutagenesis studies reveal the molecular basis of endogenous hormones recognition specificity and selectivity to PTH1R. These results provide a rational template for the clinical use of PTH and PTHrP analogs as an anabolic therapy for osteoporosis and other disorders.

INTRODUCTION

Parathyroid hormone (PTH) and PTH-related protein (PTHrP) are two endogenous peptide hormones that play key and distinct biological roles in skeletal development, calcium- and phosphate- regulating actions, and bone turn over[1,2]. Lack of functional PTH results in hypoparathyroidism[3], while overexpression of PTHrP derived from tumor cells is the most common reason for cancer-associated hypercalcemia[4]. Both PTH and PTHrP also have potent cardiovascular effects, including stimulating heart rate increase independent of autonomic reflexes[5]. Both PTH and PTHrP exert their effects on bone by activating the PTH type 1 receptor (PTH1R), primarily via G_s-cAMP signaling[1,6]. High-resolution structures of PTH1R-G_s complexes bound to these endogenous hormones are important for both pharmacological research as well as clinical development aiming at PTH1R.

Osteoporosis is a disease of decreased bone mass, microarchitectural deterioration, and fragility fractures. It is widespread and can affect all ethnic groups. Osteoporosis becomes a major clinical problem in older women and men, especially in postmenopausal women[7]. Clinical trials were proved that both PTH and PTHrP can stimulate bone anabolism in patients with osteoporosis[6,8]. Teriparatide, recombinant human parathyroid hormone encompassing residues 1-34 (PTH1-34), and abaloparatide, recombinant human PTH-related protein encompassing residues 1-34 (PTHrP1-34), are two synthetic analogs of PTH and PTHrP. Although they exhibit distinct pharmacology, they were the only two approved drugs by FDA for the treatment of osteoporosis[6]. Recent studies suggest that PTH and PTHrP differ in their relative capacities to bind to two pharmacologically distinguishable high-affinity PTH1R conformations, R^0 (the G protein-uncoupled PTH1R conformation) and RG (the G protein-coupled PTH1R conformation)[9,10]. PTH (1-34) can bind to the R^0 conformation with the higher affinity than PTHrP (1-36) and prolonged cAMP responses[10]. Whereas PTHrP (1-36) binds weakly to R^0, but it preferentially binds to the RG, promoted by the over expression of a high affinity of $G\alpha_s$ and produce shorter cAMP responses[6,10].

Hattersley et al.[11] found that abaloparatide has similar high affinity to the RG as

teriparatide, and abaloparatide shows more selectivity for the RG and a shorter cAMP

signaling response compared to teriparatide. Abaloparatide and teriparatide different

selectivity for the RG conformation can explain the different increase in bone mineral

density (BMD) and different incidence of hypercalcemia in the clinical trials treated by

abaloparatide and teriparatide[6]. Such conformational selectivity of receptor binding can

be a key regulator of that ligand's biological activity, which provides an important clue

to optimize conformational selective PTH and PTHrP analogs for a new treatment

option for PTH-related disorders[12].

Up to now, the most effective treatments of osteoporosis are based on daily injections

teriparatide or abaloparatide. These therapies are both costly and inconvenient in

clinical application due to the therapy by injection. However, there are no available

small-molecule agonist drugs for the treatment of osteoporosis, in part due to lack of

detailed structure information of an intact endogenous peptide-PTH1R-G-protein

complex[13,14]. The only available structure of the full-length PTH1R is in complex with

a synthetic long-acting PTH (LA-PTH), which reveals extensive interactions of LA-

PTH with PTH1R[2]. The detailed structural and mechanistic understanding of the

receptor activation by endogenous peptide hormones is critical for explaining how PTH

and PTHrP mediate their distinct biologic actions via a common receptor and for aiding

the rational development of new drug therapies targeting PTH1R. Therefore, we

determined two cryo-electron microscopy (cryo-EM) structures of the active PTH1R–

G_s complexes bound to PTH and PTHrP to provide insights into the pharmacological

mechanisms for the clinical use of endogenous and clinical ligands.

Materials and Methods

Constructs

The human PTH1R (residues 27-502) with G188A and K484R mutations was cloned

into pFastBac vector (Invitrogen) with the haemagglutinin signal peptide (HA)

followed by a TEV protease cleavage site and a double MBP (2MBP) tag to facilitate

expression and purification[2]. To facilitate a stable complex, the above PTH1R construct

was added the LgBiT subunit (Promega) at the C terminus of PTH1R. A bovine

dominant-negative human $G\alpha_s$ (DNGα_s) with mutations G226A and A366S, is a

chimera that the AHD of $G\alpha_s$ (69-205) was replaced with $G\alpha_i$ (62-182) to bind to the

Fab_G50[2]. Based on the published DNGα_s, a modified bovine $G\alpha_s$ (mDNGα_s), its N

terminus (M1–K25) and α-helical domain (AHD, F68–L203) of $G\alpha_s$ were replaced by

the N terminus (M1–M18) and AHD (Y61–K180) of the human $G\alpha_i$, which can bind

scFv16 and Fab_G50[18] and the residues N254-T263 of $G\alpha_s$ were deleted. In addition,

eight mutations (G49D, E50N, L63Y, A249D, S252D, L272D, I372A, and V375I). To

facilitate the folding of the G protein, DNGα_s and mDNGα_s was co-expressed with

GST-Ric-8B[19].   Rat   Gβ1   with   an   N-terminal

MHHHHHHSSGLVPRGSHMASHHHHHHHHHH-tag (His16) was fused with the

SmBiT subunit (peptide 86, Promega)[20] via a 15 amino acid (GSSGGGGSGGGGSSG)

linker at its C terminus. In addition, to clone the constructs into the pcDNA3.1 vector

(Promega) for cAMP accumulation, all constructs were cloned using homologous

recombination (Clone Express One Step Cloning Kit, Vazyme).

Protein expression

The PTH1R and G proteins were co-expressed in *Sf9* insect cells (Invitrogen). When

the cells grew to a density of 3.5×10^6 cells/mL in ESF 921 cell culture medium

(Expression Systems), we infected the cells with five separate virus preparations at a

ratio of 1:2:2:2:2 for PTH1R(27-502)-LgBiT-2MBP, DNGα_s or mDNGα_s, His16-Gβ1-

peptide 86, Gγ2, and GST-Ric-8B. The infected cells were cultured at 27°C for 48 h

before collection by centrifugation and the cell pellets were stored at −80°C.

Expression and purification of Nb35

Nanobody-35 (Nb35) was expressed in *E. coli* BL21 cells, the cultured cells were

grown in 2TB media with 100 μg/mL ampicillin, 2 mM MgCl$_2$, 0.1% glucose at 37°C

for 2.5 h until OD600 of 0.7-1.2 was reached. Then the culture was induced with 1 mM

IPTG at 37°C for 4-5 h, and harvested and frozen at -80°C until use. Nb35 was purified

by nickel affinity chromatography and followed by size-exclusion chromatography

using HiLoad 16/600 Superdex 75 column following overnight dialysis against 20 mM

HEPES, pH 7.4, 100 mM NaCl, 10% glycerol. The quality of purified protein was

verified by SDS-PAGE and store at -80 °C.

Complex purification

The complexes were purified according to previously described methods[21,22]. It was

resuspended in 20 mM HEPES pH 7.4, 100 mM NaCl, 10 mM MgCl$_2$, 10 mM CaCl$_2$,

2 mM MnCl$_2$, 10% glycerol, 0.1 mM TCEP, 15 µg/mL Nanobody (Nb35), 25 mU/mL

apyrase (Sigma), 12 µM PTH1-34 (Synpeptide Co., Ltd) and 12 µM PTHrP1-36

(TGpeptide Biotechnology Co., Ltd), supplemented with Protease Inhibitor Cocktail

(TargetMol, 1 mL/100 mL suspension). The lysate was incubated for 1 h at room

temperature and complex from membranes solubilized by 0.5% (w/v) lauryl maltose

neopentylglycol (LMNG, Anatrace) supplemented with 0.1% (w/v) cholesteryl

hemisuccinate TRIS salt (CHS, Anatrace) for 2 h at 4°C. The supernatant was isolated

by centrifugation at 65,000 × g for 40 min, and the solubilized complex was incubated

with Amylose resin (NEB) for 2 h at 4°C. The resin was loaded onto a plastic gravity

flow column and washed with 15 column volumes of 20 mM HEPES, pH 7.4, 100 mM

NaCl,10% glycerol,10 mM MgCl$_2$, 1 mM MnCl$_2$, 0.01% (w/v) LMNG, 0.01% glyco-

diosgenin (GDN, Anatrace) and 0.004% (w/v) CHS, 2 µM PTH1-34 or 2 µM PTHrP1-

36, and 25 µM TCEP. After washing, the protein was treated overnight with TEV

protease on column at 4°C. Next day the flow through was collected and concentrated,

then UCN1-CRF2R-G$_o$ and UCN1-CRF2R-G$_{11}$ were loaded onto a Superdex200

10/300 GL column and Superose6 Increase 10/300GL (GE Healthcare), respectively,

with the buffer containing 20 mM HEPES, pH 7.4, 100 mM NaCl, 2 mM MgCl$_2$,

0.00075% (w/v) LMNG, 0.00025% GDN, 0.0002% (w/v) CHS, 2 µM PTH1-34 or 2

µM PTHrP1-36, and 100 µM TCEP. The complex fractions were collected and

concentrated individually for electron microscopy experiments.

Cryo-EM data acquisition

For the preparation of cryo-EM grids, 2.5 μL of the purified PTH-PTH1R-G_s and

PTHrP-PTH1R-G_s complexes at a concentration of ~8.4 mg/mL and ~5.1 mg/mL were

respectively applied to the glow-discharged Au 300 mesh holey carbon grids

(Quantifoil R1.2/1.3). The grids were blotted and then plunge-frozen in liquid ethane

using a Vitrobot Mark IV (ThermoFisher Scientific) at 4°C.

Cryo-EM images of PTH-PTH1R-G_s were collected on a Titan Krios G4 equipped with

a Gatan K3 direct electron detector with super-resolution mode and EPU were used to

acquire cryo-EM movies at Advanced Center for Electron Microscopy at Shanghai

Institute of Materia Medica, Chinese Academy of Sciences. A total of 8,002 Movies

were recorded with pixel size of 0.824 Å at a dose of 50 electron per $Å^2$ for 36 frames.

The defocus range of this dataset was -0.8 μm to -1.8 μm. For dimer complex, another

5,054 movies ware obtained with same parameters.

Cryo-EM images of the PTHrP-PTH1R-G_s complex were collected on a Titan Krios

G4 at 300KV accelerating voltage equipped with Gatan K3 direct electron detector. A

total of 7,266 movies were recorded with pixel size of 0.824 Å at a dose of 50 electron

per $Å^2$ for 36 frames. The defocus range of this dataset was -0.8 μm to -1.8 μm.

Image processing

All dose-fractionated image stacks were subjected to beam-induced motion correction

by RELION-4.0[23]. The defocus parameters were estimated by CTFFIND 4.1[24]. PTH-

PTH1R-G_s complex and dimer complex, by auto-picking yielded 3,051,218 particles

and 10,497,057 which were processed by reference-free 2D classification using

Cryosparc[25]. With initial model, after several rounds of 3D classification using

RELION, 407,467 particles and 55,858 particles were used to further refinement and

polishing, yielding two reconstructions with global resolution of 2.62 Å and 3.72 Å

respectively, and subsequently post-processed by DeepEMhancer[26].

For the PTHrP-PTH1R-G_s complex, movies were aligned with RELION-4.0[23]. Initial

contrast transfer function (CTF) fitting was performed with CTFFIND4.1[24] from

Cryosparc[25]. Auto-picking and two-dimensional (2D) Classification were processed

using Cryosparc, producing 7,700,728 particles for further processing. With initial

model, two rounds of 3D classifications were carried out. A further one round of 3D

classification was conducted with a mask on the receptor, in which 436,359 particles

were subjected to 3D auto-refinement and polishing. A map with an indicated global

resolution of 3.25 Å at a Fourier shell correlation (FSC) of 0.143 was generated from

the final 3D refinement, and subsequently post-processed by DeepEMhancer[26].

Model building and refinement

The cryo-EM structure of the LA-PTH1R-G_s-Nb35 complex (PDB code 6NBF) and

two ECD crystal structures (PDB: 3C4M and PDB: 3H3G) were used as the start for

model building and refinement against the electron microscopy map. The model was

docked into the electron microscopy density map using Chimera[27], followed by iterative

manual adjustment and rebuilding in COOT[28]. Real space and Rosetta refinements were

performed using Phenix[29]. The model statistics were validated using MolProbity[30].

Fitting of the refined model to the final map was analyzed using model-versus-map

FSC. To monitor the potential over-fitting in model building, FSC_{work} and FSC_{free} were

determined by refining 'shaken' models against unfiltered half-map-1 and calculating

the FSC of the refined models against unfiltered half-map-1 and half-map-2. The final

refinement statistics are provided in Supplementary Table 2. Structural figures were

prepared in Chimera and PyMOL (https://pymol.org/2/).

Molecular dynamics simulations

The cryo-EM structures of PTH34-PTH1R-G_s complex and PTHrP-PTH1R-G_s

complex were used for the construction of MD simulation systems. Missing ICL3 and short part before TM1 were supplied according to the loop builder program in Molecular Operation Environment. Missing N-terminal and ECL1 structures were constructed according to the AlphaFold2 models[31]. The mutation G188A was modified back and the G proteins were removed. Using CHARMM-GUI, the models were inserted in POPC (palmitoyl-2-oleoyl-sn-glycerol-3-phosphocholine) lipids[32,33]. The length of X and Y were both 130 Å. TIP3P waters and 0.15 M NaCl under a periodic boundary condition were also added to systems. The force fields for proteins and lipids were FF19SB and lipid17, respectively[34-36]. We used hydrogen mass repartitioning during system construction.

After preparation, the systems first encountered a minimization process of 5,000 steepest descent cycles with a constraint on backbone atom, sidechain atom, and lipid coordinates. Then, the constraints were generally decreased in the separated 6 steps of the equilibration process provided by CHARMM-GUI. The first three of them had a timestep of 1 fs and heated the system to 303.15 K, 1 atm in 375 ps. The last three of them had 2 fs timestep and decreased constrain in the coming 1.5 ns. We then performed the independent 400 ns × 6 MD simulations on Amber 20. During simulations, the Langevin thermostat and Berendsen barostat were used for temperature (300 K) and pressure (1 atm) control. Long-range electrostatic interactions were treated by the Particle mesh Ewald and a 10 Å cutoff was employed for short-range interactions. In the following analysis, tICA was calculated in MSMbuilder[37] and the representative structures were extracted by CPPTRAJ[38]. MMGBSA were calculated by Maestro, Schrödinger.

cAMP accumulation assay

PTH and PTHrP stimulated cAMP accumulation was measured by a LANCE Ultra cAMP kit (PerkinElmer). After 24 h culture, the transfected cells were seeded into 384-well microtiter plates at a density of 3,000 cells per well in HBSS supplemented with

5 mM HEPES, 0.1% (w/v) BSA or 0.1% (w/v) casein and 0.5 mM 3-isobutyl-1-

methylxanthine. The cells were stimulated with different concentrations of peptide

agonists for 40 min at RT. Eu-cAMP tracer and ULight^{TM}-anti-cAMP were then diluted

by cAMP detection buffer and added to the plates separately to terminate the reaction.

Plates were incubated at RT for 1 h and the fluorescence intensity measured at 620 nm

and 665 nm by an EnVision multilabel plate reader (PerkinElmer).

Statistical analysis

All functional data were displayed as means ± standard error of the mean (S.E.M.).

Statistical analysis was performed using GraphPad Prism 8.0 (GraphPad Software).

Experimental data were evaluated with a three-parameter logistic equation. The

significance was determined with either two-tailed Student's t-test or one-way ANOVA.

$P < 0.05$ was considered statistically significant.

Data availability

Cryo-EM maps have been deposited in the Electron Microscopy Data Bank under

accession codes: EMD-XXX (PTH-bound PTH1R-G$_s$ complex), EMD- XXX (PTH-

bound PTH1R-G$_s$ complex). The atomic coordinates have been deposited in the Protein

Data Bank under accession codes: XXX (PTH-bound PTH1R receptor) and XXX

(PTHrP-bound PTH1R receptor).

RESULTS

Structural determination

Stable PTH1R-G$_s$ protein complex can be assembled with LA-PTH, but not with PTH

and PTHrP. To overcome this technical obstacle, we employed the NanoBiT tethering

strategy to stabilize the PTH1R-G$_s$ protein complex assembly with PTH and PTHrP

(Supplement Fig.1). The PTH1R was co-expressed with Gα$_s$, Gβ1, Gγ2 and GST-Ric-

8B. Nanobody 35 (Nb35) was added into membranes to stabilizing the receptor-G

protein complex according to the method previously described[2] (Supplement Fig. 1).

The complex structures of the G_s-coupled PTH1R bound to PTH and PTHrP were

determined by cryo-EM to the resolutions of 2.6 Å and 3.2 Å, respectively (Fig. 1 and

Supplementary Fig. 2-3, Supplementary Table 1). In addition, we applied extensive

particle classifications that yielded a distinct subclass of a dimeric form of the PTH-

PTH1R-G_s complexes, which was determined to a resolution of 3.7 Å (Supplementary

Fig. 4). Because this dimeric form is mediated through no-physiological Nb35 and G_s

interface, no further discussion is presented in the paper. Both endogenous peptide-

PTH1R-G_s complexes are similar to other GPCR-G protein complex structures, the α-

helical domain (AHD) of G_s is flexible and invisible. Apart from extracellular domain

(ECD) of receptor, the majority of the amino acid side chains of receptor and

heterotrimeric G_s protein were well resolved in the final models, which are refined

against the EM density map with excellent geometry. Both peptides, PTH (1-34) and

PTHrP (1-36) were clearly identified, thus providing reliable models for the

mechanistic explanation of pharmacological actions of teriparatide and abaloparatide at

PTH1R (Supplement Fig. 5). In addition, like LA-PTH-PTH1R-G_s complex, an ordered

annular lipid band wrapping around the periphery of the receptor transmembrane

domain (TMD) is visible in the cryo-EM map, among which several lipid molecules

were built (Fig. 1a-e). These structural lipids possibly stabilize the PTH1R in its active

state.

Overall structures of G_s-coupled PTH1R bound to PTH and PTHrP

Both PTH-PTH1R-G_s and PTHrP-PTH1R-G_s complex structures closely resemble that

of the LA-PTH-PTH1R-G_s complex with the root mean square deviation (RMSD) value

of 0.5 Å for whole complexes, 0.7 Å for G_s alone, 0.5 Å for receptor alone (Fig. 2).

Notable conformational differences were observed in the position and orientation of the

C-terminal fragment of peptide and the ECD of receptor. The C-terminal fragment of

PTH and PTHrP in complexes are shifted away TM7 by ~9.6 Å and 2.6 Å (measured

at Cα of H^{32P}), respectively, relative to their positions in the LA-PTH-bound structure

(PDB: 6NBF) [2]. The receptor ECD-α1 helix in the PTH- and PTHrP-bound structures are shifted away TM7 by ~10 Å and 7.1 Å (measured at Cα of K50^{ECD}), respectively, relative to their positions in the LA-PTH-bound structure (Fig. 2a-c). These conformational changes allow the C-terminal fragment of hormone bind specificity to PTH1R, indicating PTH1R-associated peptide recognition specificity. Except ECD-α1 helix in PTH-PTH1R cryo-EM structure is shifted away PTH by ~2 Å, relative to X-ray structure (measured at Cα of K50^{ECD}), the overall conformation of the PTH-receptor ECD and PTHrP-receptor ECD in PTH-PTH1R cryo-EM structures resemble the previously published crystal structures of PTH-ECD and PTHrP-ECD complexes (PDB: 3C4M and 3H3G) [9,15], respectively. The structural basis for the binding specificity of PTH and PTHrP to PTH1R is relatively consistent with the crystal structures (Fig. 2d).

Molecular basis for recognition of PTH and PTHrP by PTH1R

Both PTH-PTH1R-G$_s$ and PTHrP-PTH1R-G$_s$ complexes exhibit a similar and unique peptide-receptor binding interface, which is consistent with the "two-domain" model of the peptide-PTH1R interaction mechanism of hormone recognition common to class B GPCRs (Fig. 3). In these complexes, PTH and PTHrP adopt an amphipathic helix with their N terminus dipping into the receptor TMD, while the C terminus is closely interaction with the ECD. The N-terminal fragment of PTH (residues 1-14, PTH^N) and PTHrP (residues 1-15, PTHrP^N) bind to the TMD and activate the receptor. The PTH^N and PTHrP^N adopt four helical turns and insert deeply into the TMD pocket. PTH^N and PTHrP^N are directly surrounded by TM1, TM2, TM3, TM5, TM6 and TM7 and form extensive interactions with the TMD core. PTH^N and PTHrP^N can also interact with ECL2 and ECL3 of receptor. The C-terminal fragment of PTH (residues 15-34, PTH^C) and PTHrP (residues 16-34 PTHrP^C) bind to the ECD of PTH1R to confer high affinity and specificity to the receptor (Fig. 3a, d).

In line with a similar conformation observed between the cryo-EM structures and crystal structures of the PTH1R ECD bound PTH and PTHrP, the extensive interactions

of the ECD with PTH^C and $PTHrP^C$ in cryo-EM structure complexes are similar to those observed in the previously reported X-ray structures of peptide-ECD complexes (Fig. 3b, e). In the PTH-PTH1R-G_s complex, the hydrophobic residues V21, W23, L24, L28, V31 and F34 of the PTH, form an extensive hydrophobic interactions with the hydrophobic groove of the three-layer PTH1R-ECD fold (Fig. 3b)[15]. In the PTHrP-PTH1R-G_s complex, the conserved ECD fold also shows the hydrophobic contacted with F23, L24, L27, I28, I31 and I36 of PTHrP (Fig. 3e)[9]. In addition, $N16^{PTH}$ and $Q16^{PTHrP}$ form a hydrogen bond with $E35^{ECD}$ and $T33^{ECD}$, respectively. The conserved amino acid R20 in the PTH and PTH-related ligands also forms a hydrogen bond with the backbone carbonyls of $M32^{ECD}$ of PTH1R and forms electrostatic interactions with $D137^{ECD}$. These polar interactions display the binding affinity and specificity of PTH and PTHrP to PTH1R. This amino acid R20 is conserved in all three endogenous peptide hormones of PTH1R and PTH2R but distinct from other class B GPCRs, suggesting that R20 is critical for peptide hormones recognition and specificity of PTH and PTH-related ligands[15]. Except for these conserved interaction, these two peptide hormones also interact with PTH1R in a peptide-specific mode. $E19^{PTH}$ forms electrostatic interaction with $K34^{ECD}$, $V21^{PTH}$ makes hydrophobic contact with $D137^{ECD}$ and $W23^{PTH}$ makes Pi interaction with $K34^{ECD}$ and hydrophobic interaction with $Q37^{ECD}$ and $I38^{ECD}$. But different amino acids at cognate positions of PTHrP, $R19^{PTHrP}$ and $R21^{PTHrP}$ form electrostatic interaction with $E35^{ECD}$, $D137^{ECD}$, respectively and $F23^{PTHrP}$ makes hydrophobic interaction with $I38^{ECD}$ and $L41^{ECD}$ of PTH1R (Fig. 3b, e). These interactions receive support from our mutagenesis studies by measuring cAMP responses. Alanine substitution of $D137^{ECD}$ and $Y167^{ECD}$ of PTH1R shows clearly a great reduction in the potency of both PTH and PTHrP-mediated G_s activation (Fig. 3g-h, Supplementary Table 2). In additional, both alanine substitution $M32^{ECD}$ and $E35^{ECD}$ also diminish the potency of PTHrP and slightly decreased the efficacy of PTH. These amino acid residues are important for determining the affinity and specificity of PTH and PTHrP to PTH1R.

The sequences of PTH^N, PTHrP^N and other PTH-related ligands highly resemble. Although PTHrP^N can overlapped well with LA-PTH^N, while PTH^N is different from their positions. Every one of the three N-terminal peptide helix inserts into the receptor TMD core by a analogical angle and orientation, thereby revealing a similar peptide hormones recognition and activation mechanism (Fig. 2c, f and supplementary Fig. 6). Both PTH^N and PTHrP^N run parallel to TM2, making extensive contacts with the residues in TM1, TM2, TM3, TM5, TM6, TM7 and ECL2 and ECL3 (Fig. 3c, f). Especially the first six residues of PTH^N and PTHrP^N play critical roles in the receptor activation. In contrast, PTH (7-34) and PTHrP (7-34) are potent antagonists of PTH receptor[1]. For PTH, A1 forms a hydrogen bond with T427$^{6.59b}$ and makes hydrophobic contacts with Q364$^{5.40b}$, M425$^{6.57b}$ and Q440$^{7.38b}$. For PTHrP, S1 makes more polar interactions with F424$^{6.56b}$, M425$^{6.57b}$, T427$^{6.59b}$ and Q440$^{7.38b}$. These observations provide support for our mutagenesis studies by measuring cAMP responses. Alanine substitutions of Q364$^{5.40b}$, M425$^{6.57b}$ and Q440$^{7.38b}$ notably reduce the potency PTHrP-mediated G$_s$ activation, and slightly effect on the potency of PTH in G$_s$ activation. (Fig. 3i-g, Supplementary Table 2). The same amino acid V2 in two peptides makes extensive hydrophobic contacts with L292$^{3.40b}$, Q364$^{5.40b}$ and I367$^{5.43b}$. Mutation of L292$^{3.40b}$ remarkably reduce pEC$_{50}$ and E$_{max}$ values in the PTH- and PTHrP-induced receptor activity (Fig. 3i-g, Supplementary Table 2). Although S3 in the same position, they exhibit different binding interaction with receptor. S3^{PTH} makes hydrophobic contacts with Q440$^{7.38b}$, M441$^{7.39b}$ and M445$^{7.43b}$. While S3^{PTHrP} forms polar interactions with E444$^{7.42b}$, N448$^{7.46b}$, and also makes hydrophobic interaction with M445$^{7.43b}$. alanine substitutions of M445$^{7.43b}$ indicates a great reduction in the potency of PTHrP-mediated G$_s$ activation (Fig. 3j, Supplementary Table 2), but almost no effect on the receptor activity by PTH in G$_s$ activation (Fig. 3i, Supplementary Table 2). E4 in both PTH and PTHrP, forms the stabilizing hydrogen bond interact and salt bridge with Y195$^{1.47b}$ and R233$^{2.60b}$, respectively. Both alanine mutations in Y195$^{1.47b}$ and R233$^{2.60b}$ remarkably reduce peptide-mediated PTH1R activation, indicating that E4 of the peptide is a key role for peptide-induced receptor activation (Fig. 3i-j, Supplementary Table 2). I5^{PTH}

makes hydrophobic contacts with $L289^{3.37b}$, $L292^{3.40b}$ and $Q364^{5.40b}$. While $H5^{PTHrP}$ at the same position of PTHrP not only makes extensive hydrophobic interactions with $L289^{3.37b}$, $K360^{5.36b}$ and $I363^{5.39b}$, but also forms an additional electrostatic interaction with $Q364^{5.40b}$. Q6 of PTH and PTHrP forms highly conserved hydrophobic interactions with $Y429^{ECL3}$, $W437^{7.35b}$, $Q440^{7.38b}$ and $M441^{7.39b}$. All of these residues mutations, $Y429^{ECL3}A$, $W437^{7.35b}A$, $Q440^{7.38b}A$ and $M441^{7.39b}A$ reduce both PTH and PTHrP to activate PTH1R. In additional, $N10^{PTH}$ is hydrogen-bonded with $W437^{7.35b}$, the same to PTH position, $D10^{PTHrP}$ also forms a hydrogen-bond with $W437^{7.35b}$. $W437^{7.35b}A$, markedly reduce both PTH and PTHrP to activate PTH1R, supporting the fact that $N10^{PTH}$ and $D10^{PTHrP}$ of the peptide are critical for two endogenous hormones-induced receptor activation (Fig. 3i-j, Supplementary Table 2). H9 in both PTH and PTHrP interacts with ECL2 and ECL3 of receptor, such as the hydrophobic interactions with $D353^{ECL2}$, $S355^{ECL2}$, and electrostatic interaction with $Y429^{ECL2}$. $D353^{ECL2}$ significantly decreases the PTHrP-induced PTH1R activation (Fig. 3j, Supplementary Table 2), but almost no effect on the receptor activity by PTH in G_s activation (Fig. 3i, Supplementary Table 2). In additional, $L7^{PTH/PTHrP}$, $M8^{PTH}$ or $L8^{PTHrP}$, $L11^{PTH}$ or $K11^{PTHrP}$, $G12^{PTH/PTHrP}$, $K13^{PTH/PTHrP}$, $H14^{PTH}$ or $K14^{PTHrP}$ can form extensive hydrophobic interactions with the TMD core and ECL2. The above interactions of $peptide^N$ with the TMD provide the major binding and activation energy for the formation of the signal complex. Therefore, the stability of the interface between $peptide^N$ and the TMD is well supported by an extensive network of complementary polar and non-polar interactions between the bound PTH and PTH-related ligands and the TMD. The detailed interactions listed in Fig. 2c, f and supplementary Fig. 6. These interactions help to explain the rich and distinct pharmacology of PTH and PTHrP acting through the same receptor.

Activation mechanism of PTH1R

Comparison of the PTH- and PTHrP-bound PTH1R-G_s complex structures with LA-PTH-bound PTH1R-G_s complex structure reveals a remarkably similarity in the G

protein binding interface, indicating a common mechanism for G protein coupling specificity (Fig. 4). $G\alpha_s$-$\alpha5$ helix dipped into the cytoplasmic receptor cavity and forms important interactions with TM2, TM3, TM5, TM6, H8 and intracellular loop 3(ICL3) (Fig. 4c-d). In all three PTH1R-G_s complex structures, L393 in $G\alpha_s$-$\alpha5$ helix make polar interaction with S409[6.41b]. Both D381 and Q384 of $G\alpha_s$-$\alpha5$ helix form an electrostatic interaction with K388[5.64b]. R385 of $G\alpha_s$-$\alpha5$ also makes an additional electrostatic interaction with E391[ICL3] of PTH1R in PTH-bound PTH1R-G_s complex and PTHrP-bound PTH1R-G_s complex. E392 forms polar interaction with the backbone amine of N363[8.47b] and G364[8.48b]. The bulky residue Y391 at the $G\alpha_s$-$\alpha5$ helix C terminus binds to a sub-pocket formed by R219[2.46b], H233[2.50b], Y305[3.53b] and L306[3.54b] in PTH1R. The hydrophobic residues L388, Y391, L393 and L394 of $G\alpha_s$-$\alpha5$ helix form extensive hydrophobic interaction with TMD helixes (Fig. 4c-d). F314[ICL2] in PTH- and PTHrP-bound PTH1R-G_s complexes insert into a cavity formed by αN-$\beta1$ junction and $\alpha5$ of $G\alpha_s$ and make hydrophobic interactions for stabilizing the interface, but they are different from LA-PTH-bound PTH1R-G_s complex, which F314[ICL2] not directly interact with the hinge region of the $G\alpha_s$ protein (Fig. 4e). Both in PTH- and PTHrP-bound PTH1R-G_s complex, the electrostatic interactions presented between R214[ICL1] and D312 of $G\beta1$, and in PTH- and LA-PTH- bound PTH1R-G_s complex, hydrogen bonds between R476[8.60b] and the backbone oxygen of G310 of $G\beta1$, while in PTHrP-bound PTH1R-G_s complex, hydrogen bonds between R472[8.56b] and the backbone oxygen of G310 of $G\beta1$. Additional contacts between R213[ICL1] and R52 of $G\beta1$ in PTH-bound PTH1R-G_s complex and in LA-PTH-bound PTH1R-G_s complex, all of these interactions may stabilize the interface of helix 8 and $G\beta1$ (Fig. 4f).

Conformational changes during PTH1R activation

PTH1R has two different conformation states, the G-protein free state R[0] and the G protein coupled state RG[10]. During peptide-bound PTH1R activation, two PTH1R conformations undergo different rearrangement of the network residues to facilitate the ligand binding and signal transduction[14]. Based on structural alignment of PTH1R:

agonist complex in the absence of G protein, with three G protein-bound PTH1R (RG):
agonist complexes shows different conformational changes, involving the ECD and the
TMD of the receptor (Fig. 5a-c). The receptor ECD-α1 in the PTH-bound and PTHrP
bound PTH1R RG structures are shifted away TM7 by ~4.4 Å and 3.5 Å (measured at
Cα of K50^{ECD}), respectively, relative to their positions in the ePTH-bound PTH1R
structure. The main conformational changes happens in TMD (Fig. 5b-c). The class B
GPCRs conserved PxxG motif (P415$^{6.48b}$, L416$^{6.48b}$, F417$^{6.49b}$ and G418$^{6.50b}$) in three
agonist-G-protein-bound PTH1R complexes caused a notable conformational
rearrangement and make TM6 kink outward ~90° in the middle (Fig. 5d). The TM6
sharp kink creating an intracellular binding pocket for the C terminus of Gα, thereby
enabling intracellular G protein activation and signal transduction. Meanwhile, the kink
of TM6 together with other conformational changes caused three class B GPCRs
conserved polar interaction networks rearrangement, including HETY network
(H223$^{2.50b}$, E302$^{3.50b}$, T410$^{6.42b}$, and Y457$^{7.57b}$), the II-VI-VII-VIII network (R219$^{2.46b}$,
K405$^{6.37b}$, N463$^{7.61b}$ and E465$^{8.49b}$) and central polar network (R233$^{2.60b}$, N295$^{3.43b}$,
H420$^{6.52b}$ and Q451$^{7.49b}$). However, this structural hallmark of class B GPCRs activation
was not observed in the ePTH-PTH1R complex structure and CLR-CGRPR-RAMP1
complex structure, which are in the absence of G protein. Their intracellular part of the
receptors still displays the hallmarks of an inactive conformation (Fig. 5b and
supplementary Fig. 7), which are called intermediate states on the transition of the
receptor toward the G protein-bound state[13,14,16]. The intermediate states of class B
GPCRs and the R^0 conformation of the PTH1R resemble in agonist binding and in the
absence of G protein. The ability of PTH and PTHrP to bind to RG is similar, but PTH
has a greater capacity to bind to R^0 state[1,10]. Such conformational selectivity mediate
their distinct biological actions[10]. From molecular dynamics (MD) simulations, the
lower R^0 binding ability of PTHrP was attributed to its higher binding free energy upon
specific states (Supplementary Fig. 8). Although the capacity of PTH and PTHrP to
bind to R^0 is different, these peptides can bind to the G protein-coupled PTH1R
complex (RG) with a highly similar G protein binding interface, suggesting a common

mechanism for G_s engagement.

DISCUSSION

The cryo-EM structures of PTH-PTH1R-G_s and PTHrP-PTH1R-G_s complexes further

reveal the structural basis for recognition of PTH and PTHrP by PTH1R and the

receptor activation and G protein-coupling mechanisms, providing an opportunity to

use these structures as modules for designing new PTH replacement therapies in the

future. Two synthetic analogs of PTH and PTHrP, teriparatide and abaloparatide were

approved drugs by FDA and used in clinic, but as conventional therapy for osteoporosis

by daily injections with high costs, probably not suitable for treating such a widespread

disease. Therefore, PTH1R-specific targeted orally agonists for osteoporosis are

urgently developed. To our knowledge, PTH and PTHrP differ in selectivity for two

pharmacologically distinguishable high-affinity PTH1R conformations, R^0 and RG[1].

Such R^0-selective analogs, including PTH, M-PTH (1-28) and M-PTH (1-34) display

high affinity binding to R^0 as well as notably mediate prolonged cAMP signaling

responses and are now being investigated as potential new drugs for treating

hypoparathyroidism[17]. While another PTH analogs, such as PTHrP (1-36) and M-PTH

(1-14) exhibit a high- affinity to bind to RG conformation, and produces a shorter-lived

and more pulsatile action at the receptor, might prove to be more efficacious in this

class of patients[1,12]. Mutational studies of the ligand-receptor binding sites indicate that

the binding specificity of PTH and PTHrP to PTH1R. According to our structures,

detailed patterns of teriparatide and abaloparatide-receptor interactions can be

identified and it can explain the binding specificity of PTH and PTHrP to PTH1R. Thus,

a detailed understanding of PTH and PTHrP binding and activation of PTH1R provide

a rational template to development of optimized therapeutics targeting PTH1R diseases.

REFERENCES

1 Gardella, T. J. & Vilardaga, J. P. International Union of Basic and Clinical Pharmacology.
XCIII. The parathyroid hormone receptors--family B G protein-coupled receptors.
Pharmacol Rev **67**, 310-337, doi:10.1124/pr.114.009464 (2015).
2 Zhao, L. H. *et al.* Structure and dynamics of the active human parathyroid hormone

508   receptor-1. *Science* **364**, 148-153, doi:10.1126/science.aav7942 (2019).

509   3   Nishimura, Y. *et al.* Lead Optimization and Avoidance of Reactive Metabolite Leading to
510   PCO371, a Potent, Selective, and Orally Available Human Parathyroid Hormone Receptor
511   1 (hPTHR1) Agonist. *J Med Chem* **63**, 5089-5099, doi:10.1021/acs.jmedchem.9b01743
512   (2020).

513   4   Kir, S. *et al.* Tumour-derived PTH-related protein triggers adipose tissue browning and
514   cancer cachexia. *Nature* **513**, 100-104, doi:10.1038/nature13528 (2014).

515   5   Wider, J., Undyala, V. V. R., Lanske, B., Datta, N. S. & Przyklenk, K. Parathyroid Hormone-
516   Related Peptide and Its Analog, Abaloparatide, Attenuate Lethal Myocardial Ischemia-
517   Reperfusion Injury. *J Clin Med* **11**, doi:10.3390/jcm11092273 (2022).

518   6   Chew, C. K. & Clarke, B. L. Abaloparatide: Recombinant human PTHrP (1-34) anabolic
519   therapy for osteoporosis. *Maturitas* **97**, 53-60, doi:10.1016/j.maturitas.2016.12.003 (2017).

520   7   Srivastava, M. & Deal, C. Osteoporosis in elderly: prevention and treatment. *Clin Geriatr*
521   *Med* **18**, 529-555, doi:10.1016/s0749-0690(02)00022-8 (2002).

522   8   Bilezikian, J. P., Rubin, M. R. & Finkelstein, J. S. Parathyroid hormone as an anabolic therapy
523   for women and men. *J Endocrinol Invest* **28**, 41-49 (2005).

524   9   Pioszak, A. A., Parker, N. R., Gardella, T. J. & Xu, H. E. Structural basis for parathyroid
525   hormone-related protein binding to the parathyroid hormone receptor and design of
526   conformation-selective peptides. *The Journal of biological chemistry* **284**, 28382-28391,
527   doi:10.1074/jbc.M109.022905 (2009).

528   10   Dean, T., Vilardaga, J. P., Potts, J. T., Jr. & Gardella, T. J. Altered selectivity of parathyroid
529   hormone (PTH) and PTH-related protein (PTHrP) for distinct conformations of the
530   PTH/PTHrP receptor. *Mol Endocrinol* **22**, 156-166, doi:10.1210/me.2007-0274 (2008).

531   11   Hattersley, G., Dean, T., Corbin, B. A., Bahar, H. & Gardella, T. J. Binding Selectivity of
532   Abaloparatide for PTH-Type-1-Receptor Conformations and Effects on Downstream
533   Signaling. *Endocrinology* **157**, 141-149, doi:10.1210/en.2015-1726 (2016).

534   12   Okazaki, M. *et al.* Prolonged signaling at the parathyroid hormone receptor by peptide
535   ligands targeted to a specific receptor conformation. *Proc Natl Acad Sci U S A* **105**, 16525-
536   16530, doi:10.1073/pnas.0808750105 (2008).

537   13   Ehrenmann, J. *et al.* High-resolution crystal structure of parathyroid hormone 1 receptor
538   in complex with a peptide agonist. *Nat Struct Mol Biol* **25**, 1086-1092,
539   doi:10.1038/s41594-018-0151-4 (2018).

540   14   Ehrenmann, J., Schoppe, J., Klenk, C. & Pluckthun, A. New views into class B GPCRs from
541   the crystal structure of PTH1R. *Febs Journal* **286**, 4852-4860, doi:10.1111/febs.15115
542   (2019).

543   15   Pioszak, A. A. & Xu, H. E. Molecular recognition of parathyroid hormone by its G protein-
544   coupled receptor. *Proc Natl Acad Sci U S A* **105**, 5034-5039,
545   doi:10.1073/pnas.0801027105 (2008).

546   16   Josephs, T. M. *et al.* Structure and dynamics of the CGRP receptor in apo and peptide-
547   bound forms. *Science* **372**, doi:10.1126/science.abf7258 (2021).

548   17   Mannstadt, M. *et al.* Efficacy and safety of recombinant human parathyroid hormone (1-
549   84) in hypoparathyroidism (REPLACE): a double-blind, placebo-controlled, randomised,
550   phase 3 study. *Lancet Diabetes Endocrinol* **1**, 275-283, doi:10.1016/S2213-
551   8587(13)70106-2 (2013).

18   Maeda, S. *et al.* Development of an antibody fragment that stabilizes GPCR/G-protein
complexes. *Nat Commun* **9**, 3712, doi:10.1038/s41467-018-06002-w (2018).

19   Chan, P. *et al.* Purification of heterotrimeric G protein alpha subunits by GST-Ric-8
association: primary characterization of purified G alpha(olf). *The Journal of biological*
chemistry **286**, 2625-2635, doi:10.1074/jbc.M110.178897 (2011).

20   Dixon, A. S. *et al.* NanoLuc Complementation Reporter Optimized for Accurate
Measurement of Protein Interactions in Cells. *ACS Chem Biol* **11**, 400-408,
doi:10.1021/acschembio.5b00753 (2016).

21   Zhao, L.-H. *et al.* Structure and dynamics of the active human parathyroid hormone
receptor-1. *Science* **364**, 148-153, doi:10.1126/science.aav7942 (2019).

22   Ma, S. *et al.* Molecular Basis for Hormone Recognition and Activation of Corticotropin-
Releasing Factor Receptors. *Mol Cell* **77**, 669-680 e664, doi:10.1016/j.molcel.2020.01.013
(2020).

23   Zivanov, J., Nakane, T. & Scheres, S. H. W. Estimation of high-order aberrations and
anisotropic magnification from cryo-EM data sets in RELION-3.1. *Iucrj* **7**, 253-267,
doi:10.1107/S2052252520000081 (2020).

24   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J Struct Biol* **192**, 216-221, doi:10.1016/j.jsb.2015.08.008 (2015).

25   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
doi:10.1038/nmeth.4169 (2017).

26   Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume
post-processing. *Commun Biol* **4**, 874, doi:10.1038/s42003-021-02399-1 (2021).

27   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

28   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*
Crystallogr D Biol Crystallogr **60**, 2126-2132, doi:10.1107/S0907444904019158 (2004).

29   Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221,
doi:10.1107/S0907444909052925 (2010).

30   Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D* **66**, 12-21, doi:10.1107/S0907444909042073 (2010).

31   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589, doi:10.1038/s41586-021-03819-2 (2021).

32   Jo, S. *et al.* CHARMM-GUI 10 years for biomolecular modeling and simulation. *Journal of*
computational chemistry **38**, 1114-1124, doi:10.1002/jcc.24660 (2017).

33   Lee, J. *et al.* CHARMM-GUI supports the Amber force fields. *The Journal of chemical*
physics **153**, 35103, doi:10.1063/5.0012280 (2020).

34   Tian, C. *et al.* ff19SB: Amino-Acid-Specific Protein Backbone Parameters Trained against
Quantum Mechanics Energy Surfaces in Solution. *J Chem Theory Comput* **16**, 528-552,
doi:10.1021/acs.jctc.9b00591 (2020).

35   Lee, J. *et al.* CHARMM-GUI supports the Amber force fields. *J Chem Phys* **153**, 035103,
doi:10.1063/5.0012280 (2020).

36   He, X., Man, V. H., Yang, W., Lee, T. S. & Wang, J. A fast and high-quality charge model

for the next generation general AMBER force field. *J Chem Phys* **153**, 114502,
doi:10.1063/5.0019056 (2020).
37 Roe, D. R. & Cheatham III, T. E. PTRAJ and CPPTRAJ: software for processing and analysis
of molecular dynamics trajectory data. *Journal of chemical theory and computation* **9**,
3084-3095 (2013).
38 Harrigan, M. P. *et al.* MSMBuilder: Statistical Models for Biomolecular Dynamics.
Biophysical journal **112**, 10-15 (2017).
603

ACKNOWLEDGMENTS

The cryo-EM data were collected at Advanced Center for Electron Microscopy at Shanghai Institute of Materia Medica, Chinese Academy of Sciences. We are grateful to W.H. and K.W. for collecting the cryo-EM data. This work was supported by National Natural Science Foundation of China (32071203 to L.H.Z.; 82073904 to M.W.W. and 81973373 to D.H.Y.), the National Key R&D Program of China (2019YFA0904200), the Young Innovator Association of CAS (2018325 to L.H.Z.) and SA-SIBS Scholarship Program to L.H.Z. and D.H.Y.; Ministry of Science and Technology (China) grants (2018YFA0507002 to H.E.X. and 2018YFA0507000 to M.W.W.), the Shanghai Municipal Science and Technology Major Project (2019SHZDZX02 to H.E.X.;18ZR1447800 to L.H.Z and 21JC1401600 to D.H.Y.), the CAS Strategic Priority Research Program (XDB08020303 to H.E.X.).

AUTHOR CONTRIBUTIONS

L.H.Z. designed the expression constructs, purified the complexes, prepared the final samples for negative stain and data collection toward the structures, participated in model building and performed structure and function data analysis, prepared figures and wrote the manuscript; L.H.Z. prepared the cryo-EM grids and Q.N.Y. collected cryo-EM images and performed map calculations, built and refined the structure models; Y.W.X. participated in model refinement; X.H.H conducted MD simulations; A.T.D., C.W.C., C.Z. and Y.Z. performed signaling experiments under the supervision of D.H.Y. and M.W.W.; L.H.Z. and H.E.X. conceived the project, wrote the manuscript.

ADDITIONAL INFORMATION
Competing interests: The authors declare that they have no competing interests.

Fig. 1 The overall cryo-EM structures of G$_s$-coupled PTH1R bound to PTH and PTHrP. (a) Schematic diagram of two endogenous hormones, PTH (salmon) and PTHrP (steel blue) are recognized by PTH1R (light sea green) and exert their effects on bone by activating PTH1R, primarily via G$_s$-cAMP signaling. G$_s$ is colored in medium purple, Gβ in light blue and Gγ in rosy brown. **(b)** Orthogonal views of the density map and **(c)** model for PTH-PTH1R-G$_s$ complex. The colored cryo-EM density map is shown at the 0.0183 threshold and light gray surface indicates a micelle diameter of 10 nm. Cadet blue, PTH1R; salmon, PTH; medium purple, G$_s$; light blue, Gβ; rosy brown, Gγ; peru, Nb35; goldenrod, CLR; orchid, PLM. **(d)** Orthogonal views of the density map and **(e)** model for PTHrP-PTH1R-G$_s$ complex. The colored cryo-EM density map is shown at the 0.136 threshold and light gray surface indicates a micelle diameter of 11 nm. Slate blue, PTH1R; steel blue, PTHrP; medium purple, G$_s$; light blue, Gβ; rosy brown, Gγ; peru, Nb35; goldenrod, CLR; orchid, PLM.

Fig. 2 Structure comparisons of PTH1R-G$_s$ complexes by different ligands. (a-c)

Structural comparison of PTH-PTH1R-G$_s$ (salmon, PTH; cadet blue, PTH1R; medium

purple, G$_s$; light blue, Gβ; rosy brown, Gγ), PTHrP-PTH1R-G$_s$ (steel blue, PTHrP; slate

blue, PTH1R; purple, G$_s$; dark khaki, Gβ; yellow green, Gγ) and LA-PTH-PTH1R-G$_s$

(PDB: 6NBF, sandy brown, LA-PTH; medium violet red, PTH1R; dark slate blue, G$_s$;

goldenrod, Gβ; dark green, Gγ) complexes. Notable conformational changes observed

in the receptor ECD and the ligand **(a-b)** in the side view and **(c)** cytoplasmic and

extracellular view. **(d)** Comparison of the PTH bound ECD in PTH-PTH1R-G$_s$ EM

structure with PTH-bound ECD crystal structure (PDB: 3C4M) and PTHrP bound ECD

in PTHrP-PTH1R-G$_s$ EM structure with PTHrP-bound ECD crystal structure (PDB:

3H3G), yellow, PTHrP; lime green, ECD; sea green, PTH; pale violet red, ECD.

Fig. 3 Molecular recognition of PTH1-34 and PTHrP1-36 by PTH1R. (a) An overall

PTH-binding pocket of PTH1R. Cutaway view showing that PTH$^{N\alpha}$ penetrates into a

pocket formed by TMD, while its C-terminal half is recognized by the ECD. **(b-c)**

Detailed interactions of PTH^{C} with the PTH1R ECD and interactions of PTH$^{N\alpha}$ with

the PTH1R TMD pocket, salmon, PTH; cadet blue, PTH1R. The polar contacts are

shown as purple dotted lines. **(d)** An overall PTHrP-binding pocket of PTH1R. Cutaway view showing that PTHrP$^{N\alpha}$ penetrates into a pocket formed by TMD, while its C-terminal half is recognized by the ECD. **(e-f)** Detailed interactions of PTHrP^{C} with the PTH1R ECD and interactions of PTHrP$^{N\alpha}$ with the PTH1R TMD pocket, steel blue, PTHrP; slate blue, PTH1R. The polar contacts are shown as purple dotted lines. **(g-j)** Effect of receptor mutation on PTH and PTHrP-induced cAMP accumulation. **(g-h)** Signaling profiles of PTH1R ECD mutants on PTH and PTHrP-induced cAMP accumulation. **(i-j)** Signaling profiles of PTH1R TMD, ECL2 and ECL3 mutants on PTH and PTHrP-induced cAMP accumulation. cAMP accumulation was measured in wild-type (WT) and single-point mutated PTH1R expressing in HEK293T cells, respectively. cAMP accumulation was normalized to the maximum response of the WT and dose-response curves were analyzed using a three-parameter logistic equation. Data were generated and graphed as means ± S.E.M. of three independent experiments (n = 3) performed in triplicate.

Fig. 4 G protein coupling of multi-agonist-bound PTH1R. (a) Comparison of G

protein coupling among PTH, PTHrP and LA-PTH bound PTH1R. The receptors and

G proteins are colored as the labels. **(b)** The Gα$_s$-α5 helix of the Gα$_s$ inserts into an

intracellular cavity of receptor's TMD. **(c-f)** Comparison of the PTH1R-G$_s$ interface, including **(c)** the Gα$_s$-α5 C terminus-TMD interface, **(d)** Gα$_s$-α5-TM3/TM5 interface, **(e)** ICL2-Gα$_s$-αN and Gα$_s$-α5 helix interface, **(f)** and helix 8-Gβ1 interface. cadet blue, PTH1R; medium purple, G$_s$; light blue, Gβ; rosy brown, Gγ in PTH-PTH1R-G$_s$ complex, slate blue, PTH1R; purple, G$_s$; dark khaki, Gβ; yellow green, Gγ in PTHrP-PTH1R-G$_s$ complex and medium violet red, PTH1R; dark slate blue, G$_s$; goldenrod, Gβ; dark green, Gγ in LA-PTH-PTH1R-G$_s$ complex The polar contacts are shown as purple dotted lines.

Fig. 5 Structures of the active PTH1R in G protein-bound state and comparison with structure of the active PTH1R in the absence of G protein. (a-c) The structural alignment of ePTH-bound PTH1R (PDB: 6FJ3, lime, ePTH; dark goldenrod, PTH1R) with PTH-bound PTH1R (salmon, PTH; cadet blue, PTH1R), PTHrP-bound PTH1R

(steel blue, PTHrP; slate blue, PTH1R) and LA-PTH-bound PTH1R (sandy brown, LA-

PTH; medium violet red, PTH1R), which are in G_s bound state, showing different

conformational changes of the receptor between PTH1R in G_s bound and no G_s bound

states **(a)** in the side view, **(b)** extracellular view and **(c)** intracellular view, the red

arrows indicate conformational changes. **(d)** The structural alignment of PTH1R in the

absence of G protein and in G protein-bound state, showing the outward bending of the

intracellular portion of TM6 of activated PTH1R in G_s-bound state, which results in a

kink at the PxxG motif in TM6 and a ~90° angle in the middle of TM6 of the activated

PTH1R in G protein-bound state. The TM6 kink in the active PTH1R structure is

indicated by a black line. The conserved PxxG motif in TM6 are shown in stick

representation. **(e)** Conformational changes during receptor activation in the absence of

G protein and in G protein-bound state in HETY network, **(f)** in the II-VI-VII-VIII

network, **(g)** and in central polar network. The residues are shown in stick

representation and conformational changes are indicated by red arrows.