TITLE: The effects of biological sex on estimates of persistent inward currents in the human lower limb

Authors: Sophia T. Jenz[1,2], James A. Beauchamp[1,3], Matheus M. Gomes [4], Francesco Negro[5], CJ Heckman[1,2,6,7], and Gregory E.P. Pearcey[1,8,9]

[1]Department of Physical Therapy and Human Movement Sciences, Feinberg School of Medicine, Northwestern University, Chicago, USA.

[2]Department of Neuroscience, Feinberg School of Medicine, Northwestern University, Chicago, USA.

[3]Department of Biomedical Engineering, McCormick School of Engineering, Northwestern University, Evanston, USA.

[4]School of Physical Education and Sport of Ribeirão Preto, University of São Paulo, Ribeirão Preto, Brazil.

[5]Department of Clinical and Experimental Sciences, Universita degli Studi di Brescia, Brescia, Italy.

[6]Department of Physical Medicine and Rehabilitation, Feinberg School of Medicine, Northwestern University, Chicago, USA.

[7]Shirley Ryan AbilityLab, Chicago, USA.

[8]School of Human Kinetics and Recreation, Memorial University of Newfoundland, St. John's, Canada.

[9]Division of BioMedical Sciences, Faculty of Medicine, Memorial University of Newfoundland, St. John's, Canada.

Running title (70 characters): Visual feedback and biological sex affect PIC magnitude

Address for correspondence:

Gregory Pearcey

Email: gpearcey@mun.ca

ABSTRACT (250 words)

Non-invasive recordings of motor unit (MU) spike trains help us understand how the nervous
system controls movement and how it adapts to various physiological conditions. The majority of
study participants in human and non-human animal physiology studies are male, and it is
assumed mechanisms uncovered in these studies are shared between males and females.
However, sex differences in neurological impairment and physical performance warrant the study
of sex as a biological variable in human physiology and performance. To begin addressing this
gap in the study of biophysical properties of human motoneurons, we quantified MU discharge
rates and estimates of persistent inward current (PIC) magnitude in both sexes by quantifying ΔF.
We decomposed MU spike trains from the tibialis anterior (TA), medial gastrocnemius (MG), and
soleus (SOL) using high-density surface electromyography and blind source separation
algorithms. Ten participants of each sex performed slow triangular (10s up and down) isometric
contractions to a peak of 30% of their maximum voluntary contraction. We then used linear mixed
effects models to determine if peak discharge rate and ΔF were predicted by the fixed effects of
sex, muscle, and their interaction. Despite a lack of significant sex-differences in peak discharge
rates across all muscles, ΔF was larger ($\chi^2_{(1)}$ = 6.26, p = 0.012) in females (4.73 ± 0.242 pps) than
males (3.81 ± 0.240 pps). These findings suggest that neuromodulatory drive, inhibitory input,
and/or biophysical properties of motoneurons differ between the sexes and may contribute to
differences in MU discharge patterns.

Keywords (3-5): Hysteresis, motor unit, sex differences, motoneuron

KEY POINTS

- *Sex differences in motor unit studies have been revealed with greater inclusion of female participants, however, mechanisms for these differences remain unclear.*

- *Estimates of persistent inward currents (i.e., ΔF) were greater in females than males in the tibialis anterior, medial gastrocnemius, and soleus muscles.*

- *This suggests that neuromodulatory drive, monoaminergic signaling, or descending control may differ between the sexes.*

- *Therefore, sex differences in estimates of PICs may provide a mechanism behind previously reported sex differences in motoneuron discharge patterns..*

INTRODUCTION

Historically, studies in physiology, and in particular neurophysiology, have excluded female participants. Woitowich and colleagues (2020) recently surveyed the literature and showed that females currently represent only 15 percent of both non-human subjects and human participants in biological studies. This has resulted in an extensive understanding of the neural control of movement in males, but a poor understanding of what mechanisms underlie differences in motor control observed in females (Jenz & Pearcey, 2022).

Regardless of biological sex, all movement is achieved through the activation of motoneurons and the muscle fibers they innervate, together known as motor units (MUs; Heckman & Enoka, 2012). Under normal conditions, motoneurons have a one-to-one spike ratio with the fibers they innervate, and therefore the discharge patterns of motoneurons and their MU are the same (Johnson *et al.*, 2017; Thompson *et al.*, 2018). This creates a unique opportunity to study the central nervous system, as motoneurons are the only neuron that can be recorded non-invasively in humans. Thus, the non-invasive assessment of MU behavior can reveal important information on how the cortex, spinal cord, and individual neurons communicate and whether sexual dimorphisms exist within these neural commands.

A contemporary understanding of the physiology underlying motor control suggests that motoneurons receive a complex combination of excitatory, inhibitory, and neuromodulatory input (Heckman et al. 2009). This neuromodulatory input involves two important monoamines (among others), serotonin (5HT) and norepinephrine (NE), that are released via long descending projections from the brainstem nuclei (raphe nucleus and locus coeruleus, respectively) onto motoneurons (Montague *et al.*, 2013). Release of 5HT and NE facilitates voltage-sensitive persistent inward currents (PICs; Hounsgaard *et al.*, 1988; Heckman *et al.*, 2005) and contributes to the gain control required for humans to perform an impressive variety of movements, from long term postural control during waking for hours, to precise targeted movements like reaching to grasp a pint in a crowded pub (Heckman *et al.*, 2008).

PICs modulate motoneuron discharge in a non-linear fashion by creating a strong initial acceleration and subsequent attenuation in discharge rate during linear increases in synaptic input (Binder *et al.*, 2020). The most robust hallmark of PICs in humans, however, manifests as discharge rate hysteresis between the recruitment and de-recruitment of MUs (Khurram *et al.*, 2022). Typically, PICs are estimated in humans using a paired MU analysis technique to quantify discharge rate hysteresis (Gorassini *et al.*, 1998, 2002*a*, 2002*b*). Although many labs have utilized

this technique with MU spike trains obtained while participants perform voluntary triangular

shaped contractions, the role of descending monoaminergic (i.e., 5HT and NE) input from the

brainstem is broadly understudied in the motor system and has not been compared between the

sexes.

In recent years, studies that have included an adequate number of females have

demonstrated sex differences in several MU discharge properties, but the results vary across

different tasks and conditions. The consensus is that MU discharge (English & Widmer, 2003;

Harwood *et al.*, 2014; Peng *et al.*, 2018; Parra *et al.*, 2020; Inglis & Gabriel, 2020; Nishikawa *et*

al., 2022) and neuromuscular fatigue (Hunter & Enoka, 2001; Celichowski & Drzymała, 2006;

Ansdell *et al.*, 2019; Inglis & Gabriel, 2021; Augsburger *et al.,* 2022) differ between the sexes

(Lulic-Kuryllo & Inglis, 2022). Recent data suggest that females use different recruitment

strategies than males at relatively low efforts due to smaller MU size and higher discharge rates

(Guo *et al.*, 2022) but the specific mechanisms underlying these differences remain unknown.

In non-human animal models, sex differences in MU physiology have been identified at

the cellular level. For example, male rats have more fast-fatigable type MUs in the gastrocnemius

muscle, while female rats have more fatigue-resistant MUs (Celichowski & Drzymała, 2006), with

male rats having a greater total number of MUs than females (Celichowski & Drzymała-

Celichowska, 2007). Most relevant to the current investigation, Kirkwood and colleagues

(Kirkwood *et al.*, 2002; Ford & Kirkwood, 2006) have shown that plateau potentials, which are

dependent on Ca^{2+} PICs and typically absent in anesthetized preparations, have been observed

only in anesthetized estrous *female* cats, suggesting that biophysical motoneuronal properties

differ between the sexes.

Therefore, to understand the effect of biological sex on motoneuronal properties in

humans, we measured MU behavior during slow isometric ramps contractions to the same relative

intensity and compared them between females and males. We hypothesized that females would

have greater PICs, which would manifest as higher peak discharge rates and more discharge rate

hysteresis – an indication of greater PIC magnitudes.

METHODS

Participants and ethical approval

Young and healthy females (n = 10; 25.5 ± 4.8 years) and males (n = 10; 27.2 ± 5.8 years) were recruited for this study. Participants had no history of muscular or neurological impairment and provided written informed consent, approved by the Institutional Review Board of Northwestern University (STU00216360).

Experimental Apparatus

Torque: During each session, participants were seated in a Biodex chair (Biodex Medical Systems, Shirley, NY) and their right foot was placed in a footplate attachment with an ankle angle of 95 degrees, a hip flexion angle of 100 degrees, and knee fully extended. The footplate was attached to a Systems 2 Dyanomometer (Biodex Medical Systems, Shirley, NY) and Velcro straps were wrapped around the foot to secure it to the plate and prevent movement. Torque was sampled at 2048 Hz and smoothed offline with a 10 Hz low-pass filter (fifth-order Butterworth filter).

High-density surface electromyography (HDsEMG): Before electrode placement, excess hair was removed and the skin overlying the muscle of interest was lightly abraded. High-density surface EMG (HDsEMG) electrodes (64 electrodes, 13x5, 8mm I.E.D., GR08MM1305, OT Bioelettronica, Inc., Turin, IT) were placed over the tibialis anterior (TA), and the medial gastrocnemius (MG) and soleus (SOL), which are dorsiflexor and plantarflexor muscles of the ankle, respectively. A reference electrode strap was placed around the right ankle, overlying the medial and lateral malleolus (Figure 1). HDsEMG data was recorded in single differential mode, sampled at 2048 Hz, amplified x150, and band-pass filtered at 10-500Hz using a Quattrocento signal amplifier (OT Bioelettronica, Inc., Turin, IT). EMG and torque were temporally synced with a 1-second TTL pulse transmitted to the Quattrocento at the onset of each trial.

***Figure 1.** Experimental setup and visual feedback conditions are shown above. Participants were seated with their right ankle secured in a device to measure isometric joint torque, with HDsEMG placed on their tibialis anterior, medial gastrocnemius, and soleus muscles. During the experiment real-time visual feedback was provided on a monitor in front of participants, with three different types of feedback, to cue triangular ramp contractions.*

Experimental Protocol

Before beginning data collection, real-time HDsEMG recordings were visually examined to ensure a high signal-to-noise ratio. First, participants were asked to perform maximum voluntary contractions (MVCs) with their plantarflexors and dorsiflexors. Three MVC trials in each direction were collected. If the last trial had the highest peak torque, additional trials were added as needed. Verbal encouragement was given to participants during MVC trials to ensure maximal performance, and participants were given 2 minutes of rest between trials to prevent fatigue. The maximum torque achieved was used for subsequent normalization of all trials.

The remaining trials consisted of a single triangular-shaped isometric dorsiflexion or plantarflexion torque ramp, with visual feedback to guide participants. For all ramps, participants increased torque (3% MVC/s) to 30% MVC over 10 seconds and then decreased at the same rate to 0% MVC. Throughout the experiment, a large monitor was positioned in front of the participants to provide real-time feedback of torque performance.

Visual feedback: This dataset was obtained from a larger ongoing project, and therefore
participants performed contractions with three different types of visual feedback, all of which had
the same goal of a 10 s linear ascending phase to a peak of 30% MVC and then a 10 s linear
descending phase to rest. During all three of these conditions, participants received real-time
torque feedback on the screen roughly two meters in front of them, which was comprised of either:
1) a standard guideline with a width of 10% of target torque (i.e., 3% MVC); 2) a precise guideline
with a width of 2.5% of target torque (i.e., 0.75% MVC); or 3) cross hairs to provide feedback
about the amplitude and timing of torque generation, along with cues about when to start and stop
contracting and a peak target (see Figure 2). At a minimum, participants completed 4 trials while
receiving each of the three types of feedback in each direction (i.e., plantarflexion and
dorsiflexion), yielding 12 ramps for both dorsiflexion and plantarflexion. Trials were grouped by
direction and randomized such that participants either performed all plantarflexion first or all
dorsiflexion first. Within each direction, the three types of feedback were randomized. Trials were
discarded if they did not exhibit a smooth increase and decrease of torque (i.e., substantial
deviations from the intended target profile), and the trial was then repeated.

Data Analysis

MU Decomposition: After data collection, EMG signals were manually inspected. Channels with
substantial artifacts, noise, or analog to digital saturation were removed (on average 1-3 channels
per array). Individual MU spike trains were decomposed using a convolutive blind-source
separation algorithm with a silhouette threshold of 0.87 (Negro *et al.*, 2016).

After decomposition, MU spike trains were manually edited by a trained technician who
was blind to the participant's sex. This inspection used a custom-made graphical user interface
in MATLAB to correct minor errors made by the decomposition algorithm using well-validated
local re-optimization methods to improve MU spike trains similar to the techniques used in recent
studies (Hug et al, 2021; Boccia *et al.*, 2019; Afsharipour *et al.*, 2020; Del Vecchio *et al.*, 2020;
Hassan *et al.*, 2020; Martinez-Valdes *et al.*, 2020). Instantaneous discharge rates of each MU
spike train were determined by computing the inverse of the interspike interval and smoothed
using support vector regression (Beauchamp *et al.*, 2022) with custom-written MATLAB scripts.
Within these scripts, the initial, peak, and final discharge rates were extracted from the smoothed
spike trains. Ascending duration was calculated as the time that a MU exhibited sustained
discharge before peak torque, and descending duration as the time a MU exhibited sustained
discharge from peak torque to derecruitment. Ascending-descending phase duration ratio was
calculated as the difference in ascending duration and descending duration, divided by the total

duration of discharge. A more negative ratio indicates greater duration on the descending portion

of the contraction, and thus more sustained discharge on the descending portion of the torque

ramp. Finally, recruitment threshold was calculated as the level of torque at the first MU spike.

Estimates of Persistent Inward Currents (PICs): Effects of PICs on MU discharge patterns can

be estimated by quantifying the amount of onset-offset hysteresis (i.e., ΔFrequency, ΔF) of a

higher-threshold (test) MU with respect to the discharge rate of a lower-threshold (reporter) MU

(Gorassini *et al.*, 1998, 2002*a*, 2002*b*). Rather than providing ΔF values for each test-reporter unit

pair, we calculated 'unit-wise' values, which gives each test unit one ΔF value based on the

average values obtained from multiple reporter units (Figure 3). Criteria for inclusion of ΔF values

from MU pairs were that 1) the test MU was recruited at least 1 second after the reporter unit to

ensure full activation of the PIC (Bennett *et al.*, 2001; Powers *et al.*, 2008; Hassan *et al.*, 2020),

test unit-reporter unit pair exhibited rate-rate correlations of $r^2 > 0.7$ to ensure MU pairs received

common synaptic input (Gorassini *et al.*, 2004; Udina *et al.*, 2010; Stephenson & Maluf, 2011;

Vandenberk & Kalmar, 2014), and 3) the reporter unit modulated its discharge rate by at least 0.5

pps while the test unit was active (Stephenson & Maluf, 2011).

Statistical Procedures

Statistical analyses were performed using R Statistical Software (v4.1.0; R Core Team

2021). To determine if variables of interest were predicted by the fixed effects of sex, muscle, and

their interactions we used linear mixed effects models with covariates of recruitment threshold

and the type of visual feedback, and a random intercept and muscle slope for participant ((Muscle

| Participant); lmer R package; v1.1.27.1; Bates *et al.*, 2015). To determine significance, we

applied Satterthwaite's method for degrees of freedom (lmerTest R package; v3.1.3; Kuznetsova

et al., 2017). Both individual participant means and group estimated marginal means (male and

female) were computed for each variable (emmeans R package, v1.8.0, Lenth *et al.*, 2022).

Results are reported as mean $\pm$ standard deviation. Effect size (Cohen's d) was calculated to

determine the standardized magnitude of the effect of sex from the estimated marginal means of

the male and female data from the model (emmeans R package, v1.8;0, Lenth *et al.* 2022). All

data were visualized in R (ggplot R package, v3.3.6, Wickham *et al.*, 2022).

Comparison between sexes with resampled MUs of matched recruitment torques: The

inclusion of recruitment torque as a covariate in the linear mixed effects model indicates that

estimates of PICs were higher in females than males across all three muscles despite the limited

number of low threshold MUs we decomposed in females. However, to further investigate the

possibility that a difference in recruitment threshold between the sexes influenced estimates of

PICs and peak discharge rate, we iteratively resampled MUs between the sexes, and compared

ΔF and peak discharge rate with similar distributions of recruitment torques. To perform this

resampling procedure, for each muscle, we randomly sampled 500 MUs from each sex such that

their distributions of recruitment threshold matched each other and represented the mean and

standard deviation of MUs across both sexes. To ensure that distributions and collected MUs

were not identical with each iteration, we offset the desired mean by a value randomly chosen

from a gaussian distribution with a mean of zero and standard deviation of one-quarter of the

population standard deviation. Within each iteration, we quantified ΔF and peak discharge rate

for each of the MUs in the new distributions and used a linear mixed effects model with a fixed

effect of sex and nested random effect of trial number within participant for sex (1 | Sex : PID :

Trial Number; fitlme MATLAB package) to determine if sex was a significant predictor of ΔF and

peak discharge rate. We repeated this process 1,000 times to determine the probability of

significant sex differences, which we report as a value from 0 (0% of runs with a significant

difference between sexes) to 1 (100% of runs with a significant difference between sexes).

RESULTS

In this study, we characterized sex differences in MU discharge properties during isometric
triangular shaped contractions to 30% of MVC. For females, we decomposed a total of 2,126 MUs
(TA: 933, MG: 691, SOL: 502), and for males a total of 3,935 MUs (TA: 1651, MG: 1484, SOL:
800). Table 1 shows the average number of units decomposed per trial for each muscle and sex.

Table 1. *The mean and standard deviation (SD) of units decomposed per trial for each muscle*
for females and males.

	Number of Units Decomposed per Trial			
	Female		Male	
	Mean (s)	*SD*	*Mean (s)*	*SD*
TA	12.61	4.28	20.22	9.06
MG	9.66	5.43	18.90	11.44
SOL	7.91	3.21	11.18	5.84

A greater number of lower threshold MUs are identified in males

When decomposed MUs were stratified by recruitment threshold, it becomes apparent
that a greater number of lower threshold MUs were decomposed in males than females (Figure
2). Thus, sex was a significant predictor of recruitment torque ($\chi^2_{(1)}$ = 13.56, p = 0.00023), and
was higher in females across all three muscles (Figure 2). To account for this in all subsequent
analyses, we included a covariate of recruitment torque in our statistical models.

Figure 2. Decomposed MUs by recruitment threshold for all participants for each muscle. Male units are shown in pink, female units overlayed in purple. For each figure, the x-axis shows the percentage of MVC the units were recruited in 1% bins. In the left panels, the y-axis represents the total number of units recruited, whereas in the right panels, the y-axis represents the percentage of total units recruited at each torque level.

Estimates of persistent inward currents are greater in females

For both sexes, estimates of PICs were in the normal range of values reported in previous studies (Afsharipour *et al.*, 2020. Kim *et al.*, 2020). Across all muscles, ΔF was larger ($\chi^2_{(1)}$ = 6.26, *p* = 0.012) in females (4.73 ± 0.242 pps) than males (3.81 ± 0.240 pps) with an effect size of d = 0.653. Figure 3 shows an example dorsiflexion trial from one male and one female, with individual ΔF values provided for the average of all test units and individually for each MU decomposed from the TA muscle. In Figure 4, within participant means (diamonds) and individual ΔF values for each test unit along with distributions across all three muscles are shown. No significant interaction between sex and muscle was identified ($\chi^2_{(2)}$ = 1.79, *p* = 0.409), and therefore ΔF is estimated 0.92 pps higher for all muscles in females based on estimated marginal means.

Figure 3. An example of a dorsiflexion contraction trial from one male (left) and one female (right)
during which motor units were decomposed from high-density surface electromyography over the
tibialis anterior muscle. Torque is plotted with a thick black line (top) and all individual motor units
recruited during that trial are plotted in the bottom plots according to recruitment threshold (lower
thresholds at the bottom). Each point represents the instantaneous discharge rate, and the thin
black line for each unit represents the smoothed discharge rate computed using support vector
regression. The test-unit mean, averaged across all suitable reporter units, is provided to the right
of each unit. The average of all these values within the trial is provided just below the torque trace.

Figure 4. *Estimates of persistent inward currents (ΔF) for TA (left), MG (middle), and SOL (right)*

for females (purple) and males (pink). Individual participant means are shown in diamonds. ΔF

measures for each unit collected across trials are shown as scatter points. Box plots represent

the 25th, 50th (median) and 75th quartiles, with whiskers for minimums and maximums at 1.5 the

interquartile range. Distributions of unit-wise ΔF values irrespective of participant are shown to

the right for each muscle.

The descending phase duration of discharge, from peak torque to derecruitment ($\chi^2_{(1)}$ =

3.316, *p* = 0.0686), and ascending duration, from recruitment to peak torque ($\chi^2_{(1)}$ = 0.0342, *p* =

0.8532), were similar between the sexes (Table 2B). Additionally, the ratio between ascending

and descending durations did not differ significantly between the sexes ($\chi^2_{(1)}$ = 2.94, *p* = 0.086,

Table 2C).

A. Ascending Phase Duration

	Female		Male	
	Mean (s)	*SD*	*Mean (s)*	*SD*
TA	6.09	0.52	6.11	0.52
MG	5.97	0.51	5.98	0.51
SOL	6.03	0.51	6.09	0.52

B. Descending Phase Duration

	Female		Male	
	Mean (s)	*SD*	*Mean (s)*	*SD*
TA	7.32	1.72	6.58	1.74
MG	5.23	1.88	4.48	1.87
SOL	5.77	1.78	5.02	1.85

C. Ascending-Descending Phase Ratio

	Female		Male	
	Mean (s)	*SD*	*Mean (s)*	*SD*
TA	-0.16	0.20	-0.02	0.20
MG	0.12	0.25	0.12	0.25
SOL	0.0369	0.22	0.0817	0.24

293 **Table 2.** *Measurements of motor unit discharge duration during the triangular ramp contractions.*
294 *Estimated marginal means and standard deviations are shown for each muscle, with females on*
295 *the left and males on the right. Abbreviations: TA; tibialis anterior, MG; medial gastrocnemius,*
296 *SOL; soleus, s; seconds, SD; standard deviation.*

MU Discharge rates did not differ between sexes

298   While peak discharge rates appeared higher in females than males (see table 3A), no
299 significant differences were detected between the two groups. Across muscles, discharge rates
300 did not differ at recruitment ($\chi^2_{(1)}$ = 1.36, p = 0.243), derecruitment ($\chi^2_{(1)}$ = 0.06, p = 0.805), or at
301 the peak ($\chi^2_{(1)}$ = 1.15, p = 0.283).

302 | | *A. Discharge Rate at Recruitment* | | | |
303

	A. Discharge Rate at Recruitment			
	Female		Male	
	Mean (pps)	*SD*	*Mean (pps)*	*SD*
TA	6.73	1.98	6.29	2.01
MG	5.01	1.91	4.57	1.96
SOL	4.04	1.83	3.60	1.88

	B. Peak Discharge Rate			
	Female		Male	
	Mean (pps)	*SD*	*Mean (pps)*	*SD*
TA	17.20	2.33	16.40	2.41
MG	13.00	2.72	12.10	2.76
SOL	11.10	2.43	10.30	2.57

	C. Discharge Rate at Derecruitment			
	Female		Male	
	Mean (pps)	*SD*	*Mean (pps)*	*SD*
TA	5.55	1.61	5.50	1.62
MG	4.60	1.63	4.56	1.64
SOL	3.70	1.58	3.65	1.60

Table 3. *Motor unit discharge rates during triangular ramp contractions. Estimated marginal*
means and standard deviations are shown for each muscle, with females on the left and males
on the right. Abbreviations: TA; tibialis anterior, MG; medial gastrocnemius, SOL; soleus, pps;
pulses per second; s; seconds, SD; standard deviation.

Resampling recruitment threshold distributions

To account for the differences in recruitment thresholds, we resampled 1000 iterations of
random MUs with a similar distribution of recruitment threshold between males and females
showed that ΔF was still significantly higher in females than males (Fig. 5B-C). Across the 1000
iterations, we found a probability of obtaining significant differences in ΔF of 0.98, 1.00, and 1.00
for the MG, SOL, and TA, respectively. Furthermore, we found the average beta coefficients for
the fixed factor of sex across all runs to be 0.63 ± 0.08, 1.01 ± 0.1, and 1.06 ± 0.1 for the MG,
SOL, and TA, respectively. Resampling also revealed that peak discharge rates were higher in
females for MG and SOL, but not TA (Fig. 5D-E), with the probability of significant differences of
0.63, 1.00, and 0.00, respectively. The fact that the resampled data with similar recruitment
thresholds revealed higher discharge rates in females likely reflects the larger numbers of low
threshold units in males compared to females (see Discussion).

Figure 5. Estimates of persistent inward currents (ΔF) and peak discharge rate between females
(F) and males (M) based on resampled MUs with equal distributions of recruitment threshold
between the sexes. **A.** Average recruitment threshold, percent maximum voluntary contraction (%

MVC), of resampled MUs for each sex across 1,000 runs for each muscle along the line of identity.

B. Unit-wise and average ΔF values from resampled MUs for each muscle and sex. **C.** Beta

coefficients (β), indicating the difference between female and male ΔF for each run, with filled

points representing significant values and unfilled indicating non-significant values. **D.** Individual

and average peak discharge rate from resampled MUs for each muscle and sex. **E.** Beta

coefficients (β), indicating the difference between female and male peak discharge rates for each

run, with filled points representing significant values and unfilled indicating non-significant values.

DISCUSSION

This study quantified the discharge patterns of several concurrently active MUs during submaximal isometric triangular-shaped contractions in the lower limb muscles of females and males. Although no significant differences in MU discharge rates were identified, estimates of PIC magnitude using a paired motor unit analysis technique revealed that discharge rate hysteresis (i.e., ΔF) was significantly larger in females than males. These findings suggest that descending neuromodulatory drive, monoaminergic signaling of 5HT and/or NE to motor units, and/or inhibitory inputs may differ between the sexes.

More lower threshold MUs decomposed in males than in females

In the current investigation, we identified a greater number of MUs in males across all three of the sampled muscles. Recent work by our colleagues (Taylor *et al.*, 2022) also documented a higher number of Mus decomposed from the TA muscle in males compared to females, and our findings extend these findings to other muscles of the lower leg. In particular, a greater proportion of lower threshold MUs were decomposed in males compared to females in the current study.

Methodological considerations that may skew comparisons of recruitment thresholds between heterogeneous groups of people (i.e., females vs males herein) should be fully considered before in-depth interpretations are made. For example, HDsEMG decomposition is biased toward larger and more superficial MUs (Hassan *et al.*, 2019) and, as shown by Guo and colleagues (Guo *et al.*, 2022), males tend to have larger MUs, which may explain why their spike times are decomposed more readily than their female counterparts (Taylor *et al.*, 2022). These sex differences in the decomposition results are important to note because they highlight a barrier to female inclusion in past HDsEMG analyses. To account for differences in recruitment threshold on ΔF and peak discharge rate, we ran a model to resample MUs with similar distributions of recruitment torque to compare these measures between the sexes. Both the comparisons of our entire dataset and the resampled data with comparable recruitment thresholds are considered in our interpretations and discussion below.

Estimates of PICs are higher in females

Linear mixed effects models revealed that biological sex was a significant predictor of estimates of PIC magnitude (ΔF) across all three muscles. Indeed, ΔF was almost 1 pps higher in females than males, indicating a greater contribution of PICs to MU discharge while performing a simple isometric ramp contraction. Resampling of MUs to compare estimates of PICs with equal

distributions of recruitment torque agreed with this finding, confirming that less low threshold MUs

decomposed in females did not affect estimates of PICs and comparisons between the sexes.

PICs are vital for motoneuronal function – without their amplification effects, repetitive

discharge cannot be achieved (Lee & Heckman, 2001), as driving currents from descending and

sensory synaptic inputs are too weak to cause repetitive MU discharge (Binder *et al.*, 2020).

Therefore, identifying and understanding differences in PIC magnitude may garner insights into

potential sex differences in motor command structure. Even though males in our experiment

certainly displayed discharge rate hysteresis (i.e., positive ΔF values), the magnitude of this

hysteresis was markedly smaller than females.

The factors that can influence PIC magnitude, and thus ΔF, include: 1) monoaminergic

input to the motoneurons, 2) Na^+ or Ca^{2+} channel function, and/or changes in 5HT/NE reuptake

and receptor function, and 3) the amount or pattern of inhibitory input to the motoneuron.

Physiologically, males and females differ most notably in their levels of sex hormones. In

eumenorrheic females (premenopausal females with a regular menstrual cycle), levels of

estradiol, one of the three endogenous estrogens, and progesterone fluctuate throughout the

month over a menstrual cycle (Sherman & Korenman, 1975). In males, the levels of these two

hormones remain low and do not fluctuate across the month. Progesterone levels in males are

similar to levels of females during the early menstrual cycle (follicular phase), but after ovulation,

progesterone levels rise much higher in females (Oettel & Mukhopadhyay, 2004; Tea *et al.*, 1975).

Throughout the entire menstrual cycle, eumenorrheic females experience fluctuations in estradiol

ten times that of males (Verdonk *et al.*, 2019). Remarkably, these two hormones have known

effects on neurons throughout the nervous system, in particular serotonin (Bethea *et al.*, 1998),

and thus, higher fluctuating levels in females may have effects on the motor system.

MU discharge rates did not differ between males and females

Despite the tendency for females to have higher discharge rates overall, there were no

significant sex differences in MU discharge rates when considering our entire dataset. Previous

studies have reported higher discharge rates in female MUs, however, during these experiments

participants performed isometric holds or trapezoid-shaped contractions (Inglis & Gabriel, 2020;

Guo *et al.*, 2022; Taylor *et al.*, 2022). In our study, discharge rates at onset, maximum, and offset

did not differ between the sexes. The greater yield of lower threshold units in the males may have

skewed the discharge rates of males to a higher level, due to the relationship of recruitment

threshold and discharge rate during submaximal isometric contractions (Pearcey & Rymer, 2022),

which would diminish statistical differences between the sexes. We did, however, attempt to deal

with this limitation by including a covariate of recruitment threshold in our models. It remains

possible that, if participants had been asked to sustain a contraction, sex differences may have

emerged as has been observed by previous studies. Moving one step beyond a covariate in our

analysis, the peak discharge rate in our resampled data with equal distributions of recruitment

torque revealed muscle specific differences between the sexes. The resampling of MUs revealed

that females had greater peak discharge rate in the MG and SOL, suggesting that peak discharge

rates may be higher in females when matched for recruitment torque in the plantarflexors, not in

the dorsiflexors (i.e., TA).

The potential role of sex hormones in mediating PIC magnitude

A possible underlying mechanism that could explain higher PICs in females is the

interaction between monoamines released from the caudal raphe nucleus (i.e., 5HT) and locus

coeruleus (i.e., NE) and the sex hormone estradiol. These monoamines facilitate PICs, and

therefore the magnitude of PICs is directly proportional to the level of brainstem monoamine

release (Lee & Heckman, 1998, 2000). Thus, sex differences in monoaminergic release could be

responsible for differences in ΔF, or perhaps the effects of estradiol on 5HT neurons in other

areas of the nervous system. In nonhuman female primates, estrogen receptor beta (Erβ) has

been identified in the dorsal and caudal raphe nuclei (Gundlah *et al.*, 2000; Bethea *et al.*, 2002).

Long-term loss of estradiol in ovariectomized female macaques also results in a reduction in the

number of 5HT neurons (Rivera & Bethea, 2012). Taken together, these data suggest that

females could have greater release of brainstem-derived monoamines than males due to greater

estradiol effects in the brainstem.

The hormones estradiol (one of the three endogenous estrogens) and progesterone

fluctuate across the female menstrual cycle in young adult females, and peak at levels much

higher than males during specific days (Sisk & Foster, 2004). Thus, since this work has revealed

potential sex differences in monoaminergic signaling, an essential next step will be to determine

the effect of fluctuating levels of estradiol and progesterone on estimates of PICs. If sex

differences were in fact caused by differences in sex hormones, and since we did not control or

measure the time of the menstrual cycle in females, we would expect greater variability in

estimates of PICs across the female group due to the fluctuating levels of sex hormones in

females in comparison to the relatively stable levels in males. Additional examination of values

obtained across individuals within each sex indeed suggests that there is more variability in ΔF

and discharge rates in the group of females compared to males. Measures of variance in ΔF

obtained from our mixed-effects models were indeed greater in females (see Figure 4).

Confidence in this sentiment, however, requires further study, in which females are studied at
multiple time points throughout their menstrual cycle, and sex hormones are quantified.
436

Methodological considerations

The non-invasive nature of surface electromyography is advantageous in that it allows us to
estimate the spike times of many MUs but since we are unable to measure from the spinal
motoneurons directly, this method is unable to directly quantify the magnitude of PICs. The paired
MU analysis technique to estimate PIC magnitude in humans has been well-validated, but it is
possible that the mechanisms behind the observed sex differences in ΔF are not solely due to
changes in monoaminergic drive. For example, differences in the patterns of inhibitory inputs,
monoamine release, monoaminergic receptor activity and/or sensitivity differences between the
sexes could possibly account for differences in MU discharge patterns (Beauchamp *et al.*, 2023).
Future studies which include a comprehensive array of contractions, that complement the
isometric triangular ramps utilized herein, and allow inferences about the biophysical properties
of motoneurons (Khurram *et al.*, 2022) or that use electrical stimulation or vibration protocols to
probe excitatory and inhibitory circuits (Mesquita *et al.*, 2021, Pearcey *et al.*, 2022, Orssatto *et*
al., 2022), may reveal further insight into the mechanism behind these sex differences. A
complimentary approach that also incorporates measures of musculoskeletal properties and/or
mechanical/passive properties of the muscle during the contraction (Martinez-Valdes *et al.*, 2022)
may further elucidate the mechanisms responsible for sex differences in discharge patterns
quantified here (Heroux *et al.* 2020).

Another limitation is that we assessed only one submaximal level of contraction (i.e., 30%
MVC). It is possible that sex differences in PICs and discharge rates are dependent on contraction
intensity. For example, previous work has shown discharge rates during plateau holds are higher
in females until 40% MVC, equal between the sexes at 60 and 80%, and higher in males at 100%
MVC (Inglis and Gabriel 2020). When data is strength matched, however, discharge rates in
females were higher across all intensities. In the current experiment, estimates of PICs were
higher in females during low intensity contractions but it is possible that increasing the contraction
intensity could reveal further insight into monoaminergic signaling between the sexes.

Another limitation of this study is that the menstrual cycle status of female participants was
not recorded and thus could not be taken into consideration during analysis. The uncontrolled
phase of the menstrual cycle in females remains a potential reason for the increased variability in
female ΔF values. Previous work has shown that various aspects of motor control, including MVC
(Sarwar *et al.*, 1996; Nicolay *et al.*, 2007), fatigability (Sarwar *et al.*, 1996; Ansdell *et al.*, 2019;

Pereira *et al.*, 2020), and initial MU discharge rate (Tenan *et al.*, 2013) all fluctuate across the

menstrual cycle. There is reason to believe that MU discharge and PIC magnitude would also be

affected by fluctuating levels of sex hormones, and thus this study provides motivation to examine

these MU properties across the female menstrual cycle.

Subcutaneous fat, or the distance between the muscle and the surface array, is a likely

contributor to the lower MU yield in females, however we cannot confirm this because we did not

perform appropriate measures for each participant. Larger distances between the electrodes and

the muscle fibers in the females recruited for this study (i.e., higher average levels of

subcutaneous fat) are likely, based on typical (i.e., population) anthropometrics (Deurenberg *et*

al., 1991). However, future analyses that include matching subcutaneous fat thickness between

the sexes will offer further insight.

Conclusion

We have revealed that estimates of PIC magnitudes are higher in the lower limb muscles

of females than males. This suggests female MUs receive a greater contribution from PICs than

males to achieve the same relative force and discharge rate, which may be due to a requirement

for more persistent discharge or a consequence of more monoamine availability. Females may

also use a different combination of synaptic excitation, inhibition, and neuromodulation than males

to achieve the same relative force levels during triangular ramp contractions. As such, this study

further highlights the necessity to include equal numbers of female participants in order to, not

only, fully understand the human physiology behind motor control, but to potentially use MU

analysis as biomarkers for neurological disease (Fajardo *et al.*, 2022).

Acknowledgements

The authors would like to thank Obaid Khurram for his fruitful discussion during planning of the experiments and Kirsten Carroll for assistance with data analysis.

Data availability statement

The data that support the findings of this study are available on request from the corresponding author.

Competing interests

The authors declare that they have no competing interests.

Author contributions

S.T.J., J.A.B., C.J.H., and G.E.P.P., conceptualized and designed the research; S.T.J., J.A.B., M.M.G., and G.E.P.P. performed the experiments; S.T.J. and J.A.B. analyzed the data; S.T.J., J.A.B., G.E.P.P., and C.J.H. interpreted the results of experiments; S.T.J., J.A.B., and G.E.P.P., prepared the figures; S.T.J. drafted the manuscript; S.T.J., J.A.B., M.M.G., G.E.P.P., F.N., and C.J.H. revised and approved the final version of the manuscript.

Funding

This work was supported by a National Institute of Health (NIH) grant 5R01NS098509-05, NIH NINDS F31 NS120500, São Paulo Research Foundation (FAPESP) grant 2020/03282-0, and Natural Sciences and Engineering Research Council of Canada Postdoctoral Fellowship.

REFERENCES

Ansdell P, Brownstein CG, Škarabot J, Hicks KM, Simoes DCM, Thomas K, Howatson G, Hunter SK & Goodall S (2019). Menstrual cycle-associated modulations in neuromuscular function and fatigability of the knee extensors in eumenorrheic women. *Journal of Applied Physiology* **126,** 1701–1712.

Augsburger GR, Soloveva A & Carr JC (2022). Sex and limb comparisons of neuromuscular function in the morning versus the evening. *Physiological Reports* **10,** e15260.

Barth C, Villringer A & Sacher J (2015). Sex hormones affect neurotransmitters and shape the adult female brain during hormonal transition periods. *Front Neurosci* **9,** 37.

Beauchamp JA, Khurram OU, Dewald JPA, Heckman C & Pearcey GEP (2022). A computational approach for generating continuous estimates of motor unit discharge rates and visualizing population discharge characteristics. *J Neural Eng* **19,** 016007.

Beauchamp JA, Pearcey GEP, Khurram OU, Chardon M, Wang YC, Powers RK, Dewald JPA & Heckman CJ (2023). A geometric approach to quantifying the neuromodulatory effects of persistent inward currents on individual motor unit discharge patterns. *J Neural Eng*; DOI: 10.1088/1741-2552/acb1d7.

Bennett DJ, Li Y & Siu M (2001). Plateau Potentials in Sacrocaudal Motoneurons of Chronic Spinal Rats, Recorded In Vitro. *Journal of Neurophysiology* **86,** 1955–1971.

Bethea CL & Reddy AP (2012). Ovarian Steroids Increase Glutamatergic Related Gene Expression in Serotonin Neurons of Macaques. *Mol Cell Neurosci* **49,** 251–262.

Bethea CL, Lu NZ, Gundlah C & Streicher JM (2002). Diverse Actions of Ovarian Steroids in the Serotonin Neural System. *Frontiers in Neuroendocrinology* **23,** 41–100.

Bethea CL, Pecins-Thompson M, Schutzer WE, Gundlah C & Lu ZN (1998). Ovarian steroids and serotonin neural function. *Mol Neurobiol* **18,** 87–123.

Binder MD, Powers RK & Heckman CJ (2020). Nonlinear Input-Output Functions of Motoneurons. *Physiology (Bethesda)* **35,** 31–39.

Bitran D, Shiekh M & McLeod M (1995). Anxiolytic effect of progesterone is mediated by the neurosteroid allopregnanolone at brain GABAA receptors. *J Neuroendocrinol* **7,** 171–177.

Celichowski J & Drzymała H (2006). Differences between properties of male and female motor units in the rat medial gastrocnemius muscle. *J Physiol Pharmacol* **57,** 83–93.

Celichowski J & Drzymała-Celichowska H (2007). The number of motor units in the medial gastrocnemius muscle of male and female rats. *J Physiol Pharmacol* **58,** 821–828.

Deurenberg P, Weststrate JA & Seidell JC (1991). Body mass index as a measure of body fatness: age- and sex-specific prediction formulas. *Br J Nutr* **65,** 105–114.

English AW & Widmer CG (2003). Sex differences in rabbit masseter motoneuron firing
behavior. *J Neurobiol* **55,** 331–340.

Fajardo ME, Jenz S & Heckman CJ (2022). The excitable motor unit: A powerful diagnostic and
pathophysiological marker for ALS? *Clin Neurophysiol* **142,** 256–257.

Ford TW & Kirkwood PA (2006). Respiratory drive in hindlimb motoneurones of the
anaesthetized female cat. *Brain Research Bulletin* **70,** 450–456.

Gorassini M, Yang JF, Siu M & Bennett DJ (2002*a*). Intrinsic Activation of Human Motoneurons:
Possible Contribution to Motor Unit Excitation. *Journal of Neurophysiology* **87,** 1850–
1858.

Gorassini M, Yang JF, Siu M & Bennett DJ (2002*b*). Intrinsic activation of human motoneurons:
reduction of motor unit recruitment thresholds by repeated contractions. *J Neurophysiol*
87, 1859–1866.

Gorassini MA, Bennett DJ & Yang JF (1998). Self-sustained firing of human motor units.
Neuroscience Letters **247,** 13–16.

Gorassini MA, Knash ME, Harvey PJ, Bennett DJ & Yang JF (2004). Role of motoneurons in the
generation of muscle spasms after spinal cord injury. *Brain* **127,** 2247–2258.

Gundlah C, Kohama SG, Mirkes SJ, Garyfallou VT, Urbanski HF & Bethea CL (2000).
Distribution of estrogen receptor beta (ERβ) mRNA in hypothalamus, midbrain and
temporal lobe of spayed macaque: continued expression with hormone replacement.
Molecular Brain Research **76,** 191–204.

Guo Y, Jones EJ, Inns TB, Ely IA, Stashuk DW, Wilkinson DJ, Smith K, Piasecki J, Phillips BE,
Atherton PJ & Piasecki M (2022). Neuromuscular recruitment strategies of the vastus
lateralis according to sex. *Acta Physiologica*; DOI: 10.1111/apha.13803.

Harwood B, Cornett KMD, Edwards DL, Brown RE & Jakobi JM (2014). The effect of tendon
vibration on motor unit activity, intermuscular coherence and force steadiness in the
elbow flexors of males and females. *Acta Physiologica* **211,** 597–608.

Hassan A, Thompson CK, Negro F, Cummings M, Powers R, Heckman CJ, Dewald J &
McPherson LM (2019). Sensitivity of the paired motor unit analysis for estimation of
motoneuron excitability to commonly used constraints and filters. 732982. Available at:
https://www.biorxiv.org/content/10.1101/732982v1 [Accessed April 1, 2022].

Hassan A, Thompson CK, Negro F, Cummings M, Powers RK, Heckman CJ, Dewald JPA &
McPherson LM (2020). Impact of parameter selection on estimates of motoneuron
excitability using paired motor unit analysis. *J Neural Eng* **17,** 016063.

Heckman CJ & Enoka R (2012). Motor Unit. *Comprehensive Physiology* **2,** 2629–2682.

Heckman CJ, Gorassini MA & Bennett DJ (2005). Persistent inward currents in motoneuron
dendrites: Implications for motor output. *Muscle & Nerve* **31,** 135–156.

Heckman CJ, Johnson M, Mottram C & Schuster J (2008). Persistent Inward Currents in Spinal
583  Motoneurons and Their Influence on Human Motoneuron Firing Patterns. *Neuroscientist*
584  **14,** 264–275.

585 Hounsgaard J, Hultborn H, Jespersen B & Kiehn O (1988). Bistability of alpha-motoneurones in
586  the decerebrate cat and in the acute spinal cat after intravenous 5-hydroxytryptophan.
587  *The Journal of Physiology* **405,** 345–367.

588 Hunter SK & Enoka RM (2001). Sex differences in the fatigability of arm muscles depends on
589  absolute force during isometric contractions. *Journal of Applied Physiology* **91,** 2686–
590  2694.

591 Hyngstrom AS, Johnson MD & Heckman CJ (2008). Summation of Excitatory and Inhibitory
592  Synaptic Inputs by Motoneurons With Highly Active Dendrites. *J Neurophysiol* **99,** 1643–
593  1652.

594 Inglis JG & Gabriel DA (2020). Sex differences in motor unit discharge rates at maximal and
595  submaximal levels of force output. *Appl Physiol Nutr Metab* **45,** 1197–1207.

596 Inglis JG & Gabriel DA (2021). Sex differences in the modulation of the motor unit discharge
597  rate leads to reduced force steadiness. *Appl Physiol Nutr Metab* **46,** 1065–1072.

598 Jenz ST & Pearcey GEP (2022). Sex matters in neuromuscular control. *Acta Physiologica* **235,**
599  e13823.

600 Johnson MD, Thompson CK, Tysseling VM, Powers RK & Heckman CJ (2017). The potential
601  for understanding the synaptic organization of human motor commands via the firing
602  patterns of motoneurons. *Journal of Neurophysiology* **118,** 520–531.

603 Khurram OU, Pearcey GEP, Chardon MK, Kim EH, García M & Heckman CJ (2022). The
604  Cellular Basis for the Generation of Firing Patterns in Human Motor Units. *Adv Neurobiol*
605  **28,** 233–258.

606 Kirkwood PA, Lawton M & Ford TW (2002). Plateau potentials in hindlimb motoneurones of
607  female cats under anaesthesia. *Exp Brain Res* **146,** 399–403.

608 Lee RH & Heckman CJ (1998). Bistability in Spinal Motoneurons In Vivo: Systematic Variations
609  in Persistent Inward Currents. *Journal of Neurophysiology* **80,** 583–593.

610 Lee RH & Heckman CJ (2000). Adjustable Amplification of Synaptic Input in the Dendrites of
611  Spinal Motoneurons *In Vivo*. *J Neurosci* **20,** 6734–6740.

612 Lee RH & Heckman CJ (2001). Essential Role of a Fast Persistent Inward Current in Action
613  Potential Initiation and Control of Rhythmic Firing. *Journal of Neurophysiology* **85,** 472–
614  475.

615 Lulic-Kuryllo T & Inglis JG (2022). Sex differences in motor unit behaviour: A review. *J
616  Electromyogr Kinesiol* **66,** 102689.

617 Martinez-Valdes E, Negro F, Botter A, Pincheira PA, Cerone GL, Falla D, Lichtwark GA &
618  Cresswell AG (2022). Modulations in motor unit discharge are related to changes in

fascicle length during isometric contractions. *Journal of Applied Physiology* **133,** 1136–1148.

Melcangi RC, Giatti S, Calabrese D, Pesaresi M, Cermenati G, Mitro N, Viviani B, Garcia-Segura LM & Caruso D (2014). Levels and actions of progesterone and its metabolites in the nervous system during physiological and pathological conditions. *Prog Neurobiol* **113,** 56–69.

Mesquita RNO, Taylor JL, Kirk B & Blazevich AJ (2021). Involuntary sustained firing of plantar flexor motor neurones: effect of electrical stimulation parameters during tendon vibration. *Eur J Appl Physiol* **121,** 881–891.

Michopoulos V, Berga SL & Wilson ME (2011). Estradiol and progesterone modify the effects of the serotonin reuptake transporter polymorphism on serotonergic responsivity to citalopram. *Exp Clin Psychopharmacol* **19,** 401–408.

Montague SJ, Fenrich KK, Mayer-Macaulay C, Maratta R, Neuber-Hess MS & Rose PK (2013). Nonuniform distribution of contacts from noradrenergic and serotonergic boutons on the dendrites of cat splenius motoneurons. *J Comp Neurol* **521,** 638–656.

Nicolay CW, Kenney JL & Lucki NC (2007). Grip strength and endurance throughout the menstrual cycle in eumenorrheic and women using oral contraceptives. *International Journal of Industrial Ergonomics* **37,** 291–301.

Nishikawa Y, Holobar A, Watanabe K, Takahashi T, Ueno H, Maeda N, Maruyama H, Tanaka S & Hyngstrom AS (2022). Detecting motor unit abnormalities in amyotrophic lateral sclerosis using high-density surface EMG. *Clinical Neurophysiology*; DOI: 10.1016/j.clinph.2022.06.016.

Oettel M & Mukhopadhyay A (2004). Progesterone: the forgotten hormone in men? *The Aging Male* **7,** 236–257.

Oliveira DS de, Casolo A, Balshaw TG, Maeo S, Lanza MB, Martin NRW, Maffulli N, Kinfe TM, Eskofier BM, Folland JP, Farina D & Del Vecchio A (2022). Neural decoding from surface high-density EMG signals: influence of anatomy and synchronization on the number of identified motor units. *J Neural Eng* **19,** 046029.

Orssatto LBR, Fernandes GL, Blazevich AJ & Trajano GS (2022). Facilitation–inhibition control of motor neuronal persistent inward currents in young and older adults. *The Journal of Physiology* **600,** 5101–5117.

Parra ME, Sterczala AJ, Miller JD, Trevino MA, Dimmick HL & Herda TJ (2020). Sex-related differences in motor unit firing rates and action potential amplitudes of the first dorsal interosseous during high-, but not low-intensity contractions. *Exp Brain Res* **238,** 1133–1144.

Pearcey GE & Rymer WZ (2022). Population recordings of human motor units often display "onion skin" discharge patterns -- implications for voluntary motor control. ; DOI: 10.48550/arXiv.2208.11783. Available at: http://arxiv.org/abs/2208.11783 [Accessed September 1, 2022].

Pearcey GEP, Khurram OU, Beauchamp JA, Negro F & Heckman CJ (2022). Antagonist tendon
vibration dampens estimates of persistent inward currents in motor units of the human
lower limb. 2022.08.02.502526. Available at:
https://www.biorxiv.org/content/10.1101/2022.08.02.502526v1 [Accessed September 1,
2022].

Peng Y-L, Tenan MS & Griffin L (2018). Hip position and sex differences in motor unit firing
patterns of the vastus medialis and vastus medialis oblique in healthy individuals.
Journal of Applied Physiology **124,** 1438–1446.

Pereira HM, Larson RD & Bemben DA (2020). Menstrual Cycle Effects on Exercise-Induced
Fatigability. *Front Physiol* **11,** 517.

Powers RK, Nardelli P & Cope TC (2008). Estimation of the Contribution of Intrinsic Currents to
Motoneuron Firing Based on Paired Motoneuron Discharge Records in the Decerebrate
Cat. *Journal of Neurophysiology* **100,** 292–303.

Rivera HM & Bethea CL (2013). Ovarian steroids increase PSD-95 expression and dendritic
spines in the dorsal raphe of ovariectomized macaques. *Synapse* **67,** 897–908.

Sarwar R, Niclos BB & Rutherford OM (1996). Changes in muscle strength, relaxation rate and
fatiguability during the human menstrual cycle. *The Journal of Physiology* **493,** 267–272.

Sherman BM & Korenman SG (1975). Hormonal characteristics of the human menstrual cycle
throughout reproductive life. ; DOI: 10.1172/JCI107979. Available at:
https://www.jci.org/articles/view/107979/pdf [Accessed September 5, 2022].

Sisk CL & Foster DL (2004). The neural basis of puberty and adolescence. *Nat Neurosci* **7,**
1040–1047.

Smith SS (1994). Female sex steroid hormones: from receptors to networks to performance--
actions on the sensorimotor system. *Prog Neurobiol* **44,** 55–86.

Stephenson JL & Maluf KS (2011). Dependence of the paired motor unit analysis on motor unit
discharge characteristics in the human tibialis anterior muscle. *J Neurosci Methods* **198,**
84–92.

Sumner BEH & Fink G (1993). Effects of Acute Estradiol on 5-Hydroxytryptamine and
Dopamine Receptor Subtype mRNA Expression in Female Rat Brain. *Molecular and
Cellular Neuroscience* **4,** 83–92.

Sumner BEH & Fink G (1997). The density of 5-hydoxytryptamine2A receptors in forebrain is
increased at pro-oestrus in intact female rats. *Neuroscience Letters* **234,** 7–10.

Taylor CA, Kopicko BH, Negro F & Thompson CK (2022). Sex differences in the detection of
motor unit action potentials identified using high-density surface electromyography.
Journal of Electromyography and Kinesiology **65,** 102675.

Tea NT, Castanier M, Roger M & Scholler R (1975). Simultaneous radioimmunoassay of
plasma progesterone and 17-hydroxyprogesterone normal values in children, in men

695  and in women throughout the menstrual cycle and in early pregnancy. *Journal of Steroid*
696   *Biochemistry* **6,** 1509–1516.

697 Tenan MS, Peng Y-L, Hackney AC & Griffin L (2013). Menstrual cycle mediates vastus medialis
698   and vastus medialis oblique muscle activity. *Med Sci Sports Exerc* **45,** 2151–2157.

699 Thompson CK, Negro F, Johnson MD, Holmes MR, McPherson LM, Powers RK, Farina D &
700   Heckman CJ (2018). Robust and accurate decoding of motoneuron behaviour and
701   prediction of the resulting force output. *The Journal of Physiology* **596,** 2643–2659.

702 Udina E, D'Amico J, Bergquist AJ & Gorassini MA (2010). Amphetamine increases persistent
703   inward currents in human motoneurons estimated from paired motor-unit activity. *J
704   Neurophysiol* **103,** 1295–1303.

705 Vandenberk MS & Kalmar JM (2014). An evaluation of paired motor unit estimates of persistent
706   inward current in human motoneurons. *J Neurophysiol* **111,** 1877–1884.

707 Verdonk SJE, Vesper HW, Martens F, Sluss PM, Hillebrand JJ & Heijboer AC (2019). Estradiol
708   reference intervals in women during the menstrual cycle, postmenopausal women and
709   men using an LC-MS/MS method. *Clinica Chimica Acta* **495,** 198–204.

710 Woitowich NC, Beery A & Woodruff T (2020). Author response: A 10-year follow-up study of sex
711   inclusion in the biological sciences. ; DOI: 10.7554/eLife.56344.sa2. Available at:
712   https://elifesciences.org/articles/56344#sa2 [Accessed October 4, 2022].

713 Woolley C & McEwen B (1992). Estradiol mediates fluctuation in hippocampal synapse density
714   during the estrous cycle in the adult rat. *J Neurosci* **12,** 2549–2554.