Leucism and stripe formation in ball pythons (*Python regius*) are associated with variants affecting endothelin signaling

Uyen M. Dao[1], Izabella Lederer[1], Ray L. Tabor[1], Basmah Shahid[1], The BIO306W Consortium[^1], Chiron W. Graves[1], and Hannah S. Seidel[1*]

[1]Department of Biology, Eastern Michigan University, Ypsilanti, MI, USA

* Corresponding author:

E-mail: hseidel@emich.edu

^The BIO306W Consortium comprises students in an undergraduate laboratory course at Eastern Michigan University. Students are listed in the Acknowledgments.

Data availability: All relevant data are within the paper and its Supporting Information files.
DNA sequences are available from GenBank, under accession numbers OP589186-OP589187.

Funding information: This work was supported by a Summer Research/Create Activity Award
from Eastern Michigan University to HSS. The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Color patterns in non-avian reptiles are beautifully diverse, but little is known about the genetics and development of these patterns. Here we investigated color patterning in pet ball pythons (*Python regius*), which have been bred to show color phenotypes that differ dramatically from the wildtype form. We report that several color phenotypes in pet animals are associated with putative loss-of-function variants in the gene encoding endothelin receptor EDNRB1: (i) frameshift variants in *EDNRB1* are associated with conversion of the normal mottled color pattern to all-white skin (leucism), (ii) missense variants affecting conserved sites of the EDNRB1 protein are associated with dorsal, longitudinal stripes, and (iii) substitutions at *EDNRB1* splice donors are associated with subtle changes in patterning compared to wildtype. We propose that these phenotypes are caused by loss of specialized color cells (chromatophores), with loss ranging from severe (leucism) to moderate (dorsal striping) to mild (subtle changes in patterning). Our study is the first to describe variants affecting endothelin signaling in a non-avian reptile and suggests that reductions in endothelin signaling in ball pythons can produce a variety of color phenotypes, depending on the degree of color cell loss.

Introduction

Skin colors in vertebrates originate from specialized color cells. These cells are known as chromatophores, and they come in three main types: melanophores synthesize the pigment melanin, whose color ranges from black to reddish brown; xanthophores store red-to-yellow pigments, including pteridines synthesized by the cell and carotenoids obtained from the diet; and iridophores contain platelets of crystallized purines, which can appear blue, white, or iridescent, depending on their structure (Mills and Patterson 2009; Kelsh *et al.* 2009; Cuthill *et al.* 2017). Skin color in non-avian reptiles and lower vertebrates is determined by layers of multiple chromatophores (Patterson and Parichy 2019; Kuriyama *et al.* 2020). Green colors, for example, are produced by yellow xanthophores layered above blue iridophores; golden brown colors, as a second example, are produced by melanophores above yellow xanthophores above white iridophores (e.g. Saenko *et al.* 2013; Brejcha *et al.* 2019). Mammals possess only one type of chromatophore (melanophores), and their skin color is limited to shades of brown and black (Mort *et al.* 2015).

Chromatophores are derived from the neural crest and migrate during development to reach their final positions in the skin (Parichy and Spiewak 2015; Mort *et al.* 2015; Haupaix and Manceau 2020). Once in the skin, chromatophores organize into patterned arrangements to produce the overall color pattern of the animal. This process has been best characterized in zebrafish, where patterning is driven by short- and long-range interactions among chromatophores themselves (Singh and Nuesslein-Volhard 2015; Patterson and Parichy 2019). Similar mechanisms likely function in mammals, to produce periodic markings such as stripes and spots (Kaelin *et al.* 2012, 2021; Mallarino *et al.* 2016). Pattern formation is less well understood in non-avian reptiles, although the idea that similar mechanisms may function in this group is supported by theoretical studies showing that Turing-like interactions can produce many of their basic color patterns (Murray and Myerscough 1991; Kondo and Shirota 2009; Kuriyama *et al.* 2020; Ascarrunz and Sanchez-Villagra 2022).

An emerging model for understanding coloration and pattern formation in non-avian reptiles is the ball python (*Python regius*) (Garcia-Elfring *et al.* 2020; Brown *et al.* 2022). Ball pythons are native to west Africa, but have become common as pets in the United States. Wild ball pythons exhibit a mottled color pattern, consisting of irregular black and golden-brown patches of skin (Fig 1). Pet ball pythons, by contrast, have been bred to include animals with dramatically different color patterns (McCurley 2005; Broghammer 2019). These variants, known as 'color morphs,' include animals with solid body colors, longitudinal stripes, or major changes in pattern patchiness. Many of these phenotypes are reminiscent of patterning

changes that have occurred within the snake lineage across evolutionary time, including patterns thought to be adaptive for certain behavioral ecologies (Brodie 1992; Olsson *et al.* 2013; Allen *et al.* 2013; Pizzigalli *et al.* 2020). Color morphs in ball pythons may therefore provide insight into the genetic and developmental mechanisms controlling patterning changes across species.

Many color morphs in ball pythons show simple dominant or recessive patterns of inheritance, suggesting single-gene causality. Some morphs belong to allelic series, with some morphs in the series showing more dramatic phenotypes than others (McCurley 2005; Broghammer 2019). Such morphs are thought to represent different molecular variants of the same gene, akin to mutants in allelic series arising from classical mutagenesis screens in model organisms.

One example of an allelic series in ball pythons is the Yellowbelly series of color morphs (Fig 1). This series is described by breeders as having five alleles (*yellowbelly*, *spark*, *specter*, *gravel*, and *asphalt*). Phenotypes in this series include a loss of coloration and coincident changes in patterning, which range from severe to mild (Fig 1A-B). Homozygotes of the strongest allele (*yellowbelly*) have skin that is almost fully white (leucism) (Fig 1B). The intermediate alleles (*spark* and *specter*) produce homozygotes with moderately lighter coloration compared to wildtype and a pair of dorsal, longitudinal stripes (Fig 1B). The weakest alleles (*gravel* and *asphalt*) produce homozygotes with mildly lighter coloration compared to wildtype and subtle alterations in patterning (Fig 1B). Compound heterozygotes show phenotypes intermediate between the parents, often including dorsal, longitudinal stripes (Fig 1D). These phenotypes are largely recessive, and heterozygotes of any of these alleles in combination with the wildtype allele are difficult to distinguish from wildtype (Fig 1F). Some heterozygotes have slightly lighter underbellies compared to wildtype (Figs 1E and 1F), but this phenotype is variable and difficult to recognize, especially for alleles other than the *yellowbelly* allele.

Phenotypes in the Yellowbelly series suggest a loss of chromatophores to varying degrees. Leucism has been reported in many groups of vertebrates and is characteristic of an absence of chromatophores (Besmer *et al.* 1993; Hosoda *et al.* 1994; Gariepy *et al.* 1996; Yang *et al.* 1998; Kelsh 2006 p. 10; Cobaleda *et al.* 2007; Miwa *et al.* 2007; Pingault *et al.* 2010; Kinoshita *et al.* 2014; Woodcock *et al.* 2017; Ullate-Agote and Tzika 2021). Dorsal striping is reminiscent of a mutant phenotype in zebrafish caused by reductions in chromatophores. The adult color pattern in wildtype zebrafish is initiated by iridophore precursors that migrate via the horizontal midline to populate the skin (Fig 2A) (Singh and Nuesslein-Volhard 2015; Patterson and Parichy 2019). In zebrafish mutants with fewer iridophores, iridophores fail to populate the skin and instead remain clustered at the horizontal midline (Parichy *et al.* 2000; Lopes *et al.* 2008; Patterson and Parichy 2013; Frohnhöfer *et al.* 2013; Mo *et al.* 2017; Spiewak *et al.* 2018). As a consequence, melanophore survival is reduced, and melanophores persist only near iridophores (Patterson and Parichy 2013; Frohnhöfer *et al.* 2013; Patterson *et al.* 2014). The resulting phenotype is a pair of midline stripes on either side of the fish's body (Fig 2B). We hypothesized that these midline stripes in mutant zebrafish might be analogous to the dorsal stripes observed for morphs in the Yellowbelly series. This hypothesis was based on the presumption that chromatophores in ball pythons migrate from the neural crest in a dorsal-to-ventral path, as they do in other higher vertebrates (Mort *et al.* 2015). Incomplete loss of chromatophores in ball pythons might therefore cause chromatophores to remain clustered near the dorsal ridge, analogous to their clustering near the horizontal midline in mutant zebrafish. This hypothesis provides a simple explanation for the relationship between phenotypes in the Yellowbelly series: leucism may reflect chromatophore loss that is severe (Fig 2F); dorsal striping may reflect chromatophore loss that is moderate (Fig 2E); and more subtle alternations in patterning may reflect chromatophore loss that is mild (Fig 2D).

Our hypothesis about chromatophore loss for phenotypes in the Yellowbelly series suggested candidate genes that might be causative for these phenotypes. At the top of this list

were genes encoding components of endothelin signaling. Endothelin signaling comprises a vertebrate-specific signaling pathway responsible for regulating blood pressure and development of neural crest cells (Saldana-Caboverde and Kos 2010; Bondurand *et al.* 2018). Components of endothelin signaling include endothelin ligands, G-protein coupled receptors (endothelin receptors), and enzymes responsible for processing the ligands (endothelin-converting enzymes) (Braasch *et al.* 2009). Endothelin signaling is known to regulate chromatophore development or pattern formation in many species: zebrafish (Parichy *et al.* 2000; Patterson and Parichy 2013; Krauss *et al.* 2014; Spiewak *et al.* 2018; Camargo-Sosa *et al.* 2019), axolotl (Woodcock *et al.* 2017), chicken (Dorshorst *et al.* 2011; Shinomiya *et al.* 2012; Kinoshita *et al.* 2014), duck (Li *et al.* 2015), quail (Dupin *et al.* 2000), mouse (Baynash *et al.* 1994; Hosoda *et al.* 1994), rat (Ceccherini *et al.* 1995; Gariepy *et al.* 1996; Yoshihara *et al.* 2017), cat (Kaelin *et al.* 2012), horse (Yang *et al.* 1998), sheep (Lühken *et al.* 2012), and human (Pingault *et al.* 2010). A role for endothelin signaling in color patterning of non-avian reptiles has not been described.

The goal of the current study was to identify the genetic cause of color phenotypes in the Yellowbelly series. We predicted that these phenotypes were caused by loss-of-function variants in a gene encoding a component of endothelin signaling. Our approach was to search for putative loss-of-function variants in two candidate genes: *EDNRB1*, which encodes an endothelin receptor, and *EDN3*, which encodes an endothelin ligand (Braasch *et al.* 2009). These genes were prioritized given their role in chromatophore regulation in zebrafish (Parichy *et al.* 2000; Spiewak *et al.* 2018) and axolotl (Woodcock *et al.* 2017). Samples of pet ball pythons (shed skins) were recruited from pet owners across the United States, and genetic variants were tested for association with phenotypes in the Yellowbelly series.

Results

The *yellowbelly* allele is associated with frameshift variants in *EDNRB1*

We hypothesized that morphs in the Yellowbelly series were caused by loss-of-function variants in *EDNRB1* or *EDN3*. To test this hypothesis, we amplified and sequenced the coding regions and adjacent splice sites of *EDNRB1* and *EDN3* in one ball python described as a *yellowbelly* homozygote by its owner. Sequences from this animal were compared to sequences from an animal having normal coloration (henceforth 'wildtype'). We observed that the *yellowbelly* homozygote was homozygous for a 1-bp deletion in the fourth coding region of *EDNRB1* (OP589186:c.1646del, Fig 3B). No coding or splice-site variants were observed in *EDN3*. The 1-bp deletion in *EDNRB1* introduces a frameshift in the transcript, which removes or alters three of the seven transmembrane helices of the EDNRB1 protein (Fig 3E). This truncation likely abolishes protein function, given that G-protein-coupled receptors are generally unable to function without an intact transmembrane structure (Rosenbaum *et al.* 2009). We conclude that the 1-bp deletion is likely a strong loss-of-function allele of *EDNRB1*.

To test whether the 1-bp deletion in *EDNRB1* was associated with phenotypes attributed to the *yellowbelly* allele, we genotyped 117 additional animals for this deletion. These animals consisted of 69 animals described as *yellowbelly* homozygotes and 48 animals described as having normal coloration or belonging to morphs other than those in the Yellowbelly series (henceforth 'Non-Yellowbelly' animals). These animals were collected from a total of 35 and 33 pet owners, respectively, and thus represent a broad sampling of the pet population. We observed that all animals described as *yellowbelly* homozygotes were either homozygous for the 1-bp deletion (60 animals) or heterozygous for it (nine animals) (Fig 3C). By contrast, none of the Non-Yellowbelly animals carried the deletion (Fig 3C). These findings demonstrate a strong but imperfect association between the 1-bp deletion and morph type.

We hypothesized that the nine animals described as *yellowbelly* homozygotes but
heterozygous for the 1-bp deletion might carry an alternate loss-of-function variant in *EDNRB1*
on the opposite chromosome. To test this hypothesis, we sequenced the coding regions and
adjacent splice sites of *EDNRB1* in one of these animals. We observed that this animal was
heterozygous for a 17-bp deletion in the fourth coding region of *EDNRB1*
(OP589186:c.1747_1763del, Fig 3B). This deletion was located on the homologous
chromosome compared to the 1-bp deletion (S1 Fig). The 17-bp deletion introduces a frameshift
in the transcript, which removes or alters two of the seven transmembrane helices of the
EDNRB1 protein (Fig 3F). The 17-bp deletion thus represents a second putative loss-of-function
allele of *EDNRB1*.

To assess the contribution of the 17-bp deletion to phenotypes attributed to the
yellowbelly allele, we genotyped all animals in our panel for the 17-bp deletion. We found that
the 17-bp deletion was present exclusively in the nine *yellowbelly* homozygotes heterozygous
for the 1-bp deletion (Fig 3C). In each case, the 17-bp deletion was carried on the opposite
chromosome compared to the 1-bp deletion (S2 Table). These results show that every animal
described as a *yellowbelly* homozygote was either homozygous for the 1-bp deletion or
compound heterozygous for the 1-bp deletion and the 17-bp deletion (Fig 3D). (The absence of
animals homozygous for the 17-bp deletion may simply reflect the low frequency of this allele in
our sample.) These data suggest that the allele known to breeders as *yellowbelly* represents the
collection of two molecular alleles of *EDNRB1*: a 1-bp deletion and a 17-bp deletion. We
propose that these deletions produce the same phenotype and were therefore not recognized
by breeders as distinct.

The *spark* and *specter* alleles are associated with *EDNRB1* missense variants

Animals described as carrying the *spark* or *specter* alleles of the Yellowbelly series show
a more moderate loss of coloration than animals described as *yellowbelly* homozygotes (Fig
1B). We therefore predicted that the *spark* and *specter* alleles might represent variants in
EDNRB1 that reduce gene function to a lesser degree than do the 1-bp and 17-bp deletions. To
test this hypothesis, we sequenced the coding regions and adjacent splice sites of *EDNRB1* in
two animals: one animal described as a *spark/yellowbelly* compound heterozygote and one
animal described as a *specter* homozygote. (Our sample did not include any animals described
as *spark* homozygotes.) We observed that the *spark/yellowbelly* animal was heterozygous for a
missense variant in the first coding region of *EDNRB1* (OP589186:c.G481C). This variant leads
to a leucine-to-phenylalanine exchange at position 152 of the protein, termed hereafter L152F
(Fig 4C). This animal was also heterozygous for the 1-bp deletion associated with the
yellowbelly allele, as expected. We observed that the *specter* homozygote was homozygous for
a missense variant in the fifth coding region of *EDNRB1* (OP589186:c.G2601C). This variant
leads to an arginine-to-proline exchange at position 315 of the protein, termed hereafter R315P
(Fig 4C). No other coding or splice-site variants were observed in either animal. Both missense
variants affected conserved sites of the EDNRB1 protein, suggesting that they likely disrupt
protein function. Position 152, which is located in a transmembrane helix (Fig 4E), is conserved
as a leucine or isoleucine across species (Fig 4G). Position 315, which is located in an
intracellular loop (Fig 4F), is conserved as an arginine across species (Fig 4G). We conclude
that *L152F* and *R315P* are candidates for the *spark* and *specter* alleles, respectively.

To test whether *L152F* and *R315P* were associated with phenotypes attributed to the
spark and *specter* alleles, we genotyped these variants in the following panel of animals: 21
additional animals described as *spark/yellowbelly* compound heterozygotes; two additional
animals described as *specter* homozygotes; 27 animals described as *specter/yellowbelly*
compound heterozygotes; and 48 Non-Yellowbelly animals. Animals were also genotyped for
the 1-bp and 17-bp deletions associated with the *yellowbelly* allele. We observed that the

missense variants were nearly perfectly associated with the *spark* and *specter* alleles: (i) both
animals described as *specter* homozygotes were homozygous for *R315P*, (ii) all but one animal
described as *spark/yellowbelly* was heterozygous for *L152F* and for one of the deletions, and
(iii) all but one animal described as *specter/yellowbelly* was heterozygous for *R315P* and for one
of the deletions (Fig 4D). The exceptions were that one animal described as *spark/yellowbelly*
carried *R315P*, rather than *L152F*, and one animal described as *specter/yellowbelly* carried
L152F, rather than *R315P* (Fig 4D). These exceptions may represent mis-labeling by owners,
given the phenotypic similarity between the *spark* and *specter* alleles (see Discussion). These
data largely support the hypothesis that *L152F* represents the *spark* allele and *R315P*
represents the *specter* allele.

The *gravel* and *asphalt* alleles are associated with variants in *EDNRB1* splice donors

Animals described as carrying the *gravel* or *asphalt* alleles of the Yellowbelly series
show the mildest loss of coloration of all morphs in the series (Fig 1B). We therefore predicted
that the *gravel* and *asphalt* alleles might represent variants in *EDNRB1* with mild effects on
gene function. To test this hypothesis, we sequenced the coding regions and adjacent splice
sites of *EDNRB1* in one animal described as a *gravel* homozygote and one animal described as
an *asphalt* homozygote. We observed that both animals were homozygous for synonymous G-
to-A nucleotide substitutions in the -1 positions of splice donors. For the *gravel* homozygote, the
substitution was located in the splice donor for the first intron (OP589186:c.G499A, Fig 5C). For
the *asphalt* homozygote, the substitution was located in the splice donor for the sixth intron
(OP589186:c.G3118A, Fig 5C). No coding variants and no other splice-site variants were
observed in either animal. The -1 position of splice donors contributes to splicing efficiency
across eukaryotes (Cartegni *et al.* 2002), and this position is a guanine in ~80% of splice donors
in vertebrate genes (Ma *et al.* 2015). In *EDNRB1*, the -1 positions of the first and sixth splice
donors are conserved as guanines across species (Fig 5C), suggesting that guanines at these
sites contribute to gene function. We conclude that the splice-donor substitutions identified
herein are candidates for the *gravel* and *asphalt* alleles.
To test whether the G-to-A substitutions in splice donors were associated with
phenotypes attributed to the *gravel* and *asphalt* alleles, we genotyped these variants in the
following panel of animals: six additional animals described as *gravel* homozygotes; three
additional animals described as *asphalt* homozygotes; 22 animals described as
gravel/yellowbelly compound heterozygotes; 23 animals described as *asphalt/yellowbelly*
compound heterozygotes; and 48 Non-Yellowbelly animals. These animals were also
genotyped for the 1-bp and 17-bp deletions associated with the *yellowbelly* allele. We observed
that the G-to-A substitutions in splice donors were nearly perfectly associated with the *gravel*
and *asphalt* alleles: (i) all animals described as *gravel* or *asphalt* homozygotes were
homozygous for the substitutions in the first or sixth splice donors, respectively, (ii) all animals
described as *gravel/yellowbelly* were heterozygous for the substitution in the first splice donor
and for one of the deletions, and (iii) all but two animals described as *asphalt/yellowbelly* were
heterozygous for the substitution in the sixth splice donor and for one of the deletions (Fig 5D).
The exceptions were that two animals described as *asphalt/yellowbelly* carried the substitution
in the first splice donor, rather than in the sixth splice donor (Fig 5D). These exceptions may
represent mis-labeling by owners, given the phenotypic similarity between the *gravel* and
asphalt alleles (see Discussion). These data largely support the hypothesis that the G-to-A
substitution in the first splice donor represents the *gravel* allele and the G-to-A substitution in the
sixth splice donor represents the *asphalt* allele.

Reliability of heterozygote descriptions

Ball pythons described as heterozygous for alleles in the Yellowbelly series sell for higher prices than wildtype animals. This price difference provides an incentive for sellers to identify such animals and label them as heterozygotes. Yet heterozygotes for alleles in the Yellowbelly series are difficult to distinguish from wildtype. Heterozygotes have subtly lighter coloration and altered patterning in the underbelly (Figs 1E and 1F), but this phenotype is variable and overlaps with the range of phenotypes observed among wildtype animals. Thus, animals described as heterozygous for alleles in the Yellowbelly series are viewed with a degree of skepticism. Many believe that some of these animals are mis-labeled, perhaps for monetary gain by sellers.

Our discovery of association between alleles in the Yellowbelly series and variants in *EDNRB1* allowed us to test the reliability of heterozygote descriptions. To do so, we genotyped 130 animals described as *yellowbelly* heterozygotes for the 1-bp and 17-bp deletions in *EDNRB1*. These animals were collected from a total of 68 pet owners across the United States and therefore represent a broad sampling of the pet population. We focused on the *yellowbelly* allele, rather than other alleles in the Yellowbelly series, because the *yellowbelly* allele is described as producing the strongest phenotype in heterozygotes. We observed that 98 of the 130 animals in our sample were heterozygous for the 1-bp or 17-bp deletions in *EDNRB1* (Fig 3C). The animals lacking these deletions were further tested to determine whether they carried any of the other variants associated with alleles in the Yellowbelly series—i.e., *L152F* (*spark*), *R315P* (*specter*), and the G-to-A substitutions in splice donors (*gravel* and *asphalt*). One animal was found to be heterozygous for *L152F* (S2 Table). The other animals did not carry any of these variants (S2 Table). Thus, of the 130 animals described as *yellowbelly* heterozygotes, 98 were correctly described, one was a *spark* homozygote, and the remainder carried only putatively wildtype copies of *EDNRB1*. These findings confirm the skepticism surrounding heterozygote descriptions for alleles in the Yellowbelly series. They suggest that approximately one in four animals described as a *yellowbelly* heterozygote is mis-labeled.

Discussion

The goal of the current study was to identify the genetic cause of phenotypes in the Yellowbelly series of ball pythons. These phenotypes were characterized by a loss of coloration and coincident changes in patterning, ranging from severe (leucism) to moderate (dorsal striping) to mild (subtle changes in patterning compared to wildtype) (Fig 1). Through careful choice of candidate genes and a bit of luck, we discovered that these phenotypes were associated with six putative loss-of-function variants in the gene encoding endothelin receptor EDNRB1 (Fig 6). Leucism was associated with frameshift variants (Fig 3). Dorsal striping was associated with missense variants (Fig 4). Subtle changes in patterning were associated with splice-donor variants (Fig 5). To our knowledge, these variants represent the first description of genetic variants affecting endothelin signaling in a non-avian reptile.

Causality and the candidate-gene approach

We propose that the *EDNRB1* variants described herein are causal for phenotypes in the Yellowbelly series. This idea is supported by a correlation between the severity of phenotypes in the Yellowbelly series and the severity of genetic lesions in *EDNRB1* (Fig 6). The most severe loss of coloration in the Yellowbelly series—leucism—was associated with frameshift variants, which plausibly represent null alleles (Fig 3). Leucism is characteristic of an absence of chromatophores (e.g. Ullate-Agote and Tzika 2021), and the association between leucism and *EDNRB1* in ball pythons is consistent with *EDNRB1* controlling chromatophore development in zebrafish (Parichy *et al.* 2000) and mammals (Hosoda *et al.* 1994; Ceccherini *et al.* 1995; Gariepy *et al.* 1996; Yang *et al.* 1998; Pingault *et al.* 2010), and with loss of the

endothelin ligand EDN3 causing leucism in axolotl (Woodcock *et al.* 2017). We propose that
leucism represents the null phenotype for *EDNRB1* in ball pythons.
The next most severe phenotype in the Yellowbelly series was dorsal striping. This
phenotype was associated with missense variants at conserved sites of the EDNRB1 protein
(Fig 4). We propose that these variants represent hypo-morphic alleles. These missense
variants might reduce EDNRB1 protein function via effects on folding or interactions with co-
factors, among other possibilities. Dorsal striping in ball pythons is reminiscent of phenotypes in
zebrafish caused by iridophores and melanophores being reduced in number and remaining
clustered near the horizontal midline, which represents the site of iridophore emergence into the
skin (Fig 2A-B) (Parichy *et al.* 2000; Lopes *et al.* 2008; Patterson and Parichy 2013; Frohnhöfer
et al. 2013; Mo *et al.* 2017; Spiewak *et al.* 2018). We propose that stripe formation in ball
pythons is analogous to this phenotype in zebrafish, and that stripes in ball pythons are caused
by chromatophores being moderately reduced in number and remaining clustered near the
dorsal ridge (Fig 2E).
The mildest phenotype in the Yellowbelly series was altered patterning. This phenotype
was associated with G-to-A nucleotide substitutions in the -1 positions of *EDNRB1* splice donors
(Fig 5). The -1 position of splice donors contributes to splicing efficiency, albeit less so than
splice-donor positions located within introns (Cartegni *et al.* 2002). We propose that the splice-
donor substitutions in *EDNRB1* reduce the fidelity of splicing and thereby reduce EDNRB1
protein levels, leading to a mild reduction in chromatophores and consequence alterations in
patterning (Fig 2D). The idea that mild reductions in chromatophores can produce noticeable
changes in patterning is consistent with the idea that patterns arise via a complex network of
cell-to-cell interactions (Singh and Nuesslein-Volhard 2015; Patterson and Parichy 2019), which
are sensitive to cell numbers (Spiewak *et al.* 2018) and other parameters (Watanabe and Kondo
2012; Kaelin *et al.* 2012, 2021; McGuirl *et al.* 2020).
The scope of our study was limited to two genes: *EDNRB1* and *EDN3*. Thus, we cannot
formally exclude the possibility that variants in other regions of the genome contribute to
phenotypes in the Yellowbelly series. However, we view this possibility as unlikely, given that six
independent haplotypes of *EDNRB1* were associated with phenotypes in the Yellowbelly series,
and that the severity of these phenotypes correlated with the severity of genetic lesions in
EDNRB1.

Pros and cons of community-sourced sampling

Our study relied on pet samples recruited from the community, rather than samples from
a single pedigree. An advantage of this approach was that it increased the genetic diversity of
our sample. An example of this diversity was the discovery that the allele known to breeders as
yellowbelly represents two distinct molecular alleles of *EDNRB1*—a 1-bp deletion and a 17-bp
deletion (Fig 3). The 17-bp deletion was approximately 16-fold less common in our sample than
the 1-bp deletion and might have been plausibly missed by our study, had we focused on a
single pedigree. The phenomenon of multiple molecular alleles corresponding to a single
breeder-defined allele is not unique to the Yellowbelly series and has been reported for two
other color morphs in ball pythons—the Albino color morph and the Ultramel color morph
(Brown *et al.* 2022). This phenomenon highlights the utility of community-sourced sampling in
ball pythons and shows that the pet population is more diverse than previously recognized.
A disadvantage of using pet samples sourced from the community is that genotype
descriptions provided by owners come with a degree of uncertainty. Certain combinations of
alleles in the Yellowbelly series produce similar phenotypes (e.g. *spark* vs. *specter* and *gravel*
vs. *asphalt*). Crosses involving these alleles therefore generate progeny whose genotypes are
uncertain. Breeders typically label such animals according to their best guess of the animal's
genotype. This practice leads to mis-labeling of genotypes by owners and likely explains the
apparent mix-ups of animals in our study: one *spark/yellowbelly* animal was mis-labeled by its

owner as *specter/yellowbelly* and vice versa (Fig 4D); two *gravel/yellowbelly* animals were mis-labeled by their owners as *asphalt/yellowbelly* (Fig 5D); and one *spark* heterozygote was mis-labeled by its owner as a *yellowbelly* heterozygote (Fig 3C). An additional cause of mis-labeling is that breeders are incentivized to label animals as heterozygous for alleles in the Yellowbelly series, even when the phenotype of heterozygotes is difficult to recognize (Figs 1E and 1F). This incentive is driven by heterozygotes selling for higher prices than wildtype animals and likely explains why approximately one in four animals described as *yellowbelly* heterozygotes carried only putatively wildtype copies of *EDNRB1* (Fig 3C). We propose that mis-labeling can be reduced in the future by genotyping animals for the *EDNRB1* variants reported in our study.

Evolution of endothelin signaling and stripe formation

The phenotypes associated with *EDNRB1* variants in ball pythons suggest that endothelin signaling may be required for the development of all three types of chromatophores in this species. Endothelin signaling in snakes may therefore play a role similar to endothelin signaling in axolotl, where the endothelin ligand EDN3 is thought to direct the development of all three types of chromatophores (Woodcock *et al.* 2017). Similar regulation has been described for mammals and birds, where endothelin signals regulate the development of melanophores and other neural crest-derived cells (Saldana-Caboverde and Kos 2010; Bondurand *et al.* 2018). These species contrast with zebrafish, where endothelin signaling directly promotes the development of iridophores, but is dispensable for xanthophore development and plays only an indirect role for melanophores (Patterson and Parichy 2013; Frohnhöfer *et al.* 2013; Patterson *et al.* 2014). These observations have led to the proposal that endothelin signaling in vertebrates was ancestrally required for all three types of chromatophores and became specialized for iridophores in teleost fish (Spiewak *et al.* 2018). Our findings support this hypothesis by adding ball pythons to the list of species in which endothelin signaling likely regulates the development of multiple types of chromatophores.

Dorsal, longitudinal stripes are not unique to ball python morphs in the Yellowbelly series. Other striped morphs in ball pythons include the Genetic Stripe morph, the Champagne morph, and some morphs in the Blue-Eyed Leucistic series (McCurley 2005; Broghammer 2019). Outside ball pythons, stripes occur naturally in many snake species, typically in species that escape predation via rapid flight (Allen *et al.* 2013). Stripes have arisen multiple times within the snake lineage (Wolf and Werner 1994), and a few species are naturally polymorphic for striped and non-striped forms (Zweifel 1981; Brodie 1992; King 2003; Westphal and Morgan 2010; Kuriyama *et al.* 2013; Murakami *et al.* 2014). Whether stripe formation in these cases involves changes to endothelin signaling or chromatophore number remains to be determined. Our study suggests that dorsal, longitudinal stripes may be easily accessible to the snake body form, via genetic changes that alter simple developmental parameters in the neural crest, such as cell number. Candidate genes for controlling these parameters include other genes in the endothelin pathway (Braasch *et al.* 2009) and genes responsible for specifying cell fates in the neural crest (Parichy and Spiewak 2015; Mort *et al.* 2015).

Methods

Recruitment of ball python sheds

Ball python sheds were recruited from pet owners and breeders by placing announcements in Twitter, Reddit, Instagram, and Facebook, and by contacting sellers having active listings on Morph Market (www.morphmarket.com). Contributors were instructed to allow sheds to dry (if wet) and to ship sheds via standard first-class mail. Contributors sending multiple sheds were instructed to package sheds individually in plastic bags during shipping. Contributors were not provided monetary compensation for sheds, although some contributors

were given pre-paid shipping envelopes to cover shipping costs. Upon receipt, sheds were
stored at -20°C to kill any insect larvae infesting the sheds.
We attempted to maximize genetic diversity within each category of morph by excluding
animals described as siblings, parents, or offspring of another animal in our sample. However,
we cannot exclude the possibility that some animals in our sample may have been close
relatives, given that full pedigree information was not available for most animals. The idea that
animals in our sample were derived from multiple lineages is supported by our discovery of
multiple molecular variants corresponding to the allele known to breeders as *yellowbelly*. Owner
codes and US states of residence for each animal are provided in S2 Table.
The total set of animals used in our study were 70 animals described as *yellowbelly*
homozygotes, 130 animals described as *yellowbelly* heterozygotes, 22 animals described as
spark/yellowbelly, three animals described as *specter* homozygotes, 27 animals described as
specter/yellowbelly, seven animals described as *gravel* homozygotes, 22 animals described as
gravel/yellowbelly, four animals described as *asphalt* homozygotes, 23 animals described as
asphalt/yellowbelly, and 48 animals described as having wildtype coloration or belonging to
morphs other than those in the Yellowbelly series (Non-Yellowbelly animals)

Performing experiments in an undergraduate laboratory course

The majority of experiments and analyses described in this study were performed by
undergraduate students as part of a laboratory course at Eastern Michigan University
(BIO306W). This practice required that our experimental design rely on simple techniques,
namely PCR, restriction digests, and Sanger sequencing. To avoid student errors in these
techniques, we implemented the following precautions. First, students never handled more than
one shed skin at the same time. Second, students performed negative and positive control
reactions for all PCR amplifications and restriction digests. Data from students having incorrect
controls were excluded from analysis. Third, all sequence analyses were performed
independently by three or more students. When the results of all students did not all agree,
sequences were re-analyzed by the instructor (HSS). Fourth, when we encountered an animal
whose molecular genotype did not match the breeder description (e.g. the single animal labeled
spark/yellowbelly whose genotype matched *specter/yellowbelly* and vice versa—see Results
and Discussion), DNA from the animal was re-extracted by the instructor (HSS), and the
molecular genotype were determined a second time. This precaution was taken to guard
against mix-ups by students, but it proved unnecessary because the second round of genotypes
always matched the first

DNA extraction

Sheds containing visible dirt or debris were rinsed in tap water until clean. Sheds were
air dried and lysed overnight at ~60˚C in ~1 ml lysis buffer (100 mM Tris-HCl pH 8.0, 100 mM
EDTA, 2% sodium dodecyl sulfate, 3 mM CaCl2, 2 mg/ml Proteinase K) per palm-sized piece of
shed. Lysate was separated from visible fragments of undigested shed and further cleared by
centrifugation at 13,000 x g for 2 min. To precipitate protein, ammonium acetate was added to
supernatant to a final concentration of 1.875 – 2.5 M. Samples were incubated on ice for 5 min
and centrifuged at 13,000 x g for 3 min at 4˚C. Supernatant was mixed with an equal volume of
magnetic bead mixture (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 1.6 M NaCl, 0.2% Tween-20,
11% polyethylene glycol, 0.04% washed SpeedBeads [Sigma #GE45152105050250]) and
shaken for 5 min. Beads were separated from supernatant using a magnet, washed twice in 0.2
ml 70% ethanol for 2 min, and air dried for ~1 min. DNA was eluted from beads in TE buffer (10
mM Tris-HCl pH 8.0, 1 mM EDTA) at 65°C for >5 min.

Primer design and PCR

Primers were designed against the genome of Burmese python, the closest relative of ball python for which genome sequence was available (Castoe *et al.* 2013). Annotations for Burmese python *EDNRB1* were XM_007436628 (mRNA) and XP_007436690 (protein). Annotations for Burmese python *EDN3* were XM_025169570 (mRNA) and XP_007432959 (protein). These annotations were manually examined for errors, by compared gene structure across vertebrates. For both genes, annotations were determined to be correct, based on the observation that gene structure matched gene structure in other vertebrates, including in species with higher quality annotations, such as mouse, chicken, and zebrafish.

Primers were designed using Primer3 (Untergasser *et al.* 2012), using default parameters and a target annealing temperature of 60°C. Amplification was performed using an annealing temperature of 57°C, to allow for occasional divergence between ball python and Burmese python genomic sequences. Coding regions of *EDNRB1* and *EDN3* were amplified and sequenced using primers given in S1 Table.

PCR was performed using OneTaq polymerase (NEB #M0480). Reactions consisted of 1X OneTaq Standard Reaction Buffer, 200 μM dNTPs, 0.2 μM of each primer, and 0.025 U/μl OneTaq polymerase. Thermocycling conditions were as follows: 94°C for 2 min; 30-35 cycles of 94°C for 30 sec, 57°C for 30 sec, and 68°C for 1-2 min; and 68°C for 5 min. Reactions used 10-100 ng template DNA per 20 μl volume.

Sanger sequencing

PCR products were purified for Sanger sequencing using magnetic beads. PCR reactions were mixed with three volumes of magnetic-bead mixture (10 mM Tris-HCl pH 8.0, 1 mM EDTA, 1.6-2.5 M NaCl, 0.2% Tween-20, 11-20% polyethylene glycol, 0.04% washed SpeedBeads [Sigma #GE45152105050250]) and agitated for 5 min. Beads were separated from supernatant using a magnet, washed twice in 0.2 ml 80% ethanol for >30 sec, and air-dried for 30 sec. PCR products were eluted from beads in 10 mM Tris-HCl pH 8.0 for >3 min at 65°C. Sanger sequencing was performed by Eton Bioscience Inc (etonbio.com). Sequencing primers are provided in S1 Table.

Protein sequence alignment

Protein sequences were aligned using Clustal Omega (Sievers *et al.* 2011), using default parameters. Accession numbers for protein sequences were as follows: elephant shark EDNRB1 (XP_007900764) and EDNRA (XP_007897347); zebrafish EDNRBb (XP_688565) and EDNRA (NP_001092915); human EDNRB1 (NP_000106) and EDNRA (NP_001948); chicken EDNRB1 (XP_015133631) and EDNRA (NP_989450); tropical clawed frog EDNRB1 (XP_012812722) and EDNRA (NP_001072275); painted turtle EDNRB1 (XP_005311368); anole lizard EDNRB1 (XP_003223432) and EDNRA (XP_003221737); Burmese python EDNRB1 (XP_007436690) and EDNRA (XP_007426479).

Genotyping assays

The 1-bp and 17-bp deletions in *EDNRB1* (*yellowbelly*) were genotyped by amplifying the fourth coding region of *EDNRB1* via PCR and digesting the PCR products with the restriction enzyme *Bsr*I (New England BioLabs, #R0527S). Both deletions eliminate cut sites for *Bsr*I and can thus be genotyped simultaneously (S1 Fig). Primers for amplification were p257F (5'-GGT TCC CAG TCT CTG CAC AT-3') and p257R (5'-TTC AAC CAA CGC ACA TCA TT-3'). Digestions consisted of 1X NEBuffer™ r3.1 and 0.05 U/μl *Bsr*I. Digestions were performed at 65°C for at least 2 hrs.

Missense variant *R315P* (*specter*) was genotyped by amplifying the fifth coding region of *EDNRB1* via PCR and digesting the PCR products with the restriction enzyme *Sma*I (New England BioLabs, #R0141S). The *R315P* variant creates a cut site for *Sma*I (S2 Fig). Primers

for amplification were p258F (5'-ATG GAG TGG ACA GGA TGG AA-3') and p258R (5'-CCA GCT AGG TGG GGT AGA CA-3'). Digestions consisted of 1X rCutSmart™ Buffer and 0.05-0.1 U/µl *Sma*I. Digestions were performed at 25°C for at least 2 hrs.

Missense variant *L152F* (*spark*) and the G-to-A substitution at the first splice donor (*gravel*) were genotyped by amplifying a fragment containing the first coding region of *EDNRB1* using primers p381F (5'-CAC CAT AAT GCT TAA CAC ACA CAA-3') and p381R (5'-TCA TAG CAA TGT GAT AAA CCC ACT-3'). PCR products were sequenced with p381F (S3 Fig).

The G-to-A substitution at the sixth splice donor (*asphalt*) was genotyped by amplifying a fragment containing the sixth and seventh coding regions of *EDNRB1* using primers p259F (5'-GGC AGG AAA ACT GCT CGA TA-3') and p260R (5'-CAA ACC AAA GTC CCA GCA TC-3'). PCR products were sequenced with p259F (S3 Fig).

Acknowledgements

We thank Bob Winning, Katy Greenwald, and David Kass for comments on the manuscript and the Educational Course Support program of New England BioLabs for reagents used in undergraduate teaching labs.

Most of the data in this study were collected by undergraduates enrolled in a laboratory course at Eastern Michigan University. These students constitute the BIO306W Consortium. These students were Bayan Abdeljalil, Garrett Bailey, John (Teddy) Belman, Keith Camac, Aaron Ellis, Amber Fatima, Delaney Garcia, Shannon Gregory, Amber Haley, Chloe Harrison, Carly Kosanovich, Corey Melcher, Lindsey Miracle, Natalia Pineda, Catherine Redding, Audrey Salsido, Caitlin Satler, Jake Sealy, Ananya Shukla, Hannah Strasser, Nehul Tanna, Evan Veenhuis, Syed Wasiuddin, Anna Watson, Madyson Weaver, Raymond Wells, and Kathleen Weymouth.

Shed skins were contributed by the following individuals: Abby Kruse, Adam and Nicole Schmid, Adam of Proper Royals, Allen Gage, Alycia Butler, Amanda Hall, Amanda Yereance, Ammar Hanif, Andelyn Czajka, Andrew Lyons, Andy Rose, Anthony of Anthill Python, AR Walters, Bankrupt Reptiles, Barbara Penyak, Big D, Brent McKelvey, Bryan Rivera, Charles Wilson, Chelsey Wheeler, Chiron Graves, Chris Hernan, Chris Shelly, Chun Ku, CPJ Reptiles, Cynthia Jones, Dale Porcher, Dave Dunn of Ball in Hand Pythons, David Burstein, Denise Jones, E & L's Livin Art Morphs 2020, Epic Vibrant Balls, Eric Chung, Everything's Better Orange, FNA Balls, Francis X. DiNino Jr, Gavin Costello, Greg and Kathee of Royally Morphadellic, Harry Roman, Jacob Rodenhausen, Jaden Christensen, Jake and Ashley Brewster, Jake Jewis, James and Angie Rehus of JAR Pythons, Jamie Palazzo, Jason Roberts, Jeff Kearns, Jeff Linton, Jeromy Shaffer, Joe and Wendy Ducos, Joe Myers, John White, Jon Bidwell, Jon Braga, Jon Godek, Justin Orbach of Just Incrediballs, Kai Li, Kandie Tucker, Kassandra Royer, Kelsi Greene, Kenneth Evans, Korie Dy, KT Creations, Laura Carter, Lauren Glynn, Leah Alexander, Leviathan Snakes, Lindsey VanOrman, Loren Morales, Lynnet Melton, Margaret Lentz, Maryann Barbon, Matt Huck, Michael Cole, Michael Nino, Michael Osmun, Miranda Joy Turpin, Morgan Evans, Nate Fitzpatrick of Baltimore Balls, Nathan Granoff, Nick Deinhart, Nicole Gallagher, OkieMom Balls, Ozzy Maldonado, Pedro Alequin of Imperial Morphs, Peter Holden, Phil Phillips, Phil Snyder, Randy Remington, Redstrom Reptiles, Reggie's Urban Jungle, Roger Gray, Royal Black Balls, Ryan Boyd and Brittney Delacruz, Sandra Frye, Sean Strait of The Morph Lab, Sergio McDole, Shomai Sanford, Steven Nordahl, Tammy Hutchinson of Hutches Clutches, The Wagner's of Texas, Tia Vogel, Tom Harbin, Troy Stallings, Tyler Grant, Wes Mosher, Wilbanks Reptiles, Zac Parpart, and several anonymous contributors.

Author Contributions

Conceived and designed the experiments: HSS. Performed the experiments: UMD IL RLT BS TBC HSS. Analyzed the data: UMD IL RLT BS TBC HSS. Contributed reagents or materials: CWG HSS. Wrote the paper: HSS.

References

Allen, W. L., R. Baddeley, N. E. Scott-Samuel, and I. C. Cuthill, 2013 The evolution and function of pattern diversity in snakes. Behav. Ecol. 24: 1237–1250.

Ascarrunz, E., and M. R. Sanchez-Villagra, 2022 The macroevolutionary and developmental evolution of the turtle carapacial scutes. Vertebr. Zool. 72: 29–46.

Baynash, A. G., K. Hosoda, A. Giaid, J. A. Richardson, N. Emoto *et al.*, 1994 Interaction of endothelin-3 with endothelin-B receptor is essential for development of epidermal melanocytes and enteric neurons. Cell 79: 1277–1285.

Besmer, P., K. Manova, R. Duttlinger, E. J. Huang, A. Packer *et al.*, 1993 The kit-ligand (steel factor) and its receptor c-kit/W: pleiotropic roles in gametogenesis and melanogenesis. Dev Suppl 125–137.

Bondurand, N., S. Dufour, and V. Pingault, 2018 News from the endothelin-3/EDNRB signaling pathway: Role during enteric nervous system development and involvement in neural crest-associated disorders. Dev. Biol. 444: S156–S169.

Braasch, I., J.-N. Volff, and M. Schartl, 2009 The endothelin system: evolution of vertebrate-specific ligand-receptor interactions by three rounds of genome duplication. Mol Biol Evol 26: 783–799.

Brejcha, J., J. V. Bataller, Z. Bosáková, J. Geryk, M. Havlíková *et al.*, 2019 Body coloration and mechanisms of colour production in Archelosauria: the case of deirocheline turtles. R Soc Open Sci 6: 190319.

Brodie, E. D., 1992 CORRELATIONAL SELECTION FOR COLOR PATTERN AND ANTIPREDATOR BEHAVIOR IN THE GARTER SNAKE THAMNOPHIS ORDINOIDES. Evolution 46: 1284–1298.

Broghammer, S., 2019 *Python regius: Atlas of Colour Morphs Keeping and Breeding*. NTV Natur und Tier-Verlag.

Brown, A. R., K. Comai, D. Mannino, H. McCullough, Y. Donekal *et al.*, 2022 A community-science approach identifies genetic variants associated with three color morphs in ball pythons (Python regius). bioRxiv 2021.05.19.444867.

Camargo-Sosa, K., S. Colanesi, J. Müller, S. Schulte-Merker, D. Stemple *et al.*, 2019 Endothelin receptor Aa regulates proliferation and differentiation of Erb-dependent pigment progenitors in zebrafish. PLoS Genet 15: e1007941.

Cartegni, L., S. L. Chew, and A. R. Krainer, 2002 Listening to silence and understanding nonsense: exonic mutations that affect splicing. Nat Rev Genet 3: 285–298.

Castoe, T. A., A. P. J. de Koning, K. T. Hall, D. C. Card, D. R. Schield *et al.*, 2013 The Burmese python genome reveals the molecular basis for extreme adaptation in snakes. Proc. Natl. Acad. Sci. U. S. A. 110: 20645–20650.

Ceccherini, I., A. L. Zhang, I. Matera, G. Yang, M. Devoto *et al.*, 1995 Interstitial deletion of the endothelin-B receptor gene in the spotting lethal (sl) rat. Hum Mol Genet 4: 2089–2096.

Cobaleda, C., M. Pérez-Caro, C. Vicente-Dueñas, and I. Sánchez-García, 2007 Function of the zinc-finger transcription factor SNAI2 in cancer and development. Annu Rev Genet 41: 41–61.

Cuthill, I. C., W. L. Allen, K. Arbuckle, B. Caspers, G. Chaplin *et al.*, 2017 The biology of color. Science 357: eaan0221.

Dorshorst, B., A.-M. Molin, C.-J. Rubin, A. M. Johansson, L. Strömstedt *et al.*, 2011 A complex genomic rearrangement involving the endothelin 3 locus causes dermal hyperpigmentation in the chicken. PLoS Genet 7: e1002412.

Dupin, E., C. Glavieux, P. Vaigot, and N. M. Le Douarin, 2000 Endothelin 3 induces the reversion of melanocytes to glia through a neural crest-derived glial-melanocytic progenitor. Proc Natl Acad Sci U S A 97: 7882–7887.

Frohnhöfer, H. G., J. Krauss, H.-M. Maischein, and C. Nüsslein-Volhard, 2013 Iridophores and their interactions with other chromatophores are required for stripe formation in zebrafish. Development 140: 2997–3007.

Garcia-Elfring, A., H. L. Roffey, A. P. Hendry, and R. D. H. Barrett, 2020 A nonsense mutation in TFEC is the likely cause of the recessive piebald phenotype in ball pythons (Python regius):, 2020.10.30.362970 p.

Gariepy, C. E., D. T. Cass, and M. Yanagisawa, 1996 Null mutation of endothelin receptor type B gene in spotting lethal rats causes aganglionic megacolon and white coat color. Proc Natl Acad Sci U S A 93: 867–872.

Haupaix, N., and M. Manceau, 2020 The embryonic origin of periodic color patterns. Developmental Biology 460: 70–76.

Hosoda, K., R. E. Hammer, J. A. Richardson, A. G. Baynash, J. C. Cheung *et al.*, 1994 Targeted and natural (piebald-lethal) mutations of endothelin-B receptor gene produce megacolon associated with spotted coat color in mice. Cell 79: 1267–1276.

Kaelin, C. B., K. A. McGowan, and G. S. Barsh, 2021 Developmental genetics of color pattern establishment in cats. Nat Commun 12: 5127.

Kaelin, C. B., X. Xu, L. Z. Hong, V. A. David, K. A. McGowan *et al.*, 2012 Specifying and sustaining pigmentation patterns in domestic and wild cats. Science 337: 1536–1541.

Kelsh, R. N., 2006 Sorting out Sox10 functions in neural crest development. Bioessays 28: 788–798.

Kelsh, R. N., M. L. Harris, S. Colanesi, and C. A. Erickson, 2009 Stripes and belly-spots—A review of pigment cell morphogenesis in vertebrates. Seminars in Cell & Developmental Biology 20: 90–104.

King, R. B., 2003 Mendelian Inheritance of Melanism in the Garter Snake Thamnophis sirtalis. Herpetologica 59: 484–489.

Kinoshita, K., T. Akiyama, M. Mizutani, A. Shinomiya, A. Ishikawa *et al.*, 2014 Endothelin receptor B2 (EDNRB2) is responsible for the tyrosinase-independent recessive white (mo(w)) and mottled (mo) plumage phenotypes in the chicken. PLoS One 9: e86361.

Kondo, S., and H. Shirota, 2009 Theoretical analysis of mechanisms that generate the pigmentation pattern of animals. Seminars in Cell & Developmental Biology 20: 82–89.

Krauss, J., H. G. Frohnhöfer, B. Walderich, H.-M. Maischein, C. Weiler *et al.*, 2014 Endothelin signalling in iridophore development and stripe pattern formation of zebrafish. Biol Open 3: 503–509.

Kuriyama, T., H. Misawa, K. Miyaji, M. Sugimoto, and M. Hasegawa, 2013 Pigment cell mechanisms underlying dorsal color-pattern polymorphism in the Japanese four-lined snake. J Morphol 274: 1353–1364.

Kuriyama, T., A. Murakami, M. Brandley, and M. Hasegawa, 2020 Blue, Black, and Stripes: Evolution and Development of Color Production and Pattern Formation in Lizards and Snakes. Frontiers in Ecology and Evolution 8:.

Li, L., D. Li, L. Liu, S. Li, Y. Feng *et al.*, 2015 Endothelin Receptor B2 (EDNRB2) Gene Is Associated with Spot Plumage Pattern in Domestic Ducks (Anas platyrhynchos). PLoS One 10: e0125883.

Lopes, S. S., X. Yang, J. Müller, T. J. Carney, A. R. McAdow *et al.*, 2008 Leukocyte Tyrosine Kinase Functions in Pigment Cell Development. PLOS Genetics 4: e1000026.

Lühken, G., K. Fleck, A. Pauciullo, M. Huisinga, and G. Erhardt, 2012 Familiar
Hypopigmentation Syndrome in Sheep Associated with Homozygous Deletion of the
Entire Endothelin Type-B Receptor Gene. PLOS ONE 7: e53020.
Ma, S. L., V. Vega-Warner, C. Gillies, M. G. Sampson, V. Kher *et al.*, 2015 Whole Exome
Sequencing Reveals Novel PHEX Splice Site Mutations in Patients with
Hypophosphatemic Rickets. PLOS ONE 10: e0130729.
Mallarino, R., C. Henegar, M. Mirasierra, M. Manceau, C. Schradin *et al.*, 2016 Developmental
mechanisms of stripe patterns in rodents. Nature 539: 518–523.
McCurley, K., 2005 *Complete Ball Python, A Comprehensive Guide to Care, Breeding, and*
Genetic Mutations: Kevin McCurley: 9780976733409: Amazon.com: Books. ECO /
Serpent's Tale NHBD.
McGuirl, M. R., A. Volkening, and B. Sandstede, 2020 Topological data analysis of zebrafish
patterns. Proceedings of the National Academy of Sciences 117: 5113–5124.
Mills, M. G., and L. B. Patterson, 2009 Not just black and white: pigment pattern development
and evolution in vertebrates. Semin Cell Dev Biol 20: 72–81.
Miwa, M., M. Inoue-Murayama, H. Aoki, T. Kunisada, T. Hiragaki *et al.*, 2007 Endothelin
receptor B2 (EDNRB2) is associated with the panda plumage colour mutation in
Japanese quail. Anim Genet 38: 103–108.
Mo, E. S., Q. Cheng, A. V. Reshetnyak, J. Schlessinger, and S. Nicoli, 2017 Alk and Ltk ligands
are essential for iridophore development in zebrafish mediated by the receptor tyrosine
kinase Ltk. Proc Natl Acad Sci U S A 114: 12027–12032.
Mort, R. L., I. J. Jackson, and E. E. Patton, 2015 The melanocyte lineage in development and
disease. Development 142: 620–632.
Murakami, A., M. Hasegawa, and T. Kuriyama, 2014 Identification of Juvenile Color Morphs for
Evaluating a Heredity Model of Stripe/Non-Stripe Pattern Polymorphism in the Japanese
Four-Lined Snake Elaphe quadrivirgata. jche 33: 68–74.
Murray, J. D., and M. R. Myerscough, 1991 Pigmentation pattern formation on snakes. J Theor
Biol 149: 339–360.
Olsson, M., D. Stuart-Fox, and C. Ballen, 2013 Genetics and evolution of colour patterns in
reptiles. Semin. Cell Dev. Biol. 24: 529–541.
Omasits, U., C. H. Ahrens, S. Müller, and B. Wollscheid, 2014 Protter: interactive protein feature
visualization and integration with experimental proteomic data. Bioinformatics 30: 884–
886.
Parichy, D. M., E. M. Mellgren, J. F. Rawls, S. S. Lopes, R. N. Kelsh *et al.*, 2000 Mutational
analysis of endothelin receptor b1 (rose) during neural crest and pigment pattern
development in the zebrafish Danio rerio. Dev Biol 227: 294–306.
Parichy, D. M., and J. E. Spiewak, 2015 Origins of adult pigmentation: diversity in pigment stem
cell lineages and implications for pattern evolution. Pigment Cell Melanoma Res. 28:.
Patterson, L. B., E. J. Bain, and D. M. Parichy, 2014 Pigment cell interactions and differential
xanthophore recruitment underlying zebrafish stripe reiteration and Danio pattern
evolution. Nat Commun 5: 5299.
Patterson, L. B., and D. M. Parichy, 2013 Interactions with iridophores and the tissue
environment required for patterning melanophores and xanthophores during zebrafish
adult pigment stripe formation. PLoS Genet 9: e1003561.
Patterson, L. B., and D. M. Parichy, 2019 Zebrafish Pigment Pattern Formation: Insights into the
Development and Evolution of Adult Form. Annu Rev Genet 53: 505–530.
Pingault, V., D. Ente, F. Dastot-Le Moal, M. Goossens, S. Marlin *et al.*, 2010 Review and update
of mutations causing Waardenburg syndrome. Hum Mutat 31: 391–406.
Pizzigalli, C., F. Banfi, G. F. Ficetola, M. Falaschi, M. Mangiacotti *et al.*, 2020 Eco-geographical
determinants of the evolution of ornamentation in vipers. Biol. J. Linnean Soc. 130: 345–
358.

Rosenbaum, D. M., S. G. F. Rasmussen, and B. K. Kobilka, 2009 The structure and function of
696   G-protein-coupled receptors. Nature 459: 356–363.
697   Saenko, S. V., J. Teyssier, D. van der Marel, and M. C. Milinkovitch, 2013 Precise colocalization
698   of interacting structural and pigmentary elements generates extensive color pattern
699   variation in Phelsuma lizards. BMC Biol. 11: 105.
700   Saldana-Caboverde, A., and L. Kos, 2010 Roles of endothelin signaling in melanocyte
701   development and melanoma. Pigment Cell Melanoma Res. 23: 160–170.
702   Shinomiya, A., Y. Kayashima, K. Kinoshita, M. Mizutani, T. Namikawa *et al.*, 2012 Gene
703   duplication of endothelin 3 is closely correlated with the hyperpigmentation of the internal
704   organs (Fibromelanosis) in silky chickens. Genetics 190: 627–638.
705   Sievers, F., A. Wilm, D. Dineen, T. J. Gibson, K. Karplus *et al.*, 2011 Fast, scalable generation
706   of high-quality protein multiple sequence alignments using Clustal Omega. Mol. Syst.
707   Biol. 7: 539.
708   Singh, A. P., and C. Nuesslein-Volhard, 2015 Zebrafish Stripes as a Model for Vertebrate
709   Colour Pattern Formation. Curr. Biol. 25: R81–R92.
710   Spiewak, J. E., E. J. Bain, J. Liu, K. Kou, S. L. Sturiale *et al.*, 2018 Evolution of Endothelin
711   signaling and diversification of adult pigment pattern in Danio fishes. PLoS Genet 14:
712   e1007538.
713   Ullate-Agote, A., and A. C. Tzika, 2021 Characterization of the Leucistic Texas Rat Snake
714   Pantherophis obsoletus. Frontiers in Ecology and Evolution 9:.
715   Untergasser, A., I. Cutcutache, T. Koressaar, J. Ye, B. C. Faircloth *et al.*, 2012 Primer3--new
716   capabilities and interfaces. Nucleic Acids Res 40: e115.
717   Watanabe, M., and S. Kondo, 2012 Changing clothes easily: connexin41.8 regulates skin
718   pattern variation. Pigment Cell Melanoma Res 25: 326–330.
719   Westphal, M. F., and T. J. Morgan, 2010 Quantitative genetics of pigmentation development in 2
720   populations of the common garter snake, Thamnophis sirtalis. J Hered 101: 573–580.
721   Wolf, M., and Y. Werner, 1994 The Striped Color Pattern and Striped Non-Striped
722   Polymorphism in Snakes (reptilia, Ophidia). Biol. Rev. Cambridge Philosophic. Soc. 69:
723   599–610.
724   Woodcock, M. R., J. Vaughn-Wolfe, A. Elias, D. K. Kump, K. D. Kendall *et al.*, 2017
725   Identification of Mutant Genes and Introgressed Tiger Salamander DNA in the
726   Laboratory Axolotl, Ambystoma mexicanum. Sci Rep 7: 6.
727   Yang, G. C., D. Croaker, A. L. Zhang, P. Manglick, T. Cartmill *et al.*, 1998 A dinucleotide
728   mutation in the endothelin-B receptor gene is associated with lethal white foal syndrome
729   (LWFS); a horse variant of Hirschsprung disease. Hum Mol Genet 7: 1047–1052.
730   Yoshihara, M., T. Sato, D. Saito, O. Ohara, T. Kuramoto *et al.*, 2017 A deletion in the intergenic
731   region upstream of Ednrb causes head spot in the rat strain KFRS4/Kyo. BMC Genetics
732   18: 29.
733   Zweifel, R. G., 1981 Genetics of color pattern polymorphism in the California kingsnake. Journal
734   of Heredity 72: 238–244.
735

Fig 1. Phenotypes of color morphs in the Yellowbelly series. (A) Schematic of homozygote
phenotypes in the Yellowbelly series. (B) Ball pythons described as homozygous for each allele in the
Yellowbelly series. (C) Wildtype ball python. (D) Ball pythons described as compound heterozygous for
alleles in the Yellowbelly series. (E) Underbelly of a wildtype ball python. (F) Ball python described as
heterozygous for the *yellowbelly* allele. The underbelly is subtly lighter and differently patterned compared
to wildtype. (B-F) Images are representative examples. Variation exists within each genotype. Photos,
Chiron Graves, Gavin Costello, Ball Python Shed, Ball In Hand Pythons, Barbara Penyak, Freedom
Breeder, Patrick Faulkner, Off The Wall Balls, Colorado Cold Bloods, Laura Carter, Brent McKelvey,
Travis Wyman.

Fig 2. Phenotypes in the Yellowbelly series are hypothesized to arise from loss of chromatophores. (A) Simplified model of chromatophore migration and stripe formation during the larval-to-adult transition in zebrafish (Singh and Nuesslein-Volhard 2015; Patterson and Parichy 2019). Xanthophore precursors (yellow) persist in the epidermis from larvalhood. Melanophore and iridophore precursors (brown and blue) migrate from the neural crest to the epidermis along nerve tracks. The primary route for iridophores is through the horizontal midline. The adult pattern consists of dark stripes (melanophores and sparse iridophores) and light inter-stripes (xanthophores and dense iridophores). Stripes and inter-stripes form iteratively, beginning with the first inter-stripe at the horizontal midline. (B) Schematic of mutant zebrafish with a primary loss of iridophores and a secondary loss of melanophores (Parichy *et al.* 2000; Lopes *et al.* 2008; Patterson and Parichy 2013; Frohnhöfer *et al.* 2013; Mo *et al.* 2017; Spiewak *et al.* 2018). In such mutants, iridophores fail to populate the skin and instead remain clustered near the horizontal midline. Melanophores require iridophores for survival and are reduced as a secondary consequence of iridophore loss. Melanophores persist only near the horizontal midline. (C) Hypothesized model for chromatophore migration in ball python. Chromatophores are presumed to migrate in a dorsal-to-ventral path, as they do in other higher vertebrates (Mort *et al.* 2015). (D-F) Phenotypes in the Yellowbelly series are hypothesized to be caused by reductions in chromatophores.

761 Hypothesized reductions range from mild (altered patterning) to moderate (dorsal striping) to severe
762 (leucism).

Breeder description	EDNRB1 genotype						Total	# Owners	# US states
	1-bp Δ / 1-bp Δ	1-bp Δ / 17-bp Δ	1-bp Δ / +	17-bp Δ / +	+/+				
yellowbelly / yellowbelly	61	9					70	35	21
+/+					48		48	33	16
yellowbelly / +			92	6	32*		130	68	27

Fig 3. The *yellowbelly* allele is associated with frameshift variants in *EDNRB1*. (A) Ball python described as homozygous for the *yellowbelly* allele. Photo, Barbara Penyak. (B) Schematic of molecular *EDNRB1* alleles associated with the *yellowbelly* allele. One allele contains a 1-bp deletion. The other allele contains a 17-bp deletion. (C) Table showing association between the *yellowbelly* allele and the 1-bp and 17-bp deletions. This table includes the original *yellowbelly* homozygote in which the 1-bp deletion was identified. Numbers of owners and US states indicate the total numbers of owners and US states from which the animals were collected. *, presumed mis-labeling by owner, including one animal heterozygous for *L152F* (see Discussion). (D) Schematic of wildtype EDNRB1 protein, created using Protter (Omasits *et al.* 2014). (E-F) Schematics of truncated EDNRB1 proteins resulting from the 1-bp and 17-bp deletions. Schematics are intended to show premature terminations and do not imply that the truncated proteins fold properly and are inserted into the cell membrane.

Fig 4. The *spark* and *specter* alleles are associated with missense variants in *EDNRB1*. (A-B) Ball
pythons described as homozygous for the *spark* allele or the *specter* allele. Photos, Freedom Breeder,
Patrick Faulkner. (C) Schematic of molecular *EDNRB1* alleles associated with the *spark* and *specter*
alleles. The *spark* allele contains missense variant *L152F*. The *specter* allele contains missense variant
R315P. (D) Table showing associations between the *spark* and *specter* alleles and the missense variants.
This table includes the original *spark/yellowbelly* compound heterozygote and the original *specter*
homozygote in which the missense variants were identified. Numbers of owners and US states indicate
the total numbers of owners and US states from which the animals were collected. *, presumed mis-
labeling by owner (see Discussion). (E-F) Schematics of EDNRB1 proteins containing missense variants,
created using Protter (Omasits *et al.* 2014). (G) Alignment of EDNRB1 and EDNRA protein sequences
surrounding missense variants. EDNRA is another endothelin receptor and is included to demonstrate
conservation outside EDNRB1. Full EDNRA sequences were not available for ball python or painted
turtle.

D — *EDNRB1* genotype

Breeder description	$G\text{-}to\text{-}A^{SD1}$ / $G\text{-}to\text{-}A^{SD1}$	$G\text{-}to\text{-}A^{SD1}$ / 1-bp Δ	$G\text{-}to\text{-}A^{SD6}$ / $G\text{-}to\text{-}A^{SD6}$	$G\text{-}to\text{-}A^{SD6}$ / 1-bp Δ	$G\text{-}to\text{-}A^{SD6}$ / 17-bp Δ	+ / +	Total	# Owners	# US states
gravel / gravel	7						7	4	4
gravel / yellowbelly		22					22	14	11
asphalt / asphalt			4				4	4	4
asphalt / yellowbelly		2*		18	3		23	13	10
+ / +						48	48	33	16

Fig 5. The *gravel* and *asphalt* alleles are associated with nucleotide substitutions in *EDNRB1* splice donors. (A-B) Ball pythons described as homozygous for the *gravel* allele or the *asphalt* allele. Photos, Off The Wall Balls, Ball Python Shed. (C) Schematic of molecular *EDNRB1* alleles associated with the *gravel* and *asphalt* alleles. The *gravel* and *asphalt* alleles contain G-to-A nucleotide substitution in the -1 positions of the first and sixth splice donors, respectively (SD1 and SD6). (D) Table showing associations between the *gravel* and *asphalt* alleles and the nucleotide substitutions in splice donors. This table includes the original *gravel* and *asphalt* homozygotes in which the nucleotide substitutions were identified. Numbers of owners and US states indicate the total numbers of owners and US states from which the animals were collected. *, presumed mis-labeling by owner (see Discussion). (E) Alignment of nucleotide sequences for the first and sixth splice donors of *EDNRB1*.

Fig 6. Summary of genetic variants in *EDNRB1* associated with phenotypes in the Yellowbelly series. The most severe phenotype in the Yellowbelly series was leucism, which was associated with frameshift variants (1-bp and 17-bp deletions). The next most severe phenotype was dorsal striping, which was associated with missense variants *L152F* and *R315P*. The mildest phenotype was altered patterning, which was associated with G-to-A nucleotide substitutions in the -1 positions of splice donors.

802 # Supporting Tables

803 **S1 Table. Primers used to amplify and sequence coding regions of *EDNRB1* and *EDN3*.**

804 **S2 Table. Genotypes of all animals.**

Breeder description	Genotype	Band sizes
+ / +	+ / +	6 bp, 103 bp, 111 bp, 173 bp
yellowbelly / +	+ / *1-bp Δ*	6 bp, 103 bp, 111 bp, 173 bp, 213 bp, 213 / 214 bp heteroduplex
yellowbelly / +	+ / *17-bp Δ*	6 bp, 103 bp, 111 bp, 173 bp, 259 bp, 259 / 276 bp heteroduplex
yellowbelly / *yellowbelly*	*1-bp Δ* / *1-bp Δ*	6 bp, 173 bp, 213 bp
yellowbelly / *yellowbelly*	*1-bp Δ* / *17-bp Δ*	6 bp, 111 bp, 173 bp, 213 bp, 213 / 214 bp heteroduplex, 259 bp, 370 / 386 bp heteroduplex

S1 Fig. Assay for genotyping the 1-bp and 17-bp deletions in *EDNRB1*. (A) Schematic of the wildtype
allele, the 1-bp deletion allele, and the 17-bp deletion allele of *EDNRB1*. The wildtype allele contains
three cut sites for *Bsr*I. The deletions each eliminate one of these sites. (B) Band sizes observed for each
genotype, demonstrating that the 1-bp and 17-dp deletions were carried on opposite chromosomes.
Heteroduplexes were formed because digests were performed at 65°C. (C) Agarose gel electrophoresis
of *Bsr*I digests.

A EDNRB1

PCR primers:

wildtype allele — 376 bp

specter allele — 96 bp / *Sma*I / R315P / 280 bp

B

Breeder description	Genotype	Band sizes
+/+	+/+	376 bp
specter/+	+/R315P	96 bp, 280 bp, 376 bp
specter/specter	R315P/R315P	96 bp, 280 bp

C

811 **S2 Fig. Assay for genotyping the *R315P* missense variant in *EDNRB1*.** (A) Schematic of the wildtype
812 allele and the *R315P* allele of *EDNRB1*. The *R315P* variant creates a cut site for *Sma*I. (B) Band sizes
813 observed for each genotype. (C) Agarose gel electrophoresis of *Sma*I digests.

Variant	Breeder description	Genotype	Chromatogram
L152F	spark / yellowbelly	L152F / 1-bp Δ	TCATTGCTTTNCCCATTAATG
	Non-Yellowbelly	+ / +	TCATTGCTTTGCCCATTAATG
G-to-A^SD1	gravel / gravel	G-to-A^SD1 / G-to-A^SD1	ATGTGTATAAAGTATTTATTT
	Non-Yellowbelly	+ / +	ATGTGTATAAGGTATTTATTT
G-to-A^SD6	asphalt / asphalt	G-to-A^SD6 / G-to-A^SD6	ACTGCTTTCAAGTAAGAGAGT
	Non-Yellowbelly	+ / +	ACTGCTTTCAGGTAAGAGAGT

814 **S3 Fig. Chromatogram examples.** Examples of chromatograms from Sanger sequencing reads used to
815 genotype the *L152F* missense variant and the G-to-A substitutions in splice donors. SD1, splice donor for
816 the first intron. SD6, splice donor for the sixth intron.

A Phenotype of homozygote

Most severe — Least severe — Wildtype

Allele(s): *yellowbelly* — *spark, specter* — *gravel, asphalt*

B
yellowbelly / yellowbelly — *spark / spark* — *specter / specter* — *gravel / gravel* — *asphalt / asphalt*

C
+ / +

D
spark / yellowbelly — *specter / yellowbelly* — *gravel / yellowbelly* — *asphalt / yellowbelly* — *spark / gravel* — *specter / asphalt*

E
+ / +

F
yellowbelly / + — *yellowbelly / +*

A Zebrafish

Wildtype

B Mutant with fewer iridophores and melanophores

■ Melanophore
■ Xanthophore
■ Iridophore
* Neural crest
▷ Horizontal midline

Ventral

C Ball python

Wildtype

D Altered patterning

E Striped

F Leucistic

Ventral

A *yellowbelly / yellowbelly*

B *yellowbelly* alleles

EDNRB1 1-bp Δ

EDNRB1 17-bp Δ

1 kb

C

	EDNRB1 genotype						
Breeder description	*1-bp Δ /* *1-bp Δ*	*1-bp Δ /* *17-bp Δ*	*1-bp Δ /* +	*17-bp Δ /* +	+ / +	Total	# Owners	# US states
yellowbelly / yellowbelly	61	9				70	35	21
+ / +					48	48	33	16
yellowbelly / +			92	6	32*	130	68	27

D EDNRB1^Wildtype

E EDNRB1^1-bp Δ

F EDNRB1^17-bp Δ

Extracellular

Cytoplasmic

○ Amino acid upstream of frameshift
● Amino acid altered by frameshift

A *spark / spark*

B *specter / specter*

C

spark allele

EDNRB1 — L152F

specter allele

EDNRB1 — R315P

1 kb

D

Breeder description	*EDNRB1* genotype					Total	# Owners	# US states
	L152F / 1-bp Δ	*L152F / 17-bp Δ*	*R315P / R315P*	*R315P / 1-bp Δ*	*+ / +*			
spark / yellowbelly	19	2		1*		22	14	11
specter / specter			3			3	3	2
specter / yellowbelly	1*			26		27	17	11
+ / +					48	33	33	16

E EDNRB1^L152F

N — L152F — C

F EDNRB1^R315P

N — R315P — C

G

Protein sequence

Position: 152   315

EDNRA

Elephant shark	...YIAIDIPIHVF...EHLKQRREVAK...
Zebrafish	...YITIDIPINVY...EHLKQRREVAK...
Human	...YVVIDIPINVF...EHLKQRREVAK...
Chicken	...YIVIDIPIIVY...EHLKQRREVAK...
Tropical clawed frog	...YIVIDIPIIVF...EHLKQRREVAK...
Anole lizard	...YIIIDIPINVY...EHLKQRREVAK...
Burmese python	...YIIIDIPINVY...KHRKQRREVAR...

EDNRB1

Elephant shark	...HIAIDIPINVY...DHLKQRREVAK...
Zebrafish	...HIVIGLPVNVY...KHRKQRREVAR...
Human	...HIVIDIPINVY...DHLKQRREVAK...
Chicken	...YILIALPINVY...DHMKRRREVAK...
Tropical clawed frog	...HIIIDIPITTY...DTLRQRREIAK...
Painted turtle	...YILIALPINVY...DHMKRRREVAK...
Anole lizard	...YIIIGLPTNVY...DHMKRRREVAK...
Burmese python	...YILIALPINVY...DHMKRRREVAK...
Ball python	...YILIALPINVY...DHMKRRREVAK...

Ball python *L152F*	...YILIAFPINVY...DHMKRRREVAK...
Ball python *R315P*	...YILIALPINVY...DHMKRPREVAK...

A *gravel / gravel*   **B** *asphalt / asphalt*   **C**

gravel allele

G-to-A^{SD1}

EDNRB1

asphalt allele

G-to-A^{SD6}

EDNRB1

1 kb

D

EDNRB1 genotype

Breeder description	*G-to-A^{SD1} / G-to-A^{SD1}*	*G-to-A^{SD1} / 1-bp Δ*	*G-to-A^{SD6} / G-to-A^{SD6}*	*G-to-A^{SD6} / 1-bp Δ*	*G-to-A^{SD6} / 17-bp Δ*	+ / +	Total	# Owners	# US states
gravel / gravel	7						7	4	4
gravel / yellowbelly		22					22	14	11
asphalt / asphalt			4				4	4	4
asphalt / yellowbelly		2*		18	3		23	13	10
+ / +						48	48	33	16

E

EDNRB1

	Exon 1	Intron 1		Exon 6	Intron 6
Elephant shark	...A **G**	G T...		...A **G**	G T...
Zebrafish	...A **G**	G T...		...G **G**	G T...
Human	...A **G**	G T...		...A **G**	G T...
Chicken	...A **G**	G T...		...A **G**	G T...
Tropical clawed frog	...A **G**	G T...		...A **G**	G T...
Painted turtle	...A **G**	G T...		...A **G**	G T...
Anole lizard	...A **G**	G T...		...A **G**	G T...
Ball python	...A **G**	G T...		...A **G**	G T...
Ball python *G-to-A^{SD1}*	...A **A**	G T...		...A **G**	G T...
Ball python *G-to-A^{SD6}*	...A **G**	G T...		...A **A**	G T...
Position:	-2 -1	+1 +2		-2 -1	+1 +2