Title: Coupled functional physiological phenotyping and simulation model to estimate dynamic water use efficiency and infer transpiration sensitivity traits

Author: Ting Sun[1,2#], Rui Cheng[3#], Yudong Sun[3], Rujia Jiang[1], Zhuoyi Wang[1], Pingping Fang[1,2], Xinyang Wu[1,2], Kang Ning[1,2], Pei Xu[1,2*]

Authors' affiliation: [1] College of Life Sciences, China Jiliang University, Hangzhou 310018, P.R. China; [2] Key Laboratory of Specialty Agri-product Quality and Hazard Controlling Technology of Zhejiang Province, Hangzhou 310018, P.R. China; [3]Vegetable Research Centre, Huaiyin Institute of Agricultural Sciences of Xuhuai Region in Jiangsu, Huai'an 223001, P.R. China

[#] Equal contribution.

***Correspondence:** Pei Xu, Email: peixu@cjlu.edu.cn

Abstract

As agricultural drought becomes more frequent worldwide, it is essential to improve crop productivity whilst reducing the water consumption to achieve a sustainable production. Plant transpiration rate and water use efficiency (WUE) collectively determine the yield performance, yet it is challenging to balance the two in breeding programs due to still insufficient mechanistic understanding of the traits. Here we demonstrate the feasibility and effectiveness of calculating dynamic and momentary WUE by coupling WUE model and the state-of-the-art functional physiological phenotyping (FPP). We also present the method of quantifying genotype-specific traits reflecting sensitivity of transpiration to radiation (S_{Tr-Rad}) and vapor pressure deficit (S_{Tr-VPD}), under evolving developmental stage and water availability. Using these methods, we revealed the genotypic difference of S_{Tr-Rad} and S_{Tr-VPD} in three watermelon accessions, the dramatic change in each of them across the drought treatment phases, and the quantitative impacts of them on dynamic WUE patterns. Based on our results and computational simulations, a general principle for transpiration ideotype design is proposed, which highlights the benefits of lowering S_{Tr-VPD} to increase WUE and increasing S_{Tr-Rad} to offset the decline of Tr. FPP-enabled phenomic selection will help screen for elite crops lines with desired transpiration sensitivities.

1. Introduction

Global farming land area subject to drought is increasing due to the enhanced evapotranspiration associated with increased temperature, net radiation and decreased relative humidity (IPCC, 2021). Improving crop productivity, particularly under deficit water condition, is essential to safeguard food security under the context of global climate change. Stomata is the gateway of both water efflux (transpiration) and CO_2 intake (assimilation). The plant transpiration rate (Tr) and the ratio of CO_2 assimilation to Tr, which is known as water use efficiency (WUE), collectively determine biomass production and yield (Steduto *et al.*, 2012; Vadez *et al.*, 2014). Large efforts are being devoted to developing drought-resistant crops by adopting a strategy of restricting stomatal closure and thereby water loss; however, a tradeoff between increased WUE and decreased total transpiration as a consequence of stomatal restraint has long impeded breeding for agronomically useful varieties (Dalal *et al.*, 2019). Previous studies have stressed that high-WUE crops usually had a smaller leaf area index (LAI), shorter development stage and thus lower Tr and carbon assimilation (Blum, 2005; Blum, 2009).

Traditionally, WUE is considered to be a conservative genotype-dependent trait (Steduto *et al.*, 2012) and is measured by carbon-isotope discrimination (Moghaddam *et al.*, 2013) or canopy gas exchange measurement systems (Song *et al.*, 2016). It is now increasingly recognized that WUE is highly variable in response to environment change, developmental stages, and abiotic stress (Nelson *et al.*, 2018; Peddinti *et al.*, 2019; Yang *et al.*, 2021). Monitoring dynamic WUE is therefore becoming routine in physiological studies and crop cultivation practices, even though it is presently very challenging due to technical constraints in the continuous nondestructive measurement of carbon assimilation and transpiration processes. WUE can change with the pattern of Tr within a day even if the daily cumulative Tr remains unchanged (Taylor *et al.*, 1983; Vadez *et al.*, 2014; Ghanem *et al.*, 2020). Sinclair and Vadez (2012) decoded daily WUE as a function of vapor pressure deficit (VPD) weighted by diurnal Tr, implying the feasibility of selecting high-WUE varieties by harnessing

genotypic variations in the dynamic Tr profile. It has been confirmed that restricted

maximum Tr at noon under high VPD could increase the WUE and yield of maize

(Jafarikouhini *et al.*, 2022), sorghum (Sinclair *et al.*, 2005) and soybean (Sinclair *et

al.*, 2010).

Mechanistically, Tr is not only governed by internal factors such as LAI and

structure (e.g. cuticle thickness, stomata number), but external factors including

temperature, relative humidity, light intensity, wind speed and soil water availability

(Kubota, 2020). The interaction between Tr and the environment confounds

phenotypic selection for Tr. According to Pieruschka *et al.* (2010), transpiration is

mainly controlled by the diffusion gradient of water vapor inside and outside the

stoma driven by VPD, and internal evaporation driven by latent heat from Rad. Since

the diurnal patterns of Rad and VPD are inconsistent (Gosa *et al.*, 2019), the

sensitivities of Tr to VPD and Rad would have a strong impact on the dynamic Tr

profile. Simulation models, which combine many biological functional hypotheses in

mathematical frameworks, describe the development of plant traits as a consequence

of environmental and genetic interaction (Génard *et al.*, 2016). Several simulation

models for Tr have been established (Katsoulas & Stanghellini, 2019). Recently,

simulation models have found their new use in inferring parameters representing

genotype-specific traits or "hidden traits" that are not intuitively extractable from the

measured phenotypes. Such an approach is referred to as "inverse modeling" (Chenu

et al., 2017).

Compared to morphological traits, plant physiological traits are notoriously

difficult to phenotype. The emerging functional physiological phenotyping (FPP) has

offered revolutionary paths for phenotyping plant water relations (Xu *et al.*, 2015;

Halperin *et al.*, 2017; Chenu *et al.*, 2018; Li *et al.*, 2020). One of such platforms

named "Plantarray" allows for high-resolution (every 3 minutes), non-destructive,

simultaneous and continuous recording of meteorological variables and physiological

traits such as diurnal Tr and daily biomass gain based on the lysimeter principle. The

acquired massive dynamic data from the soil-plant-atmosphere continuum provides

unprecedent opportunities for deeply understand the mechanisms of plant water

relations. Here, we report on the estimation of dynamic WUE, a traditionally

difficult-to-measure trait, by coupling existing WUE models with FPP, as well as the

quantification of the genotype-specific dynamic traits of "Tr sensitivity to Rad/VPD"

($S_{Tr\text{-}Rad}$ and $S_{Tr\text{-}VPD)}$ by coupling FPP and the modified Penman–Monteith model

(Medrano *et al.*, 2005; Jo & Shin, 2021). This study demonstrates the great promise of

leveraging advanced phenotyping techniques and existing modeling methods to tackle

the long-standing complex trait challenge. The implications for transpiration ideotype

design are discussed.

2. Materials and methods

2.1. Plant materials and experimental design

Three genotypes of watermelon, i.e., a wild watermelon accession "PI-296341-FR"

(*Citrullus lanatus* var. citroides), a cultivated watermelon variety "Jincheng No. 5"

(*Citrullus lanatus* var. lanatus, hereafter "JC5") popular in Northwest China, and a

drought-tolerant inbred line of *Citrullus lanatus* var. lanatus "HA" were used in this

study. The experiments were carried out in the spring and summer seasons of 2021 in

a semi-controlled greenhouse (L×W×H: 10m×5m×4m) in Huai'an (119°01′E,

33°35′N), China. One seedling at the three-leaf stage was transplanted per pot

(16cm×13cm×18cm, 1.5 L) with Profile Porous Ceramic substrate (diameter: 0.2mm,

pH: 5.5±1, porosity: 74, and CEC: 33.6mEq/100g).

Four replicates for each genotype were set in a completely randomized design on

"Plantarray" (Plant-Ditech, Israel, Fig.1a). This system provides a simultaneous

measurement and detailed physiological response profile for plants in each pot (Dalal

et al., 2020). Water and nutrient supplements (Yamazaki nutrient solution) were

controlled by the automatic irrigation system of Plantarray. Experiment in each

growth season included three treatment periods, i.e., well-irrigation (WI), progressive

water deficit (WD) and water recovery (WR). Under the WI and WR phase, nutrient

solution was provided by irrigation for 240 seconds (oversaturated) at 23:00, 1:00,

2:00, and 3:00 of a day, respectively, and no nutrient solution was supplied during the

WD stage.

2.2.Functional physiological phenotyping (FPP)

The Plantarray system is a weighing type lysimeter that consists of highly sensitive,

temperature compensated load cells (Halperin *et al.*, 2017). A series of functional

physiological traits relative to the water loss by plant transpiration and plant growth

could be obtained by the weight change throughout the day using the SPAC software

implemented in the Plantarray system (Fig. 1a & b). Briefly, the pre-dawn system

weight (averaged over 4:00-4:30 a.m.) was recoded as W_m at the end of the free

drainage after saturated irrigation, and system weight in the evening (averaged over

9:00-9:30 p.m.) was recoded as W_e before irrigation, both of which were stable due to

no water loss by plant transpiration and water input from irrigation during this period.

As each pot on the system was covered with plastic film, soil evapotranspiration was

prevented. The whole-plant daily transpiration (E, g d^{-1}) was calculated as the

difference between the W_m and W_e for each day. The daily plant growth (PG, g FW

d^{-1}) was determined as the difference of W_m between consecutive days, as the soil

weight at field capacity in different days remained constant. The whole-plant WUE

during the well irrigation period was thus determined by the ratio between the

accumulated PG and E. The momentary whole-plant transpiration (Tr, g min^{-1}) at

3-min step was calculated by multiplying the first derivative of the measured load-cell

time series by -1, assuming that the plant's weight gain during the short time interval

used to calculate the transpiration rate was negligible (Halperin *et al.*, 2017).

The air temperature (T_{air}, $\square$) and relative humidity (RH, %), photosynthetically

active radiation (PAR, μmol m^{-2} s^{-1}) above the canopy, and soil volumetric water

content (VWC, m^3 m^{-3}) in each pot were measured by the VP-4, 5TM, and PYR solar

radiation sensor (Decagon Devices, Pullman, Wash, USA), respectively (Fig. 1c-e,

Fig. S1). VPD (kPa) was calculated according to Halperin *et al.* (2017).

2.3.Simulation of WUE and biomass gain

148 The initial definition of WUE considers the resistances to diffusion resulting from

149 stomata, leaf aerodynamic boundary layer, and leaf mesophyll resistance, and can be
150 ultimately simplified to:

$$WUE = \frac{K_d}{VPD_d} \quad (1)$$

where VPD_d is the mean VPD over a day. K_d is a coefficient that reflects the CO_2
concentration in the stomatal chamber, which has been validated among different
crops (Tanner & Sinclair, 1983; Sinclair *et al.*, 1984). K_d was estimated as 6.5 kPa in
this study for watermelon plant with the data of FPP and WUE obtained from
destructive sampling during a long growth period.

To estimate WUE over a long period, VPD needs to be integrated over a growth
season and weighted by the transpiration over the course (Taylor *et al.*, 1983; Vadez *et*
al., 2014; Ghanem *et al.*, 2020). Applying the same logic to a single day, Sinclair and
Vadez (2012) proposed an assumption that the weighting of this component based on
the transpiration rate throughout the day leads to the possibility of obtaining altered
values if there are genotypic variations in the transpiration profile throughout the day.
Therefore, during a computation interval (i), the mean VPD (VPD_i) is weighted for
mean transpiration rate (Tr_i) to calculate the estimated WUE (WUE_e) as in the
following integrating equation:

$$WUE_e = \frac{\int K_d Tr_i /VPD_i}{\int Tr_i} \quad (2)$$

165   And biomass gain during this period could be estimated as:

$$Biomass_e = \frac{\int K_d Tr_i}{VPD_i} \quad (3)$$

2.4.Canopy gas exchange parameters for validation
Gas exchange parameters were obtained by the classical but low-throughput canopy
photosynthesis and transpiration system (CAPTS 100, China) from wheat plants
which were cultivated in the field in 2021. The gas exchange parameters of the 1 m ×
1 m canopies were measured automatically every 30 min using CAPTS 100
consisting of a cubic transparent chamber, which can be open and closed
automatically with programming and equipped with CO_2, humidity, air pressure, and

temperature sensors in it (Song *et al.*, 2016). Canopy gas exchange rate was measured

according to the change rate of CO_2 and H_2O concentrations in the chamber during

the closure of the chamber (Fig. S2a). The parameter K_d for wheat plants in this study

was estimated using the least square method by fitting the estimated and measured

daily WUE obtained during the growth period of March; then estimated and measured

WUE at half-hour intervals and daily intervals during the days between April 5 and

April 10 of 2021 were shown in Fig. S2b.

2.5. Transpiration model

We choose the simulation models which account for the response of Tr to the major

meteorological variables and plant growth, following the principle that model obtains

simple structure using as few parameters as possible. A modified Penman–Monteith

model which is appropriate for greenhouse environment, is employed to simulate the

Tr of watermelons in this study, which has been test for cucumber and tomato plants

(Medrano *et al.*, 2005; Jo & Shin, 2021). The diurnal Tr could be described as a

function of radiation, VPD and leaf area index (LAI):

$$\lambda \mathrm{Tr}_i = A(1 - e^{-K\mathrm{LAI}_d})\mathrm{Rad}_i + B\mathrm{LAI}_d * \mathrm{VPD}_i \qquad (4)$$

where λ is the vaporization heat of water (J kg^{-1}), Tr_i is the instantaneous transpiration

rate, K is the light extinction coefficient (0.86), Rad_i is the instantaneous solar

radiation (W m^{-2}, converted from PAR), LAI_d is the daily leaf area index (m^{-2} m^{-2}).

VPD_i is the instantaneous vapor pressure deficit. The LAI was estimated based on

daily transpiration in each experiment unit by considering the individual difference

(detailed as follows).

2.6. LAI estimation and validation

Step 1, dynamic Tr recorded in a frequency of every 3 minutes across a day was fitted

to the observed radiation and VPD, thus optimum *A*, *B*, and LAI in the modified

Penman–Monteith equation (Eq. 4) were estimated according to nonlinear least square

(NLS) method and trust-region algorithm via the trial version of MATLAB 2018b

(MathWorks, Inc., Cambridge, MA, USA). Only daily LAI with robust goodness of fit

(adjusted $R^2 > 0.85$) were picked for subsequent calculations. Step 2, we checked the

accumulated daily transpiration (E) measured by Plantarray and the estimated LAI

from daily fitting of step 1, where good linear relationship was found between them.

Therefore, a constant LAI growth rate per E was estimated for individual experiment

units, and then daily LAIs of individual experiment units were re-calculated according

to daily measured E and estimated LAI growth rate per E. Step 3, daily parameter A

and B were determined with observed Tr, RAD, VPD and re-calculated LAI during

Step 2 for WI, WS, and WR stage, respectively. Parameter fitting was performed also

with MATLAB 2018b.

Independent experiment was conducted to verify the aforementioned method,

where estimated LAI was compared to the LAI measured by an LA-S leaf area

analyzer system (Wanshen Technology Co., Hangzhou, China, Fig. 2).

3. Results and discussion

3.1. Coupled FPP and simulation models to estimate dynamic and momentary

WUE

Our idea was to extend Sinclair and colleagues' model (Eq. 1), a theoretical

expression of WUE (Sinclair *et al.*, 1984; Sinclair & Ghanem, 2020), to FPP data for

dynamic WUE estimation (Eq. 2). Prior to doing so, we tested the reliability of this

approach by using a dataset obtained from wheat via the traditional low-throughput

canopy gas exchange method (Song *et al.*, 2016). During a 5-day period, canopy

photosynthesis rate was found to peak much earlier than canopy transpiration rate

before noon (Fig. S2a), which led to a higher WUE during the whole forenoon (square

in Fig. S2b). The WUE_e estimated from the canopy transpiration rate and VPD at

half-hourly intervals could well reproduce the measured WUE during the daytime

(line in Fig. S2). At the daily step, day-to-day variations of both measured and

estimated WUE were observed (triangles in Fig. S2), which were highly consistent as

indicated by the tiny root mean square error (RMSE) of 0.00137 g g^{-1}. These results

proved that the chosen WUE model was suitable for dynamic WUE estimation.

In our high-throughput FPP data, the daily estimated WUE (WUE_e, g DW/g

H$_2$O) calculated by Eq.2 based on the Tr and VPD continuously acquired by the

Plantarray system, was found to fluctuate across the course of the experiment and

varied significantly by genotype (Fig. 3a, similar result during spring growth season

was shown in Fig. S3). During the WI phase, WUE (g FW/g H$_2$O) within this period

could be measured based on traditional method, i.e., accumulated daily transpiration

dividing by accumulated biomass gain. The measured WUE based on fresh weight

returned similar result of genetic difference to estimated WUE$_e$ based on dry weight

(Fig. 3b). Among the three genotypes, "HA" possessed the highest WUE across all

phases, in particular late WD stage, followed by "JC5" and "PI296341-FR" (Fig. 3a,

Fig. S3), in accord with the empirical knowledge that wild varieties tend to possess a

higher level of drought resistance by sacrificing more biomass production (Kawasaki

et al., 2000).

When shortening the interval i in Eq. 2, the higher-resolution WUE (in the

extreme case, the instantaneous or momentary WUE) and its diurnal variation could

be estimated. In Fig. 3c (red solid line), we show that the hourly WUE$_e$ calculated

from measured Tr and VPD peaked in the early morning (6:00 am) and decreased

until reaching a stable level at noon, and then rebounded slightly in the afternoon,

which was consistent with the gas-exchange measurements elsewhere at the single

leaf scale (Peddinti *et al.*, 2019; Yang *et al.*, 2019) or at the canopy scale.

3.2. Inferred sensitivity of Tr to Rad/VPD through inverse modeling

With the diurnal data of Tr and VPD, we were also able to conduct a simulation

analysis by artificially advancing the dynamic Tr for 1 hour (black solid line shift to

black dotted line) across a whole day (Fig. 3c), which revealed an increase of daily

WUE$_e$ and Biomass$_e$ by 29.9 % and 20.0 %, respectively. The hourly WUE$_e$ (red line)

and Biomass$_e$ (blue shadows) indicated that enhancing WUE during the early morning

contributed greatly to the biomass gain (Fig. 3c). According to Eq. 1 and Eq. 2, the

interpretation of this result is straightforward: without sacrificing daily accumulative

Tr, more Tr in the low-VPD morning time decreases the integral value of daily VPD

weighted by Tr, thus contributing to the increase of whole-day WUE and biomass gain.

Such relatively small, daily improvement will accumulatively lead to an improved

seasonal yield. Therefore, the pattern of diurnal Tr, even under the same daily

cumulative transpiration conditions, is considered to reverse daily WUE.

To disentangle the intrinsic traits related to sensitivity of Tr to major

environmental factors that were hidden within the daily Tr profiles, the modified

Penman–Monteith model (Eq. 4) was employed to fit the dynamic Tr with the

measured Rad and VPD, as well as the estimated daily LAI. LAI is often estimated as

a function of thermal time, i.e. growing degree days (Medrano *et al.*, 2005; Jo & Shin,

2021); however, such approch fell short of the need in our study, as LAI of individual

plants differed remarkably which would influence the parameterization of Eq. 4. Here,

daily transpiration of individual plants monitored by Plantarray in a non-destructive

way was used to calibrate LAI in Eq. 4, a strategy that has been widely applied in

coupling remote sensing and crop modeling (Zhao et al., 2013). Meanwhile, we

measured the dynamic LAI of watermelon plants destructively during canopy

development. As shown in Fig. 2, the estimated and measured LAI at each sampling

time were similar (RMSE= 0.073), corroborating the robustness of our methods.

Following that, two genotype-specific parameters in this model, *A* and *B*, which were

essentially the S_{Tr-Rad} and S_{Tr-VPD}, were estimated from curve fitting of transpiration

rate with the input of measured Rad, VPD and estimated LAI, respectively. The fitted

transpiration dynamics well reproduced the measured diurnal transpirations, as

evidenced by the high correlation (R^2=0.94) between them and the low total RMSE of

only 0.05 g/min, taking the commercial cultivar "JC5" for example (Fig. 4a). S_{Tr-Rad}

and S_{Tr-VPD} were thus inferred via inverse modeling in a daily step as shown in Fig 4b

& c (similar result during spring growth season was shown in Fig. S4). We conducted

two-way ANOVA for the daily S_{Tr-Rad} and S_{Tr-VPD} within each phase of WI, WD and

WR, and the result indicated genotypic variation and dramatic change in each of them

across the treatment phases ($P < 0.01$, Fig. 4b, c, Fig. S4). In general, S_{Tr-Rad} decreased

as the water deficit progressed and rapidly recovered when water was resumed (Fig.

4b, Fig. S4a); S_{Tr-VPD}, however, decreased quickly on the initial days of water deficit,

and continued to decrease gradually on the following days until reaching a rather

stable level even when irrigation was resumed (Fig. 4c, Fig. S4b). Among the three

genotypes, "HA" exhibited high S_{Tr-Rad} only at the WD phase and remarkably low

S_{Tr-VPD} particularly at the WI stage in comparison to the remaining two genotypes (Fig.

4b&c, Fig. S4), indicating that plasticity of S_{Tr-Rad} and S_{Tr-VPD} in response to the

environment is important in shaping the genotypic-specific dynamic WUE patterns.

3.3. Transpiration ideotype design based on S_{Tr-Rad} and S_{Tr-VPD}

Based on the results above, we propose that an optimized combination of S_{Tr-Rad} and

S_{Tr-VPD} is important for balanced Tr, WUE, and biomass gain. Simulation models are

powerful tools to support breeding because of their capability to reveal the

interactions between each component within a system. Therefore, we conducted

simulations with the inputs of 5-day meteorological data from each of the WI and WD

phases (Fig. 5a, c) and the artificial combinations of S_{Tr-Rad} and S_{Tr-VPD} extremes (i.e.,

highest parameter A × lowest parameter B, lowest A × lowest B, highest A × highest B,

and lowest A × highest B in the model) adopted from the 3 watermelon genotypes

averaged over each 5-day course (Fig. 5b, d), where LAI was fixed to 3 to maintain a

fair canopy size. Since Rad peaked much earlier than VPD in the morning (Fig. 5a, c),

high A and low B resulted in high Tr during the low VPD intervals within a day and

consequently a low integrated value of daily VPD weighted by Tr (Eq. 1), leading to

high WUE (red line in Fig. 5b, d).

Further, the estimated biomass gains of different combinations were shown in Fig

5b, d. During the WI phase, highest A × highest B and lowest A × highest B both

resulted in much larger Tr, which produced approximately double the biomass gain

compared to the remaining combinations despite their lower WUE (Fig. 5b). While

during the WD phase, the highest A × highest B still obtained the highest biomass

gain benefitting from the highest Tr (blue line in Fig. 5d), but the highest A × lowest B

obtained a second highest biomass gain. Therefore, by considering both WUE and Tr,

we put forward the following principles for transpiration ideotype design: in the

well-irrigated agricultural areas, crop genotypes with high S_{Tr-Rad} and high S_{Tr-VPD} is

favorable by producing superior yield; in the arid agricultural areas where

water-saving is a necessity, genotypes with high S_{Tr-Rad} and low S_{Tr-VPD} is desired for

its marvelously reduced transpiration yet still considerable yield; in the intermittent

drought-prone areas where irrigation and water shortage alternate, genotypes showing

strong plasticity in S_{Tr-Rad} and S_{Tr-VPD} that are suited to the optimized balance of yield

and water consumption will be desired.

4. Conclusions

We demonstrate that daily to hourly or even instantaneous WUE can be estimated

by coupling existing WUE models with the dynamically recorded transpiration and

VPD from high-throughput FPP, which breaks the bottleneck of dynamic and

momentary WUE acquisition. We further demonstrate that daily WUE is affected by

the diurnal transpiration pattern of a plant, which is tightly linked to the

genotype-dependent sensitivity of Tr to Rad and VPD. A quantitative approach was

developed to infer these transpiration sensitivity parameters. We propose that it is

beneficial to increase WUE by lowering S_{Tr-VPD} while offsetting its negative impact on

Tr by increasing S_{Tr-Rad}. Although in previous studies high-WUE crops selected by

carbon isotope discrimination generally failed to match the demand of luxuriant

growth (Blum, 2005; Blum, 2009), the high-WUE trait is not necessarily linked to

growth limitation (Marguerit *et al.* (2014), which lays the foundation for synergistic

improvement of WUE and yield. FPP-enabled phenomic selection will help catch the

desired elite individuals from enormous germplasm lines, e.g. those with substantially

high S_{Tr-Rad} under both well-watered and water scarcity conditions.

Acknowledgments

340 We thank the supports of National Key Research & Development Program of China

341 (2021YFE19800), the Research & Development Program of Ningbo (2021Z006) and

342 the Key Scientific and Technological Grant of Zhejiang for Breeding New

343 Agricultural Varieties (2021C02066-5, 2021C02067-7).

Figure legends

Figure 1 Parameters of soil–plant–atmosphere continuum monitored by high-throughput phenotyping platform "Plantarray". a, overview of the automated physiological phenotyping array loaded with watermelon seedlings. b, system weights that were recorded every 3 minutes during the experimental course. Shadow areas show the observed variation. The experiment was divided into the well-irrigation (WI), progressive water deficit (WD), and water recovery (WR) phases. c-e, meteorological parameters, i.e., air temperature (T_{air}, □, c), relative humidity (RH, %, d), and photosynthetic active radiation (PAR, $\mu mol\ m^{-2}\ s^{-1}$, e) during the summer-season experiment in 2021.

Figure 2 Comparison of estimated and measured leaf area index (LAI). LAI was estimated by the methods described in the "Leaf area index (LAI) estimation" section of SI (shown as lines). At 13,17, 21, 25 and 28 days after transplanting, leaf tissues from 3 experiment units of Plantarray were destructively removed from watermelon plants, respectively. Leaf areas were measure by LA-S leaf area analyzer system (shown as circles). RMSE represents root mean square error.

Figure 3 Measured and estimated water use efficiency (WUE), and estimated biomass gain based on models. a, daily estimated WUE (WUEe, g DW g-1 H2O) using mechanism models based on continuously recorded transpiration rate and VPD by Plantarray during the summer-season experiment in 2021. Multiple comparison results among three genotypes were shown as mean values with lowercase letters to show statistically significant differences for the well-irrigation (WI), progressive water deficit (WD) and water recovery (WR) phases, respectively. b, measured WUE (g FW g-1 H2O) based on accumulated biomass and transpiration during the summer-season experiment in 2021. Accumulated biomass (g FW) and transpiration (g H2O) were calculated according to daily biomass gain and daily transpiration measured by "Plantarray" system during WI phase. Accumulated biomass of PI296341-FR, JC5 and HA were fitted to a linear function of accumulated transpiration for individual experiment units, and multiple comparison results among

375　three genotypes were shown. c, dynamic Tr (every 3 minutes), hourly WUE and

376　biomass gain estimated based on models, taking the data from an experiment unit in

377　June 19 for instance. A left-ward shift of dynamic Tr was applied to artificially

378　advance it for 1 h. Corresponding WUE and biomass gain were estimated and shown

379　as dotted lines.

Figure 4 Simulation of diurnal transpiration by using the modified Penman-Monteith model. a, observed and simulated diurnal variation of transpiration rate during summer-season experiment in 2021, taking an experiment unit of JC5 for instance. The observations and simulations were both 3-min steps. b-c, fitted daily parameter A or S_{Tr-Rad} (b) and parameter B or S_{Tr-VPD} (c) by modified Penman-Monteith model during summer-season experiment in 2021. Multiple comparison results among three genotypes (PI296341-FR, JC5, and HA) were shown as mean values with lowercase letters to show statistically significant differences for the well-irrigation (WI), progressive water deficit (WD) and water recovery (WR) phases, respectively.

Figure 5 Diurnal variation of Rad and VPD, and the responsive patterns of simulated Tr to different combinations of parameter A and B. a, c five-day Rad and VPD (normalized to 0-1 scale) recorded at 3-min intervals, which were selected from the well-irrigation (WI, 6.17-6.21, a) and progressive water deficit (WD, 6.25-6.29, c) phase, respectively. The LAI was set to a constant of 3 for all simulations. Mean values of parameters A and B for different genotypes (PI296341-FR, JC5, and HA) were calculated over each 5-day course, and the maxima and minima formed the four combinations as shown in the graph. Mean estimated WUE and biomass gain over 5-day course were denoted as $WUE_{e,5d}$ and $Biomass_{e,5d}$, respectively.

References

Blum A. 2005. Drought resistance, water-use efficiency, and yield potential—are they compatible, dissonant, or mutually exclusive? *Australian Journal of Agricultural Research* **56**: 1159-1168.

Blum A. 2009. Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. *Field Crops Research* **112**: 119-123.

Chenu K, Porter JR, Martre P, Basso B, Chapman SC, Ewert F, Bindi M, Asseng S. 2017. Contribution of crop models to adaptation in wheat. *Trends in Plant Science* **22**: 472-490.

Chenu K, Van Oosterom EJ, McLean G, Deifel KS, Fletcher A, Geetika G, Tirfessa A, Mace ES, Jordan DR, Sulman R, et al. 2018. Integrating modelling and phenotyping approaches to identify and screen complex traits: transpiration efficiency in cereals. *Journal of Experimental Botany* **69**: 3181-3194.

Dalal A, Bourstein R, Haish N, Shenhar I, Wallach R, Moshelion M. 2019. Dynamic Physiological Phenotyping of Drought-Stressed Pepper Plants Treated With "Productivity-Enhancing" and "Survivability-Enhancing" Biostimulants. *Frontiers in Plant Science* **10**: 905.

Dalal A, Shenhar I, Bourstein R, Mayo A, Grunwald Y, Averbuch N, Attia Z, Wallach R,
Moshelion M. 2020. A Telemetric, Gravimetric Platform for Real-Time Physiological
Phenotyping of Plant-Environment Interactions. *Journal of Visualized Experiments*.
Génard M, Memmah M-M, Quilot-Turion B, Vercambre G, Baldazzi V, Le Bot J, Bertin N,
Gautier H, Lescourret F, Pagès L 2016. Process-Based Simulation Models Are Essential
Tools for Virtual Profiling and Design of Ideotypes: Example of Fruit and Root. *Crop Systems*
Biology, 83-104.
Ghanem ME, Kehel Z, Marrou H, Sinclair TR. 2020. Seasonal and climatic variation of weighted
VPD for transpiration estimation. *European Journal of Agronomy* **113**: 125966.
Gosa SC, Lupo Y, Moshelion M. 2019. Quantitative and comparative analysis of whole-plant
performance for functional physiological traits phenotyping: New tools to support
pre-breeding and plant stress physiology studies. *Plant Science (Amsterdam, Netherlands)* **282**:
49-59.
Halperin O, Gebremedhin A, Wallach R, Moshelion M. 2017. High-throughput physiological
phenotyping and screening system for the characterization of plant-environment interactions.
The Plant Journal **89**: 839-850.
IPCC. 2021. Climate change 2021: the physical science basis. Contribution of working group I to the
sixth assessment report of the intergovernmental panel on climate change.
Jafarikouhini N, Sinclair TR, Resende MJCS. 2022. Limited-transpiration rate and plant
conductance in a diverse sweet corn population. *Crop Science* **62**: 374-381.
Jo WJ, Shin JH. 2021. Development of a transpiration model for precise tomato (Solanum
lycopersicum L.) irrigation control under various environmental conditions in greenhouse.
Plant Physiology and Biochemistry **162**: 388-394.
Katsoulas N, Stanghellini C. 2019. Modelling crop transpiration in greenhouses: different models for
different applications. *Agronomy* **9**: 392.
Kawasaki S, Miyake C, Kohchi T, Fujii S, Uchida M, Yokota A. 2000. Responses of Wild
Watermelon to Drought Stress: Accumulation of an ArgE Homologue and Citrulline in Leaves
during Water Deficits. *Plant and Cell Physiology* **41**: 864-873.
Kubota C 2020. Growth, development, transpiration, and translocation as affected by abiotic
environmental factors. In: Kozai T, Niu G, Takagaki M eds. *Plant Factory (Second Edition)*:
Academic Press, 207-220.
Li Y, Wu X, Xu W, Sun Y, Wang Y, Li G, Xu P. 2020. High-throughput physiology-based stress
response phenotyping: Advantages, applications and prospective in horticultural plants.
Horticultural Plant Journal **7**: 181-187.
Marguerit E, Bouffier L, Chancerel E, Costa P, Lagane F, Guehl JM, Plomion C, Brendel O. 2014.
The genetics of water-use efficiency and its relation to growth in maritime pine. *Journal of*
Experimental Botany **65**: 4757-4768.
Medrano E, Lorenzo P, Sánchez-Guerrero MC, Montero JI. 2005. Evaluation and modelling of
greenhouse cucumber-crop transpiration under high and low radiation conditions. *Scientia*
Horticulturae **105**: 163-175.
Moghaddam A, Raza A, Vollmann J, Ardakani MR, Wanek W, Gollner G, Friedel JK. 2013.
Carbon isotope discrimination and water use efficiency relationships of alfalfa genotypes
under irrigated and rain-fed organic farming. *European Journal of Agronomy* **50**: 82-89.
Nelson JA, Carvalhais N, Migliavacca M, Reichstein M, Jung M. 2018. Water-stress-induced

459   breakdown of carbon–water relations: indicators from diurnal FLUXNET patterns.
460   *Biogeosciences* **15**: 2433-2447.

461 **Peddinti SR, Kambhammettu BVNP, Rodda SR, Thumaty KC, Suradhaniwar S. 2019.** Dynamics
462   of Ecosystem Water Use Efficiency in Citrus Orchards of Central India Using Eddy
463   Covariance and Landsat Measurements. *Ecosystems* **23**: 511-528.

464 **Pieruschka R, Huber G, Berry JA. 2010.** Control of transpiration by radiation. *Proceedings of the*
465   *National Academy of Sciences* **107**: 13372-13377.

466 **Sinclair TR, Ghanem ME. 2020.** Plant□based predictions of canopy transpiration instead of
467   meteorological approximations. *Crop Science* **60**: 1133-1141.

468 **Sinclair TR, Hammer GL, van Oosterom EJ. 2005.** Potential yield and water-use efficiency benefits
469   in sorghum from limited maximum transpiration rate. *Functional Plant Biology* **32**: 945-952.

470 **Sinclair TR, Messina CD, Beatty A, Samples M. 2010.** Assessment across the United States of the
471   Benefits of Altered Soybean Drought Traits. *Agronomy Journal* **102**: 475-482.

472 **Sinclair TR, Tanner CB, Bennett JM. 1984.** Water-Use Efficiency in Crop Production. *Bioscience*:
473   36-40.

474 **Sinclair TR, Vadez V. 2012.** The future of grain legumes in cropping systems. *Crop and Pasture*
475   *Science* **63**.

476 **Song Q, Xiao H, Xiao X, Zhu X-G. 2016.** A new canopy photosynthesis and transpiration
477   measurement system (CAPTS) for canopy gas exchange research. *Agricultural and Forest*
478   *Meteorology* **217**: 101-107.

479 **Steduto P, Hsiao TC, Fereres E, Raes D. 2012.** *Crop yield response to water*: FAO Rome.

480 **Tanner CB, Sinclair TRJltewuicp 1983.** Efficient water use in crop production: research or re-search?
481   *Limitations to efficient water use in crop production*.

482 **Taylor HM, Jordan WR, Sinclair TR, Unger PW, Sneed TV, Jordan WR, Jensen R. 1983.**
483   *Limitations to efficient water use in crop production*: Limitations to efficient water use in crop
484   production.

485 **Vadez V, Kholova J, Medina S, Kakkera A, Anderberg H. 2014.** Transpiration efficiency: new
486   insights into an old story. *Journal of Experimental Botany* **65**: 6141-6153.

487 **Xu P, Moshelion M, Wu X, Halperin O, Wang B, Luo J, Wallach R, Wu X, Lu Z, Li G. 2015.**
488   Natural variation and gene regulatory basis for the responses of asparagus beans to soil
489   drought. *Frontiers in Plant Science* **6**: 891.

490 **Yang H, Shukla MK, Mao X, Kang S, Du T. 2019.** Interactive Regimes of Reduced Irrigation and
491   Salt Stress Depressed Tomato Water Use Efficiency at Leaf and Plant Scales by Affecting Leaf
492   Physiology and Stem Sap Flow. *Frontiers in Plant Science* **10**: 160.

493 **Yang YJ, Bi MH, Nie ZF, Jiang H, Liu XD, Fang XW, Brodribb TJ. 2021.** Evolution of stomatal
494   closure to optimize water-use efficiency in response to dehydration in ferns and seed plants.
495   *New Phytologist* **230**: 2001-2010.

496 **Zhao Y, Chen S, Shen S. 2013.** Assimilating remote sensing information with crop model using
497   Ensemble Kalman Filter for improving LAI monitoring and yield estimation. *Ecological*
498   *Modelling* **270**: 30-42.