Structural and biochemical insights into heterotetramer formation of
oncogenic K-Ras4B^{G12V} and Rgl2, a RalA/B activator

Mishal Tariq[1], Teppei Ikeya[2], Naoyuki Togashi[2], Louise Fairall[1,3], Carlos Bueno-Alejo[3], Shun
Kamei[2], Beatriz Romartinez-Alonso[1,3], Miguel Angel Muro Campillo[1], Andrew J. Hudson[3,4], Yutaka
Ito[2], John W.R. Schwabe[1,3], Cyril Dominguez[1,3], and Kayoko Tanaka[1]

1. Department of Molecular and Cell Biology, University of Leicester

2. Department of Chemistry, Tokyo Metropolitan University

3. Leicester Institute of Structure and Chemical Biology, University of Leicester

4. Department of Chemistry, University of Leicester

*Paste corresponding author name(s) here.

*Kayoko Tanaka

Email: kt96@le.ac.uk

Author Contributions: M.T., T.I., Y.I., C.D. and K.T. designed research. M.T., T.I., N.T., C.B.-A.,
S.K., M.A.M.C., C.D. and K.T. performed research. M.T., T.I., N.T., L.F., C.B.-A., S.K., B.R.-A, A.J.H.,
Y.I., J.W.R.S., C.D. and K.T. analyzed data. M.T., T.I., C.D. and K.T. wrote the manuscript with
feedback from all the authors.

Competing Interest Statement: The authors declare that they have no conflicts of interest with
the contents of this work.

Classification: Biochemistry, Structural Biology

Keywords: Ras signaling, RalA/B pathway, Structural Biology

1 Abstract

About a quarter of total human cancers carry mutations in Ras isoforms. Accumulating evidence
suggests that small GTPases, RalA and RalB, and their activators, Ral guanine nucleotide exchange
factors (RalGEFs), play an essential role in oncogenic Ras-induced signalling. We studied the
interaction between human K-Ras4B and the Ras binding domain (RBD) of Rgl2 (Rgl2^{RBD}), one of
the RBD-containing RalGEFs. We show that the G12V oncogenic K-Ras4B mutation increases the
affinity with Rgl2^{RBD}. The crystal structure of the K-Ras4B^{G12V}: Rgl2^{RBD} complex shows a 2:2
heterotetramer where the Switch I and Switch II regions of each K-Ras^{G12V} interact with both
Rgl2^{RBD} molecules. This structural arrangement is highly similar to the H-Ras^{E31K}:RALGDS crystal
structure and is distinct from the well-characterised Ras:RAF^{RBD} complexes. Importantly, the G12V
mutation was found at the dimer interface of K-Ras4B^{G12V} with its partner. Solution state NMR
and mass photometry analyses support the heterotetramer formation. Our study reveals a
distinct mode of Ras:effector complex formation by RalGEFs, and offers a possible mechanistic
explanation for how the oncogenic K-Ras4B^{G12V} hyperactivates the RalA/B pathway.

Main Text

Introduction

Ras belongs to a family of small G protein that switches between two states, GDP-bound form

(Ras[GDP]) and GTP-bound form (Ras[GTP]), and regulates a wide range of cellular activities (Cox and

Der, 2010). Guanine nucleotide exchange factors (GEFs), typically activated by growth factor

signalling, mediate the conversion of Ras[GDP] to Ras[GTP], whilst the GTP-bound status lasts only

transiently as the intrinsic GTPase activity, aided by the GTPase activating Proteins (GAPs),

hydrolyses the bound GTP into GDP (Vetter and Wittinghofer, 2001). Ras acts as a signalling hub

where the Ras[GTP], but not Ras[GDP], physically interacts with multiple Ras effectors, which then

transmit the signal to downstream molecules, including ERK, Akt and small G proteins RalA and

RalB (Simanshu et al., 2017).

There are three human *RAS* genes, *K-RAS*, *N-RAS* and *H-RAS*, and as *K-RAS* is alternatively spliced

at exon 4, the three *RAS* genes produce four Ras isoforms K-Ras4A, K-Ras4B, N-Ras and H-Ras

(Castellano and Santos, 2011). Among them, *K-RAS4B* typically represents more than half of all

RAS transcripts (Newlaczyl et al., 2017). The Catalogue Of Somatic Mutations In Cancer

(COSMIC) shows that about 25% of all cancers carry mutations in *RAS* genes, and *K-RAS* is

responsible for about 70% of these mutations (COSMIC, v.95). Hence, it is vital to obtain more

insights into how the oncogenic K-Ras signalling leads to cancers.

Extensive earlier biochemical studies revealed that oncogenic *RAS* mutations cause a reduction

of the GTP hydrolysis rate, generating an increased population of Ras[GTP] (Bollag and McCormick,

1991; Der et al., 1986; Gibbs et al., 1984; Manne et al., 1985; McGrath et al., 1984; Scheffzek et

al., 1997). This likely leads to an overactivation of the downstream signalling pathways and

therefore is considered the major cause of the K-Ras oncogenicity. In addition, recent

biochemical, structural and molecular modelling studies indicate that oncogenic *RAS* mutations

affect the interaction kinetics with the effector molecules and may trigger a biased

overactivation of a set of effectors (Hunter et al., 2015; Mazhab-Jafari et al., 2015; Pantsar et al.,

2018; Smith and Ikura, 2014). Therefore, obtaining more insights into the Ras-effector

interaction mechanism is essential to understanding the oncogenic-Ras mediated tumorigenesis

process.

Among Ras effectors, Raf kinase and PI3K, which ignite ERK and Akt signalling respectively, have

attracted much attention, especially because they were found not only to be activated by

oncogenic Ras but also to be able to carry oncogenic mutations themselves (Chalhoub and

Baker, 2009; Maurer et al., 2011). However, accumulating evidence suggests that the

misregulation of Ral GTPases, RalA and RalB, rather than ERK and Akt signalling pathways, may

be the initial trigger of oncogenic-Ras induced tumorigenesis. For example, in the oncogenic K-

ras(G12D) knock-in mouse model, although neoplastic and developmental defects were

observed, hyper-activation of ERK or Akt was undetected (Tuveson et al., 2004). In humans, ERK

hyper-activation is often missing in cancer cell lines and tissues with oncogenic Ras mutations,

whereas RalA and RalB are essential in oncogenic-Ras induced cell proliferation, motility and

tumorigenesis (Campbell et al., 2007; Lim et al., 2005; Lim et al., 2006; Luo and Sharif, 1999;

Miller et al., 1997; Yip-Schneider et al., 1999; Yip-Schneider et al., 2001; Zago et al., 2018).

Therefore, a biased misregulation of the Ral GTPases may be the critical feature behind Ras

oncogenicity. Indeed, NMR-based effector competition assays suggested the intriguing

possibility that the oncogenic Ras molecules may develop an altered effector preference leading

to a biased hyperactivation of RalA (Smith and Ikura, 2014).

Activation of RalA and RalB is mediated by guanine nucleotide exchange factors (RalGEFs). There

are eight RalGEFs reported for humans, and four of them, RALGDS, Rgl1, Rgl2 and Rgl3, have a

Ras-association (RA) domain (in this work, we generically call it Ras Binding Domain (RBD))

responsible for Ras-binding (Apken and Oeckinghaus, 2021). RALGDS was the first to be

identified as a Ras effector among these RalGEFs and is by far the most studied (Neel et al.,

2011). RALGDS is one of the first effector molecules that was crystallised together with an active

Ras. Rat RALGDS was co-crystalised with human H-Ras harbouring an E31K mutation, which

helped complex formation (Huang et al., 1998). A recent modelling approach, integrating

proteomic data of Ras, its 56 effectors and their interaction affinities, predicts that Rgl2 would

generate the highest concentration of Ras:effector complex among RalGEFs in 28 out of 29

healthy human tissues (Catozzi et al., 2021). Furthermore, a critical contribution of Rgl2 in

oncogenic Ras induced tumour phenotypes has been reported (Vigil et al., 2010). However, the

Ras:Rgl2 complex interface has not been studied.

In this report, we examined the mode of interaction between human K-Ras4B and the RBD of

Rgl2 (Rgl2^{RBD}). We observed striking affinity increase of the interaction of K-Ras4B with

theRgl2^{RBD} upon introduction of G12V oncogenic mutation. Our crystal structure of K-

Ras4B^{G12V}:Rgl2^{RBD} complex shows a heterotetramer formation, highly similar to the reported H-

Ras^{E31K}:RALGDS^{RBD} complex, but distinct from other Ras:effector complexes including Ras:Raf1

kinase. Strikingly, the G12V oncogenic mutation is located at the dimer interface of K-Ras4B^{G12V}

with its homodimeric partner. Solution NMR analyses support the heterotetramer formation

and mass photometry confirms the presence of the heterotetramer. Our findings provide an

interesting possibility that K-Ras4B^{G12V} oncogenicity might be due to its capability to form a

heterotetramer with Rgl2.

11
12
13   **Results**

14
Active K-Ras4B^{G12V} has a stronger affinity to Rgl2^{RBD} than K-Ras4B^{WT}

Throughout this study, we used recombinant bacteria constructs of human K-Ras4B lacking the

C-terminal hyper-variable region and the Rgl2 RBD (Rgl2^{RBD}) consisting of amino acids 648 – 735

of human Rgl2. The binding of Rgl2^{RBD} and K-Ras4B^{G12V} was confirmed for the active K-Ras4B^{G12V}

loaded with a non-hydrolysable GTP analogue (Guanosine-5'-[(β,γ)-imido]triphosphate

(GMPPNP)), by GST-pulldown assays (Figure 1-figure supplement 1A and 1B). To examine

whether the binding mode of Rgl2^{RBD} differs between K-Ras4B^{WT} and K-Ras4B^{G12V}, the binding

affinities were determined by biolayer interferometry (BLI). We noticed that both K-Ras4B^{WT} and

K-Ras4B^{G12V}, purified at 4^{o}C at all times as described in the Materials and Methods, retained GTP

as the major bound guanine nucleotide, and we confirmed that the K-Ras4B^{WT} and K-Ras4B^{G12V}

samples that were subjected to the incubation condition of the BLI assay were still associated

with GTP (Figure 1A). Therefore, we used these GTP-bound samples without loading GMPPNP so

that we could minimize possible artefacts caused by GMPPNP. The on/off kinetics between K-

Ras4B and Rgl2 were fast for both wildtype and G12V mutant (Figure 1B lower panels), and the

sensorgram curves could not be fitted to the pre-programmed curve fitting models (Data

Analysis HT Software, Sartorius). Therefore, we deduced the equilibrium dissociation constant

(K_D) using steady-state analysis (Figure 1B upper panels). Strikingly, the affinity between Rgl2^{RBD}

and K-Ras4B was substantially increased when K-Ras4B carried an oncogenic mutation, G12V,

where Glycine at position 12 is replaced with Valine. This is consistent with the size-exclusion

chromatography analysis conducted under the same condition, showing that Rgl2^{RBD} does not

form a stable complex with K-Ras4B^{WT} in contrast to K-Ras4B^{G12V} (Figure 1-figure supplement 1C

and 1D). The circular dichroic (CD) spectra for K-Ras4B^{WT} and K-Ras4B^{G12V} samples across the

increasing temperature from 20^{o}C to 90^{o}C also showed improved structural stability for K-

Ras4B^{G12V}, which might contribute to the higher affinity towards Rgl2^{RBD} (Figure 1 C-E).

A 2:2 tetramer of K-Ras4B^{G12V}:Rgl2^{RBD} complex in the crystal structure

We conducted crystallization trials to obtain structural insights into the K-Ras4B:Rgl2^{RBD}

complex. To purify the complex, a mixture of K-Ras4B and Rgl2^{RBD} was loaded onto a size-

exclusion column. We could obtain the K-Ras4B^{G12V}:Rgl2^{RBD} complex in a peak that eluted earlier

than K-Ras4B^{G12V} alone (Figure 1-figure supplement 1C). However, we were unable to obtain an

elution peak for K-Ras4B^{WT}:Rgl2^{RBD} complex (Figure 1-figure supplement 1D). The inability of K-

Ras4B^{WT} to form a stable complex that survives the gel filtration chromatography is consistent

with the relatively low affinity between K-Ras4B^{WT} and Rgl2^{RBD} seen in Figure 1B.

Crystals of the K-Ras4B^{G12V}:Rgl2^{RBD} complex were obtained which diffracted at a resolution of

3.07Å (Table 1). After molecular replacement with the structure of the human H-Ras^{E31K}:rat

RALGDS^{RBD} complex (PDB ID 1LFD) (Huang et al., 1998), the K-Ras4B^{G12V}:Rgl2^{RBD} structure was

manually built using Coot (Emsley et al., 2010) and was refined with PHENIX (Torices and

Munoz-Pajares, 2015)(Figure 2A). The overall arrangement of K-Ras4B^{G12V}:Rgl2^{RBD} crystal

structure is similar to H-Ras^{E31K}:RALGDS^{RBD} crystal structure (PDB ID 1LFD (Huang et al., 1998))(

Figure 2-figure supplement 1A-C). The complex forms a 2:2 heterotetramer where β2 (within

Switch I) of Ras and β2 of Rgl2^{RBD} generate a continuously extended β-sheet, along with

interaction at Switch II of the same Ras molecule with the second Rgl2^{RBD} (Figure 2B, Figure 2-

figure supplement 2A). In both K-Ras4B^{G12V}:Rgl2^{RBD} and H-Ras^{E31K}:RALGDS^{RBD} cases, the Ras-Ras

interface is formed within the region spanning amino acids 1-90 where K-Ras4B^{G12V} and H-Ras^{E31K}

share 100% amino acid sequence identity except for the point mutations G12V and E31K.

The K-Ras4B^{G12V}:Rgl2^{RBD} heterotetramer complex is stabilized by a network of hydrogen bonds

and hydrophobic interactions (Figure 2B, C and D, Figure 2-figure supplement 2A), leading to a

complimentary surface charges (Figure 2-figure supplement 2B). K-Ras4B^{G12V} residues within the

Switch I interact with residues in β1, β2 and α1 of Rgl2^{RBD}. These interactions include a salt

bridge between E37 of K-Ras4B[G12V] Switch I and R653 of Rgl2[RBD] (Figure 2B and Figure 2-figure

supplement 2A). The Switch II of the same K-Ras4B[G12V] molecule also contributes to complex

formation by interacting with the second Rgl2[RBD] through residues in β2 and α1 (Figure 2B and

Figure 2-figure supplement 2A). Meanwhile, two K-Ras4B[G12V] molecules have direct contact

through the Switch I, Switch II and α3 (Figure 2C and Figure 2-figure supplement 2A). Strikingly,

the V12 residue of the oncogenic G12V mutation is within 4 Å of the ring of Y32 of the

neighboring K-Ras4B[G12V] contributing to the K-Ras4B[G12V]: K-Ras4B[G12V] interface (Figure 2C and

Figure 2-figure supplement 2A). The β1 of both Rgl2[RBD] molecules run anti-parallel to each other,

interacting through various hydrophobic interactions and hydrogen bonds at both side-chain

and backbone levels (Figure 2D).

Solution NMR analyses support the K-Ras4B[G12V]:Rgl2[RBD] heterotetramer formation

The K-Ras4B[G12V]:Rgl2[RBD] complex was further analyzed in solution. First, the solution structure of

the Rgl2[RBD] construct was determined by solution NMR (Figure 3A) (Table 2). The structure

retained the ββαββαβ ubiquitin-fold structure, a common feature for the RBDs (Kiel and

Serrano, 2006). Overall, it is similar to Rlf, the mouse homologue of human Rgl2 (PDB ID

1RLF)(Esser et al., 1998). However some differences can be observed, even though the primary

sequences are similar (Figure 3-figure supplement 1A). The structures of the Rgl2[RBD] molecules

in the K-Ras4B[G12V]:Rgl2[RBD] complex crystal structure are highly comparable with the structure of

the NMR solution structure of free Rgl2[RBD] indicating that the complex formation causes

relatively small structural changes (Figure 3B).

The K-Ras4B[G12V]:Rgl2[RBD] complex was next analyzed in solution by NMR chemical shift

perturbation. Two-dimensional (2D) ^{15}N–^{1}H-heteronuclear single quantum coherence (HSQC)

spectrum were measured of ^{15}N-labelled Rgl2[RBD] sample, in the absence or presence of

increasing amount of non-labelled K-Ras4B[G12V] (Figure 3C, Figure 3-figure supplement 1 B and C).

Chemical shift perturbations were observed for most of the Rgl2[RBD] residues, in agreement with

K-Ras4B[G12V]:Rgl2[RBD] complex formation in solution (Figure 3-figure supplement 1 B and C). Most

of the Rgl2[RBD] residues at the K-Ras4B[G12V]:Rgl2[RBD] and Rgl2[RBD]:Rgl2[RBD] interfaces in the crystal

structure showed a greater change (either decreased or increased) in their NMR signal

intensities, indicating their participation to the complex formation (Figure 3C). Notably, signals

from the Rgl2[RBD] residues in β1 at the N-terminal end made the largest increase. An explanation

may be that the highly flexible N-terminal region of free Rgl2 (Figure 3A) becomes confined into

a more defined structure upon complex formation as observed in the crystal structure.

It has been shown that a lysine residue of RALGDS, that corresponds to K686 of Rgl2, plays an

essential role in H-Ras[E31K]:RALGDS complex formation (Huang et al., 1998)(Figure 3-figure

supplement 2 A). In the K-Ras4B[G12V]:Rgl2[RBD] complex crystal structure, K686 forms a hydrogen

bond with D33 of K-Ras4B[G12V] (Figure 3-figure supplement 2 B). K686 was also among the

residues that showed a substantial signal decrease in the NMR titration assays (Figure 3C). We

therefore mutated K686 into an alanine and as expected the K686A mutant lost the ability to

interact with K-Ras4B[G12V] (Figure 3-figure supplement 2 C), confirming the important role of

K686 in the interaction.

We next examined the NMR chemical shift perturbation of K-Ras4B[G12V] in solution by 2D ^{15}N–^{1}H-

HSQC, in absence or presence of non-labelled Rgl2[RBD]. We noticed that when loaded with

GMPPNP, the number of detectable peaks of the ^{15}N labelled K-Ras4B[G12V] signal was

substantially decreased compared to the number of signals obtained from the GDP-loaded

sample (Figure 3-figure supplement 3) as previous studies reported (Ito et al., 1997; Menyhard

et al., 2020). Consequently, most of the K-Ras4B[G12V] residues at the K-Ras4B[G12V]:Rgl2[RBD] complex

interface could not be detected, except for K42 and K88, both of which were showing

substantially decreased signal intensities, in agreement with the expected participation in the

complex formation (Figure 3-figure supplement 4 C). Furthermore, upon Rgl2[RBD] titration, most

of the ^{15}N K-Ras4B[G12V] signals were reduced (Figure 3-figure supplement 4 A-C). This overall

signal reduction was consistent with broadening arising as a result of the large complex size.

Mass photometry confirms the presence of a heterotetramer of K-Ras4B[G12V]:Rgl2[RBD] complex

To confirm that K-Ras4B[G12V]:Rgl2[RBD] forms a heterotetramer in solution, we analyzed the

complex by mass photometry, a label free technique that has recently been adapted to measure

the mass of biomolecules in solution (Young et al., 2018). Our commercially-available instrument

(see Materials and methods) can measure accurately masses in the range 30-5000 kDa, and is

suitable to differentiate between K-Ras4B[G12V]:Rgl2[RBD] heterodimer (31kDa) or heterotetramer

(62kDa). The K-Ras4B[G12V]:Rgl2[RBD] complex was freshly prepared by size exclusion

chromatography and the peak fraction that contained both proteins at 1:1 ratio was used for

the measurement (Figure 4A). The histogram of the frequency counts for this sample showed a

peak, that after calibration with the proper standard (Native Mark) , corresponds to a

biomolecule with molecular weight of 68 KDa (Figure 4B). As the expected molecular weight of

the K-Ras4B^{G12V}:Rgl2^{RBD} heterotetramer is about 62 KDa, the result strongly indicates that at

least part of the complex population exists as the 2:2 heterotetramer in solution.

5
6
7   **Discussion**
8
We conducted interaction studies of human K-Ras4B and Rgl2^{RBD} to obtain structural insights

into the oncogenic Ras-dependent activation of the RalA/B pathway. Using X-ray crystallography

and the crystal structure of human H-Ras^{E31K} and rat RALGDS complex (PDB ID 1LFD) as a

molecular replacement template (Huang et al., 1998), we reveal a heterotetramer formation

stabilised by the oncogenic G12V mutation. The presence of heterotetramer in solution was

further confirmed by mass photometry.

Mode of interaction between Ras and Rgl2/RALGDS is distinct from other Ras:RBD

interactions

The key contributor of the heterotetramer formation seen in this Ras:RalGEF complex, K-

Ras4B^{G12V}:Rgl2^{RBD}, and the previously published H-Ras^{E31K}:RALGDS^{RBD} is the interaction between

the Switch II region of Ras and the second RBD molecule (Figure 2B and Figure 2-figure

supplement 3A). This feature is distinct from the previously reported Ras:RBD crystal structures,

where the Switch II region is either not participating in the interaction (as seen in Ras:Raf1^{RBD}

(PDB ID 6VJJ)(Tran et al., 2021)) or is interacting with an RBD that also interacts with the Switch I

region of the same Ras molecule (as seen in Ras:PI3Kγ^{RBD} (PDB ID 1HE8)(Pacold et al., 2000), Ras:

PLCε^{RBD} (PDB ID 2C5L)(Bunney et al., 2006), Ras:RASSf5^{RBD} (PDB ID 3DDC)(Stieglitz et al., 2008)),

hence forming a Ras:RBD heterodimer, rather than a heterotetramer, in each crystal lattice

(Figure 2-figure supplement 3B). One interesting case is a recently reported structural study of

Rgl1, yet another RalGEF (Eves et al., 2022). Crystal structures of Rgl1 in complex with either K-

Ras^{WT} or K-Ras^{G12V} (PDB ID 7SCW and 7SCX, respectively), showed that one Rgl1^{RBD} interacts with

one K-Ras molecule through both Switch I and Switch II, as other heterodimer examples shown

in Figure S8B. It will be interesting to further examine whether the K-Ras:Rgl1^{RBD} may have the

capacity to form a heterotetramer as a common feature for Ras:RalGEF complexes.

Oligomerisation of K-Ras:effector complexes

Accumulating evidence suggests that K-Ras is capable of forming a dimer in solution even in the

absence of effectors (Andreadelis et al., 2022; Ingolfsson et al., 2022; Jang et al., 2016; Lee et al.,

2021; Lee et al., 2020; Muratcioglu et al., 2015; Ozdemir et al., 2022; Packer et al., 2021; Prakash

et al., 2017; Sarkar-Banerjee et al., 2017). The physiological importance of Ras dimerisation and

its potential to be a therapeutic target was highlighted by a G12D-specific inhibitor, BI2852,

which may cause artificial dimerisation and block the protein function (Kessler et al., 2019; Tran

et al., 2020). Furthermore, Ras forms nanoclusters in the membrane *in vivo* and *in vitro*

(Lakshman et al., 2019; Plowman et al., 2005; Prior et al., 2003; Weise et al., 2011; Zhou et al.,

2014). The modes of these Ras oligomer formation are dependent on various parameters

including membrane lipid compositions, Ras nucleotide binding status, availability of Ras

effectors and actin cytoskeleton and are expected to be context dependent.

Previously it has been suggested that dimerisation status may be categorised into four classes

based on the utilized α helices and β sheets as follows; (i) $\alpha4/\alpha5$ (Andreadelis et al., 2022; Jang

et al., 2016; Lee et al., 2021; Lee et al., 2020; Packer et al., 2021; Prakash et al., 2017), (ii) $\alpha3/\alpha4$

(Jang et al., 2016; Muratcioglu et al., 2015; Prakash et al., 2017), (iii) α/β (Lee et al., 2021), and

(iv) β/β (Jang et al., 2016; Muratcioglu et al., 2015) (Figure 2-figure supplement 4). Importantly,

the K-Ras4B^{G12V}:K-Ras4B^{G12V} interface that we observe in the K-Ras4B^{G12V}:Rgl2^{RBD} heterotetramer

does not belong to any of these categories and uses unstructured regions of Switch I, Switch II,

V12 in the P-loop and K88 (Figure 2, Figure 2-figure supplement 1 and 4). Interestingly, this K-

Ras4B^{G12V}:K-Ras4B^{G12V} interface is similar to the spatial arrangement of symmetry mates of

human H-Ras^{WT} crystal (PDB ID 5P21) (Pai et al., 1990) (Figure 2-figure supplement 5), indicating

that this Ras:Ras contact is likely stabilized by Rgl2^{RBD}.

Currently, we do not have information as to the orientation of the K-Ras4B^{G12V}:Rgl2^{RBD} in relation

to the plasma membrane (PM). The Rgl2 molecules could be away from the plasma membrane

(PM-away conformation), or could be in direct contact with the plasma membrane (PM-

proximity conformation) (Figure 5). In either case, the complex status is distinct from the case

for Ras:Raf1 kinase complex which can be formed onto the existing Ras:Ras homodimer of the

$\alpha4/\alpha5$ interface (Packer et al., 2021). If the majority of the cellular K-Ras4B^{G12V} molecules exist

as the $\alpha4/\alpha5$ homo-dimer at the PM, what drives the K-Ras4B^{G12V}:Rgl2^{RBD} heterotetramer

formation by causing a major structural change is an important future question.

Direct involvement of V12 in the K-Ras4B:Rgl2^{RBD} complex formation and oncogenicity

Strikingly, the K-Ras4B^{G12V}:K-Ras4B^{G12V} interface of the K-Ras4B^{G12V}:Rgl2^{RBD} tetramer complex

involves the residue V12, an oncogenic amino-acid substitution occurring in about 28% of

mutated *KRAS* cases (COSMIC). This G12V substitution substantially increases the affinity

towards Rgl2^{RBD} compared to the wildtype (Figure 1B). In agreement with this observation, our

attempt to biochemically purify the K-Ras4B^{WT}:Rgl2^{RBD} complex has been unsuccessful as K-

Ras4B^{WT} and Rgl2^{RBD} eluted in separate fractions when subjected to size exclusion

chromatography (data not shown). In the K-Ras4B^{G12V}:Rgl2^{RBD} tetramer complex, the presence of

V12 creates a larger hydrophobic pocket together with Y32, compared to the wildtype case of

G12, seen in the H-Ras^{WT} crystal (PDB ID 5P21) or in the H-Ras^{E31K}:RALGDS^{RBD} crystal (PDB ID

1LFD) (Figure 2-figure supplement 5 and 6). To our knowledge, the K-Ras4B^{G12V}:Rgl2^{RBD} complex

serves as the first example where a structural explanation is provided for an oncogenic-Ras

driven stabilization of a Ras:effector complex. Interestingly, the E31K substitution, used in the H-

Ras^{E31K}:RALGDS crystal structure analysis to stabilize the complex (Huang et al., 1998) (Figure 2-

figure supplement 6E), has been reported in cancer samples with *HRAS* mutation (COSMIC).

Therefore, the capability of Ras to form a stable complex with RalGEFs may be directly linked

with Ras oncogenicity. It is interesting to note that in the previously proposed Ras:Raf1 kinase

heterotetramer complex derived from SAXS data (Packer et al., 2021), the position of the amino

acid residue 12 is relatively distant from the interacting surfaces. An interesting speculation

could be that the oncogenic substitution mutations at glycine 12 might have a greater impact on

RalGEF-mediated signalling than Raf kinase-mediated signalling (Figure 5).

Conclusion

To summarize, our work demonstrates an increased affinity of K-Ras4B^{G12V} oncogenic mutant

with a RalGEF, Rgl2, and provides a structural explanation by revealing the complex interface.

The information will open the way to target oncogenic-*KRAS*-induced tumorigenesis by novel

strategies, including interfering molecules for the newly identified interfaces.

Materials and Methods

Plasmid constructs

The RBD of human Rgl2 was obtained by amplifying a cDNA fragment encoding the position 643

– 740 of the human Rgl2 by PCR with a pair of primers (5'

TACTTCCAATCCATGGGGCCAGGGGCCTCTGATTGCCG3') and (5' TATCCACCTTTACTGTCA

TGTAGCAGTAGAGGACCTTCGCCGCTGC 3') using human cDNA prepared from hTERT RPE-1 cells

using GoScript Reverse Transcription System (Promega) following the manufacturer's

instruction. The amplified Rgl2^{RBD} fragment was cloned into pLEICS2 vector (PROTEX, University

of Leicester), which contains a glutathione S-transferase (GST) affinity tag and a Tobacco Etch

virus (TEV) potease cleavage site at the N-terminal end of the Rgl2^{RBD}, using In-Fusion HD EcoDry

enzyme (Takara Bio, #638915), following the manufacturer's instruction. A point mutation,

K686A, was introduced by site-directed mutagenesis using a pair of primers (5'

ATCAGTCGTGTCCTTAAGGCAAACAATCGTGACTCT 3') and (5'

CAGAGTCACGATTGTTTGCCTTAAGGACACGACTGAT 3') by following the QuikChange^{TM} method

(Agilent Technologies, La Jolla, CA, USA).

A cDNA fragment encoding the position 1 – 169 (C-terminal truncated) of the human wildtype K-

Ras4B isoform was amplified by PCR with a pair of primers (5' TACTTCCAATCCATG

ACTGAATATAAACTTGTGGTAGTTGGAGCTG') and (5' TATCCACCTTTACTGTCA

CTTTTCTTTATGTTTTCGAATTTCTCGAACTAATGTATAG 3') using human cDNA prepared from

hTERT RPE-1 cells as described above. The produced fragment was cloned in pLeics1 plasmid

(PROTEX, University of Leicester), which introduces a His6 tag and a TEV cleavage site at the N-

terminus. Site directed mutagenesis was conducted to introduce the oncogenic G12V mutation

using a pair of DNA oligos (5' AGTTGGAGCTGTTGGCGTAGGCAAGAGTGCC 3') and (5'

GTCAAGGCACTCTTGCCTACGCCAACAGCTCCAACTAC 3') by following the QuikChange^{TM} method

(Agilent Technologies, La Jolla, CA, USA).

Protein expression

DE3 Rosetta cells (Novagen) carrying expression plasmids were grown at 37^{o}C in TY media until

OD600 reached about 0.6. Then protein expression was induced by adding Isopropyl β-D-1-

thiogalactopyranoside (IPTG) to a final concentration of 0.5 mM and keeping the culture at 18^{o}C

overnight in a shaking incubator. Cells were collected by centrifugation and resuspended in

either Sl1 buffer (Tris 20mM (pH 7.65), NaCl 150 mM, 5 mM imidazole) for His-tagged KRas or in

PBS-NaCl buffer (237 mM NaCl, 2.7 mM KCl, 8 mM Na_2HPO_4, and 2 mM KH_2PO_4, pH 7.4) for Rgl2

and RalGDS. The cell suspensions were then stored at -80^{o}C.

Protein expression of stable isotope labelling K-Ras4B^{G12V} and RBD of Rgl2 for NMR measurement

The gene encoding human K-Ras4B^{G12V} and Rgl2^{RBD} were constructed into the expression vector

pGHL9 and pLEICS2 over-expressed in *Escherichia coli* (*E.coli*) strain BL21 (DE3) and Rosetta,

respectively. Uniformly ^{13}C, ^{15}N-labeled protein was obtained by growing bacteria at 37 °C in M9

minimal media, containing [$^{13}C_6$]-glucose and $^{15}NH_4Cl$ (Isotec) as the sole carbon and nitrogen

source, supplemented with 20 mM $MgSO_4$, 0.1 mM $CaCl_2$, 0.4 mg/ml thiamin, 20 μM $FeCl_3$, salt

mix [4 μM $ZnSO_4$, 0.7 μM $CuSO_4$, 1 μM $MnSO_4$, 4.7 μM H_3BO_3], and 50 mg/L ampicilin. K-

Ras4B^{G12V} NMR sample was prepared essentially as described previously (Ito et al., 1997).

Protein expression of Rgl2^{RBD} was induced by adding 119mg/L IPTG at an OD 600 nm of 0.5.

After 18 h of further growth, cells were harvested, and washed with a pH 7.5 lysis buffer [50 mM

Tris-HCl, 25% sucrose, and 0.01% NP-40]. Uniformly ^{15}N-labeled K-Ras4B^{G12V} and Rgl2^{RBD} were

produced by the identical steps unless growing cells in M9 medium containing [$^{12}C_6$]-glucose and

$^{15}NH_4Cl$ (Isotec).

Purification of GST-tagged Rgl2^{RBD}

Bacteria cell suspensions were thawed and supplemented with Triton X-100 to a final

concentration of 0.1 % (v/v). Cells were broken by a probe sonicator and insoluble materials

were removed by centrifugation. The supernatant was mixed with glutathione (GSH) beads (GE

Healthcare, #17-0756-01) and incubated for 20 min at 4^{o}C. The GSH beads were washed three

times with the PBS-NaCl buffer. the GST-Rgl2^{RBD} fusion protein was either eluted by the elution

buffer (50mM Tris-Cl (pH 8.0), 100mM NaCl, 5mM GSH) for GST pulldown experiments or the

Rgl2^{RBD} was separated from the GST tag through cleavage by tobacco etch virus (TEV) protease,

which was prepared as previously described (Kapust et al., 2001; Tropea et al., 2009), for

structural analysis. The obtained Rgl2^{RBD} or GST-Rgl2^{RBD} fusion proteins were concentrated using

a concentrator (10 kDa MWCO, Merck, Amicon Ultra centrifugal filters, #UFC901024) and

filtrated using a centrifugal filter unit (Milipore, Ultrafree, #UFC30GV00) before conducting size

exclusion chromatography (SEC) with a gel-filtration column (GE Healthcare, HiLoad Superdex

75) attached to an FPLC (Akta) system. SEC was carried out in a gel filtration buffer (20 mM Tris-

Cl (pH 7.65), 100 mM NaCl, 5mM MgCl2, 1mM tris (2-carboxyethyl) phosphine (TCEP)).

Purification of His-tagged K-Ras4B wildtype and oncogenic G12V mutant proteins

Bacteria cell suspensions were supplemented with Triton X-100 to a final concentration of 0.1 %

(v/v). Cells were broken by a probe sonicator and insoluble materials were removed by

centrifugation. The soluble cell lysates were applied on a Ni-sepharose excel (GE Healthcare,

#17-3712-01), packed in a column of 4 ml bed volume with Sl1 buffer buffer (20mM Tris, pH6.5,

150mM NaCl, 5 mM imidazole). The column was washed with 20 ml of SL1 Buffer, then with 20

ml of Sl3 Buffer (20mM Tris, pH6.5, 150mM NaCl, 6mM Imidazole) and finally with 15 ml of Sl4

Buffer (20mM Tris, pH6.5, 150mM NaCl, 10mM Imidazole). The His-tagged Kras protein was

eluted from the column by applying 10 ml of Elution buffer (50mM Tris, pH 7.65, 150mM NaCl,

200mM Imidazole), followed by 10 ml of 1 M imidazole. In order to remove the His6-tag at the

N-terminal end, TEV protease, prepared as previously described (Tropea et al., 2009), was added

to the elution fraction to about a 2 % molar ratio of the His6-K-Ras4B preparation and incubated

overnight at 4^{o}C. Nucleotide exchange of the purified K-Ras4B wildtype or G12V proteins was

carried out essentially as previously described (Ito et al., 1997). The proteins were diluted 10

times by the exchange buffer (20mM Tris-Cl (pH7.5) 1mM EDTA, 1mM TECP) and denatured by

the addition of EDTA to a final concentration of 5 mM. The sample was mixed with about 10

time molar excess of guanosine 5'-[β,γ-imido]triphosphate (GMMPPNP, SIGMA G0635) or

guanosine 5' –diphosphate (GDP, SIGMA G7127). The reaction was incubated at 37℃ for 20

minutes then put it on ice for 20 minutes. The protein was renatured by adding ice cold MgCl$_2$ to

a final concentration of 20 mM.

Purification of isotope labelling K-Ras4B^{G12V}

All the procedures described below were carried out at 4 °C unless otherwise stated. All the

isotope-labelled K-Ras4B^{G12V} samples were purified by the same step. The cells dispersed in the

lysis buffer was disrupted by sonication for 30 min on ice with hen egg lysozyme (0.1 mg/mL).

The cell debris was clarified by centrifugation at 14,000 g for 1 h. The supernatant was loaded

onto a 25 mL of DEAE-Sepharose Fast Flow (Cytiva) anion exchange equilibrated with buffer A

[50 mM Tris-HCl (pH 7.5), 1 mM MgCl$_2$, 1 mM dithiothreitol (DTT), 0.1 mM APMSF (FUJIFILM

Wako)]. After washing the column with buffer A until sufficiently low of UV absorption at 280

nm, the K-Ras4B^{G12V} protein was eluted by linearly increasing the concentration of KCl from 0 to

350 mM with a flow rate of 0.5 mL/min in buffer A. The fractions containing the target protein

were concentrated to 5 mL with Amicon Ultra-15 10 kDa (Merck). The concentrated sample was

loaded onto a 320 mL of HiLoad Superdex 75 (GE Healthcare Life Science) gel filtration with a

flow rate of 0.8 mL/min using FPLC systems (AKTA pure 25, GE Healthcare Life Science). The 5

mL sample concentrated from the fractions involving the target proteins with Amicon Ultra-15

10 kDa was loaded on Resource Q (GE Healthcare Life Science) anion-exchange column

equilibrated with buffer A using the FPLC systems. After washing the column with 30 mL of

buffer A, K-Ras4B^{G12V} was eluted by an KCl, the K-Ras4B^{G12V} protein was eluted by linearly

increasing the concentration of KCl from 0 mM to 350 mM with a flow rate of 1 mL/min in buffer

A. The purification of isotope labelled Rgl2^{RBD} was pefromed by the same step described above.

The purity of the K-Ras4B^{G12V} and Rgl2^{RBD} samples in each step was confirmed by SDS-PAGE.

Protein concentrations were determined by NanoDrop 2000 (ThermoFisher) measuring UV

absorption at 280 nm. K-Ras4B^{G12V} samples for NMR measurements were concentrated and

dissolved in NMR buffer A (90% ^1H_2O/ 10% ^2H_2O containing 20 mM Tris-HCl (pH 7.5), 100 mM

NaCl, 5 mM MgCl$_2$, 1 mM β-mercaptoethanol). Rgl2^{RBD} samples for NMR meassruments were

NMR buffer B (90% ^1H_2O/ 10% ^2H_2O containing 1 mM Na$_2$HPO$_4$-NaH$_2$PO$_4$ (pH 7.4), 150 mM NaCl).

K-Ras4B-Rgl2^{RBD} binding measurement using BLI

Octet Red96 (Sartorius) was used for Biolayer interferometry assays of KRas4B (G12V or WT)

and Rgl2^{RBD} interactions. Anti-GST biosensors (Sartorius #18-5096) were used to immobilise GST-

Rgl2^{RBD} (provided as a 13 µM solution in the binding reservoir well), and the baseline was

stabilised in BLI buffer (20mM Tris pH7.65, 0.5mM NaCl, 5mM MgCl, 1mM TCEP) for 200sec. As

a negative control, GST only (provided as a 13 µM solution in the binding reservoir well) was

used. Protein concentrations were determined by absorbance at 280 nm using the following

values; the extinction coefficient (ε) for KRas (WT) and KRas(GV) was estimated to be 19685 cm-

^{1} M $^{-1}$, by taking into account that the bound GTP adds 7765 cm^{-1} M $^{-1}$ (Smith and Rittinger,

2002), and the molecular weights (including GTP) were estimated to be 19856 and 19898,

respectively. The association of KRas4B was measured for 300sec in 6 serial dilutions

concentrations. For KRas (WT) the concentrations used ranged from 625 nm - 10 µM and for

KRas4B (G12V) the concentrations ranged from 200 nM – 6.4 µM. The dissociation steps were

measured in fresh BLI buffer for 300sec. For each assay a biosensor immobilised with GST only

and a sample well with only buffer instead of KRas4B (WT/G12V) was set up for double

referencing. The experiments were conducted at 23°C. The resulting data were processed using

Data Analysis HT Software (ver. 12.0.2.59)(FORTE BIO) and reference biosensor and reference

wells were subtracted from sample wells (double reference). Because of the fast on- and off-

rates, full kinetic analysis using a global 1:1 model resulted in low quality fitting. Therefore,

steady state curves were used to determine the K_D values.

Analysis of bound nucleotide

The nucleotide-binding status of K-Ras4B[WT] and K-Ras4B[G12V] were examined by denaturating the

proteins and detecting the released nucleotides (Smith and Rittinger, 2002). About 2 nmoles of

K-Ras4B molecules were adjusted to the volume of 200 µl with AKTA buffer. Add 12.5 µl of 10%

perchloric acid to denature the protein. Add 8.75 µl of 4M CH_3COONa, pH 4.0, to neutralise the

sample and centrifuge it to precipitate the denatured protein. The supernatant was analysed by

High-performance liquid chromatography (HPLC) using an ion-exchange column (Partisil 10 SAX

column, Whatman) equilibrated with 10 mM NH_4H2PO_4 (buffer A). The column was run with the

following gradient condition with 0.6 M NH_4H2PO_4 (buffer B) at a flow rate of 0.8 ml/min. Step1:

100% buffer A (0% buffer B) for 11 min, Step2: a gradient increase to 40% buffer B over 6 min,

Step3: a gradient increase to 50% buffer B over 23 min, Step4: a gradient increase to 100 %

buffer B over 1 min, Step 5: 100 % buffer B for 19 min, Step 6: a gradient decrease to 0% buffer

B (100 % buffer A) over 1 min, and Step 7: 100% Buffer A for 14 min. The nucleotides were

detected by 254 nm absorption. As a reference control, 1 µl of 10 mM GTP or GDP was diluted

to 200 µl AKTA buffer and was processed in the same manner as protein samples.

Crystallography

The purified and GMPPNP-loaded KRas4B (G12V) and Rgl2[RBD] were mixed in gel filtration buffer

and the complex was purified on SEC using a gel-filtration column (GE Healthcare, HiLoad

Superdex 75). The peak fractions containing both proteins in the 1:1 ratio were collected and

concentrated to set up crystallization screenings. Crystals of KRas4B (G12V) and Rgl2[RBD] were

obtained using sitting-drop vapour diffusion at room temperature, with 100nl of protein

(11.6mg/ml) against 100nl of crystallisation buffer (0.2M sodium/potassium phosphate ph 7.5,

0.1M HEPES pH 7.5, 22.5% v/v PEG smear Medium (Molecular Dimensions MD2-100-259), 10%

v/v glycerol. The crystals were frozen in liquid nitrogen with 20% glycerol as cryoprotectant.

Data were collected at Diamond beamline I04. AIMLESS (Evans and Murshudov, 2013) was used

for data reduction before obtaining phaser solution using the HRas-RALGDS[RBD] complex

structure (PDB ID 1LFD) as search model with PhaserMR (McCoy et al., 2007). The structure was

built using multiple rounds of refinements using PDBredo, REFMAC, PHENIX and COOT (Emsley

et al., 2010; Joosten et al., 2014; Murshudov et al., 2011; Torices and Munoz-Pajares, 2015). The

coordinates of the complex have been deposited to the Protein Data Bank (PDB) under access

code 8B69.

Circular dichroism (CD) spectroscopy

K-Ras4B(WT) and (G12V) proteins at a concentration of 20 µM were prepared in the CD buffer

(50 mM phosphate (pH 7.6), 1.5 mM Tris (pH 7.6), 5mM MgSO4, 7.5 mM NaCl, 0.375 mM

MgCl2), placed in a quartz cuvette of 0.1 cm path length and CD spectra were recorded at

wavelengths ranging from 195 to 250 nm using a Chirascan[TM]-plus CD spectrometer (Applied

Photophysics) at 20 °C. The melting curves of these proteins were examined at 220 nm at

temperatures ranging from 20°C to 90°C. Measurements were conducted at 1 °C increment.

NMR spectroscopy

K-Ras4B[G12V] NMR sample was prepared essentially as described previously (Ito et al., 1997). The

bacteria expression plasmid for H-Ras was modified to encode K-Ras4B[G12V] by site-directed

mutagenesis and used to produce K-Ras4B[G12V]. Loading of a GTP analogue, GMPPNP (Jena

Bioscience) was conducted essentially as previously described (Ito et al., 1997). Rgl2[RBD] sample

was prepared as described above. All NMR samples were measured in 20 mM Tris-Cl (pH 7.65),

100 mM NaCl, 5mM $MgCl_2$, and 0.1% β-mercaptoethanol at 303 K. All spectra were analysed

with the CcpNmr Analysis 2.5.1 software. Backbone chemical shifts of K-Ras4B[G12V] and Rgl2 have

been deposited to the BioMagResBank (BMRB) with accession ID 34754 and PDB with accession

ID 8AU4, respectively.

All NMR experiments were performed at 25°C probe temperature in a triple-resonance

cryoprobe fitted with a z-axis pulsed field gradient coil, using Bruker AVANCE-III HD 600 MHz

spectrometers. All spectra were processed with the Azara software package (Boucher). For the

3D data, the two-dimensional maximum entropy method (2D MEM) or Quantitative Maximum

Entropy (QME) (Hamatsu et al., 2013) were applied to obtain resolution enhancement for the

indirect dimensions. All NMR spectra were visualized and analyzed using the CcpNmr Analysis

2.5.0 software (Vranken et al., 2005). All of the 3D triple-resonance experiments used for the

assignments of K-Ras4B and Rgl2^{RBD} were performed on ^{13}C/^{15}N samples in NMR buffer A and B,

respectively. The backbone ^{1}H^N, ^{13}C', and ^{15}N for K-Ras4B and Rgl2^{RBD}, and side-chain ^{13}C$^\alpha$ and

^{13}C$^\beta$ resonance assignments for Rgl2^{RBD} were achieved by analyzing six types of 3D triple-

resonance experiments, HNCO, HN(CA)CO, HNCA, HN(CO)CA, CBCANNH, and CBCA(CO)NNH. 3D

HBHA(CBCACO)NH, H(CCCO)NH, (H)CC(CO)NH, HCCH-COSY, and HCCH-TOCSY experiments on

the ^{13}C/^{15}N -labeled Rgl2^{RBD} were performed for side-chain ^{1}H and ^{13}C resonance assignments. A

15 ms ^{13}C isotropic mixing time was employed for the (H)CC(CO)NH, H(CCCO)NH and HCCH-

TOCSY experiments. For the collection of NOE-derived distance restraints of Rgl2^{RBD}, 3D ^{15}N-

separated and 3D ^{13}C-separated NOESY-HSQC spectra were measured on the ^{13}C/^{15}N -labeled

Rgl2^{RBD}. A 100 ms NOE mixing period was employed for the 3D NOESY experiments. All 2D and

3D NMR data were recorded using the States-TPPI protocol for quadrature detection in

indirectly observed dimensions. Water flip-back ^{1}H pulses and the WATERGATE pulse sequence

were used for solvent suppression in the experiments performed on ^{15}N-labeled, and ^{13}C/^{15}N-

labeled samples, whereas presaturation and gradient-spoil pulses were used for ^{13}C-labeled

samples.

A series of 2D ^{1}H-^{15}N HSQC spectra were measured for titration experiments of ^{15}N-labelled K-

Ras4B^{G12V} with in the presence of non-labelled Rgl2^{RBD}. The experiments were performed in the

NMR buffer at 25 °C and the peptide concentration was increased stepwise (for the ^{15}N-K-

Ras4B^{G12V} / Rgl2^{RBD}, its molar ratio of 1:0.25, 1:0.5, 1:0.75, 1:1, 1:1.5, and 1:2 were used, while

for the ^{15}N-Rgl2^{RBD} / K-Ras4B^{G12V}, its molar ratio of 1:0.25, 1:0.5, 1:1, 1:1.5, 1:2, and 1:3 were

used). The mean chemical shift difference $\Delta\delta_{ave}$ for each amino acid was calculated as $[(\Delta\delta^1$H$^N)^2$

$+ (\Delta\delta^{15}$N$)^2]^{1/2}$ where $\Delta\delta^1$H^N and $\Delta\delta^{15}$N are the chemical shift differences (Hz) between K-

Ras4B^{G12V} or Rgl2^{RBD} on their own and the proteins in the presence of the other side.

NMR structure calculation

Intra-residual and long-range NOEs were automatically assigned by the program CYANA with the

use of automated NOE assignment and torsion angle dynamics (Güntert and Buchner, 2015).

The peak position tolerance was set to 0.03 ppm for the 1H dimension and to 0.3 ppm for the

13C and 15N dimensions. Hydrogen-bond and dihedral angle restraints were not used. CYANA

calculations were started from 100 conformers with random torsion angle values, simulated

annealing with 50,000 torsion angle dynamics steps was applied. The 20 conformers with the

lowest final target-function values of CYANA were selected and optimised with OPALp 2.1

[22,23] using the AMBER force field [24,25]

Mass photometry measurement

Freshly purified KRas4B (G12V) and Rgl2[RBD] were mixed in AKTA buffer, concentrated to about

500 μl using a concentrator (Merck, Amicon Ultra centrifugal filters, #UFC901024), and loaded to

a gel-filtration column (GE Healthcare, HiLoad Superdex 75). The peak fractions containing both

proteins in the 1:1 ratio were collected and the concentration was estimated by OD_{280} and the

predicted extinction coefficient of 24155 $M^{-1}cm^{-1}$. The sample was diluted to 40 nM in gel

filtration buffer and a 20 μL aliquot was subjected to mass photometry (OneMP, Refeyn). The

measurement was done in conventional microscope cover glass (Marienfeld, nº 1.5H) cleaned

by rinsing with deionized water (×5) and isopropanol (×5) followed by drying under a N_2 flow,

using a silicon gasket (Grace Bio-labs) in order to confine the sample. Adsorption of individual

molecules of complex was detected across an imaging area of 10.8 μm by 2.9 μm.

Video recordings from interferometric scattering microscopy for a duration of 60s were

obtained and the single events corresponding to surface adsorption of the complex were

identified using AcquireMP software (Refeyn). Data analysis was performed using DiscoverMP

software (Refeyn) and OriginPro 2021 (OriginLab).

Graphical representation of protein structures

Protein structure images were generated using Pymol (The PyMOL Molecular Graphics System,

Version 1.2r3pre, Schrödinger, LLC). Electrostatic surface charge potential images were

produced using Pymol vacuum electrostatics function. Amino acid residues in the interaction

surfaces of protein complexes in a PDB format were predicted using LIGPLOT (Wallace et al.,

1995).

28
29 **Materials availability statement**

1 Bacteria expression plasmids to generate His_6-K-Ras4B^{G12V}(1-169), untagged K-Ras4B^{G12V} (1-169)

2 and GST-Rgl2^{RBD}(643-740) are freely available upon request.

3
4 **Acknowledgments**
5

We thank the help and expert advice from the University of Leicester colleagues Peter Moody,

Thomas Schalch, Ian Eperon, Mohammed Bhogadia, Luke Bailey, Kyle Daynes, Idir Malki, and

Ziad Ibrahim. Biolayer interferometry experiments were conducted with the guidance of Holly

Birchenough and Thomas Jowitt (Biomolecular Analysis Core Facility, University of Manchester).

We gratefully acknowledge financial support from Wellcome Trust ISSF COVID career support

and inclusion scheme 204801/Z/16/Z and University of Leicester MSc program M32DF74 (to

K.T.), University of Leicester College PhD programme M32RT742 (to M.T.), the Funding

Programs for Core Research for Evolutional Science and Technology (CREST; JPMJCR13M3 to Y.I.,

JPMJCR21E5 to T.I.) the Japan Science and Technology Agency (JST), Grants-in-Aid for Scientific

Research (JP15K06979 to T.I., JP19H05645 to Y.I.) and Scientific Research on Innovative Areas

(JJP15H01645, JP16H00847, JP17H05887 and JP19H05773 to Y.I. P26102538, JP25120003,

JP16H00779 and JP21K06114 to T.I.) from the Japan Society for the Promotion of Science (JSPS),

Shimazu foundation, and the Precise Measurement Technology Promotion Foundation.

19
20

References

Andreadelis, I., Kiriakidi, S., Lamprakis, C., Theodoropoulou, A., Doerr, S., Chatzigoulas, A., Manchester, J., Velez-Vega, C., Duca, J.S., and Cournia, Z. (2022). Membrane Composition and Raf[CRD]-Membrane Attachment Are Driving Forces for K-Ras4B Dimer Stability. J Phys Chem B *126*, 1504-1519.

Apken, L.H., and Oeckinghaus, A. (2021). The RAL signaling network: Cancer and beyond. Int Rev Cell Mol Biol *361*, 21-105.

Bollag, G., and McCormick, F. (1991). Differential regulation of rasGAP and neurofibromatosis gene product activities. Nature *351*, 576-579.

Boucher, W. Azara, v2.7 http://www.bio.cam.ac.uk/azara/.

Bunney, T.D., Harris, R., Gandarillas, N.L., Josephs, M.B., Roe, S.M., Sorli, S.C., Paterson, H.F., Rodrigues-Lima, F., Esposito, D., Ponting, C.P.*, et al.* (2006). Structural and mechanistic insights into ras association domains of phospholipase C epsilon. Mol Cell *21*, 495-507.

Campbell, P.M., Groehler, A.L., Lee, K.M., Ouellette, M.M., Khazak, V., and Der, C.J. (2007). K-Ras promotes growth transformation and invasion of immortalized human pancreatic cells by Raf and phosphatidylinositol 3-kinase signaling. Cancer Res *67*, 2098-2106.

Castellano, E., and Santos, E. (2011). Functional specificity of ras isoforms: so similar but so different. Genes Cancer *2*, 216-231.

Catozzi, S., Halasz, M., and Kiel, C. (2021). Predicted 'wiring landscape' of Ras-effector interactions in 29 human tissues. NPJ Syst Biol Appl *7*, 10.

Chalhoub, N., and Baker, S.J. (2009). PTEN and the PI3-Kinase Pathway in Cancer. Annual Review of Pathology: Mechanisms of Disease *4*, 127-150.

Cox, A.D., and Der, C.J. (2010). Ras history: The saga continues. Small GTPases *1*, 2-27.

Der, C.J., Finkel, T., and Cooper, G.M. (1986). Biological and biochemical properties of human rasH genes mutated at codon 61. Cell *44*, 167-176.

Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501.

Esser, D., Bauer, B., Wolthuis, R.M., Wittinghofer, A., Cool, R.H., and Bayer, P. (1998). Structure determination of the Ras-binding domain of the Ral-specific guanine nucleotide exchange factor Rlf. Biochemistry *37*, 13453-13462.

Evans, P.R., and Murshudov, G.N. (2013). How good are my data and what is the resolution? Acta Crystallogr D Biol Crystallogr *69*, 1204-1214.

Eves, B.J., Gebregiworgis, T., Gasmi-Seabrook, G.M.C., Kuntz, D.A., Prive, G.G., Marshall, C.B., and Ikura, M. (2022). Structures of RGL1 RAS-Association Domain in Complex with KRAS and the Oncogenic G12V Mutant. J Mol Biol *434*, 167527.

Gibbs, J.B., Sigal, I.S., Poe, M., and Scolnick, E.M. (1984). Intrinsic GTPase activity distinguishes normal and oncogenic ras p21 molecules. Proc Natl Acad Sci U S A *81*, 5704-5708.

Güntert, P., and Buchner, L. (2015). Combined automated NOE assignment and structure
calculation with CYANA. J Biomol NMR *62*, 453-471.

Hamatsu, J., O'Donovan, D., Tanaka, T., Shirai, T., Hourai, Y., Mikawa, T., Ikeya, T., Mishima, M.,
Boucher, W., Smith, B.O.*, et al.* (2013). High-resolution heteronuclear multidimensional NMR of
proteins in living insect cells using a baculovirus protein expression system. J Am Chem Soc *135*,
1688-1691.

Huang, L., Hofer, F., Martin, G.S., and Kim, S.H. (1998). Structural basis for the interaction of Ras
with RalGDS. Nat Struct Biol *5*, 422-426.

Hunter, J.C., Manandhar, A., Carrasco, M.A., Gurbani, D., Gondi, S., and Westover, K.D. (2015).
Biochemical and Structural Analysis of Common Cancer-Associated KRAS Mutations. Mol Cancer
Res *13*, 1325-1335.

Ingolfsson, H.I., Neale, C., Carpenter, T.S., Shrestha, R., Lopez, C.A., Tran, T.H., Oppelstrup, T.,
Bhatia, H., Stanton, L.G., Zhang, X.*, et al.* (2022). Machine learning-driven multiscale modeling
reveals lipid-dependent dynamics of RAS signaling proteins. Proc Natl Acad Sci U S A *119*.

Ito, Y., Yamasaki, K., Iwahara, J., Terada, T., Kamiya, A., Shirouzu, M., Muto, Y., Kawai, G.,
Yokoyama, S., Laue, E.D.*, et al.* (1997). Regional polysterism in the GTP-bound form of the
human c-Ha-Ras protein. Biochemistry *36*, 9109-9119.

Jang, H., Muratcioglu, S., Gursoy, A., Keskin, O., and Nussinov, R. (2016). Membrane-associated
Ras dimers are isoform-specific: K-Ras dimers differ from H-Ras dimers. Biochem J *473*, 1719-
1732.

Joosten, R.P., Long, F., Murshudov, G.N., and Perrakis, A. (2014). The PDB_REDO server for
macromolecular structure model optimization. IUCrJ *1*, 213-220.

Kapust, R.B., Tozser, J., Fox, J.D., Anderson, D.E., Cherry, S., Copeland, T.D., and Waugh, D.S.
(2001). Tobacco etch virus protease: mechanism of autolysis and rational design of stable
mutants with wild-type catalytic proficiency. Protein Eng *14*, 993-1000.

Kessler, D., Gmachl, M., Mantoulidis, A., Martin, L.J., Zoephel, A., Mayer, M., Gollner, A., Covini,
D., Fischer, S., Gerstberger, T.*, et al.* (2019). Drugging an undruggable pocket on KRAS. Proc Natl
Acad Sci U S A *116*, 15823-15829.

Kiel, C., and Serrano, L. (2006). The ubiquitin domain superfold: structure-based sequence
alignments and characterization of binding epitopes. J Mol Biol *355*, 821-844.

Lakshman, B., Messing, S., Schmid, E.M., Clogston, J.D., Gillette, W.K., Esposito, D., Kessing, B.,
Fletcher, D.A., Nissley, D.V., McCormick, F.*, et al.* (2019). Quantitative biophysical analysis
defines key components modulating recruitment of the GTPase KRAS to the plasma membrane.
J Biol Chem *294*, 2193-2207.

Lee, K.Y., Enomoto, M., Gebregiworgis, T., Gasmi-Seabrook, G.M.C., Ikura, M., and Marshall, C.B.
(2021). Oncogenic KRAS G12D mutation promotes dimerization through a second,
phosphatidylserine-dependent interface: a model for KRAS oligomerization. Chem Sci *12*, 12827-
12837.

Lee, K.Y., Fang, Z., Enomoto, M., Gasmi-Seabrook, G., Zheng, L., Koide, S., Ikura, M., and
Marshall, C.B. (2020). Two Distinct Structures of Membrane-Associated Homodimers of GTP-

and GDP-Bound KRAS4B Revealed by Paramagnetic Relaxation Enhancement. Angew Chem Int
Ed Engl *59*, 11037-11045.

Lim, K.H., Baines, A.T., Fiordalisi, J.J., Shipitsin, M., Feig, L.A., Cox, A.D., Der, C.J., and Counter,
C.M. (2005). Activation of RalA is critical for Ras-induced tumorigenesis of human cells. Cancer
Cell *7*, 533-545.

Lim, K.H., O'Hayer, K., Adam, S.J., Kendall, S.D., Campbell, P.M., Der, C.J., and Counter, C.M.
(2006). Divergent roles for RalA and RalB in malignant growth of human pancreatic carcinoma
cells. Curr Biol *16*, 2385-2394.

Luo, W., and Sharif, M. (1999). Stable expression of activated Ki-Ras does not constitutively
activate the mitogen-activated protein kinase pathway but attenuates epidermal growth factor
receptor activation in human astrocytoma cells. Int J Oncol *14*, 53-62.

Manne, V., Bekesi, E., and Kung, H.F. (1985). Ha-ras proteins exhibit GTPase activity: point
mutations that activate Ha-ras gene products result in decreased GTPase activity. Proc Natl Acad
Sci U S A *82*, 376-380.

Maurer, G., Tarkowski, B., and Baccarini, M. (2011). Raf kinases in cancer-roles and therapeutic
opportunities. Oncogene *30*, 3477-3488.

Mazhab-Jafari, M.T., Marshall, C.B., Smith, M.J., Gasmi-Seabrook, G.M., Stathopulos, P.B.,
Inagaki, F., Kay, L.E., Neel, B.G., and Ikura, M. (2015). Oncogenic and RASopathy-associated K-
RAS mutations relieve membrane-dependent occlusion of the effector-binding site. Proc Natl
Acad Sci U S A *112*, 6625-6630.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J.
(2007). Phaser crystallographic software. J Appl Crystallogr *40*, 658-674.

McGrath, J.P., Capon, D.J., Goeddel, D.V., and Levinson, A.D. (1984). Comparative biochemical
properties of normal and activated human ras p21 protein. Nature *310*, 644-649.

Menyhard, D.K., Palfy, G., Orgovan, Z., Vida, I., Keseru, G.M., and Perczel, A. (2020). Structural
impact of GTP binding on downstream KRAS signaling. Chem Sci *11*, 9272-9289.

Miller, M.J., Prigent, S., Kupperman, E., Rioux, L., Park, S.H., Feramisco, J.R., White, M.A.,
Rutkowski, J.L., and Meinkoth, J.L. (1997). RalGDS functions in Ras- and cAMP-mediated growth
stimulation. J Biol Chem *272*, 5600-5605.

Muratcioglu, S., Chavan, T.S., Freed, B.C., Jang, H., Khavrutskii, L., Freed, R.N., Dyba, M.A.,
Stefanisko, K., Tarasov, S.G., Gursoy, A.*, et al.* (2015). GTP-Dependent K-Ras Dimerization.
Structure *23*, 1325-1335.

Murshudov, G.N., Skubak, P., Lebedev, A.A., Pannu, N.S., Steiner, R.A., Nicholls, R.A., Winn,
M.D., Long, F., and Vagin, A.A. (2011). REFMAC5 for the refinement of macromolecular crystal
structures. Acta Crystallogr D Biol Crystallogr *67*, 355-367.

Neel, N.F., Martin, T.D., Stratford, J.K., Zand, T.P., Reiner, D.J., and Der, C.J. (2011). The RalGEF-
Ral Effector Signaling Network: The Road Less Traveled for Anti-Ras Drug Discovery. Genes
Cancer *2*, 275-287.

Newlaczyl, A.U., Coulson, J.M., and Prior, I.A. (2017). Quantification of spatiotemporal patterns
of Ras isoform expression during development. Sci Rep *7*, 41297.

Ozdemir, E.S., Koester, A.M., and Nan, X. (2022). Ras Multimers on the Membrane: Many Ways
for a Heart-to-Heart Conversation. Genes (Basel) *13*.

Packer, M.R., Parker, J.A., Chung, J.K., Li, Z., Lee, Y.K., Cookis, T., Guterres, H., Alvarez, S.,
Hossain, M.A., Donnelly, D.P*., et al.* (2021). Raf promotes dimerization of the Ras G-domain with
increased allosteric connections. Proc Natl Acad Sci U S A *118*.

Pacold, M.E., Suire, S., Perisic, O., Lara-Gonzalez, S., Davis, C.T., Walker, E.H., Hawkins, P.T.,
Stephens, L., Eccleston, J.F., and Williams, R.L. (2000). Crystal structure and functional analysis of
Ras binding to its effector phosphoinositide 3-kinase gamma. Cell *103*, 931-943.

Pai, E.F., Krengel, U., Petsko, G.A., Goody, R.S., Kabsch, W., and Wittinghofer, A. (1990). Refined
crystal structure of the triphosphate conformation of H-ras p21 at 1.35 A resolution:
implications for the mechanism of GTP hydrolysis. EMBO J *9*, 2351-2359.

Pantsar, T., Rissanen, S., Dauch, D., Laitinen, T., Vattulainen, I., and Poso, A. (2018). Assessment
of mutation probabilities of KRAS G12 missense mutants and their long-timescale dynamics by
atomistic molecular simulations and Markov state modeling. PLoS Comput Biol *14*, e1006458.

Plowman, S.J., Muncke, C., Parton, R.G., and Hancock, J.F. (2005). H-ras, K-ras, and inner plasma
membrane raft proteins operate in nanoclusters with differential dependence on the actin
cytoskeleton. Proc Natl Acad Sci U S A *102*, 15500-15505.

Prakash, P., Sayyed-Ahmad, A., Cho, K.J., Dolino, D.M., Chen, W., Li, H., Grant, B.J., Hancock, J.F.,
and Gorfe, A.A. (2017). Computational and biochemical characterization of two partially
overlapping interfaces and multiple weak-affinity K-Ras dimers. Sci Rep *7*, 40109.

Prior, I.A., Muncke, C., Parton, R.G., and Hancock, J.F. (2003). Direct visualization of Ras proteins
in spatially distinct cell surface microdomains. J Cell Biol *160*, 165-170.

Sarkar-Banerjee, S., Sayyed-Ahmad, A., Prakash, P., Cho, K.J., Waxham, M.N., Hancock, J.F., and
Gorfe, A.A. (2017). Spatiotemporal Analysis of K-Ras Plasma Membrane Interactions Reveals
Multiple High Order Homo-oligomeric Complexes. J Am Chem Soc *139*, 13466-13475.

Scheffzek, K., Ahmadian, M.R., Kabsch, W., Wiesmuller, L., Lautwein, A., Schmitz, F., and
Wittinghofer, A. (1997). The Ras-RasGAP complex: structural basis for GTPase activation and its
loss in oncogenic Ras mutants. Science *277*, 333-338.

Simanshu, D.K., Nissley, D.V., and McCormick, F. (2017). RAS Proteins and Their Regulators in
Human Disease. Cell *170*, 17-33.

Smith, M.J., and Ikura, M. (2014). Integrated RAS signaling defined by parallel NMR detection of
effectors and regulators. Nat Chem Biol *10*, 223-230.

Smith, S.J., and Rittinger, K. (2002). Preparation of GTPases for structural and biophysical
analysis. Methods Mol Biol *189*, 13-24.

Stieglitz, B., Bee, C., Schwarz, D., Yildiz, O., Moshnikova, A., Khokhlatchev, A., and Herrmann, C.
(2008). Novel type of Ras effector interaction established between tumour suppressor NORE1A
and Ras switch II. EMBO J *27*, 1995-2005.

Torices, R., and Munoz-Pajares, A.J. (2015). PHENIX: An R package to estimate a size-controlled
phenotypic integration index. Appl Plant Sci *3*.

Tran, T.H., Alexander, P., Dharmaiah, S., Agamasu, C., Nissley, D.V., McCormick, F., Esposito, D.,
Simanshu, D.K., Stephen, A.G., and Balius, T.E. (2020). The small molecule BI-2852 induces a
nonfunctional dimer of KRAS. Proc Natl Acad Sci U S A *117*, 3363-3364.

Tran, T.H., Chan, A.H., Young, L.C., Bindu, L., Neale, C., Messing, S., Dharmaiah, S., Taylor, T.,
Denson, J.P., Esposito, D.*, et al.* (2021). KRAS interaction with RAF1 RAS-binding domain and
cysteine-rich domain provides insights into RAS-mediated RAF activation. Nat Commun *12*,
1176.

Tropea, J.E., Cherry, S., and Waugh, D.S. (2009). Expression and purification of soluble His(6)-
tagged TEV protease. Methods Mol Biol *498*, 297-307.

Tuveson, D.A., Shaw, A.T., Willis, N.A., Silver, D.P., Jackson, E.L., Chang, S., Mercer, K.L.,
Grochow, R., Hock, H., Crowley, D.*, et al.* (2004). Endogenous oncogenic K-ras(G12D) stimulates
proliferation and widespread neoplastic and developmental defects. Cancer Cell *5*, 375-387.

Vetter, I.R., and Wittinghofer, A. (2001). The guanine nucleotide-binding switch in three
dimensions. Science *294*, 1299-1304.

Vigil, D., Martin, T.D., Williams, F., Yeh, J.J., Campbell, S.L., and Der, C.J. (2010). Aberrant
overexpression of the Rgl2 Ral small GTPase-specific guanine nucleotide exchange factor
promotes pancreatic cancer growth through Ral-dependent and Ral-independent mechanisms. J
Biol Chem *285*, 34729-34740.

Vranken, W.F., Boucher, W., Stevens, T.J., Fogh, R.H., Pajon, A., Llinas, M., Ulrich, E.L., Markley,
J.L., Ionides, J., and Laue, E.D. (2005). The CCPN data model for NMR spectroscopy:
development of a software pipeline. Proteins *59*, 687-696.

Wallace, A.C., Laskowski, R.A., and Thornton, J.M. (1995). LIGPLOT: a program to generate
schematic diagrams of protein-ligand interactions. Protein Eng *8*, 127-134.

Weise, K., Kapoor, S., Denter, C., Nikolaus, J., Opitz, N., Koch, S., Triola, G., Herrmann, A.,
Waldmann, H., and Winter, R. (2011). Membrane-mediated induction and sorting of K-Ras
microdomain signaling platforms. J Am Chem Soc *133*, 880-887.

Yip-Schneider, M.T., Lin, A., Barnard, D., Sweeney, C.J., and Marshall, M.S. (1999). Lack of
elevated MAP kinase (Erk) activity in pancreatic carcinomas despite oncogenic K-ras expression.
Int J Oncol *15*, 271-279.

Yip-Schneider, M.T., Lin, A., and Marshall, M.S. (2001). Pancreatic tumor cells with mutant K-ras
suppress ERK activity by MEK-dependent induction of MAP kinase phosphatase-2. Biochem
Biophys Res Commun *280*, 992-997.

Young, G., Hundt, N., Cole, D., Fineberg, A., Andrecka, J., Tyler, A., Olerinyova, A., Ansari, A.,
Marklund, E.G., Collier, M.P.*, et al.* (2018). Quantitative mass imaging of single biological
macromolecules. Science *360*, 423-427.

Zago, G., Veith, I., Singh, M.K., Fuhrmann, L., De Beco, S., Remorino, A., Takaoka, S., Palmeri, M.,
Berger, F., Brandon, N.*, et al.* (2018). RalB directly triggers invasion downstream Ras by
mobilizing the Wave complex. Elife *7*.

Zhou, Y., Liang, H., Rodkey, T., Ariotti, N., Parton, R.G., and Hancock, J.F. (2014). Signal
integration by lipid-mediated spatial cross talk between Ras nanoclusters. Mol Cell Biol *34*, 862-
876.

 Figures and Tables

Figure 1. K-Ras4B^{G12V} is structurally more stable and has a stronger affinity towards Rgl2^{RBD}

than K-Ras4B^{WT}.

Analyses of binding kinetics of GST-Rgl2^{RBD} and K-Ras4B^{WT} or K-Ras4B^{G12V}. Biolayer

Interferometry (BLI) was used to directly measure the steady state binding of K-Ras4B (analyte)

to immobilized GST-Rgl2^{RBD} (ligand). (A and C) K-Ras4B^{WT} and K-Ras4B^{G12V} samples used for BLI

and Circular Dichroism (CD) were confirmed to be fully loaded with GTP. The nucleotide binding

status of K-Ras4B^{WT} and K-Ras4B^{G12V} were examined by denaturating the proteins and detecting

the released nucleotides by anion exchange chromatography. Samples of pure GDP or GTP were

used as references. (B) BLI results of binding between K-Ras4B^{WT}/K-Ras4B^{G12V} and Rgl2^{RBD}. GST-

Rgl2^{RBD} was immobilised on the bio-sensors, and varying concentrations of free K-Ras4B^{WT} (left

panels) and K-Ras4B^{G12V} (right panels) were provided and the interactions were measured at

23^{o}C. The kinetics diagrams (lower panels, association at t=0(s), dissociation at t=300(s))

revealed a fast on/off binding. Therefore, steady state curves (upper panels) were used to

estimate the K_D values; about 6.1 µM for K-Ras4B^{WT}:Rgl2^{RBD} and 1.4 µM for K-Ras4B^{G12V}:Rgl2^{RBD},

respectively. (D) Circular dichroism (CD) spectra of K-Ras4B^{WT} and K-Ras4B^{G12V} (20 µM) at 20^{o}C.

(E) CD signal intensity at 220 nm as a function of temperature from 20^{o}C to 90^{o}C.

11
12

Figure 1-figure supplement 1. Rgl2 Ras binding domain interacts with active K-Ras4B^{G12V}.

(A) A schematic diagram of domains organization of human Rgl2 as defined in the UniProt

database (Uniprot number: O15211). A domain spanning amino acid residues 648-735 is

annotated as "Ras-associating" by PROSITE annotation rule PRU00166. In this work, we

generically call this domain Rgl2 Ras Binding Domain (Rgl2^{RBD}). (B) Rgl2^{RBD} interacts with active K-

Ras4B^{G12V}. Bacteria recombinant Rgl2^{RBD} fragment spanning the amino acid residues 643-740 of

Rgl2 was fused with GST and the fusion protein was fixed on glutathione beads. Recombinant K-

Ras4B^{G12V} 1-169, loaded with either GDP or non-hydrolisable GTP analogue, GMPPNP, was

applied on the beads to examine the protein-protein interaction. GST-Rgl2^{RBD} interacted only

with the GMPPNP-loaded K-Ras4B^{G12V}. (C) K-Ras4B^{G12V}:Rgl2^{RBD} complex was purified using size

exclusion chromatography. The fractions were analyzed by 15% or 12% SDS-PAGE gel (lower

panel) according to the elution profile (upper panel). Fractions indicated by the green double-

arrow line were used for crystallization trials. (D) K-Ras4B^{WT} and Rgl2^{RBD} preparations were

mixed and applied to size exclusion chromatography. The elution profile (upper panel) shows

little complex formation. Peak fractions were analyzed by 15% SDS-PAGE gel (lower panel).

1
2
3   **Figure 2. Crystal structure of the K-Ras4B^{G12V}:Rgl2^{RBD} 2:2 heterotetramer**

(A) Cartoon representation of the structure of the heterotetramer complex of K-Ras4B^{G12V} and

Rgl2^{RBD} with top view and side view; the two K-Ras4B^{G12V} molecules are shown in dark and pale

cyan and the two molecules of Rgl2^{RBD} in pink and violet. Switch-I and Switch-II regions of K-

Ras4B^{G12V} are shown in green and yellow, respectively, and the α-helix and β-sheets are

numbered for each chain. The Mg^{2+} is shown as a grey sphere. The structure shows that each K-

Ras4B^{G12V} molecule interacts with two Rgl2^{RBD} molecules (referred to as 1 and 2) at Switch-I and

Switch-II individually. (B) The interacting interface of K-Ras4B^{G12V} and Rgl2^{RBD}, highlighting the

residues involved in hydrogen and hydrophobic interactions between β1, β2 and α1 of Rgl2^{RBD1}

(violet sticks) and K-Ras4B^{G12V} Switch-I (green sticks, enlarged in the green box) or Switch-II

(yellow sticks, enlarged in the yellow box). An orange dotted line shows a salt bridge formed

between E37 of K-Ras4B^{G12V} and R653 of Rgl2^{RBD}. K686 of Rgl2^{RBD}, which plays an essential role in

the complex formation as shown in Figure 3-figure supplement 2B, is annotated in purple

letters. (C) The interacting interface of K-Ras4B^{G12V}: K-Ras4B^{G12V}, highlighting the residues

involved in hydrogen and hydrophobic interactions between two K-Ras4B^{G12V} molecules. The

interface comprises Switch-I, Switch-II and α3 of the two K-Ras4B^{G12V} molecules. The oncogenic

mutation, V12, is annotated in red letters. (D) The interacting interface of Rgl2^{RBD}: Rgl2^{RBD},

highlighting the residues involved in hydrogen and hydrophobic interactions between two

Rgl2^{RBD} molecules, shaded in pink and violet. The interaction occurs at the N-terminal between

the two anti-parallel β1.

```
Human Rgl2  SDCRIIRVQMELGEDGSVYKSILVTSQDKAPSVISRVLKKNNRDSAVASEYELVQLLPGERELTIPASANVFYAMDGASH-DFLLRQRRRS 736
Rat RALGDS  -DCCIIRVSLDV-DNGNMYKSILVTSQDKAPTVIRKAMDKFNLDEDEPEDVEILQIISEDHKLKIPENANVFYAMNSLANYDFILKKR--- 864
            **  ****.::: ::*.:**************:** :.:.*:* *.  ..:***:*::. :. :*.** .*******:.*:: **;*::*
```

Figure 2-figure supplement 1. K-Ras4B^{G12V}:Rgl2^{RBD} complex is highly similar to H-Ras^{E31K}:RALGDS^{RBD} complex

(A) The structures of the K-Ras4B^{G12V}:Rgl2^{RBD} complex and the H-Ras^{E31K}:RALGDS^{RBD} complex (PDB: 1LFD) were superimposed by aligning the RBD chains. (B) and (C) Superimposition of each

corresponding chain of the K-Ras4B^{G12V}: Rgl2^{RBD} complex and H-Ras^{E31K}:RALGDS^{RBD} complex

individually shows that each chain is similar to its respective counterpart structure. (B) K-

Ras4B^{G12V} and H-Ras^{E31K} chains are in cyan and grey. (C) Rgl2^{RBD} and RALGDS^{RBD} are in purple and

light pink. Amino acid sequence similarities of human Rgl2^{RBD} and rat RALGDS^{RBD} are shown

below the RBD cartoon representations, and the conserved amino acids in Rgl2^{RBD} and

RALGDS^{RBD} are shaded in light green and dark green, respectively. The structural similarity

between Rgl2^{RBD} and RALGDS^{RBD} was seen beyond the regions composed of the conserved

amino acids. The RALGDS^{RBD} sequence is derived from PDB 1LFD, which is rat RALGDS^{RBD}, and the

numbering was done according to Uniprot Q03386. Root-mean-square deviation of atomic

positions (RMSD) values were generated using Pymol.

A

B

2 **Figure 2-figure supplement 2. Schematic representation of the intermolecular contacts**

3 **observed in the X-ray structure**

(A) The residues in K-Ras4B^{G12V} Switch-I region are shown in green, in the Switch-II region are

shown in yellow background and those for Rgl2^{RBD} molecules 1 and 2 are shown in pink. V12

oncogenic mutation is in red. Hydrogen bonds, salt bridges and hydrophobic interactions were

predicted by LIGPLOT and represented by green dashed lines, solid orange lines and blue dotted

lines, respectively. (B) The electrostatic surface charge shows that the negatively charged

Switch-I and Switch-II regions of KRas4B^{GV} interact with the positively charged surface of Rgl2^{RBD}-

1 and Rgl2^{RBD}-2. Surface charge potential was computed using the Pymol vacuum electrostatics

function, and the negatively charged and positively charged areas are shown in red and blue,

respectively.

A

K-Ras4B^{G12V} : Rgl2^{RBD}

H-Ras^{E31K} : RALGDS^{RBD}
(1LFD)

RBD1
RBD2

B

K-Ras4B^{WT}:Raf1^{RBD} (6VJJ)	H-Ras^{G12V}:PI3Kγ^{RBD} (1HE8)	H-Ras^{G12V}:PLCε^{RBD} (2C5L)	H-Ras$^{D30E/E31K}$:RASSF5^{RBD} (3DDC)	K-Ras4B^{G12V}:Rgl1^{RBD} (7SCX)	K-Ras4B^{WT}:Rgl1^{RBD} (7SCW)
K-Ras4B (Rgl2 complex) vs K-Ras4B (Raf1 complex)	K-Ras4B (Rgl2 complex) vs H-Ras (PI3Kγ complex)	K-Ras4B (Rgl2 complex) vs H-Ras (PLCε complex)	K-Ras4B (Rgl2 complex) vs H-Ras (RASSF5 complex)	K-Ras4B (Rgl2 complex) vs K-Ras4B (Rgl1 complex)	K-Ras4B (Rgl2 complex) vs K-Ras4B (Rgl1 complex)
RMSD: 0.494Å (139 to 139 atoms)	RMSD: 0.457Å (151 to 151 atoms)	RMSD: 0.383Å (137 to 137 atoms)	RMSD: 0.400Å (140 to 140 atoms)	RMSD: 0.305Å (132 to 132 atoms)	RMSD: 0.376Å (138 to 138 atoms)

Figure 2-figure supplement 3. Ras:RalGEF^{RBD} complex formation is distinct from other

Ras:effector complexes.

K-Ras4B^{G12V}:Rgl2^{RBD} complex was compared with representative Ras:effector complexes. The

spectrum colour feature illustrates the orientation and ubiquitin super-fold status for one of the

RBD(s). When two RBDs interact with one Ras molecule, the second RBD is coloured pink

(Rgl2^{RBD}) or blue (RALGDS^{RBD}). The K-Ras4B^{G12V} in the Rgl2^{RBD} complex is in cyan, and other Ras

molecules from previously published structures are in grey. The interacting residues involved in

Ras Switch-II and RBDs are depicted with magenta sticks and are labelled. (A) The two

Ras:RalGEF complexes (K-Ras4B^{G12V}:Rgl2^{RBD} complex and H-Ras^{E31K}:RALGDS^{RBD} complex PDB:

1LFD) show similar structural features, including the orientation of the RBDs, the Switch-I

contact and the mode of usage of Switch-II. Two RBDs (RBD-1 in the spectrum colour and RBD-2

in either pink or in blue) interact separately with a single Ras molecule at Switch-I and Switch-II

regions. (B) Examples of Ras:effector complex crystal structures. K-Ras4B^{WT}:Raf1 complex (PDB:

6VJJ) shows that although Raf1^{RBD} has similar orientations of the ubiquitin fold structure to

Rgl2^{RBD}-1, it does not utilise Ras Switch-II in complex formation. Superimposition of the K-

Ras4B^{G12V} of the Rgl2^{RBD} complex onto the K-Ras4B^{WT} of the Raf1^{RBD} complex shows some

differences in Switch-II, β1 and α3 (arrows). PI3Kγ^{RBD} (PDB:1HE8) and PLCε^{RBD} (PDB:2C5L)

complexed with Ras utilise Switch-II region differently to Ras:RalGEF complexes despite a similar

orientation of the ubiquitin fold structure of RBDs. A single RBD interacts at both Switch-I and

Switch-II regions of one Ras molecule. Superimposed H-Ras molecules in these complexes to K-

Ras4B^{G12V} in the Rgl2^{RBD} complex gave lower RMSD values compared to K-Ras4B^{WT} in the Raf1^{RBD}

complex. H-Ras$^{D30E/E31K}$:RASSF5^{RBD} complex (3DDC) shows yet another unique interaction where

the additional α-helix at the N-terminal of the RBD interacts with the Switch-II region of the Ras

molecule, instead of the RBD region itself. Rgl1 is one of the RalGEFs, as Rgl2 and RALGDS are.

However, unlike Rgl2 or RALGDS, K-Ras4B^{G12V}:Rgl1^{RBD} complex (7SCX) and K-Ras4B^{WT}:Rgl1^{RBD}

complex (7SCW) show one Rgl1^{RBD} to interact with both Switch-I and Switch-II of one Ras

molecule.

2 **Figure 2-figure supplement 4. Various Ras:Ras interfaces proposed by experimental and**

3 **computational prediction studies.**

4 The top panel on the left shows the K-Ras4B[G12V]:K-Ras4B[G12V] interface of the K-Ras4B[G12V]:Rgl2[RBD]

5 heterotetramer complex (this study). The K-Ras4B[G12V]:K-Ras4B[G12V] contacts (shaded in purple)

6 are located in loop regions of Switch-I, Switch-II and α3. The K-Ras4B[G12V]:K-Ras4B[G12V] interface is

distinct from previously proposed dimerization interfaces, which can be classified into four

categories based on the relevant structural elements; α5-α4, α4-α3, α-β and β-β, as listed in (a)-

(n). Images of a single Ras molecule at the Ras:Ras interface are presented with the interacting

residues highlighted in purple, and the involved α-helices and β-sheets are annotated. PDB IDs

of the image templates are indicated next to each image. **(a)-(f)** Examples of Ras:Ras interfaces

that involve α4-α5. **(g)-(i)** Examples of Ras:Ras interfaces that involve α3-α4. **(j)** A K-Ras4B^{G12D}:K-

Ras4B^{G12D} interface that lies in α4-β2 and α1-β6. The two K-Ras4B^{G12D} in the dimer are in two

different orientations for this interaction to occur. **(k)-(m)** Examples of Ras:Ras interfaces that

involve β-β. The interfaces overlap with the effector binding region and are thought to impair

effector interaction. **(n)** The dimer formation mediated by the β-β contact was aided by a small

molecule inhibitor BI2852 (shown in purple). The Mg^{2+} is depicted as a dark-grey sphere and the

bound nucleotide is shown as a ball-and-stick model.

Figure 2-figure supplement 5. Spatial Ras:Ras arrangements of K-Ras4B^{G12V}:K-Ras4B^{G12V} in the

K-Ras4B^{G12V}:Rgl2^{RBD} complex and H-Ras^{WT} crystal structure 5P21

Neighboring Ras molecules are shown to reveal comparable spatial arrangements. The side view

(top panel), top view (middle panel) and the blow-up of the area Y32 (bottom panel) are shown.

Switch I and Switch II regions are indicated in green and yellow. (A) K-Ras4B^{G12V}:K-Ras4B^{G12V} in

the K-Ras4B^{G12V}:Rgl2^{RBD} complex. The Rgl2^{RBD} is shown as a ribbon in light magenta. (B) H-

Ras^{WT}:H-Ras^{WT} in the H-Ras^{WT} crystal (5p21). Spatial molecular arrangements around Y32 are

similar in both cases, but the oncogenic V12 provides a larger hydrophobic pocket.

2 **Figure 2-figure supplement 6. V12 contributes to Ras:Ras interaction**

Ras residues at position 12 and Y32 are shown to demonstrate the V12 contribution to stabilise

the K-Ras4B^{G12V}:K-Ras4B^{G12V} interface of the K-Ras4B^{G12V}:Rgl2^{RBD} heterotetramer. The Ras:Ras

interface is indicated by a grey dotted line in (A)-(C). (A) A cartoon diagram of the K-Ras4B^{G12V}:K-

Ras4B^{G12V} interface of the K-Ras4B^{G12V}:Rgl2^{RBD} crystal structure where V12 generates a

hydrophobic pocket together with Y32. (B) A simulation of V12 -> G12 replacement in the

structure shown in (A). The hydrophobic pocket is reduced by the G12 replacement. (C) A

cartoon diagram of the H-Ras^{E31K}:H-Ras^{E31K} interface of H-Ras^{E31K}:RALGDS^{RBD} crystal structure

(PDB ID 1LFD). Y32 orientation is different from (A). Nonetheless, with G12, the hydrophobic

area around Y32 looks less robust compared to the G12 case shown in (A). (D and E) Extended

views of (A) and (C) where the neighbouring E31 (A) and K31 (D) are shown. (D) E31 of K-

Ras4B^{G12V} interacts with K-Ras4B^{G12V} K88 and Rgl2^{RBD} K685. (E) The hydrophobic pocket created

by Y32 is not reinforced by G12, but K31 of H-Ras^{E31K} interacts with rat RALGDS^{RBD} D815, N818

and D820 (the numbering was done according to Uniprot Q03386), facilitating the complex

formation.

15
16

2　**Figure 3. NMR analysis of K-Ras4B^{G12V}:Rgl2^{RBD} complex formation in solution**

3　(A) Rgl2^{RBD} solution structure determined by NMR. The 2D ^{1}H-^{15}N-HSQC spectrum of the assigned

4　residues of the Rgl2^{RBD} is shown (left). The Rgl2^{RBD} retains the ββαββαβ ubiquitin-fold structure

5　(right). 20 Rgl2 NMR structures are superimposed. (B) Comparison between the Rgl2^{RBD}

structures identified in solution NMR (shown in magenta) and in crystal complex with K-

Ras4B^{G12V} (shown in cyan). (C) NMR signal intensity changes of ^{15}N-labelled Rgl2^{RBD} upon addition

of non-labelled K-Ras4B^{G12V} as a function of Rgl2^{RBD} amino acid sequence. The original HSQC

spectra of the titration experiment are presented in Figure 3-figure supplement 1B. The signal

intensities of Rgl2^{RBD} residues in the presence of three times molar excess of K-Ras4B^{G12V} were

divided by the signal intensities in the absence of K-Ras4B^{G12V} and plotted as a bar-chart graph.

Proline and unassigned Rgl2^{RBD} residues are shaded in grey. Red-dotted lines are drawn at the

fold-change values of 0.5 and 1.5 to highlight the residues that show substantial increase or

decrease of the signals upon addition of K-Ras4B^{G12V}. Rgl2^{RBD} residues involved in the

Rgl2^{RBD}:Rgl2^{RBD} interface of the K-Ras4B^{G12V}:Rgl2^{RBD} crystal structure are highlighted in pink on

the amino acid sequence, and residues involved in the K-Ras4B^{G12V}:Rgl2^{RBD} interface are

highlighted in blue. Rgl2^{RBD} residue position numbers according to the UniProt are indicated at

the bottom of the diagram. K686, which plays a crucial role in the K-Ras4B^{G12V} complex

formation, is annotated in orange.

A

Human Rgl2 (This study)   MGPGASDCRIIRVQMELGEDGSVYKSILVTSQDKAPSVISRVLKKNNRDSAVASEYELVQLLPGERELTIPASANVFYAMDGASHDFLLRQRRRSSTAT
Mouse Rlf1 (1RLF)   ---GSSDCRIIRVQMELGEDGSVYKSILVTSQDKAPSVISRVLKKNNRDSAVASEFELVQLLPGDRELTIPHSANVFYAMDGASHDFLLRQRR------
                              *:**:********:****** ********************

B

¹⁵N-Rgl2^{RBD}:K-Ras4B^{G12V} Titration

C

■ Residues in the Rgl2^{RBD}:Rgl2^{RBD} binding interface of the K-Ras4B^{G12V}:Rgl2^{RBD} crystal strucutre
■ Residues in the K-Ras4B^{G12V}:Rgl2^{RBD} binding interface of the K-Ras4B^{G12V}:Rgl2^{RBD} crystal strucutre

The ratio of the ¹⁵N Rgl2^{RBD} and K-Ras4B^{G12V} in the samples

Figure 3-figure supplement 1. K-Ras4B^{G12V}:Rgl2^{RBD} complex formation in solution

(A) Comparison between the human Rgl2^{RBD} and mouse Rlf^{RBD} (PDB 1RLF) solution structures.

Rgl2^{RBD} is presented in magenta, and 20 NMR structures are superimposed. Rlf^{RBD} is in cyan, and

10 NMR structures are superimposed. The primary structures of the two constructs are listed

below. (B) and (C) ^{1}H-^{15}N HSQC titration analysis of ^{15}N-labelled Rgl2^{RBD} upon addition of non-

labelled K-Ras4B^{G12V} supports the K-Ras4B^{G12V}:Rgl2^{RBD} tetramer formation. (B) Overlays of 2D ^{1}H-

^{15}N HSQC NMR spectra from multipoint titrations of ^{15}N-labelled Rgl2^{RBD} with non-labelled K-

Ras4B^{G12V}.. Rgl2^{RBD}:K-Ras4B^{G12V} molar ratios of the titration samples are colour-coded as follows;

1:0 - black, 1:0.25 - blue, 1:0.25 – cyan, 1:0.5 – green, 1:1 – yellow, 1:2 – orange and 1:3 – red.

(C) The chemical shift perturbation of backbone ^{1}H^{N} and ^{15}N nuclei of Rgl2^{RBD} with non-labelled

K-Ras4B^{G12V}, presented in (B), is summarised as a column diagram. The mean shift difference

$\Delta\delta_{ave}$ was calculated as $[(\Delta\delta^{1}H^{N})^{2} + (\Delta\delta^{15}N/10)^{2}]^{1/2}$ where $\Delta\delta^{1}H^{N}$ and $\Delta\delta^{15}N$ are the chemical shift

differences between Rgl2^{RBD} on its own and in the presence of non-labelled K-Ras4B^{G12V}. The bar

graphs are colour-coded according to the Rgl2^{RBD}- K-Ras4B^{G12V} concentration ratio and are

overlaid. Proline and unassigned residues are shaded in grey. Rgl2^{RBD} residues in the

Rgl2^{RBD}:Rgl2^{RBD} interface of the K-Ras4B^{G12V}:Rgl2^{RBD} crystal structure is highlighted in pink and

residues in the K-Ras4B^{G12V}:Rgl2^{RBD} interface of the crystal structure is highlighted in blue.

Rgl2^{RBD} residue positions according to the UniProt are indicated at the bottom of the diagram.

A

B

C

Figure 3-figure supplement 2. Rgl2 K686 plays a key role in the complex formation

3 **(A)** Multiple sequence alignment of Ras binding domain (RBD) of human BRAF (residues 155-

4 227) and Ras Associating (RA) domains of human RALGDS (residues 798-885), Rgl1 (residues

648-735) and Rgl2 (residues 648-735), as defined by UniProt. The alignment reveals that

sequences are relatively divergent between BRaf^{RBD} and RalGDS family, although all RBD/RA

domains share the common ubiquitin fold $\beta\beta\alpha\beta\beta\alpha\beta$ structure. The RalGDS family shares a high

degree of sequence homology, including the red-arrowhead-indicated key lysine 686 residue

(highlighted in red, K686 in Rgl2^{RBD}), which is involved in the Ras:RA complex interface. **(B)** The

conserved K686 residue in Rgl2^{RBD} interacts via a hydrogen bond and hydrophobic interaction

with D33 in the Switch-I region of KRas4B^{G12V}. **(D)** Rgl2^{RBD}-K686A mutant hinders the interaction

between Rgl2^{RBD} and KRas4B. The Biolayer interferometry analysis of Rgl2^{RBD}-K686A mutant and

KRas4B^{G12V} did not give a conclusive Kd, likely due limitation of the technique in measuring low

binding affinity. The reduced affinity of the K686A mutant can be illustrated by a comparison

between Rgl2^{RBD}-K686A mutant and Rgl2^{RBD}-WT amplitudes during the association step with

KRas4B^{G12V}, which shows a decrease in the case of K686A mutant.

Figure 3-figure supplement 3. Backbone resonance assignment of GDP-bound and GMPPNP-bound K-Ras4B[G12V]

The 2D ^{1}H-^{15}N-HSQC spectra of GDP-Bound and GMPPNP-bound K-Ras4B[G12V]. Cross peaks are labelled with their corresponding backbone assignments. Residues which could not be assigned are shaded in grey in the primary sequence shown at the bottom.

2   **Figure 3-figure supplement 4. K-Ras4B^{G12V}:Rgl2^{RBD} complex formation in solution**

^{1}H-^{15}N-HSQC titration analysis of ^{15}N-labelled K-Ras4B^{G12V} upon addition of non-labelled Rgl2^{RBD}

supports the K-Ras4B^{G12V}:Rgl2^{RBD} tetramer formation. (A) and (B) The 2D ^{1}H-^{15}N-HSQC NMR

spectra of K-Ras4B^{G12V}. ^{15}N-labelled K-Ras4B^{G12V} was titrated with non-labelled Rgl2^{RBD}. (A) The

2D ^{1}H-^{15}N-HSQC spectra of K-Ras4B^{G12V}: Rgl2^{RBD} complex mixed with the molar ratio of 1:0 (black,

left panel) and 1:2 (red, right panel) are shown. Many signals from ^{15}K-Ras4B^{G12V} residues

disappeared upon the addition of Rgl2^{RBD}. (B) Superimposed 2D ^{1}H-^{15}N-HSQC NMR spectra of

^{15}N-labelled K-Ras4B^{G12V}:Rgl2^{RBD} titration experiments. The titration samples are colour-coded as

follows; 1:0-black, 1:0.25 - blue, 1:0. 5 – cyan, 1:0.75 –green, 1:1 – yellow, 1:1.5 – orange and 1:2

– red. (C) Fold changes of the signal intensities of ^{15}N-labelled K-Ras4B^{G12V} upon addition of non-

labelled Rgl2^{RBD}. The signal intensities of K-Ras4B^{G12V} residues in the presence of two times

molar excess of Rgl2^{RBD} were divided by the signal intensities in the absence of Rgl2^{RBD}, and the

obtained values were plotted as a column graph. Undetectable residues are shaded in grey. The

residue T2 was also shaded grey, as the chemical shift after the addition of Rgl2^{RBD} overlapped

with other signals. A red-dotted line is drawn at the fold-change values of 0.5 to indicate that

the most residues show substantial decrease of the signals upon addition of Rgl2^{RBD}. K-Ras4B^{G12V}

residues in the K-Ras4B^{G12V}: K-Ras4B^{G12V} interface of the K-Ras4B^{G12V}:Rgl2^{RBD} crystal structure is

highlighted in orange, and residues in the K-Ras4B^{G12V}:Rgl2^{RBD} interface of the crystal structure is

highlighted in blue. K-Ras4B^{G12V} residue positions according to the UniProt are indicated at the

bottom of the diagram.

Figure 4. K-Ras4B^{G12V}:Rgl2^{RBD} complex can form a heterotetramer in solution

Mass photometry analysis of KRas4B^{G12V}:Rgl2^{RBD} complex in solution. (A) K-Ras4B^{G12V}:Rgl2^{RBD} complex was purified using size exclusion chromatography. The fractions were analysed by 15% SDS-PAGE gel (upper panel) according to the elution profile (lower panel). Fraction 5 was chosen and diluted to a concentration of 40nM for measurement using mass photometry (OneMP, Refeyn). (B) Cartoon representing the different complexes configurations possibilities indicating that only heterodimers (~31kDa) and heterotetramers (~62kDa) can be detected the system. Histogram of the frequency counts against of the purified KRas4B^{GV}:Rgl2^{RBD} complex with fitting for Gaussian distribution (red). The duration of the video analysed was 60s and it shows that the population identified has an average mass of ~68kDa, which is in good agreement with the expected MW of the heterotetramer (61.7kDa).

Figure 5. Possible K-Ras4B^G12V:Rgl2^RBD complex orientations relative to the plasma membrane

Possible conformational arrangements of the K-Ras4B^G12V:Rgl2^RBD complex at the plasma membrane (PM). The Rgl2^RBD molecules may be facing the cytoplasm (PM-away conformation) or the PM (PM-proximity conformation). Cartoons are prepared based on the previously proposed α4-α5 mediated K-Ras4B^G12D:K-Ras4B^G12D homodimer at the PM (PDB ID 7RSC) (Lee et al., 2021). The Ras:Raf1 kinase heterotetramer cartoon was generated by merging the K-Ras4B^G12D:K-Ras4B^G12D homodimer (PDB ID 7RSC) and K-Ras4B^WT:Raf1 crystal structure (PDB ID 6Vjj) (Tran et al., 2021; Packer et al., 2020). K-Ras4B is depicted in cyan, and domains are colour coded as follows; Switch-I in yellow, Switch-II in green, α4 and α5 helices in blue, C-terminal hypervariable region (HVR) in purple and the residue at position 12 in red. For the K-Ras4B^G12V:Rgl2^RBD structure, the HVR is in grey, as the K-Ras4B^G12V in the original crystal structure lacked the HVR.

1 **Table 1. Data collection and refinement statistics for human K-Ras4B^{G12V}:Rgl2^{RBD} complex.**
2 Statistics for the highest-resolution shell are shown in parentheses.

Wavelength	0.9795
Resolution range	28.68 - 3.071 (3.18 - 3.071)
Space group	P 32 2 1
Unit cell	77.843 77.843 163.728 90 90 120
Total reflections	22133 (1819)
Unique reflections	11107 (949)
Multiplicity	2.0 (1.9)
Completeness (%)	98.22 (86.51)
Mean I/sigma(I)	10.15 (2.11)
Wilson B-factor	95.59
R-merge	0.02682 (0.1774)
R-meas	0.03794 (0.2509)
R-pim	0.02682 (0.1774)
CC1/2	0.999 (0.973)
CC*	1 (0.993)
Reflections used in refinement	11065 (936)
Reflections used for R-free	582 (41)
R-work	0.2367 (0.3229)
R-free	0.3204 (0.3572)
CC(work)	0.943 (0.893)
CC(free)	0.849 (0.490)
Number of non-hydrogen atoms	4116
macromolecules	4050
ligands	66
solvent	0
Protein residues	514
RMS(bonds)	0.007
RMS(angles)	1.40
Ramachandran favored (%)	95.22
Ramachandran allowed (%)	3.98
Ramachandran outliers (%)	0.80
Rotamer outliers (%)	0.89
Clashscore	14.47
Average B-factor	113.50
macromolecules	113.74
ligands	98.55

3
4

1 **Table 2. NMR structure statistics for Rgl2[a]**

Quantity	
Assigned $^1H/^{13}C/^{15}N$ chemical shifts	305/256/56
NOE distance restraints[b]	866/281/777
Max. distance restraint violation (Å)	0.13 ± 0.01
Deviations from idealised geometry:	
Bond lengths (Å)	0.0138 ± 0.0001
Bond angles (º)	1.81 ± 0.03
AMBER energy (kcal/mol)	−3624.25 ± 78.09
AMBER vdW energy (kcal/mol)	−279.71 ± 12.49
Ramachandran plot statistics[c] (%)	89.5/10.5/0.1/0
Backbone RMSD (Å)[d]	0.42 ± 0.05
All heavy atom RMSD (Å)[d]	0.84 ± 0.06
Backbone RMSD to the reference (Å)[e]	0.91
All heavy atom RMSD to the reference (Å)[e]	1.68

[a]Where applicable, the average value and the standard deviation over the 20 energy-refined
conformers obtained by the program OPALp 2.1 in the presence of the experimental restraints.
CYANA calculations were started from 100 conformers with random torsion angle values,
simulated annealing with 10,000 torsion angle dynamics steps was applied.
[b]Short/medium/long-range distance restraints derived from NOESY spectra.
[c]Percentage of residues in the most favoured/additionally allowed/generously
allowed/disallowed regions of the Ramachandran plot according the program PROCHECK.

[d]RMSD to the mean structure for residues 649-657, and 665–732.

[e]RMSD between the closest structure to the reference among the ensemble conformations and
the reference structure (the crystal structure of the K-Ras4B^{G12V}:Rgl2 complex).

14
15