Bidirectional changes in postmitotic H3K27me3 distributions underlie cerebellar

granule neuron maturation dynamics

Vijyendra Ramesh[1,2], Fang Liu[1,3], Melyssa S. Minto[3], Urann Chan[3] and Anne E. West[2,3]

[1]These authors contributed equally

[2]Molecular Cancer Biology Program, Duke University School of Medicine, Durham, NC-27705

[3]Department of Neurobiology, Duke University School of Medicine, Durham, NC-27705

19
20

Abstract

The functional maturation of neurons is a prolonged process that extends beyond mitotic exit and requires the dynamic orchestration of gene transcription programs. We find that the postnatal maturation of cerebellar granule neurons (CGNs) is associated with dynamic changes in the genomic distribution of histone H3 lysine 27 trimethylation (H3K27me3), a chromatin modification best known for its role in cell fate specification. The developmental loss of H3K27me3 in postmitotic neurons is facilitated by the lysine demethylase KDM6B, and target genes induced in maturing CGNs show impaired expression with conditional KDM6B knockout. Interestingly, inhibition of the H3K27 methyltransferase EZH2 in newborn CGNs not only blocks the repression of progenitor genes but also impairs the induction of mature CGN genes, showing the importance of bidirectional regulation. These data demonstrate that H3K27me3 functions in developing postmitotic neurons to regulate the temporal coordination of gene expression programs that underlie functional neuronal maturation.

Introduction

Cell identity is determined during differentiation by the coordinated actions of
chromatin regulatory factors, which serve to establish the accessibility of gene regulatory
elements that control transcription of cell-type specific genes. Histone proteins play a key role
in determining the architecture of chromatin, and specific post-translational modifications of
the histones are tightly associated with the transcriptional state of a given locus or gene [1].
Among these modifications, histone methylation occurs on lysine and arginine residues of
specific histones, and has roles in both gene activation and repression [2], and acts to alter gene
transcription by recruiting the site-specific binding of effector proteins that carry out
specialized functions [3]. The trimethylation of lysine 27 on histone H3 (H3K27me3) is a
repressive histone modification that is best known for its role during cell-fate determination [4].
H3K27 methylation is deposited by the Polycomb repressive complex 2 (PRC2) lysine
methyltransferases EZH1 and EZH2 [5, 6]. H3K27me3 can be enzymatically removed by the
H3K27-specific lysine demethylases KDM6A (UTX) and KDM6B (JMJD3) [7]. In mice, null
mutations in *Ezh2* result in early embryonic lethality [5], *Kdm6a* knockouts show severely
impaired mid-gestational development [8], and *Kdm6b* knockouts die at birth [9] suggesting the
physiological importance of H3K27me3 regulation for mammalian embryonic development.
67
At the cellular level, the functions of H3K27me3-dependent gene regulation in cell fate
determination are well established [10]. In stem cells, the colocalization of H3K27me3 with the
active histone marks H3K4me3 and H3K27ac at regulatory elements of cell-type specific genes
is thought to poise these genes for subsequent activation or silencing upon terminal
differentiation depending on the chosen fate of the cell. Activation of poised genes that
function in the chosen fate is associated with the permanent loss of H3K27me3, whereas
persistent silencing of genes that define alternative fates is mediated by recruitment of the
PRC1 complex. However, the enzymes that regulate H3K27me3 remain expressed in fate-
committed, postmitotic neurons both during postnatal stages of development and in the adult
brain [11, 12], where their functions are less well understood. Genetic evidence that ongoing
regulation of H3K27me3 after fate commitment might be important emerged when conditional

deletion of both *Ezh1* and *Ezh2* in neurons of the adult mouse striatum was shown to cause a

slow global loss of H3K27me3 as well as progressive neurodegeneration and death [13].

H3K27me3 may be required in terminally differentiated neurons to persistently repress genes

that have the potential to drive trans-differentiation to alternate cell fates. For example, in a

longitudinal study of postmitotic dopamine neurons undergoing directed differentiation in

culture, poised RNA Pol-II was found to colocalize with H3K27me3 at promoters of genes

encoding non-neuronal transcription factors (TFs) [14]. A separate study showed knockout of

the PRC2 component EED resulted in transcriptional de-repression of a set of TFs that have the

capacity to drive cellular reprogramming [13]. A non-exclusive possibility is that H3K27me3 is

required to maintain the expression of genes that define the terminal neuronal fate. Consistent

with this model, deleting EED in developing postmitotic dopamine or serotonin neurons *in vivo*

leads not only to loss of H3K27me3 but also to reduced expression of tyrosine hydroxylase (TH)

and tryptophan hydroxylase 2 (TPH2), respectively, which, as the key enzymes that mediate

dopamine and serotonin synthesis, define the transmitter phenotype of these

neuromodulatory neurons [15].

These data suggest that dynamic changes in H3K27me3 in postmitotic neurons could

play a role in plasticity of cell fate or function and raise the question of the molecular

mechanisms that can control this distribution. The global depletion of H3K27me3 in neurons

takes months to appear after deletion of EZH1/2 [13] revealing that this histone modification is

remarkably stable across the bulk of the genome. Furthermore, whereas histone replacement is

a major mechanism of H3K27me3 removal in dividing cells [16], histone turnover is very slow in

postmitotic neurons [17]. Instead, lysine demethylases may have a privileged role in the local

regulation of histone methylation patterns in postmitotic neurons [18]. Both *Kdm6a* and *Kdm6b*

are expressed in neurons over the course of brain development, though only *Kdm6b* shows

enhanced expression following exposure of cells to neuronal differentiation cues [12, 19-21].

Germline knockout of *Kdm6b* does not impair the gross architecture of the embryonic brain,

but it does result in perinatal lethality caused by disruption of respiratory circuit maturation [9].

To identify functions of histone H3K27 demethylation in postmitotic stages of

neurodevelopment, we previously conditionally deleted *Kdm6a* and *Kdm6b* in *Atoh1*-expressing

cerebellar granule neuron (CGN) progenitors of the mouse. We used CGNs to model

neurodevelopment because they comprise more than 99% of all neurons [22] and 85% of cells

[23] in the cerebellum, enabling the *in vivo* comparison of chromatin across CGN

developmental states with limited contamination from other cell types. When purified and

placed in *ex vivo* culture, CGN progenitors coordinately exit the cell cycle, minimizing

heterogeneity of cell developmental stage across the population [24]. In both preparations,

gene transcription programs change over time coincident with changes in CGN biology (*e.g.,*

proliferation, migration, synapse formation, synapse maturation [25, 26]). Loss of *Kdm6a/b* had

no effect on postnatal morphogenesis of the cerebellum, demonstrating these enzymes are not

required *in vivo* for regulation of CGN progenitor proliferation. However knockdown of *Kdm6b*

in cultured CGNs impaired the developmental induction of genes during the period of

postmitotic neuronal maturation [12]. Based on these data, we hypothesized that H3K27me3

may serve as a temporal gatekeeper of these maturation genes, delaying their activation

beyond the immediate period of terminal differentiation to the neuronal fate.

Here we investigated the contribution of H3K27me3 dynamics to changing programs of

gene transcription in maturing CGNs. We used ChIP-seq and CUT&RUN to identify the genomic

location and timing of differential sites of H3K27me3 enrichment across the genome of CGNs

developing *in vivo* and in culture. We observed that H3K27me3 distributions continue to

change even after fate committed CGN progenitors exit the cell cycle, and that the loss of

H3K27me3 at gene promoters is associated with the activation of genes induced as neurons

mature. We used both genetic and pharmacological approaches to disrupt the action of

H3K27me3 regulatory enzymes and show evidence of roles for both the lysine demethylase

KDM6B and the lysine methyltransferase EZH2 in the induction of a postmitotic program of

gene transcription that underlies functional maturation of CGNs. Taken together, our data

advance understanding of how programs of gene expression are temporally orchestrated in

maturing neurons.

Materials and Methods

Animal Husbandry

We performed all procedures under an approved protocol from the Duke University Institutional Animal Care and Use Committee. *Kdm6b* floxed mice described in [27] were obtained from Jackson Labs (Stock #029615, RRID #IMSR_JAX:029615). These mice contain *loxP* sites flanking exon 14-20, encoding the catalytic Jumonji-C domain, of the mouse *Kdm6b* gene on chromosome 11. *Atoh1-cre* mice were obtained from Jackson Labs (Stock #011104, RRID #IMSR_JAX:011104). Mice were genotyped by ear clipping at the time of weaning using protocols described on the Jackson Labs website. CD-1 IGS mice (#Strain 022, #IMSR_CRL:022) and C57BL/6NCrl mice (#Strain 027, RRID #IMSR_CRL:027) were obtained from Charles River Laboratories. Both male and female mice were used for all experiments in this study.

CGN Culture

Cerebella of CD1-IGS mice were dissected at P6-P8. Cells were dissociated and run on a Percoll gradient to isolate cerebellar granule neuron progenitors (GNPs) as previously described [28]. GNPs were plated at a density of 1 million cells/well in a 24-well plate for RNA or protein isolation. At day *in vitro* 2 (DIV2) CGNs were treated with 1 µM Cytosine Arabinoside (AraC) (Sigma #C1768) to block division of any non-neuronal cells. KDM6 and EZH2 inhibitors were added at the doses and days described in the text.

RT-qPCR

CGNs were lysed in TRIzol Reagent (Thermo #15596026) at a volume of 0.25 mL per 1 million cells for RNA purification. Isolated RNA was analyzed using a nanodrop to determine concentration and purity, after which it was treated with DNase I (NEB #M0303S). We made Oligo(dT) primed cDNA using SuperscriptII Reverse Transcriptase (Invitrogen #19064014). qPCR was carried out on an Applied Biosystems Quantstudio 3 Thermal cycler using Power SYBR Green PCR Master Mix (Thermo #4367659) and PCR primers targeting the gene/locus of interest (**Supplementary Information, Table S1**).

RNA-seq

RNA for RNA-seq was obtained by processing cerebellar tissue or CGN cultures with Trizol reagent followed by clean up on the Zymo Direct-zol RNA miniprep kit (Zymo #R2052). RNA purity was measured to ensure samples had A260/280 and A260/230 values > 1.9. RNA was then polyA enriched and 150 bp paired-end sequencing was performed by Novogene, Inc on an Illumina Hi-seq 2000 machine.

Chromatin Immunoprecipitation

Cerebellar Tissue: We pooled cerebellum from three C57BL/6NCrl (Charles River Laboratories IMSR Cat# CRL:027, RRID: IMSR_CRL:027) P7 mice, two P14 mice and one P60 mouse for each biological replicate. Both male and female mice were used. Cerebellum samples were dounced in a glass homogenizer containing 2 ml of 1% formaldehyde (w/v) PBS buffer per sample and kept at 25 °C for 15 min, washed twice with cold PBS, lysed in 600 µl lysis buffer (1% SDS (w/v), 10 mM EDTA, and 50 mM Tris, pH 8.1). The cross-linked material was sonicated with a Bioruptor (Diagenode), with output set to "high" with 30 s on/off cycles to an average size range of 150–350 bp which was then determined by agarose gel electrophoresis. Sonicated chromatin was centrifuged for 10 minutes at 14000 RPM at 4°C, after which the supernatant was diluted 10-fold in dilution buffer (0.01% SDS, 1.1% Triton X-100, 1.2 mM EDTA, 16.7 mM Tris-HCl, pH 8.1, 167 mM NaCl). Prior to immunoprecipitation, 6 µl of antibody (anti-H3K27Me3, Active Motif 39155, RRID: AB_2561020) was incubated with 100 µl of Dynabeads Protein G (Invitrogen 10004D) in PBS for 4-6h at 4 °C. The antibody-bead conjugate was then washed twice with cold PBS, and added to 6 ml of cell lysis for overnight immunoprecipitation at 4°C. The following day, bead-bound DNA-protein complexes were washed, eluted, and purified using a PCR purification kit (Cat# 28104, Qiagen). ChIP-seq libraries were made using the MicroPlex Library Preparation kit V2 (C05010012, Diagenode); and 50 bp single-end sequencing was performed at the Duke Sequencing and Analysis Core Resource on a Hi-Seq 2000 machine. Three independent biological replicates were performed for each antibody and developmental time point and ChIP reagents were prepared according to the Millipore 17–295 ChIP kit.

Nuclear Isolation of CGNs for CUT&RUN

CGNs were scraped into 1X DPBS at 0.25 mL per 1 million cells and then 'pop-spun' according to the REAP method [29], until the rotor reached 7000 rpm. Pellets were then washed once with 1X DPBS and pop-spun again, after which they were resuspended in Nuclei Isolation Buffer (20 mM HEPES pH 7.9, 10 mM KCl, 2 mM Spermidine, 0.1% v/v Triton X-100, 20% v/v glycerol), incubated on ice for 5 minutes and then spun at 2,000g for 5 min at 4˚C. After this step the supernatant was removed, and pelleted nuclei were then resuspended in Nuclei Storage Buffer (20 mM Tris-HCl pH 8.0, 75 mM NaCl, 0.5 mM EDTA, 50% v/v glycerol, 1 mM DTT, 0.1 mM PMSF) and stored in -80˚C until ready to process.

CUT&RUN

CUT&RUN was performed on nuclei isolated from CGN cultures using the CUTANA ChIC/CUT&RUN kit (EpiCypher #14-1408) as per manufacturer guidelines. Specific changes made to the protocol are noted here. Nuclei in Nuclei Storage Buffer were pelleted and resuspended in Nuclei Isolation Buffer. Nuclei were then incubated with activated ConA beads. Antibodies used for CUT&RUN were: H3K27me3 (Active Motif 39155, RRID: AB_2561020), H3K4me3 and IgG (positive and negative controls included in kit). CUT&RUN libraries were made using the NEB Ultra II DNA Library Prep Kit for Illumina (NEB #E7645L), and NEBNext Multiplex Oligos for Illumina (96 Unique Dual Index Primer Pairs) (NEB #E6440S). Library cleanup was performed prior to and after PCR amplification using 0.8X Kapa Hyperpure beads (Roche #08963851001). PCR amplification was performed with the following parameters as described in the EpiCypher CUT&RUN kit: 1) 98˚C, 45 sec; 2) 98˚C, 15 sec; 60˚C, 10 sec x 14 cycles; 3) 72˚C, 60 sec. Libraries were then pooled and 50 bp paired-end sequencing was performed at the Duke Sequencing and Analysis Core Resource on a NovaSeq 6000 S-Prime flow cell.

Sequencing Data Analysis

ChIP-seq:

ChIP-seq reads were quality scored and screened through fastQC, trimmed for adapter

sequences using Trimmomatic 0.38 and then aligned to the mouse GRCm38.p4 reference

genome using STAR 2.7.2b. Alignments were filtered for mapped reads using samtools and then

normalized using the bamCoverage function of the deepTools2.0 suite. ChIP-seq tracks were

generated using bamCoverage, to generate continuous BigWig files. Peak calling was performed

using Macs2, with parameters for broad peaks and an FDR threshold of 0.01. The R package

DESeq2 [30] was used for read-count normalization and differential binding analysis. R package

ChIPseeker [31] was used to annotate ChIP-seq peaks using a window of TSS $\pm$ 3000.

'Promoters' is a group of ChIPseeker terms 'Promoter (<=1kb), Promoter (1-2 kb) and Promoter

(2-3 kb). Gene Body is a group of terms 5' UTR, 3' UTR, Intron and Exon.

CUT&RUN:

CUT&RUN reads were processed the same as ChIP-seq reads until peak-calling. Peak calling was

performed using Macs2, with parameters for broad peaks and an FDR threshold of 0.1.

Metagene plots were created using SeqPlots [32], by inputting bigwig and BED files and setting

a window of TSS +/- 5000 bp.

RNA-seq

RNA-seq reads were processed similarly to ChIP-seq reads until and excluding peak calling. Read

counts for genes were generated using HTseq2 count, with feature type set to 'exon', and the

Gencode GRCm38.p4 GTF annotation file. DESeq2 was used for read-count normalization and

differential expression analysis.

Data Processing

Heatmaps were generated using the Broad Institute's Morpheus tool:

https://software.broadinstitute.org/Morpheus, with the following clustering parameters:

hierarchical clustering, complete linkage, one minus spearman rank correlation, cluster by rows.

VST-transformed counts used to generate heatmaps and clusters are provided in Table S2. Gene

ontology analysis was performed using the GO consortium GO tool [33, 34] using terms related

to 'biological process'.

Western Blot

Cerebellar Tissue: For preparation of protein lysates from cerebellar tissue, we pooled

cerebellum from two P7 mice, one P14 and one P60 mouse for each biological replicate.

Cerebellum samples were dounced 50 times with PBS in glass homogenizer, centrifuged at 8000

x g for 5 minutes, and washed twice by cold PBS, then pellets were re-suspended in 300 μL

extraction buffer (10 mM HEPES, pH 7.9, 1.5 mM $MgCl_2$, 10 mM KCl), supplemented with 1X

protease inhibitor cocktail (Roche, Indianapolis, IN). Thereafter, 60μL of 1N HCl was added to

the cell lysates and kept on ice for 30 minutes, following which the cell lysates were centrifuged

at 11,000 x g for 10 minutes at 4 ºC. Protein concentration of acid-soluble supernatants was

measured using Pierce BCA protein assay kit (Cat # 23227, Thermo). 20 μg of acid-soluble

supernatant was used for western blots. Samples were loaded onto wells of a 10-20% gradient

gel and run at 150V for 30-40 min in running buffer (25 mM Tris-HCl, 192 mM Glycine, 0.1% w/v

SDS) after which protein was transferred onto a PVDF membrane (BioRad #1620177) in transfer

buffer (25 mM Tris-HCl, 192 mM Glycine, 20% v/v Methanol) for 1 hour at 100V or overnight at

30V. The membrane was blocked for 1 hour in 5% (w/v) BSA (Sigma #A3059) in TBST. After the

blocking step, the membrane was incubated with primary antibody overnight at 4˚C. The

following day, primary antibody was removed, after which the membranes were washed three

times with TBST for 5 minutes each, incubated with secondary antibody for 1 hour, and then

again washed three times with TBST for 5 minutes each. At the end of this, membranes were

imaged using the Li-Cor Odyssey using appropriate exposure times for the target antibodies.

CGN Culture: Cells were scraped into PBS from 6 well plates at the relevant endpoints,

centrifuged at 200 x g for 5 minutes, and washed twice with PBS. Pellets were then

resuspended in 100 μL of extraction buffer (described above), after which 20 μL of 1N HCl was

added. Samples were hereafter processed as described above.

Antibodies: The antibodies used were H3K27Ac (Abcam ab4729), H3K27Me3 (Active Motif

39155, RRID: AB_2561020), and Total H3 (Tissue: Millipore 05-499, Culture: Cell Signaling 9715).

Western blot signals were detected by Li-Cor Odyssey InfraRed Imaging System (Lincoln, NE)

with the secondary antibodies, CF770 goat anti-Rabbit, (Biotium 20078) and CF 680 goat anti-

mouse, (Biotium 20065). Protein bands were quantified using ImageJ.

GEO Accession Codes

H3K27me3 ChIP-seq data for P7, P14 and P60 cerebellum, WT and *Kdm6b*-cKO cerebellum;

RNA-seq data for WT and *Kdm6b*-cKO cerebellum; CUT&RUN and RNA-seq data for cultured

CGNs can be accessed at GEO: GSE212441. RNA-seq data for P7, P14 and P60 cerebellum and

DIV0 and DIV7 CGNs, H3K27ac, ZIC1/2 ChIP-seq data and DHS-seq data for P7 and P60

cerebellum were adapted from Frank et. al 2015 [26].

Author Contributions

Conceptualization: V.R., F.L. and A.E.W.; Methodology: V.R., F.L., U.C., and A.E.W.; Formal

Analysis: V.R., M.M.; Investigation V.R.; Resources, A.E.W.; Writing: V.R., A.E.W.; Funding

Acquisition: A.E.W.

Acknowledgements

Fig schematics in Fig 1A, 3A and 5A were created using biorender.com. This work was

supported by NIH grant R01NS0988804.

Results

<u>Cerebellar Maturation Involves Dynamic, Genome-Wide Changes in H3K27me3 Distribution</u>

To measure H3K27me3 dynamics during cerebellar maturation, we harvested cerebellar

cortex from C57BL/6NCrl mice at postnatal days 7, 14, and 60 (Fig. 1A). At P7, about half of all

cells in the mouse cerebellum are proliferating granule neuron precursors (GNPs) whereas the

rest are immature, postmitotic cerebellar granule neurons (CGNs). By P14 GNP cell division has

ceased, and the cerebellum is composed of immature postmitotic CGNs that are migrating to the
inner granule layer (IGL) where they receive synapses and fully mature by P60 [23, 26]. Western
blot of acid extracted histones showed a significant global increase in total H3K27me3 levels from
P7 to P14 that was maintained at P60 (Fig 1B), whereas total H3K27ac fell in a similar temporal
pattern (Fig. S1A). These data are consistent with the PRC2-dependent restriction of cell potential
upon terminal differentiation [4].

However, when we assessed the genomic location of H3K27me3 by ChIP-seq (Fig S1B) we
observed a more complex pattern. We found local losses as well as gains in H3K27me3
enrichment at different sites between developmental stages (Fig 1C). This was true even when
comparing P14 with P60 (Fig 1C), demonstrating that bidirectional changes in H3K27me3
distributions occur during postmitotic stages of neuronal maturation, well after the period of
terminal cell fate commitment has ended (Fig S1C). We stratified differential H3K27me3 ChIP-
seq peaks by their genomic location with respect to proximal gene promoters, gene bodies or
intergenic regions (Fig 1D), and found a consistent overrepresentation of proximal gene
promoters among developmentally demethylated peaks for all three comparisons. By contrast,
H3K27me3 ChIP-seq peaks differentially gained between all three comparisons did not differ
from the global distribution of H3K27me3, which was distributed across gene-body and distal
intergenic regions as well as proximal promoters. The percentage of total H3K27me3-positive
promoters gradually decreased by ~2-fold from P7 to P60, while that of H3K27me3-positive distal
intergenic regions increased, and that of gene bodies stayed about the same over time (Fig 1E).
Additionally, the total number of H3K27me3 peaks called centered within a 5 kb window around
the TSS appeared to gradually decrease from P7 to P60 (Fig 1E). These data demonstrate that in
CGNs H3K27me3 patterns are locally regulated in postmitotic neurons, suggesting a function for
H3K27me3-dependent gene regulation in developing neurons that extends beyond the
restriction of cell fate commitment.

H3K27me3 Dynamics at Promoters During Cerebellar Maturation Inversely Correlate with Gene
Expression

We filtered differential H3K27me3-peaks for those occurring at promoters because these

regions can be directly mapped to their likely target gene for transcriptional regulation. We then

asked whether peaks that were up- and downregulated over stages of CGN differentiation were

found at genes that can be stratified according to cellular function. Examples of genes with

promoter-associated peaks lost or gained from P7 to P60, respectively, were the genes encoding

the mature NMDA-type glutamate receptor *Grin2c* which is strongly expressed in mature CGNs

[12] and the cell-proliferation factor *Mycn,* which is expressed only in GNPs [35] (Fig 1F). This is

consistent with the hypothesis that changes in the levels of promoter H3K27me3 inversely

correlate with changes in gene expression over development.

To determine if the correlation of H3K27me3 demethylation with transcriptional

maturation held more broadly, we first performed hierarchical clustering on all differential

H3K27me3 ChIP-seq peaks and identified 3 major clusters with differing direction and timing of

change (Fig 2A). The first cluster, which we termed 'H3K27me3 Up,' contained peaks that gained

methylation as early as P14 and maintained methylation during CGN maturation. Gene Ontology

(GO) analysis demonstrated that these peaks were enriched for genes associated with progenitor

functional terms related to cell proliferation and morphogenesis. The next two clusters included

demethylated peaks, that we termed 'H3K27me3 Down (Fast)' and 'H3K27me3 Down (Slow)'

respectively, depending on whether they changed between P7 and P14 or between P14 and P60.

Both clusters were enriched for neuronal GO terms. Because developmentally gained peaks

showed a broader distribution than just promoters, we also performed the same type of filtering

and hierarchical clustering for differential H3K27me3 peaks overlying gene bodies (Fig S2). This

resulted in three clusters, two of which contained peaks that gained and lost methylation

respectively at gene-bodies. Both these clusters were enriched for neuronal GO terms including

early CGN-maturation genes. Together, these data show that bidirectional H3K27me3 dynamics

are distinct across distinct genomic locations.

We then asked if H3K27me3 dynamics at gene promoters were associated with the

expression of the corresponding RNA transcript across CGN maturation (Fig 2B, C). For this

purpose, we compared our H3K27me3 ChIP-seq to a dataset of gene expression at the same time

points from our previous study [26]. We found that 'H3K27me3 Up' peaks at gene promoters

were significantly associated with a decrease in expression of the associated gene over the same

period of developmental time and 'H3K27me3 Down' peaks were associated with an increase in

gene expression. Among the H3K27me3 Down peaks, 'Fast' peaks were associated with a

significantly greater magnitude of gene induction than 'Slow' peaks between P7 and P14, but

'Slow' peaks were associated with a significantly greater magnitude of gene induction than 'Fast'

peaks between P14 and P60. There was no significant difference between 'Fast' and 'Slow' peaks

from P7 to P60 indicating that it is the timing not the magnitude of differential developmental

gene expression that defines the two clusters. These data show that the kinetics of H3K27me3

demethylation at gene promoters are associated with the timing of gene expression changes in

postmitotic neurons, which allows for multiple postmitotic waves of gene expression (Fig. 2C).

Kdm6b-cKO Impairs CGN Maturation via Promoter Hypermethylation

The developmental loss of H3K27me3 at a subset of gene promoters in CGNs could occur

through changes in expression of the enzymes that maintain or remove these histone

modifications. At the level of gene expression, RNA-seq counts of the H3K27 methyltransferases

Ezh1/2 and the H3K27 demethylases *Kdm6a*/b showed that although there was a developmental

switch between the dominant reader and writer enzymes, the overall expression levels of each

class of enzyme did not change over time (Fig S3). However, previous *in situs* from our lab

demonstrated that *Kdm6b* expression is transiently induced in newborn CGNs in the inner EGL,

and we further showed that shRNA-mediated knockdown of *Kdm6b* in cultured CGNs impairs the

expression of a subset of developmentally induced genes [12]. Thus, we considered the

possibility that KDM6B might mediate postmitotic loss of H3K27me3 at a subset of

developmentally induced gene promoters in the developing cerebellum *in vivo* to promote their

transcriptional induction. To test this hypothesis, we conditionally knocked out *Kdm6b* in

cerebellar granule neuron precursor cells (GNPs) *in vivo* by crossing heterozygous transgenic

Atoh1-Cre mice with *Kdm6b*^{fl/fl} mice. We then harvested cerebellar tissue from littermates that

were either *Kdm6b*^{fl/fl} (WT) or *Atoh1-Cre; Kdm6b*^{fl/fl} (*Kdm6b*-cKO) at P14 as an intermediate

developmental time point for CGN maturation [26] (Fig 3A). We processed this tissue to perform

either RNA-seq (Fig S4A) or ChIP-seq for H3K27me3 (Fig S4B). We used DESeq2 to compute

differentially enriched H3K27me3 ChIP peaks between WT and *Kdm6b*-cKO tissue at P14, and

obtained 250 upregulated peaks, and 327 downregulated peaks (FDR < 0.05, |Log2FC| > 1) (Fig

3B). We then mapped these differential peaks to their genomic location (Fig 3C). We found peaks

with elevated levels of H3K27me3 in *Kdm6b*-cKO mice to be significantly more likely to be within

gene promoters compared with the overall distribution of peaks or the distribution of peaks that

had reduced H3K27me3 in *Kdm6b*-cKO mice. We observed that the percentage of promoter

associated H3K27me3 peaks decreased between WT and *Kdm6b*-cKO mice, while peaks in distal

intergenic regions and gene-bodies increased (Fig S5C). The genes containing promoters with

increased H3K27me3 in the *Kdm6b*-cKO were enriched for neuronal GO terms (Fig 3D), whereas

those containing promoters with decreased H3K27me3 in the *Kdm6b*-cKO mice failed to pull any

significant GO terms. Overall, these observations mirror our findings of H3K27 demethylation at

the promoters of neuronal maturation genes and suggests that KDM6B mediates that process.

To determine the relationship between H3K27me3 levels at promoters and gene

expression in the *Kdm6b*-cKO mice, we computed the correlation between *Kdm6b*-cKO

associated differential H3K27me3 enrichment at promoter regions, with the expression of the

associated gene in WT and *Kdm6b*-cKO mice. These data showed that the gain of H3K27me3 is

associated with a significant loss of gene expression (Fig 3E) as exemplified by the synaptic genes

Grm4 and *Gabbr2* (Fig 3F). By contrast, promoters that lost H3K27me3 in *Kdm6b*-cKO mice

showed no significant difference in gene expression (Fig 3E). These data suggest that

developmental loss of H3K27me3 at the promoters of genes induced in CGNs during postmitotic

maturation is facilitated by KDM6B, and that this loss promotes the developmental induction of

gene expression.

H3K27me3 Removal at CGN-Maturation Gene Promoters is Associated with H3K27ac Enrichment

and Increased Chromatin Accessibility

Previously we showed CGN maturation to occur in concert with broad, genome-wide

changes in H3K27ac [26] (Fig S5A). To ask whether H3K27me3 removal at promoters overlapped

with the gain of H3K27ac, we computed the overlap between genes containing promoter peaks

that were H3K27ac Up (|L2FC| > 0, FDR < 0.05) and H3K27me3 Down (Fig S5B). We observed that

only a subset of genes that gained H3K27ac also lost H3K27me3 at P60. H3K27ac Up genes that

were H3K27me3 Down (912 genes), or not H3K27me3 Down (4208 genes) were both highly

enriched for neuronal GO terms (Fig S5C).

To determine what characterizes the subset of promoters that become demethylated we

sought to ask what transcription factors might be found at these gene promoters compared to

all H3K27me3-positive promoters. To conduct an unbiased screen, we first utilized the tool called

Binding Analysis for Regulation of Transcription (BART) [36], which calculates enrichments of

previously published ChIP-seq TF binding profiles within a defined set of genomic regions. We

first filtered our ChIP-seq data for genes associated with the 'H3K27me3-Down' peak cluster

described in Fig 2A, combining 'Down (Fast)', and 'Down (Slow)' clusters, and obtained predicted

transcription factors (TF) whose binding significantly overlaps these gene promoters. We

compared the enrichment statistics of TFs enriched in these clusters to those derived from all

H3K27me3– promoter associated genes (Fig 4A). As expected, we found PRC2 components EZH2,

JARID2, SUZ12 to be significantly enriched across both the H3K27me3-Down cluster and all

H3K27me3-promoter associated genes. However, we found only the H3K27me3-Down cluster to

be significantly enriched for the putative binding of transcription factors ZIC1 and ZIC2, which

previous work from our group has shown to be a driver of CGN maturation [26]. Importantly,

when we assessed the 'H3K27me3-Down (Fast)', and 'H3K27me3-Down (Slow)' clusters

separately (Fig S6A), they were also enriched for the binding of ZIC1 and ZIC2, which is consistent

with evidence [26] that these TFs function at multiple stages of CGN differentiation. By contrast,

when we performed the same analysis for the 'H3K27me3-Up' cluster (Fig S6A), we found this

set of promoters was significantly enriched for the binding of a distinct set of TFs involved in cell

differentiation, namely SOX3, FOXF1, SHOX2, POU3F2 as well as histone deacetylation (HDAC1)

[37].

To directly compare changes in ZIC1/2 TF binding to changes in H3K27me3 across CGN

development, we re-analyzed our previously published ZIC1/2 ChIP-seq data [26] and measured

ZIC1/2 binding at All Genes, Genes with H3K27me3 Up, and Genes with H3K27me3 Down (Fig

4A). We then identified gene promoters with increased ZIC binding at P60 compared with P7. We

termed these genes 'ZIC Up' and first compared them to the H3K27me3-Down cluster. We found

that 56% of H3K27me3-Down genes were also ZIC-Up, and that these genes were enriched for

neuronal GO terms (Fig 4B). As a control, we compared 'ZIC Up' genes to H3K27me3-Up genes

(Fig. S6A) and only found an overlap of only 11%. We repeated this analysis with the H3K27me3

ChIP-seq peaks that were differential at P14 between WT and *Kdm6b*-cKO (Fig. 3) and found a

similar enrichment of putative ZIC1 and ZIC2 binding within gene promoters associated with

H3K27me3 peaks 'Up in *Kdm6b*-cKO' (Fig S6B), but not 'Down in *Kdm6b*-cKO' (Fig S6C). 72% of

the 'H3K27me3 Up in *Kdm6b*-cKO' genes were also 'ZIC Up' (Fig S6B) and were enriched for

neuronal GO terms. Only 9% of 'H3K27me3 Down in *Kdm6b*-cKO' were also 'ZIC Up' (Fig S6C). As

examples of gene promoters that belong to both the H3K27me3-Down cluster, and H3K27me3

Up peaks due to *Kdm6b*-cKO, we show tracks for *Grin2c* and *Grm4,* that are strongly enriched for

ZIC1/2 at their promoters at P60 (Fig 4C).

We have shown previously [26] that increased ZIC1/2 binding strongly correlates with the

developmental enrichment of H3K27ac and increased chromatin accessibility during cerebellar

maturation. Given the strong enrichment of ZIC1/2 binding within the H3K27me3-Down cluster

during development, and H3K27me3 Up peaks due to *Kdm6b*-cKO, we wanted to ask whether

H3K27me3 turnover during cerebellar maturation is associated with H3K27ac enrichment and

changes in chromatin accessibility. As such we computed the distribution of Log2FC values from

our previously published H3K27ac-ChIP-seq and DNAse Hypersensitivity data [26] comparing P7

and P60 cerebellum and filtered these distributions for those related to peaks belonging to our

differential H3K27me3 promoter clusters. Indeed, H3K27me3 Down peaks at P60 were

significantly associated with an increase in H3K27ac enrichment (Fig 4D) and DNAse

hypersensitivity (Fig 4E). H3K27me3-Up peaks due to *Kdm6b*-cKO conversely were associated

with genes that became enriched with H3K27ac, and DNAse hypersensitive regions at P60 (Fig

S6D). As examples, we show H3K27ac ChIP-seq and DNAse-seq tracks for 'H3K27me3 Down' gene

Grin2c, and 'H3K27me3 Up' gene *Atoh1*. Thus, these data provide further evidence that the

postmitotic removal of H3K27me3 from a subset of gene promoters is important for the

coordinated induction of gene expression that underlies neuronal maturation.

<u>Modulation of H3K27me3 Turnover Mediates CGN Maturation Gene-Regulation in Culture</u>

To determine the functional consequences of modulating H3K27me3 at CGN-maturation

gene promoters or gene bodies we used cultures of primary mouse CGNs as context for

pharmacological manipulation of histone regulatory enzymes. When GNPs isolated from P6-8

mouse cerebellum are placed in culture they show changes in gene expression over the course

of 7 days [26] (Fig 5A, S7A, B). We performed CUT&RUN-seq to measure genome-wide changes

in H3K27me3 after 1, 3, 5 and 7 days *in vitro* (DIV) (Fig 5A). We found methylation to be gained

quickly after cell cycle exit, with 1688 peaks significantly up between DIV1 and DIV3 CGNs,

followed by 109 peaks between DIV3 and DIV5, and 2 peaks between DIV5 and DIV7 CGNs (Fig

5B). Like our observations *in vivo* (Fig 1, 2), we saw that demethylation occurred more gradually,

with 477 peaks significantly down between DIV1 and DIV3 CGNs, followed by 227 and 25 peaks

between DIV3 and DIV5, and DIV5 and DIV7 CGNs respectively. Again, consistent with our

findings *in vivo*, we found demethylation to occur primarily at promoters (Fig 5C, Fig S7C), and

gains in methylation to occur more uniformly, including at gene-bodies and distal intergenic

regions. When we filtered for differential H3K27me3 peaks at gene promoters, they clustered in

two groups that gain H3K27me3 or lose H3K27me3 at DIV7 (Fig 5D). Genes whose promoters

were in the 'Gain H3K27me3 at DIV7' cluster were associated with cell fate specification GO terms

as predicted by canonical PRC functions. Conversely, genes whose promoters were in the 'Lose

H3K27me3 at DIV7' cluster were associated with functional neuronal GO terms. We observed

mature synaptic genes such as *Grin2c* and *Grm4* to lose H3K27me3 from DIV1 to DIV7 (Fig 5E

lower), and proliferation genes expressed in GNPs such as *Myc* and *Atoh1* to gain H3K27me3

from DIV1 to DIV7 in culture (Fig 5E upper). Finally, we stratified our previously published RNA-

seq data between DIV0 and DIV7 CGNs by their respective methylation status ('Gain H3K27me3

at DIV7' or 'Lose H3K27me3 at DIV7' cluster) and calculated the distribution of Log2FC (Fig 5F).

We found peaks associated with the 'Gain H3K27me3 at DIV7' cluster to have a significant

reduction in gene expression over time in culture, and those associated with the 'Lose H3K27me3

at DIV7' cluster to have a significant gain in gene expression by DIV7 (Fig 5F, S7D, E). These data

show that we can use this CGN culture system to explore mechanisms and causality with respect

to the enzymes mediating H3K27me3 turnover and gene regulation during CGN maturation.

To determine the functional contribution of H3K27me3 regulation to CGN-maturation

gene expression, we treated CGN cultures at DIV1 with small molecule inhibitors of either EZH2

or KDM6A/B, called GSK-126 and GSK-J4 respectively, and measured the consequences for cell

viability, histone modification, and gene expression at DIV5. Because these drugs have not been

widely used in primary neurons, we first performed a cell viability assay and measured the IC_{50}

values of both drugs. We determined the IC_{50} values to be ~3.07 µM for GSK-J4 and ~7.32 µM for

GSK-126, respectively (Fig S8A). For the KDM6B inhibitor, the IC_{50} was nearly 10-fold below the

dose (25 µM) used in the original characterization studies to block the histone demethylase

activity of these enzymes [38]. Thus, we sought to determine if these drugs could block enzymatic

function in viable cells. Choosing concentrations below the measured IC_{50} values, we first

measured global levels of H3K27me3 by western blot (Fig 6A), after treating with 0, 0.5, 1.0 and

2.0 µM of either drug. Treatment of CGNs with GSK-126 caused a dose-dependent loss of

H3K27me3 validating the method. GSK-126 treatment also appeared to cause a global increase

(Fig S8B) in H3K27ac, which is consistent with the idea that H3K27me3 and H3K27ac are mutually

exclusive modifications that oppose each other's functions [39]. By contrast, GSK-J4 treatment

failed to cause any dose-dependent change in H3K27me3 or H23K27ac detected by western.

Given our evidence that KDM6B facilitates demethylation of a subset of gene promoters

over development (Fig 2), even in the absence of global reductions in H3K27me3 (Fig 1), we asked

whether there might be local changes in H3K27me3 after GSK-J4 treatment. We treated CGNs

daily with 1 µM of GSK-J4 between DIV1-DIV5 and then assessed H3K27me3 peaks across the

genome by CUT&RUN-seq to determine if there were local increases in H3K27me3 at a subset of

genes (Fig S7C-D). We failed to observe any significant changes in H3K27me3 enrichment across

the genome at this concentration of GSK-J4 (Fig S8E), with biological replicate being the primary

factor for variation in this dataset (Fig S8D). We did find 1µM GSK-J4 treatment to slightly

decrease mRNA levels of specific neuronal function genes (Fig S8F), suggesting that the drug may

act to regulate genes in a demethylation independent manner in CGNs. We conclude that the

therapeutic range of GSK-J4 is too small in our CGNs cultures to permit its use for inhibition of

KDM6-dependent demethylation.

Our data showed that EZH2-inhibiting doses of GSK-126 were compatible with CGN

viability, thus we then used CUT&RUN-seq for H3K27me3 on CGNs treated with 5 µM GSK-126

(Fig 6B, Fig S9A). Most of the regions that showed differential methylation following GSK-126

treatment had lower levels of H3K27me3 in GSK-126 treated neurons, consistent with the

inhibition of this histone methyltransferase (Fig 6B). A much smaller number of peaks showed

increases in H3K27me3, which may arise from secondary effects of EZH2 inhibition on other

processes that impact H3K27me3 distributions such as EZH1 function [40] or histone turnover

[41]. Regions that lost H3K27me3 after GSK-126 treatment showed a similar genomic distribution

to the overall distribution of H3K27me3 peaks in CGNs (Fig 9B-C). Demethylated peaks were

enriched for GO terms associated with several non-neuronal cell types, consistent with the

general hypothesis that PRC functions in postmitotic cells at least in part to maintain the chosen

cell fate [14] (Fig S9D).

To determine how EZH2 activity influences H3K27me3 distribution at early genes, we

created metagene plots (Fig 6C) describing H3K27me3 coverage across all H3K27me3 peaks

compared with the differential peaks from Fig 5D. We observed GSK-126 treatment to

significantly reduce H3K27me3 coverage across the 'Gain H3K27me3 at DIV7' cluster but not the

'Lose H3K27me3 at DIV7' cluster or at all H3K27me3 peaks. To determine the effects of GSK-126

treatment on gene expression, we performed RNA-seq and identified 259 significantly

downregulated and 393 significantly upregulated genes (Fig 6D, (upper)). Upregulated genes

included *Mycn*, *Ebf3* and *Jag1* and were enriched for GO terms including 'regulation of cell

proliferation', while downregulated genes included *Gabrd*, *Gabra6*, *Grm4* and were enriched for

synaptic GO terms (Fig 6D, (lower)). We then computed gene expression profiles of CGNs

maturing in culture (from DIV0 to DIV7) to CGNs treated with GSK-126 (Fig 6 left). We found

genes repressed due to GSK-126 treatment to be strongly induced at DIV7 compared to DIV0.

Conversely, we saw genes upregulated at DIV0 such as *Mycn* and *Ebf3* to be induced upon GSK-

126 treatment. Furthermore, we computed changes to gene expression in response to GSK-126

treatment as a function of clusters described in Fig 5E – 'All peaks', 'Gain H3K27me3 at DIV7' and

'Lose H3K27me3 at DIV7' (Fig 6E right). We observed GSK-126 treatment to significantly

upregulate the expression of genes belonging to the 'Gain H3K27me3 at DIV7' cluster, and

significantly downregulate the expression of genes belonging to the 'Lose H3K27me3 at DIV7'

cluster (Fig 6D, (upper right)). Visualizing H3K27me3 and RNA-seq tracks at early gene *Ebf3*, we

observed GSK-126 treatment to completely block H3K27me3 and induce gene expression

compared to DMSO treatment (Fig 6F upper). The late synaptic gene *Gabrd*, conversely,

remained unchanged with respect to H3K27me3 enrichment, but was strongly downregulated in

response to GSK-126 treatment (Fig 6F lower). These data show that EZH2 functions in newly

postmitotic neurons to add H3K27me3 to genes that were not methylated at the time of cell fate

commitment, and that this methylation helps to repress expression of progenitor genes [42] in

neurons. By contrast, EZH2 is not required over this time course for the maintenance of

H3K27me3 at genes that possessed the modification at the time of terminal differentiation,

consistent with the slow turnover of methylated histones in postmitotic neurons. Why EZH2

catalytic activity early on during CGN maturation is critical for the induction of late genes is a

point we discuss below.

Discussion

590
591

In this study, we investigated the role of H3K27me3 turnover in postmitotic gene

regulation, using the maturation of mouse CGNs as a model system. PRC2 activity has long been

studied as a mechanism of early cell-fate specification in mammals through the deposition of

H3K27me3 and silencing of fate defining genes during the differentiation of embryonic stem cells

(ESCs) into other cell types including neural progenitor cells (NPCs) and early neurogenesis [5, 43,

44]. Our study demonstrates that dynamic changes in H3K27me3 continue in postmitotic

neurons at specific sites in the genome to coordinate the temporal induction of gene expression

programs that underlie functional maturation. Increases in H3K27me3 occurred rapidly after cell

cycle exit, whereas promoter demethylation occurred gradually and these turnover kinetics

strongly correlated with the kinetics of induced gene expression. Promoters and gene bodies that

gained H3K27me3 after cell cycle exit were enriched for genes involved in proliferation and cell-

fate specification such as *Atoh1* and *Myc*. By contrast, developmentally demethylated promoters

included those for genes that confer mature function on synapses such as the NMDA receptor

subunit *Grin2c* and the metabotropic glutamate receptor *Grm4*. This suggested to us that there

are temporally coordinated mechanisms of enzymatic H3K27me3 turnover during CGN

maturation, and that postmitotic H3K27me3 demethylation may be a key step for timing

functional synaptic maturation.

Our *in vivo* cerebellar data show that developmental loss of H3K27me3 from gene

promoters is facilitated by the H3K27me3 demethylase KDM6B, and further that conditional

knockout of KDM6B in CGNs impaired the expression of genes that are induced as CGNs mature.

Kdm6b expression is upregulated in newborn CGNs in the inner EGL [12], and we speculate that

the temporal increase in expression of this enzyme may tip the balance of H3K27me3 regulation

to favor histone demethylation at specific target sites. Our evidence that genes with increased

H3K27me3 in the *Kdm6b*-cKO mice overlap those that show developmental loss of H3K27me3 is

consistent with the possibility that KDM6B functions enzymatically to demethylate histones at

these gene promoters. However, chromatin regulators including KDM6B have scaffolding

functions as well as enzymatic functions [45-49], thus we attempted to use pharmacological

inhibition of KDM6B to validate the role of its enzymatic activity in developing CGN gene

regulation. The small molecule GSK-J4 is a highly selective inhibitor of the KDM6 family compared

with other histone demethylases, and it was previously demonstrated in primary human

macrophages to block LPS-mediated histone demethylation and induction of TNFα expression

with an IC_{50} of 9 μM and no apparent toxicity up to 30 μM [38]. By contrast we found 50%

cytotoxicity in CGNs at a GSK-J4 concentration of just 3.1 μM and no evidence by western blot or

CUT&RUN for any global or local changes in H3K27me3 when the drug was applied at the non-

toxic dose of 1 μM. Moving forward, as an alternative to pharmacological inhibition of KDM6B,

genetic mutation of the JmjC domain of KDM6B may be a more effective means to test its

functional requirement in neuronal H3K27me3 demethylation [50]. Indeed, a prior study showed

that in a BAC transgene, enzymatically dead KDM6B was unable to rescue perinatal death in

KDM6B knockout mice, which is consistent with a requirement for the enzymatic function of

KDM6B in maturation of the respiratory circuit [9].

Our observation that H3K27me3 removal occurs at only a subset of genes that are

developmentally induced during CGN maturation raises the question of how these promoters are

marked for methylation and demethylation. Through an unbiased approach, we showed that

promoters that lose H3K27me3 during CGN maturation are strongly enriched for binding of the

transcription factors ZIC1/2; thus, one possibility is that an interaction between KDM6B and the

ZICs selects these genes for induction in maturing neurons. ZIC1/2 are strong drivers of CGN

maturation [26], and mutation of these genes is associated with cerebellar disorders, showing

their importance in CGN development [51, 52]. ZIC1 and 2 are zinc-finger binding proteins that

exhibit dynamic DNA binding properties during cerebellar maturation, and they regulate the

expression of both early and late CGN maturation genes [26]. Given the strong association of

H3K27ac enrichment at these promoters, however, it is likely that ZIC1/2 binding is a

consequence and not a cause of H3K27-demethylation at promoters. It would therefore be

essential to measure the genome-wide enrichment patterns of KDM6B during CGN maturation,

in order to identify its binding partners and direct targets. Existing literature investigating KDM6B

dynamics across the epigenome is sparse owing to the lack of well validated, commercially

available KDM6B antibodies. Nevertheless, KDM6B has been shown to localize to promoters and

enhancers of neurogenic genes in NSCs [53, 54], and more recently to induce muscle

regeneration gene *Has2*, by removing H3K27me3 from its promoter, to initiate muscle repair in

response to injury [50]. KDM6B targeting to these specific loci might be facilitated through

interactions with TFs [54, 55], and chromatin remodeling complexes [56]. For example, KDM6B

has been shown to strongly associate with the ATP-dependent chromatin remodeler mSWI/SNF

or BAF complex, by interacting with its subunits which in turn potentiate its activity [57]. This

interaction has been shown to play a role in neuronal development [57, 58].

Interestingly, we find that maturation of CGN transcriptional programs not only depends

on the loss of H3K27me3 at genes that turn on late, but also requires the addition of H3K27me3

at regions that gain this mark in newly postmitotic neurons. Using CUT&RUN in cultured CGNs,

we discovered that the rapid increases in H3K27me3 that accumulate in newly postmitotic

neurons are highly sensitive to inhibition of EZH2 with GSK-126. Failure to gain H3K27me3 at

these sites in differentiating neurons not only prevented the developmental downregulation of

CGN progenitor markers like *Mycn* and *Ebf3* but also impaired the developmental induction of

late genes like *Gabrd*, *Grm4* and *Grin2c*. The impact of GSK-126 on the mature CGN

transcriptional program was not due to changes in H3K27me3 regulation at the late genes

themselves. This shows that the postmitotic addition of H3K27me3 at specific genomic sites is

crucial in regulating the induction of genes that turn on well after neuronal cell fate is

determined. However, these data also raise the question of how PRC2 complex function can

indirectly regulate the activation of late gene expression programs over a period of days. One

possibility is that the persistent expression of early TFs like MYCN and EBF3 interferes with the

expression or function of the TFs that activate late gene promoters even once these promoters

lose H3K27me3. However, another possibility may be via the effects of H3K27me3 on chromatin

architecture. Though H3K27me3 is an acutely repressive mark, PRC2 dependent deposition of

H3K27me3 has been shown in some cases to contribute to physical interactions between

enhancers and their target genes by anchoring chromatin loops [59, 60]. These loops may then

poise genes for enhancer activation later upon receiving a developmental stimulus. The gain of

H3K27me3 in early differentiating CGNs could serve a similar function to establish such enhancer

contacts with the promoters of late genes such that when KDM6B demethylates these

promoters, the enhancer can permit activation. In the future, investigating dynamics in higher-

order chromatin structure through low-input techniques such as HiCAR [61] might help inform

how H3K27me3 dynamics contribute to gene regulation in postmitotic neurons.

References

1. Hyun, K., et al., *Writing, erasing and reading histone lysine methylations.* Experimental & Molecular Medicine, 2017. **49**(4): p. e324-e324.

2. Jambhekar, A., A. Dhall, and Y. Shi, *Roles and regulation of histone methylation in animal development.* Nature Reviews Molecular Cell Biology, 2019. **20**(10): p. 625-641.

3. Daniel, J.A., M.G. Pray-Grant, and P.A. Grant, *Effector proteins for methylated histones: an expanding family.* Cell Cycle, 2005. **4**(7): p. 919-26.

4. van Mierlo, G., et al., *The Complexity of PRC2 Subcomplexes.* Trends in Cell Biology, 2019. **29**(8): p. 660-671.

5. O'Carroll, D., et al., *The polycomb-group gene Ezh2 is required for early mouse development.* Mol Cell Biol, 2001. **21**(13): p. 4330-6.

6. Voncken, J.W., et al., *Rnf2 (Ring1b) deficiency causes gastrulation arrest and cell cycle inhibition.* Proceedings of the National Academy of Sciences, 2003. **100**(5): p. 2468.

7. Swigut, T. and J. Wysocka, *H3K27 Demethylases, at Long Last.* Cell, 2007. **131**(1): p. 29-32.

8. Welstead, G.G., et al., *X-linked H3K27me3 demethylase Utx is required for embryonic development in a sex-specific manner.* Proc Natl Acad Sci U S A, 2012. **109**(32): p. 13004-9.

9. Burgold, T., et al., *The H3K27 demethylase JMJD3 is required for maintenance of the embryonic respiratory neuronal network, neonatal breathing, and survival.* Cell Rep, 2012. **2**(5): p. 1244-58.

10. Margueron, R. and D. Reinberg, *The Polycomb complex PRC2 and its mark in life.* Nature, 2011. **469**(7330): p. 343-9.

11. Wijayatunge, R., et al., *The histone lysine demethylase Kdm6b is required for activity-dependent preconditioning of hippocampal neuronal survival.* Mol Cell Neurosci, 2014. **61**: p. 187-200.

12. Wijayatunge, R., et al., *The histone demethylase Kdm6b regulates a mature gene expression program in differentiating cerebellar granule neurons.* Molecular and Cellular Neuroscience, 2018. **87**: p. 4-17.

13. von Schimmelmann, M., et al., *Polycomb repressive complex 2 (PRC2) silences genes responsible for neurodegeneration.* Nature Neuroscience, 2016. **19**(10): p. 1321-1330.

14. Ferrai, C., et al., *RNA polymerase II primes Polycomb-repressed developmental genes throughout terminal neuronal differentiation.* Molecular Systems Biology, 2017. **13**(10): p. 946.

15. Toskas, K., et al., *PRC2-mediated repression is essential to maintain identity and function of differentiated dopaminergic and serotonergic neurons.* Science Advances. **8**(34): p. eabo1543.

16. Shpargel, K.B., et al., *KDM6 demethylase independent loss of histone H3 lysine 27 trimethylation during early embryonic development.* PLoS Genet, 2014. **10**(8): p. e1004507.

17.   Maze, I., et al., *Critical Role of Histone Turnover in Neuronal Transcription and Plasticity.*
Neuron, 2015. **87**(1): p. 77-94.

18.   Swahari, V. and A.E. West, *Histone demethylases in neuronal differentiation, plasticity,*
and disease. Curr Opin Neurobiol, 2019. **59**: p. 9-15.

19.   Jepsen, K., et al., *SMRT-mediated repression of an H3K27 demethylase in progression from*
neural stem cell to neuron. Nature, 2007. **450**(7168): p. 415-9.

20.   Burgold, T., et al., *The histone H3 lysine 27-specific demethylase Jmjd3 is required for*
neural commitment. PLoS One, 2008. **3**(8): p. e3034.

21.   De Santa, F., et al., *Jmjd3 contributes to the control of gene expression in LPS-activated*
macrophages. Embo j, 2009. **28**(21): p. 3341-52.

22.   Consalez, G.G., et al., *Origins, Development, and Compartmentation of the Granule Cells*
of the Cerebellum. Frontiers in Neural Circuits, 2021. **14**: p. 88.

23.   Altman, J., *Development of the Cerebellar System.* Relation to its Evolution, Structure, and
Functions., 1997.

24.   Fogarty, M.P., et al., *Fibroblast growth factor blocks Sonic hedgehog signaling in neuronal*
precursors and tumor cells. Proceedings of the National Academy of Sciences, 2007.
104(8): p. 2973-2978.

25.   Sathyanesan, A., et al., *Emerging connections between cerebellar development, behaviour*
and complex brain disorders. Nature Reviews Neuroscience, 2019. **20**(5): p. 298-313.

26.   Frank, C.L., et al., *Regulation of chromatin accessibility and Zic binding at enhancers in the*
developing cerebellum. Nature Neuroscience, 2015. **18**(5): p. 647-656.

27.   Manna, S., et al., *Histone H3 Lysine 27 demethylases Jmjd3 and Utx are required for T-cell*
differentiation. Nat Commun, 2015. **6**: p. 8152.

28.   Bilimoria, P.M. and A. Bonni, *Cultures of cerebellar granule neurons.* CSH Protoc, 2008.
2008: p. pdb.prot5107.

29.   Suzuki, K., et al., *REAP: A two minute cell fractionation method.* BMC Research Notes,
2010. **3**(1): p. 294.

30.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion*
for RNA-seq data with DESeq2. Genome Biology, 2014. **15**(12): p. 550.

31.   Yu, G., L.-G. Wang, and Q.-Y. He, *ChIPseeker: an R/Bioconductor package for ChIP peak*
annotation, comparison and visualization. Bioinformatics, 2015. **31**(14): p. 2382-2383.

32.   Stempor, P. and J. Ahringer, *SeqPlots - Interactive software for exploratory data analyses,*
pattern discovery and visualization in genomics [version 1; peer review: 2 approved, 1
approved with reservations]. Wellcome Open Research, 2016. **1**(14).

33.   Ashburner, M., et al., *Gene ontology: tool for the unification of biology. The Gene Ontology*
Consortium. Nat Genet, 2000. **25**(1): p. 25-9.

34.   *The Gene Ontology resource: enriching a GOld mine.* Nucleic Acids Res, 2021. **49**(D1): p.
D325-d334.

35.   Ma, M., et al., *N-myc is a key switch regulating the proliferation cycle of postnatal*
cerebellar granule cell progenitors. Scientific Reports, 2015. **5**(1): p. 12740.

36.   Wang, Z., et al., *BART: a transcription factor prediction tool with query gene sets or*
epigenomic profiles. Bioinformatics, 2018. **34**(16): p. 2867-2869.

37.   Wang, T.-W., et al., *Sox3 expression identifies neural progenitors in persistent neonatal*
768   *and adult mouse forebrain germinative zones.* Journal of Comparative Neurology, 2006.
769   **497**(1): p. 88-100.
770   38.   Kruidenier, L., et al., *A selective jumonji H3K27 demethylase inhibitor modulates the*
771   *proinflammatory macrophage response.* Nature, 2012. **488**(7411): p. 404-408.
772   39.   Lavarone, E., C.M. Barbieri, and D. Pasini, *Dissecting the role of H3K27 acetylation and*
773   *methylation in PRC2 mediated control of cellular identity.* Nature Communications, 2019.
774   **10**(1): p. 1679.
775   40.   Shen, X., et al., *EZH1 mediates methylation on histone H3 lysine 27 and complements EZH2*
776   *in maintaining stem cell identity and executing pluripotency.* Mol Cell, 2008. **32**(4): p. 491-
777   502.
778   41.   Chory, E.J., et al., *Nucleosome Turnover Regulates Histone Methylation Patterns over the*
779   *Genome.* Mol Cell, 2019. **73**(1): p. 61-72.e3.
780   42.   Consalez, G.G., et al., *Origins, Development, and Compartmentation of the Granule Cells*
781   *of the Cerebellum.* Front Neural Circuits, 2020. **14**: p. 611841.
782   43.   Mohn, F., et al., *Lineage-Specific Polycomb Targets and De Novo DNA Methylation Define*
783   *Restriction and Potential of Neuronal Progenitors.* Molecular Cell, 2008. **30**(6): p. 755-766.
784   44.   Pereira, J.D., et al., *Ezh2, the histone methyltransferase of PRC2, regulates the balance*
785   *between self-renewal and differentiation in the cerebral cortex.* Proceedings of the
786   National Academy of Sciences, 2010. **107**(36): p. 15957-15962.
787   45.   Kim, E., et al., *Phosphorylation of EZH2 activates STAT3 signaling via STAT3 methylation*
788   *and promotes tumorigenicity of glioblastoma stem-like cells.* Cancer Cell, 2013. **23**(6): p.
789   839-52.
790   46.   Kim, J., et al., *Polycomb- and Methylation-Independent Roles of EZH2 as a Transcription*
791   *Activator.* Cell Rep, 2018. **25**(10): p. 2808-2820.e4.
792   47.   Miller, S.A., S.E. Mohn, and A.S. Weinmann, *Jmjd3 and UTX play a demethylase-*
793   *independent role in chromatin remodeling to regulate T-box family member-dependent*
794   *gene expression.* Mol Cell, 2010. **40**(4): p. 594-605.
795   48.   Xun, J., et al., *JMJD3 suppresses stem cell-like characteristics in breast cancer cells by*
796   *downregulation of Oct4 independently of its demethylase activity.* Oncotarget, 2017.
797   **8**(13): p. 21918-21929.
798   49.   Ohguchi, H., et al., *KDM6B modulates MAPK pathway mediating multiple myeloma cell*
799   *growth and survival.* Leukemia, 2017. **31**(12): p. 2661-2669.
800   50.   Nakka, K., et al., *JMJD3 activated hyaluronan synthesis drives muscle regeneration in an*
801   *inflammatory environment.* Science, 2022. **377**(6606): p. 666-669.
802   51.   Twigg, S.R., et al., *Gain-of-Function Mutations in ZIC1 Are Associated with Coronal*
803   *Craniosynostosis and Learning Disability.* Am J Hum Genet, 2015. **97**(3): p. 378-88.
804   52.   Blank, M.C., et al., *Multiple developmental programs are altered by loss of Zic1 and Zic4*
805   *to cause Dandy-Walker malformation cerebellar pathogenesis.* Development, 2011.
806   **138**(6): p. 1207-16.
807   53.   Park, Dae H., et al., *Activation of Neuronal Gene Expression by the JMJD3 Demethylase Is*
808   *Required for Postnatal and Adult Brain Neurogenesis.* Cell Reports, 2014. **8**(5): p. 1290-
809   1299.

54.   Estarás, C., et al., *Genome-wide analysis reveals that Smad3 and JMJD3 HDM co-activate*
the neural developmental program. Development, 2012. **139**(15): p. 2681-2691.

55.   Dai, J.P., et al., *Jmjd3 activates Mash1 gene in RA-induced neuronal differentiation of P19*
cells. J Cell Biochem, 2010. **110**(6): p. 1457-63.

56.   De Santa, F., et al., *The Histone H3 Lysine-27 Demethylase Jmjd3 Links Inflammation to*
Inhibition of Polycomb-Mediated Gene Silencing. Cell, 2007. **130**(6): p. 1083-1094.

57.   Narayanan, R., et al., *Loss of BAF (mSWI/SNF) Complexes Causes Global Transcriptional*
and Chromatin State Changes in Forebrain Development. Cell Reports, 2015. **13**(9): p.
1842-1854.

58.   Nguyen, H., et al., *Epigenetic Regulation by BAF Complexes Limits Neural Stem Cell*
Proliferation by Suppressing Wnt Signaling in Late Embryonic Development. Stem Cell
Reports, 2018. **10**(6): p. 1734-1750.

59.   Cruz-Molina, S., et al., *PRC2 Facilitates the Regulatory Topology Required for Poised*
Enhancer Function during Pluripotent Stem Cell Differentiation. Cell Stem Cell, 2017. **20**(5):
p. 689-705.e9.

60.   Kraft, K., et al., *Polycomb-mediated genome architecture enables long-range spreading of*
H3K27 methylation. Proceedings of the National Academy of Sciences, 2022. **119**(22): p.
e2201883119.

61.   Wei, X., et al., *HiCAR is a robust and sensitive method to analyze open-chromatin-*
associated genome organization. Molecular Cell, 2022. **82**(6): p. 1225-1238.e6.

Figures and Captions

Fig 1. The maturing mouse cerebellum undergoes a genome-wide redistribution of H3K27me3 dominated by loss at promoters

A) CD1 mice were sacrificed at postnatal days 7, 14 and 60 after which cerebellar tissue was dissected out and processed to harvest histones, chromatin and RNA. B) (Left) Western blot of acid extracted histone from cerebellar tissue for H3K27me3 and total Histone H3 (n=3 biological replicates), (Right) Quantification of Western Blot, one-way ANOVA, * indicates $p < 0.05$. C) MA plot showing H3K27me3 peaks gained (Up) and lost (Down) throughout cerebellar maturation *in vivo* (n=3 biological replicates, differential enrichment calculated using DESeq2 package. Up and Down are loci with FDR < 0.05 and |L2FC| > 1). D) Percentage of differential H3K27me3 peaks between P7-P14, P14-P60 and P7-P60, annotated by genomic region (Annotation performed using ChIPseeker package, TSS +/- 3000 bp). E) (Left) Percentage of H3K27me3 peaks annotated by genomic region, (Right) Heatmap of H3K27me3 peaks centered around TSS +/- 5000 *bp*. F)

ChIP-seq tracks of example 'Down' gene *Grin2c* (Upper) and 'Up' gene *Myc* (lower) (y-axes for

ChIP-seq tracks are autoscaled according to data range in current view).

Fig 2. Differentially methylated promoters cluster by turnover kinetics and correlate with CGN

maturation gene expression

A) (Left) Heatmap of hierarchically clustered, VST-transformed DESeq2-normalized counts of

differential H3K27me3 promoter peaks from P7-P60, (Right) Corresponding Gene Ontology (GO)

Terms associated with nearest gene and FDR. B) Violin plot showing the distribution of Log2FC of

gene expression as a function of clustering performed in A, across developmental time (one-way

ANOVA, * indicates $p < 0.05$). C) Example ChIP-seq tracks of genomic loci belonging to clusters

described in A, at *Mycn*, *Grm4* and *Grin2c* loci and corresponding RNA-seq tracks (y-axes for ChIP-

seq and RNA-seq tracks are autoscaled separately, according to data range in current view).

Fig 3. *Kdm6b* knockout in CGN precursors impairs CGN maturation via H3K27me3

hypermethylation of CGN maturation gene promoters

A) *Atoh1-Cre*; *Kdm6b^{fl/fl}* (cKO) and *Kdm6b^{fl/fl}* (WT) mice were generated by crossing *Atoh1-Cre*;

Kdm6b^{fl/+} mice to *Kdm6b^{fl/fl}* mice and sacrificed at P14 for cerebellar tissue harvest and

subsequent performing of H3K27me3 ChIP-seq (WT n=2, cKO n=3 biological replicates) and RNA-

seq (n=2 biological replicates). B) MA plot showing differential enrichment of genome wide

H3K27me3 peaks between WT and cKO mice (|Log2FC| > 1, FDR < 0.05). C) Differentially

methylated H3K27me3 between WT and cKO annotated by genomic location using ChIPseeker

package, TSS +/- 3000 bp. D) GO Terms and FDR for genes nearest genomic promoters within

H3K27me3 Up and Down (due to *Kdm6b*-cKO) peaks. E) Distribution of Log_2Fold Change

Expression (WT v cKO, computed using DESeq2) as a function of unchanged, H3K27me3 Up and

H3K27me3 Down genomic promoters between WT and cKO (one-way ANOVA, * indicates $p <$

0.05). F) Representative genes containing H3K27me3 Up genomic promoter peaks *Grm4* and

Gabbr2 and corresponding RNA-seq tracks between WT and cKO mice (y-axes for ChIP-seq and

RNA-seq tracks are autoscaled separately, according to data range in current view).

Fig 4. H3K27me3 removal is followed by gain of H3K27me3 and chromatin accessibility at gene

promoters

A) (Left) Binding Analysis for Regulation of Transcription (BART) for genes near H3K27me3 Down

peaks (x-axis) and near all H3K27me3 peaks (y-axis) plotted by -$\log_{10}$p-adj. (Right) Metagene plots

for ZIC ChIP-seq signal at genes near all H3K27me3 peaks, genes with H3K27me3 Up and genes

with H3K27me3 down. B) (Left) Percentage of genes near H3K27me3 Down promoter peaks that

are also ZIC1/2 Up ($\log_2$Fold Change > 1, P60/P7, ZIC1/2 ChIP-seq). (Right) GO Terms and FDR for

genes nearest to H3K27me3 Down promoter peaks that are also ZIC1/2 Up, and not ZIC1/2 Up.

C) ZIC ChIP-seq tracks at P7 and P60 at synaptic genes *Grm4* and *Grin2c* (y-axes for ChIP-seq are

autoscaled separately, according to data range in current view). D) (Left) Distribution of $\log_2$Fold

Change of H3K27ac enrichment as a function of H3K27me3 Down and H3K27me3 Up described

in Fig 2; and (Right) Metagene plots showing H3K27ac ChIP-seq signal at genes with H3K27me3

Up and H3K27me3 Down. E) (Left) Distribution of $\log_2$Fold Change of DNase Hypersensitivity as

a function of H3K27me3 Down and H3K27me3 Up described in Fig 2; and (Right) Metagene plots

showing DNAse-seq signal at genes with H3K27me3 Up and H3K27me3 Down. F) H3K27ac ChIP-

seq and DNAse-seq tracks at P7 and P60 at synaptic gene *Grin2c* (upper), and early gene *Atoh1*

(lower) (y-axes for ChIP-seq are autoscaled separately, according to data range in current view).

Fig 5. CGN Culture System refines temporal dynamics of H3K27me3 changes at gene promoters

A) Schematic describing CGN culture protocol over 7 days *in vitro* annotated with early and late

CGN genes. B) Differentially methylated H3K27me3 CUT&RUN peaks annotated by genomic

location, between DIV1-3, 3-5, 5-7 and 1-7 time points. C) Percentage of differential H3K27me3

CUT&RUN peaks between DIV1-3, DIV3-5, DIV5-7 and DIV1-7 annotated by genomic region

(Annotation performed using ChIPseeker package, TSS +/- 3000 bp). D) (Left) Heatmap of

hierarchical clustered VST-transformed DESeq2-normalized counts of differential H3K27me3

CUT&RUN promoter peaks from DIV1-7 (Right) Corresponding GO Terms and FDR associated with

nearest gene. E) (Left) Metagene plots for H3K27me3 CUT&RUN signal at 'Gain H3K27me3 at

DIV7' and 'Lose H3K27me3 at DIV7' clusters. (Right) CUT&RUN tracks for H3K27me3 for (top)

early genes *Myc*, and *Atoh1* and (bottom) late genes *Gabrd* and *Grin2c* at DIV1, 3, 5 and 7 (y-axes

for CUT&RUN tracks are autoscaled according to data range in current view). F) Violin plot

showing the distribution of Log2FC of gene expression measured by RNA-seq between DIV0 and

DIV7 CGNs as a function of clustering performed in E (one-way ANOVA, * indicates $p < 0.05$).

Fig 6. EZH2 catalytic activity temporally regulates CGN maturation in culture by depositing

H3K27me3 at early CGN genes

A) (Left) Western blot of acid extracted histones from CGNs treated for H3K27me3 and total

Histone H3 (n=3 biological replicates). CGNs were treated with GSK-J4 or GSK-126 at DIV1 and

harvested at DIV5, (Right) Quantification of Western Blot. B) MA plot describing differential

methylation due to GSK126 treatment in cultured CGNs (FDR < 0.05 and |L2FC| > 1) (n=3

biological replicates). C) Metagene plots for H3K27me3 peaks of DMSO and GSK-126 treated

CGNs at genes near (Left) All H3K27me3 promoter peaks, (Center) genes that gain H3K27me3 at

DIV7, and (Right) genes that lose H3K27me3 at DIV7, described in Fig 5E. D) (Upper) Volcano plot

for differentially expressed genes measured by RNA-seq between DMSO and GSK-126 treated

CGNs, (Lower) corresponding GO Terms and FDR for up- and downregulated genes. E) (Left)

Relationship between gene expression changes during CGN maturation in culture and due to

GSK-126 treatment. (Right) Violin plot showing the distribution of Log2FC of gene expression

between DMSO and GSK-126 treated CGNs as a function of clustering performed in Fig 5D (one-

way ANOVA, * indicates $p < 0.05$. F) (Upper) CUT and RUN tracks and (Lower) RNA-seq tracks for

early genes *Ebf3* and synaptic gene *Gabrd*, for DMSO and GSK126 treated CGNs.

Figure 1. The maturing mouse cerebellum undergoes a genome-wide redistribution of H3K27me3 dominated by loss at promoters

Figure 2. Differentially methylated promoters cluster by turnover kinetics and correlate with CGN maturation gene expression

A. ChIP-seq | H3K27me3+ Promoters

GO Terms	FDR
regulation of neuron projection arborization	2.48E-02
regulation of neural precursor cell proliferation	2.73E-02
neuron projection development	2.62E-02
neuronal signal transduction	2.52E-02
regulation of synapse maturation	9.76E-03
regulation of axon guidance	3.59E-02
gamma-aminobutyric acid transport	3.57E-02
calcium ion import	1.29E-03
calcium ion transmembrane transport	1.51E-02
GABAergic neuron differentiation	4.00E-03
calcium ion transport into cytosol	4.84E-02
calcineurin-NFAT signaling cascade	4.82E-02
postsynaptic density organization	2.23E-02

Figure 3. *Kdm6b* knockout in CGN precursors impairs CGN maturation via hypermethylation of CGN maturation gene promoters

Figure 4. H3K27me3 removal is followed by gain of H3K27me3 and chromatin accessibility at gene promoters

Figure 5. CGN Culture System refines temporal dynamics of H3K27me3 changes at gene promoters

Figure 6. EZH2 catalytic activity temporally regulates CGN maturation in culture by depositing H3K27me3 at early CGN genes