Co-option of ancestral signal elements in the evolution of a
cooperative parental behavior

Jeanette B. Moss[1], James P. Tumulty[2], and Eva K. Fischer[1*]

[1] Department of Evolution, Ecology, and Behavior, University of Illinois, Urbana-Champaign,
Urbana, IL 61801, U.S.A.

[2] Department of Biology, College of William and Mary, Williamsburg, VA, 23185, U.S.A.

*Eva K. Fischer

Email: efischer@illinois.edu

Author Contributions: J.B.M., J.P.T., and E.K.F. conceived of and designed the study. J.B.M.
collected the data. J.B.M. and J.P.T. performed the analyses. J.B.M., J.P.T., and E.K.F. wrote the
paper.

Keywords: Animal communication, anuran, parental care, cooperation, social context

Abstract

The emergence of complex social interactions is predicted to be an important selective force in the

diversification of communication systems. Parental care presents a key social context in which to

study the evolution of novel signals, as care often requires communication and behavioral

coordination between parents and is an evolutionary stepping-stone towards increasingly complex

social systems. Anuran amphibians (frogs and toads) are a classic model of acoustic

communication and the vocal repertoires of many species have been characterized in the contexts

of advertisement, courtship, and aggression; yet quantitative descriptions of calls elicited in the

context of parental care are lacking. The biparental poison frog, *Ranitomeya imitator*, exhibits a

remarkable parenting behavior in which females, cued by the calls of their male partners, feed

tadpoles unfertilized eggs. Here, we characterized and compared calls across three social contexts,

for the first time including a parental care context. We found that egg feeding calls share some

properties with both advertisement and courtship calls but also had unique properties. Multivariate

analysis revealed high classification success for advertisement and courtship calls but misclassified

nearly half of egg feeding calls as either advertisement or courtship calls, suggesting additional

signal modalities play a role in parental communication. Egg feeding and courtship calls both

contained less identity information than advertisement calls, as expected for signals used in close-

range communication where uncertainty about identity is low. Taken together, egg feeding calls

likely borrowed and recombined elements of both ancestral call types to solicit a novel, context-

dependent parenting response.

Significance Statement

Parental care has evolved independently in every major animal lineage and represents a major

step in the evolution of complex sociality. Communication systems may need to increase in

complexity. To explore these ideas, we characterized calls associated with trophic egg feeding, a

unique cooperative parental behavior in the biparental mimic poison frog and compared them to

calls associated with mate attraction (advertisement and courtship calls). Our analysis revealed

some distinct, but many shared properties of signals elicited during egg feeding, suggesting that

signals deployed in a novel social context evolve via modification and recombination of existing

signals. These findings deepen our understanding of the relationship between complexity of social

and communication systems.

Main Text

Introduction

Complexity of communication systems is predicted to co-evolve with complexity of social systems (1, 2). Transitions in social organization introduce novel contexts for interaction between individuals, which in turn require more specialized communication systems. While traditionally examined through the lens of group size (3–5), there is increasing interest in understanding how these co-evolutionary dynamics play out across multiple indices of social complexity (6–8). Parental care is a major component of a species' social system (9) and represents a key step in the evolution of complex sociality (10). When care responsibilities are shared between multiple caregivers, ensuring offspring survival requires coordinated behaviors that rely on effective communication. One solution is to expand the size of the signal repertoire by adding novel, context-specific communication features (7, 11), as has been shown in the case of avian cooperative breeding (8). Alternatively, existing signals may be co-opted to function in the new context via context-dependent regulation of responses (12, 13). Quantitative descriptions of variation in signal components provide an important first step for generating hypotheses about signal function and evolution.

One of the most influential models for studying animal communication is anuran amphibian vocal communication and auditory processing. Across anurans, vocal repertoires of differing sizes and compositions have been meticulously characterized along with the diversity of behavioral and neural responses they elicit reviewed in 14–16). This rich literature illustrates how greater complexity of communication systems may evolve through the specialization of signal components for a given behavioral context, evolution of receiver bias, or both. In a classic example, the common coquí frog (*Eleutherodactylus coqui*), advertisement call is composed of two syllables – 'co' and 'qui' – which are differentially specialized for aggressive versus mate attraction contexts (17). In this case, regulation of responses is further facilitated by the evolution of sex-specific auditory sensitivities (17). Conversely, modulation of the túngara frog's (*Physalaemus pustulosus*) characteristic 'whine-chuck' call has similar motivating effects on potential mates and rivals, but with clearly distinct social and behavioral outcomes (18).

Beyond serving as models for the study of animal communication, anurans also exhibit remarkable diversity of parental care (19). The mimic poison frog, *Ranitomeya imitator*, is a species characterized by monogamy, pair bonding, and biparental care of eggs and tadpoles (20). Terrestrial eggs are cared for by both parents and tadpoles are transported "piggy-back" to pools of water upon hatching. Remarkably, parents visit and care for their developing tadpoles until metamorphosis. Specifically, females feed tadpoles with unfertilized 'trophic' eggs, a fascinating behavior coordinated by males calling to females and leading them to hungry, begging tadpoles . Egg feeding has evolved multiple times in association with biparental care (20) and functions

crucially in offspring development (21), suggesting that selection on signalers and receivers during

these interactions is strong. It has been hypothesized that egg feeding behaviors are evolutionarily

derived from courtship behaviors (21, 23, 24); however, the signals that coordinate and elicit egg

feeding have yet to be quantitatively described for any species, including *R. imitator*.

Here, we characterize and compare calls of *R. imitator* across three social contexts to test

whether the emergence of coordinated parental care within this lineage has promoted the

incorporation of novel signals for communication. Given that several call types in frogs are known

to serve dual functions, we predicted that egg feeding calls were likely co-opted and modified from

calls adapted for the more ancestral functions of advertisement and courtship (described in 29 and

30).

Results

Calls vary across social contexts

To quantify and compare advertisement, courtship, and egg feeding signals, we recorded

representative calls of each type for captive male *Ranitomeya imitator* across the breeding cycle.

Briefly, advertisement and courtship calls – both deployed for attracting females but over different

distances (i.e., long vs. short range, respectively; 6, 7, 33) were distinguished based on published

descriptions of *R. imitator* vocalizations (25, 26), whereas egg feeding calls were identified using

video surveillance of parenting frogs (e.g., Movie S1). Visual inspection of waveforms and

spectrograms of putative call types revealed distinct characteristics (Fig. 1).

Because signals are composed of multiple components, which may differ in salience to

different receivers and be under different forms of selection , we measured nine acoustic properties

to quantify call variation (Table 1; Fig. S1). All properties except for inter-call interval and pulse rate

showed significant variation depending on the social context in which the call was measured

(ANOVA $p < 0.05$). In terms of temporal properties, courtship calls were significantly shorter than

advertisement (t = 6.59, $p < 0.001$) or egg feeding calls (t = 4.56, $p < 0.001$) and contained fewer

pulses (advertisement-courtship: t = 7.24, $p < 0.001$; courtship-egg feeding: t = 4.98, $p < 0.001$).

Pulse durations were longest in advertisement calls (advertisement-courtship: t = 3.06, $p = 0.003$;

advertisement-egg feeding: t = 4.65, $p < 0.001$) and shortest in egg feeding calls (egg feeding-

courtship: t = 2.52, $p = 0.013$; Fig. 2A), while pulse intervals were shortest in advertisement calls

(advertisement-egg feeding: t = 4.22, $p < 0.001$; advertisement-courtship: t = 2.72, $p = 0.007$;) and

longest in egg feeding calls (courtship-egg feeding: t = 2.33, $p = 0.021$) (Fig. 2B). In terms of

spectral properties, egg feeding calls had significantly lower dominant frequencies than either

advertisement (t = 3.67, $p < 0.0001$) or courtship calls (t = 2.88, $p = 0.004$; Fig. 2C), and

advertisement calls had significantly narrower call bandwidths than courtship (t = 6.93, *p* < 0.001)

or egg feeding calls (t = 5.77, *p* < 0.001; Fig. 2D).

Individual identity information varies across call types

We next examined the degree to which acoustic properties of each call type convey

information about signaler identity. We found that putative call types and acoustic properties within

those call types varied in their individual distinctiveness (Table 2). A key criterion for conveying

information about signaler identity is that signal properties exhibit high levels of variation among

individuals (CV_a) but low levels of variation within individuals (CV_w; 36). For calls classified as

advertisement calls, most acoustic properties showed greater variation among individuals than

within individuals (CV_a / CV_w, or potential for individual coding (PIC) > 1). The greatest individual

distinctiveness was found in spectral properties, in particular dominant frequency (PIC = 5.50).

Courtship and egg feeding calls showed less individual distinctiveness overall, with most acoustic

properties showing as much or more variation within individuals as among individuals, and

dominant frequency being a less reliable indicator of identity (courtship PIC = 0.63; egg feeding PIC

= 0.82). When considering all acoustic properties together, courtship calls contained the least

identity information (H_s = 1.48) of the three types (advertisement H_s = 3.82; egg feeding calls H_s =

2.81).

High call type classification success

To describe variation among call types at the population level, we performed a principal

components analysis (PCA). This returned three primary axes of variation (eigenvalues > 1) which

together explained 67.8% of variation in acoustic properties (Table S2). Pulse duration, pulse

interval, dominant frequency, and frequency of the 2nd harmonic all loaded heavily on the first

principal component (PC1; 37.8%), while call duration and number of pulses loaded heavily on the

second principal component (PC2; 18.2%) (Table S2). The third principal component (PC3; 11.8%)

contributed only subtly to variation among call types, with variation explained by call interval and

pulse duration. Overlap between call types was evident across all individuals assayed (Fig. S2–3),

suggesting that population-level patterns were not obscured by variation at the individual level.

To evaluate the predictive capacity of the PCs in assigning calls to their predefined social

contexts, we next performed a discriminant function analysis (DFA) using PCs 1–3 as inputs. This

approach correctly classified 72.2% of calls to their assigned social context. The chance-corrected

classification success was significant (Weighted Cohen's kappa = 0.566, *p* < 0.001). The strongest

coefficients of the first linear discriminant were held by PC1 (the 'frequency and pulse period'

factor), whereas PC2 (the 'call length' factor) contributed more to the second linear discriminant.

The highest accuracy of classification according to this model was for advertisement calls (82.9%)

followed by courtship calls (73.5%) and egg feeding calls (57.1%). Courtship calls were equally

likely to be misclassified as advertisement (13.2%) or egg feeding calls (13.2%), and egg feeding

calls were equally likely to be misclassified as advertisement (21.4%) or courtship calls (21.4%).

While overall misclassification rates did not vary significantly across males (27.7 $\pm$ 9.31%; χ^2 = 21,

p = 0.279), over half of all misclassified egg feeding calls (16/24) were attributed to three individuals:

Ri.0181, Ri.0187, and Ri.0442 (Fig. S3). Given that egg feeding calls were sampled at the same

grain (10 calls from a single bout of egg feeding) for each male, variation in parents or specific

parenting conditions may explain variation in call distinctiveness.

Discussion

Increased complexity of communication systems is predicted to co-evolve with the

emergence of new forms of social interaction (1, 2, 6, 7), but this relationship has yet to be explored

in the context of signals specialized for parental care. In this study, we characterized and compared

calls of the biparental poison frog, *Ranitomeya imitator*, across social contexts in which calls either

serve an ancestral communication function – advertisement or courtship – or a derived social

function – parental care via coordinated, trophic egg feeding. We found that vocalizations produced

in distinct social contexts differ in both acoustic properties and the amount of identity information

contained in those properties. Crucially, calls elicited during egg feeding possessed unique spectral

and temporal properties that were not observed in advertisement or courtship contexts and retained

moderate identity information, suggesting that egg feeding calls were likely co-opted from existing

elements but have since undergone evolutionary modification in form. Despite this, nearly half of

all egg feeding calls were not sufficiently distinct to be distinguished from ancestral call types based

on acoustic properties alone, implicating a role for multimodal communication in the coordination

of parental behavior.

Studies of anuran acoustic communication recognize multiple common types of calls that

function in distinct social contexts (15). Advertisement calls are by far the most studied call type

because they serve multiple social functions (i.e., mate attraction, territory defense, and species

delimitation) and are therefore taxonomically widespread (15). Virtually no attention has been paid

to characterizing vocalizations deployed during parental care, in which communication between the

sexes holds a special significance beyond simple mate attraction. We quantified nine acoustic

properties of vocalizations elicited during egg feeding in *R. imitator* and compared these to the

same properties quantified for advertisement and courtship calls. We found that egg feeding calls

shared some properties with advertisement calls (i.e., longer than courtship calls with more pulses)

and some with courtship calls (i.e., broader call bandwidths), supporting the prediction that egg-

feeding calls evolved via co-option of elements from ancestral call types. This is not surprising, as

male advertisement and courtship calls already function to elicit female responses. However, we
also identified properties of egg feeding calls that were distinct from either ancestral call type (i.e.,
low dominant frequencies and especially short pulse durations coupled with long pulse intervals).
Some of these differences in acoustic properties could be physiologically related to egg feeding
calls being produced at lower amplitudes (i.e., "quieter") because pairs are typically very close in
space (<0.5 m) when males are leading females to tadpoles and stimulating them to feed (Movie
S1). However, courtship calls are also "quiet" close-range signals, so the unique elements of egg
feeding calls cannot be entirely explained as physiological by-products of low amplitude calls.
Modification of temporal patterning is a common theme in the evolution of context-dependent
signals (e.g., 22, 31) and we suggest the subtle differences observed in pulse durations and
intervals reflect novel elements that are selectively maintained for the coordination of egg feeding.
A second major finding of our study was that the identity information conveyed in *R. imitator*
vocalizations varies according to the social context in which they are deployed. By measuring calls
repeatedly and across multiple social behaviors in the same males, we showed that advertisement
calls contain the most individual identity information overall, with dominant frequency being the
most reliable signature of male caller. Dominant frequency is relatively 'static' within males of many
species (e.g., 11, 32) presumably because it is constrained by the morphology of the sound
producing structures (35) and thus correlated with body size. Advertisement calls may be a signal
or cue of individual identity used by rival males to recognize territory neighbors. In the case of mimic
poison frogs, which are monogamous and pair bonding, individually distinctive advertisement calls
could also be used by females to recognize their mate. If males benefit from being recognized by
either territory neighbors or mates, selection could favor greater identity information in these calls
to facilitate recognition. A complementary explanation is that courtship and egg feeding calls are
typically produced over shorter distances than advertisement calls, such that other signal modalities
(e.g., visual and/or chemosensory cues) may interact with auditory cues in mediating individual
recognition. Indeed, the majority of acoustic properties examined were more dynamic (i.e., showed
more within-individual variation) in courtship and egg feeding calls than in advertisement calls,
which could reflect stronger selection for extreme values to enhance the motivational effects of
these signals (33).
Finally, we evaluated whether subtle variation in the spectral range, temporal patterning,
and individual distinctiveness was sufficient to quantitatively distinguish egg feeding calls from
ancestral call types. In a discriminant function analysis, all three call types were classified above
chance level (33%) from their acoustic properties alone; however, overlap between call types was
considerable (Fig. 3A) and calls elicited during egg feeding were misclassified in nearly half of all
instances. Thus, egg feeding calls may not be distinct enough to afford their own category within
the vocal repertoire of *R. imitator* despite eliciting a novel, complex behavior. An intriguing

observation can be made, however, in the patterns of misclassification. Egg feeding calls in our
sample were equally likely to be misclassified as advertisement or courtship calls, suggesting that
egg feeding calls borrow and recombine elements from multiple call types. Perhaps this complexity
arises from the unique function of egg feeding calls, which may align selection on the response
with both advertisement (i.e., must attract females to a specific location) and courtship calls (i.e.,
must motivate females to lay eggs). Additionally, females deciding to lay trophic eggs (which are
typically laid in the water) versus eggs that will be fertilized (which are typically attached to a leaf
above the water line or in a location away from a pool) are likely responding to their own
physiological state and other contextual cues and signals as well as male signals. For one, tadpoles
of this species vibrate vigorously against parental frogs to signal their hunger, which appears to be
important in eliciting trophic eggs (22). Use of multimodal cues would also be consistent with
acoustic studies of other frogs, which have shown that behavioral responsiveness to male calls is
enhanced when females are also presented with visual (36) or contextual cues (37). Future studies
should focus on female perception of male call variants in relation to female state and other
contextual cues to better understand how selection on signalers and receivers could synergistically
drive the co-option and modification of acoustic signals for trophic egg feeding.

In conclusion, our study examined acoustic signals produced in a parental coordination
context in poison frogs and contrasted these against ancestral signal types adapted to function in
other facets of male-female communication. We found support for the prediction that extended
functionality of the communication system in association with increased social complexity likely
arose through co-option of pre-existing signals followed by evolutionary modification resulting in
novel elements. Our results highlight the complexity of anuran communication systems and the
need to characterize vocal repertoires across a greater diversity of social contexts. More generally,
this study contributes to understanding how communication systems are evolutionarily fine-tuned
to convey context-dependent information in increasingly complex social systems.

Materials and Methods

Animal Husbandry

A captive colony of *Ranitomeya imitator* descended from five localities in Peru (Veradero,
Tarapoto, Huallaga, Sauce, Southern, and hybrids between these; Table S1) was maintained in
temperature- (71.86 ± 2.79°C) and light- (12:12 h cycle) controlled rooms at the University of Illinois
Urbana-Champaign (UIUC). Breeding pairs were housed in 12x12x18 inch glass terraria (©Exo
Terra, Mansfield, MA, U.S.A) containing live plants, driftwood, sphagnum moss, leaf litter, and
water-filled film cannisters for egg laying and tadpole deposition. Tank humidity (84.95 ± 3.08%)

was maintained using a misting system (©Mist King, Ontario, Canada). Frogs were fed wingless
Drosophila fruit flies dusted with vitamin supplements three times weekly. Only sexually mature
individuals were used as focal subjects. All procedures were approved by the UIUC Animal Care
and Use committee (Protocol #20147).
262
Acoustic Recordings
Between July 2021 and May 2022, seven male *R. imitator* were recorded in their home
terraria across the breeding cycle. A shotgun microphone (K6/ME66, ©Sennheiser, Wennebostel,
Germany) was positioned above the screen covering the focal tank and connected to a handheld
recorder (H4n Pro, Zoom, Tokyo, Japan); 44.1kHZ sampling rate, 16-bit resolution. For each
recording, we documented the time of day and reproductive stage of the focal male by the following
categories: (1) nonbreeding (pair did not currently have eggs or tadpoles); (2) brooding (frog had
at least one egg but hatching had not yet occurred); or (3) tadpole (at least one tadpole was present
in a deposition pool). Across all three stages, advertisement calls were distinguished from shorter
courtship calls based on a published characterization of advertisement calls for this species (26).
While advertisement calls produce a characteristic 'trill' sound and function in long-range mate
attraction and territory defense, courtship calls are typically shorter and softer and function in close-
range interactions with females (25, 26). To isolate vocalizations elicited in the unique social context
of egg feeding and test whether these can be quantitatively distinguished from advertisement and
courtship calls, we took video recordings (RLC-510A, ©Reolink, New Castle, DE, U.S.A.)
concurrently with audio recordings during the tadpole stage. Male *R. imitator* stimulate female
partners to lay trophic eggs by leading them to tadpole deposition pools and maintaining close
and/or tactile contact while calling continually (21). Thus, egg feeding calls were identified as calls
coinciding with males moving towards tadpole deposition pools, leading females to pools, sitting
on or near the pools, and culminating with the arrival of females at pools (e.g., Movie S1). Calls
recorded during the tadpole stage while males were away from pools and/or oriented towards egg
laying cannisters were identified as either advertisement or courtship calls, based on the same
criteria used during nonbreeding and brooding stages.
286
Call Filtering and Analysis
To examine variation within and among the three call types, ten calls of each type for each
individual were selected from available recordings (Dataset S1). Very low-amplitude or low-quality
calls, or calls that could not be clearly classified between the three types on the basis of social
context or previous published accounts were excluded. Raw recordings of calls were trimmed in
Audacity v. 3.0.2 (Audacity Team, 2021). Analysis of calls was performed in Raven Pro v. 1.6.1
(Cornell Lab of Ornithology, 2022). In total, we measured nine acoustic properties of each call. We

measured six temporal properties using the entire call or part of the call (i.e., a single pulse or inter-
pulse interval) as the unit of measurement in the waveform view in Raven. These included call
duration (time in seconds from the first pulse to the last note; Fig. S1A), inter-call interval (time in
seconds to the next call within the same bout, with bout defined as calls occurring within one minute
of each other; Fig. S1A), number of pulses, pulse duration (time in seconds from the beginning of
a note to the end; Fig. S1B), pulse interval (time in seconds from the end of one note to the start of
the next note), and pulse rate (the reciprocal of pulse period, or the full time from the beginning of
one note to the beginning of next consecutive note; Fig. S1B). Pulse duration, pulse interval, and
pulse rate were calculated by taking the average of three notes within the call, excluding
introductory notes. Three spectral properties were also quantified using the entire call as the unit
of measurement, including the dominant frequency of the call (in Hz; Fig. S1C) and its bandwidth
(90%; Fig. S1C). In most cases the dominant frequency corresponded to the second harmonic (5-
6 kHz) but occasionally the fundamental frequency (<3 kHz) was dominant and in a few cases, a
broad frequency peak in the range of 3-4 kHz was observed and the harmonic structure was unclear
(Fig. 1; Fig. S1). Thus, the frequency of the second harmonic was also quantified for each call.
309
Statistical Analyses
All statistical analyses were performed in R Studio v. 1.1.318 (R Core Development Team,
2020). We first summarized each of nine acoustic properties by their assigned call type. We tested
for differences between call types using Type III ANOVAs followed by post-hoc testing. Model
testing was carried out using the packages 'lmerTest' (38) and 'emmeans' (39), with reproductive
stage specified as a covariate and individual ID specified as a random effect in all models. For
metrics that were collected in multiples per call (i.e., pulse duration, pulse interval, and pulse rate),
we also specified call ID nested within individual ID as a random effect. Some variables (inter-call
interval, pulse interval, pulse rate, and all spectral properties) were log-transformed prior to analysis
to better meet assumptions of normality and homogeneity of variance, which we evaluated with
analyses of random effects and residuals.
We next evaluated within- and among-individual variation for each call type for each of the
nine acoustic properties. For each male, we calculated a within-individual coefficient of variation
(CV_w = individual SD/individual mean × 100) and report the mean CV_w across males. We computed
the among-individual coefficient of variation (CV_a = grand SD/grand mean × 100) and report the
ratio of among-individual variability (CV_a) to within-individual variability (CV_w) as the potential for
individual coding (PIC; 43). Values of PIC greater than one indicate that a call property varies more
among individuals than within individuals. To further describe individual distinctiveness of calls, we
calculated Beecher's Information Statistic (H_s; 44), which expresses the overall amount of individual
identity information contained in calls. To control redundancy due to correlations among

measurements, data were transformed into uncorrelated variables using principal components
analysis (PCA) and all principal components were retained to calculate H_s (30). PCAs were
implemented with the 'prcomp' function in R after transforming pooled variables (pulse duration,
pulse interval, and pulse rate) into averages to ensure equality of sample sizes and scaling all input
variables.

335   Finally, to describe variation among call types at the population-level and evaluate the
success rate of classifying calls to the correct call type (advertisement, courtship, or egg feeding),
we took a two-step multivariate approach. First, we performed PCA as described above. Input
variables were prepared by converting acoustic properties into z-scores scaled to the mean of the
individual, which controlled for variation among males. Second, principal components with
eigenvalues > 1 were extracted as inputs into a discriminant function analysis (DFA). We
implemented this test using the linear discriminant analysis ('lda' function) in the package 'MASS'
(41). Chance-corrected classification success (Cohen's kappa) was calculated following the
approach of (42) in the R package 'vcd' (43). Differences in classification success among males
were statistically evaluated using a Chi-Squared test of independence.

Acknowledgments

348   We thank Mark Bee for helpful advice on bioacoustics equipment and approaches in
early project design stages. We thank the Fischer Lab for constructive feedback on earlier drafts
of our manuscript. This study was supported by a National Science Foundation Postdoctoral
Research Fellowship in Biology to J.B.M. (#2010649), at University of Illinois Research Board
Grant to E.K.F. (RB21025), and University of Illinois Urbana-Champaign laboratory start-up funds
to E.K.F.

References

1. T. M. Freeberg, R. I. M. Dunbar, T. J. Ord, Social complexity as a proximate and ultimate
factor in communicative complexity. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 1785–1801
(2012).
2. K. B. Sewall, Social complexity as a driver of communication and cognition. *Integr. Comp.*
Biol. **55**, 384–395 (2015).
3. T. Freeberg, Social complexity can drive vocal complexity: Group size and information in
chickadee calls in Carolina chickadees. *J. Acoust. Soc. Am.* **119**, 3222–3222 (2006).
4. H. Bouchet, C. Blois-Heulin, A. Lemasson, Social complexity parallels vocal complexity: a
comparison of three non-human primate species. *Front. Psychol.* **4** (2013).

5. M. B. Manser, *et al.*, "Vocal complexity in meerkats and other mongoose species" in
Advances in the Study of Behavior, (Elsevier, 2014), pp. 281–310.

6. L. Peckre, P. M. Kappeler, C. Fichtel, Clarifying and expanding the social complexity
hypothesis for communicative complexity. *Behav. Ecol. Sociobiol.* **73**, 11 (2019).

7. T. M. Freeberg, I. Krams, Does social complexity link vocal complexity and cooperation?
J. Ornithol. **156**, 125–132 (2015).

8. G. M. Leighton, T. Birmingham, Multiple factors affect the evolution of repertoire size
across birds. *Behav. Ecol.* **32**, 380–385 (2021).

9. P. M. Kappeler, T. Clutton-Brock, S. Shultz, D. Lukas, Social complexity: patterns,
processes, and evolution. *Behav. Ecol. Sociobiol.* **73**, 5 (2019).

10. L. Socias-Martínez, P. M. Kappeler, Catalyzing transitions to sociality: Ecology builds on
parental care. *Front. Ecol. Evol.* **7**, 1–17 (2019).

11. E. D. Broder, *et al.*, Evolutionary novelty in communication between the sexes. *Biol. Lett.*
16, 20200733 (2021).

12. M. J. West-Eberhard, *Developmental plasticity and evolution* (Oxford University Press,
2003).

13. R. L. Rodríguez, Trait duplication by means of sensory bias. *Behav. Ecol.* **20**, 1376–1381
(2009).

14. K. D. Wells, *The Ecology and Behavior of Amphibians* (University of Chicago Press,
2010).

15. L. F. Toledo, *et al.*, The anuran calling repertoire in the light of social context. *Acta
Ethologica* **18**, 87–99 (2015).

16. M. A. Bee, M. S. Reichert, J. Tumulty, "Assessment and recognition of rivals in anuran
contests" in *Advances in the Study of Behavior*, (Elsevier, 2016), pp. 161–249.

17. P. M. Narins, R. R. Capranica, Sexual differences in the auditory system of the tree frog
Eleutherodactylus coqui. *Science* **192**, 378–380 (1976).

18. X. E. Bernal, K. L. Akre, A. T. Baugh, A. S. Rand, M. J. Ryan, Female and male
behavioral response to advertisement calls of graded complexity in túngara frogs,
Physalaemus pustulosus. *Behav. Ecol. Sociobiol.* **63**, 1269–1279 (2009).

19. A. I. Furness, I. Capellini, The evolution of parental care diversity in amphibians. *Nat.
Commun.* **10**, 4709 (2019).

20. J. L. Brown, V. Morales, K. Summers, A key ecological trait drove the evolution of
biparental care and monogamy in an amphibian. *Am. Nat.* **175**, 436–446 (2010).

21. J. Tumulty, V. Morales, K. Summers, The biparental care hypothesis for the evolution of
monogamy: Experimental evidence in an amphibian. *Behav. Ecol.* **25**, 262–270 (2014).

22. M. Yoshioka, C. Meeks, K. Summers, Evidence for begging as an honest signal of
offspring need in the biparental mimic poison frog. *Anim. Behav.* **113**, 1–11 (2016).
23. J. P. Caldwell, V. R. L. de Oliveira, V. R. L. de Oliveira, Determinants of biparental care in
the spotted poison frog, *Dendrobates vanzolinii* (Anura: Dendrobatidae). *Copeia* **1999**,
565 (1999).
24. E. H. Poelman, M. Dicke, Offering offspring as food to cannibals: oviposition strategies of
Amazonian poison frogs (*Dendrobates ventrimaculatus*). *Evol. Ecol.* **21**, 215–227 (2007).
25. M. Mayer, L. M. Schulte, E. Twomey, S. Lötters, Do male poison frogs respond to
modified calls of a Müllerian mimic? *Anim. Behav.* **89**, 45–51 (2014).
26. E. Twomey, M. Mayer, K. Summers, Intraspecific call variation in the mimic poison frog
Ranitomeya imitator. *Herpetologica* **71**, 252–259 (2015).
27. A. P. Moller, A. Pomiankowski, Why have birds got multiple sexual ornaments? *Behav.*
Ecol. Sociobiol. **32**, 167–176 (1993).
28. E. A. Hebets, D. R. Papaj, Complex signal function: developing a framework of testable
hypotheses. *Behav. Ecol. Sociobiol.* **57**, 197–214 (2005).
29. J. C. Tanner, J. L. Ward, R. G. Shaw, M. A. Bee, Multivariate phenotypic selection on a
complex sexual signal. *Evolution* **71**, 1742–1754 (2017).
30. M. D. Beecher, Signaling systems for individual recognition: an information theory
approach. *Anim. Behav.* **38**, 248–261 (1989).
31. A. S. Bhat, V. A. Sane, K. S. Seshadri, A. Krishnan, Behavioural context shapes vocal
sequences in two anuran species with different repertoire sizes. *Anim. Behav.* **184**, 111–
129 (2022).
32. T. Q. Gentner, K. M. Fenn, D. Margoliash, H. C. Nusbaum, Recursive syntactic pattern
learning by songbirds. *Nature* **440**, 1204–1207 (2006).
33. H. C. Gerhardt, Female mate choice in treefrogs: static and dynamic acoustic criteria.
Anim. Behav. **42**, 615–635 (1991).
34. M. A. Bee, H. C. Gerhardt, Habituation as a mechanism of reduced aggression between
neighboring territorial male bullfrogs (*Rana catesbeiana*). *J. Comp. Psychol.* **115**, 68–82
(2001).
35. W. F. Martin, Mechanics of sound production in toads of the genus *Bufo*: Passive
elements. *J. Exp. Zool.* **176**, 273–293 (1971).
36. R. C. Taylor, B. A. Klein, J. Stein, M. J. Ryan, Faux frogs: multimodal signalling and the
value of robotics in animal behaviour. *Anim. Behav.* **76**, 1089–1097 (2008).
37. C. Martin, L. Guy, R. C. Taylor, Male position relative to foam nests influences female
mate choice in the Túngara frog, *Physalaemus pustulosus*. *J. Herpetol.* **45**, 178–180
(2011).

38. A. Kuznetsova, P. B. Brockhoff, R. H. B. Christensen, lmerTest Package: Tests in Linear
Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26 (2017).

39. R. V. Lenth, *et al.*, Package "emmeans": Estimated marginal means, aka least-squares
means. *R Package Version 340* (2022) https:/doi.org/10.1080/00031305.1980.10483031
(June 17, 2022).

40. P. Linhart, *et al.*, Measuring individual identity information in animal signals: Overview
and performance of available identity metrics. *Methods Ecol. Evol.* **10**, 1558–1570
(2019).

41. W. N. Venables, B. D. Ripley, *Modern Applied Statistics with S-PLUS* (Springer, 2002)
(June 17, 2022).

42. K. Titus, J. A. Mosher, B. K. Williams, Chance-corrected classification for use in
discriminant analysis: ecological applications. *Am. Midl. Nat.* **111**, 1–7 (1984).

43. M. Friendly, Working with categorical data with R and the vcd and vcdExtra packages
(2016) (July 27, 2022).

Figures and Tables

Figure 1. Vocalizations have unique spectral profiles across social contexts. Spectrograms and associated power spectra of a representative (A) advertisement call; (B) courtship call; and (C) egg feeding call of *Ranitomeya imitator* (photo, top). Calls were visualized in Raven Pro v. 1.6.1 (Cornell Lab of Ornithology, 2022) using the Hann window, 256 pt frequency resolution. The courtship and egg feeding call power spectra show the two forms of increased frequency bandwidth: broader dominant frequency peak (B) or a fundamental frequency that is at a higher relative amplitude (C). Photo by Anton Sorokin, antonsrkn@gmail.com.

461
462
Figure 2. Acoustic properties of *Ranitomeya imitator* vocalizations vary between advertisement,
courtship, and egg feeding calls. Plots depict distributions of four acoustic properties by call type:
(A) Pulse duration; (B) log10(Pulse interval); (C) log10(Dominant frequency); and (D)
log10*Bandwidth 90%). Color of dots corresponds to seven unique males in the sample.
Significance levels are shown for Tukey-HSD post-hoc contrasts. Pairwise contrasts without
significant differences are not shown.

469
470 **Figure 3.** Classification success of vocalizations based on acoustic properties exceeds chance
471 levels, despite considerable overlap. Plot of *Ranitomeya imitator* calls in two-dimensional space
472 defined by the first two discriminant functions of a linear discriminant function analysis (DFA).
473 Calls are clustered by call type (advertisement, courtship, and egg feeding) with centroids (large
474 points) and 95% confidence intervals (ellipses) depicted for each type.

Table 1: Summary of acoustic properties of 70 advertisement calls, 70 courtship calls, and 60 egg feeding calls across seven males. Pulse duration, interval, and rate were measured three times for each call, and summary statistics are thus calculated based on 210 or 180 measurements, respectively. Results of pairwise contrasts between call types are also reported. Tukey test statistics (t) and adjusted *p*-values are reported for each contrast and significant test-statistics are bolded.

Acoustic Property	Advertisement	Courtship	Egg feeding	Advertisement-Courtship		Advertisement-Egg feeding		Courtship-Egg feeding	
	Mean (± SD)	Mean (± SD)	Mean (± SD)	t	*p*	t	*p*	t	*p*
Temporal Properties									
Call Duration (s)	1.14 ± 0.23 (0.66–1.73)	0.75 ± 0.29 (0.14–1.92)	1.04 ± 0.35 (0.48–1.97)	**6.589**	**<0.001**	1.072	0.285	**-4.557**	**<0.001**
Call Interval (s)	10.53 ± 7.93 (2.53–40.89)	8.15 ± 5.42 (0.41–34.96)	7.40 ± 5.27 (1.46–28.94)	1.568	0.119	1.424	0.156	0.198	0.843
No. of Pulses	31.33 ± 6.02 (18–46)	20.93 ± 7.28 (5–39)	27.08 ± 9.22 (7–48)	**7.235**	**<0.001**	1.164	0.246	**-4.979**	**<0.001**
Pulse Duration (s)	0.019 ± 0.005 (0.005–0.031)	0.016 ± 0.005 (0.006–0.035)	0.014 ± 0.005 (0.005–0.032)	**3.064**	**0.003**	**4.645**	**<0.001**	**2.519**	**0.013**
Pulse Interval (s)	0.016 ± 0.005 (0.007–0.030)	0.02 ± 0.01 (0.006–0.120)	0.027 ± 0.023 (0.009–0.247)	**-2.722**	**0.007**	**-4.222**	**<0.001**	**-2.330**	**0.021**
Pulse Rate (notes/s)	28.55 ± 2.16 (19.8–37.5)	28.16 ± 3.67 (7.6–37.9)	26.73 ± 5.02 (3.9–38.8)	0.990	0.324	1.907	0.058	1.267	0.207
Spectral Properties									
Dominant Frequency (Hz)	5249.1 ± 496.2 (4048.2–5857)	5040.3 ± 613 (2842.4–5857)	4783.3 ± 779.1 (2625–6201.6)	1.403	0.162	**3.676**	**<0.001**	**2.879**	**0.004**
Frequency of 2nd Harmonic (Hz)	5220.8 ± 601.1 (4048.2–5857)	5122.1 ± 495.2 (3789.9–5857)	4954.8 ± 542.1 (3617.6–6201.6)	1.520	0.130	**3.946**	**<0.001**	**3.080**	**0.002**
Bandwidth 90% (Hz)	678.8 ± 675.8 (258.4–2928.5)	1544.1 ± 1193.4 (281.3–4823.4)	1662.7 ± 1419.7 (281.3–5426.4)	**-6.932**	**<0.001**	**-5.767**	**<0.001**	-0.413	0.680

Table 2: Individual distinctiveness of nine acoustic properties of 70 advertisement calls, 68 courtship calls, and 56 egg feeding calls across seven males. Potential for individual coding (PIC) is calculated as the ratio of the variation among individuals to the mean variation within individuals. Values greater than one indicate that a call property varies more among individuals than within individuals (shown in bold).

Acoustic Property	Advertisement			Courtship			Egg feeding		
	CV_a	Mean CV_w	PIC	CV_a	Mean CV_w	PIC	CV_a	Mean CV_w	PIC
Temporal Properties									
Call Duration (s)	11.76	16.74	0.70	13.06	34.36	0.38	20.67	26.24	0.79
Interval to next call (s)	29.56	65.90	0.45	36.30	50.91	0.71	44.70	57.25	0.78
No. of Pulses	**15.39**	**13.43**	**1.15**	18.77	30.32	0.62	**26.43**	**23.62**	**1.12**
Pulse Duration (s)	**20.29**	**19.48**	**1.04**	18.88	26.50	0.71	20.22	27.81	0.73
Pulse Interval (s)	22.21	23.45	0.95	8.43	38.56	0.22	**40.16**	**35.16**	**1.14**
Pulse Rate (notes/s)	5.88	7.59	0.77	5.84	11.02	0.53	9.34	13.95	0.67
Spectral Properties									
Dominant Frequency (Hz)	**9.94**	**1.81**	**5.50**	6.21	10.09	0.63	9.94	12.10	0.82
Frequency of 2nd Harmonic (Hz)	**11.35**	**3.81**	**2.98**	**7.37**	**6.49**	**1.14**	**9.6**	**6.58**	**1.46**
Bandwidth 90% (Hz)	**70.83**	**47.50**	**1.49**	48.57	62.18	0.78	58.74	62.49	0.94