1

Is shape in the eye of the beholder? Assessing landmarking error in geometric morphometric analyses on live fish

Paolo Moccetti[1,2,3], Jessica R. Rodger[4], Jonathan D. Bolland[2], Phoebe Kaiser-Wilks[5], Rowan
Smith[5], Andy D. Nunn[2], Colin E. Adams[5], Jen A. Bright[6], Hannele M. Honkanen[5], Angus J.
Lothian[4], Matthew Newton[5], Domino A. Joyce[1]

[1] Evolutionary and Ecological Genomics Group, School of Natural Sciences, University of
Hull, UK
[2] Hull International Fisheries Institute, School of Natural Sciences, University of Hull, UK
[3] Energy and Environment Institute, University of Hull, UK
[4] Atlantic Salmon Trust Fellow, Scottish Centre for Ecology and the Natural Environment,
Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,
UK
[5] Scottish Centre for Ecology and the Natural Environment, Institute of Biodiversity, Animal
Health and Comparative Medicine, University of Glasgow, UK
[6] School of Natural Sciences, University of Hull, UK

Corresponding Author:
Paolo Moccetti[1,2,3]
School of Natural Sciences, University of Hull
Cottingham Road, Hull, HU6 7RX, UK
Email address: p.m.moccetti-2019@hull.ac.uk, moccetti.paolo@gmail.com

Abstract

Geometric morphometrics is widely used to quantify morphological variation between
biological specimens, but the fundamental influence of operator bias on data reproducibility
is rarely considered, particularly in studies using photographs of live animals taken under
field conditions. We examined this using four independent operators that applied an identical
landmarking scheme to replicate photographs of 291 live Atlantic salmon (*Salmo salar* L.)
from two rivers. Using repeated measures tests, we found significant inter-operator
differences in mean body shape, suggesting that the operators introduced a systematic error
despite following the same landmarking scheme. No significant differences were detected
when the landmarking process was repeated by the same operator on a random subset of
photographs. Importantly, in spite of significant operator bias, small but statistically
significant morphological differences between fish from the two rivers were found
consistently by all operators. Pairwise tests of angles of vectors of shape change showed that
these between-river differences in body shape were analogous across operator datasets,

suggesting a general reproducibility of findings obtained by geometric morphometric studies.
In contrast, merging landmark data when fish from each river are digitised by different
operators had a significant impact on downstream analyses, highlighting an intrinsic risk of
bias. Overall, we show that, even when significant inter-operator error is introduced during
digitisation, following an identical landmarking scheme can identify morphological
differences between populations. This study indicates that operators digitising at least a sub-
set of all data groups of interest may be an effective way of mitigating inter-operator error
and potentially enabling data sharing.
50

Introduction

51
Landmark-based geometric morphometrics (GM) is a quantitative approach widely used to
describe the shape of biological specimens and its covariation with other biological and
environmental factors (Zelditch et al., 2004; Webster & Sheets, 2010). Morphological
variables are quantified using a set of Cartesian landmarks located on distinct homologous
anatomical points, and observed body shape variations are then displayed through user-
friendly graphical representations (Zelditch et al., 2004; Mitteroecker & Gunz, 2009; Adams,
Rohlf & Slice, 2013). GM is a powerful technique capable of detecting even tiny
morphological differences among groups of specimens (Mitteroecker & Gunz, 2009; Webster
& Sheets, 2010), but is highly sensitive to measurement errors introduced during data
acquisition, which can affect subsequent analyses and produce inaccurate results (von
Cramon-Taubadel, Frazier & Lahr, 2007; Fruciano, 2016; Robinson & Terhune, 2017; Fox,
Veneracion & Blois, 2020). This is particularly problematic when such morphological
differences are erroneously regarded as biologically meaningful variation (Fruciano, 2016).
65
Surprisingly, despite GM being a widely used technique, researchers rarely consider
measurement error in their study design and statistical analyses (Fruciano, 2016; Fox,
Veneracion & Blois, 2020). Measurement error can be introduced at different stages of the
data acquisition process, i.e. when positioning specimens in front of the imaging device
(camera or scanner), during image capture and landmark digitisation (Arnqvist &
Mårtensson, 1998; Muir, Vecsei & Krueger, 2012; Fruciano et al., 2020; Fox, Veneracion &
Blois, 2020). Indeed, the so-called inter-operator (or inter-observer) error during landmarking
has been found to be one of the most critical factors affecting GM analyses because different
operators tend to position what should be homologous landmarks in slightly different
locations (Ross & Williams, 2008; Dujardin, Kaba & Henry, 2010; Campomanes-Álvarez et
al., 2015; Fruciano, 2016; Fruciano et al., 2020; Fox, Veneracion & Blois, 2020).
Importantly, inter-operator error can be substantial and potentially obscure biological
variation, making data sharing and comparisons of landmarked datasets difficult (Shearer et
al., 2017).
80
Intra-operator (or intra-observer) error has also been shown to significantly affect GM
analyses (Wilson, Cardoso & Humphrey, 2011; Fox, Veneracion & Blois, 2020). Intra-
operator error is introduced when specimens are inconsistently digitised by a single operator
and can be influenced by several factors, including landmarking experience or time between

landmarking sessions (Fox, Veneracion & Blois, 2020). However, the magnitude of intra-
operator error is invariably modest compared to inter-operator discrepancies (Cardoso &
Saunders, 2008; Dujardin, Kaba & Henry, 2010; Wilson, Cardoso & Humphrey, 2011;
Robinson & Terhune, 2017; Shearer et al., 2017; Thoma et al., 2018; Fox, Veneracion &
Blois, 2020), indicating a general good precision in digitisation by individual operators (but
see Engelkes et al., 2019).
91

The degree and impacts of operator error in GM studies have been tested for a range of
organisms, anatomical structures, preservation methods and image acquisition devices
(Fruciano, 2016; Fruciano et al., 2020; Fox, Veneracion & Blois, 2020). Nevertheless, most
studies have focussed on images of specific human, bone or plant structures acquired under
identical (laboratory) conditions (e.g., Ross & Williams, 2008; Cardoso & Saunders, 2008;
Gonzalez, Bernal & Perez, 2011; Wilson, Cardoso & Humphrey, 2011; Viscosi & Cardini,
2011; Shearer et al., 2017; Carayon et al., 2019; Engelkes et al., 2019; Messer et al., 2021).
Few have investigated images of live animals (but see Fruciano et al., 2020), despite
commonly being used when it is not possible to euthanise samples for ethical reasons or
research purposes. Undeniably, such photographs, especially if taken under field conditions,
are more likely to result in subsequent measurement error (relative to preserved specimens)
(Muir, Vecsei & Krueger, 2012), thereby restricting the utility of such datasets (Webster &
Sheets, 2010). Understanding the prevalence, magnitude and implications of inter- and intra-
operator error during the landmark digitisation process for photographs of live animals could
facilitate data sharing and open science practices.
107

With the increasing focus on reproducibility in science (Baker, 2016), and an
acknowledgment that sharing data can accelerate scientific progress, assessing whether live
animals digitised repeatedly by single versus multiple operators produce consistent results
and conclusions is essential. Data exchange, such as crowdsourcing, is opening new frontiers
in GM research, enabling large-scale studies, which use unprecedented sample sizes, to be
conducted within a short time frame (Thomas, Bright & Cooney, 2016; Chang & Alfaro,
2016). Such studies, involving several operators collecting shape data, can potentially address
key questions in evolutionary biology and other disciplines (Cooney et al., 2017; Hughes et
al., 2022). However, pooling landmarked datasets from multiple operators can increase the
degree of measurement error (Fruciano et al., 2017; Evin, Bonhomme & Claude, 2020), but
the consequences of inter-individual operator error when sharing datasets remain poorly
understood.
120

The aim of this study was therefore to determine whether GM analyses on photographs of
live animals are reproducible. To accomplish this, four independent operators digitised the
same photographs of sedated Atlantic salmon (*Salmo salar* L.) sampled in two rivers,
following a shared landmarking scheme. The shape data and results obtained by the four
operators were then compared and contrasted to assess the magnitude of inter- and intra-
operator error, and infer the potential for meaningful data sharing.
127

Material & Methods

Study design

Salmon were captured from the River Spey (57° 24.960' N 3° 22.602' W) and River Oykel (57° 59.640' N 4° 48.282' W) in Scotland using a 1.5 m diameter Rotary Screw trap during their smolt stage, i.e. on their first migration to sea. The sampling occurred in the context of a tracking study aiming to identify areas and causes of smolt mortality during their seaward migration (see Whelan, Roberts & Gray, 2019; and https://atlanticsalmontrust.org/our-work/morayfirthtrackingproject/). Fish were photographed in the field under anaesthetic before being tagged and released to the river after recovery. Photographs of the left side of each fish were taken freehand from approximately 30 cm directly above the fish, with a Fujifilm FinePix XP130 Compact Digital Camera with fish on a background reference scale. Photographs were taken by a team of eight people who met prior to field work to standardise methods as far as possible. Our study here focusses on inter-operator variation downstream of photography, but variation caused by variation between individual photographers would be worthy of future study. The care and use of experimental animals complied with the UK Home Office animal welfare laws, guidelines and policies (UK Home Office Licence PPL 70/8794) and was approved by the University of Glasgow Animal Welfare and Ethics Review Board (AWERB). Field permits were provided verbally by: Keith Williams: Kyle of Sutherland Fisheries Trust (River Oykel) and Brian Shaw: Spey District Salmon Fisheries Board (River Spey).

The GM analyses were based on photographs of 291 salmon (Spey $n = 144$, Oykel $n = 147$). The images were imported into tpsUtil v. 1.78 (Rohlf, 2019) and randomly shuffled using the relevant function so that operators were blinded to the river-of-origin of the specimens. Fifteen fixed and seven semi- landmarks (Bookstein, 1997) were digitised on each image by four independent operators (Op.1, Op.2, Op.3 and Op.4) using tpsDig v. 2.31 (Rohlf, 2017) and following an identical scheme (Fig. 1). The landmark positions chosen were those commonly used in studies on salmonids (e.g., Boulding et al., 2008; Muir, Vecsei & Krueger, 2012; Simonsen et al., 2017; Goerig et al., 2019; Dermond, Sperlich & Brodersen, 2019). In addition, the first ten fish for each river, after using the *randomly order specimens* function in tpsUtil, were consecutively landmarked a further two times (i.e. three times in total) by each operator to evaluate the intra-operator consistency in digitisation.

Landmark coordinates from all operators were imported as unique files into R (R Core Team, 2021) and analysed using the 'geomorph' and 'RRPP' v. 4.0.4 (Adams et al., 2021; Baken et al., 2021; Collyer & Adams, 2021), 'Morpho' v. 2.8 (Schlager, 2017), and 'GeometricMorphometricsMix' v. 0.0.8.4 (Fruciano, 2018) packages. Plots were produced with the 'ggplot2' package (Wickham, 2016), while projections of body shape variation between groups were generated with the *plotRefToTarget* function in 'geomorph'.

170　The landmark data were then used to test if: (1) similar mean body shapes were obtained by
171　all operators; (2) any morphological differences between salmon from the two different rivers
172　were detected by all operators; (3) identified between-river differences were consistent across
173　all operators; (4) divergent datasets from different operators could be merged; and (5) the
174　magnitude of intra-operator error was similar across operators.
175
176　*Preliminary analyses*
177
First, a generalised Procrustes analysis (GPA) with sliding of semi-landmarks was performed
to remove effects not related to body shape through translation, scaling and rotation of the
landmark configurations (Rohlf & Slice, 1990). A preliminary principal component analysis
(PCA) conducted on superimposed coordinates revealed body bending as a major source of
shape variation, a known issue in morphometric studies on fish (Valentin et al., 2008). To
remove the bending effect, the *unbend* function in tpsUtil was used, employing landmarks 1,
21, 22 and 17, which normally lie in a straight line in salmonids (Arbour, Hardie &
Hutchings, 2011; Dermond, Sperlich & Brodersen, 2019). All subsequent analyses were
performed on landmarks 1-20 only. A new GPA on coordinates with the bending deformation
removed was then executed and outlier specimens were investigated for each operator using
the *plotOutliers* function in 'geomorph'. Two specimens digitised by one operator were
found to be very different to the other individuals and were therefore removed from the
dataset of all four operators, leaving 289 samples for analyses (Spey $n = 144$; Oykel $n = 145$).
Another GPA using the landmark data without outliers was then implemented.
192
193　*Test 1. Were similar mean body shapes obtained by all operators?*
194
To investigate whether results produced by a single operator are accurate and reproducible,
we tested differences in the mean body shapes of fish digitised by independent operators.
First, a between-group PCA (Boulesteix, 2004) was computed to explore variations between
the four operators. Between-group PCA is a type of discriminant analysis used to maximise
segregation between known groups which, unlike canonical variate analysis (CVA), does not
overestimate the degree of distinction among groups (Mitteroecker & Bookstein, 2011). The
leave-one-out cross-validation operation was implemented to quantify the proportion of fish
specimens correctly assigned to the operator who digitised them.
203
To investigate whether landmarking by multiple operators introduced bias, i.e. systematic
error affecting body shape (*sensu* Fruciano, 2016), differences in the mean body shapes of the
fish digitised by the four independent operators were tested using Hotelling's T^2 as
implemented by the *repeated_measures_test* function in 'GeometricMorphometricsMix'. To
compute the differences in mean body shapes, a PCA was performed on all Procrustes-
aligned coordinates of all operators, and the scores for all the PC axes (i.e. 100% variance
explained) of each operator were then used in a repeated measures test as an approximation
of shape.
212

213 *Test 2. Were morphological differences between salmon from different rivers detected by all*
214 *operators?*
215
We next tested whether there was a difference in body shape between rivers, and whether the
operators were consistent in identifying any differences. The following analyses were
performed separately for each operator. First, a GPA was computed on landmark coordinate
datasets obtained by each operator with outliers removed (see end of *Preliminary analyses*).
The effect of fish size on body shape was tested using Procrustes ANOVAs (*procD.lm*
function in 'geomorph'), with Procrustes coordinates used as an outcome variable, the log
value of centroid size and 'River' as independent variables and a randomised residual
permutation procedure (10,000 iterations). A small but significant effect of size on shape was
found for all operators (*P*-values < 0.0001, r^2 = 0.022–0.034). Procrustes coordinates were
therefore adjusted for allometry by using residuals from a regression of shape against
centroid size + 'River'. Procrustes ANOVAs were then used to compare mean body shape
between rivers, while another between-group PCA was implemented to quantify the
proportion of fish correctly assigned to the river of origin, for each of the four datasets.
229
230 *Test 3. Were identified between-river differences consistent across all operators?*
231
To assess if body shape differences between rivers were analogous across operators, pairwise
angles (Li, 2011) of vectors of shape change between fish from the rivers Spey and Oykel
were computed. The *TestOfAngle* function in 'GeometricMorphometricsMix' based on the
analogous function implemented in 'Morpho' was used, as performed by Fruciano et al.
(2020). Specifically, we calculated the pairwise angles among between-group principal
components obtained using 'River' as the grouping factor within each operator subset of
digitisations (one between-group PC axis - herein bwgPC - per operator) to test if they
followed the same "direction", i.e. if the shape differences between rivers were
approximately the same for all operators.
Furthermore, bwgPC1 vectors of between-river differences for each operator were compared
(test of angles) with vectors of inter-operator differences obtained by subtracting
corresponding Procrustes corrected coordinates of each specimen (eg. [coordinates of
specimen 1 digitised by Op.1] – [coordinates of specimen 1 digitised by Op.2]) and then
calculating mean shapes for all specimens. In this way, it was possible to determine whether
or not biological body shape differences between rivers and artefactual variation among
operators were similar (following the same "direction").
Finally, the magnitude of shape differences between rivers obtained by each operator was
examined with the *dist_mean_boot* function in 'GeometricMorphometricsMix'. This function
was used to perform a bootstrap estimate of the shape distance between the two rivers and
allowed us to test if the amount of shape difference between the rivers Spey and Oykel was
consistent across different operators or, on the contrary, one or more operators detected larger
or smaller between-river differences than the others.
254
255 *Test 4. Can divergent datasets from different operators be merged?*

257 The two operators producing the most dissimilar mean shapes were used to simulate a worst-

258 case-scenario process of data pooling, in which two independent researchers perform their

259 own GM study each on different rivers, but following the same landmarking scheme. Inter-

260 operator analysis showed that Op.2 and Op.4 produced the most dissimilar body shapes

261 (greatest Euclidean distance), so from these, two datasets were created: one comprising shape

262 data from the River Oykel digitised by Op.2 (herein Op.2-Oykel) and the River Spey data

263 digitised by Op.4 (herein Op.4-Spey) and *vice versa*, i.e. the River Spey data digitised by

264 Op.2 (herein Op.2-Spey) and the River Oykel data digitised by Op.4 (herein Op.4-Oykel).

265 For both datasets, differences between rivers were tested with Procrustes ANOVA, as

266 described earlier. Then, a between-group PCA was performed and the resulting bwgPC1

267 separating the two rivers was used to run a test of angles to compare between-river

268 differences detected by these two separate datasets. We also compared these latter between-

269 river differences with those found when using the complete datasets of all four operators

270 including both rivers (see section above). This enabled us to test whether any between-river

271 differences as a result of different operators outweighed any biological differences between

272 rivers found when using the complete intra-operator datasets.

274 *Test 5. Quantifying intra-operator error*

276 A GPA was computed separately on landmark coordinates obtained by each operator re-

277 digitising a sub-sample of 20 fish (ten per river). Individual consistency in landmarking was

278 then investigated using PCA and tested using repeated measures tests. To test for differences

279 in mean body shapes between digitisation trials, a PCA was performed on the Procrustes-

280 aligned coordinates of each operator separately and the PC scores of each trial were then used

281 in the repeated measures tests as an approximation of shape. Repeatability among digitisation

282 trials was also calculated for each operator using the intraclass correlation coefficient (Fisher,

283 1958). A one-way Procrustes ANOVA was computed using individual fish as a categorical

284 variable (Fruciano, 2016). The resulting mean squares were used to calculate repeatability by

285 applying equations presented in Arnqvist & Mårtensson (1998) and Fruciano (2016). Here,

286 repeatability measured variation in the three independent digitisations of the sub-sample of

287 20 salmon relative to the variability among specimens, i.e. the biological variation among all

288 fish samples. Repeatability assumes a value of between zero and one, with one indicating

289 100% repeatability and an absence of measurement error (Arnqvist & Mårtensson, 1998;

290 Fruciano, 2016). Finally, a Procrustes ANOVA with individual fish specimens ('ID') as the

291 main factor and 'operator' nested within 'ID' was run to test the relative contributions of

292 biological variation ('ID') and variation introduced by inter-operator ('ID:operator') and

293 intra-operator (residual) error.

295 **Results**

297 *Test 1. Were similar mean body shapes obtained by all operators?*

Despite digitising replicate photographs with homologous landmarks, fish specimens were
correctly assigned to their operator based on body shape with 83.0% accuracy by the
exploratory between-group PCA (Fig. 2, Supplementary Table 1). There was a significant
operator effect on mean body shape, with all pairwise tests displaying highly significant
differences between operators (P-value < 0.001 for all comparisons; Table 1), supporting the
exploratory between-group PCA (Fig. 2). The Euclidean distances between means, i.e. the
measure of the extent of shape change, highlighted different distances among pairs of
operators, with the smallest difference (0.00771) occurring between Op.1 and Op.3 and the
greatest between Op.2 and Op.4 (0.01577). The between-group PCA scatterplot (Fig. 2)
broadly reflected these results along axis 1 (61.9% of variance), with Op.1 and Op.3
overlapping extensively and Op.2 and Op.4 being furthest apart. The anatomical differences
among operators were concentrated mainly on the head (Fig. 3, Supplementary Fig. 1), with
major areas of disagreement being the snout, eye, mouth and posterior of the cranium.
Morphological dissimilarities were more or less pronounced depending on the operator
comparisons (Fig. 3).

Test 2. Were morphological differences between salmon from different rivers detected by all
operators?

There were significant differences in body shape between fish from different rivers (Spey and
Oykel; Table 2), with between-group PCA (Supplementary Table 2) separating them for all
operators (70.9% mean classification success rate). The fish from the River Oykel had a
greater body depth, more pronounced caudal peduncle, larger eye, longer mouth and more
pointed snout than those from the River Spey (Fig. 4).

Test 3. Were identified between-river differences consistent across all operators?

All comparisons of the "direction" of body shape variation between rivers were significant
for all operators, meaning that the way in which shape differed between the rivers Spey and
Oykel was approximately the same for all operators (Table 3). In contrast, pairwise
comparisons of between-river and between-operator differences were mostly non-significant,
with nine of 24 tests generating P-values < 0.05 (Supplementary Table 3). This indicates that
the shape variation between rivers and operators were mainly divergent and not collinear.
Estimated mean distances between rivers computed through bootstrapping were similar
across operators, as shown by the widely overlapping confidence intervals (Table 4),
suggesting that different operators did not influence the magnitude of shape difference
detected between the rivers Spey and Oykel.

Test 4. Can divergent datasets from different operators be merged?

There were significant differences in body shape between fish from the rivers Spey and
Oykel digitised separately by Op.2 and Op.4 (Table 5). Notably, shape variation explained by
the rivers was markedly higher for these merged datasets compared to the between-river

differences detected by single operators ($r^2 = 0.11$-0.21 vs. 0.03-0.04, respectively; Tables 5
and 2). Similarly, the between-group PCA separated fish from different rivers with a higher
accuracy than the analogous analysis performed on individual operator datasets (93.6% vs.
70.9% mean classification success rate, respectively; Supplementary Tables 4 and 2).
The comparison of the "direction" of between-river body shape variation detected by the two
merged datasets from Op.2 and Op.4 was highly significant (P-value < 0.0001;
Supplementary Table 5), meaning that the way in which shape differed between the rivers
Spey and Oykel was approximately the same regardless of the selected dataset. However,
only five of eight comparisons were found to be significant when comparing the two Op.2
and Op.4 merged datasets with the complete within-operator datasets including both rivers
(Supplementary Table 5), indicating that the river differences detected by combined and
individual operator datasets were only partly similar.
354
355   *Test 5. Quantifying intra-operator error*
356
There was extensive overlap among landmarking trials, suggesting a high consistency in
digitisation across all operators (Fig. 5). Pairwise comparisons supported this since none of
the mean body shapes differed significantly between repeated digitisations (P-values > 0.82;
Table 6). All four operators achieved the highest landmarking consistency between trials 2
and 3, as indicated by the smallest Euclidean distance values (0.001-0.003; Table 6).
Repeatability was also high for all operators (0.925-0.977), indicating high landmarking
precision (Table 7). Nested Procrustes ANOVA showed that 67.3% of the morphological
variation within the subset of 20 fish was explained by individual fish ('ID'), while 25.7%
and 7.0% of the variation was attributable, respectively, to inter- ('ID:operator') and intra-
(residual) operator digitisation error (Supplementary Table 6).
367

Discussion
We show here that independent operators applying an identical landmarking scheme to
replicate photographs of live Atlantic salmon taken in field conditions yielded significantly
different mean body shapes (*Test 1*). However, morphological differences between salmon
from different rivers were detected by all operators (*Test 2*), and these were consistent
differences across all operators (*Test 3*), provided they landmarked both rivers and not one
each (*Test 4*). Furthermore, intra-operator error calculated on a subset of samples was
minimal, suggesting that it did not have a significant influence on the body shape results
obtained by the different operators (*Test 5*).
377

Despite digitising replicate photographs with homologous landmarks, all the operators
produced significantly different mean body shapes. The high rate (83.0%) of specimens
assigned to the correct operator by the between-group PCA suggests that the operators
introduced a systematic error, which created four identifiable body shapes despite following
the same landmarking scheme. This digitisation bias is likely to have been introduced by
operators consistently applying personal, fine-scaled landmarking rules in addition to the
general scheme. The fact that the differences among operators were localised mainly in the

head region (landmarks 1-12) may be explained by the less discrete and recognizable nature
of these landmarks compared to those located on well-defined anatomical loci, such as fin
intersections (landmarks 13-20). This suggests that the use of unambiguous landmarks can be
an effective way of reducing measurement error in GM (Fagertun et al., 2014; Campomanes-
Álvarez et al., 2015, Fruciano et al., 2017).
390

In GM studies, digitisation is typically performed by a single operator, leaving the question
of whether multiple operators digitising the same set of images would generate different
results. This could undermine the reliability of findings presented by many GM
investigations, particularly those using images of live animals taken in field conditions,
which are potentially more prone to measurement error (Webster & Sheets, 2010; Muir,
Vecsei & Krueger, 2012). In our study, however, we found that inconsistencies between
operators did not mask small, but significant morphological differences between fish from the
rivers Spey and Oykel, which were consistent across operators. The fact that, as shown by
tests of angles and bootstrapped estimates of mean distances, all the operators detected
analogous between-river differences, strongly suggests that they were biologically authentic.
Similarly, Fruciano et al. (2020) found that preservation methods significantly affected the
body shape of brown trout (*Salmo trutta* L.), but the subsequent between-groups
classification was similar regardless of preservation method. As suggested by Fruciano et al.
(2020), this could be because the shape variation detected by the operators between the rivers
Spey and Oykel was not significantly affected by inter-operator differences in landmarking
since they were not collinear (i.e. they followed different "directions", as shown by the angle
comparisons).
408

Conversely, merging landmark data of fish from the rivers Spey and Oykel digitised by two
distinct operators (Op.2 and Op.4, *Test 4*) had a significant impact on subsequent analyses
and produced contrasting results. As shown by Procrustes ANOVA and between-group PCA
classification rate, shape differences between rivers in the merged datasets were greater than
those detected by single operators, suggesting they were artificially inflated by inter-operator
digitisation error. Angle comparisons showed that the river differences detected by combined
and individual operator datasets were partly dissimilar. Overall, these findings point towards
a potential risk in pooling datasets from multiple operators when there are confounding
biological factors, as highlighted by other studies (Fruciano et al., 2017; Evin, Bonhomme &
Claude, 2020). Distinct operators obtained analogous results when they landmarked both
rivers (and not one river each as in *Test 4*). This suggests that operators digitising at least a
sub-set of all data groups of interest (rivers in this case) may be an effective way of
mitigating inter-operator error and potentially enabling data sharing.
422

In contrast to the inter-operator effects described in this study, we found no statistical
evidence of intra-operator effects on the quantification of fish morphology. On the contrary,
we found a very high level of repeatability across trials for all operators. This corroborates
previous studies that showed intra-operator error to be limited (e.g. Cardoso & Saunders,
2008; Dujardin, Kaba & Henry, 2010; Wilson, Cardoso & Humphrey, 2011; Robinson &

Terhune, 2017; Shearer et al., 2017; Thoma et al., 2018; Fox, Veneracion & Blois, 2020).
Interestingly, for all operators, landmarking consistency was highest between their last two
trials, suggesting that they 'learnt' where to place the landmarks with increasing experience
of the images. However, it should be noted that the first trial was performed while digitising
all specimens, whereas trials 2 and 3 were performed consecutively after digitising the full
dataset, which may have artificially inflated precision, with operators "remembering" their
landmarking choices in trial 2 during trial 3.
The negligible impact of intra- compared to inter-operator error was also clearly shown by
the percentage of variance explaining shape variation in the sub-sample of 20 fish (*Test 5*,
7.0% vs. 25.7%, respectively). Interestingly, the percentage of variance explained by inter-
operator error (25.7%) is similar to that reported by Fruciano et al. (2020) for brown trout
photographed in the field (30.1%), and supports previous studies that identified inter-operator
effects as the major source of error in GM analyses (Ross & Williams, 2008; Dujardin, Kaba
& Henry, 2010; Campomanes-Álvarez et al., 2015; Fruciano, 2016; Shearer et al., 2017; Fox,
Veneracion & Blois, 2020).
443

Conclusions

Overall, we show that, even when significant inter-operator error is introduced through
digitisation, following an identical landmarking scheme can be an effective tool to obtain
robust and reliable results, even without accounting for variation introduced by the
photography process, which was not quantified here. This implies that GM studies based on
common landmarking schemes are potentially reproducible, even when analyses are based on
images of live specimens taken in the field, as in the current study. Nevertheless, since
operator error can vary between studies and is impossible to determine *a priori*, we
recommend assessing the magnitude and effects of landmarking error by using multiple
operators for a sub-set of samples, as here, to improve confidence in study results. If
landmark data merging is required, we recommend that all the operators involved digitise at
least a sub-set of all data groups of interest (rivers in this case) to mitigate inter-operator
error.
457

Acknowledgements

459   We would like to thank the Spey Fisheries Board and the Kyle of Sutherland Fisheries Trust
460   for their help and support with this project. We would also like to thank Jonathan Archer,
461   Georgios Kyriakou, Fraser Brydon, Jessica Whitney and Mustafa Soganci for their help
462   collecting the data for this study. We also thank Carmelo Fruciano, Michelle C. Gilbert and
463   Kiran Liversage for their constructive reviews.
464

References

Adams DC, Rohlf FJ, Slice DE. 2013. A field comes of age: geometric morphometrics in the
21st century. *Hystrix, the Italian Journal of Mammalogy* 24:7–14. DOI:
10.4404/hystrix-24.1-6283.
Adams DC, Collyer M, Kaliontzopoulou A, Baken E. 2021. "Geomorph: Software for
geometric morphometric analyses. R package version 4.0.2." https://cran.r-

project.org/package=geomorph.

Arbour JH, Hardie DC, Hutchings JA. 2011. Morphometric and genetic analyses of two
sympatric morphs of Arctic char (*Salvelinus alpinus*) in the Canadian high Arctic.
Canadian Journal of Zoology 89:19–30. DOI: 10.1139/Z10-100.

Arnqvist G, Mårtensson T. 1998. Measurement error in geometric morphometrics: empirical
strategies to assess and reduce its impact on measures of shape. *Acta Zoologica*
Academiae Scientiarum Hungaricae 44:73–96.

Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC. 2021. geomorph v4.0 and gmShiny:
Enhanced analytics and a new graphical interface for a comprehensive morphometric
experience. *Methods in Ecology and Evolution* 12:2355–2363. DOI:
https://doi.org/10.1111/2041-210X.13723.

Baker M. 2016. 1,500 scientists lift the lid on reproducibility. *Nature* 533:452–454. DOI:
10.1038/533452a.

Bookstein FL. 1997. Landmark methods for forms without landmarks: morphometrics of
group differences in outline shape. *Medical Image Analysis* 1:225–243. DOI:
https://doi.org/10.1016/S1361-8415(97)85012-8.

Boulding EG, Culling M, Glebe B, Berg PR, Lien S, Moen T. 2008. Conservation genomics
of Atlantic salmon: SNPs associated with QTLs for adaptive traits in parr from four
trans-Atlantic backcrosses. *Heredity* 101:381–391. DOI: 10.1038/hdy.2008.67.

Boulesteix A. 2004. A note on between-group PCA. *International Journal of Pure and*
Applied Mathematics 19:359–366. DOI: 10.5282/ubm/epub.1767.

Campomanes-Álvarez BR, Ibáñez O, Navarro F, Alemán I, Cordón O, Damas S. 2015.
Dispersion assessment in the location of facial landmarks on photographs. *International*
Journal of Legal Medicine 129:227–236. DOI: 10.1007/s00414-014-1002-4.

Carayon D, Adhikari K, Monsarrat P, Dumoncel J, Braga J, Duployer B, Delgado M,
Fuentes-Guajardo M, de Beer F, Hoffman JW, Oettlé AC, Donat R, Pan L, Ruiz-Linares
A, Tenailleau C, Vaysse F, Esclassan R, Zanolli C. 2019. A geometric morphometric
approach to the study of variation of shovel-shaped incisors. *American Journal of*
Physical Anthropology 168:229–241. DOI: https://doi.org/10.1002/ajpa.23709.

Cardoso HF V, Saunders SR. 2008. Two arch criteria of the ilium for sex determination of
immature skeletal remains: a test of their accuracy and an assessment of intra- and inter-
observer error. *Forensic Science International* 178:24–29. DOI:
10.1016/j.forsciint.2008.01.012.

Chang J, Alfaro ME. 2016. Crowdsourced geometric morphometrics enable rapid large-scale
collection and analysis of phenotypic data. *Methods in Ecology and Evolution* 7:472–
482. DOI: https://doi.org/10.1111/2041-210X.12508.

Collyer ML, Adams DC. 2021. "RRPP: Linear Model Evaluation with Randomized
Residuals in a Permutation Procedure, R package version 1.1.2." https://cran.r-
project.org/package=RRPP.

Cooney CR, Bright JA, Capp EJR, Chira AM, Hughes EC, Moody CJA, Nouri LO, Varley
ZK, Thomas GH. 2017. Mega-evolutionary dynamics of the adaptive radiation of birds.
Nature 542:344–347. DOI: 10.1038/nature21074.

Dermond P, Sperlich N, Brodersen J. 2019. Heritable morphological differentiation in

salmonids from two distinct stream types. *Journal of Fish Biology* 95:1215–1222. DOI: 10.1111/jfb.14121.

Dujardin J-PAL, Kaba D, Henry AB. 2010. The exchangeability of shape. *BMC Research Notes* 3:266. DOI: 10.1186/1756-0500-3-266.

Engelkes K, Helfsgott J, Hammel JU, Büsse S, Kleinteich T, Beerlink A, Gorb SN, Haas A. 2019. Measurement error in μCT-based three-dimensional geometric morphometrics introduced by surface generation and landmark data acquisition. *Journal of Anatomy* 235:357–378. DOI: https://doi.org/10.1111/joa.12999.

Evin A, Bonhomme V, Claude J. 2020. Optimizing digitalization effort in morphometrics. *Biology Methods and Protocols* 5:bpaa023. DOI: 10.1093/biomethods/bpaa023.

Fagertun J, Harder S, Rosengren A, Moeller C, Werge T, Paulsen RR, Hansen TF. 2014. 3D facial landmarks: Inter-operator variability of manual annotation. *BMC Medical Imaging* 14:35. DOI: 10.1186/1471-2342-14-35.

Fisher RA. 1958. *Statistical Methods for Research Workers*. Oliver & Boyd, Edinburgh.

Fox NS, Veneracion JJ, Blois JL. 2020. Are geometric morphometric analyses replicable? Evaluating landmark measurement error and its impact on extant and fossil *Microtus* classification. *Ecology and Evolution* 10:3260–3275. DOI: https://doi.org/10.1002/ece3.6063.

Fruciano C. 2016. Measurement error in geometric morphometrics. *Development Genes and Evolution* 226:139–158. DOI: 10.1007/s00427-016-0537-4.

Fruciano C. 2018. GeometricMorphometricsMix: miscellaneous functions useful for geometric morphometrics. Version 0.0.3. https://github.com/fruciano/GeometricMorphometricsMix.

Fruciano C, Celik MA, Butler K, Dooley T, Weisbecker V, Phillips MJ. 2017. Sharing is caring? Measurement error and the issues arising from combining 3D morphometric datasets. *Ecology and Evolution* 7:7034–7046. DOI: https://doi.org/10.1002/ece3.3256.

Fruciano C, Schmidt D, Ramírez Sanchez MM, Morek W, Avila Valle Z, Talijančić I, Pecoraro C, Schermann Legionnet A. 2020. Tissue preservation can affect geometric morphometric analyses: a case study using fish body shape. *Zoological Journal of the Linnean Society* 188:148–162. DOI: 10.1093/zoolinnean/zlz069.

Goerig E, Wasserman BA, Castro-Santos T, Palkovacs EP. 2019. Body shape is related to the attempt rate and passage success of brook trout at in-stream barriers. *Journal of Applied Ecology* 57:91–100. DOI: 10.1111/1365-2664.13497.

Gonzalez PN, Bernal V, Perez SI. 2011. Analysis of sexual dimorphism of craniofacial traits using geometric morphometric techniques. *International Journal of Osteoarchaeology* 21:82–91. DOI: https://doi.org/10.1002/oa.1109.

Hughes EC, Edwards DP, Bright JA, Capp EJR, Cooney CR, Varley ZK, Thomas GH. 2022. Global biogeographic patterns of avian morphological diversity. *Ecology Letters* 25:598–610. DOI: https://doi.org/10.1111/ele.13905.

Li S. 2011. Concise formulas for the area and volume of a hyperspherical cap. *Asian Journal of Mathematics & Statistics* 4:66–70. DOI: 10.3923/ajms.2011.66.70.

Messer D, Svendsen MS, Galatius A, Olsen MT, Dahl VA, Conradsen K, Dahl AB. 2021. Measurement error using a SeeMaLab structured light 3D scanner against a Microscribe

3D digitizer. *PeerJ* 9:e11804. DOI: 10.7717/peerj.11804.

Mitteroecker P, Bookstein F. 2011. Linear discrimination, ordination, and the visualization of
selection gradients in modern morphometrics. *Evolutionary Biology* 38:100–114. DOI:
10.1007/s11692-011-9109-8.

Mitteroecker P, Gunz P. 2009. Advances in geometric morphometrics. *Evolutionary Biology*
36:235–247. DOI: 10.1007/S11692-009-9055-X/FIGURES/7.

Muir AM, Vecsei P, Krueger CC. 2012. A perspective on perspectives: methods to reduce
variation in shape analysis of digital images. *Transactions of the American Fisheries*
Society 141:1161–1170. DOI: 10.1080/00028487.2012.685823.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Robinson C, Terhune CE. 2017. Error in geometric morphometric data collection: combining
data from multiple sources. *American Journal of Physical Anthropology* 164:62–75.
DOI: https://doi.org/10.1002/ajpa.23257.

Rohlf F. 2017. tpsUtil v. 1.78. Stony Brook, NY: Department of Ecology and Evolution,
State University of New York.

Rohlf F. 2019. tpsDig v. 2.31, Stony Brook, NY: Department of Ecology and Evolution, State
University of New York.

Rohlf FJ, Slice D. 1990. Extensions of the Procrustes method for the optimal superimposition
of landmarks. *Systematic Biology* 39:40–59. DOI: 10.2307/2992207.

Ross AH, Williams S. 2008. Testing repeatability and error of coordinate landmark data
acquired from crania. *Journal of Forensic Sciences* 53:782–785. DOI:
https://doi.org/10.1111/j.1556-4029.2008.00751.x.

Schlager S. 2017. Morpho and Rvcg – Shape Analysis in R. In: *Statistical Shape and*
Deformation Analysis (eds Zheng G, Li S, Szekely G), pp. 217–256. Academic Press,
London. ISBN 9780128104934.

Shearer BM, Cooke SB, Halenar LB, Reber SL, Plummer JE, Delson E, Tallman M. 2017.
Evaluating causes of error in landmark-based data collection using scanners. *PLOS ONE*
12:e0187452.

Simonsen MK, Siwertsson A, Adams CE, Amundsen PA, Præbel K, Knudsen R. 2017.
Allometric trajectories of body and head morphology in three sympatric Arctic charr
(*Salvelinus alpinus* (L.)) morphs. *Ecology and Evolution* 7:7277–7289. DOI:
10.1002/ece3.3224.

Thoma C, Makridou K, Bakaloudis D, Vlachos C. 2018. Evaluating the potential of three-
dimensional laser surface scanning as an alternative method of obtaining morphometric
data. *Annales Zoologici Fennici* 55:55–66. DOI: 10.5735/086.055.0106.

Thomas GH; Bright JA; Cooney CR. 2016. *Mark My Bird* [Data set]. Natural History
Museum. DOI: 10.5519/0005413

Valentin AE, Penin X, Chanut JP, Sévigny JM, Rohlf FJ. 2008. Arching effect on fish body
shape in geometric morphometric studies. *Journal of Fish Biology* 73:623–638. DOI:
10.1111/j.1095-8649.2008.01961.x.

Viscosi V, Cardini A. 2011. Leaf Morphology, Taxonomy and Geometric Morphometrics: A
Simplified Protocol for Beginners. *PLOS ONE* 6:e25630.

von Cramon-Taubadel N, Frazier BC, Lahr MM. 2007. The problem of assessing landmark
error in geometric morphometrics: theory, methods, and modifications. *American*
Journal of Physical Anthropology 134:24–35. DOI: https://doi.org/10.1002/ajpa.20616.
Webster M, Sheets HD. 2010. A practical introduction to landmark-based geometric
morphometrics. *Quantitative methods in Paleobiology* 16:168–188. DOI:
10.1017/S1089332600001868.
Whelan K, Roberts D, Gray J. 2019. The SAMARCH Project International Salmonid Coastal
and Marine Telemetry Workshop. Available at https://www.samarch.org/wp-
content/uploads/2020/05/SAMARCH-Tracking-Conference-Nov-2019-
final_compressed.pdf (accessed 11 March 2022).
Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New
York. ISBN 978-3-319-24277-4.
Wilson LAB, Cardoso HFV, Humphrey LT. 2011. On the reliability of a geometric
morphometric approach to sex determination: a blind test of six criteria of the juvenile
ilium. *Forensic Science International* 206:35–42. DOI: 10.1016/j.forsciint.2010.06.014.
Zelditch ML, Swiderski DL, Sheets HD, Fink WL. 2004. Introduction. In: *Geometric*
Morphometrics for Beginners: a Primer (eds Zelditch ML, Swiderski DL, Sheets HD,
Fink WL), pp. 1–20. Academic Press, San Diego. DOI: https://doi.org/10.1016/B978-

Figure 1. Fixed (blue) and semi- (green) landmarks used for the geometric morphometrics

analyses of Atlantic salmon smolts. Landmarks 21 and 22 (red) were used to correct for body

arching and not included in the analyses. (1) Tip of snout; (2) Midpoint between 1 and 3; (3)

Directly above middle of eye; (4) Perpendicular to 3; (5) Midpoint of top of eye (directly

below 3); (6) Midpoint of bottom of eye (directly below 3); (7) Midpoint of posterior of eye;

(8) Midpoint of anterior of eye; (9) End of maxillary bone; (10) Posterior tip of bony

operculum; (11) Dorsal surface posterior of cranium; (12) Perpendicular to 11; (13) Anterior

insertion point of dorsal fin; (14) Anterior insertion point of adipose fin; (15) Dorsal insertion

point of caudal fin; (16) Versal insertion point of caudal fin; (17) Posterior midpoint of

hypural plate; (18) Anterior insertion point of anal fin; (19) Anterior insertion point of ventral

fin; (20) Anterior insertion point of pectoral fin; (21) Lateral line - perpendicular to 13; (22)

Lateral line - perpendicular to 18. Anterior insertion point of ventral fin; (20) Anterior

insertion point of pectoral fin; (21) Lateral line - perpendicular to 13; (22) Lateral line -

perpendicular to 18.

645
646 **Figure 2.** Between-operator PCA scatterplot showing the cross-validated scores along the
647 first two between-group principal components (bwgPCs). Dots represent individual Atlantic
648 salmon (*n* = 289) landmarked by four independent operators (different colours). Between-
649 operator variance (%) explained by the first and second axes is shown.

Op.1 - Op.2 Op.1 - Op.3

Op.1 - Op.4 Op.2 - Op.3

Op.2 - Op.4 Op.3 - Op.4

Figure 3. Pairwise comparisons of the mean body shape of 289 Atlantic salmon landmarked by four independent operators. Morphological differences were magnified five times to aid visualisation.

Op.1 Op.2

Op.3 Op.4

Figure 4. Comparisons of the mean body shape of 289 Atlantic salmon in the rivers Spey (black) and Oykel (blue) landmarked by four independent operators. Morphological differences were magnified six times to aid visualisation.

Figure 5. Principal components analysis scatterplots of Procrustes-aligned coordinates for 20
Atlantic salmon in three landmarking trials by four independent operators. Dots represent
individual fish. Variance (%) explained by the first and second axes and 95% confidence
ellipses are shown.

679 **Tables**

681 **Table 1.** Pairwise comparisons of the body shape of 289 Atlantic salmon landmarked by four
682 independent operators based on Hotelling's T^2.

Comparison	Euclidean dist.	Hotelling's T^2	F	P-value
Op.1 vs. Op.2	0.01023	7100.15	162.84	$< 1 \times 10^{-6}$
Op.1 vs. Op.3	0.00771	3422.47	78.49	$< 1 \times 10^{-6}$
Op.1 vs. Op.4	0.01135	5130.73	117.67	$< 1 \times 10^{-6}$
Op.2 vs. Op.3	0.01215	9760.50	223.86	$< 1 \times 10^{-6}$
Op.2 vs. Op.4	0.01577	7511.23	172.27	$< 1 \times 10^{-6}$
Op.3 vs. Op.4	0.01114	5910.24	135.55	$< 1 \times 10^{-6}$

685 **Table 2.** Procrustes ANOVA summary statistics of effect of river-of-origin on the body
686 shape of 289 Atlantic salmon landmarked by four independent operators.

Operator	Df	SS	r^2	F	Z	P-value
Op.1	1	0.002411	0.03208	9.5117	4.7621	0.0001
Op.2	1	0.003243	0.03744	11.163	5.7723	0.0001
Op.3	1	0.003394	0.04069	12.172	5.2423	0.0001
Op.4	1	0.003090	0.03388	10.063	5.4314	0.0001

687 Df = Degrees of freedom, SS = Sum of squares, F = F statistics, Z = Effect size

688
689
690
691

692 **Table 3.** Pairwise tests of angles between body shape differences among rivers detected by
693 four different operators. Measurements of angles (degrees) between bwgPC1 vectors (below
694 the diagonal) and P-values (above the diagonal) are shown. Significant P-values (in bold)
695 indicate that shape change vectors are similar to each other.

	Op.1 (Spey vs. Oykel)	Op.2 (Spey vs. Oykel)	Op.3 (Spey vs. Oykel)	Op.4 (Spey vs. Oykel)
Op.1 (Spey vs. Oykel)		**5.3×10^{-20}**	**1.8×10^{-14}**	**6.0×10^{-17}**
Op.2 (Spey vs. Oykel)	22.7°		**1.1×10^{-22}**	**1.9×10^{-19}**
Op.3 (Spey vs. Oykel)	28.4°	17.0°		**1.9×10^{-19}**
Op.4 (Spey vs. Oykel)	24.3 °	20.8°	20.8°	

696
697
698

699 **Table 4.** Estimated mean and median shape distance (with confidence intervals) between the
700 rivers Spey and Oykel obtained by each operator.

Operator	Mean distance	Median distance	Lower CI extreme	Upper CI extreme
Op.1	0.00606	0.00608	0.00491	0.00729
Op.2	0.00697	0.00695	0.00573	0.00821
Op.3	0.00707	0.00706	0.00571	0.00845
Op.4	0.00684	0.00684	0.00553	0.00821

701
702

Table 5. Procrustes ANOVA summary statistics of effect of river-of-origin on the body shape of 289 Atlantic salmon based on combined datasets of Op.2 and Op.4.

Dataset	Df	SS	r^2	F	Z	P-value
Op.2-Oykel Op.4-Spey	1	0.011268	0.11434	37.051	8.872	0.0001
Op.2-Spey Op.4-Oykel	1	0.023561	0.21159	77.022	8.2704	0.0001

Df = Degrees of freedom, SS = Sum of squares, F = F statistics, Z = Effect size

Table 6. Pairwise comparisons of the body shape of 20 Atlantic salmon in three landmarking trials by four independent operators.

Operator	Trials	Euclidean dist.	Hotelling's T^2	F	P-value
Op.1	1 vs. 2	0.00240	67.48440	0.1869	0.97
	1 vs. 3	0.00315	141.9397	0.3932	0.87
	2 vs. 3	0.00153	22.45716	0.0622	0.99
Op.2	1 vs. 2	0.00358	47.28936	0.1310	0.99
	1 vs. 3	0.00377	69.33009	0.1921	0.97
	2 vs. 3	0.00236	56.09318	0.1554	0.98
Op.3	1 vs. 2	0.00353	49.37887	0.1368	0.99
	1 vs. 3	0.00360	50.734703	0.1405	0.98
	2 vs. 3	0.00127	15.02887	0.0416	0.99
Op.4	1 vs. 2	0.00922	184.9322	0.5123	0.82
	1 vs. 3	0.00731	128.5921	0.3562	0.89
	2 vs. 3	0.00344	58.91748	0.1632	0.98

Table 7. Repeatability values for the three landmarking trials on 20 Atlantic salmon by four independent operators.

Operator	Repeatability	Procrustes ANOVA r^2 (%)
Op.1	0.977	95.3
Op.2	0.961	92.4
Op.3	0.956	91.5
Op.4	0.925	86.3