Macro-scale patterns in functional connectivity associated with ongoing thought patterns and dispositional traits

Samyogita Hardikar [1], Brontë Mckeown [2], H. Lina Schaare [3, 4], Raven Wallace [2], Ting Xu [5],

Mark Edgar Lauckner, [6], Sofie L. Valk [3, 4, 7], Daniel S. Margulies [8], Adam Turnbull [9 10], Boris

Bernhardt [11], Reinder Vos de Wael [11], Arno Villringer [1, 12-14], Jonathan Smallwood [2].

[1] Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[2] Department of Psychology, Queens University, Kingston, Ontario, Canada.

[3] Otto Hahn Group Cognitive Neurogenetics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[4] Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich, Jülich, Germany

[5] Center for the Developing Brain, Child Mind Institute, New York, New York.

[6] Max Planck Research Group: Adaptive Memory, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[7] Institute of Systems Neuroscience, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

[8] Frontlab, Institut du Cerveau et de la Moelle épinière, UPMC UMRS 1127, Inserm U 1127, CNRS UMR 7225 Paris, France

[9] Department of Psychiatry and Behavioral Sciences, Stanford University, CA 94305, USA

[10] Department of Brain and Cognitive Sciences, University of Rochester, NY 14627, USA

[11] McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Montreal, Canada

[12] Day Clinic of Cognitive Neurology, Universitätsklinikum Leipzig, Leipzig, Germany.

[13] MindBrainBody Institute, Berlin School of Mind and Brain, Humboldt-Universität zu Berlin, 10099 Berlin, Germany;

[14] Center for Stroke Research Berlin (CSB), Charité - Universitätsmedizin Berlin, 10117 Berlin, Germany

Abstract

Complex macro-scale patterns of brain activity that emerge during periods of wakeful rest provide insight into the organisation of neural function, how these differentiate individuals based on their traits, and the neural basis of different types of self-generated thoughts. Although brain activity during wakeful rest is valuable for understanding important features of human cognition, its unconstrained nature makes it difficult to disentangle neural features related to personality traits from those related to the thoughts occurring at rest. Our study builds on recent perspectives from work on ongoing conscious thought that highlight the interactions between three brain networks - ventral and dorsal attention networks, as well as the default mode network. We combined measures of personality with state-of-the-art indices of ongoing thoughts at rest and brain imaging analysis and explored whether this 'tri-partite' view can provide a framework within which to understand the contribution of states and traits to observed patterns of neural activity at rest. To capture macro-scale relationships between different brain systems, we calculated cortical gradients to describe brain organisation in a low dimensional space. Our analysis established that for more introverted individuals, regions of the ventral attention network were functionally more aligned to regions of the somatomotor system and the default mode network. At the same time, a pattern of detailed self-generated thought was associated with a decoupling of regions of dorsal attention from regions in the default mode network. Our study, therefore, establishes that interactions between attention systems and the default mode network are important influences on ongoing thought at rest and highlights the value of integrating contemporary perspectives on conscious experience when understanding patterns of brain activity at rest.

Introduction

Macro-scale patterns of brain activity at rest have the potential for elucidating the organisation of neural function, different types of psychiatric conditions (Cao et al., 2018; Koban et al., 2021), developmental changes including those during adolescence and old age (Dosenbach et al., 2010; Cui et al., 2020; Gratton et al., 2020; Wen et al., 2020), neurological disorders (Zhang et al., 2021) and are important for revealing the neural basis behind the landscape of self-generated experiences (Karapanagiotidis et al., 2019; Mckeown et al., 2020; Kucyi et al., 2021). However, compared to controlled experimental conditions, interpreting neural activity recorded during resting-state functional magnetic resonance imaging (rs-fMRI) is challenging, partly because both trait-level aspects of the individual, and, the inherently complex and dynamic nature of ongoing experience at rest are both contributory factors to the observed brain activity (Smallwood et al., 2021b). It has recently been suggested that the meaning of different patterns of neural activity can be usefully constrained by pairing imaging data at rest with additional measures (Finn, 2021), for example by accounting for the patterns of thoughts individuals experience at rest (Karapanagiotidis et al., 2020, 2021; Mckeown et al., 2020; Gonzalez-Castillo et al., 2021; Kucyi et al., 2021), trait variation in how people think during tasks (Smallwood et al., 2016) or features of their personality (Hsu et al., 2018). While this methodological perspective is invaluable, we currently lack a theoretical framework within which to understand the brain-cognition relationships that these observations will establish. To address this gap in the literature, our study explores whether contemporary theories of the neural basis of ongoing conscious thought can provide a framework within which to interpret associations between macro-scale patterns of neural activity observed at rest, and measures of traits and self-reports of ongoing experience.

Emerging views of how the brain supports patterns of ongoing conscious thought highlight interactions between three large scale networks (Menon, 2011; Huang et al., 2021; Smallwood et al., 2021b): the ventral attention network (VAN), the dorsal attention network (DAN) and the

default mode network (DMN)⊕. Traditionally, it was argued that the DMN was thought to have

an antagonistic relationship with systems linked to external processing (Fox et al., 2005).

However, according to the 'tri-partite' network accounts the relationship between the DMN and

other brain systems is more nuanced. From this perspective key hubs of the ventral attention

network, such as the anterior insula and dorso-lateral prefrontal cortex, help gate access to

conscious experience regardless of the focus of attention. This is hypothesised to occur

because the VAN influences interactions between the DAN, which is more important for

external mental content (Corbetta and Shulman, 2002), and the DMN which is important when

states (including tasks) rely more on internal representations (Smallwood et al., 2021a). For

example, Huang and colleagues (Huang et al., 2021) established that activity levels in the

anterior insula determine whether stimuli presented at perceptual threshold are consciously

perceived, and this gating of external input emerged as a consequence of changes in the

normal interactions between the DAN and DMN. They also found that disruptions to activity in

the insula through anaesthesia resulted in reductions in self-generated mental imagery.

Coming from a different perspective, Turnbull and colleagues (Turnbull et al., 2019b) used

experience sampling during task performance to link patterns of neural activity to different

features of ongoing thought. For example, they found activity in the dorsolateral prefrontal

cortex (a member of the VAN according to a parcellation by Yeo, Krienen and colleagues,

(Yeo et al., 2011) was correlated with apparently contradictory patterns of ongoing thought –

(i) self-generated episodic thoughts during periods of low demands, and (ii) patterns of detailed

task focus when individuals were engaged in demanding external task. In the same study,

however, neural activity within regions of the dorsal parietal cortex within the DAN, was

exclusively reduced when participants engaged in self-generated thinking, highlighting a

parallel neglect of external input seen by Huang and colleagues. Finally, in a second study,

Turnbull and colleagues found that at rest, trait variance in the ability to focus on self-

generated experience in laboratory situations with lower task demands is associated with

⊕ In this paper we refer to the networks using the taxonomy provided by Yeo, Krienen and colleagues (Yeo et al., 2011)

decoupling of signals arising from the DAN and DMN (Turnbull et al., 2019a). Summarising

this emerging evidence, studies focused on understanding ongoing thought patterns from

different perspectives converge on the view that regions of the VAN may be important for

gating conscious access to different types of content by biasing interactions between the DAN

and the DMN (Huang et al., 2021; Smallwood et al., 2021b).

Our current study explored whether this "tri-partite network" view of ongoing conscious thought

derived from studies focused on understanding conscious experience, provides a useful

organising framework for understanding the relation between observed brain activity at rest

and patterns of cognition/ personality traits. Such analysis is important because at rest there

are multiple features of brain activity that can be identified via complex analyses that include

regions that show patterns of coactivation (which are traditionally viewed as forming a

cohesive network, (Biswal et al., 1995) as well as patterns of anti-correlation with other regions

(e.g. Fox et al., 2005). However, it is unclear which of these relationships reflect aspects of

cognition or behaviour or are in fact aspects of the functional organisation of the cortex (Fox

and Raichle, 2007). Consequently, our study builds on foundational work (e.g.

Vanhaudenhuyse et al., 2011) in order to better understand which aspects of neural function

observed at rest are mostly likely linked to cognition and behaviour. With this aim in mind, we

examined links between macro-scale neural activation and both (i) trait descriptions of

individuals and (ii) patterns of ongoing thought. Our sample was a cohort of 144 participants

who underwent a one-hour resting-state scan. Across this one-hour period, participants were

interrupted on four occasions to answer a set of questions about their experiences at rest

using multidimensional experience sampling (MDES), similar to a number of prior

neuroimaging studies (Smallwood et al., 2016; Poerio et al., 2017; Wang et al., 2018). During

a different session, the same participants also completed a battery of measures assessing

features of their personality (such as the Big five (Costa and McCrae, 2008)) as well as

subclinical/ psychiatric traits such as trait anxiety and depression (Zigmond and Snaith, 1983).

Since our research question depends on understanding the hypothesised relationship

between large scale networks (VAN, DMN and DAN), we used the BrainSpace toolbox (Vos

de Wael et al., 2020) to provide whole brain low dimensional representations of functional

brain organisation, generating maps which represent the similarities and differences in the

activity within different systems. We used R version 4.2.0 (R Core Team, 2021) to produce

low dimensional representations of both traits and thoughts using principal component

analysis (PCA). Using these two sets of data, we performed multiple regression to identify how

brain network organisation varies with traits and states. In these analyses, the low dimensional

representations of brain organisation were the dependent measures, and the components of

traits and states were explanatory variables.

Methods

Data

The dataset used here is part of the MPI-Leipzig Mind-Brain-Body (MPILMBB) database

(Mendes et al., 2019). The complete dataset consists of a battery of self-reported personality

measures, measures of spontaneous thought, task data, and structural and resting-state

functional MRI from participants between 20 and 75 years of age. Data were collected over a

period of five days, with the MRI sessions always falling on day 3. The questionnaires were

completed by participants before and after this day, using Limesurvey

(https://www.limesurvey.org: version 2.00+) at their own convenience and using pen-and-

paper on-site. A detailed description of the participants, measures, and data acquisition

protocol has been previously published along with the dataset (Mendes et al., 2019).

Participants

We limited our investigation to personality and thought self-reports, and rs-fMRI from

participants under 50 years of age, who had complete data from at least three resting-state

scans. The resulting sample included 144 participants (74 men, mean age= 26.77 years, SD=

4.03; 70 women, mean age = 26.93 years, SD = 5.55).

<u>Resting state fMRI with Multidimensional Experience Sampling (MDES)</u>

The current sample includes one hour of fully pre-processed rs-fMRI data from 144

participants (four scans from 135 participants, and three scans from nine participants whose

data were missing or incomplete). The rs-fMRI was performed in four adjacent 15-minute

sessions each immediately followed by MDES which retrospectively measured various

dimensions of spontaneous thought during the scan. Images were acquired in axial orientation

using T2∗-weighted gradient-echo echo planar imaging (GE-EPI) sensitive to blood oxygen

level-dependent (BOLD) contrast. Sequences were identical across the four runs, except for

alternating slice orientation and phase-encoding direction, to vary the spatial distribution of

distortions and signal loss. Motion correction parameters were derived by rigid-body

realignment of the time-series to the first (after discarding the first five volumes) volume with

FSL MCFLIRT (Jenkinson et al., 2002). Parameters for distortion correction were calculated

by rigidly registering a temporal mean image of this time series to the fieldmap magnitude

image using FSL FLIRT (Jenkinson and Smith, 2001) which was then unwarped using FSL

FUGUE (Jenkinson et al., 2012). Transformation parameters were derived by coregistering

the unwarped temporal mean to the subject's structural scan using FreeSurfer's boundary-

based registration algorithm (Greve and Fischl, 2009). All three spatial transformations were

then combined and applied to each volume of the original time series in a single interpolation

step. The time series was residualised against the six motion parameters, their first

derivatives,   "outliers"   identified   by   Nipype's   rapidart   algorithm

(https://nipype.readthedocs.io/en/latest/interfaces/ A CompCor (Behzadi et al., 2007)

approach was implemented to remove physiological noise from the residual time-series- which

included first six principal components from all the voxels identified as white-matter

cerebrospinal fluid. The denoised time series were temporally filtered to a frequency range

between 0.01 and 0.1 Hz using FSL, mean centered and variance normalized using Nitime

(Rokem et al., 2009). Imaging and pre-processing protocols are described in detail in Mendes

et al (Mendes et al., 2019).

The MDES battery included 12 statements (Table 2) which participants rated on a visual

analog scale with 5% response increments that go from 0% = "describes my thoughts not at

all" to 100% = "describes my thoughts completely". The current analysis sample includes

MDES data for all available instances of rs-fMRI scans for each participant.

Table 1. Multidimensional Experience Sampling (MDES) Statements

Dimension	Statement
Positive	"My thoughts were positive"
Negative	"My thoughts were negative."
Future	"I thought about future events."
Past	"I thought about past events."
Myself	"I thought about myself."
People	"I thought about other people."
Surroundings	"I thought about my present environment/ surrounding."
Wakeful	"I was completely awake."
Images	"My thoughts were in the form of images."
Words	"My thoughts were in the form of words"
Specific	"My thoughts were more specific than vague"
Intrusive	"My thoughts were intrusive"

196 <u>Personality measures</u>

197 To provide a broad description of individual traits we included data from the following 21

198 questionnaires:

200 **Table 2. List of personality/ dispositional trait questionnaires**

Abbreviation	Behavioural Measure
ACS	Attention Control Scale (Derryberry and Reed, 2002)
ASR	Adult Self Report (Achenbach and Rescorla, 2003)
BDI-II	Beck Depression Inventory -II (Beck et al., 1993)
BIS/BAS	Behavioural Inhibition and Approach System (Carver and White, 1994)
BP	Boredom Proneness Scale (Farmer and Sundberg, 1986)
ESS	Epworth Sleepiness Scale (Johns, 1991)
Gold-MSI	Goldsmiths Musical Sophistication Index (Müllensiefen et al., 2014)

HADS	Hospital Anxiety and Depression Scale (Zigmond and Snaith, 1983)
IAT	Internet Addiction Test (Young, 1998)
IMIS	Involuntary Musical Imagery Scale (Floridou et al., 2015)
MMI	Multimedia Multitasking Index (Ophir et al., 2009)
NEO PI-R	NEO Personality Inventory-Revised (Costa and McCrae, 2008)
PSSI	Personality Style and Disorder Inventory (Kuhl and Kazén, 2009)
SCS	Brief Self-Control Scale(Tangney et al., 2004)
SDS	Social Desirability Scale-17 (Crowne and Marlowe, 1960)
SES	Self-Esteem Scale (O'Malley and Bachman, 1979)
SD3	Short Dark Triad (Jones and Paulhus, 2014)
S-D-MW	Spontaneous and Deliberate Mind-Wandering (Carriere et al., 2013; Golchert et al.,
STAXI	State-Trait Anger Expression Inventory
TPS	Tuckman Procrastination Scale (Tuckman, 2016)
UPPS-P	UPPS-P Impulsive Behaviour Scale (Lynam et al., 2006; Schmidt et al., 2008)

Analyses

Dimension reduction for questionnaire and MDES data

We performed two separate principal component analyses (PCA) to obtain low dimensional summaries of the 71 trait variables from 21 questionnaires, and 12 thought variables from MDES.

71 scores from the personality questionnaires of 144 participants were included and missing data (3.15% of including all variables) were imputed by the variable mean. PCA was performed on this matrix and five "trait" components (henceforth referred to as "traits") were selected on the basis of eigenvalues >1, using the Kaiser-Guttman criterion (Joliffe, 2002) and their congruence with the previously well-established "Big Five" personality traits (Digman, 1990; Cobb-Clark and Schurer, 2012). For the MDES data, separate instances of responding for each participant were concatenated, resulting in a matrix with 576 observations of 12 variables. PCA was performed on this matrix, and five "thought" components (henceforth referred to as "thought patterns") were selected on the basis of eigenvalues >1. Varimax rotation was applied to both solutions to optimize the distinctiveness of each component. The five thought pattern scores were then averaged across the four scans, resulting in one score

for each thought pattern for each participant, describing their location on a particular thought

dimension.

<u>Dimension reduction for whole-brain functional connectivity</u>

Functional time-series for each participant was extracted using the Schaefer 400 parcellation

(Schaefer et al., 2018) using the fully pre-processed data from all four resting-state scans. The

data from separate scans were concatenated, and a 400x400 connectivity matrix was

calculated from the resulting time series for each participant using Pearson correlation. A

group connectivity matrix of the whole sample was calculated by averaging the 144 individual

matrices.

In order to summarize whole brain connectivity in a low-dimensional space, we performed

gradient analysis using the BrainSpace toolbox (Vos de Wael et al., 2020). 10 macro-scale

gradients were calculated for the group (Figure 2). First, we applied fisher's z transform to the

group matrix, building an affinity matrix (kernel= normalized angle, sparsity= 0.9) and then

decomposed it using PCA. We chose the PCA approach for gradient calculation, as Hong and

colleagues (Hong et al., 2020) have shown that compared to non-linear decomposition

methods, PCA provides better reliability and higher phenotypic predictive value for

connectivity gradients. For ease of interpretation and comparability, group gradients were

aligned to a subsample of the HCP dataset (Van Essen et al., 2013) included in BrainSpace.

Finally, following Mckeown et al. (Mckeown et al., 2020) 10 gradients were calculated in order

to maximize the gradient fit for all individuals during alignment. Individual gradients were

calculated for each participant, aligned to the group-level gradients, resulting in a 400x10

matrix for each participant. 10 gradients were calculated to facilitate alignment across

individuals irrespective of differences in rank order of individual gradients (to control for the

possibility, for example, that the pattern summarised by group-level Gradient 2 is Gradient 4

for some participants). Subsequent regression analyses were limited to the first three group-

aligned gradients, which have been relatively well-characterized in previous work (Margulies et al., 2016; Mckeown et al., 2020; Turnbull et al., 2020b). To visualize the functional axis captured by each gradient, we performed Neurosynth (Yarkoni et al., 2011) decoding on the group gradient maps (Figure 2). Further, we calculated the average gradient score for all parcels within each of the seven connectivity networks described by Yeo and colleagues (Yeo et al., 2011) (Figure 2).

Stability of thoughts-patterns and gradients

To quantify the stability of thoughts and connectivity gradients over the whole scanning period, intra-class correlation coefficients (ICC) were calculated for the thought patterns and following Hong and colleagues (Hong et al., 2020), discriminability indices (Bridgeford et al., 2021) were calculated for whole gradients by treating the 4 scans and subsequent thought probes as separate instances. We used the two-way mixed effects model (i.e. type 3 ICC) used for quantifying test-retest reliability, where samples cannot be considered independent (Koo and Li, 2016). Only the 135 participants who had four full-length resting state scans were included in this analysis. As this analysis found reasonable levels of reliability (see below), the averages of the four separate thought scores were used as regressors in subsequent analysis. This allowed for both a robust measure of thought patterns over the whole testing period, and the inclusion of all 144 participants in the analysis.

Multiple Multivariate Regression

To investigate the relationship between individual differences in traits, thoughts, and macro-scale cortical gradients, we used multiple multivariate regression as implemented in the MATLAB SurfStat Toolbox (Worsley et al., 2009) [http://www.math.mcgill.ca/keith/surfstat/]. 400 separate linear models were estimated for 400 parcels, with the gradient scores from the first three gradients as the dependent variables, and with five trait scores (Figure 1A) and five thought scores (Figure 1B), as well as nuisance variables age, motion, and gender included

as independent variables. The resulting significant effects from 400 parcels were corrected for

False Discovery Rate (FDR, q < .05) (Storey, 2003) at the multivariate (three gradients) level.

In order to test which gradient score was affected, follow-up univariate analyses were

performed on the resulting parcels for each gradient separately, and effects were further

Bonferroni corrected (p_{bonf} < .025) for the total number of comparisons performed for all parcels

(including the analyses of all three gradients) for each variable. Additionally, to see how the

trait components related to the thought components, we performed multiple multivariate

regression with the thoughts as dependent variables and traits as independent variables.

Results

Traits and thought patterns

Application of PCA to the battery of personality questionnaires resulted in five "traits" (Figure

1 A, Supplementary Figure 1) with eigenvalues > 1, explaining 48.4% of the variance. The five

trait components, independent of the direction of loadings, largely map onto the "big five"

personality factors: neuroticism, conscientiousness (positive loading on "procrastination" in

our PCA result), extraversion (positive loading on "introversion" in our PCA result),

agreeableness (positive loading on anti-social in our results) and openness to experience,

respectively. Application of PCA to the MDES questions revealed five "thought patterns"

(Figure 1B, Supplementary Figure 2) with eigenvalues>1, explaining 65.4% of the variance.

Based on the most heavily loaded dimensions within each pattern, we named these: "modality"

(image vs words), "positive episodic social", "specific internal", "self-relevant" and

"prospective".

<u>Intraclass correlation – thoughts</u>

Our first analysis established the reliability of thought components across the one hour of

scanning. The five thought patterns showed low to moderate agreement between individual

scores from single sessions (modality = 0.5856, positive episodic-social= 0.4531, specific

internal= 0.5226, self-relevant = 0.5832, prospective= 0.3118), indicating a degree of

variability between sessions. The average of all scores had high ICCs for the first four

components (modality = 0.8497, positive episodic social = 0.7783, specific internal = 0.8141,

self-relevant = 0.8484, prospective = 0.6444). The average scores from 4 sessions were used

as regressors in subsequent analyses.

Next, we examined the relationship between the low dimensional representations of both

personality and thoughts. Multiple multivariate regression using traits as predictor variables of

thought patterns established that "negative affect" had a significant effect on thoughts $(_{5,134})$ =

3.88, p = 0.003, partial η^2 = 0.127). Univariate follow-up showed that a high score on trait

neuroticism was significantly associated with less "positive episodic social" thought (pattern 2;

ß = -0.229, 95% CI = [-0.389 -0.07], p = 0.005, partial η^2 = 0.055) as well as greater "self-

relevant" thought (pattern 4; ß = 0.229, 95% CI = [0.066 0.391], p = 0.006, partial η^2 = 0.053),

(Supplementary Fig. 4).

Figure 1. Principal components of traits and thoughts

A) First five trait components derived from PCA after varimax rotation are represented as

word-clouds with negative loadings shown in cold colours and positive loadings in warm

colours; the relative loading of each variable within a component is represented by the relative

font size (See Supp Figures 1 and 2 for numerical loading values). In the bottom left panel,

scree-plot showing the percentage of trait variance explained by the each of the first 10

components in grey, and the first five components after varimax rotation in red. **B)** Results of the application of PCA to the MDES data, depicted in the same way.

Macro-scale cortical gradients

The first three group level gradients are shown in Figure 2 along with their Neurosynth meta-analytic associations and relationships to the Yeo-networks (Yeo et al., 2011) (7-network solution). The first gradient (G1) differentiates between visual regions at one end and DMN at the other. The second gradient (G2) describes the dissociation between somatomotor and visual cortices. The third gradient (G3) captures the segregation between different transmodal systems (the default mode network versus the fronto-parietal system). The three gradients are largely similar to the ones reported by Margulies et al.(Margulies et al., 2016) and subsequent literature (Hong et al., 2019; Paquola et al., 2019; Bethlehem et al., 2020; Mckeown et al., 2020; Turnbull et al., 2020b). Due to the difference in the decomposition provided by PCA vs. DM approach, the endpoints of G1 are different from those first reported by Margulies et al (Margulies et al., 2016) in that one end is anchored by the visual network alone, as opposed to visual and somatomotor, while in G2, the somatomotor network is separated from both the visual and default mode networks, as opposed to the visual network alone in the (Margulies et al., 2016) study. However, like those reported by Margulies and colleagues, the first two gradients together describe network-level connectivity space anchored at three ends by the visual, somatomotor and default mode network, respectively (Figure 3). Stability of results irrespective of the decomposition approach is shown in supplementary Figure 4. Single gradients tended to be stable over the 4 sessions, with a discriminability index of 0.964 for Gradient 1, 0.918 for Gradient 2, and 0.983 for Gradient 3 over four adjacent scans from 135 participants. Discriminability indices are similar to those previously reported by (Hong et al., 2020).

Figure 2. Group-level gradients of functional connectivity

On the left are the first three group-averaged gradients, represented in left lateral and medial

views. Regions with similar whole-brain connectivity profiles are shown in similar colours, with

yellow and purple regions indicating most dissimilar connectivity patterns. Loading ranges and

directions are arbitrary. In the middle, word clouds representing the top 10 positively (warm

colours) and negatively correlated (cold colours) Neurosynth decoding topic terms for each

gradient map. The relative strength of correlation is represented by the relative font size. On

the right, radar-plots showing the Yeo-network profile of each group-level gradient depicted in

the left column. Each radar-plot shows the mean gradient loadings for all parcels within the

seven Yeo networks.

Figure 3. Comparison of group-level gradients to BrainSpace HCP template

The first scatterplot shows 400 parcel positions along G1 and G2 in the template calculated

from the HCP subsample included in BrainSpace toolbox (Vos de Wael et al., 2020). The

second scatterplot shows parcel positions in the group-level gradients G1 and G2 after

Procrustes alignment to the HCP template. Parcels are color-coded according to their

respective Yeo network. Yeo networks are shown as color-coded brain maps on the right.

Relationship between state- trait variability and cortical gradients

Having established low dimensional representations of thought, personality, and brain

organisation, we next examined associations between different types of personality and

ongoing thought experienced during the scan and our metrics of functional brain organisation.

To this end, we performed a multiple multivariate regression with thoughts, traits, and nuisance

variables (motion, age and gender) as independent variables, with whole brain functional

organisation, as captured by the first three gradients, as dependent variables. In this analytic

approach relationships between cognition along one gradient but not along another help

identify which relationships between brain systems are mostly likely to relate to the feature of

cognition in question (i.e. each gradient acts as a control for the other). In these analyses

both trait "introversion" and a pattern describing "specific internal" thought showed significant

effects at the multivariate level. Results from the univariate follow-up of effects within each

gradient are shown in Figures 4 and 5, and Table 3.

 Trait Introversion

Along the first gradient, a parcel within the right orbitofrontal cortex (within the executive

control network, shown in orange) showed more similarity with transmodal regions for

individuals high on introversion. Six parcels within the ventral attention network, including

anterior insula, operculum and cingulate cortex were closer to the somatomotor end along

gradient two (shown in purple). The same regions showed lower scores along the third

gradient in participants with higher introversion scores, indicating stronger integration with the

default mode network. A parcel within posterior cingulate cortex (control) was also more

segregated from the visual end of gradient two in participants with higher introversion scores.

Associations between trait "introversion" and brain-wide activity are shown in Figure 4.

Figure 4. Relationship between trait "Introversion" on the first three connectivity gradients

On the left, parcels within the first three gradients that show significant ($p_{bonf}<0.025$) differences related to trait "introversion", orange indicating regions within the "frontoparietal control network", and violet indicating regions within the "ventral attention". Scatter plots depict the relationship between individual scores for "introversion" thought (x-axis) and average gradient score of all affected parcels (y-axis) within each gradient. Each datapoint is a participant. Both axes show standardized scores. Detailed results from individual parcels are reported in Table 3. The right column shows Neurosynth decoding of ROI maps of affected parcels within each gradient, showing top ten positively correlated topic terms in warm colours, and top ten negatively associated topic terms in cold colours.

Specific internal cognition

Relationships with patterns of more specific internal cognition were confined to the dorsal attention and visual networks. A region within the superior parietal lobule (DAN) had lower scores on the first gradient (more transmodal) and higher scores on the second gradient (less visual), indicating less similarity with visual regions whose ongoing experience was more "specific" and "internal". Along the third gradient, higher "specific internal" thought scores were associated with greater separation between these regions and the default mode network. Finally, a parcel within the parahippocampal gyrus/ extrastriate (visual network) showed a broad spread along gradient one, with participants with higher "specific internal" thought scores falling on the transmodal/ DMN side and participants with lower scores (higher "surroundings") closer to the visual system side. These findings are shown in Figure 5.

Figure 5. Relationship between specific internal thought and the first three connectivity gradients

On the left, parcels within the first three gradients that show significant differences ($p_{bonf}<0.025$) related to "specific internal" thought, green indicating regions within "dorsal attention network" (DAN), and purple indicating regions within the "visual network". Scatter plots depict the relationship between individual scores for "specific internal" thought (x-axis) and average gradient score of all affected parcels (y-axis) within each gradient. Each datapoint is a participant. Both axes show standardized scores. Detailed results from individual parcels are reported in table 1. The right column shows Neurosynth decoding of ROI maps of affected parcels within each gradient, showing top ten positively correlated topic terms in red, and top ten negatively associated topic terms in blue.

Table 3. Relationships between first three connectivity gradients and introversion and

specific internal thought

IV	DV	Yeo Network	Parcel	$t_{(130)}$	$\eta^2_{partial}$	p_{uncorr}	p_{bonf}
Introversion	G1	Control	OFC (R)	-4.2214	0.11	0.00002	<0.0012
	G2	VAN	A Ins (L)	3.7888	0.10	0.00012	
			A Ins (R)	4.3110	0.12	0.00002	
			Fr Oper (R)	3.8081	0.10	0.00011	
			A Cing (R)	3.1879	0.06	0.00090	
		Control	P Cing (L)	4.0504	0.07	0.00004	
	G3	VAN	A Ins (L)	-3.9539	0.11	0.00006	
			A Ins (L)	-3.6784	0.09	0.00017	
			A Ins (R)	-4.2031	0.13	0.00002	
			Fr Oper (R)	-4.1767	0.12	0.00003	
			A Cing (R)	-3.9732	0.12	0.00006	
Specific Internal Thought	G1	Visual	PHC/ ExStr	-2.9105	0.05	0.00212	<0.0028
		DAN	SPL (L)	-4.5433	0.12	0.00001	
	G2	DAN	SPL (L)	4.1217	0.09	0.00003	
			SPL (R)	3.3542	0.09	0.00052	
	G3	DAN	SPL (L)	4.4548	0.10	0.00001	

IV = Independent Variable, **DV** = Dependent Variable, **G** = Gradient, **VAN** = Ventral Attention Network, **DAN** = Dorsal Attention Network, **OFC** = Orbitofrontal Cortex, **A** = Anterior, **P** = Posterior, **Fr** = Frontal, **Ins** = Insula, **L**= Left, **R**= Right, **Oper** = Operculum, **Cing** = Cingulate Cortex, **PHC** = Parahippocampal Cortex, **ExStr** = Extrastriate cortex, **SPL** = Superior Parietal Lobule. Results reported in the table are from univariate (single-gradient) follow-up tests for parcels showing a significant effect for each IV at the multivariate (3-gradient) level. Univariate tests are Bonferroni corrected for the total number of parcels (all 3 gradients) where tests were performed (21 parcels for Introversion, 9 for Specific Internal Thought).

Discussion

Our study investigated whether an emerging "tri-partite" perspective from contemporary views of ongoing conscious thought can provide a framework that can account for the relationship between dispositional traits, self-reports of ongoing thought, and individual differences in large scale patterns of neural connectivity at rest. This tri-partite network account emphasises the roles of regions embedded within three large scale networks as important for ongoing experience: the ventral attention network, the dorsal attention network and the default mode network (Huang et al., 2021; Smallwood et al., 2021b). We calculated macro-scale connectivity gradients from one hour of resting state fMRI for 144 participants. The variability of these gradients was then analysed as a function of self-reports of ongoing thought patterns (captured by the principal components of MDES) and personality traits (described by the principal components of a battery of personality and habit measures). Given the tendency of certain traits to be correlated with frequency of specific patterns of thought (e.g. depression level with intrusive and negative thought (Konu et al., 2021) we also looked for possible dependencies between trait and thought components through multivariate regression using traits as predictors.

Our analyses confirmed that both patterns of thought, and indices of traits, contribute to patterns of brain organisation in a manner that converges with the emerging tri-partite view of ongoing conscious thought. For example, it has been hypothesised that the ventral attention network (VAN) helps adjudicate between internal and external influences on ongoing thought (Huang et al., 2021; Smallwood et al., 2021b) and we found that individuals who were high on dispositional "introversion" showed variation in anterior insula, overlying operculum and ACC: all regions making up part of the VAN. For more introverted people these regions showed greater alignment with somatomotor regions and less with visual cortex (gradient 2), and greater alignment with default mode network than the fronto-parietal network (gradient 3). Notably, prior studies have found that regions of sensorimotor cortex are linked to deliberate

mind-wandering (Golchert et al., 2017) and individuals who tend to generate patterns of episodic social cognition during periods of low task demands show greater temporal correlation between the VAN and sensori-motor cortex (Turnbull et al., 2019a). Introversion reflects a predisposition towards internal subjective states rather than external objects (Jung, 1995), and so our analysis adds to an emerging literature (Golchert et al., 2017; Turnbull et al., 2019a) that suggests that there may be an important relationship between sensorimotor cortex and the ventral attention network in patterns of internal self-generated thought.

Our study also highlighted that associations between observed patterns of neural organisation at rest and patterns of ongoing thought are consistent with the hypothesized tripartite view (Huang et al., 2021; Smallwood et al., 2021b). For example, we found a pattern of detailed thinking during wakeful rest that was correlated with stronger decoupling between a region within dorsal parietal cortex from the visual network (as indexed by changes in both gradients 1 and 2), and greater separation from the default mode network (gradient 3). This region overlaps with a region within the DAN identified by (Turnbull et al., 2019b) in which brain activity was reduced during self-generated thoughts relative to external task focus, suggesting an important role for the DAN in external facing attention. Furthermore, using a technique known as "echoes" analysis (see Leech et al., 2012), Turnbull and colleagues (Turnbull et al., 2019b) established that individuals who engaged in self-generated thought during situations of low external demand, at rest showed greater separation between the dorsal attention network and lateral regions of the default mode network in a region of the dLPFC, also a member of the VAN. Thus, our study confirms prior studies that highlight that greater segregation between the DAN and the DMN in the capacity to engage in thoughts that are less linked to the external environment. Together, the convergence between the current analysis and perspectives from research on conscious experience highlight a high degree of overlap in both the regions identified and the hypothesised functions. Our observations are important, therefore, because they help establish that with the appropriate methodology (Mckeown et al., 2020; Finn, 2021; Gonzalez-Castillo et al., 2021) neural accounts of

conscious experience (Huang et al., 2021; Smallwood et al., 2021b) provide an important valuable way to make sense of brain-cognition links observed at rest. Second, our data provides evidence for the "decoupling" hypothesis of self-generated experience (Smallwood, 2013a). This perspective emerged from observations that cortical processing of external inputs is reduced when individuals focus on internal self-generated thought (Smallwood et al., 2008; Kam et al., 2011; Baird et al., 2014), and assumes that this reduced processing of external input allows an internal train of thought to persist in a more detailed manner (p.524,Smallwood, 2013 Smallwood, 2013b)). Our data is consistent with this view since both personality traits linked to internal focus ("introversion"), and patterns of detail experience that are not directed externally ("specific internal thought"), are linked to reductions in the similarity between neural activity in regions linked to attention and cognition, with regions of visual cortex. In this way our study provides novel insight into how the macro-scale functional patterns across the cortex support the emergence of detailed patterns of internal experience. Critically, in our study there was no external task from which thinking needed to be decoupled from, ruling out accounts of this process as a "lapse" in the normal upregulation of task relevant material needed for task completion (for discussion see Franklin et al., 2013, Smallwood 2013ab).

Although our study establishes how contemporary work on conscious experience can help understand patterns of brain organisation observed at rest and highlights how these approaches can be leveraged to understand the neural correlates of both an individuals' traits, and their thoughts, there are nonetheless important questions that our study leaves open. For example, contemporary work on ongoing conscious thought highlights time and context as key variables necessary for understanding the neural correlates of different features of thinking (Smallwood et al., 2021b). Since the aim of our study was to focus on the brain at rest, interpretations of our results should bear in mind that under different task conditions, neural correlates between thinking, personality and neural activity may be different. For example, prior studies have established that posterior elements of the default mode network can

become integrated into task positive systems (Krieger-Redwood et al., 2016; Vatansever et al., 2017) and under demanding task conditions the default mode network is linked to patterns of task focused cognition (Sormaz et al., 2018). Similarly, our analysis focused on 'static' indices of neural activity rather than dynamic measures. We chose to focus on static indices of neural activity because our prior studies have shown that brain-behaviour correlations can be relatively stable over time (Wang et al., 2020), and as we establish in our study these patterns show reasonable stability across a one hour session. Furthermore, our study, particularly with respect to the findings relating to DAN, map closely onto studies that use experience sampling to identify momentary correlations between neural activity and experience (Turnbull et al., 2019b, 2019a, 2020b). Nonetheless, there are likely to be important dynamic features of ongoing experience that our analysis of static brain organisation cannot capture (Kucyi, 2018). Recent discussions in cognitive neuroscience have highlighted the link between sample size and the reproducibility of brain-wide associations with behavioural phenotypes (Marek et al., 2022; Spisak et al., 2023). Our analysis of the trait and thought data alone revealed that "neuroticism" was related to high negative and episodic thoughts, however, we did not find any other significant relationships among traits and thought patterns. In the current data, neuroticism was the most prominent out of all five traits included in the analysis, accounting for 29% of the total variance explained by them. It is therefore likely that more extensive and robust correlations between thoughts patterns and other traits, as well as thoughts, traits and macroscale connectivity patterns would emerge with data sets with larger sample sizes. In the future it may also be important to consider measures of traits that could have relationships to both neural activity and or experience at rest (e.g. self-consciousness de Caso et al., 2017, or autistic tendencies, Turnbull et al., 2020a). It is worth noting that mapping momentary changes between ongoing experience and neural activity will likely depend on a data set tailored to this question, in particular in which (i) experience sampling measures are collected more frequently as well as (ii) methodological advances that allow patterns of activity to be mapped without using temporal correlation. and that measure thinking across multiple contexts.

References

Achenbach TM, Rescorla LA (2003) No TitleManual for the ASEBA adult forms & profiles. Research Center for Children, Youth, & Families. University of Vermont, Burlington, VT, USA.

Baird B, Smallwood J, Lutz A, Schooler JW (2014) The Decoupled Mind: Mind-wandering Disrupts Cortical Phase-locking to Perceptual Events. J Cogn Neurosci 26:2596–2607.

Beck A, Steer R, Brown G (1993) Beck Depression Inventory-II (BDI-II). … Beck Depress Invent.

Behzadi Y, Restom K, Liau J, Liu TT (2007) A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. Neuroimage 37:90–101 Available at: /pmc/articles/PMC2214855/ [Accessed July 5, 2021].

Bethlehem RAI, Paquola C, Seidlitz J, Ronan L, Bernhardt B, Consortium C-CCAN, Tsvetanov KA (2020) Dispersion of functional gradients across the adult lifespan. Neuroimage 222:117299 Available at: https://linkinghub.elsevier.com/retrieve/pii/S1053811920307850 [Accessed November 3, 2021].

Biswal B, Zerrin Yetkin F, Haughton VM, Hyde JS (1995) Functional connectivity in the motor cortex of resting human brain using echo-planar mri. Magn Reson Med 34:537–541 Available at: https://onlinelibrary.wiley.com/doi/full/10.1002/mrm.1910340409 [Accessed April 11, 2024].

Bridgeford EW, Wang S, Yang Z, Wang Z, Xu T, Craddock C, Dey J, Kiar G, Gray-Roncal W, Colantuoni C, Douville C, Noble S, Priebe CE, Caffo B, Milham M, Zuo X-N, Reproducibility C for R and, Vogelstein JT (2021) Eliminating accidental deviations to minimize generalization error and maximize replicability: applications in connectomics and genomics. bioRxiv:802629 Available at: https://www.biorxiv.org/content/10.1101/802629v8 [Accessed September 24, 2021].

Cao H et al. (2018) Cerebello-thalamo-cortical hyperconnectivity as a state-independent functional neural signature for psychosis prediction and characterization. Nat Commun 9 Available at: /pmc/articles/PMC6155100/ [Accessed March 22, 2022].

Carriere JSA, Seli P, Smilek D (2013) Wandering in both mind and body: Individual differences in mind wandering and inattention predict fidgeting. Can J Exp Psychol Can Psychol expérimentale 67:19–31 Available at: http://doi.apa.org/getdoi.cfm?doi=10.1037/a0031438 [Accessed April 4, 2019].

Carver CS, White TL (1994) Behavioral Inhibition , Behavioral Activation , and Affective Responses to Impending Reward and Punishment : The BIS / BAS Scales. J Pers Soc Psychol 67:319–333.

Cobb-Clark DA, Schurer S (2012) The stability of big-five personality traits. Econ Lett 115:11–15.

Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the brain. Nat Rev Neurosci 3:201–215 Available at: https://pubmed.ncbi.nlm.nih.gov/11994752/ [Accessed March 21, 2022].

Costa PT, McCrae RR (2008) The revised NEO personality inventory (NEO-PI-R). SAGE Handb Personal Theory Assess Vol 2 - Personal Meas Test:179–198.

Crowne DP, Marlowe D (1960) A new scale of social desirability independent of

psychopathology. J Consult Psychol 24:349–354.

Cui Z et al. (2020) Individual Variation in Functional Topography of Association Networks in
Youth. Neuron 106:340-353.e8.

de Caso I, Poerio G, Jefferies E, Smallwood J (2017) That's me in the spotlight: neural basis
of individual differences in self-consciousness. Soc Cogn Affect Neurosci 12:1384
Available at: /pmc/articles/PMC5629813/ [Accessed April 11, 2024].

Derryberry D, Reed MA (2002) Anxiety-related attentional biases and their regulation by
attentional control. J Abnorm Psychol 111:225–236 Available at: /record/2002-12652-
002 [Accessed September 24, 2021].

Digman JM (1990) PERSONALITY STRUCTURE: EMERGENCE OF THE FIVE-FACTOR
MODEL. Annu Rev Psychol 41:417–457 Available at: www.annualreviews.org
[Accessed October 10, 2023].

Dosenbach NUF, Nardos B, Cohen AL, Fair DA, Power JD, Church JA, Nelson SM, Wig GS,
Vogel AC, Lessov-Schlaggar CN, Barnes KA, Dubis JW, Feczko E, Coalson RS, Pruett
JR, Barch DM, Petersen SE, Schlaggar BL (2010) Prediction of individual brain maturity
using fMRI. Science (80-) 329:1358–1361 Available at:
https://www.science.org/doi/abs/10.1126/science.1194144 [Accessed March 21, 2022].

Farmer R, Sundberg ND (1986) Boredom proneness--the development and correlates of a
new scale. J Pers Assess 50:4–17 Available at:
https://europepmc.org/article/MED/3723312 [Accessed September 24, 2021].

Finn ES (2021) Is it time to put rest to rest? Trends Cogn Sci Available at:
https://linkinghub.elsevier.com/retrieve/pii/S1364661321002345 [Accessed October 14,
2021].

Floridou GA, Williamson VJ, Stewart L, Müllensiefen D (2015) The Involuntary Musical
Imagery Scale (IMIS). Psychomusicology Music Mind, Brain 25:28–36 Available at:
/record/2015-22418-001 [Accessed September 24, 2021].

Fox MD, Raichle ME (2007) Spontaneous fluctuations in brain activity observed with
functional magnetic resonance imaging. Nat Rev Neurosci 2007 89 8:700–711
Available at: https://www.nature.com/articles/nrn2201 [Accessed April 11, 2024].

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005) The human
brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc
Natl Acad Sci U S A 102:9673–9678 Available at:
https://www.pnas.org/doi/abs/10.1073/pnas.0504136102 [Accessed April 11, 2024].

Golchert J, Smallwood J, Jefferies E, Seli P, Huntenburg JM, Liem F, Lauckner ME,
Oligschläger S, Bernhardt BC, Villringer A, Margulies DS (2017) Individual variation in
intentionality in the mind-wandering state is reflected in the integration of the default-
mode, fronto-parietal, and limbic networks. Neuroimage 146:226–235 Available at:
https://www.sciencedirect.com/science/article/pii/S1053811916306425?via%3Dihub
[Accessed July 31, 2018].

Gonzalez-Castillo J, Kam JWYY, Hoy CW, Bandettini PA (2021) How to Interpret Resting-
State fMRI: Ask Your Participants. J Neurosci 41:1130–1141 Available at:
https://doi.org/10.1523/JNEUROSCI.1786-20.2020 [Accessed July 19, 2021].

Gratton C, Smith DM, Dorn M (2020) Digging Deeper to Chart the Landscape of Human
Brain Development. Neuron 106:209–211 Available at:
https://doi.org/10.1016/j.neuron.2020.03.030 [Accessed February 15, 2022].

Greve DN, Fischl B (2009) Accurate and robust brain image alignment using boundary-

based registration. Neuroimage 48:63–72.

Hong SJ, de Wael RV, Bethlehem RAI, Lariviere S, Paquola C, Valk SL, Milham MP, Di Martino A, Margulies DS, Smallwood J, Bernhardt BC (2019) Atypical functional connectome hierarchy in autism. Nat Commun 10.

Hong SJ, Xu T, Nikolaidis A, Smallwood J, Margulies DS, Bernhardt B, Vogelstein J, Milham MP (2020) Toward a connectivity gradient-based framework for reproducible biomarker discovery. Neuroimage 223:117322.

Hsu WT, Rosenberg MD, Scheinost D, Constable RT, Chun MM (2018) Resting-state functional connectivity predicts neuroticism and extraversion in novel individuals. Soc Cogn Affect Neurosci 13:224–232 Available at: https://academic.oup.com/scan/article/13/2/224/4823668 [Accessed March 22, 2022].

Huang Z, Tarnal V, Vlisides PE, Janke EL, McKinney AM, Picton P, Mashour GA, Hudetz AG (2021) Anterior insula regulates brain network transitions that gate conscious access. CellReports 35:109081 Available at: https://doi.org/10.1016/j.celrep.2021.109081 [Accessed May 27, 2021].

Jenkinson M, Bannister P, Brady M, Smith S (2002) Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. Neuroimage 17:825–841.

Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM (2012) FSL. Neuroimage 62:782–790 Available at: https://linkinghub.elsevier.com/retrieve/pii/S1053811911010603 [Accessed April 16, 2024].

Jenkinson M, Smith S (2001) A global optimisation method for robust affine registration of brain images. Med Image Anal 5:143–156.

Johns MW (1991) A new method for measuring daytime sleepiness: the Epworth sleepiness scale. Sleep 14:540–545 Available at: https://europepmc.org/article/MED/1798888 [Accessed September 24, 2021].

Joliffe IT (2002) Principal Component Analysis. New York: Springer-Verlag. Available at: http://link.springer.com/10.1007/b98835 [Accessed February 15, 2022].

Jones DN, Paulhus DL (2014) Introducing the Short Dark Triad (SD3): A Brief Measure of Dark Personality Traits. Assessment 21:28–41 Available at: /record/2014-10974-004 [Accessed September 24, 2021].

Jung CG (1995) Memories, Dreams, Reflections. In: Memories, Dreams, Reflections, pp 414–415. London: Fontana Press.

Kam JWY, Dao E, Farley J, Kevin F, Smallwood J, Schooler JW, Handy TC (2011) Slow fluctuations in attentional control of sensory cortex. J Cogn Neurosci 23:460–470 Available at: https://pubmed.ncbi.nlm.nih.gov/20146593/ [Accessed October 22, 2021].

Karapanagiotidis T, Jefferies E, Smallwood J (2021) Interactions between the neural correlates of dispositional internally directed thought and visual imagery. Philos Trans R Soc B 376:20190691 Available at: https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2019.0691 [Accessed August 11, 2021].

Karapanagiotidis T, Vidaurre D, Quinn AJ, Vatansever D, Poerio GL, Turnbull A, Ho NSP, Leech R, Bernhardt BC, Jefferies E, Margulies DS, Nichols TE, Woolrich MW, Smallwood J (2020) The psychological correlates of distinct neural states occurring during wakeful rest. Sci Rep 10 Available at: /pmc/articles/PMC7712889/ [Accessed

August 11, 2021].

Karapanagiotidis T, Vidaurre D, Quinn AJ, Vatansever D, Poerio GL, Turnbull A, Leech R, Bernhardt B, Jefferies E, Margulies DS, Nichols TE, Woolrich MW, Smallwood J (2019) Emergence of neural dynamics within a co-ordinate space of large-scale neural hierarchies. bioRxiv:2019.12.21.885772 Available at: https://www.biorxiv.org/content/10.1101/2019.12.21.885772v1 [Accessed January 20, 2020].

Koban L, Gianaros PJ, Kober H, Wager TD (2021) The self in context: brain systems linking mental and physical health. Nat Rev Neurosci 2021 225 22:309–322 Available at: https://www.nature.com/articles/s41583-021-00446-8 [Accessed March 10, 2022].

Konu D, Mckeown B, Turnbull A, Siu Ping Ho N, Karapanagiotidis T, Vanderwal T, McCall C, Tipper SP, Jefferies E, Smallwood J (2021) Exploring patterns of ongoing thought under naturalistic and conventional task-based conditions. Conscious Cogn 93:103139.

Koo TK, Li MY (2016) A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. J Chiropr Med 15:155 Available at: /pmc/articles/PMC4913118/ [Accessed August 11, 2021].

Krieger-Redwood K, Jefferies E, Karapanagiotidis T, Seymour R, Nunes A, Ang JWA, Majernikova V, Mollo G, Smallwood J (2016) Down but not out in posterior cingulate cortex: Deactivation yet functional coupling with prefrontal cortex during demanding semantic cognition. Neuroimage 141:366–377 Available at: https://pubmed.ncbi.nlm.nih.gov/27485753/ [Accessed September 24, 2021].

Kucyi A (2018) Just a thought: How mind-wandering is represented in dynamic brain connectivity. Neuroimage 180:505–514.

Kucyi A, Esterman M, Capella J, Green A, Uchida M, Biederman J, Gabrieli JDE, Valera EM, Whitfield-Gabrieli S (2021) Prediction of stimulus-independent and task-unrelated thought from functional brain networks. Nat Commun 12:1–17 Available at: https://doi.org/10.1038/s41467-021-22027-0 [Accessed May 13, 2021].

Kuhl J, Kazén M (2009) Persönlichkeits-Stil-und-Störungs-Inventar (PSSI): Manual (Hogrefe, ed).

Leech R, Braga R, Sharp DJ (2012) Echoes of the Brain within the Posterior Cingulate Cortex. J Neurosci 32:215–222 Available at: https://www.jneurosci.org/content/32/1/215 [Accessed September 24, 2021].

Lynam DR, Smith GT, Whiteside SP, Cyders MA (2006) The UPPS-P: Assessing five personality pathways to impulsive behavior. West Lafayette, Purdue Univ 10.

Marek S et al. (2022) Reproducible brain-wide association studies require thousands of individuals. Nature 603:654 Available at: /pmc/articles/PMC8991999/ [Accessed December 14, 2022].

Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G, Bezgin G, Eickhoff SB, Castellanos FX, Petrides M, Jefferies E, Smallwood J (2016) Situating the default-mode network along a principal gradient of macroscale cortical organization. Proc Natl Acad Sci U S A 113:12574–12579 Available at: http://www.ncbi.nlm.nih.gov/pubmed/27791099 [Accessed December 11, 2019].

Mckeown B, Strawson WH, Wang HT, Karapanagiotidis T, Vos de Wael R, Benkarim O, Turnbull A, Margulies D, Jefferies E, McCall C, Bernhardt B, Smallwood J (2020) The relationship between individual variation in macroscale functional gradients and distinct aspects of ongoing thought. Neuroimage 220:117072.

Mendes N et al. (2019) A functional connectome phenotyping dataset including cognitive state and personality measures. Sci Data 6:180307 Available at: http://www.nature.com/articles/sdata2018307 [Accessed April 4, 2019].

Menon V (2011) Large-scale brain networks and psychopathology: a unifying triple network model. Trends Cogn Sci 15:483–506 Available at: https://pubmed-ncbi-nlm-nih-gov.browser.cbs.mpg.de/21908230/ [Accessed March 10, 2022].

Müllensiefen D, Gingras B, Musil J, Stewart L (2014) Measuring the facets of musicality: The Goldsmiths Musical Sophistication Index (Gold-MSI). Pers Individ Dif 60:S35.

O'Malley PM, Bachman JG (1979) Self-esteem and education: Sex and cohort comparisons among high school seniors. J Pers Soc Psychol 37:1153–1159 Available at: /record/1980-33361-001 [Accessed September 24, 2021].

Ophir E, Nass C, Wagner AD (2009) Cognitive control in media multitaskers. Proc Natl Acad Sci 106:15583–15587 Available at: https://www.pnas.org/content/106/37/15583 [Accessed September 24, 2021].

Paquola C, Vos De Wael R, Wagstyl K, Bethlehem RAI, Hong S-J, Seidlitz J, Bullmore ET, Evans AC, Misic B, Margulies DS, Smallwood J, Bernhardt BC (2019) Microstructural and functional gradients are increasingly dissociated in transmodal cortices. Kennedy H, ed. PLoS Biol 17:e3000284 Available at: http://dx.plos.org/10.1371/journal.pbio.3000284 [Accessed November 12, 2019].

Poerio GL, Sormaz M, Wang H-T, Margulies D, Jefferies E, Smallwood J (2017) The role of the default mode network in component processes underlying the wandering mind. Soc Cogn Affect Neurosci 12:1047–1062 Available at: http://www.ncbi.nlm.nih.gov/pubmed/28402561 [Accessed July 31, 2018].

R Core Team (2021) R: A language and environment for statistical computing. Available at: https://www.r-project.org/.

Rokem A, Trumpis M, Pérez F (2009) In Proceedings of the 8th Python in Science Conference (SciPy 2009). In: SciPy Conference, pp 68–75.

Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo X-N, Holmes AJ, Eickhoff SB, Yeo BTT (2018) Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. Cereb Cortex 28:3095–3114 Available at: https://academic-oup-com.browser.cbs.mpg.de/cercor/article/28/9/3095/3978804 [Accessed August 11, 2021].

Schmidt RE, Gay P, D'Acremont M, Van Der Linden M (2008) A German adaptation of the upps impulsive behavior scale: Psychometric properties and factor structure. Swiss J Psychol 67:107–112 Available at: http://psycontent.metapress.com/openurl.asp?genre=article&id=doi:10.1024/1421-0185.67.2.107 [Accessed June 5, 2014].

Smallwood J (2013a) Distinguishing how from why the mind wanders: A process-occurrence framework for self-generated mental activity. Psychol Bull 139:519–535.

Smallwood J (2013b) Searching for the elements of thought: Reply to franklin, mrazek, broadway, and schooler (2013). Psychol Bull 139:542–547.

Smallwood J, Beach E, Schooler JW, Handy TC (2008) Going AWOL in the Brain: Mind Wandering Reduces Cortical Analysis of External Events. J Cogn Neurosci 20:458–469.

Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS (2021a) The default mode network in cognition: a topographical perspective. Nat Rev Neurosci

2021:1–11 Available at: https://www-nature-com.browser.cbs.mpg.de/articles/s41583-
021-00474-4 [Accessed July 7, 2021].

Smallwood J, Karapanagiotidis T, Ruby F, Medea B, de Caso I, Konishi M, Wang H-T,
Hallam G, Margulies DS, Jefferies E (2016) Representing Representation: Integration
between the Temporal Lobe and the Posterior Cingulate Influences the Content and
Form of Spontaneous Thought Biagini G, ed. PLoS One 11:e0152272 Available at:
http://dx.plos.org/10.1371/journal.pone.0152272 [Accessed May 28, 2018].

Smallwood J, Turnbull A, Wang H ting, Ho NSP, Poerio GL, Karapanagiotidis T, Konu D,
Mckeown B, Zhang M, Murphy C, Vatansever D, Bzdok D, Konishi M, Leech R, Seli P,
Schooler JW, Bernhardt B, Margulies DS, Jefferies E (2021b) The neural correlates of
ongoing conscious thought. iScience 24:102132.

Sormaz M, Murphy C, Wang H, Hymers M, Karapanagiotidis T, Poerio G, Margulies DS,
Jefferies E, Smallwood J (2018) Default mode network can support the level of detail in
experience during active task states. Proc Natl Acad Sci U S A 115:9318–9323
Available at: http://www.ncbi.nlm.nih.gov/pubmed/30150393 [Accessed April 17, 2019].

Spisak T, Bingel U, Wager TD (2023) Multivariate BWAS can be replicable with moderate
sample sizes. Nature 615:E4–E7 Available at: https://www.nature.com/articles/s41586-
023-05745-x [Accessed March 31, 2023].

Storey JD (2003) The positive false discovery rate: a Bayesian interpretation and the q-
value. https://doi.org/101214/aos/1074290335 31:2013–2035 Available at:
https://projecteuclid.org/journals/annals-of-statistics/volume-31/issue-6/The-positive-
false-discovery-rate--a-Bayesian-interpretation-and/10.1214/aos/1074290335.full
[Accessed November 8, 2021].

Tangney JP, Baumeister RF, Boone AL (2004) High self-control predicts good adjustment,
less pathology, better grades, and interpersonal success. J Pers 72:271–324 Available
at: https://pubmed.ncbi.nlm.nih.gov/15016066/ [Accessed September 24, 2021].

Tuckman BW (2016) The Development and Concurrent Validity of the Procrastination Scale:
http://dx.doi.org/101177/0013164491512022 51:473–480 Available at:
https://journals.sagepub.com/doi/10.1177/0013164491512022 [Accessed September
24, 2021].

Turnbull A, Garfinkel SN, Ho NSP, Critchley HD, Bernhardt BC, Jefferies E, Smallwood J
(2020a) Word up – Experiential and neurocognitive evidence for associations between
autistic symptomology and a preference for thinking in the form of words. Cortex
128:88–106.

Turnbull A, Karapanagiotidis T, Wang HT, Bernhardt BC, Leech R, Margulies D, Schooler J,
Jefferies E, Smallwood J (2020b) Reductions in task positive neural systems occur with
the passage of time and are associated with changes in ongoing thought. Sci Rep 10
Available at: /pmc/articles/PMC7303126/?report=abstract [Accessed November 11,
2020].

Turnbull A, Wang H-T, Schooler JW, Jefferies E, Margulies DS, Smallwood J (2019a) The
ebb and flow of attention: Between-subject variation in intrinsic connectivity and
cognition associated with the dynamics of ongoing experience. Neuroimage 185:286–
299 Available at:
https://www.sciencedirect.com/science/article/pii/S1053811918319414?via%3Dihub
[Accessed January 18, 2019].

Turnbull A, Wang H, Murphy C, Ho NSP, Wang X, Sormaz M, Karapanagiotidis T, Leech
RM, Bernhardt B, Margulies DS, Vatansever D, Jefferies E, Smallwood J (2019b) Left
dorsolateral prefrontal cortex supports context-dependent prioritisation of off-task

thought. Nat Commun 10:3816 Available at: http://www.nature.com/articles/s41467-
019-11764-y [Accessed November 12, 2019].

Van Essen DC, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K, Consortium for
the W-MH (2013) The WU-Minn Human Connectome Project: An Overview.
Neuroimage 80:62 Available at: /pmc/articles/PMC3724347/ [Accessed August 12,
2021].

Vanhaudenhuyse A, Demertzi A, Schabus M, Noirhomme Q, Bredart S, Boly M, Phillips C,
Soddu A, Luxen A, Moonen G, Laureys S (2011) Two distinct neuronal networks
mediate the awareness of environment and of self. J Cogn Neurosci 23:570–578
Available at: https://pubmed.ncbi.nlm.nih.gov/20515407/ [Accessed April 11, 2024].

Vatansever D, Menon DK, Stamatakis EA (2017) Default mode contributions to automated
information processing. Proc Natl Acad Sci U S A 114:12821–12826 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/29078345 [Accessed April 29, 2019].

Vos de Wael R, Benkarim O, Paquola C, Lariviere S, Royer J, Tavakol S, Xu T, Hong S-J,
Langs G, Valk S, Misic B, Milham M, Margulies D, Smallwood J, Bernhardt BC (2020)
BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and
connectomics datasets. Commun Biol 2020 31 3:1–10 Available at:
https://www.nature.com/articles/s42003-020-0794-7 [Accessed November 3, 2021].

Wang H-T, Ho NSP, Bzdok D, Bernhardt BC, Margulies DS, Jefferies E, Smallwood J (2020)
Neurocognitive patterns dissociating semantic processing from executive control are
linked to more detailed off-task mental time travel. Sci Reports 2020 101 10:1–14
Available at: https://www.nature.com/articles/s41598-020-67605-2 [Accessed
September 24, 2021].

Wang H-TT, Poerio G, Murphy C, Bzdok D, Jefferies E, Smallwood J (2018) Dimensions of
Experience: Exploring the Heterogeneity of the Wandering Mind. Psychol Sci 29:56–71
Available at: http://www.ncbi.nlm.nih.gov/pubmed/29131720 [Accessed July 31, 2018].

Wen X, He H, Dong L, Chen J, Yang J, Guo H, Luo C, Yao D (2020) Alterations of local
functional connectivity in lifespan: A resting-state fMRI study. Brain Behav 10:e01652
Available at: https://onlinelibrary.wiley.com/doi/full/10.1002/brb3.1652 [Accessed March
21, 2022].

Worsley KJ, Taylor JE, Carbonell F, Chung MK, Duerden E, Bernhardt B, Lyttelton O,
Boucher M, Evans AC (2009) SurfStat: A Matlab toolbox for the statistical analysis of
univariate and multivariate surface and volumetric data using linear mixed effects
models and random field theory. Neuroimage 47 Available at:
http://www.stat.uchicago.edu/~worsley/surfstat/poster.htm [Accessed August 11, 2021].

Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD, T Y, RA P, TE N, DC VE,
TD W (2011) Large-scale automated synthesis of human functional neuroimaging data.
8:665–670 Available at: https://pubmed-ncbi-nlm-nih-
gov.browser.cbs.mpg.de/21706013/ [Accessed August 11, 2021].

Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, Roffman JL,
Smoller JW, Zöllei L, Polimeni JR, Fisch B, Liu H, Buckner RL (2011) The organization
of the human cerebral cortex estimated by intrinsic functional connectivity. J
Neurophysiol 106:1125–1165 Available at:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3174820/ [Accessed March 30, 2021].

Young KS (1998) Internet Addiction: The Emergence of a New Clinical Disorder.
CyberPsychology Behav 1:237–244 Available at:
https://www.liebertpub.com/doi/abs/10.1089/cpb.1998.1.237 [Accessed September 24,
2021].

Zhang J, Kucyi A, Raya J, Nielsen AN, Nomi JS, Damoiseaux JS, Greene DJ, Horovitz SG,
Uddin LQ, Whitfield-Gabrieli S (2021) What have we really learned from functional
connectivity in clinical populations? Neuroimage 242:118466 Available at:
https://linkinghub.elsevier.com/retrieve/pii/S1053811921007394 [Accessed August 25,
2021].

Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. Acta Psychiatr
Scand 67:361–370 Available at: https://pubmed.ncbi.nlm.nih.gov/6880820/ [Accessed
September 24, 2021].

Supplementary Figures

Supplementary Figure 1. Heatmap showing variable component loadings for the first 5 principal components derived from trait questionnaires

Supplementary Figure 2. Heatmap showing variable component loadings for the first 5 principal components derived from MDES

882 **Supplementary figure 3. Ten group-level cortical gradients shown from the left lateral**

883 **and medial views.**

886 **Supplementary Figure 4. Scatter-plots showing the relationship between trait "Negative**

887 **affect", and "Positive Episodic Social" and "Self-relevant" thought.**