Phylogenetic and biogeographical traits predict unrecognized hosts of zoonotic

leishmaniasis

Caroline K. Glidden[1], Aisling Murran[1], Rafaella Albuquerque Silva[2], Adrian A. Castellanos[3],

Barbara A. Han[3], Erin Mordecai[1]

[1]Department of Biology, Stanford University, Stanford, California, USA 94305

[2] Secretaria de Vigilância em Saùde, Ministério da Saúde do Brasil, Distrito Federal, Brasília

[3]Cary Institute of Ecosystem Studies, Millbrook, New York, USA 12545

Abstract

The spatio-temporal distribution of leishmaniasis, a parasitic vector-borne zoonotic disease, is

significantly impacted by land-use change and climate warming in the Americas. However,

predicting and containing outbreaks is challenging as the zoonotic *Leishmania* system is highly

complex: leishmaniasis (visceral, cutaneous and muco-cutaneous) in humans is caused by up to

14 different *Leishmania* species, and the parasite is transmitted by dozens of sandfly species and

is known to infect almost twice as many wildlife species. Despite the already broad known host

range, new hosts are discovered almost annually and *Leishmania* transmission to humans occurs

in absence of a known host. As such, the full range of *Leishmania* hosts is undetermined,

inhibiting the use of ecological interventions to limit pathogen spread and the ability to accurately

predicting the impact of global change on disease risk. Here, we employed a machine learning

approach to generate trait profiles of known zoonotic *Leishmania* wildlife hosts (mammals that

are naturally exposed and susceptible to infection) and used trait-profiles of known hosts to

identify potentially unrecognized hosts. We found that biogeography, phylogenetic distance, and

study effort best predicted *Leishmania* host status. Traits associated with global change, such as

agricultural land-cover, urban land-cover, and climate, were among the top predictors of host

status. Most notably, our analysis suggested that zoonotic *Leishmania* hosts are significantly

undersampled, as our model predicted just as many unrecognized hosts as unknown hosts.

Overall, our analysis facilitates targeted surveillance strategies and improved understanding of

the impact of environmental change on local transmission cycles.

Author summary

Leishmaniasis is a zoonotic, vector borne disease of poverty with a high burden throughout the

Americas: within Latin America there are an estimated 58,500 new cases per year and 54,050

years of life lost due to disability. Although the World Health Organization has targeted

leishmaniasis for elimination and control by 2030, the disease remains a persistent threat. Across

the Americas, particularly in Central America, the southeastern United States, and perimeters of

the Amazon Basin, risk of infection is increasing in geographic extent and elevation. While it is

known that *Leishmania* parasites, the causative agent of leishmaniasis, are maintained in the

environment via a mammalian host, the full suite of wildlife hosts has yet to be documented, which

significantly hinders control efforts. Here, we use machine learning and ecological and

evolutionary trait profiles of known hosts to identify unrecognized potential wildlife hosts of

Leishmania. We identify 136 mammals in the Americas that are likely to be exposed to and

infected by zoonotic *Leishmania* in the wild. The high number of unrecognized potential hosts

emphasizes a need to better invest in studying the ecological epidemiology of leishmaniasis. The

study provides information and tools to support targeted intervention and management of this

important poverty-associated disease.

Introduction

Leishmaniasis, a debilitating and sometimes fatal parasitic disease, is one of the 20 neglected

tropical diseases targeted for control and elimination by the United Nations by 2030 (1). However,

leishmaniasis remains a persistent threat throughout the tropics and subtropics. In the Americas,

leishmaniasis cases occur from the southern United States to Argentina, including the Caribbean,

with an estimated 58,500 new cases each year (2). Further, incidence is rapidly shifting in

geographic extent with land-use and climate change (3–7). For instance, hotspots of

leishmaniasis have significantly expanded throughout Costa Rica in the last two decades, in

concert with agricultural intensification, deforestation, and reforestation (3).

Treatment for leishmaniasis has many side effects and are extremely expensive (2,8), thus control

and elimination hinges on limiting transmission. *Leishmania* parasites are transmitted by sandflies

(Order: Diptera; Family: Psychodidae) and maintained in the environment via a multitude of

sylvatic and domestic mammalian hosts. As such, controlling transmission requires ecological

interventions targeting either component of the transmission cycles. For instance, clearing vector

habitat proximate to households, supporting populations of natural enemies that regulate

populations of vectors or small mammals (e.g., fish and mosquito larvae), and/or modifying

distribution of wildlife food sources are sustainable and effective interventions in preventing

transmission of flaviviruses, henipaviruses, *Plasmodium* spp. (malaria), and *Brucella abortus*

(brucellosis) (9,10). Similar interventions could be employed against *Leishmania,* however,

successful application requires species identification of vectors, hosts, and their corresponding

interactions with their biotic and abiotic environment.

Leishmaniasis was first diagnosed by Western medicine in the Americas in 1909, at the time dogs

and foxes were believed to be the zoonotic hosts (11). Continued research throughout the century

hypothesized that rodents, and in some cases opossums and sloths, were the primary zoonotic

hosts of these pathogens (11). However, especially with the advancement of molecular diagnostic

techniques, the number and taxonomic diversity of hosts has since expeditiously increased. In

the Americas, there are now 137 species of wildlife that are recognized as potential hosts (i.e.,

animals that are naturally exposed and become infected once exposed), of which around 60 may

act as competent hosts (i.e., animals can become infectious once infected, and may maintain the

pathogen in the environment as reservoir hosts) (12,13). Wild hosts are now known to include

eight orders of mammals including Chiropetra (bats), Carnivora, Cingulata (armadillos),

Didelphimorphia (opossums), Lagomorpha (rabbits), Pilosa (sloths and anteaters), Primates, and

Rodentia. However, even with progress in host discovery, the full suite of *Leishmania* hosts is not

yet described: *Leishmania* spillover occurs in absence of known hosts and new hosts are

discovered almost annually (14–17) .

Limited knowledge on the transmission cycle makes it difficult to understand the mechanisms

driving *Leishmania* distribution and spillover. As previously stated, unknown transmission cycles

impede effective ecological interventions to reduce spillover risk to people. Additionally, global

change can alter host dynamics to amplify spillover risk. For example, land-use change may

increase host densities and/or push hosts into closer proximity to anthropophilic vectors and

humans (9). Climate warming may also shrink or expand the range of hosts, further changing the

rate of contact among hosts, vectors, and humans (18). As such, the incomplete classification of

possible *Leishmania* hosts could lead to unexpected emergence and hotspots of human

leishmaniasis.

Study designs optimal for detecting *Leishmania* hosts, such as systematic sampling of large

populations of a diversity of species through space and time, are limited by logistics and cost of

sampling. *Leishmania* host discovery could benefit from strong *a priori* hypotheses about which

mammals to target for sampling effort; taxa specific sampling could help to efficiently use time

and monetary resources, ultimately expediting the full classification of the *Leishmania*

transmission cycle. Through the last decade, machine learning approaches have been used to

generate predictions of hosts of a multitude of different pathogens, including betacoronaviruses

(19), flaviviruses (20), Rabies virus (21), and *Borrelia burgdoferi* (causative agent of Lyme

disease) (22). These methods have facilitated more targeted laboratory and field work of novel

pathogens, and they have great potential to improve our understanding of diseases of poverty

with broad host and vector ranges.

Host status for any parasite or pathogen is driven by two processes: exposure and physiological

susceptibility (23,24). Exposure depends on the environmental conditions needed to support

pathogen transmission and, in the case of vector-borne disease, vector reproduction, contact with

competent hosts, and vector survival (25). Following exposure, physiological susceptibility then

depends on pathogen interaction with host cells to gain entry and, in some cases, avoid immune

attack and replicate (25). Host traits related to exposure and/or susceptibility likely interact in non-

linear and higher-order combinations to delineate zoonotic *Leishmania* hosts from non-hosts. In

brief, the effect of a trait on host status may vary in magnitude or direction depending on trait

value and traits may become more or less important under different conditions. Hypothetically, for

example, the average temperature of a species range may have a positive effect on host status

between 25-30 °C, but a negative effect outside of this range. Depending on biting activity of

vector species, animals that forage on the ground may be more likely to be hosts if they are also

nocturnal, whereas animals that are arboreal may be more likely to be hosts if they are

crepuscular but less likely to be hosts if they are nocturnal. Thus, when viewing the zoonotic

Leishmania system in one or two dimensions (e.g., only considering phylogeny or habitat use), it

is difficult to determine unifying traits useful for identifying hosts. To date, it has been challenging

to combine these processes across scales from physiological to evolutionary to ecological to

biogeographic to predict their overall impact on infection potential. Machine learning, particularly

tree-based methods, offers a set of new tools that can be used to incorporate and dissect this

complexity by increasing the dimensionality in which the *Leishmania* system can be studied,

allowing for multiple predictors that may have nonlinear and interactive effects on host status. The

flexibility of these models allows for pinpointing combinations of mammalian traits unique to the

confluence of *Leishmania* exposure and susceptibility, by learning from patterns of known hosts.

With this information, we can identify likely host species as targets for *Leishmania* surveillance

efforts and the key traits that distinguish host species from non-hosts.

Here we use tree-based machine learning (extreme gradient boosted regression, 'Xgboost') and

mammalian traits, including but not limited to biogeographical, phylogenetic, and life-history

features, to: (i) describe trait profiles of hosts for zoonotic *Leishmania (Viannia)* and *Leishmania*

(Leishmania) parasites; and (ii) use the trait profiles to predict unrecognized wild hosts of zoonotic

Leishmania within the subgenera *Leishmania (Viannia)* and *Leishmania (Leishmania)*.

We aim for this analysis to identify animal species that are likely to be exposed and infected in

the wild; due to extreme data sparsity, we do not employ our methods to identify competent hosts

(i.e., animals that become infectious once infected and may act as reservoirs given the density

and distribution of their populations). Our findings can be used to identify hosts that can then be

surveyed at larger spatio-temporal scales, investigated for competence, or used as sentinel

species. As such, while our modeling approach does not necessarily identify competent hosts

that can transmit zoonotic *Leishmania* on to sandflies and eventually humans, it does identify

animals likely to be exposed to and susceptible to zoonotic *Leishmania* infection, facilitating the

first step of determining epidemiological importance and fine-tuning surveillance efforts.

Methods

Data collection

Leishmaniasis is caused by protists within the *Leishmania* genus. There are 14 known species of

zoonotic *Leishmania* (i.e., they spill over from animals to humans) in the Americas. These *Leishmania*

species are divided between two subgenera: *Leishmania* (*Viannia*) and *Leishmania (Leishmania)*. In

the Americas, *Leishmania (Viannia)* consists of ten species, nine of which are zoonotic; *Leishmania*

(Leishmania) consists of seven species, five of which are zoonotic (26). *Leishmania (Viannia)* spp.

cause cutaneous and mucocutaneous leishmaniasis, which manifests as skin lesions that are

susceptible to painful secondary bacterial infections, and/or destruction of the mucus membranes of the

nose, mouth, and throat (2). In addition to cutaneous and mucocutaneous leishmaniasis, species within

the *Leishmania (Leishmania)* subgenera (*L. (L.) infantum* and, less frequently, *L. (L.) mexicana*) cause

fatal visceral leishmaniasis. Visceral leishmaniasis affects internal organs, typically causing

enlargement of the spleen and liver (2). The following analyses were conducted for each subgenus (*L.*

(Viannia) and *L. (Leishmania)* so to better described the ecological associations of each taxonomic

group of parasites and their corresponding disease manifestations.

Zoonotic Leishmania *host status:* We gathered *Leishmania* host status from all wild, endemic,

and invasive terrestrial mammals with ranges in Mexico, Central America, and South America

using recent reviews(12,13) in addition to a Web of Science query on February 12, 2021 and

February 22, 2021 (see supplementary text 1 for the specific search strings). Host status was

defined as a binary trait: an animal has been naturally infected by one of the 14 *Leishmania*

species known to cause leishmaniasis in humans (1), or there is no record of infection by a

Leishmania species known to cause leishmaniasis in humans (0). The latter category includes

animals that have been tested for *Leishmania* and have not been found to be infected as well as

those that have not been tested for *Leishmania* infection. Infection was determined via detection

of *Leishmania* species specific antigens or genetic material. Species identification was performed

either using animal tissue samples directly or using live parasites cultured from animal tissue. To

ensure that all relevant mammals were included in our study, a list of species names of endemic

Latin American mammals was retrieved from the International Union for Conservation of Nature

(IUCN) Red List Inventory (27) and a list of invasive mammals was retrieved from the Global

Invasive Species Database (28). We cross-referenced our host status data table with the Global

Infectious Disease and Epidemiology Network (29). We then limited our analysis to wild mammals

that had at least a 10% range overlap with reported human cases of leishmaniasis for inclusion

in our analysis. In total, 86.64% (1460/1685) of species were retained for the *L. Viannia* analysis

and 87.24% (1470/1685) of species were retained for the *L. Leishmania* analysis. Zoonotic

Leishmania ranges were constructed by outlining a concave polygon around cases of disease

occurrence in humans from (30) (Figure 1a, 1c). We conducted all data filtering and downstream

analyses separately for each subgenus.

Traits associated with Leishmania exposure and susceptibility: Our analysis leverages patterns

of traits of known hosts to predict identities of unrecognized hosts. We used host traits related to

Leishmania exposure (via sandfly bites) and susceptibility, including life-history, biogeographical,

and phylogenetic traits (supplementary table 1).

For life-history and some habitat use traits, we collected traits from panTHERIA (31),

mammalDIET (32), and EltonTraits (33). We additionally used IUCN range maps and satellite

imagery to extract data associated with species biogeography, including variables related to

climate (34,35), land-cover type (36), and global human modification index, an aggregate

measure of anthropogenic pressure on a landscape (37), within each species range (see

supplementary table 1 for full description of traits and satellite imagery used). Ranges of endemic

species were retrieved from the IUCN database and ranges of invasive species in Mexico, Central

America, and South America were constructed by building a concave polygon around species

occurrence points from (38). *Leishmania* and invasive mammal species ranges were built using

the R (v 4.0.2) packages *speciesgeocodeR* (39) and *concaveman* (40). We additionally included

IUCN-designated main habitat and habitat breadth (IUCN) as traits. To quantify phylogenetic

traits, we downloaded pairwise divergence time (phylogenetic distance) from TimeTree.org (41)

and reduced the dimensionality of this matrix by mapping species in ordination space using

principal coordinate analysis in base R. TimeTree is a knowledge-base that has collected and

synthesized species divergence times from > 3000 peer-reviewed studies. We included whether

an animal is invasive or endemic in the given *Leishmania* range and zoonotic host status using

GISD and GIDEON. Finally, we accounted for sampling effort by downloading the number of

citations found on PubMed per species using the R package *easyPubmed* (42). We choose to

use PubMed as this indicates the biomedical study effort, as opposed to a more general measure

of study effort that would be estimated using the number of citations on Web of Science.

Biomedical study effort better represents the number of times a species was studied in the context

of testing for pathogen assemblages.

Data analysis

We applied extreme gradient boosted regression (XGboost) in the R package *xgboost* (43) to use

mammal trait data to predict the probability that a species is a *Leishmania (Leishmania)* or

Leishmania (Viannia) host. Extreme gradient boosted regression is a machine learning algorithm

that creates an ensemble of weak decision trees to form a stronger prediction model by iteratively

learning from weak classifiers and combining them into a strong classifier (i.e., boosting) (44).

Gradient boosted regression is flexible in that it allows for non-linearity, both among features (i.e.,

interactions) and between features and predictions, collinearity between features, and non-

random patterns of missing data. Further, XGboost handles extremely unbalanced data well by

weighting positive labels to increase class separability and allowing for a regularization parameter

to prevent overfitting to a few positive labels – an advantage when analyzing our data set with

relatively few known hosts compared to the number of mammal species examined. The ability to

handle non-random patterns of missing data and extremely unbalanced data makes the XGBoost

algorithm best suited for our study as compared to other machine learning methods, such as

random forest and support machine vector, which severely overfit unbalanced data (random

forest) and/or cannot handle missing data without use of imputation (random forest, support

machine vector). Prior to analysis we removed traits with > 0.7 correlation to increase

interpretability of variable importance. For the *L. Viannia* analysis 32/66 were retained in the final

analysis; for the *L. Leishmania* analysis 33/66 were retained. To maximize the number of traits

retained in the model, out of a pair of correlated traits, the one with the highest number of

correlations to other traits and/or the lowest data coverage was dropped. We used a box-cox

transformation of extremely skewed variables to reduce the influence of extreme values on model

performance. Categorical traits were one-hot-encoded so that each category was represented as

a binary trait. Finally, traits with < 10% coverage (i.e., less than 10% of the species had data on

the trait) were removed and not used in the analysis (both for training the model and prediction).

Model performance and hyperparameter tuning: The accuracy of our model was evaluated using nested cross-validation. Nested cross-validation, as opposed to K-fold cross-validation, produces the least biased evaluation of model performance when using a small sample size (45). Three-fold cross validation was used to generate model predictions and evaluate overall model performance, while 5-fold cross validation was performed to train hyperparameters within each 3-fold cross validation step. In other words, the full dataset was split into three outer folds. Hyperparameters were tuned using Bayesian optimization, optimizing over a parameter space aimed to reduce overfitting (low training rate, low ratio of samples used in trees, low ratio of features used in trees, low maximum tree depth, and high regularization). Using the hyperparameters that yielded the best performance (minimum logloss), a final model was then trained using the full data within these two folds and model predictions were calculated on the third hold-out fold. This process was repeated until out-of-sample predictions were made for each fold. As model output may be dependent on the distribution of the data across each fold, nested cross validation was repeated 25 times using 25 unique splits of the data. Our dataset was extremely unbalanced (i.e., many more 0s than 1s), which makes our model vulnerable to overfitting. To determine if our model was simply fitting spurious correlations in the data, we conducted a target shuffling analysis (46,47). We repeated the nested cross-validation analysis but randomized the response variables (*Leishmania* host status) for each iteration. We then calculated the mean AUC from the target shuffling experiment. AUC, the area under the receiver operator characteristic curve, is a classification metric that measures the probability that model output for a randomly chosen positive label (known host) will be higher than a randomly chosen negative label (animal with no record of zoonotic *Leishmania* infection). AUC ranges from 0 to 1, with an AUC of 1 indicating that the model perfectly classifies all samples, while an AUC < 0.5 indicates the model performs no better than a coin flip. We found that our model performed minutely better than a coin flip in the target shuffling experiment (average *L. (Leishmania)* target-

shuffled AUC = 0.53; average *L. (Viannia)* target-shuffled AUC = 0.54), thus we adjusted our final

AUC values by $AUC_{final\ model}$ - ($AUC_{target\ shuffled\ model}$ - 0.5).

In our final model, citation count was among the top 10 most important variables. To ensure that

we were not simply predicting well-studied animals, we repeated the analysis above while

replacing the response variable with study effort (the number of citations on PubMed). To reduce

computational time, we ran ten iterations instead of 25. We concluded that our model was not

simply predicting well-studied species by evaluating how well the traits in our model predict study

effort, measured as pseudo-R^2. Additionally, we checked that trait responses (i.e., partial

dependence plots) that predicted publication number were not overly similar to the functional form

of the trait responses that predicted host status.

Leishmania *host trait profiles:* To assess *Leishmania* host trait profiles, we first identified important

features using SHAP (Shapley Additive Explanations) calculated via the R package

SHAPforxgboost (48). Shapley scores represent the average marginal contribution of a feature to

a prediction across all combinations of features; the contribution of the feature is interpreted as

the change in prediction associated with that feature in relation to the average model prediction

(44). SHAP calculates local feature contribution by evaluating the contribution of each feature to

the prediction for each observation in a dataset. Here, local contribution refers to the contribution

of each feature to the prediction for each individual species (local corresponds to the level of a

single sample, which, for our analysis, is a species). Global feature contribution can be obtained

by aggregating the Shapley scores for each feature across all observations; global feature

contribution is the average contribution of each feature across all species (global corresponds to

the level of the entire dataset). To generate a measure of uncertainty in feature contribution, we

trained the model with 70% of the data, and calculated Shapley scores for the remaining 30% of

observations. We repeated the above procedure 100 times, using different subsets of data with

each iteration. We then described trait profiles using SHAP partial dependence plots, which map

the relationship between each value of a feature in the validation set and the contribution of the

feature to model predictions.

Leishmania host predictions: Following (21), we identified animals as unrecognized zoonotic

Leishmania spp. hosts using the Shapley value classification criterion. The sum of the Shapley

scores across features for each species sums to the prediction for each species, relativized by

the average model prediction across species. A summed Shapley value of 0 represents the

average prediction in the dataset (49,50). To account for model uncertainty, we used a

bootstrapping procedure to calculate Shapley scores for models fit with 70% of the data 100x. We

classified an animal as a newly predicted host if > 95% of Shapley scores were > 0; in other

words, we classified an animal as a newly predicted host if predicted probability of being a host

was greater than the average probability for that model iteration for > 95% or model iterations

(21).

The full record of species infection status and references, trait data-base and R code can be found

at https://github.com/ckglidden/leish_hosts.

Results

Figure 1. The ranges of zoonotic *Leishmania* and their known and predicted hosts extend throughout the Americas. The number of overlapping ranges of known *Leishmania* hosts are depicted in (a & c), with the range of human cases of *L. (Viannia)* and *L. (Leishmania)* outlined in grey. The number of overlapping ranges of newly predicted hosts are outlined in (b & d). *Hylaeamys megacephalus* (e) and *Calomys callosus* (f) are unrecognized hosts with the highest and second highest predictions for *L. (Viannia)* according to Shapley values. Their ranges are outlined in dark green and light green, respectively, in (b). *Dasypus novemcinctus* (g) and

Leopardus wiedii (h) are currently hosts with the highest and second highest Shapley values for

L. (Leishmania). Their ranges are outlined in dark blue and light blue, respectively, in (d). An

exhaustive list of newly predicted hosts is listed in supplementary table 2.

<u>*Leishmania (Viannia):*</u> Through our literature search, we found 96 known *L. (Viannia)* hosts out of

1460 mammals that occur within the geographic range of *L. (Viannia)* parasites (Figure 1a; Figure

2a). After accounting for target shuffling, our model performed moderately well with an average

out of sample AUC of 0.81 (95% CI: 0.80 - 0.82) and in sample AUC of 0.94.

L. (Viannia) *trait profiles:* After removing highly correlated traits, 30 features were used to predict

L. (Viannia) hosts. A mix of biogeography, life-history, phylogeny, and study effort covariates were

among the top important model features when estimating global feature contribution (Figure 3).

Phylogenetic distances, reduced into six axes of a PCoA ordination, were, on average, the most

important trait for *Leishmania (Viannia)* hosts. Specifically, animals were more likely to be hosts

if they were rodents and opossums, and less likely to be hosts if they were non-human primates,

carnivores, and even toed ungulates (supplementary figure 1-2).   Similar to other zoonotic

pathogen systems, *L. (Viannia)* hosts live at high population densities and demonstrate signatures

of fast-paced life-histories, indicated by short gestation periods (Figure 4a). Further, *L. (Viannia)*

host ranges are large, have high proportions of crop and urban land cover, and, on average, peak

at ~28°C during the warmest quarter of the year (Figure 4a). Geographic coordinates of the

species' native ranges were also an important predictors: animals were more likely to be hosts if

the minimum longitude of their native range was around western Central America. Notably,

although study effort (number of PubMed citations, normalized with a Box-Cox transformation)

was among the most important traits, our supplementary analyses indicated that the host models

were not simply predicting well-studied hosts. Specifically, the supplementary model predicting

study effort (number of PubMed citations) performed poorly (average out-of-sample R^2 = 0.19).

Further, traits that predict citation count, and the response curves of those traits, greatly differed

from traits that predict *L. (Viannia)* host status (supplementary figure 3-4). As such, while *L.*

(*Viannia*) hosts are likely understudied, we are not simply reporting the trait profiles of well-studied

mammals.

L. (Viannia) *hosts:* Eighty-three animals were identified as likely hosts (Figure 2, supplementary

table 2). Out of these newly predicted hosts, the majority were rodents, bats, and opossums.

Further, 14 of these hosts are known hosts of *L. (Leishmania)* (supplementary table 2)*.* The newly

predicted hosts with the highest summed Shapley scores were the large headed rice rat

(*Hylaeamys megacephalus*) and the large vesper mouse, *Calomys callosus* (Figure 1e,f). In

contrast to known hosts, the range of newly predicted hosts extends to the southern tip of South

America and there is are is high density of species ranges that overlap in the southern United

States (Figure 1a, b).

Figure 2. The number of known and predicted hosts per order for *L. (Viannia)* (a) and *L.*
(Leishmania) (b). Grey bars represent the number of known hosts within each order, colored
bars represent the number of newly predicted hosts per order. Animals were classified as newly
predicted if > 95% of mean SHAP values were greater than 0 (a-b). Bars in (c) represent the
proportion of newly classified hosts within each order; colors match (a-b).

Leishmania (Leishmania): Through our literature search, we found 110 known hosts out of 1470
mammals that have overlapping ranges with the zoonotic *L. (Leishmania)* range (Figure 1c). Of
these, 40 animals were hosts for both *Leishmania* subgenera. After accounting for target shuffling,

our *L. (Leishmania)* hosts model performed moderately well with an average out of sample AUC

of 0.84 (95% CI: 0.83 - 0.85) and in sample AUC of 0.94.

L. (Leishmania) *trait profiles:* After removing highly correlated traits, 32 features were used to

predict *L. (Leishmania)* hosts. On average, study effort was the most important trait (i.e., had the

highest global feature contribution), followed by the land cover/land-use composition of the

species range, average temperature in the warmest quarter of the species range, range area,

phylogenetic distance, litter size, gestation length, and longitudinal extent of the species range

(Figure 3). Similarly, to *L. (Viannia)* hosts, *L. (Leishmania)* hosts show signatures of fast-paced

life histories, indicated by short gestation lengths and large litter sizes, and have ranges with high

degrees of crop and urban cover (Figure 3b). Further, the range of *L. (Leishmania)* hosts are

similar to *L. (Viannia)* hosts in that the mean temperature in the warmest quarter peaks around

28°C (Figure 4b). On average, geographic coordinates of the species' native range were also an

important predictor: Animals were more likely to be hosts if the minimum longitude of the range

was around western Central America, while the maximum longitude of the range was western

Europe, which indicates a role of invasive species in the *L. (Leishmania)* transmission cycle.

While study effort was among the most important traits, host traits explained little variation in

number of PubMed citations (average out-of-sample R^2 = 0.17) and traits that predict citation

count, and the response curves of those traits, greatly differed from traits that predict *L.*

(Leishmania) host status suggesting that traits important for discriminating host status were not

just those that predicted study effort (supplementary figures 5-6).

L. (Leishmania) *hosts*: Using the Shapley classification criterion, 98 animal species that are

currently unrecognized hosts were labeled as newly predicted hosts (Figure 2, supplementary

table 2). Out of these newly predicted hosts, the majority were rodents, bats, and carnivores. Nine

newly predicted hosts are known hosts of *L. (Viannia)*. Forty-five newly predicted hosts were

newly predicted hosts for both the *L. (Leishmania)* and *L. (Viannia)* subgeneras (supplementary

table 2). The newly predicted hosts for the *L. (Leishmania)* with the top summed Shapley scores

included the nine-banded armadillo (*Dasypus novemcinctus*) and margay (*Leopardus wiedii*)

(Figure 1g,h). The nine-banded armadillo is a known host of *L. (Viannia)* parasites, with a range

that extends from northern Argentina to the central United States (Figure 1d).

Figure 3. Biogeography, life-history, and phylogenetic traits all significantly contributed to model predictions for host status. Biogeographical traits (minimum longitude of the range, maximum longitude of the range, % cover of land-use/land cover in the species range, average temperature in the warmest quarter in the species range, range area) are colored in blue, life-history traits (population density, gestation length, litter size) are colored in red, phylogenetic distance (location along PCoA ordination axes) is colored in orange, and study effort is colored in grey. Points are the absolute value of the mean Shapley importance (mean |SHAP value|) for the trait across all mammals (i.e., global feature contribution), bars represent the absolute values of the 0.05-0.95 percentiles. Only features with 0.05 percentiles > 0 are shown.

Figure 4. Hosts have fast-paced life-histories and live in proximity to humans. Shapley

partial dependence plots showing the effect of each feature on *L. (Viannia)* (a) and *L. (Leishmania)*

(b) host status after accounting for the average effect of the other features in the model. Colored

lines represent the average effect across model iterations, while grey lines show each individual

model iteration (model fit with 80% of data) (blue = biogeographical traits; red = life-history).

Features with global mean feature contribution scores > 0 for > 95% of model iterations are

shown. Rug plots on the x-axis indicate the distribution of the data.

Discussion

Our literature search identified 137 wild mammals in the Americas as known zoonotic *Leishmania*

hosts, and our model predicted there to be an additional 136 likely hosts (animals that are likely

to be exposed and infected in the wild, but not necessarily infectious upon infection). The majority

of the predicted hosts for both subgenera fell within the Rodentia order (*L. (Leishmania)* N = 27;

L. (Viannia) N: = 34), with a similar number of predicted *Leishmania (Leishmania)* hosts within the

Chiroptera order (N = 32). While their contribution to model predictions varied between

subgenera, key traits included phylogenetic distance, human associated land-use, climate,

population density, and study effort.

For both *Leishmania* subgenera, biogeographical features encompassing descriptors of species

ranges were among the top five contributors to host predictive accuracy. Notably, zoonotic *L.*

(*Leishmania*) hosts are more likely to be animals with larger ranges than non-hosts. Geographic

range area was also reported to be a shared predictive feature among zoonotic hosts and

reservoirs in other systems (21,51). Further, ranges of *L. (Leishmania)* and *L. (Viannia)* hosts are

made up of higher proportions of crop cover and urban cover than ranges of mammals that are

not *Leishmania* hosts. These results are in accordance with previous findings that, in general,

zoonotic host richness and relative abundance increases with anthropogenic pressure (52).

Further, our results align with recent literature showing that risk of leishmaniasis is associated

with agricultural intensification and urbanization (4,53). *L. (Viannia)* hosts are marginally more

likely to have ranges with moderate forest cover, than non-hosts whereas there is little effect of

forest cover on *L. (Leishmania)* host status. Vectors of cutaneous leishmaniasis, which is caused

by all species of zoonotic *Leishmania* except *L. infantum*, are more likely to live in forests with

high integrity and low human modification (54), while hosts are more common in disturbed

landscapes (55). Our analysis adds support to the latter finding, highlighting that transmission of

parasites associated with cutaneous leishmaniasis likely occurs at forest-human settlement

interfaces. The incomplete overlap between vector ranges and host ranges may explain the highly

localized dynamics of cutaneous leishmaniasis spillover (i.e., transmission from wildlife to humans

via the sandfly vector) (6,56). Spillover occurs when reservoir hosts, vectors, and humans overlap

in space and time so that forward transmission occurs from animal to sandfly to human. If vector

and host distributions only overlap at habitat interfaces, area of reservoir host, sandfly, and human

co-occurrence may be low and variable through time. Finally, average temperature of the warmest

quarter was an important predictor of host status, with the feature contribution to model

predictions peaking around 28°C. In the case of zoonotic leishmaniasis, temperature is likely a

predictor of *Leishmania* exposure as this trait profile reflects temperatures associated with vector

survival and parasite development (57,58).

The profiles of important life history traits for *Leishmania* hosts resembled those observed in other

comparative analyses of zoonotic diseases. Specifically, for both *Leishmania* subgenera, the

probability of host status peaked at low to median lengths of gestation, a pattern similar to that of

rodent reservoirs of zoonotic pathogens and competent hosts of zoonotic tick-borne pathogens

(51,59). Further, the probability of zoonotic *L. (Leishmania)* host status increased with litter size,

a trend that parallels that of both rodents reservoirs of zoonotic pathogens and bat and carnivore

reservoirs of rabies virus (21,51). Altogether, our findings suggest that, like other zoonotic

reservoirs and hosts, zoonotic *Leishmania* hosts may have fast-paced life-histories. Animals with

fast-paced life-histories are believed to maximize fitness by expeditiously investing in short-term

reproductive effort -- such as through younger ages at sexual maturity, short gestation lengths,

greater litter sizes -- at the cost of long-term survival (60). As a result, animals with fast-paced life

histories are thought to divert energetic resources away from effective adaptive immunity (61), or

utilize responses conducive to tolerance but not resistance (62), allowing them to allocate

maximal resources to short term reproductive effort. As such, animals with fast-paced life-histories

may be more susceptible to infections.

However, *L. (Viannia)* hosts only had one trait (gestation length) clearly associated with a fast-

paced life history, suggesting weaker evidence for the "fast-paced" host hypothesis for this

subgenus. Phylogenetic distance was the most important trait for *L. (Viannia)* hosts. We

hypothesize that phylogenetic relationships may better account for host physiological defenses

against *Leishmania* than life-history traits. If this is the case, inclusion of specific genetic markers

associated with *Leishmania*-host cell interactions may improve model predictions.

Notably, study effort significantly contributed to host predictions for both subgenera. While our

follow up analyses do not indicate that model results are biased by uneven study effort, this finding

may suggest that zoonotic *Leishmania* hosts are generally under sampled. Though leishmaniasis

is a neglected tropical disease primarily impacting resource-poor communities, state-of-the-art

zoonotic *Leishmania* diagnostic and surveillance techniques are available (16,55,63–67). As

such, under sampling *Leishmania* hosts primarily stems from both the lack of funding to employ

these tools during large scale surveillance efforts and the large number of animal species that are

potentially hosts. The main leishmaniasis research funding mechanism within the USA, the

National Institute of Health (NIH), primarily focuses on the molecular biology of leishmaniasis work

and does not prioritize research elucidating the ecology of the sandfly-wildlife transmission cycle.

In fact, a simple search for "leishmania" on the NIH and National Science Foundation grants

database yields a combined total of three awards aiming to elucidate transmission cycles of

Leishmania in the Americas (searched October 10, 2022). Researchers across Latin America are

faced with similar, if not more severe, funding constraints. For example, in Brazil, a major hotspot

of leishmaniasis, the Bolsonaro administration cut 90% of federal science funding (68). While our

model results facilitate more targeted field sampling for particular species, optimized surveillance

should consist of an iterative feedback between local knowledge and model predictions to guide

field work, and updating the model with new results from field sampling to guide ongoing

surveillance and field studies (16,19). Importantly, to reduce study effort bias going forward, care

should be taken to estimate and survey under studied areas and species in addition to newly

predicted species, particularly in areas with high rates of human spillover with poorly known hosts.

A lack of investment in understanding the basic ecology of *Leishmania* slows research progress

by obscuring the major drivers of disease dynamics and spillover transmission to humans.

Leishmaniasis surveillance and management would benefit from larger funding mechanisms

prioritizing understanding the ecology of this group of pathogens.

In some cases, our host predictions are supported by multiple independent but complementary

lines of evidence (summarized below and in supplementary table 3). Non-specific *Leishmania*

antigens, antibodies, or DNA has been detected in the crab-eating racoon (*Procyon cancrivorus*),

neotropical river otter (*Lontra longicaudis*), six-banded armadillo (*Euphractus sexcinctus*), coyote

(Canis latrans), and the grey short-tailed opposum (*Monodelphis domestica*) (12,66,69,70). As

tests were not parasite species specific, it is unconfirmed if they were infected with zoonotic

Leishmania species, but our model suggests they may be. Further, *Leishmania*-like flagellates

were discovered in Derby's wooly opossum (*Caluromys derbianus*) (69); however, to our

knowledge this species has not been tested for zoonotic *Leishmania* species.

A number of animals predicted to be hosts for *L. (Leishmania)* were also found to co-occur with

vectors of *L. (L.) mexicana* in Mexico: the hairy big-eyed bat (*Chiroderma villosum*), Brazilian

brown bat (*Eptesicus brasiliensis*), great sac-winged bat (*Saccopteryx bilineata*), tent-making bat

(*Uroderma bilobatum*), Mexican woodrat (*Neotoma mexicana*), black footed pygmy rice rat

(Oligoryzomys nigripes), Coue's rice rat (*Oryzomys couesi*), fulvous harvest mouse

(*Reithrodontomys fulvescens*), nine-banded armadillo (*Dasypus novemcinctus*), lowland paca

(*Cuniculus paca*), Virginia oppossum (*Didlephis virginiana*), kinkajou (*Potos flavus*), racoon

(*Procyon lotor*), white-nose coati (*Nasua narica*), and the margay (*Leopardus wiedii*) (16,71).

Notably, the nine-banded armadillo and margay had the two highest summed Shapley scores as

potential *L. (Leishmania)* hosts. The nine-banded armadillo is a known host of *L. (Viannia)* as well

as other zoonotic pathogens including the causative agent of leprosy (72).

Host competence (i.e., the ability to become infected and infectious) has been demonstrated in

the lab for four of the predicted species: the Virginia opossum (*Didlephis virginiana)*, the large

vesper mouse (*Calomys callosus*), spix's yellow toothed cavy (*Galea spixii*), and the Mexican

free-tailed bat (*Tadarida brasiliensis*) (69,73–75). Although host competence for these animals

has been tested in the laboratory, infection/exposure has not been tested in the wild or animals

have not tested positive in the wild (possibly due to minimal sampling of the species). Our model

suggests that these animals are also likely to be naturally infected in the wild and thus warrant

further investigation. The large vesper mouse is one of the top predicted species for *L. (Viannia).*

Surprisingly, we found no field studies testing wild large vesper mice for zoonotic *Leishmania*

infection. Finally, our model suggests that a handful of species naturally infected with *L. infantum*

in Europe are also likely to be exposed to and susceptible to zoonotic *Leishmania* in the Americas:

the red fox (*Vulpes vulpes*), the European hare (*Lepus europaeus*), and the European rabbit

(*Oryctolagus cuniculus*) (76–78).

Importantly, while there are 14 species of zoonotic *Leishmania* parasites in the Americas, our

analysis divides these species into two subgenera but does not identify *Leishmania* species

specific hosts. Future work could leverage the information presented by the current analysis to

identify *Leishmania* species specific hosts. Additionally, our analysis identifies animals that are

likely to be infected in the wild but does not narrow the host pool down to competent hosts

(animals that are likely to be both infected *and* infectious in the wild). Next steps of model

development could identify trait profiles of competent hosts that delineate them from non-

competent hosts. Notably, (54) used machine learning to predict unrecognized vectors of

leishmaniasis. Further, metabarcoding of arthropods has started to identify sandfly feeding

networks (64,79) that may be useful in linking model predictions of vectors and hosts to local

transmission cycles.

Overall, models for both subgenera identified a number of new hosts, including but not limited to

the nine-banded armadillo, the margay, the large vesper mouse, and the large headed rice rat,

that can be targeted for future surveillance and research to determine their ecological significance

in leishmaniaisis transmission and spillover to humans. Next steps should include ground truthing

the role of newly identified potential hosts in the zoonotic *Leishmania* transmission cycle, through

field sampling and laboratory experiments, through engagement with local communities, and

iteratively updating the current analysis as new information is collected, taking careful

consideration to not perpetuate the effects of study bias. Although a cautious first step in host

discovery work, our study can immediately inform surveillance efforts and study design. For

instance, host predictions may provide a starting point for implication of hosts during local

outbreaks of human leishmaniasis by providing a targeted list of likely suspects to sample. While

the study of leishmaniasis has historically focused on rodents and small marsupials, our study

suggests that capture methods should more commonly be supplemented by bat mist-netting

and/or techniques to capture small to medium sized carnivores. Finally, many newly predicted

hosts have ranges that extend throughout the southeastern USA. Ecological niche models

suggest that climate warming may increase the range of *Leishmania* vectors as far north as

Canada, and leishmaniasis endemicity within the US was recently confirmed (80–82). Similarly,

the range of predicted hosts for *L. (Viannia)* extends farther south through Argentina than currently

known hosts. Our analysis provides tools for monitoring leishmaniasis risk under global change

as these hosts could be used as sentinel species for tracking leishmaniasis range expansion.

Testing these hosts for leishmaniasis infection beyond the current putative boundaries of zoonotic

Leishmania ranges would help to evaluate if range expansion has already occurred as well as

provide an updated baseline for future monitoring.

Identifying local transmission cycles is key for reducing leishmaniasis spillover and disease

burden in people, especially in the face of land-use and climate change, which likely influence the

geographic distribution and extent of vector—host--human contact and were strongly associated

with novel host predictions in our models. We estimate that there are likely many unknown hosts

of leishmaniasis throughout the Americas, highlighting the complexity of the transmission cycle

and a need to increase study effort on the ecology and epidemiology of this system.

Acknowledgements

This project was supported by the National Science Foundation (DEB-2011147 CKG, EAM;

EEID-1717282 BAH), the National Institutes of Health (R35GM133439, R01AI102981,

R0AI168097; EAM), the Stanford King Center for Global Development (EAM), the Stanford

Woods Institute for the Environment (EAM), the Stanford Center for Innovation in Global Health

(EAM), and the Office of the Vice Provost for Undergraduate Education at Stanford (AM). We

thank the Mordecai Lab for providing feedback on manuscript drafts.

610
References

1.   WHO. The road map targets for 2030 [Internet]. The road map targets for 2030. 2022 [cited 2022
Jun 12]. Available from: https://www.who.int/teams/control-of-neglected-tropical-diseases/ending-
ntds-together-towards-2030/targets

2.   PAHO. Leishmaniasis - PAHO/WHO | Pan American Health Organization [Internet]. 2022 [cited
2022 Jun 12]. Available from: https://www.paho.org/en/topics/leishmaniasis

3.   Bayles BR, Rusk AE, Pineda MA, Chen B, Dagy K, Hummel T, et al. Spatiotemporal trends of
cutaneous leishmaniasis in Costa Rica. Trans R Soc Trop Med Hyg. 2021 May 8;115(5):569–71.

4.   Bayles BR, Rusk A, Christofferson R, Agar G, Pineda MA, Chen B, et al. Spatiotemporal dynamics
of vector-borne disease risk across human land-use gradients: examining the role of agriculture,
indigenous territories, and protected areas in Costa Rica. Lancet Glob Health. 2020 Apr 1;8:S32.

5.   Dutari LC, Loaiza JR. American Cutaneous Leishmaniasis in Panama: a historical review of
entomological studies on anthropophilic *Lutzomyia* sand fly species. Parasit Vectors. 2014 May
11;7(1):218.

6.   Gonçalves NV, Miranda C do SC, Costa RJF da, Guedes JA, Matsumura ES de S, Costa SBN da,
et al. Cutaneous leishmaniasis: Spatial distribution and environmental risk factors in the state of
Pará, Brazilian Eastern Amazon. J Infect Dev Ctries. 2019 Oct 31;13(10):939–44.

7.   Portella TP, Kraenkel RA. Spatial-temporal pattern of cutaneous leishmaniasis in Brazil. Infect Dis
Poverty. 2021 Jun;10(03):47–57.

8.   Chakravarty J, Sundar S. Current and emerging medications for the treatment of leishmaniasis.
Expert Opin Pharmacother. 2019 Jul 3;20(10):1251–65.

9.   Glidden CK, Nova N, Kain MP, Lagerstrom KM, Skinner EB, Mandle L, et al. Human-mediated
impacts on biodiversity and the consequences for zoonotic disease spillover. Curr Biol. 2021 Oct
11;31(19):R1342–61.

10.   Sokolow SH, Nova N, Pepin KM, Peel AJ, Pulliam JRC, Manlove K, et al. Ecological interventions
to prevent and manage zoonotic pathogen spillover. Philos Trans R Soc Lond B Biol Sci. 2019 Sep
30;374(1782):20180342.

11.   Lainson R, Shaw JJ. Epidemiology and ecology of leishmaniasis in Latin-America. Nature. 1978
Jun;273(5664):595–600.

12.   Azami-Conesa I, Gómez-Muñoz MT, Martínez-Díaz RA. A Systematic Review (1990–2021) of Wild
Animals Infected with Zoonotic *Leishmania*. Microorganisms. 2021 May 20;9(5):1101.

13.   Roque ALR, Jansen AM. Wild and synanthropic reservoirs of *Leishmania* species in the Americas.
Int J Parasitol Parasites Wildl. 2014 Aug 29;3(3):251–62.

14.   López M, Erazo D, Hoyos J, León C, Fuya P, Lugo L, et al. Measuring spatial co-occurrences of
species potentially involved in *Leishmania* transmission cycles through a predictive and fieldwork
approach. Sci Rep. 2021 Mar 24;11(1):6789.

15. Medkour H, Davoust B, Levasseur A, Mediannikov O. Molecular Evidence of *Leishmania infantum*
and *Leishmania guyanensis* in Red Howler Monkey (Alouatta seniculus) from French Guiana.
Vector-Borne Zoonotic Dis. 2019 Dec;19(12):896–900.

16. Stephens CR, González-Salazar C, Sánchez-Cordero V, Becker I, Rebollar-Tellez E, Rodríguez-
Moreno Á, et al. Can You Judge a Disease Host by the Company It Keeps? Predicting Disease
Hosts and Their Relative Importance: A Case Study for Leishmaniasis. PLoS Negl Trop Dis. 2016
Oct 7;10(10):e0005004.

17. Vieira TM, Silva S de O, Lima L, Sabino-Santos G, Duarte ER, Lima SM, et al. *Leishmania* diversity
in bats from an endemic area for visceral and cutaneous leishmaniasis in Southeastern Brazil. Acta
Trop. 2022 Apr 1;228:106327.

18. Carlson CJ, Albery GF, Merow C, Trisos CH, Zipfel CM, Eskew EA, et al. Climate change increases
cross-species viral transmission risk. Nature. 2022 Jul;607(7919):555–62.

19. Becker DJ, Albery GF, Sjodin AR, Poisot T, Bergner LM, Chen B, et al. Optimising predictive
models to prioritise viral discovery in zoonotic reservoirs. Lancet Microbe [Internet]. 2022 Jan 10
[cited 2022 Jun 12]; Available from:
https://www.sciencedirect.com/science/article/pii/S2666524721002457

20. Han BA, Majumdar S, Calmon FP, Glicksberg BS, Horesh R, Kumar A, et al. Confronting data
sparsity to identify potential sources of Zika virus spillover infection among primates. Epidemics.
2019 Jun 1;27:59–65.

21. Worsley-Tonks KEL, Escobar LE, Biek R, Castaneda-Guzman M, Craft ME, Streicker DG, et al.
Using host traits to predict reservoir host species of rabies virus. PLoS Negl Trop Dis. 2020 Dec
8;14(12):e0008940.

22. Becker DJ, Han BA. The macroecology and evolution of avian competence for *Borrelia burgdorferi*.
Glob Ecol Biogeogr. 2021;30(3):710–24.

23. Combes C. The Art of Being a Parasite [Internet]. Chicago, IL: University of Chicago Press; 2005
[cited 2023 Jan 5]. 280 p. Available from:
https://press.uchicago.edu/ucp/books/book/chicago/A/bo3534652.html

24. Sweeny AR, Albery GF. Exposure and susceptibility: The Twin Pillars of infection. Funct Ecol. 2022
May 18;1365-2435.14065.

25. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to zoonotic
spillover. Nat Rev Microbiol. 2017 Aug;15(8):502–10.

26. Akhoundi M, Kuhls K, Cannet A, Votýpka J, Marty P, Delaunay P, et al. A Historical Overview of the
Classification, Evolution, and Dispersion of *Leishmania* Parasites and Sandflies. PLoS Negl Trop
Dis. 2016 Mar 3;10(3):e0004349.

27. The IUCN Red List of Threatened Species [Internet]. IUCN Red List of Threatened Species. [cited
2022 Aug 31]. Available from: https://www.iucnredlist.org/en

28. GISD [Internet]. [cited 2023 Jan 5]. Available from: http://www.iucngisd.org/gisd/

29. Global Infectious Diseases and Epidemiology Network [Internet]. GIDEON. [cited 2023 Jan 5].
Available from: https://www.gideononline.com/

30. Herrera G, Barragán N, Luna N, Martínez D, De Martino F, Medina J, et al. An interactive database
of *Leishmania* species distribution in the Americas. Sci Data. 2020 Apr 3;7(1):110.

31. Jones KE, Bielby J, Cardillo M, Fritz SA, O'Dell J, Orme CDL, et al. PanTHERIA: a species-level
database of life history, ecology, and geography of extant and recently extinct mammals. Ecology.
2009;90(9):2648–2648.

32. Kissling WD, Dalby L, Fløjgaard C, Lenoir J, Sandel B, Sandom C, et al. Establishing
macroecological trait datasets: digitalization, extrapolation, and validation of diet preferences in
terrestrial mammals worldwide. Ecol Evol. 2014;4(14):2913–30.

33. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. EltonTraits 1.0:
Species-level foraging attributes of the world's birds and mammals. Ecology. 2014;95(7):2027–
2027.

34. Brun P, Zimmermann NE, Hari C, Pellissier L, Karger DN. Global climate-related predictors at
kilometre resolution for the past and future. Earth Syst Sci Data Discuss. 2022 Jun 27;1–44.

35. Karger DN, Schmatz DR, Dettling G, Zimmermann NE. High-resolution monthly precipitation and
temperature time series from 2006 to 2100. Sci Data. 2020 Jul 23;7(1):248.

36. Buchhorn M, Lesiv M, Tsendbazar NE, Herold M, Bertels L, Smets B. Copernicus Global Land
Cover Layers—Collection 2. Remote Sens. 2020 Jan;12(6):1044.

37. Kennedy CM, Oakleaf JR, Theobald DM, Baruch-Mordo S, Kiesecker J. Managing the middle: A
shift in conservation priorities based on the global human modification gradient. Glob Change Biol.
2019;25(3):811–26.

38. Rosa CA da, Ribeiro BR, Bejarano V, Puertas FH, Bocchiglieri A, Barbosa AL dos S, et al.
NEOTROPICAL ALIEN MAMMALS: a data set of occurrence and abundance of alien mammals in
the Neotropics. Ecology. 2020;101(11):e03115.

39. Töpel M, Zizka A, Calió MF, Scharn R, Silvestro D, Antonelli A. SpeciesGeoCoder: Fast
Categorization of Species Occurrences for Analyses of Biodiversity, Biogeography, Ecology, and
Evolution. Syst Biol. 2017 Mar 1;66(2):145–51.

40. Gombin J, Vaidyanathan R, Agafonkin V. concaveman. R. 2020;

41. Kumar S, Stecher G, Suleski M, Hedges SB. TimeTree: A Resource for Timelines, Timetrees, and
Divergence Times. Mol Biol Evol. 2017 Jul 1;34(7):1812–9.

42. Fantini D. easyPubMed – Data Pulse [Internet]. 2019 [cited 2022 Aug 31]. Available from:
https://www.data-pulse.com/dev_site/easypubmed/

43. Chen T, Guestrin C. XGBoost: A Scalable Tree Boosting System. In: Proceedings of the 22nd ACM
SIGKDD International Conference on Knowledge Discovery and Data Mining [Internet]. San
Francisco California USA: ACM; 2016 [cited 2022 Aug 31]. p. 785–94. Available from:
https://dl.acm.org/doi/10.1145/2939672.2939785

44. Molnar C. Interpretable Machine Learning [Internet]. 2nd ed. 2022 [cited 2022 Aug 31]. Available
from: https://christophm.github.io/interpretable-ml-book/

45. Vabalas A, Gowen E, Poliakoff E, Casson AJ. Machine learning algorithm validation with a limited
sample size. PLOS ONE. 2019 Nov 7;14(11):e0224365.

46. Fischhoff IR, Castellanos AA, Rodrigues JPGLM, Varsani A, Han BA. Predicting the zoonotic
capacity of mammals to transmit SARS-CoV-2. Proc Biol Sci. 2021 Nov 24;288(1963):20211651.

47. Martin JT, Fischhoff IR, Castellanos AA, Han BA. Ecological Predictors of Zoonotic Vector Status
Among *Dermacentor* Ticks (Acari: Ixodidae): A Trait-Based Approach. Gaff H, editor. J Med
Entomol. 2022 Sep 5;tjac125.

48. Liu Y, Just A. SHAPforxgboost: SHAP Plots for "XGBoost". R Package Version 010. 2020;

49. Lundberg S, Lee SI. A Unified Approach to Interpreting Model Predictions [Internet]. arXiv; 2017
[cited 2022 Sep 21]. Available from: http://arxiv.org/abs/1705.07874

50. Lundberg SM, Erion G, Chen H, DeGrave A, Prutkin JM, Nair B, et al. Explainable AI for Trees:
From Local Explanations to Global Understanding [Internet]. arXiv; 2019 [cited 2022 Sep 21].
Available from: http://arxiv.org/abs/1905.04610

51. Han BA, Schmidt JP, Bowden SE, Drake JM. Rodent reservoirs of future zoonotic diseases. Proc
Natl Acad Sci. 2015 Jun 2;112(22):7039–44.

52. Gibb R, Redding DW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, et al. Zoonotic host
diversity increases in human-dominated ecosystems. Nature. 2020 Aug;584(7821):398–402.

53. Lima ÁLM, de Lima ID, Coutinho JFV, de Sousa ÚPST, Rodrigues MAG, Wilson ME, et al.
Changing epidemiology of visceral leishmaniasis in northeastern Brazil: a 25-year follow-up of an
urban outbreak. Trans R Soc Trop Med Hyg. 2017 Oct 1;111(10):440–7.

54. Vadmal GM, Glidden CK, Han BA, Carvalho B, Castellanos AA, Mordecai EA. Data-Driven
Predictions of Potential *Leishmania* Vectors in Latin America [Internet]. bioRxiv; 2022 [cited 2022
Oct 10]. p. 2022.08.22.504897. Available from:
https://www.biorxiv.org/content/10.1101/2022.08.22.504897v1

55. Kocher A, Cornuault J, Gantier JC, Manzi S, Chavy A, Girod R, et al. Biodiversity and vector-borne
diseases: Host dilution and vector amplification occur simultaneously for Amazonian leishmaniases.
Mol Ecol. 2022 Jan 9;

56. Silva J, Queiroz A, Moura I, Sousa RS, Guimarães LH, Machado PRL, et al. Dynamics of American
tegumentary leishmaniasis in a highly endemic region for Leishmania (Viannia) braziliensis
infection in northeast Brazil. PLoS Negl Trop Dis. 2017 Nov 2;11(11):e0006015.

57. Guzmán H, Tesh RB. Efectos de la temperatura y la dieta sobre el crecimiento y la longevidad de
flebotomíneos (Diptera: Psychodidae). Biomédica. 2000 Sep 1;20(3):190.

58. Hlavacova J, Votypka J, Volf P. The effect of temperature on *Leishmania* (Kinetoplastida:
Trypanosomatidae) development in sand flies. J Med Entomol. 2013 Sep;50(5):955–8.

59. Ostfeld RS, Levi T, Jolles AE, Martin LB, Hosseini PR, Keesing F. Life history and demographic
drivers of reservoir competence for three tick-borne zoonotic pathogens. PloS One.
2014;9(9):e107387.

60. Demas N, Nelson R. Ecoimmunology. Oxford, New York: Oxford University Press; 2011. 656 p.

61. Martin LB, Weil ZM, Nelson RJ. Immune defense and reproductive pace of life in *Peromyscus* mice.
Ecology. 2007 Oct;88(10):2516–28.

62. Schwanz LE. Persistent Effects of Maternal Parasitic Infection on Offspring Fitness: Implications for
Adaptive Reproductive Strategies When Parasitized. Funct Ecol. 2008;22(4):691–8.

63. Caldart ET, Pinto-Ferreira F, Matos AMRN de, Pascoal ATP, Bertão-Santos A, Mitsuka-Breganó R,
et al. Evaluation of an active and early surveillance methodology for visceral leishmaniasis by
molecular detection in road-killed wild fauna. Rev Bras Parasitol Veterinária. 2021;30(2):e027920.

64. Massey AL, Bronzoni RV de M, da Silva DJF, Allen JM, de Lázari PR, dos Santos-Filho M, et al.
Invertebrates for vertebrate biodiversity monitoring: Comparisons using three insect taxa as iDNA
samplers. Mol Ecol Resour. 2022;22(3):962–77.

65. Patiño LH, Castillo-Castañeda AC, Muñoz M, Jaimes JE, Luna-Niño N, Hernández C, et al.
Development of an Amplicon-Based Next-Generation Sequencing Protocol to Identify Leishmania
Species and Other Trypanosomatids in Leishmaniasis Endemic Areas. Microbiol Spectr. 2021 Oct
31;9(2):e0065221.

66. Richini-Pereira VB, Marson PM, Hayasaka EY, Victoria C, da Silva RC, Langoni H. Molecular
detection of *Leishmania spp.* in road-killed wild mammals in the Central Western area of the State
of São Paulo, Brazil. J Venom Anim Toxins Trop Dis. 2014 Jun 16;20:27.

67. Shender LA, De Los Santos M, Montgomery JM, Conrad PA, Ghersi BM, Razuri H, et al. Native
Rodent Species Are Unlikely Sources of Infection for *Leishmania (Viannia) braziliensis* along the
Transoceanic Highway in Madre de Dios, Peru. PLoS ONE. 2014 Jul 25;9(7):e103358.

68. Kowaltowski AJ. Brazil's scientists face 90% budget cut. Nature. 2021 Oct 25;598(7882):566–566.

69. Araujo Carreira JC, Magalhães M de AFM, Brazil RP, da Silva AVM. *Leishmania* in Marsupials—An
Overview of Infection Records in the Americas and Australia. Open J Anim Sci. 2017;07(03):315–
43.

70. Rosypal AC, Alexander A, Byrd D, Weaver M, Stewart R, Gerhold R, et al. SURVEY OF
ANTIBODIES TO *LEISHMANIA SPP*. IN WILD CANIDS FROM PENNSYLVANIA AND
TENNESSEE. J Zoo Wildl Med. 2013;44(4):1131–3.

71. Stephens CR, Heau JG, González C, Ibarra-Cerdeña CN, Sánchez-Cordero V, González-Salazar
C. Using biotic interaction networks for prediction in biodiversity and emerging diseases. PloS One.
2009 May 28;4(5):e5725.

72. Silva MB da, Portela JM, Li W, Jackson M, Gonzalez-Juarrero M, Hidalgo AS, et al. Evidence of
zoonotic leprosy in Pará, Brazilian Amazon, and risks associated with human contact or
consumption of armadillos. PLoS Negl Trop Dis. 2018 Jun 28;12(6):e0006532.

73. Barbosa PBBM, de Queiroz PVS, Jerônimo SMB, Ximenes M de FF de M. Experimental infection
parameters in *Galea spixii* (Rodentia: Caviidae) with *Leishmania infantum chagasi*. Mem Inst
Oswaldo Cruz. 2008 Sep;103(6):545–8.

74. Berzunza-Cruz M, Rodríguez-Moreno Á, Gutiérrez-Granados G, González-Salazar C, Stephens
CR, Hidalgo-Mihart M, et al. *Leishmania (L.) mexicana* Infected Bats in Mexico: Novel Potential
Reservoirs. PLoS Negl Trop Dis. 2015 Jan 28;9(1):e0003438.

75. Rosa RB, Costa MS da, Teixeira SC, Castro EF de, Dantas WM, Ferro EAV, et al. Calomys
callosus: An Experimental Animal Model Applied to Parasitic Diseases Investigations of Public
Health Concern. Pathog Basel Switz. 2022 Mar 17;11(3):369.

76. García N, Moreno I, Alvarez J, de la Cruz ML, Navarro A, Pérez-Sancho M, et al. Evidence of
Leishmania infantum Infection in Rabbits (*Oryctolagus cuniculus*) in a Natural Area in Madrid,
Spain. BioMed Res Int. 2014;2014:318254.

77. Karayiannis S, Ntais P, Messaritakis I, Tsirigotakis N, Dokianakis E, Antoniou M. Detection of
Leishmania infantum in red foxes (*Vulpes vulpes*) in Central Greece. Parasitology. 2015
Nov;142(13):1574–8.

78. Tsokana CN, Sokos C, Giannakopoulos A, Mamuris Z, Birtsas P, Papaspyropoulos K, et al. First
evidence of *Leishmania* infection in European brown hare (*Lepus europaeus*) in Greece: GIS
analysis and phylogenetic position within the *Leishmania spp*. Parasitol Res. 2016 Jan;115(1):313–
21.

79. Kocher A, Cornuault J, Gantier JC, Manzi S, Chavy A, Girod R, et al. Biodiversity and vector-borne
diseases: Host dilution and vector amplification occur simultaneously for Amazonian leishmaniases.
Mol Ecol [Internet]. 2022 [cited 2022 Aug 6];n/a(n/a). Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.16341

80. Curtin JM, Aronson NE. Leishmaniasis in the United States: Emerging Issues in a Region of Low
Endemicity. Microorganisms. 2021 Mar 11;9(3):578.

81. González C, Wang O, Strutz SE, González-Salazar C, Sánchez-Cordero V, Sarkar S. Climate
change and risk of leishmaniasis in North America: predictions from ecological niche models of
vector and reservoir species. PLoS Negl Trop Dis. 2010 Jan 19;4(1):e585.

82. McIlwee BE, Weis SE, Hosler GA. Incidence of Endemic Human Cutaneous Leishmaniasis in the
United States. JAMA Dermatol. 2018 Sep 1;154(9):1032–9.

834 **Supplemental materials**
835
836 **Supplementary text 1. Search methods for recording *Leishmania* host status:** We used the search
837 string listed below to run search Web of Science of *Leishmania* hosts. We ran the initial search on
838 February 12, 2021. We ran additional searches that included the species names of zoonotic *Leishmania*
839 on February 22, 2021. Our host database was then updated in June 2021 after the publication of (1).
840
Search string:
AB=(("leishmania*" OR "uta" OR "chiclero ulcer" OR "pian bois") AND ("PCR" OR "sero*" OR "serum" OR
"infected" OR "detected" OR "RT-PCR" OR "qPCR" OR "PCR" OR "chain reaction" OR "xenodiagnosis" OR
"culture" OR "isolat*" OR "reservoir") AND ("Central America" OR "South America" OR "Latin America" OR
"America*" OR "Antigua" OR "Barbuda" OR "Aruba" OR "Bahamas" OR "Barbados" OR "Grand Cayman" OR
"Cayman Islands" OR "Little Cayman" OR "Cayman Brac" OR "Cuba" OR "Dominica" OR "Dominican
Republic" OR "Grenada" OR "Guadeloupe" OR "Haiti" OR "Jamaica" OR "Martinique" OR "Puerto Rico" OR
"Saint Barthelemy" OR "Saint Kitts" OR "Nevis" OR "Saint Vincent" OR "Grenadines" OR "Canouan" OR
"Bequia" OR "Trinidad" OR "Tobago" OR "Virgin Islands" OR "Belize" OR "Costa Rica" OR "El Salvador" OR
"Guatemala" OR "Honduras" OR "Mexico" OR "Nicaragua" OR "Panama" OR "Argentina" OR "Bolivia" OR
"Brazil" OR "Chile" OR "Colombia" OR "Ecuador" OR "Guyana" OR "Paraguay" OR "Peru" OR "Suriname"
OR "Uruguay" OR "Venezuela") AND ("animal" OR "vertebrate" OR "host" OR "wildlife" OR "livestock" OR
"domestic"))
854
855 **References**

856   1. Azami-Conesa I, Gómez-Muñoz MT, Martínez-Díaz RA. A Systematic Review (1990–2021) of
857   Wild Animals Infected with Zoonotic *Leishmania*. Microorganisms. 2021 May 20;9(5):1101.

860   **Supplementary Table 1.** Final traits and sources of traits used in the analysis.

trait	definition	source
neonate body mass (g)	mass of neonates	panTHERIA
adult forearm length (mm)	total length from elbow to wrist	panTHERIA
gestation length (d)	length of time of non-inactive fetal growth	panTHERIA
litter size	average # of offspring in a litter	panTHERIA
litters per year	average # of litters per year	panTHERIA
max longevity (m)	maximum adult age	panTHERIA
social group size (n)	# of individuals in a group	panTHERIA
weaning age (d)	age when primary nutritional dependency on the mother ends and independent foraging begins to make a major contribution to the offspring's energy	panTHERIA
activity cycle	activity cycle (nocturnal, crepuscular, diurnal)	panTHERIA
dominate diet	primary food group consumed by the animal, including groups related to sandfly habitat and exposure (insectivore, frugivore, granivore, folivore)	mammalDiet
diet breadth	number of dietary categories eaten by each species	panTHERIA
home range size (km^2)	size of the area within which everyday activities of individuals or groups (of any type) are typically restricted	panTHERIA
population density (n/km^2)	population size per km^2	panTHERIA
foraging strata	assignment to one of five foraging stratum categories (ground level, scansorial, arboreal, aerial)	EltonTraits
trophic strata	assignment to one of three trophic levels: carnivore, omnivore, herbivore	EltonTraits
temperature seasonality		CHELSA
temperature in the warmest quarter		CHELSA
precipitation in the driest quarter		CHELSA
humidity range		CHELSA

climate moisture index range		CHELSA
main habitat	Main habitat defined by IUCN, includes forest, savanna, shrubland, grassland, wetland, rocky area, desert, artificial	IUCN
habitat breadth	number of IUCN defined main habitats	
percent crop cover	average percent of the range covered by crop land from 2015-2019	IUCN, gee[a]: COPERNICUS/Landcover/100m/Proba-V-C3/Global
percent tree cover	average percent of the range covered by trees from 2015-2019	IUCN, gee[a]: COPERNICUS/Landcover/100m/Proba-V-C3/Global
percent urban cover	average percent of the range covered by urban landscape from 2015-2019	IUCN, gee[a]: COPERNICUS/Landcover/100m/Proba-V-C3/Global
global human modification index	aggregate measure of the human modification; mean gHm within an animal's range in 2016	IUCN, gee[a]: CSP/HM/GlobalHumanModification
range area (km^2)		panTHERIA
maximum longitude of range (dd)		panTHERIA
minimum longitude of range (dd)		panTHERIA
median latitude of range (dd)		panTHERIA
phylogenetic distance	Pairwise divergence time, averaged across all available sources for species pairs, reduced into five dimensions via PCoA	TimeTree
invasive	invasive in Mexico, Central America, or South America	GISD
zoonotic host	reservoir host of any zoonotic pathogen	GIDEON
citation count	number of citations found on pubmed	pubmed
[a] google earth engine		

861
862

867 **Supplementary Table 2.** Summed Shapley Scores per species for newly predicted species, averages
864 across 100 model iterations (0.05 percentile - 0.95 percentile). Grey boxes represent animals that are
865 already known hosts for the subgenera.

866
867

Binomial	Order	*L. (Viannia)* scores	*L. (Leishmania)* scores	Reference
Odocoileus virginianus	Artiodactyla	1.40 (0.01-2.58)	1.46 (0.54-2.17)	
Pecari tajacu	Artiodactyla	--	2.28 (1.84-2.83)	(1)
Sus scrofa	Artiodactyla	--	1.49 (0.38-2.34)	
Tayassu pecari	Artiodactyla	--	2.22 (1.70-2.92)	
Bassariscus astutus	Carnivora	--	0.74 (0.31-1.75)	
Canis latrans	Carnivora	--	1.39 (0.75-1.92)	
Canis lupus	Carnivora	0.86 (0.18-1.45)	1.84 (1.02-2.73)	
Herpailurus yagouaroundi	Carnivora	0.86 (0.24-1.57)	2.20 (1.57-2.82)	
Leopardus wiedii	Carnivora	0.89 (0.26-1.61)	2.74 (2.22-3.14)	
Lontra longicaudis	Carnivora	--	1.74 (1.05-2.35)	
Mustela frenata	Carnivora	--	1.64 (0.88-2.44)	
Nasua narica	Carnivora	1.29 (0.10-2.30)	1.48 (0.70-2.19)	
Potos flavus	Carnivora	1.77 (1.02-2.37)	--	(2)
Procyon cancrivorus	Carnivora	--	2.34 (1.57-3.01)	
Procyon lotor	Carnivora	1.02 (0.19-2.25)	1.95 (1.21-2.65)	
Puma concolor	Carnivora	0.89 (0.23-1.69)	--	(3)
Vulpes vulpes	Carnivora	--	1.91 (1.31-2.63)	
Anoura caudifer	Chiroptera	--	1.49 (0.43-2.23)	
Anoura geoffroyi	Chiroptera	1.23 (0.21-2.31)	2.15 (1.48-2.86)	
Artibeus phaeotis	Chiroptera	1.05 (0.54-1.73)	0.97 (0.27-1.45)	

Carollia perspicillata	Chiroptera	2.16 (1.43-2.87)	--	(4)
Chiroderma villosum	Chiroptera	--	0.98 (0.22-1.63)	
Choeronycteris mexicana	Chiroptera	0.60 (0.10-1.29)	--	
Chrotopterus auritus	Chiroptera	--	1.75 (0.57-2.55)	
Diclidurus albus	Chiroptera	1.11 (0.37-1.94)	1.43 (0.45-2.34)	
Diphylla ecaudata	Chiroptera	--	1.79 (0.82-2.42)	
Eptesicus brasiliensis	Chiroptera	--	0.95 (0.23-1.73)	
Eptesicus furinalis	Chiroptera	1.01 (0.09-1.87)	--	
Eptesicus fuscus	Chiroptera	1.11 (0.31-1.86)	1.41 (0.72-1.94)	
Lasionycteris noctivagans	Chiroptera	1.41 (0.73-2.09)	1.10 (0.28-1.70)	
Lasiurus borealis	Chiroptera	1.04 (0.21-1.99)	1.24 (0.26-2.17)	
Lasiurus cinereus	Chiroptera	--	1.33 (0.64-1.89)	(5)
Lasiurus ega	Chiroptera	--	2.11 (1.36-2.86)	
Lasiurus intermedius	Chiroptera	--	0.84 (0.18-1.78)	
Leptonycteris yerbabuenae	Chiroptera	0.73 (0.07-1.38)	--	
Macrophyllum macrophyllum	Chiroptera	--	1.04 (0.59-1.64)	
Macrotus waterhousii	Chiroptera	--	0.83 (0.15-1.49)	
Micronycteris buriri	Chiroptera	--	0.86 (0.38-1.39)	
Mormoops megalophylla	Chiroptera	--	1.16 (0.51-1.93)	
Myotis albescens	Chiroptera	--	1.66 (0.56-2.38)	
Myotis riparius	Chiroptera	--	0.88 (0.31-1.64)	
Natalus stramineus	Chiroptera	--	1.08 (0.23-1.97)	
Noctilio albiventris	Chiroptera	--	1.86 (1.12-2.54)	
Noctilio leporinus	Chiroptera	--	1.21 (0.48-2.12)	

Nyctinomops aurispinosus	Chiroptera	--	1.07 (0.51-1.87)	
Perimytotis subflavus	Chiroptera	1.41 (0.65-1.98)	1.25 (0.08-1.88)	
Peropteryx macrotis	Chiroptera	--	1.14 (0.29-1.78)	
Phyllostomus hastatus	Chiroptera	1.34 (0.54-2.08)	--	(6)
Pteronotus parnellii	Chiroptera	1.43 (0.66-2.19)	--	(7)
Pteronotus quadridens	Chiroptera	0.82 (0.07-1.52)	--	
Saccopteryx bilineata	Chiroptera	--	1.88 (1.08-2.51)	
Sturnira lilium	Chiroptera	2.19 (1.19-3.02)	--	(8)
Sturnira tildae	Chiroptera	--	1.12 (0.25-2.11)	
Tadarida brasiliensis	Chiroptera	1.70 (0.86-2.53)	2.33 (1.93-2.73)	
Thyroptera tricolor	Chiroptera	--	1.25 (0.20-2.25)	
Trachops cirrhosus	Chiroptera	--	0.77 (0.13-1.58)	
Uroderma bilobatum	Chiroptera	1.28 (0.31-2.06)	1.82 (0.93-2.54)	
Dasypus novemcinctus	Cingulata	--	3.05 (2.57-3.42)	(9)
Euphractus sexcinctus	Cingulata	0.90 (0.22-1.61)	1.98 (1.32-2.69)	
Priodontes maximus	Cingulata	0.82 (0.08-1.40)	--	
Caluromys derbianus	Didelphimorphia	2.04 (1.06-2.85)	1.23 (0.67-1.82)	
Caluromys philander	Didelphimorphia	--	2.06 (1.33-2.68)	(10)
Chironectes minimus	Didelphimorphia	--	1.28 (0.37-1.99)	
Didelphis virginiana	Didelphimorphia	2.28 (1.52-3.21)	1.77 (0.91-2.62)	
Gracilinanus agilis	Didelphimorphia	--	1.17 (0.28-2.00)	(11)
Lutreolina crassicaudata	Didelphimorphia	--	1.41 (0.18-2.22)	
Marmosa murina	Didelphimorphia	1.51 (0.32-2.38)	--	(10)
Metachirus nudicaudatus	Didelphimorphia	1.67 (0.76-2.54)	--	(10)

Species	Order	Col1	Col2	Ref
Monodelphis brevicaudata	Didelphimorphia	1.53 (0.38-2.54)	--	
Monodelphis domestica	Didelphimorphia	1.56 (0.55-2.57)	1.85 (0.89-2.59)	
Thylamys karimii	Didelphimorphia	0.95 (0.00-1.89)	--	
Tlacuatzin canescens	Didelphimorphia	1.03 (0.14-1.91)	--	
Lepus europaeus	Lagomorpha	--	1.25 (0.24-2.07)	
Oryctolagus cuniculus	Lagomorpha	0.95 (0.20-2.07)	0.95 (0.18-1.66)	
Romerolagus diazi	Lagomorpha	0.96 (0.13-1.90)	--	
Sylvilagus audubonii	Lagomorpha	1.36 (0.54-2.12)	1.20 (0.49-2.06)	
Sylvilagus brasiliensis	Lagomorpha		1.67 (0.63-2.29)	(7)
Sylvilagus palustris	Lagomorpha	--	0.79 (0.11-1.56)	
Tapirus terrestris	Perissodactyla	--	1.22 (0.54-1.81)	
Bradypus variegatus	Pilosa		1.85 (1.15-2.81)	(12)
Cyclopes didactylus	Pilosa	2.14 (1.15-2.99)	1.66 (0.86-2.37)	
Myrmecophaga tridactyla	Pilosa	1.36 (0.42-2.22)		(7)
Ateles geoffroyi	Primates	1.88 (1.00-2.61)	0.97 (0.16-1.69)	
Callithrix jacchus	Primates	1.10 (0.24-1.84)		(7)
Cebus capucinus	Primates	1.52 (0.30-2.63)	--	
Erythrocebus patas	Primates	--	1.11 (0.33-2.16)	
Leontocebus fuscicollis	Primates	0.92 (0.14-1.67)	--	
Macaca mulatta	Primates	--	1.10 (0.31-2.09)	
Saguinus oedipus	Primates	1.67 (0.72-2.65)	1.04 (0.12-2.20)	
Saimiri oerstedii	Primates	1.15 (0.12-2.30)	--	
Saimiri sciureus	Primates	1.00 (0.38-1.68)	--	
Sapajus apella	Primates	--	0.91 (0.07-1.99)	

Calomys callosus	Rodentia	2.32 (1.61-3.09)	2.15 (1.24-3.02)	
Calomys expulsus	Rodentia	0.84 (0.09-1.69)	--	
Cavia aperea	Rodentia	1.11 (0.52-2.03)	--	(7)
Cerradomys subflavus	Rodentia	--	1.40 (0.13-2.27)	(13)
Cuniculusi paca	Rodentia	--	2.45 (1.77-3.18)	
Dasyprocta leporina	Rodentia	0.87 (0.03-2.09)	1.46 (0.57-2.22)	
Dasyprocta punctata	Rodentia	1.54 (0.57-2.82)	0.73 (0.07-1.41)	
Dipodomys merriami	Rodentia	--	1.68 (0.86-2.41)	
Dipodomys ordii	Rodentia	1.46 (0.59-2.22)	1.45 (0.63-2.43)	
Galea spixii	Rodentia	--	2.34 (1.76-2.96)	
Glaucomys volans	Rodentia	1.37 (0.64-2.07)	0.77 (0.01-1.68)	
Heteromys salvini	Rodentia	1.53 (0.73-2.53)	1.77 (1.02-2.45)	
Holochilus brasiliensis	Rodentia	--	0.43 (0.07-0.89)	
Hydrochoerus hydrochaeris	Rodentia	1.74 (0.63-2.59)	--	(7)
Hylaeamys megacephalus	Rodentia	2.64 (1.72-3.29)	2.44 (1.90-3.09)	
Kerodon rupestris	Rodentia	--	1.30 (0.50-2.01)	
Myocastor coypus	Rodentia	0.78 (0.18-1.48)	0.79 (0.22-1.45)	
Myoprocta acouchy	Rodentia	--	0.94 (0.03-1.96)	
Myoprocta pratti	Rodentia	1.54 (0.52-2.29)	--	
Neotoma mexicana	Rodentia	1.17 (0.29-1.78)	--	
Neotomodon alstoni	Rodentia	1.11 (0.41-1.96)	--	
Ochrotomys nuttalli	Rodentia	2.08 (1.38-2.73)	1.66 (1.22-2.33)	
Oligoryzomys fornesi	Rodentia	1.21 (0.54-2.00)	--	
Oligoryzomys fulvescens	Rodentia	1.63 (0.84-2.48)	1.52 (0.90-2.25)	

Oligoryzomys nigripes	Rodentia	0.75 (0.13-1.36)	--	
Oryzomys couesi	Rodentia	1.42 (0.72-2.24)	1.22 (0.39-2.00)	
Oryzomys palustris	Rodentia	2.16 (1.40-2.83)	1.70 (1.13-2.35)	
Otonyctomys hatti	Rodentia	0.83 (0.12-1.74)	--	
Peromyscus aztecus	Rodentia	0.91 (0.09-2.08)	--	
Peromyscus gossypinus	Rodentia	2.11 (1.75-2.65)	1.62 (1.20-2.09)	
Proechimys guairae	Rodentia	1.23 (0.52-1.92)	1.69 (0.82-2.30)	
Reithrodontomys fulvescens	Rodentia	1.12 (0.26-1.94)	1.01 (0.29-1.65)	
Reithrodontomys humulis	Rodentia	1.79 (1.10-2.74)	1.43 (0.74-2.38)	
Reithrodontomys megalotis	Rodentia	1.39 (0.65-2.03)	0.89 (0.43-1.69)	
Rhipidomys mastacalis	Rodentia	1.39 (0.81-2.24)	--	(11)
Sciurus granatensis	Rodentia	1.43 (0.60-2.40)	--	
Sciurus variegatoides	Rodentia	--	0.93 (0.14-1.76)	
Sciurus vulgaris	Rodentia	1.48 (0.46-2.25)	--	(2)
Scotinomys teguina	Rodentia	0.94 (0.29-1.92)	--	
Spermophilus variegatus	Rodentia	0.91 (0.16-1.84)	--	
Tylomys nudicaudus	Rodentia	1.58 (0.50-2.29)	0.84 (0.12-1.67)	
Zygodontomys brevicauda	Rodentia	1.87 (1.16-2.68)	1.07 (0.27-1.84)	
Cryptotis parva	Soricomorpha	1.40 (0.50-2.51)	0.99 (0.31-1.59)	
Scalopus aquaticus	Soricomorpha	1.78 (1.19-2.46)	1.64 (0.84-2.25)	

Supplementary Table 2 References

869
1. Rojas-Jaimes JE, Correa-Nuñez GH, Rojas N, Cáceres-Rey O. Detection of *Leishmania (V) guyanensis* in Rhipicephalus
(Boophilus) microplus (Acari: Ixodidae) collected from Pecari tajacu. Biomed Rev Inst Nac Salud. 2017 Mar 29;37(0):208–
14.

2. Mimori T, Grimaldi G, Kreutzer RD, Gomez EA, McMahon-Pratt D, Tesh RB, et al. Identification, using isoenzyme
electrophoresis and monoclonal antibodies, of *Leishmania* isolated from humans and wild animals of Ecuador. Am J Trop
Med Hyg. 1989 Feb 1;40(2):154–8.

3. Dahroug MAA, Almeida ABPF, Sousa VRF, Dutra V, Turbino NCMR, Nakazato L, et al. *Leishmania (Leishmania) chagasi*
in captive wild felids in Brazil. Trans R Soc Trop Med Hyg. 2010 Jan 1;104(1):73–4.

4. De Lima H, Rodríguez N, Barrios MA, Avila A, Cañizales I, Gutiérrez S. Isolation and molecular identification of
Leishmania chagasi from a bat (*Carollia perspicillata*) in northeastern Venezuela. Mem Inst Oswaldo Cruz. 2008
Jun;103(4):412–4.

5. Castro LS, Dorval MEC, Matheus LMD, Bednaski AV, Facco GG, Silveira M, et al. *Leishmania* presence in bats in areas
endemic for leishmaniasis in central-west Brazil. Int J Parasitol Parasites Wildl. 2020 Apr 1;11:261–7.

6. Gómez-Hernández C, Bento EC, Rezende-Oliveira K, Nascentes G a. N, Barbosa CG, Batista LR, et al. *Leishmania*
infection in bats from a non-endemic region of Leishmaniasis in Brazil. Parasitology. 2017 Dec;144(14):1980–6.

7. Azami-Conesa I, Gómez-Muñoz MT, Martínez-Díaz RA. A Systematic Review (1990–2021) of Wild Animals Infected with
Zoonotic *Leishmania*. Microorganisms. 2021 May 20;9(5):1101.

8. Savani ESMM, de Almeida MF, de Oliveira Camargo MCG, D'Auria SRN, Silva MMS, de Oliveira ML, et al. Detection of
Leishmania (Leishmania) amazonensis and *Leishmania (Leishmania) infantum chagasi* in Brazilian bats. Vet Parasitol.
2010 Feb 26;168(1):5–10.

9. Lainson R, Shaw JJ, Ward RD, Ready PD, Naiff RD. Leishmaniasis in Brazil: XIII. Isolation of *Leishmania* from armadillos
(*Dasypus novemcinctus*), and observations on the epidemiology of cutaneous leishmaniasis in north Pará State. Trans R
Soc Trop Med Hyg. 1979 Jan 1;73(2):239–42.

10. Araujo Carreira JC, Magalhães M de AFM, Brazil RP, da Silva AVM. *Leishmania* in Marsupials—An Overview of Infection
Records in the Americas and Australia. Open J Anim Sci. 2017;07(03):315–43.

11. Quaresma PF, Rêgo FD, Botelho HA, da Silva SR, Moura AJ Júnior, Neto RGT, et al. Wild, synanthropic and domestic
hosts of *Leishmania* in an endemic area of cutaneous leishmaniasis in Minas Gerais State, Brazil. Trans R Soc Trop Med
Hyg. 2011 Oct 1;105(10):579–85.

12. Muñoz-García CI, Sánchez-Montes S, Villanueva-García C, Romero-Callejas E, Díaz-López HM, Gordillo-Chávez EJ, et
al. The role of sloths and anteaters as *Leishmania spp.* reservoirs: a review and a newly described natural infection of
Leishmania mexicana in the northern anteater. Parasitol Res. 2019 Apr;118(4):1095–101.

13. Tonelli GB, Tanure A, Rego FD, Carvalho GM de L, Stumpp R, Ássimos GR, et al. *Leishmania (Viannia) braziliensis*
infection in wild small mammals in ecotourism area of Brazil. PLoS ONE. 2017 Dec 28;12(12):e0190315.

Supplementary Table 3. Complementary evidence of predicted hosts. Complementary evidence is defined as published literature that suggests animals may be exposed to and/or infected by zoonotic *Leishmania*, but was not robust enough to include the animal in the analysis as a known host. P = predicted by our model; K = known host (the animal has been naturally infected by a *Leishmania* species known to cause leishmaniasis in humans.

Binomial	Common name	*L. (V)* host	*L. (L)* host	complementary evidence	References
Calomys callosus	Large vesper mouse	P	P		(1)
Galea spixii	Spix's yellow-toothed cavy	--	P	experimental infection	(2)
Tadarida brasiliensis	Mexican free-tailed bat	P	P		(3)
Didelphis virginiana	Virginia opossum	P	P	experimental infection; significant overlap w/ *L. mexicana* vectors	(4–6)
Vulpes vulpes	Red fox	--	P		(7)
Lepus europaeus	European hare	--	P	*L. infantum* host in Europe	(8)
Oryctolagus cuniculus	European rabbit	P	P		(9)
Sus scrofa	Wild boar	--	P		(10)
Canis latrans	Coyote	--	P		(11)
Lontra longicaudis	Neotropical otter	--	P	*Leishmania* infection or exposure but not confirmed as zoonotic	(12)
Procyon cancrivorus	Crab-eating racoon	--	P		(13)

Monodelphis domestica	Gray short-tailed opossum	P	P		(4)
Euphractus sexcinctus	Six-banded armadillo	P	P		(14)
Cavia aperea	Brazilian guinea pig	P	K		(15)
Hydrochoerus hydrochaeris	Capybara	P	K		(15)
Sciurus granatensis	Red-tailed squirrel	P	--		(12)
Caluromys derbianus	Derby's wooly opossum	P	P	*Leishmania* infection but not confirmed as zoonotic; vector feeds on animal	(4)
Procyon lotor	Common racoon	P	P	significant overlap w/ *L. mexicana* vectors	(5,6)
Chiroderma villosum	Hairy big-eyed bat	--	P		
Eptesicus brasiliensis	Brazilian brown bat	--	P		
Saccopteryx bilineata	Greater sac-winged bat	--	P		
Uroderma bilobatum	Tent-making bat	P	P		
Cuniculus paca	Lowland paca	K	P		
Neotoma mexicana	Mexican woodrat	P	--		
Oligoryzomys fulvescens	Fulvous pygmy rice rat	P	P		
Oryzomys nigripes		P	--		
Oryzomys couesi	Coues's rice rat	P	P		

Reithrodontomys fulvescens	Fulvous harvest mouse	P	P		
Potos flavus	Kinkajou	P	K		
Nasua narica	White-nosed coati	P	P		
Leopardus wiedii	Margay	P	P		
Dasyprocta leporina	Red-rumped agouti	P	P	vector feeds on animal	(16)

Supplementary Table 3 References

1. Rosa RB, Costa MS da, Teixeira SC, Castro EF de, Dantas WM, Ferro EAV, et al. *Calomys callosus*: An Experimental Animal Model Applied to Parasitic Diseases Investigations of Public Health Concern. Pathog Basel Switz. 2022 Mar 17;11(3):369.

2. Barbosa PBBM, de Queiroz PVS, Jerônimo SMB, Ximenes M de FF de M. Experimental infection parameters in *Galea spixii* (Rodentia: Caviidae) with *Leishmania infantum chagasi*. Mem Inst Oswaldo Cruz. 2008 Sep;103(6):545–8.

3. Berzunza-Cruz M, Rodríguez-Moreno Á, Gutiérrez-Granados G, González-Salazar C, Stephens CR, Hidalgo-Mihart M, et al. *Leishmania (L.) mexicana* Infected Bats in Mexico: Novel Potential Reservoirs. PLoS Negl Trop Dis. 2015 Jan 28;9(1):e0003438.

4. Araujo Carreira JC, Magalhães M de AFM, Brazil RP, da Silva AVM. Leishmania in Marsupials—An Overview of Infection Records in the Americas and Australia. Open J Anim Sci. 2017;07(03):315–43.

5. Stephens CR, González-Salazar C, Sánchez-Cordero V, Becker I, Rebollar-Tellez E, Rodríguez-Moreno Á, et al. Can You Judge a Disease Host by the Company It Keeps? Predicting Disease Hosts and Their Relative Importance: A Case Study for Leishmaniasis. PLoS Negl Trop Dis. 2016 Oct 7;10(10):e0005004.

6. Stephens CR, Heau JG, González C, Ibarra-Cerdeña CN, Sánchez-Cordero V, González-Salazar C. Using biotic interaction networks for prediction in biodiversity and emerging diseases. PloS One. 2009 May 28;4(5):e5725.

7. Karayiannis S, Ntais P, Messaritakis I, Tsirigotakis N, Dokianakis E, Antoniou M. Detection of *Leishmania infantum* in red foxes (*Vulpes vulpes*) in Central Greece. Parasitology. 2015 Nov;142(13):1574–8.

8. Tsokana CN, Sokos C, Giannakopoulos A, Mamuris Z, Birtsas P, Papaspyropoulos K, et al. First evidence of *Leishmania infection* in European brown hare (*Lepus europaeus*) in Greece: GIS analysis and phylogenetic position within the *Leishmania spp*. Parasitol Res. 2016 Jan;115(1):313–21.

9. García N, Moreno I, Alvarez J, de la Cruz ML, Navarro A, Pérez-Sancho M, et al. Evidence of *Leishmania infantum* Infection in Rabbits (*Oryctolagus cuniculus*) in a Natural Area in Madrid, Spain. BioMed Res Int. 2014;2014:318254.

10. Brazil RP, Desterro MD, Nascimento SB, Macau RP. Natural infection of a pig (*Sus scrofa*) by *Leishmania* in a recent focus of cutaneous leishmaniasis on the Island of São Luis, Maranhão. Mem Inst Oswaldo Cruz. 1987 Mar;82(1):145.

11. Rosypal AC, Alexander A, Byrd D, Weaver M, Stewart R, Gerhold R, et al. Survey Of Antibodies To *Leishmania spp.* In Wild Canids From Pennsylvania And Tennessee. J Zoo Wildl Med. 2013;44(4):1131–3.

12. Azami-Conesa I, Gómez-Muñoz MT, Martínez-Díaz RA. A Systematic Review (1990–2021) of Wild Animals Infected with Zoonotic *Leishmania*. Microorganisms. 2021 May 20;9(5):1101.

13. Richini-Pereira VB, Marson PM, Hayasaka EY, Victoria C, da Silva RC, Langoni H. Molecular detection of *Leishmania spp*. in road-killed wild mammals in the Central Western area of the State of São Paulo, Brazil. J Venom Anim Toxins Trop Dis. 2014 Jun 16;20:27.

14. de Oliveira Barbosa W, Coelho TG, da Costa TO, Paiz LM, Fornazari F, Langoni H, et al. Antibodies to *Toxoplasma gondii*, *Leishmania spp.,* and *Leptospira spp*. in Free-Ranging Six-Banded Armadillos (*Euphractus sexcinctus*) from Northeastern Brazil. J Wildl Dis. 2020 Apr;56(2):486–8.

15. Caldart ET, Pinto-Ferreira F, Matos AMRN de, Pascoal ATP, Bertão-Santos A, Mitsuka-Breganó R, et al. Evaluation of an active and early surveillance methodology for visceral leishmaniasis by molecular detection in road-killed wild fauna. Rev Bras Parasitol Veterinária. 2021;30(2):e027920.

16. Kocher A, de Thoisy B, Catzeflis F, Valière S, Bañuls AL, Murienne J. iDNA screening: Disease vectors as vertebrate samplers. Mol Ecol. 2017;26(22):6478–86.

Supplementary figure 1. Importance of each PCoA phylogenetic dimension in predicting *L. (Viannia)* host status. Points are the mean Shapley importance for the PCoA dimension across all mammals, bars represent the 0.05-0.95 percentiles.

Supplementary figure 2. (a) Shapley partial dependence plots showing the effect of PCoA dimensions on *L.* (*Viannia*) host status. Blue non-linear lines represent the average effect across model iterations , while grey lines show each individual model iteration (model fit with 70% of data). Dark blue vertical line represents a threshold value, where the dimension switches from having a negative effect on host status to a positive effect on host status. Only dimensions with importance scores > 0 for > 95% of model iterations are shown. **(b)** distribution of hosts within each order with PCoA dimension values above or below the threshold.

Supplementary figure 3. Features inconsistently contribute to predictions for study effort of mammals included in the *L. (Viannia)* model. Global feature contribution for traits predicting study effort of mammals included in the *L. (Viannia)* model Points are the absolute value of the mean Shapley importance (mean |SHAP value|) for the trait across all mammals, bars represent the absolute values of the 0.05-0.95 percentiles. No trait consistently contributes to model predictions across all model bootstrapping iterations (i.e., there is no trait where the 0.05 percentile is greater than 0). Further, traits with average values > 0 do not correspond to traits that consistently predict *L. (Viannia)* host status.

Supplementary figure 4. Response of study effort predictions to mammal traits are inconsistent and do not mirror functional forms of response of *L. (Viannia)* host status to mammal traits. Shapley partial dependence plots showing the effect of mammal traits on study effort of mammals included in the *L. (Viannia)* model. Blue non-linear lines represent the average effect across model iterations, while grey lines show each individual model iteration (model fit with 70% of data). Functional form varies greatly across model iterations. Only traits where the absolute value of average contribution was > 0 are shown. Rug plots on the x-axis indicate the distribution of the data.

Supplementary figure 5. Features inconsistently contribute to predictions for study effort of mammals included in the *L. (Leishmania)* model. Global feature contribution for traits predicting study effort of mammals included in the *L. (Leishmania)* model. Points are the absolute value of the mean Shapley importance (mean |SHAP value|) for the trait across all mammals, bars represent the absolute values of the 0.05-0.95 percentiles. No trait consistently contributes to model predictions across all model bootstrapping iterations (i.e., there is no trait where the 0.05 percentile is greater than 0). Further, traits with average values > 0 do not correspond to traits that consistently predict *L. (Leishmania)* host status.

Supplementary figure 6. Response of study effort predictions to mammal traits are inconsistent and do not mirror functional forms of response of *L. (Leishmania)* host status to mammal traits. Shapley partial dependence plots showing the effect of mammal traits on study effort of mammals included in the *L. (Leishmania)* model. Blue non-linear lines represent the average effect across model iterations, while grey lines show each individual model iteration (model fit with 70% of data). Functional form varies greatly across model iterations. Only traits where the absolute value of average contribution was > 0 are shown. Rug plots on the x-axis indicate the distribution of the data.