Quantitative dose-response analysis untangles host bottlenecks to enteric infection

Ian W. Campbell, Karthik Hullahalli, Jerrold R. Turner, and Matthew K. Waldor

Division of Infectious Diseases, Brigham & Women's Hospital, Boston

Division of Pathology, Brigham & Women's Hospital, Boston

Department of Microbiology, Harvard Medical School, Boston

For correspondence: Matthew K. Waldor mwaldor[at]research.bwh.harvard.edu or Ian W. Campbell

icampbell3[at]bwh.harvard.edu

18 **Abstract**
19
Host bottlenecks prevent many infections before the onset of disease by eliminating invading pathogens.
Monitoring the diversity of a barcoded population of the diarrhea causing bacterium *Citrobacter rodentium*
during colonization of its natural host, mice, allowed us to determine the number of cells that found the
infection by establishing a replicative niche. The size of the pathogen's founding population scaled with dose
and was controlled by a severe yet slow-acting bottleneck. Reducing stomach acid or changing host genotype
modestly relaxed the bottleneck without breaking the fractional relationship between dose and founders. In
contrast, disrupting the microbiota caused the founding population to no longer scale with the size of the
inoculum and allowed the pathogen to infect at almost any dose, indicating that the microbiota creates the
dominant bottleneck. Further, in the absence of competition with the microbiota, the diversity of the pathogen
population slowly contracted as the population was overtaken by bacteria that lost the critical virulence island,
the locus of enterocyte effacement (LEE). Collectively, our findings reveal that the mechanisms of protection by
colonization bottlenecks are reflected in and can be generally defined by the impact of dose on the pathogen's
founding population.

Introduction

Diarrhea-causing pathogens remain a global threat to human health, with billions of cases impacting over half of the world's population every year (Troeger *et al.*, 2018). These pathogens are typically transmitted in food, water, and between infected individuals by the fecal-oral route. However, exposure does not always result in the pathogen establishing a replicative niche in the host, a prerequisite for colonization and thus most disease.

As a natural mouse pathogen, the gram-negative bacterium *Citrobacter rodentium* provides a particularly valuable experimental model to identify the factors governing the earliest events that result in colonization of the mammalian gastrointestinal tract. This enteric pathogen proliferates in the cecum and colon, causes diarrhea, and naturally transmits between mice via coprophagy (Mundy *et al.*, 2005). *C. rodentium* shares an infection strategy with the human extracellular pathogens enteropathogenic and enterohemorrhagic *Escherichia coli* (EPEC and EHEC). To colonize the gastrointestinal tract these pathogens depend on a conserved pathogenicity island known as the locus of enterocyte effacement (LEE), which encodes a type three secretion system that mediates pathogen adherence through creation of attaching and effacing (A/E) lesions on host epithelial cells in the colon (Collins *et al.*, 2014).

While extensive work has focused on the molecular functions of the LEE and the disease caused by A/E pathogens, less is known about the early events that result in establishment of a replicative niche. Dosing studies indicate that ingestion of between a million and a billion bacteria are required for *C. rodentium* to infect mice (Wiles *et al.*, 2005), indicating that one or more barriers likely impede colonization. These barriers collectively create an infection bottleneck by causing a stochastic contraction the pathogen population. However, the bottlenecks restricting A/E pathogens have not yet been fully characterized because the dramatic expansion of the pathogen population that follows establishment of a replicative niche makes it experimentally difficult to quantify the earliest events required for infection.

Here we inoculated mice with a barcoded, but otherwise isogenic population of *C. rodentium* to differentiate the effects of the initial colonization bottleneck from subsequent population expansion. By monitoring the diversity of barcodes in the population we determined the number of unique cells from the inoculum (founders) which expand to cause infection (schematized in Figure 1A). Quantifying the size of the pathogen founding population allowed us to define the bottleneck to colonization and determine when/where the invading pathogen establishes a replicative niche. We discovered that a severe bottleneck restricts colonization and results in a very small number of founders that increase with the size of the inoculum. Multiple factors contribute to the infection bottleneck with the greatest restriction occurring after the pathogen reaches the sites of infection. Notably, the magnitude of the infection bottleneck is largely explained by a combination of stomach acid and microbiota, where the latter is approximately a million times more protective.

Results

A small number of *C. rodentium* founders initiates enteric infection

To enable monitoring of the pathogen population's diversity during infection we introduced short, random, ~20 nucleotide DNA tags (barcodes) at a neutral location in the *C. rodentium* genome. As previously described (Abel *et al.*, 2015a), monitoring barcode diversity using high-throughput DNA sequencing and the STAMP (Sequence Tag-based Analysis of Microbial Populations) computational framework can quantify the constriction of the pathogen population that often occurs during establishment of infection (schematized in Figure 1A). We created two independent STAMP libraries of barcoded bacteria. Library "STAMP-CR253" contains 253 unique barcodes integrated in the intergenic region between genes ROD_05521 and *selU*. The neutrality of the barcode insertions was confirmed by measuring growth in lysogeny broth (LB; Supplemental figure 1A). Library "STAMP-CR69K" contains approximately 69,000 unique barcodes inserted into the genome on a Tn7 vector, which integrates at a neutral site downstream of the *glmS* gene (Choi *et al.*, 2005; Fakoya *et al.*, 2022). While the libraries were not directly compared, both yielded similar results in our studies.

To validate that these barcoded STAMP libraries can quantify the population effects of a bottleneck, we created *in vitro* bottlenecks by plating serial dilutions of the libraries grown in culture. The number of colony forming units (CFU) per plate provides a true measure of the number of founders, i.e., the number of cells from the initial population (culture) that gave rise to the observed population (plated colonies). Bacteria were harvested from the plates, the barcodes were amplified and sequenced, and barcode frequencies were analyzed using the recently updated STAMP analysis pipeline "STAMPR" (Hullahalli *et al.*, 2021). The size of the founding population (founders) was calculated by comparing the diversity and frequency of barcodes recovered from plated samples to those in the initial cultures. There was a strong correlation between the counted founders (CFU) and the calculated founders (N_r for STAMP-CR253 and N_s for STAMP-CR69K) up to 10^4 founders (Supplemental figure 1B). These data were also used as standard curves to increase the resolution of the experiments described below to approximately 10^6 founders.

Contraction of a barcoded population during colonization changes the frequency and number of barcodes relative to the inoculum. To determine when a *C. rodentium* infection is founded, C57BL/6J (B6) mice were orally gavaged with 4×10^8 CFUs. Remarkably, despite this relatively large dose, within 24 hours there was an average of only 9 founders (geometric mean), and as few as one founder per mouse (a single barcode; Figure 1B). Thus, only ~1 of every 4×10^7 cells in the inoculum establishes infection, revealing that host bottlenecks result in a massive constriction of the pathogen population. Beyond 24 hours, the founding population remained stable at ~10 founders. The diminutive *C. rodentium* founding population indicates that the vast majority of the inoculum does not survive to give rise to detectable offspring and is thus either killed by the host or passes through the intestine and is excreted in feces. Consistent with the latter possibility, 5 hours after inoculation there were 1×10^7 CFU *C. rodentium* per gram of feces with 9×10^4 founders, suggesting that at this early point a numerous and diverse population has already reached the colon and cecum but failed to become founders. Surprisingly, the contraction of the pathogen population continued beyond 5 hours, when the pathogen had already reached the principal sites of colonization. Despite the profound bottleneck to infection, the ~10 founders were capable of replication, and by 5 days post inoculation the *C. rodentium* burden in the feces was on average 9×10^8 CFU/gram. Together, these observations reveal that there is a severe bottleneck to infection with this natural, mouse enteric pathogen; however, even though the restrictive bottleneck leaves a founding population that is a miniscule fraction of the inoculum, the founders robustly replicate, creating a total pathogen burden that ultimately exceeds the inoculum (Figure 1B).

The size of the founding population increases with dose

We reasoned that determining how dose impacts the number of founders could provide insight into the mechanisms underlying the bottleneck (Abel *et al.*, 2015b; Gillman *et al.*, 2021). For example, one explanation proposed for the *C. rodentium* bottleneck is that it is created by finite niches or resources (e.g., sugar or amino acids) whose scarcity limits the size of the population (Kamada *et al.*, 2012; Caballero-Flores *et al.*, 2020). At doses where the pathogen saturates this limited resource the 'finite resource' hypothesis predicts that increasing dose will not increase the number of founders (schematized in Figure 2A). An alternate possibility is that the bottleneck eliminates potential founders through a mechanism such as acid killing in the stomach

125 (Tennant *et al.*, 2008; Woodward *et al.*, 2022), which is expected to result in a founding population that
126 increases with dose.
127
To characterize the bottleneck, we orally inoculated B6 mice with *C. rodentium* doses ranging 1,000-fold from
10^7 to 10^{10} CFUs. Doses $\geq 10^8$ CFUs led to infection, with the founding population decreasing for ~2 days
before reaching a steady value that persisted until the infection began clearing (Figure 2B). Lower doses of *C.*
rodentium resulted in fewer founders and a longer period to reach peak shedding, with a correspondingly
longer time from inoculation to pathogen elimination. As the delay in shedding at lower doses correlated with
the delay in clearance, all mice were infected for a similar number of days and had similar total fecal burdens,
regardless of dose (Supplemental Figure 2).
135
The founding population was small in number. Even at the maximum inoculum of 10^{10} CFUs relatively few
founders were detected (83, geometric mean; Figure 2B). While founders were never numerous, increasing the
size of the inoculum always increased the size of the founding population. The bottleneck eliminated a
proportion of the *C. rodentium* population, resulting in a founding population that scaled with dose (increasing
dose 100-fold also increased founders ~100-fold). These observations indicate that the number of founders is
likely not dictated by limited space or resources, contradicting the finite resource hypothesis (Figure 2A).
142
As an increase in dose resulted in a proportional increase in the number of founders, we represented their
relationship as a line by plotting $\log_{10}$-transformed dose and founding population data from 5 days post
inoculation (Figure 2C). This line indicates that the bottleneck is not fixed, but rather functions by eliminating a
fraction of potential founders, as schematized in Figure 2A ('elimination bottleneck'). Since our findings
conform to a simple fractional relationship between dose and founding population, we will use this relationship
to define the bottleneck: in B6 mice 1 of every ~10^8 inoculated *C. rodentium* establish a replicative niche.
149
The x-intercept of the log-linear relationship between dose and founders can be used to calculate the dose at
which we expect 1 founder. This dose corresponds to the ID_{50} - the dose that leads to infection of ~50% of
animals. Thus, for *C. rodentium* infection, the ID_{50}, a critical parameter describing a pathogen's infectivity, is a
property biologically defined by the infection bottleneck. For B6 mice, the x-intercept of this line is between
$10^{7.2}$ and $10^{7.9}$ CFU (95% confidence-intervals) and explains why infection did not result from an inoculum of
10^7 CFU (Fig 2B). Surprisingly, even though *C. rodentium* is a natural mouse pathogen, at least ~100-million
organisms are required to routinely establish infection.
157
158 <u>Stomach acid contributes a 10- to 100-fold bottleneck to *C. rodentium* colonization</u>
159
We next probed the contribution of stomach acid to the highly restrictive B6 enteric colonization bottleneck.
The acidity of the stomach is thought to be a potent barrier against ingested bacteria; human studies find that
taking stomach acid reducing drugs increases the risk of contracting multiple enteric pathogens (Bavishi and
DuPont, 2011). Notably, it has been observed that eliminating stomach acid decreases the minimum infectious
dose for *C. rodentium* and increases the size of the founding population (Tennant *et al.*, 2008; Woodward *et*
al., 2022). Further, acid is mechanistically consistent with the fractional relationship which we observe between
dose and founding population (Figure 2). To test the role of stomach acid in restricting *C. rodentium* enteric
colonization, we treated mice with the fast-acting, irreversible H2-antagonist Loxtidine (aka
Lavoltidine)(Maradey-Romero and Fass, 2014). 3-5 hours after Loxtidine treatment, the pH of the stomach
rose from 2.5 to 4.7 (Figure 3A). Importantly, a pH of 2.5 sterilized 10^{10} CFUs of *C. rodentium* in under 15
minutes, whereas pH 4.7 did not kill *C. rodentium* even after a 1 hour exposure (Figure 3B).
171
Loxtidine treatment prior to inoculating B6 mice resulted in infection at a lower dose, a higher pathogen burden
in the feces 1 day post inoculation, and more founders on day 5 (Figure 3C-D). The fractional relationship
between dose and founding population was also observed in the absence of stomach acid, but the line
depicting this relationship was shifted upward. Loxtidine treatment increased the number of *C. rodentium*
founders approximately 10-fold at every dose, reducing the ID_{50} computed from the founding population from
$10^{7.3}$ to $10^{5.4}$ CFUs. Thus, stomach acid significantly contributes to the bottleneck restricting *C. rodentium*
colonization. However, the magnitude of stomach acid's contribution is relatively small, between 10- and 100-
fold of the observed ~10^8-fold B6 bottleneck to *C. rodentium* colonization. In the absence of stomach acid, the

C. rodentium population constricts >10^6-fold prior to establishing a replicative niche, indicating that other
factors must more potently contribute to the bottleneck.
182
183 Constriction of the *C. rodentium* inoculum occurs distal to the stomach, at the sites of infection
184
To further define the *C. rodentium* population dynamics and host barriers that accompany establishment of
infection, we probed where and when the bottleneck occurs. Five days post-inoculation, the largest pathogen
burdens were detected in the cecum and distal colon, with less numerous populations in the small intestine (SI)
(Figure 4A), consistent with previous observations (Wiles *et al.*, 2004). Within individual mice (intra-mouse) the
cecum, colon, and feces contained related populations of *C. rodentium*, with approximately the same number
of founders and similar barcodes (Figure 4B-E). Importantly, the near identity of the barcodes found in the fecal
population to those in the cecum and colon indicates that fecal samples can be used to report on the pathogen
population at these primary infection sites, facilitating longitudinal monitoring. While intra-mouse populations
were related, comparisons of barcodes between cohoused mice (inter-mouse) inoculated with the same
inoculum revealed that each mouse contained a distinct *C. rodentium* population (Figure 4C-E). The distinct
identities of the founding populations in each of five cohoused, co-inoculated mice was apparent when
comparing pathogen barcode frequencies with principal component analysis (PCA), where intra-mouse
samples formed their own tight clusters (Figure 4C). Similarly, analysis of barcode genetic distances showed
that the intra-mouse pathogen populations were highly similar (low genetic distance), whereas they were
dissimilar to the populations in cohoused, co-inoculated mice (Figure 4D-E). A notable exception were the *C.*
rodentium populations from some SI samples that were more closely related to cage-mates than other intra-
mouse samples, likely reflecting recent inter-mouse exchange via coprophagy (Supplemental figure 3). These
data suggest that despite the consumption of *C. rodentium*-laden feces, *C. rodentium* infection leads to super-
colonization resistance at the primary infection sites in the cecum and colon, preventing transmission to
cohoused, co-infected mice.
205
To test this super-colonization resistance hypothesis, we separately infected two groups of 'seed' mice with
different sets (A and B) of barcoded *C. rodentium* (Supplemental figure 4A). At the peak of colonization in the
seed mice, 7 days post-inoculation, they were co-housed for 16 hours along with an uninfected 'contact'
mouse, three mice per cage. After 16 hours, the mice were separated back into 3 cages containing mice
originally inoculated with the A barcodes, inoculate with the B barcodes, or uninoculated. No transmission of
barcodes was detected between the animals originally inoculated with the A and B barcodes (Supplemental
figure 4B), confirming that *C. rodentium* infection prevents super-colonization. In marked contrast, the contact
mice became infected with founders from seed A and/or B, demonstrating the ready transmission of *C.*
rodentium from infected to uninfected mice. Furthermore, the co-infection of contact mice with barcodes from A
and B confirms a previous report from super-infection experiments in mice lacking a microbiota (Buschor *et al.*,
2017) that immunity to super-colonization takes time, providing a window for co-colonization.
217
Based on the high burdens of *C. rodentium* in the cecum but not the colon during the first 3 days following
inoculation, prior studies proposed that infection begins with pathogen expansion in the cecum, followed by
subsequent spread to the colon (Wiles *et al.*, 2004); a hypothesis that is consistent with the closely related
intra-mouse *C. rodentium* populations that we observe in the cecum and colon 5 days after inoculation (Figure
4A-E). To determine when and where *C. rodentium* initiates infection, we monitored the luminal and adherent
C. rodentium populations in the cecum and colon. Within the first 5 hours a large burden (>10^7 CFU) and
numerous founders (>10^5 N_r') were detected in both locations (Figure 4F-G). Since a large founding population
was observed in the cecum and colon early after inoculation, we can discount the model that the primary
bottleneck occurs proximal to these locations (e.g., stomach acid or bile). The number of founders and the total
burden contracted over the first 24 hours, resulting in small (<10^5 CFU) dissimilar (genetic distance >0.4)
populations in the cecum and colon one day post inoculation. Expansion was detected first in the cecum, on
day 2. Concomitant with cecal expansion, the populations in the cecum and colon became increasingly similar;
i.e., the genetic distance between the populations became smaller. The most plausible model to fit these data
(depicted in Figure 4H) is that (1) within hours many bacteria pass through the stomach, reaching the cecum
and colon, and then (2) these populations diverge as they separately constrict, and finally, (3) spread to both
locations when a small number of founders begin to replicate. We propose that the initial population expansion
begins in the cecum and then spreads to the colon, but we cannot rule out the opposite directionality because
we were unable to serially sample the internal populations from a single mouse. However, displacement of the

cecal population by bacteria from the colon seems unlikely because it would require non-flagellated *C.*
rodentium to move against the bulk flow of the gut and thus we favor the model that infection initiates in the
cecum.
239
240 <u>C3H/HeOuJ mice have a less restrictive bottleneck than C57BL/6J</u>
241
While *C. rodentium* causes self-limited diarrhea in B6 mice, infection leads to a lethal diarrheal disease in
C3H/HeOuJ (C3Ou) mice (Figure 5A-B)(Vallance *et al.*, 2003). We found that increased vulnerability to
disease correlated with a less restrictive bottleneck. *C. rodentium* is 10- to 100-fold more infectious in C3Ou
than B6 mice, infecting at a ~10-fold lower dose and producing ~10-times more founders at every dose (Figure
5C). While the bottleneck was relaxed in C3Ou mice, a fractional relationship remained between dose and
founding population, suggesting a similar underlying mechanism restricts colonization in both mouse
genotypes. Also, as in B6 animals, higher doses and more founders accelerated the dynamics of pathogen
shedding in C3Ou mice (Figure 5A). These observations demonstrate that in addition to dose, the size of the
founding population is determined by host genetics, which may impact the bottleneck through several
mechanisms.
252
253 <u>The bottleneck to *C. rodentium* enteric colonization is microbiota dependent</u>
254
As shown above, a large portion of the restrictive, fractional, B6 bottleneck to *C. rodentium* colonization occurs
distal to the stomach, at the chief sites of infection in the cecum and colon. These data strongly suggest that
the principal step limiting colonization occurs during the pathogen's establishment of a replicative niche in the
cecum and/or colon. One factor present at these sites and previously linked to limiting *C. rodentium*
colonization is the microbiota (Kamada *et al.*, 2012; Buschor *et al.*, 2017). We therefore tested whether acute
microbiota depletion eliminated the bottleneck to *C. rodentium* colonization. Treating mice with the antibiotic
streptomycin for the 3 days prior to inoculation with streptomycin-resistant *C. rodentium* greatly accelerated
pathogen population expansion, with mice shedding >10^9 CFUs per gram of feces within the first day (Figure
6A). Further, streptomycin pretreatment almost completely ablated the bottleneck, with colonization at doses
as low as ~100 CFUs; at this low dose, we measured an average of 25 founders 5 days post inoculation,
indicating that *C. rodentium* experiences less than a 10-fold bottleneck following microbiota depletion (Figure
6B). Since an ~10-fold bottleneck remains after microbiota depletion and an ~10-fold bottleneck is stomach
acid dependent (Figure 3D), these data suggest that the combination of the microbiota and stomach acid can
account for the majority of factors restricting *C. rodentium* colonization.
269
To confirm that streptomycin's ablation of the bottleneck to *C. rodentium* colonization occurs because of
microbiota depletion rather than an off-target effect, we also determined the bottleneck in B6 mice lacking a
microbiota (germ-free). In germ-free mice, like streptomycin pretreated animals, there was almost no
bottleneck to *C. rodentium* colonization (Figure 7A-B). Animals lacking a microbiota were colonized at a dose
of 150 CFU and shed numerous *C. rodentium* within 1 day of inoculation. Together, experiments with germ-
free and streptomycin-pretreated mice reveal that the primary barrier to enteric colonization is linked to the
microbiota.
277
Microbiota disruption also impaired the capacity of mice to clear *C. rodentium* infection (Figure 6A,
7A)(Kamada *et al.*, 2012). Pathogen burden in the feces of germ-free mice did not decrease over time, in
marked contrast to mice with an intact microbiota (specific pathogen free; SPF). Similarly, most cages of
streptomycin-pretreated mice failed to clear the pathogen, with heterogeneity presumably caused by variation
in the rebound of the microbiota after streptomycin treatment (Figure 6A). Despite high fecal burdens, germ-
free animals only exhibited mild diarrhea and did not lose weight for the 30 days of observation. These data
indicate that the microbiota is the primary impediment to *C. rodentium* replication in the gastrointestinal tract,
antagonizing the pathogen's capacity to initiate a replicative niche and promoting its clearance.
286
287 In germ-free and streptomycin pretreated animals the number of *C. rodentium* founders ceased to be
288 fractionally related to dose; doses ranging 10,000-fold, from 10^6 to 10^{10} CFUs, all yielded a similar number of
289 founders 5-days post-inoculation (Figure 6B, 7B). These data suggest that there is an upper limit to the size of
290 the *C. rodentium* founding population of ~10^5 on day 5 (i.e., a bottleneck caused by limited resources as
291 illustrated in Figure 2A). Furthermore, in the absence of a microbiota dependent bottleneck the maximum size

of the founding population continuously decreased for the 20 days of observation (Fig 6C, 7C). Although there was no contraction in the *C. rodentium* burden following infection of germ-free animals, the maximum number of founders decreased from ~10^5 on day 5 to ~10^2 on day 20 (Figure 7A, C). A decrease in diversity without a decrease in abundance suggests that *C. rodentium* adapts to the germ-free environment, introducing a new bottleneck caused by intra-pathogen competition.

To test the hypothesis that *C. rodentium* evolved during colonization of germ-free animals, we sequenced the genomes of single *C. rodentium* colonies isolated from infected SPF or germ-free mice 5 or 20 days post inoculation. 5 days post-inoculation, the *C. rodentium* genomes isolated from SPF and germ-free mice were similar to the inoculum, with 6/10 colonies lacking detectable variations (Supplemental figure 5). These data indicate that the initial contraction of the *C. rodentium* population observed during establishment of infection is not caused by selection of a genetically distinct subpopulation of the inoculum. By contrast, 20 days growth in the absence of a microbiota was always accompanied by changes in the *C. rodentium* genome. Notably, *C. rodentium* with structural variations in the LEE pathogenicity island became dominant in 4 out of 5 cages of infected germ-free animals (Figure 7D, Supplemental figure 5). These variations included large deletions of up to 97,691-bps (Figure 7E, isolate from mouse F1) that eliminated the entire island, which is essential for colonization of SPF mice (Schauer and Falkow, 1993). These genome alterations suggest that in the absence of a microbiota, a common mechanism for *C. rodentium* adaption to the host environment is to lose the LEE pathogenicity island. Thus, we conclude that competition among *C. rodentium* constricts the diversity of the population in the absence of a microbiota-dependent bottleneck, with organisms that lose the LEE virulence island outcompeting bacteria possessing the LEE. Additional mutations were also detected in the *C. rodentium* isolated on day 20 from germ-free animals, including in the galactonate operon, which have previously been observed in *Escherichia coli* colonizing microbiota depleted mice (Lescat *et al.*, 2017) (Supplemental figure 5). Thus, there may be common evolutionary strategies for pathogenic and non-pathogenic bacteria to adapt to growth without competition in the host intestine.

Collectively these experiments show that of the multiple host factors protecting against enteric infection, the microbiota is by far the most restrictive. Diminution of the microbiota markedly increases host susceptibility, permitting infection at almost any dose. In the absence of competition with the microbiota, a new slow-acting bottleneck constricted the *C. rodentium* population as the pathogen evolved increased fitness, notably through loss of the LEE pathogenicity island.

Discussion

Here, we investigated the host bottlenecks that protect against *C. rodentium* colonization. We found that the pathogen population undergoes a severe and slow-acting bottleneck. The size of the founding population scaled proportionally with dose: increasing the inoculum 100-fold increased the number of founders ~100-fold. Since increasing pathogen dose also increased the number of founders, we conclude that the bottleneck to colonization is not caused by the limited availability of finite resources or niches. We used the relationship between dose and founding population to interrogate the factors that create the bottleneck to *C. rodentium* colonization and discovered that the bottleneck occurs in at least two distinct sites: an initial stomach acid barrier kills 90-99% of the inoculum and the microbiota removes an additional 99.99999% after the pathogen reaches the cecum. Based on these results we propose a microbiota dependent factor creates the primary bottleneck by inhibiting *C. rodentium's* infectivity. Overall, our findings suggest that quantifying the impact of dose on founding population is a powerful framework for understanding the protection provided by infection bottlenecks.

While *C. rodentium* is capable of remarkable expansion after establishing a replicative niche, this pathogen has unexpectedly low infectivity in experimental conditions and requires a high dose to colonize. In contrast, some human adapted enteric pathogens are highly infectious, with Shigella and EHEC colonizing people after exposure to as few as 100 CFU (Porter *et al.*, 2013; Koutsoumanis *et al.*, 2020). We found that the high dose required for *C. rodentium* colonization is largely attributable to the microbiota. We can contextualize this observation by considering that the microbiota of wild mice, presumably the ancestral host of *C. rodentium*, varies between individuals and differs drastically from the microbiota observed in laboratory animals (Weldon *et al.*, 2015). Based on these observations, we speculate there could be two non-mutually exclusive reasons why *C. rodentium* was inefficient at overcoming the microbiota in our experiments: (1) this pathogen may utilize a generalist strategy, able to colonize in the context of diverse microbiotas, but not optimal to any, and/or (2) *C. rodentium* may have adapted to colonize mice whose microbiotas differ from the laboratory mice tested here. Importantly, the microbiota has been implicated in impeding enteric colonization by many human pathogens (Buffie and Pamer, 2013). In general, heterogeneity of the microbiota within host populations may explain why the microbiota has remained a potent barrier preventing colonization by multiple pathogens.

Our results suggest that the LEE pathogenicity island is a specific adaptation strategy for *C. rodentium* (and presumably other A/E pathogens) to overcome barriers to colonization created by pressures from the microbiota. LEE activity allows attaching effacing pathogens, such as *C. rodentium*, to create a replicative niche that is in close apposition to the epithelial surface and relatively free of microbiota (Collins *et al.*, 2014). While the LEE is required for colonization in the presence of the microbiota, we found that this island is detrimental in the microbiota's absence. During long term colonization of germ-free mice, bacteria with genomic deletions of the region containing the LEE took over the population (Figure 7). A similar phenomenon was previously observed in mice treated with dietary iron (Sanchez *et al.*, 2018), which increased intestinal glucose and presumably decreased competition with the microbiota. These examples of selection for *C. rodentium* containing genomic deletions of the LEE suggest that the LEE has a significant fitness cost. Kamada *et al.* described one cost of the LEE, proposing that LEE-encoded proteins are targets of the host's antibody response, leading to the elimination of LEE expressing bacteria (Kamada *et al.*, 2015). We propose that the LEE permits attaching and effacing pathogens to overcome microbiota-dependent pressures, but at the cost of increased stimulation of host immunity.

One way the gut microbiota is thought to inhibit intestinal colonization of invading bacteria is by competing for resources. For example, measurement of total pathogen burden has shown that *C. rodentium* colonization is limited by the availability of simple sugars and amino acids (Kamada *et al.*, 2012; Caballero-Flores *et al.*, 2020). However, it remains unclear whether limitation of resources contributes directly to the bottleneck or impacts downstream processes such as replication. Although we did not directly address the mechanisms of the microbiota dependent bottleneck, our discovery that the number of founders increases with pathogen dose strongly suggests that the microbiota does not create a bottleneck by limiting nutrients or niches. If the microbiota creates a bottleneck by depleting an essential resource, we would anticipate an upper limit to the size of the founding population (e. g., only enough iron or binding sites to support a limited number of founders). However, we found that the size of the founding population is extremely small and increased with dose; the number of founders increased from 2 to 83 when the dose increased 100-fold from 10^8 to 10^{10} CFU

379 (Figure 2). It seems unlikely that increasing the size of the inoculum by ~10^{10} CFU would also increase the
380 quantity of a limited resource to accommodate 81 additional founders. Thus, our findings suggest that the
381 primary mechanism by which the microbiota prevents infection is not through limitation of a finite resource.
382 Nonetheless, limited resources likely control other components of colonization, including pathogen expansion,
383 carrying capacity, and/or clearance.
384
Our dosing studies strongly suggest that the bottleneck to *C. rodentium* infection acts by eliminating potential
founders. Broadly, 'elimination' includes many mechanisms such as killing incoming bacteria or regulating their
ability to establish a replicative niche. Since constriction of the founding population is slow and *C. rodentium*
cells spend hours in proximity to their replicative niche without becoming founders, we favor a mechanism that
impedes infectivity, rather than direct pathogen killing. We speculate that one or more soluble, microbiota-
dependent metabolites are the primary factor that determines the size of the founding population. Indeed,
expression of the *C. rodentium* genes required for colonization is regulated by several soluble metabolites,
including bicarbonate, indole, and short-chain fatty acids (Hart *et al.*, 2008; Curtis *et al.*, 2014; Connolly *et al.*,
2018; Pifer *et al.*, 2018; Kumar and Sperandio, 2019; Osbelt *et al.*, 2020).
394
395 In conclusion, we found that multiple factors contribute to the dose at which infection occurs. We propose that
396 the mechanisms creating a bottleneck are reflected in and can largely be defined by the relationship between
397 dose and founding population. In the case of *C. rodentium*, the most restrictive infection bottleneck is created
398 by the microbiota, an observation that likely applies to many enteric pathogens.

399 **Figures**
400
401

Figure 1

402
Figure 1. A constrictive bottleneck impedes *C. rodentium* enteric colonization.
(**A**) Experimental strategy. To quantify the contraction (bottleneck) of the *C. rodentium* population during
colonization a barcoded library of bacteria was orally administered to mice. The number of unique cells from
the inoculum (founders) that expand to produce the observable population (burden) was determined from the
frequency and diversity of barcodes.
(**B**) 5 cohoused, adult, female, C57BL/6J mice were administered 4×10^8 CFU STAMP-CR253 by oral gavage
and the *C. rodentium* population was monitored in the feces for 5 days post inoculation (p. i.). Lines and error
are geometric means and standard deviation. Bacteria not detected (ND) counted as 0.5. Colony forming units
(CFU), grams (g). Validation of STAMP libraries in **Supplemental figure 1**.

Figure 2

412
Figure 2. The *C. rodentium* bottleneck is defined by a fractional relationship between dose and founders.

(**A**) Models for the relationship between dose and founding population. In the absence of a bottleneck all bacteria from the inoculum become founders. If the inoculum contracts due to the limited availability of finite nutrients or niches (e.g. iron, sugar, binding sites), the diversity of the population and thus the size of the founding population will remain fixed once those nutrients are saturated. If increasing dose increases the number of founders, then the underlying mechanism is not due to a limited resource; instead, the bottleneck acts proportionally on the inoculum by eliminating potential founders.

(**B-C**) C57BL/6J mice were inoculated with doses ranging from 10^7 to 10^{10} CFU of STAMP-CR253 and the *C. rodentium* population was monitored in the feces (geometric means and standard deviations; not detected counted as 0.5). Additional shedding analysis in **Supplemental figure 2**. (**C**) The bottleneck impeding B6 colonization is described 5 days post inoculation by comparing dose and founders with a linear regression of the log$_{10}$-transformed data (regression line with 95% confidence intervals; not detected counted as 0.8; x-intercept "ID$_{50}$" $10^{7.2}$-$10^{7.9}$ CFU). 4-8 animals per dose.

Figure 3

Figure 3. Stomach acid constricts the *C. rodentium* population by 10- to 100-fold.
(**A**) The effect of Loxtidine on stomach acid in C57BL/6J mice 3-5 hours after intraperitoneal administration of 1mg in 0.1ml PBS. pH determined post-mortem in aspirated stomach fluid. Boxes arithmetic mean (2.5 for mock and 4.7 for Loxtidine). Two-tailed t test with p-value 0.0002. n is 7 (mock) and 5 (Loxtidine).
(**B**) The acid tolerance of STAMP-CR69K in culture measured by diluting cells in LB at pH 2.5 or 4.7 and incubating at 37°C with shaking. pH 2.5 sterilized 10^{10} CFU in 15 minutes.
(**C-D**) 3-5 hours after intraperitoneal administration of PBS (vehicle) or Loxtidine (1mg), C57BL/6J mice were orally gavaged with doses ranging from 10^7 to 10^{10} CFU of STAMP-CR69K. (**C**) Bacterial burden monitored in

437 the feces for 5 days following inoculation (geometric means and standard deviations; not detected counted as
438 0.5). (**D**) Bottleneck to colonization measured 5 days post inoculation by comparing dose and founders with
439 linear regression of the $\log_{10}$-transformed data (regression line and 95% confidence intervals; significance
440 compares elevation with p-value <0.0001; not detected counted as 0.8). 4 animals per dose per group.

Figure 4

Figure 4. Infection is initiated by related populations of *C. rodentium* in the cecum and colon.

(**A-E**) *C. rodentium* populations in whole organ homogenates from 5 cohoused (intra-cage) C57BL/6J mice 5
days post inoculation with 4×10^8 CFU of STAMP-CR253. Within a mouse (intra-mouse) the *C. rodentium*
populations at the primary sites of colonization (cecum, proximal colon, distal colon, feces) share founders
(number, identity, and frequency of barcodes). (**A-B**) Lines connect intra-mouse samples. (**C**) Clustering of
barcode populations by principal component analysis (PCA). (**D-E**) Relatedness determined by comparing the
barcode frequencies by genetic distance (arithmetic means) with zero indicating no difference between
populations (identical). (**E**) Two-tailed t test with p-value <0.0001. Proximal (p), mid (m), distal (d), small
intestine (SI). Heatmap depicting genetic distance relationships of all intra-cage populations in **Supplemental**
figure 3.
(**F-G**) To determine when/where *C. rodentium* establishes a replicative niche, C57BL/6J mice were orally
gavaged with between 3×10^9 and 6×10^9 CFU STAMP-CR253. Following dissection, the cecum and colon were
flushed to separate organ adherent (**F**) and luminal (**G**) bacteria. Burden and founders display geometric
means and standard deviations. Bacteria not detected (ND) counted as 0.5. Relatedness of populations was
determined by comparing the barcode frequencies of colon and cecal populations from within the same animal
(intra-mouse) by genetic distance (arithmetic mean and standard deviation). 22 animals.
(**H**) Model depicting how related *C. rodentium* populations could initiate infection in both the cecum and colon:
(1) the inoculum minorly constricts passing through the stomach and SI to deposit diverse populations in the
cecum and colon, (2) the populations in the cecum and colon contract separately over the first 24-48 hours
becoming dissimilar, and then (3) expansion occurs in either the cecum or colon moving to both locations. We
depict the movement from cecum to colon as we judge this to be more likely, but the opposite is possible.
463

Figure 5

464
Figure 5. Host genotype impacts the bottleneck to *C. rodentium* colonization.
C3H/HeOuJ (**A**) or C57BL/6J (**B**) mice were inoculated with doses ranging from 10^6 to 10^{10} CFU of STAMP-
CR253. The *C. rodentium* population was monitored in the feces (geometric means and standard deviations;
not detected counted as 0.5) and animal health assessed by weight loss (percent compared to pre-inoculation;
arithmetic means and standard deviations) and body condition. For survival, lines are percent of initial animals
not moribund. (**C**) The bottleneck to C3Ou and B6 colonization is described 5 days post inoculation by
comparing dose and founders with a linear regression of the $\log_{10}$-transformed data (regression line with 95%
confidence intervals; significance compares elevation with p-value <0.0001; not detected counted as 0.8). B6
bottleneck data is repeated from **Figure 2**. 4 animals per dose.

Figure 6

Figure 6. Streptomycin treatment ablates most of the bottleneck preventing *C. rodentium* colonization.
The microbiota of conventional C57BL/6J mice was reduced by treatment with the antibiotic streptomycin for
the 3 days prior to inoculation with a streptomycin resistant library of STAMP-CR69K. (**A**) Founding population
and bacterial burden monitored in the feces, and (**B-C**) the bottleneck to colonization measured by comparing
dose and founders (geometric means and standard deviations; resolution limit is ~10^6 founders). (**A**) Animal
health monitored by weight loss (percent compared to pre-inoculation; arithmetic means and standard
deviations) and body condition. For survival, lines are percent of initial animals not moribund. Untreated B6
bottleneck data is repeated from **Figure 2** for comparison. 4 animals per dose.

Figure 7

A

Germ-free

Dose (CFU): 10^{10}, 10^8, 10^6, 10^4, 10^2

Founders (fecal N_s')

~resolution (10^6)

Founders (N_s')

Burden (CFU/g feces)

Weight (% initial)

days p.i.

100% survival: 10^{10}, 10^8, 10^6, 10^4, 10^2 (all doses)

B

Bottleneck (fecal 5 days p. i.)

~res.

No bottleneck (theoretical)

Founders (N_r)

Germ-free

~10^8-fold

SPF

Dose (CFU)

C

Germ-free bottleneck (fecal)

~res.

1 day

2 days

5 days

10 days

Founders (N_r')

20 days

Dose (CFU)

D

LEE variants identified by *C. rodentium* genomic sequencing

SPF 5 days p.i.	Germ-free 5 days p.i.	Germ-free 20 days p.i.
A 1 wt co-housed 2 wt 3 wt 4 wt	C 1 wt 2 wt 3 wt 4 wt	C 1 del 2 del 3 del 4 del
B 1 wt co-ho. 2 wt	7 cages (A-G)	wt - D1, del - E1, del - F1, ins - G1

E

3,100,000

annotation

LEE pathogenicity island

3,200,000

inoculum

germ-free 20-days p. i.

C1   80,204bp-deletion

D1

E1   97,469bp-deletion

F1   97,691bp-deletion

G1   insertion

total mapped reads overlaid with non-specific mapped reads

483
484 **Figure 7. The bottleneck to *C. rodentium* colonization is microbiota dependent.**
485 Germ-free C57BL/6J mice were orally inoculated with doses ranging from 10^2 to 10^{10} CFU of STAMP-CR69K.
486 4 animals per dose, cohoused with animals receiving the same dose in sterile cages.

(**A**) Bacterial burden and founding population monitored in the feces (geometric means and standard deviations; resolution limit is ~10^6 founders). Animal health monitored by weight loss (percent compared to pre-inoculation; arithmetic means and standard deviations) and body condition. No animals became moribund.

(**B-C**) Bottleneck to colonization measured by comparing dose and founders (geometric means and standard deviations). SPF B6 bottleneck data is repeated from **Figure 2** for comparison.

(**D**) To determine if colonization was accompanied by changes in the *C. rodentium* genome, whole genome sequencing was performed on 3 clones (colonies) from the STAMP-CR69K input library and compared to clones isolated from feces of infected mice (1 colony per mouse). Boxes represent the genome status of the LEE pathogenicity island. No LEE genomic changes (wild-type; wt), deletion of the entire LEE region (del), insertion within the LEE (ins). Mice with a conventional microbiota (specific pathogen free; SPF). Other genomic changes listed in **Supplemental figure 5**.

(**E**) Depiction of a ~100,000 base-pair region of the *C. rodentium* genome containing the LEE pathogenicity island. Read depth from STAMP-CR69K inoculum and 5 clones isolated after 20 days passage in otherwise germ-free animals. Large deletions in 3 of 5 clones revealed by lack of specifically mapped reads in regions of up to 97,691 base-pairs. Non-specific reads map to multiple loci in the genome (primarily transposons).

Supplement 1

502
Supplemental figure 1. Validation of *Citrobacter rodentium* STAMP libraries.
(**A**) Growth of STAMP-CR253 library compared to the parental, untagged strain. Growth curve performed in 96
well plate reader at 37°C, shaking, in LB. 18 clones (colonies) from STAMP library and 2 from parent. Lines are
means of triplicate samples.
(**B**) The ability to determine the size of the founding population from STAMP libraries was validated in culture
by comparing the number of plated colonies (CFU; known number of founders) to the size of the founding
population following processing of those plated populations (calculated founders). Without calibration, STAMP
accurately determined the number of founders up to ~10^4 CFU and plateaued at ~10^6 CFU. Therefore, these
values can be used to calibrate future experiments to determine the size of founding populations up to 10^6,
with increased uncertainty above 10^4. STAMP-CR253 (253 unique barcodes) founders (N_r) measured for 4
cultures across 7 doses. STAMP-CR69K (~69,000 unique barcodes) founders (N_s) measured for 3 cultures
across 10 doses.

Supplement 2

516
517 **Supplemental figure 2. Additional data related to shedding of C57BL/6J mice in Figure 2.**
518 In C57BL/6J mice, dose impacts the lag between inoculation and peak shedding (**A**) and the lag between
519 inoculation and clearance (**D**) but does not change the days of peaks shedding (**B**) or the total fecal burden
520 (**C**). n is 4 animals per dose. (**A, B, D**) Arithmetic mean, (**C**) geometric mean.

Supplement 3

521
Supplemental figure 3. All intra-cage relationships from Figure 4A-E.
Comparison by genetic distance of *C. rodentium* populations 5 days post inoculation in 5 cohoused (intra-cage)
C57BL/6J mice inoculated with STAMP-CR253. Zero indicates no difference between populations (identical).
From proximal (p) to distal (d): p small intestine (SI), mid (m) SI, dSI, cecum, pcolon, dcolon, feces.

Supplement 4

Supplemental figure 4. Super-colonization resistance prevents cohoused mice from sharing *C. rodentium* populations

To test transmission of *C. rodentium* to already infected mice, two cages of 4 C57BL/6J mice (seeds A and B) were separately inoculated with mixed pools of barcoded strains (barcodes A and B). Barcodes A and B are each a group of 3 separate strains with non-redundant barcodes. Seed animals A and B were separated until peak colonization. On the 6th day after seed inoculation, the mice were re-arranged to co-house 1 mouse each of seed A, seed B, and uninfected contact C overnight for 16 hours. The next morning, they were separated, housing the A seeds together, the B seeds together, and the C contacts together. 12 days post seed inoculation (5 days post cohousing) the fecal *C. rodentium* populations were analyzed to determine the contribution from pools A and B. <1% of reads from seed A mice were contributed by the B barcodes and vice versa.

Supplement 5

mouse	days p. i.	cage	mouse	variation	genomic change	annotation	coding change	amino acid change
SPF C57BL/6J	5 days	A	A1	complex	g.3081146..3081467	intergenic		
			A2	none				
			A3	SNV	g.388509C>G	lamB	c.766G>C	p.Val256Leu
			A4	none				
		B	B1	none				
			B2	none				
germ-free C57BL/6J		C	C1	SNV	g.527543C>T	panE	c.232G>A	p.Asp78Asn
			C2	none				
			C3	SNV	g.330872T>C	ROD_03001	c.557A>G	p.Glu186Gly
				SNV	g.4274290A>G	dgoT	c.442A>G	p.Arg148Gly
			C4	none				
	20 days	C	C1	deletion	g.3100349…3180553	multiple (including LEE)		
				SNV	g.4029101C>T	glpK	c.1000C>T	p.Pro334Ser
				SNV	g.5295383C>T	intergenic		
			C2	deletion	g.3100349…3180553	multiple (including LEE)		
				SNV	g.4029101C>T	glpK	c.1000C>T	p.Pro334Ser
				SNV	g.5295383C>T	intergenic		
			C3	deletion	g.3100349…3180553	multiple (including LEE)		
				SNV	g.4029101C>T	glpK	c.1000C>T	p.Pro334Ser
				SNV	g.5295383C>T	intergenic		
			C4	deletion	g.3100349…3180553	multiple (including LEE)		
				SNV	g.4029101C>T	glpK	c.1000C>T	p.Pro334Ser
				SNV	g.5295383C>T	intergenic		
		D	D1	SNV	g.4270455T>A	dgoR	c.143T>A	p.Val48Glu
				SNV	g.4274290A>G	dgoT	c.442A>G	p.Arg148Gly
				SNV	g.4634927C>T	gntU	c.895C>T	non-coding
				insetion	g.5407322^5407357	intergenic		
		E	E1	deletion	g.3100354…3197823	multiple (including LEE)		
				insertion	g.4989831^4989834	gatZ		
		F	F1	deletion	g.3100357…3198048	multiple (including LEE)		
				SNV	g.3937003C>T	purH	c.837C>T	non-coding
				SNV	g.4274290A>G	dgoT	c.442A>G	p.Arg148Gly
		G	G1	deletion	g.2391346…2391356	galS		
				complex	g.3174563...3174662	grlR		

Supplemental figure 5. Changes in the *C. rodentium* genome following mouse expansion.
Whole genome sequencing was used to detect variations in the *C. rodentium* genome after expansion and
selection in mice. 3 clones (colonies) from the STAMP-CR69K library were compared to isolates from the feces
of infected mice (1 colony per mouse). No variants ("none") detected in 4/6 isolates from SPF (specific
pathogen free) animals 5 days p. i., 2/4 isolates from germ-free animals 5 days p. i., and 0/8 isolates from
germ-free animals 20 days p.i. 4 cohoused germ-free animals 20 days p. i. (C1-4) had identical (clonal)
isolates. These data were filtered to only include high confidence variants (present in >90% of reads).

Resource Availability

Lead contact
Requests for further information, resources, or reagents will be fulfilled by the Lead Contact, Matthew K. Waldor (mwaldor@research.bwh.harvard.edu).

Data and code availability
STAMP scripts and barcode read counts will be deposited in an online database prior to final publication at https://github.com/hullahalli/stampr_rtisan. Sequencing reads from STAMP experiments and bacterial whole genome sequencing will be made available in the Sequencing Read Archive.

Experimental Model and Subject Details

Mice
SPF C57BL/6J and C3H/HeOuJ mice were purchased from Jackson Laboratory. Mice were acclimated for at least 72 hours prior to experimentation. Germ-free C57BL/6J mice were provided by the Massachusetts Host-Microbiome Center and following inoculation were maintained with autoclave-sterilized cages, food, and water. Female mice 9-14 weeks of age at the start of experimentation were used. Animal studies were conducted in a biosafety level 2 (BSL2) facility at the Brigham and Women's Hospital. All experiments involving mice were performed according to protocols reviewed and approved by the Brigham and Women's Hospital Institutional Animal Care and Use Committee (protocol 2016N000416) and in compliance with the Guide for the Care and Use of Laboratory Animals.

Bacterial Strains
A spontaneous streptomycin-resistant (SmR) mutant of *C. rodentium* strain ICC168, previously known as *Citrobacter freundii* biotype 4280 (ATCC 51459) was used to create the barcoded strains used throughout this manuscript. The growth of this SmR mutant in lysogeny broth (LB) was indistinguishable from the wild-type strain from which it was derived by growth curves and competition in liquid media. Whole genome sequencing indicates that this SmR mutant contains 2 variations from the published (Petty *et al.*, 2010) ICC168 genome: (1) a non-coding single nucleotide variation in *tolB*, and (2) a coding single nucleotide variation in *rpsL* (which presumably confers SmR). Unless noted, *C. rodentium* was grown at 37°C in LB broth with shaking or on solid LB agar. As necessary, media was supplemented with streptomycin (200 µg/ml) and/or kanamycin (50 µg/ml).

Method Details

Generation of STAMP libraries
The library "STAMP-CR253" was created by integrating a barcoded variant of the plasmid pDS132 (Philippe *et al.*, 2004) into the *C. rodentium* genome in the intergenic region between ROD_05521 and *selU*. To create the donor plasmid, we first inserted an ~800 base-pair fragment of *C. rodentium* genomic intergenic sequence into plasmid pKD4 (addgene Plasmid #45605)(Datsenko and Wanner, 2000) using New England Biolabs (NEB) HiFi DNA Master Mix. A fragment containing both the *C. rodentium* intergenic region and a kanamycin-resistance cassette was then amplified from the modified version of pKD4 using primers containing 20 random nucleotides (Integrated DNA Technologies). This fragment was cloned into pDS132 and a library of ~2,000 assembled plasmids were transformed into the donor strain MFDλpir, creating a barcoded donor plasmid library capable of integrating into the intergenic region of the *C. rodentium* genome. The donor library was conjugated to SmR *C. rodentium*, with selection for streptomycin and kanamycin resistant colonies. A tiled library of transconjugants were pooled in phosphate buffed saline plus glycerol (PBSG) and frozen at -80°C in aliquots to create STAMP-CR253. Sequencing of the pooled library revealed the presence of 253 unique ~20 nucleotide barcodes.

The library "STAMP-CR69K" was created using the plasmid donor library pSM1 (Fakoya *et al.*, 2022). The pSM1 donor library is composed of ~70,000 unique plasmids transformed into the donor strain MFDλpir. Each pSM1 plasmid carries a site specific Tn7 transposon containing a random ~25 nucleotide barcode adjacent to a kanamycin resistance cassette. The Tn7 transposon system has previously been characterized to integrate at a neutral site in the genome downstream of the gene *glmS* (Choi *et al.*, 2005). Conjugation was used to introduce the pSM1 library into SmR *C. rodentium* and transconjugants containing the transposon were

603 selected using streptomycin and kanamycin. Transconjugant colonies were pooled in PBSG and frozen at -
604 80°C in aliquots to create the library STAMP-CR69K. Sequencing indicated that the library contains ~69,000
605 unique barcodes.
606
<u>STAMP sample processing</u>
As previously described (Hullahalli and Waldor, 2021; Hullahalli *et al.*, 2021), bacteria from cultures or animal
samples were plated on LB agar containing streptomycin and/or kanamycin. Colonies were resuspended in
PBSG and stored at -80°C. Genomic templates were created by diluting samples in water and boiling at 95°C
for 15 minutes. Barcodes were amplified from the genomic templates using either Phusion DNA polymerase or
OneTaq HS Quick-Load (New England Biolabs) and amplified by PCR for 25 cycles. The presence of
amplicons was verified by agarose gel electrophoresis. Samples were pooled, purified using the GeneJet PCR
purification kit (Fisher), and sequenced with an Illumina MiSeq. Sequencing reads were subsequently
processed using CLC Genomics Workbench (Qiagen) and Geneious (Biomatters) to determine the number of
reads from each sample that map to the barcodes present in the respective libraries. The size of the founding
population was determined by comparing the frequency of barcodes to the inoculum. STAMP-CR253 (N_r) was
analyzed using the STAMP analysis pipeline (Abel *et al.*, 2015a). To accommodate an increased number of
barcodes, STAMP-CR69K (N_s) was analyzed using the STAMPR analysis pipeline (Hullahalli *et al.*, 2021;
Fakoya *et al.*, 2022). Standard curves were used to refine the measurements of founding population (N_r' and
N_s'). Cavalli-Sforza chord distance was used to compare the genetic distance of barcoded samples (Abel *et al.*,
2015a).
623
C. rodentium <u>whole genome sequencing</u>
Genomic DNA was extracted from *C. rodentium* colonies using the GeneJet genomic DNA purification kit
(Thermo Fisher). Genomic sequencing was performed by the Microbial Genome Sequencing Center LLC
(MiGS; Pittsburgh PA) using an Illumina NextSeq 2000. CLC Genomics Workbench (Qiagen) was used to map
sequencing reads to the *C. rodentium* ICC168 reference genome (NCBI ASM2708v1, RefSeq sequence: Chr.
NC_013716.1, pCROD1 NC_013717.1, pCROD2 NC_013718.1, pCROD3 NC_013719.1) and basic variant
detection and structural variant detection functions were used to compare between clones. We manually sorted
detected variants and only report high confident variants, defined as being present in >90% of all reads.
632
633 <u>Mouse infection</u>
634 Unless otherwise noted, libraries of barcoded bacteria were prepared for oral challenge by resuspending
635 frozen aliquots in liquid LB broth and expanding at 37°C with shaking for 3-5 hours. Following growth bacteria
636 were pelleted, resuspended in phosphate buffered saline (PBS), diluted to the desired concentration, and
637 stored at room-temperature until inoculation.
638
639 Mice were deprived of food for 3-4 hours prior to inoculation. For intra-gastric gavage mice were lightly sedated
640 with isoflurane inhalation; 100 µL of bacteria were inoculated into the stomach using a 1 mL syringe and a
641 sterile, 18G, 1.5 inch, 2 mm ball, feeding needle (Cadence Science). Following inoculation, dose was
642 determined by serial dilution and plating. Mice weight, body condition, and fecal appearance were monitored
643 during infection.
644
645 *C. rodentium* populations were isolated from fecal pellets and/or organs. Organ "Adherent" and "Content"
646 populations from the cecum and colon were separated by washing the respective organs with PBS. Samples
647 were homogenized in PBS using a bead beater (BioSpec Products, Inc) and 2 stainless-steel 3.2 mm beads.
648 To determine the total burden and size of the founding population *C. rodentium* were expanded from
649 homogenized animal samples on solid media containing streptomycin and/or kanamycin.
650
<u>Loxtidine treatment</u>
Loxtidine was generously provided by James R. Goldenring, M.D., Ph.D. (Vanderbilt University School of
Medicine). Loxtidine was dissolved in PBS and 1 mg was administered intraperitoneal 3-5 hours prior to
inoculation with *C. rodentium*. The pH of stomach content was determined by aspirating liquid from the
stomach immediately post-mortem and measuring the pH using an Orion Micro pH Electrode (Thermo
Scientific).
657
658 <u>Germ-free infection</u>

659 Germ-free mice were provided by the Massachusetts Host-Microbiome Center in autoclave-sterilized cages
660 and immediately inoculated with *C. rodentium*. Mice were maintained and handled using sterile equipment.
661
662 <u>Streptomycin-treatment</u>
663 Streptomycin-treated mice were given filter-sterilized water containing 5 mg/mL streptomycin sulfate (Teknova)
664 for the 3 days prior to inoculation. During streptomycin treatment feces were monitored on LB agar with
665 streptomycin for streptomycin-resistant microbes, which did not appear. Mice were switched back to regular
666 water at the same time as they were inoculated with *C. rodentium*.
667
668 **Quantification and Statistical Analysis**
669
670 Statistical analyses were performed using GraphPad Prism version 9. Information regarding the number of
671 samples and statistical tests are described in the figure legends. Geometric means, geometric standard
672 deviations, and non-parametric tests were used when quantifying bacterial populations. As the bottom of our
673 detection limit was near 1 we substituted a value between 0.1 and 0.8 when no bacteria were detected.
674
675 **Acknowledgments**
676

We acknowledge members of the Waldor lab for their discussions and input. We thank Dr. James R.
Goldenring for providing Loxtidine and Ms. Tiffany S. Davanzo (Slaybaugh Studios) for assisting with
illustrations. Gnotobiotic mouse studies were supported by the Massachusetts Host-Microbiome Center, with
funding from the NIH (P30DK034854), and the Massachusetts Life Sciences Center (MLSC). This work was
supported by NIH grants to J.R.T. (R01DK61931, R01DK68271) and M.K.W (R01AI042347), a US Department
of Defense CDMRP contract to J.R.T. (PR181271), the Howard Hughes Medical Institute (HHMI) to M.K.W,
and fellowships from the NIH to K.H. (F31AI156949) and I.W.C (T32DK007477-37). This article is subject to
HHMI's Open Access to Publications policy. HHMI lab heads have previously granted a nonexclusive CC BY
4.0 license to the public and a sublicensable license to HHMI in their research articles. Pursuant to those
licenses, the author-accepted manuscript of this article can be made freely available under a CC BY 4.0
license immediately upon publication.
688

References:

Abel, S, Abel Zur Wiesch, P, Chang, HH, Davis, BM, Lipsitch, M, and Waldor, MK (2015a). Sequence tag-based analysis of microbial population dynamics. Nat Methods 12, 223–226.

Abel, S, Abel zur Wiesch, P, Davis, BM, and Waldor, MK (2015b). Analysis of Bottlenecks in Experimental Models of Infection. PLOS Pathog 11, e1004823.

Bavishi, C, and DuPont, HL (2011). Systematic review: the use of proton pump inhibitors and increased susceptibility to enteric infection. Aliment Pharmacol Ther 34, 1269–1281.

Buffie, CG, and Pamer, EG (2013). Microbiota-mediated colonization resistance against intestinal pathogens. Nat Rev Immunol 2013 1311 13, 790–801.

Buschor, S, Cuenca, M, Uster, SS, Schären, OP, Balmer, ML, Terrazos, MA, Schürch, CM, and Hapfelmeier, S (2017). Innate immunity restricts Citrobacter rodentium A/E pathogenesis initiation to an early window of opportunity. PLOS Pathog 13, e1006476.

Caballero-Flores, G, Pickard, JM, Fukuda, S, Inohara, N, and Núñez, G (2020). An Enteric Pathogen Subverts Colonization Resistance by Evading Competition for Amino Acids in the Gut. Cell Host Microbe 28, 526-533.e5.

Choi, KH, Gaynor, JB, White, KG, Lopez, C, Bosio, CM, Karkhoff-Schweizer, RAR, and Schweizer, HP (2005). A Tn7-based broad-range bacterial cloning and expression system. Nat Methods 2005 26 2, 443–448.

Collins, JW, Keeney, KM, Crepin, VF, Rathinam, VAK, Fitzgerald, KA, Finlay, BB, and Frankel, G (2014). Citrobacter rodentium: infection, inflammation and the microbiota. Nat Rev Microbiol 2014 129 12, 612–623.

Connolly, JPR et al. (2018). Host-associated niche metabolism controls enteric infection through fine-tuning the regulation of type 3 secretion. Nat Commun 9.

Curtis, MM, Hu, Z, Klimko, C, Narayanan, S, Deberardinis, R, and Sperandio, V (2014). The Gut Commensal Bacteroides thetaiotaomicron Exacerbates Enteric Infection through Modification of the Metabolic Landscape. Cell Host Microbe 16, 759–769.

Datsenko, KA, and Wanner, BL (2000). One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A 97, 6640–6645.

Fakoya, B, Hullahalli, K, Rubin, DHF, Leitner, DR, Chilengi, R, Sack, DA, and Waldor, MK (2022). Nontoxigenic Vibrio cholerae Challenge Strains for Evaluating Vaccine Efficacy and Inferring Mechanisms of Protection. MBio 13.

Gillman, AN, Mahmutovic, A, Abel zur Wiesch, P, and Abel, S (2021). The Infectious Dose Shapes Vibrio cholerae Within-Host Dynamics. MSystems 6.

Hart, E, Yang, J, Tauschek, M, Kelly, M, Wakefield, MJ, Frankel, G, Hartland, EL, and Robins-Browne, RM (2008). RegA, an AraC-like protein, is a global transcriptional regulator that controls virulence gene expression in Citrobacter rodentium. Infect Immun 76, 5247–5256.

Hullahalli, K, Pritchard, JR, and Waldor, MK (2021). Refined Quantification of Infection Bottlenecks and Pathogen Dissemination with STAMPR. MSystems 6.

Hullahalli, K, and Waldor, MK (2021). Pathogen clonal expansion underlies multiorgan dissemination and organ-specific outcomes during murine systemic infection. Elife 10.

Kamada, N, Kim, YG, Sham, HP, Vallance, BA, Puente, JL, Martens, EC, and Núñez, G (2012). Regulated

virulence controls the ability of a pathogen to compete with the gut microbiota. Science (80-) 336, 1325–1329.

Kamada, N, Sakamoto, K, Seo, SU, Zeng, MY, Kim, YG, Cascalho, M, Vallance, BA, Puente, JL, and Núñez,
G (2015). Humoral immunity in the gut selectively targets phenotypically virulent attaching-and-effacing
bacteria for intraluminal elimination. Cell Host Microbe 17, 617–627.

Koutsoumanis, K et al. (2020). Pathogenicity assessment of Shiga toxin-producing Escherichia coli (STEC)
and the public health risk posed by contamination of food with STEC. EFSA J 18, e05967.

Kumar, A, and Sperandio, V (2019). Indole signaling at the host-microbiota-pathogen interface. MBio 10.

Lescat, M, Launay, A, Ghalayini, M, Magnan, M, Glodt, J, Pintard, C, Dion, S, Denamur, E, and Tenaillon, O
(2017). Using long-term experimental evolution to uncover the patterns and determinants of molecular
evolution of an Escherichia coli natural isolate in the streptomycin-treated mouse gut. Mol Ecol 26, 1802–1817.

Maradey-Romero, C, and Fass, R (2014). New and Future Drug Development for Gastroesophageal Reflux
Disease. J Neurogastroenterol Motil 20, 6–16.

Mundy, R, MacDonald, TT, Dougan, G, Frankel, G, and Wiles, S (2005). Citrobacter rodentium of mice and
man. Cell Microbiol 7, 1697–1706.

Osbelt, L et al. (2020). Variations in microbiota composition of laboratory mice influence Citrobacter rodentium
infection via variable short-chain fatty acid production. PLOS Pathog 16, e1008448.

Petty, NK et al. (2010). The Citrobacter rodentium genome sequence reveals convergent evolution with human
pathogenic Escherichia coli. J Bacteriol 192, 525–538.

Philippe, N, Alcaraz, JP, Coursange, E, Geiselmann, J, and Schneider, D (2004). Improvement of pCVD442, a
suicide plasmid for gene allele exchange in bacteria. Plasmid 51, 246–255.

Pifer, R, Russell, RM, Kumar, A, Curtis, MM, and Sperandio, V (2018). Redox, amino acid, and fatty acid
metabolism intersect with bacterial virulence in the gut. Proc Natl Acad Sci U S A 115, E10712–E10719.

Porter, CK, Thura, N, Ranallo, RT, and Riddle, MS (2013). The Shigella human challenge model. Epidemiol
Infect 141, 223–232.

Sanchez, KK, Chen, GY, Schieber, AMP, Redford, SE, Shokhirev, MN, Leblanc, M, Lee, YM, and Ayres, JS
(2018). Cooperative Metabolic Adaptations in the Host Can Favor Asymptomatic Infection and Select for
Attenuated Virulence in an Enteric Pathogen. Cell 175, 146-158.e15.

Schauer, DB, and Falkow, S (1993). The eae gene of Citrobacter freundii biotype 4280 is necessary for
colonization in transmissible murine colonic hyperplasia. Infect Immun 61, 4654–4661.

Tennant, SM, Hartland, EL, Phumoonna, T, Lyras, D, Rood, JI, Robins-Browne, RM, and Van Driel, IR (2008).
Influence of gastric acid on susceptibility to infection with ingested bacterial pathogens. Infect Immun 76, 639–
645.

Troeger, C et al. (2018). Estimates of the global, regional, and national morbidity, mortality, and aetiologies of
diarrhoea in 195 countries: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Infect
Dis 18, 1211–1228.

Vallance, BA, Deng, W, Jacobson, K, and Finlay, BB (2003). Host susceptibility to the attaching and effacing
bacterial pathogen Citrobacter rodentium. Infect Immun 71, 3443–3453.

Weldon, L, Abolins, S, Lenzi, L, Bourne, C, Riley, EM, and Viney, M (2015). The Gut Microbiota of Wild Mice.
PLoS One 10, e0134643.

Wiles, S, Clare, S, Harker, J, Huett, A, Young, D, Dougan, G, and Frankel, G (2004). Organ specificity,
colonization and clearance dynamics in vivo following oral challenges with the murine pathogen Citrobacter
rodentium. Cell Microbiol 6, 963–972.

Wiles, S, Dougan, G, and Frankel, G (2005). Emergence of a "hyperinfectious" bacterial state after passage of
Citrobacter rodentium through the host gastrointestinal tract. Cell Microbiol 7, 1163–1172.

Woodward, SE et al. (2022). Gastric acid and escape to systemic circulation represent major bottlenecks to
host infection by Citrobacter rodentium. ISME J 2022, 1–11.