**Comparative genomics and phylogenomics of *Campylobacter* unveil**

**potential novel species and provide insights into niche segregation**

Sarah Henaut-Jacobs[1], Hemanoel Passarelli-Araujo[1,2,*], Thiago M. Venancio[1,*]

[1]Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e

Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos

Goytacazes, RJ, Brazil.

[2]Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade

Federal de Minas Gerais, Belo Horizonte, MG, Brazil.

*Corresponding authors

Short running title: Genomic diversity of *Campylobacter* spp.

Av. Alberto Lamego 2000, P5 sala 217; Parque Califórnia

Campos dos Goytacazes, RJ, Brazil

CEP: 28013-602

HPA: hemanuel.passarelli@gmail.com, thiago.venancio@gmail.com.

**Abstract**
*Campylobacter* is a bacterial genus associated with community outbreaks and gastrointestinal
symptoms. Studies on *Campylobacter* generally focus on specific pathogenic species such as
*C. coli* and *C. jejuni*. Currently, there are thousands of publicly available *Campylobacter*
genomes, allowing a more complete assessment of the genus diversity. In this work, we report
a network-based analysis of all available *Campylobacter* genomes to explore the genus
structure and diversity, revealing potentially new species and elucidating genus features. We
also hypothesize that the previously established Clade III of *C. coli* is in fact a novel species
(referred here as *Campylobacter spp12*). Finally, we found a negative correlation between
pangenome fluidity and saturation coefficient, with potential implications to the lifestyles of
distinct *Campylobacter* species. Since pangenome analysis depends on the number of
available genomes, this correlation could help estimate pangenome metrics of
*Campylobacter* species with less sequenced genomes, helping understand their lifestyle and
niche adaptation. Together, our results indicate that the *Campylobacter* genus should be re-
evaluated, with particular attention to the interplay between genome structure and niche
segregation.
35
36  **Keywords:** Network analysis; *Campylobacter* diversity; Pangenome.
37
38
39

## INTRODUCTION

*Campylobacter* is a Gram-negative Epsilonproteobacteria genus of motile bacteria that can infect humans and other animals. *Campylobacter* species are the leading cause of bacterial foodborne infections, especially transmitted via the consumption of undercooked chicken meat, contaminated water, or dairy products[1]. The genus comprises 38 validly published and non-synonymous species and 7 subspecies according to LPSN (List of Prokaryotic names with Standing in Nomenclature)[2] (on March 2023).

Among all known *Campylobacter* species, *C. coli* and *C. jejuni* are responsible for over 90% of human campylobacteriosis[1]. The ease of adaptation of these two species to multiple hosts led the World Health Organization to define *Campylobacter* as a pathogen of critical priority for the study and development of new antibiotic treatments[3]. Because of their clinical relevance, *C. coli* and *C. jejuni* have the largest number of publicly available genomes among *Campylobacter* species, creating an unintended bias for genus-wide population structure and ecological studies.

*Campylobacter* species occupy a wide range of niches[4]. For example, *C. coli* is differentiated into three clades (I, II, and III)[5]. While isolates from Clade I are usually associated with acute diarrhea in humans, those belonging to clades II and III are mainly found in environmental sources (e.g., lakes)[5,6]. In contrast, species such as *C. jejuni* present a more clonal population structure with host-specific clonal complexes (e.g., CC-42 and CC-61), which are associated with cattle and sheep[7]. Importantly, these niche preferences can be related to genome sequences, with bacteria with similar lifestyles sharing a greater number of genes.

While 16S rRNA sequencing has been widely used to study microbial communities, it is often unable to discriminate closely related genomes and can exhibit bias towards certain species[8]. The development of second and third generation sequencing technologies fueled the use of whole-genome based methods, such as Average Nucleotide Identity (ANI) and genome distance estimation[9]. Here, we employ a network-based approach using genome-wide comparative metrics to explore *Campylobacter* diversity and taxonomic classification.

## Results

### Dataset collection and type strains

We retrieved 3,319 *Campylobacter* genomes from the RefSeq database[10] in August 2020. After removing genomes with completeness below 90% and duplication levels (a proxy for contamination) greater than 5%, we created a dataset with 3261 *Campylobacter* genomes (Table S1). According to the NCBI classification, 46.24% and 29.9% of the genomes belong to *C. jejuni* and *C. coli*, respectively (Table S1), highlighting an expected overrepresentation of these species. These results also uncovered possible misclassification problems, since overrepresented species might overshadow less studied ones. The LPSN comprises 45 validly named and published *Campylobacter* representative genomes (Table S2), encompassing 38 species and 7 subspecies. We conducted an ANI analysis with all type strain genomes and confirmed that all pairs of type strains from different species had ANI values lower than 95%

81  (the threshold to define bacterial species[11]). We also identified three subspecies with ANI
82  lower than 95% with their respective species' type strains (*C. fetus* subsp*. testudinum*, *C. lari*
83  subsp*. Concheus* and *C. pinnipediorum* subsp*. Caledonicus*), suggesting that the communities
84  containing these subspecies might be from distinct novel species. These subspecies were
85  further evaluated using a network-based approach, as described below.
86
**Network analysis identifies potentially new *Campylobacter* species**
We conducted a network analysis based on Mash distances (see methods for details). In this
network, nodes and edges represent genomes and the estimated identities (1 – Mash;
minimum threshold of 95%), respectively. Each community in the resulting network is
considered a different species, as described elsewhere[9]. Species networks inferred with
genomic distance or identity are highly structured, with few overlapping communities[9,12]. In
our *Campylobacter* network, we found 65 genome communities (Figure 1), which is above the
expected number considering the 34 type genomes listed in LPSN. Out of the 65 communities,
only 36 contained type strain genomes. We found no communities with more than one
species type genome, while two subspecies formed separated communities (*C. lari* subsp.
*concheus* and *C. fetus* subsp. *testudinum*). The communities with a type genome (from species
or subspecies) were named after their , propagating this classification to all the genomes of
the community.
We named the 29 communities without a type genome as *Campylobacter "sppX"*,
where X represents an arbitrary community number (see methods). We found three large
unnamed communities: *Campylobacter spp5* (134 genomes), corresponding to the *C. concisus*
GSII; *Campylobacter spp12* (78 genomes)*,* representing the well-known *C. coli Clade III*; and
*Campylobacter spp33* (37 genomes), comprising genomes from isolates exclusively obtained
from swine feces[13] that form a sister group with *C. lanienae* (Figure 2). These results support
that a significant part of the *Campylobacter* diversity remains largely unexplored.
Besides investigating potentially new species, we also constructed a distance tree to
infer the proximity of their respective communities to type species (Figure 2). We observed
that *C. coli* is the sister group of *C. jejuni*, as previously described[5]. The *C. coli* population is
structured in three main clades[14]. We observed that the *C. coli* Clade III is an independent
community in the distance network and is paraphyletic in relation to the other *C. coli* genomes
(Figure 2). Although *C. coli* Clade II is a monophyletic sister group of Clade I in the distance
tree, the high number of network connections with other *C. coli* genomes does not support
its separation as an independent community.
We also corroborated the split of *C. concisus* into two distinct genomospecies (GSI and
GSII), as previously proposed[15]. Within the 216 *C. concisus* genomes available in RefSeq, 82
belong to *C. concisus* GSI, while 134 genomes formed a distinct community named as
*Campylobacter spp5* (*C. concisus* GSII). The two communities also differ in GC content (37.22%
in *C. concisus* and 39.19% *in Campylobacter spp5*; Wilcoxon test, p-value < 0.001) and genome
sizes (1.9 Mbp in *C. concisus* and 2.0 Mbp in *Campylobacter spp5*; Wilcoxon test, p-value <

0.001), supporting their split into two different species. Accordingly, it has been proposed
that the GC content within *Campylobacter* species should not vary by more than 1%[16].
The distance tree also provides insights into novel species and niche segregation. For
example, *C. fetus* subsp*. testudinum* is probably a new species, typically found in mammals.
Furthermore, we also observed a clear division of *C. lari* into two clades, corresponding to the
two subspecies previously proposed based on 16S rRNA: *C. lari* subsp. *Lari* and *C. lari* subsp.
*concheus*[17]. These results imply that the *C. lari* phylogeny also warrants revisions.
128
### *Campylobacter* phylogeny of representative genomes and species reclassification
We used all core single-copy genes from 65 representative genomes (Table 2) to reconstruct
the *Campylobacter* phylogenetic tree. This phylogenetic reconstruction uncovered a
remarkable difference in GC content across clades. (Figure 3). For example, the clade
comprising *C. showae* and *C. rectus* have GC content greater than 40%, far greater than the
30.4% found in the *C. jejuni* clade.
We also used the representative genomes to estimate a misclassification rate of 7.57%
in *Campylobacter spp* (247 misclassified genomes; Table S3). We considered as misclassified
only those genomes previously annotated at the species level and re-assigned either to a
different validly published species or to a new community (*Campylobacter sppX*). Therefore,
genomes identified as "*Campylobacter sp.*" were not considered misclassified if assigned to a
community.
We found extreme misclassification levels in eight known species (Table 1),
particularly *C. showae* (81.82%, n=11). Given the small number of *C. showae* genomes, it
would be premature to reach a conclusion on this species. However, as most of the genomes
formed a cluster apart from the type species, we hypothesize that the type genome may not
satisfactorily represent the group. Species such as *C. concisus* and *C. coli* showed significant
reclassification levels due to their known heterogeneity[5,15] (61.86%, n=215 and 7.96%, n=980,
respectively). Further, genomes submitted to the RefSeq database as *C. aviculae*, *C.*
*estrildidarum*, *C. taeniopygiae,* and *C. troglodytis* were marked as reclassified because of the
absence of their respective representative genomes in LPSN at the time of the analysis.
Finally, 92 genomes previously identified as *Campylobacter sp.* (54.76%)*.* were assigned to
communities composed by known species, particularly *C. jejuni*.
We also compared the results obtained by the genome analysis with the
*Campylobacter* classification provided by GTDB (Release 207, downloaded in August 2022).
GTDB has a standardized framework to classify genomes in higher taxonomic ranks (e.g.,
genus)[18]  by following a phylogenomic approach based on a set of conserved single-copy
proteins. This scheme divides *Campylobacter* in five different genera (Table S1):
*Campylobacter*, *Campylobacter A*, *Campylobacter B*, *Campylobacter D*, and *Campylobacter E*.
Importantly, this distribution is coherent with the phylogenetic reconstruction and
community analysis described here. The largest genus, *Campylobacter E*, comprised the *C. coli*
and *C. jejuni* communities, whereas the second largest genus, *Campylobacter A*, harbored the
monophyletic group of *C. concisus*, *Campylobacter spp5*, *C. curvus*, and *C. pinnipediorum*.

A comparison with the TYGS (Type Strain Genome Server)[19,20] was also conducted
using the representative genomes of each community (Table S4). TYGS corroborated our
findings, with most of the unknown communities being identified as potentially new species.
We also found that *Campylobacter spp55* (n=1), and *Campylobacter spp38* (n=2) were
identified as *C. bilis* and *C. vulpis* respectively, both found in the most recent LPSN update[2]
(Table S2). Finally, the TYGS classification of representative genomes of the most well studied
communities were consistent with our observations.
169

**Pangenome analysis**
The pangenome analysis of a bacterial species provides insights into its lifestyle[21]. A
pangenome is the total set of non-redundant genes in a species. It comprises the core
genome, the set genes present in all isolates; the accessory genome, with genes present in
more than one but not in all isolates; and the isolate-exclusive genes[22]. We computed the
genomic fluidity and pangenome openness to investigate the overall architecture of the
*Campylobacter* pangenome (Table 3, Figure 4). The genomic fluidity measures how dissimilar
genomes are at the gene level[23], while pangenome openness reflects the propensity to detect
new genes, as new genomes are included in the analysis. Since sample size can bias
pangenome estimation, we analyzed only those communities with more than 20 genomes.
Previous analyses on the connection between pangenome openness and genomic
fluidity have shown that these two metrics are not necessarily correlated, with both
pangenome fluidity and saturation being related to its openness[24]. We found that genomic
fluidity of *Campylobacter* is negatively correlated with saturation coefficient (α) ($\rho$ = -0.8733;
p-value = 0.0004), with greater values of α representing a closer pangenome. We found a
significant regression equation ($F_{1;9}$ = 28.927; p-value < 0.0001), with an $R^2$ of 0.73 (Figure 4b)
and predicted genomic fluidity of $0.8187 - 0.8467\alpha$, meaning that the genomic fluidity
decreased 0.8467 units for each unit of the parameter α. This result provides a model for
estimating the genomic fluidity of species without enough genomes to be included in the
pangenome analysis.
We observed that *C. hepaticus* is genetically conserved and homogeneous. This
species has the lowest diversity and the highest saturation coefficient (closed pangenome)
(Figure 4a) across *Campylobacter* communities, supporting a low genomic diversity and
reduced HGT in this species. However, it is important to mention that many *C. hepaticus*
genomes are from the same bioproject, which might result in unrealistic diversity estimates
for this species. On the other hand, species such as *C. jejuni* and *C. coli* have a high genomic
fluidity, which is in line with the generalist ecological character of both species, with isolates
from various host sources.
198

**DISCUSSION**
The *Campylobacter* genus is often associated with gastrointestinal diseases and antibiotic
resistance, having a strong study bias towards clinically relevant species such as *C. jejuni* and
*C. coli*. Here, we analyzed 3,261 *Campylobacter* genomes from several species and sources.

We identified 65 different communities within *Campylobacter*, way over the 38 species
originally listed on the LPSN. We found a portion of these novel communities (here considered
as potential new species) to be closely related to some *Campylobacter* groups such as the *C.*
*lari* and *C. fetus* groups. Our ANI-based network analysis also showed that 7.57% of the
*Campylobacter* genomes warrants phylogenetic reclassification.
*C. concisus* is an oral bacterium that consists of two genomospecies (GSI and GSII) that
are undistinguishable at the phenotypic level[15]. Our results support this division, as we
identified two *C. concisus* communities*,* the first containing the species' type strain with 82
genomes, and *Campylobacter spp5* with 134 genomes. Similarly, we also found support to
separate *C. coli* in two communities: one comprising the original type strain and other 902
genomes, and a second community with 78 genomes (referred as *Campylobacter spp12*).
We also reported strong evidence for potentially novel species into the *C. fetus* group.
*Campylobacter spp33* comprises 37 genomes of isolates obtained only from swine feces[13],
supporting a specific niche occupation. Interestingly, specific niche occupation is a common
feature of the *C. fetus* group [25]. We can further divide this clade using characteristics like the
low capacity to cause diseases in humans (*C. lanienae*-related species) and selenium
metabolism deficiency (*Campylobacter spp33* clade[13]). Finally, the identification of
*Campylobacter spp33* as a novel species should be further investigated for a more thorough
and formal description.
We found that genomic fluidity and pangenome saturation coefficient are negatively
correlated in the *Campylobacter* genus. Although these two parameters are usually
conceptually associated, previous findings have shown that there was not a direct and
unequivocal correlation between their values, but that fluidity was just one of the
determinants of genome openness[24]. *C. hepaticus* showed a close and homogeneous
pangenome, although one must consider that a high fraction of the *C. hepaticus* genomes
were from the same study, which could result in the underestimation of the *C. hepaticus*
diversity. Finally, it is worth mentioning that more in-depth investigations of the large
unnamed communities described here should be conducted to validate them as novel
species.
232
**METHODS**
**Dataset collection**
We downloaded 3,319 *Campylobacter* genomes available in RefSeq using Datasets v4.2.0
(https://www.ncbi.nlm.nih.gov/datasets/) in August 2020. We used BUSCO v4.1.0[26] and
removed the genomes with completeness lower than 90%, and duplication greater than 5%.
Genomic metadata were gathered from RefSeq using the Reutils package v. 0.2.3
(http://mirrors.ucr.ac.cr/CRAN/web/packages/R.utils/R.utils.pdf).
All the genomes were analyzed using Mash v1.1[27]. We compared different Mash
sketch values to find the parameter with the lowest error rate[9]. We used the mean distances
generated by sketch sizes of 1000, 2000, and 5000 bp, removing genome distances greater

than 0.8. We used the reciprocal of Mash distance to estimate the identity (1 – Mash). For
type strain analyses, we computed ANI using pyani v0.2.10[28].
The network analyses of the distances between the genomes were conducted using
igraph v1.2.7[29]. We generated a weighted graph with nodes and edges representing genomes
and genetic identities between them, respectively. Only edges representing identities greater
than 95% were retained.
249

**Type strains validation**

The pre-set *Campylobacter* type strains were retrieved from LPSN[2] in March 2023. The valid
accession numbers of each species and subspecies were manually gathered, and we used the
information in LPSN to identify validly reported and documented type genomes (38
representing valid species, 7 representing valid subspecies) (Table S2). Next, we conducted
an ANI analysis with Pyani v0.2.10[28] to verify that all the genome pairs are at least 95%
identical to each other.
257

**Network analysis**

We employed the label propagation algorithm to detect communities in the genome distance
network, as previously reported[9]. Communities with type genomes were named after them,
while communities without a type genome were arbitrarily identified as *Campylobacter sppX*,
where X represents the number of the community. These communities had one of its
members randomly assigned as its representative genome. The named communities were
contrasted with the RefSeq taxonomic classification to estimate the level of misclassification.
265

**Phylogeny and pangenome characterization**

After selecting a representative genome from each *Campylobacter* community, we used
Orthofinder v.2.5.2[30] to infer orthologs. Single-copy genes were used to reconstruct the
phylogenetic tree using IQ-TREE v.1.6.12[31] with the TEST argument to find the best fitting
model. One thousand bootstrap replicates were generated to assess the significance of
internal nodes. The *Helicobacter pylori* strain ATCC 43504 (Accession number:
GCF_900478295.1) were used as outgroup. We used ggtree[32] to visualize the reconstructed
tree. GTDB data (July 2022) and TYGS data (March 2023) were used to analyze different
phylogenetic groups within *Campylobacter*[19,20,33].
We conducted the pangenome analysis for those communities with more than 20
genomes with Roary v3.13.0[34]. Micropan v2.1[35] and Pagoo v0.3.9[36] were used to assess
pangenome fluidity values and openness values, respectively.
278

**Declaration of Competing Interest**

280 The authors declare no conflict of interest.
281

**Author contributions**

283 Conceived the study: HP-A and TMV. Data analysis: HP-A, SH-J. Funding, project coordination

284    and infrastructure: TMV. Manuscript writing: HP-A, SH-J, TMV.
285

**Acknowledgments**
This work was supported by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado
do Rio de Janeiro (FAPERJ; grants E-26/203.309/2016 and E-26/203.014/2018), Coordenação
de Aperfeiçoamento de Pessoal de Nível Superior- Brasil (CAPES; Finance Code 001), and
Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). The funding agencies
had no role in the design of the study and collection, analysis, and interpretation of data and
in writing.
293

**References**

1.    Heimesaat, M. M., Backert, S., Alter, T. & Bereswill, S. Human Campylobacteriosis-A
Serious Infectious Threat in a One Health Perspective. *Curr Top Microbiol Immunol*
**431**, 1–23 (2021).
2.    Parte, A. C. LPSN - List of prokaryotic names with standing in nomenclature. *Nucleic*
*Acids Res* **42**, 613–616 (2014).
3.    Geneva: World Health Organization. *Prioritization of pathogens to guide discovery,*
*research and development of new antibiotics for drug-resistant bacterial infections,*
*including tuberculosis.* (2017).
4.    Epping, L., Antão, E. M. & Semmler, T. Population Biology and Comparative
Genomics of Campylobacter Species. in *Current Topics in Microbiology and*
*Immunology* vol. 431 59–78 (Springer Science and Business Media Deutschland
GmbH, 2021).
5.    Sheppard, S. K. *et al.* Progressive genome-wide introgression in agricultural
Campylobacter coli. *Mol Ecol* **22**, 1051–1064 (2013).
6.    Sheppard, S. K., Guttman, D. S. & Fitzgerald, J. R. Population genomics of bacterial
host adaptation. *Nature Reviews Genetics* vol. 19 549–565 Preprint at
https://doi.org/10.1038/s41576-018-0032-z (2018).
7.    Colles, F. M., Jones, K., Harding, R. M. & Maiden, M. C. J. Genetic Diversity of
Campylobacter jejuni Isolates from Farm Animals and the Farm Environment. *Appl*
*Environ Microbiol* **69**, 7409–7413 (2003).
8.    Poretsky, R., Rodriguez-R, L. M., Luo, C., Tsementzi, D. & Konstantinidis, K. T.
Strengths and limitations of 16S rRNA gene amplicon sequencing in revealing
temporal microbial community dynamics. *PLoS One* **9**, (2014).
9.    Passarelli-Araujo, H., Franco, G. R. & Venancio, T. M. Network analysis of ten
thousand genomes shed light on Pseudomonas diversity and classification. *Microbiol*
*Res* **254**, 126919 (2022).
10.   Pruitt, K. D., Tatusova, T. & Maglott, D. R. NCBI Reference Sequence (RefSeq): A
curated non-redundant sequence database of genomes, transcripts and proteins.
*Nucleic Acids Res* **33**, (2005).
11.   Tettelin, H. & Medini, D. *The pangenome: Diversity, dynamics and evolution of*
*genomes. The Pangenome: Diversity, Dynamics and Evolution of Genomes* (Springer
International Publishing, 2020). doi:10.1007/978-3-030-38281-0.
12.   Passarelli-Araujo, H., Jacobs, S. H., Franco, G. R. & Venancio, T. M. Phylogenetic
analysis and population structure of Pseudomonas alloputida. *Genomics* **113**, 3762–
3773 (2021).
13.   Miller, W. G. *et al.* Comparative genomic analysis identifies a campylobacter clade
deficient in selenium metabolism. *Genome Biol Evol* **9**, 1843–1858 (2017).
14.   Sheppard, S. K., McCarthy, N. D., Falush, D. & Maiden, M. C. J. Convergence of

Campylobacter species: Implications for bacterial evolution. *Science (1979)* **320**, 237–
239 (2008).

15.  Aabenhus, R., On, S. L. W., Siemer, B. L., Permin, H. & Andersen, L. P. Delineation
of Campylobacter concisus genomospecies by amplified fragment length
polymorphism analysis and correlation of results with clinical data. *J Clin Microbiol* **43**,
5091–5096 (2005).

16.  Meier-Kolthoff, J. P., Klenk, H. P. & Göker, M. Taxonomic use of DNA G+C content
and DNA-DNA hybridization in the genomic age. *Int J Syst Evol Microbiol* **64**, 352–
356 (2014).

17.  Debruyne, L., On, S. L. W., De Brandt, E. & Vandamme, P. Novel Campylobacter lari-
like bacteria from humans and molluscs: Description of Campylobacter peloridis sp.
nov., Campylobacter lari subsp. concheus subsp. nov. and Campylobacter lari subsp.
lari subsp. nov. *Int J Syst Evol Microbiol* **59**, 1126–1132 (2009).

18.  Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat Biotechnol* **36**, 996 (2018).

19.  Meier-Kolthoff, J. P. & Göker, M. TYGS is an automated high-throughput platform for
state-of-the-art genome-based taxonomy. *Nat Commun* **10**, (2019).

20.  Meier-Kolthoff, J. P., Carbasse, J. S., Peinado-Olarte, R. L. & Göker, M. TYGS and
LPSN: A database tandem for fast and reliable genome-based classification and
nomenclature of prokaryotes. *Nucleic Acids Res* **50**, D801–D807 (2022).

21.  Brockhurst, M. A. *et al.* The Ecology and Evolution of Pangenomes. *Current Biology*
vol. 29 R1094–R1103 Preprint at https://doi.org/10.1016/j.cub.2019.08.012 (2019).

22.  Tettelin, H. *et al.* Genome analysis of multiple pathogenic isolates of Streptococcus
agalactiae: Implications for the microbial "pan-genome". *Proceedings of the National*
*Academy of Sciences* **102**, 13950–13955 (2005).

23.  Kislyuk, A. O., Haegeman, B., Bergman, N. H. & Weitz, J. S. Genomic fluidity: An
integrative view of gene diversity within microbial populations. *BMC Genomics* **12**,
(2011).

24.  Maistrenko, O. M. *et al.* Disentangling the impact of environmental and phylogenetic
constraints on prokaryotic within-species diversity. *ISME Journal* **14**, 1247–1259
(2020).

25.  Mourkas, E. *et al.* Host ecology regulates interspecies recombination in bacteria of the
genus Campylobacter. *Elife* **11**, (2022).

26.  Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

27.  Ondov, B. D. *et al.* Mash: Fast genome and metagenome distance estimation using
MinHash. *Genome Biol* **17**, (2016).

28.  Pritchard, L., Cock, P., Esen, Ö. & Yt, ; Pyani. *Research Software. Software* (2019)
doi:10.5281/zenodo.2584238.

29.  Han, W.-S., Lee, J., Pham, M.-D. & Yu, J. X. iGraph. *Proceedings of the VLDB*
*Endowment* **3**, 449–459 (2010).

30.  Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for
comparative genomics. *Genome Biol* **20**, (2019).

31.  Nguyen, L. T., Schmidt, H. A., Von Haeseler, A. & Minh, B. Q. IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
*Evol* **32**, 268–274 (2015).

32.  Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. Y. ggtree: an r package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol Evol* **8**, 28–36 (2017).

33.  Parks, D. H. *et al.* GTDB: An ongoing census of bacterial and archaeal diversity
through a phylogenetically consistent, rank normalized and complete genome-based
taxonomy. *Nucleic Acids Res* **50**, D785–D794 (2022).

34.  Page, A. J. *et al.* Roary: Rapid large-scale prokaryote pan genome analysis.
*Bioinformatics* **31**, 3691–3693 (2015).

35.   Snipen, L. & Liland, K. H. micropan: An R-package for microbial pan-genomics. *BMC*
*Bioinformatics* **16**, 1–8 (2015).
36.   Ferrés, I. & Iraola, G. An object-oriented framework for evolutionary pangenome
analysis. *Cell Reports Methods* **1**, (2021).

394
395
396
397

398 **Tables**

*Table 1: Rate of misclassified genomes per species in the RefSeq database. Genomes originally assigned as belonging to a species and appeared an unexpected community (different from that of their respective type strain genome) were considered misclassified.*

| Original classification | Reclassification rate (%) |
|---|---|
| *C. coli* | 7.96 (78/980) |
| *C. concisus* | 61.86 (133/215) |
| *C. fetus* | 27.27 (10/35) |
| *C. hepaticus* | 3.85 (1/26) |
| *C. jejuni* | 0.39 (6/1551) |
| *C. lanienae* | 6.45 (2/31) |
| *C. lari* | 50.00 (9/18) |
| *C. showae* | 81.82 (9/11) |

*Table 2: Representative genomes of each community detected in this study. The communities were named after their type strains. Communities without type strains were arbitrarily named with a sequential suffix and received a random representative genome.*

| Community name | Refseq accession | Community name | Refseq accession |
|---|---|---|---|
| *C. armoricus* | GCF_013372105.1 | *Campylobacter spp57* | GCF_012978795.1 |
| *C. avium* | GCF_002238335.1 | *Campylobacter spp58* | GCF_012978835.1 |
| *C. blaseri* | GCF_013201895.1 | *Campylobacter spp62* | GCF_013416015.1 |
| *C. canadensis* | GCF_013177655.1 | *Campylobacter spp63* | GCF_900573945.1 |
| *C. coli* | GCF_003590975.1 | *Campylobacter spp64* | GCF_900573985.1 |
| *C. concisus* | GCF_001298465.1 | *Campylobacter spp65* | GCF_902795495.1 |
| *C. corcagiensis* | GCF_013201645.1 | *C. sputorum* | GCF_008802035.1 |
| *C. cuniculorum* | GCF_002104335.1 | *C. subantarcticus* | GCF_000816305.1 |
| *C. curvus* | GCF_013372125.1 | *C. upsaliensis* | GCF_900637395.1 |
| *C. fetus* | GCF_900475935.1 | *C. ureolyticus* | GCF_013372225.1 |
| *C. fetus testudinum* | GCF_002973655.1 | *C. volucris* | GCF_008801985.1 |
| *C. geochelonis* | GCF_013201685.1 | | |
| *C. gracilis* | GCF_001190745.1 | | |
| *C. helveticus* | GCF_002080395.1 | | |
| *C. hepaticus* | GCF_001687475.2 | | |
| *C. hominis* | GCF_900446395.1 | | |
| *C. hyointestinalis* | GCF_013372145.1 | | |
| *C. iguaniorum* | GCF_000736415.1 | | |
| *C. insulaenigrae* | GCF_900637885.1 | | |
| *C. jejuni* | GCF_000254515.1 | | |
| *C. lanienae* | GCF_002139935.1 | | |
| *C. lari* | GCF_013372185.1 | | |
| *C. lari concheus* | GCF_008245025.1 | | |
| *C. mucosalis* | GCF_013372205.1 | | |
| *C. novaezeelandiae* | GCF_004323845.1 | | |
| *C. ornithocola* | GCF_013201605.1 | | |
| *C. peloridis* | GCF_000816785.1 | | |
| *C. pinnipediorum* | GCF_002021925.1 | | |
| *C. portucalensis* | GCF_009690845.1 | | |
| *C. rectus* | GCF_004803795.1 | | |
| *C. showae* | GCF_004803815.1 | | |
| *Campylobacter spp11* | GCF_000313615.1 | | |
| *Campylobacter spp12* | GCF_001489835.1 | | |
| *Campylobacter spp13* | GCF_000344295.2 | | |
| *Campylobacter spp23* | GCF_000816245.1 | | |
| *Campylobacter spp33* | GCF_900446565.1 | | |
| *Campylobacter spp34* | GCF_002139105.1 | | |
| *Campylobacter spp35* | GCF_002139895.1 | | |
| *Campylobacter spp36* | GCF_002179165.1 | | |
| *Campylobacter spp38* | GCF_002738235.1 | | |
| *Campylobacter spp41* | GCF_004357905.1 | | |
| *Campylobacter spp42* | GCF_004378855.1 | | |
| *Campylobacter spp44* | GCF_012978635.1 | | |
| *Campylobacter spp45* | GCF_004803855.1 | | |
| *Campylobacter spp46* | GCF_005406205.1 | | |
| *Campylobacter spp47* | GCF_005406215.1 | | |
| *Campylobacter spp48* | GCF_005406225.1 | | |
| *Campylobacter spp49* | GCF_010092865.1 | | |
| *Campylobacter spp5* | GCF_902460635.1 | | |
| *Campylobacter spp51* | GCF_006864365.1 | | |
| *Campylobacter spp52* | GCF_006864425.1 | | |
| *Campylobacter spp53* | GCF_006864455.1 | | |
| *Campylobacter spp54* | GCF_008633875.1 | | |
| *Campylobacter spp55* | GCF_009362725.1 | | |

*Table 3: Pangenome features for each community with 20 or more genomes.*

| Community name | Fluidity | Openness (α value) | Gene families | Core genes |
|---|---|---|---|---|
| *C. coli* | 0.209 | 0.724 | 12,477 | 619 |
| *C. concisus* | 0.216 | 0.770 | 6682 | 671 |
| *C. fetus* | 0.125 | 0.799 | 4079 | 1097 |
| *C. fetus testudinum* | 0.080 | 0.936 | 2730 | 1239 |
| *C. hepaticus* | 0.031 | 1.021 | 1753 | 1321 |
| *C. hyointestinalis* | 0.236 | 1.091 | 4255 | 1003 |
| *C. jejuni* | 0.255 | 0.629 | 18,350 | 469 |
| *C. lanienae* | 0.116 | 0.943 | 2766 | 1146 |
| *Campylobacter spp12* | 0.074 | 0.717 | 3113 | 1126 |
| *Campylobacter spp33* | 0.140 | 0.879 | 3345 | 1136 |
| *Campylobacter spp5* | 0.230 | 0.706 | 9431 | 869 |

**Figures**

**Figure 1.** *Campylobacter* network structure. Nodes represent genomes and edges connect those with at least 95% identity. Type strains are highlighted in red. Communities without type strains represent potentially novel species. *C. jejuni* and *C. coli* communities have a large number of genomes, which can make it more difficult to understand the network as a whole, they were omitted to improve clarity.

**Figure 2.** Mash distance tree of *Campylobacter*. Main *Campylobacter* groups are represented in different colors. Branches with many taxa were collapsed to improve visibility.

**Figure 3.** Phylogenetic reconstruction and GC content in *Campylobacter*. The Maximum likelihood tree was reconstructed using core single-copy genes from one representative genome from each of the 65 *Campylobacter* communities. *H. pylori* was used as outgroup.

**Figure 4.** Pangenome openness and fluidity of *Campylobacter* species. **a)** Pangenome cumulative curve from eleven *Campylobacter* species from communities with at least 20 genomes. The number of genomes of each species are indicated within parentheses. Due to the high number *C. jejuni* and *C. coli* genomes, we limited the x-axis to 200 genomes. **b)** Simple linear regression to predict *Campylobacter* pangenome fluidity (φ) based on pangenome openness (α). This analysis was conducted with the same communities from panel a.

*C. jejuni*

*C. coli*

Clade II

Clade III

GSII

GSI

| | | | |
|---|---|---|---|
| ■ *C. helveticus* | ■ *C. hepaticus* | ■ *C. spp33* | ■ *C. curvus* |
| ■ *C. volucris* | ■ *C. coli (clade II)* | ■ *C. hyointestinalis* | ■ *C. concisus (GSI)* |
| ■ *C. armonicus* | ■ *C. spp12 (clade III)* | ■ *C. fetus testudinum* | ■ *C. spp5 (GSII)* |
| ■ *C. lari concheus* | ■ *C. sputorum* | ■ *C. fetus* | |
| ■ *C. lari* | ■ *C. lanienae* | ■ *C. pinnipediorum* | |

**a**

Gene Families

Genomes

**b**

Fluidity (φ)

Openness (α)

$\varphi = -0.847\alpha + 0.819$
$R^2\ \text{adj} = 0.73$

*C. spp5* (134)   *C. coli* (903)   *C. spp33* (37)   *C. spp12* (78)
*C. jejuni* (1638)   *C. hyointestinalis* (41)   *C. lanienae* (29)   *C. hepaticus* (25)
*C. concisus* (82)   *C. fetus* (46)   *C. fetus testudinum* (35)