Single Cell Transcriptomics Reveals the Hidden Microbiomes of Human Tissues

Gita Mahmoudabadi[1], Sheela Crasta[2], Tabula Sapiens Consortium[3], Stephen R. Quake[1,2,4,*]

[1]Department of Bioengineering, Stanford University, Stanford CA, USA
[2]Authors and their affiliations are listed at the end
[3]Department of Applied Physics, Stanford University, Stanford CA, USA
[4]Chan-Zuckerberg Initiative, Redwood City CA, USA
*Corresponding author. Email: steve@quake-lab.org

Abstract

The human microbiome has been studied extensively across those sites in the body that are readily accessible to sampling. Internal organs and tissues, however, have remained largely unexplored and, in the absence of infectious disease, are widely assumed to be free of microorganisms. Using single-cell transcriptomic data from the Tabula Sapiens spanning 11 human organ donors, 19 tissue types, 400,000+ annotated cells, 100+ cell types, and ~70 billion sequences, we created an atlas of the human tissue microbiome with cell type resolution across three domains of life which we refer to as the Tabula Sapiens Microbiome. Together with data from 8 additional donors derived from a different set of experimental techniques for validation, we identified sequences from numerous bacterial, viral and fungal species in human tissues. We mapped the likely microbial flow routes in the body from external-facing microbiomes and natural environments to internal tissues and tumors, demonstrating the existence of several unexpected routes. In the case of tumor microbiomes, we find a considerable number of bacterial genera found in tumors are detectable across tumor-free tissues, suggesting that tumor microbiomes are in part sourced from adjacent normal tissues, even those from distant sites. Finally, by increasing the resolution of sampling from tissues to cell types, we quantified the microbial load and diversity across different human cell types to reveal a network of host cell type and microbe associations. For example, we identified traces of both latent and active Epstein Barr Virus infections in various cell types such as splenic plasma cells. Broad exploration of the human tissue microbiomes may provide insights which ultimately are of clinical importance.

Introduction

Microbial life contains a vast repertoire of biochemical pathways that enable it to colonize even the most extreme environments on the planet. It is no wonder that the human body, an energy-rich environment for trillions of our own cells, would be host to many highly diverse microbes[1,2]. Historically, understanding of microbes in connection to human health has emerged from waves of encounters with devastating pathogens such as *Mycobacterium tuberculosis*, *Vibrio cholerae*, and most recently, SARS-CoV2. In the last two decades our view of human-microbe interactions has been transformed through increasingly powerful sequencing technologies which have enabled broad surveys of the microbial world[3–8]. As such, previously held dogma has been upended; one such example of this is the assertion that internal organs such as the healthy human lung are sterile environments[9].

The study of the human microbiome has seen an explosion of data connecting communities of microbes or individual species to a variety of metabolic[10,11], neurological[12,13], cardiovascular[14,15], and autoimmune[16,17] diseases, among others. Most

microbiome studies, however, have delineated microbial communities at sites easily
accessible to sampling such as the human intestinal tract via feces, the mouth, and the
skin. Less accessible internal sites, on the other hand, are largely assumed to be free of
microorganisms, as they are protected by layers of epithelial and endothelial tissue.
However, a few recent studies have challenged this assumption. For example, human
tumors were shown to contain tumor-specific microbial sequences, providing a possible
diagnostic approach for cancer detection[18]. Other studies similarly identified the presence
of tumor-specific mycobiomes and microbiomes including intracellular bacteria[19–24].
Microbiomes in both tumor-free and tumor-derived breast, esophageal, oropharyngeal,
gastrointestinal, and colorectal tissues have also been shown[25–27].
49
We assembled the first atlas of human tissue and cell type microbiomes by mining the
raw sequences from one of the largest single-cell transcriptomic datasets of human
tissues called the Tabula Sapiens[28] which encompasses 400,000+ annotated human
cells, spanning 100+ cell types and 19 tissues, originating from 11 human organ donors.
We identified high-confidence viral, bacterial, and fungal sequences originating from
human single-cell transcriptomes, together forming the Tabula Sapiens Microbiome
(TSM) dataset. Using a series of experimental and bioinformatic filters including a
machine learning model for decontamination and a Large Language Model for habitat
classification, we rigorously decontaminated and quality controlled this dataset.
59
Moreover, we combined our data with 16 external datasets (**SI Figure 1.a-b**), 10 of which
were used for creating and benchmarking a pipeline for the identification of microbial
sequences. Some examples of these databases include the Human Gastrointestinal
Bacteria Culture Collection (HBC)[29], the expanded Human Oral Microbiome Dataset
(HOMD)[30], Human Microbiome Project (HMP)[4], and others. The other external databases,
including Narunsky-Haziza *et al*.[20] (tumor mycobiomes), Nejman *et al.* (tumor
microbiomes)[21], the Unified Human Gastrointestinal Genome[5] (UHGG) collection,
Genomes from Earth's Microbiomes (GEM)[7], Pathosystems Resource Integration Center
(PATRIC)[31], and others were used for decontamination and analysis. For example, we
explored the intersection of species in TSM tissues and those found at tumor sites and at
sites accessible to sampling, or what we will refer to as "external-facing microbiomes"
(e.g. the gastrointestinal, oral and skin microbiomes), as well as natural environments, to
chart out possible microbial flow routes throughout the body. We report the inter-and intra-
donor variability of tissue microbiomes and characterize microbial load and diversity
across human cell types. Moreover, using an additional cohort of 8 donors, we devised
an assay to extract virus-like and bacterial-like particles from human tissues and explored
them using imaging and DNA sequencing. We will refer to this dataset as the Extracted
Human Tissue Microbiome (EHTM) dataset. We further show that the human tissue and

cellular microbiomes, beyond cell-free or capsid-free nucleic acids, may also include
intact organisms.
80

81 Results

Computational pipeline for identification of microbial sequences from multiple
biomes in single-cell datasets

We created a computational pipeline written in Snakemake[32] to identify viral, bacterial and
fungal sequences in single-cell transcriptomic data. Briefly, we use UMI-tools[33] to filter
and extract cell barcodes and UMIs from fastqs, and then use STAR[34] to align resulting
sequences against a reference library containing the human genome and ERCCs. After
the removal of low-complexity sequences from the unmapped reads using a custom
script, we use BLASTn[35] to align reads against several custom databases followed by the
nt database, which is one of largest publicly available databases. We refer to this pipeline
as SIMBA. For a more detailed description of this pipeline see **SI: SIMBA, a**
computational pipeline for the identification of bacteria, viruses and fungi in single
cells, along with **SI Figure 1.a.** We also benchmarked this pipeline against several truth
datasets to show that it is highly precise **(**see **SI: Benchmarking SIMBA's precision**
and recall as well as **SI Tables 1-8).** We provide a full explanation for why we chose not
to rely on faster aligners or smaller reference databases given their higher false positive
rates in samples that have low ratios of non-host to host sequences (See **SI: SIMBA, a**
computational pipeline for the identification of bacteria, viruses and fungi in single
cells, as well as **SI: Benchmarking Kraken2 precision and recall,** as we all as **SI**
Tables 9-11 and **SI Figure 2)**. A common theme in our study is that in the tug-of-war
between precision and recall, we choose precision as we decided it was more important
to us to identify microbial reads precisely and with high confidence than it was to capture
more of them.
104

Decontamination and quality filtering of the Tabula Sapiens Microbiome dataset

In the previous section and its supplementary material, we address a major source of
false positives in microbial datasets, which are false positives generated during
alignments. Another important source of false positives is physical contamination. We
discuss in depth the measures taken during tissue procurement and transport to prevent
potential contaminants (see **SI: Measures against contamination during tissue**
recovery and transport). We acquired samples of reagents and instruments that were
used to process tissues and sequence single cells. Even when the same reagent was

used across different labs, we obtained each lab's specific sample of that reagent to
account for lab-specific contaminants. Following the same protocols as the original
experiments, we deeply sequenced more than 200 reagent and instrument controls using
five 10X lanes (**SI Table 12**), and eliminated contaminant species (filter 1). We then
eliminated putative bacterial (filter 2) and fungal (filter 3) contaminants previously
identified by Salter *et al.*[36] and Narunsky-Haziza *et al*[20]. For a detailed description of these
three filters and their impact on the dataset, see **SI: Identification of contaminants**
based on negative controls and external datasets and **SI Tables 12-13**.
121
To further account for any potentially missed contaminants, we built 6 types of machine
learning models to learn the shared features of the known and putative contaminants
acquired from the first three filters. As our fourth filter, we applied the best performing
model type, which was the gradient boosting classifier (test AUPR and AUROC 95%
confidence intervals were 0.75-0.78 and 0.82-.84, respectively) to identify putative
contaminants in the portion of the dataset that was withheld from both training and testing
(see **SI: Machine learning models of contamination, SI Tables 14-15** and **SI Figure**
1.c-d). Additionally, we observed a bimodal distribution of alignments and removed
species that are represented primarily by the lower-quality alignment peak, which served
as the 5th filter (see **SI: Gaussian Mixture Model of alignments,** and **SI Figure 3).** To
further verify our findings, we imposed a 6th filter which selects taxa that have been
previously identified in the human microbiome by three large studies, namely **H**MP,
UHGG, or **G**EM datasets, or what we abbreviate as HUE studies (see **SI: Identifying**
putative habitats of origin for 500+ bacterial genera in the Tabula Sapiens Microbial
dataset using the Genomes from Earth's Microbiomes, the Human Microbiome
Project, and the Unified Human Gastrointestinal Genome collection and **SI Figure**
4). For the remaining taxa that had not appeared in HUE studies (7th filter), we
benchmarked (precision of 98% and a recall of 75%) and programmatically employed
openAI's Davinci Large Language Model to further select human-associated taxa (see
SI: Benchmarking the Davinci large language model for classification of microbial
species based on their habitat, SI Tables 16-18 and **SI Figure 1.e**). Habitat-based
filters were placed to select commonly known human-resident or human-associated
microbes from the rest.
145
The output of SIMBA that has been filtered to contain only sequences from the three
biomes of interest is referred to as the Tabula Sapiens Microbiome (TSM) dataset F0, to
signify that it has not been filtered based on contaminants. The dataset filtered only by
the first filter is referred to as TSM F1. The dataset filtered by the first two filters is referred
to as TSM F2 and so on, such that TSM F7 is the dataset filtered by all seven filters. The
impact of each filter on the total number of microbial hits, species, genera, cell-containing
droplets and empty droplets with significant hits is shown in **SI Table 12** and **SI Figure 5**.

With the imposition of all filters, we arrive at ~0.4% of the unique hits found in the TSM
F0 dataset, ~3% of the species and ~5% of the genera. Like the percentage of unique
hits remaining, the percentage of cell-containing and empty droplets is ~0.7% and ~0.4%,
respectively. Because the first five filters represent decontamination and quality control
filters, we will generally report our findings on TSM F5 and TSM F7, which is the most
conservative version of the TSM dataset.

The Tabula Sapiens Microbiome atlas reveals numerous species of microbes
spanning viruses, fungi and bacteria

Binning sequences by donor and tissue type, we summarize their breakdown into high-
level taxonomic categories **(Figure 1)**. Numerous species spanning three domains of life
are present across 11 donors and 19 tissue types, with bacteria having the highest
prevalence of the three domains. We show the fraction of microbial hits in TSM F5 and
F7 datasets that have resolved taxonomic identity at each classification level and
demonstrate a high percentage, typically >95% (**SI Table 19**). Unlike bacterial
sequences, which map largely to commensal species, viral sequences mapped mostly to
the Epstein Barr Virus (EBV) - a ubiquitous and persistent human pathogen with very
limited transcriptional footprint during its latency compared to its reactivation[37,38]. As we
will discuss in more detail, this dichotomy in the transcriptional load of EBV during its
latency and reactivation likely gives rise to detectable transcripts in some donors while
appearing to be missing from others. Finally, fungal sequences are generally present with
lower relative abundances compared to bacteria. This is consistent with previous
estimates of fungi at external-facing microbiomes, depicting them significantly
outnumbered by bacteria[39–41].

Bacterial sequences are derived from a range of phyla, most notably Actinobacteria,
Firmicutes, and Proteobacteria (**Figure 1**). We find that more than a quarter of all
transcripts map bacterial ribosomal RNA (rRNA) (**SI Figure 6**). This relatively large
percentage of ribosomal sequences is expected as rRNA is the most abundant form of
RNA in bacterial cells, and its abundance is tightly coupled to the bacterial growth rate[42,43],
accounting for up to 85% of the RNA pool in rapidly growing bacterial cells in culture[44,45].
Furthermore, our finding of bacterial rRNA from diverse bacterial taxa across numerous
human tissue types (**SI Figure 7)** is in agreement with a recent study of several tumor
types, wherein 16S rRNA was detected using both sequencing and imaging[21].

The single cell RNA sequencing protocols such as the one used in this study rely on a
polyA selection. Hence, at first glance, the discovery of any microbial transcripts is
surprising. However, many families of viruses have polyadenylated transcripts[46,47].
Additionally, polyadenylation plays a ubiquitous yet understudied role in prokaryotic

transcription[48,49]. Moreover, it is well known that oligo-dT primers often bind with similar
efficiencies to A-rich regions as they do to polyadenylated tails, and thus mispriming
events inadvertently provide a window into non-polyadenylated transcripts[50,51].

Mapping the possible microbial flow routes from external-facing microbiomes
and natural environments to internal tissues and tumors

To explore the human tissue microbiome at a more granular level, we created a Sankey
plot (**Figure 2.a**), wherein each one of the microbial hits is represented by a line, the path
of which can be traced to obtain the tissue type of origin, taxonomic categorization, as
well as additional information gathered from external databases. We provide this as an
interactive plot (**SI File 1**) along with the table underlying these flow paths (**SI Table 20**).
The HMP dataset was used to identify species that had previously been observed at
external-facing microbiomes from the gastrointestinal tract, oral cavity, ear, nose, airways,
urogenital tract, and the skin. Approximately 50% of bacterial genera that make up tissue
microbiomes appear in external-facing microbiomes (**SI Figure 8**).

With the assumption that microbial flow occurs from external to internal sites, we depict
possible flow routes for numerous bacterial genera and show that various external and
internal sites are connected through their shared taxa (**Figure 2.a**, **SI Figure 9**).
Surprisingly, not all species found at internal sites originate from the gastrointestinal tract.
Other external-facing microbiomes appear to contribute many species to internal organs
and tissues. To illustrate a few microbial flow paths, we detected sequences from
Treponema denticola, an oral bacterium commonly associated with periodontal disease,
in the heart. This observation is in line with studies that have linked *Treponema denticola*
to atherosclerosis through its invasion of aortic tissue[52–54]. Sequences mapping to
Aggregatibacter aphrophilus, another oral bacterium whose infection can lead to
endocarditis, was also detected in the heart (**SI Figure 9**). While various members of the
oral microbiome have been associated with cardiovascular diseases[55], we find that the
heart is certainly not the only destination for sequences from species normally found in
the oral cavity. For example, we found sequences mapping to *Porphyromonas gingivalis*
not only in the salivary gland tissue as might be expected, but also in tissues from the
thymus, liver, vasculature, fat, bone marrow and others. Sequences mapping to
inhabitants of other external-facing microbiomes were similarly found in various tissues
(**Figure 2.a**, **SI Figure 9**). It remains to be discovered to what extent species from
external-facing microbiomes can colonize these distant sites.

In addition to the HMP dataset, we compared the number of shared genera between TSM
datasets and UHGG, which represents a much larger number of gut metagenomic
samples than found in HMP (**Figure 2.a, SI Figure 10**) and includes samples collected
from various countries, thus providing a global perspective on the shared genera. We

further explored the bacterial taxa shared between TSM and tumors (Nejman *et al.*[21]) of
the pancreas, breast, colon, lung, skin, ovaries, bone, and the brain (**Figure 2.a, SI**
Figures 11-14). Nearly half of TSM F7 bacterial genera have previously appeared in
human tumors (**SI Figure 11**). We created an abundance heatmap of these genera
across TSM tissues (**SI Figure 11**). Sequences from species that were previously
identified as highly enriched in tumors including *Paracoccus marcusii* and *Roseomonas*
mucosa are also present across various TSM F5 tissues, primarily from lymphatic organs
(**SI Figure 12**). We do not identify any of the tumor-enriched species in the TSM F7
dataset. The relatively large overlap between TSM and tumor microbiomes suggests that
tumor microbiomes may be sourced from adjacent tumor-free tissue microbiomes, even
possibly from tissues at distant sites from the tumor (**SI Figure 12-14**). We provide a list
of bacterial genera found in tumor microbiomes that are missing from the TSM F5 or F7
datasets (**SI Table 21**). Some of these genera may perhaps prove to be tumor-
microbiome specific, however future studies applying the same experimental and
bioinformatic protocols to tumor-free and tumor-derived tissues are needed.
246
We mapped the possible microbial flow routes from natural environments and external-
facing microbiomes to tissues and tumors by focusing on sequences from species found
across TSM, GEM, HMP, UHGG, and the Nejman *et al.* datasets (**SI File 2, SI Table 22**).
Many interesting flow routes appear to exist with varying degrees of complexity. As an
example of a flow route, sequences from members of the *Aggregatibacter* genus appear
to originate from the oral microbiome and the gastrointestinal tract (HMP, GEM, UHGG).
They are detected in salivary gland tissue as well as thymus, blood, liver, heart, lymph
nodes and other tissues (TSM F7). They are also found in breast tumors (Nejman *et al.*).
We provide these flow routes both as an interactive plot (**SI File 2**) and as a table (**SI**
Table 22).
257
Additionally, we used the PATRIC database[31] to obtain further information about the
pathogenicity potential of bacterial species. We found the following potential bacterial
pathogens in the TSM F7 dataset: *Porphyromonas gingivalis, Campylobacter showae,*
Aggregatibacter actinomycetemcomitans, Capnocytophaga gingivalis, Campylobacter
curvus, and *Treponema denticola*. From this list, the HMP dataset contains all but one
pathogen, namely *Campylobacter curvus*. *Campylobacter curvus* and *Porphyromonas*
gingivalis appear in human gut samples from the UHGG dataset. Finally,
Capnocytophaga gingivalis appears in the Nejman et al. dataset.
266
To explore the inter-and-intra donor tissue microbiomes at finer taxonomic resolution, we
computed the shared species within and across individuals. Whether considering the
tissues of one donor or the collective tissue microbiomes of many donors, tissue
microbiomes appear to have many overlapping species **(Figure 2.b, SI Figure 15)**.
However, when accounting for all species, species unique to each sample outnumbered

those that are shared (**Figure 2.c, SI Figures 16-17**). Furthermore, we compared
species-level bacterial microbiomes in a pairwise fashion using the Spearman correlation
for both TSM F7 and F5 datasets (**SI Figures 18-19**), revealing tissue microbiomes
across and within each individual to be uncorrelated at the species level, but increasingly
more similar at higher levels of classification. For a more detailed discussion, see **SI:**
Exploring the inter-and-intra tissue microbiomes.

Increasing the resolution of tissue microbiomes to delineate cell type
microbiomes

Many of the microbial sequences can be traced back to cells that have been annotated
by the Tabula Sapiens Consortium. Merging cell type annotation data from Tabula
Sapiens objects with cellular barcodes provides an unprecedented view of each tissue's
microbiome at the level of cell types and even individual cells. We wondered how different
cell types compare in terms of their microbial load. Thus, we quantified single-cell
microbial load as $\log_2(h_c/g_c)$, where h_c represents the number of microbial sequences
detected in cell c and is normalized by the number of host genes detected in that cell, g_c,
to account for cell to cell variation in sequencing coverage. It is important to note that this
data does not indicate whether these microbes or their sequences have been internalized
by these cells; they could be binding to the external cell membrane.

Ranked distributions of single-cell microbial loads (**Figure 3.a**), reveal that cell types with
the largest microbial load include enterocytes, T cells, and plasma B, among other cell
types, especially those from the immune compartment. The over-representation of
immune cell types is expected as these cells, with the ability to survey tissues, are the
first to encounter microbial organisms. Enterocytes are also the first line of barrier against
the trillions of microbes housed in the human intestinal tract. At first glance, the finding of
microbial sequences in pancreatic acinar cells and in the pancreas more generally, may
be the most surprising observation, however the pancreas is an organ with a physical
connection to the small intestine. This connection has been previously shown to enable
the passage of intestinal fungi to the pancreas where they can promote pancreatic
adenocarcinoma[24]. Additionally, pancreatic adenocarcinoma tumors have been shown to
harbor microbiomes that are predictive of long-term survival[22].

Network of human cell types in association with microbes reveals cell types with
greatest microbial diversity

To explore the diversity of cellular microbiomes, we built a bipartite network of cell types
stratified by tissue of origin (source nodes) and their associated microbial species (target

nodes) for both the TSM F5 and TSM F7 datasets (**Figure 3.b, SI Files 3**). Cell types in
each tissue are weighted by the number of unique microbial species, normalized by the
number of annotated cells sequenced. Hence, in this network of host-microbe
associations, hubs represent cell types with the greatest microbial diversity. Examples of
these hubs include Thymus and bone marrow CD4+ T cells, lymph node B cells and
CD8+ T cells. As with microbial load measurements, immune cell types are over-
represented in their diversity of microbial associations. Cellular microbiome network
depictions such as this open a window into host cell and microbe interactions in their
native context which would not be captured in bulk tissue sequencing experiments. An
example of an important association is between EBV and splenic plasma cells. We report
statistics on droplets with hits, and identify outlier cells based on number of hits per cell
(see **SI: Single-droplet statistics, outlier cells and outlier cell types,** and **SI Tables
23-26**). Interestingly, the most prevalent species found in outlier cells is EBV, and most
outlier cells are from immune cell types.

**Detecting the transcriptional footprints of a persistent viral pathogen and an
ancient viral relic in human cells**

EBV was the first oncovirus to be discovered and has since been shown to cause
Burkitt's, Hodgkin's, NK and T cell lymphomas as well as nasopharyngeal and gastric
carcinomas, thus accounting for ~200,000 (or 2% of) cancer deaths annually worldwide[56].
In absence of an approved EBV vaccine, currently most primary infections occur in
children and adolescents, such that by adulthood a significant fraction of the population
is latently infected[57–59]. Additionally, a recent longitudinal study of 10 million young adults
has shown a ~30 fold increase in the risk of multiple sclerosis after EBV infection, placing
EBV as one of the leading potential causes of multiple sclerosis[60].

During the latent phase of its lifecycle, EBV suppresses the transcription of most of its
genome[38] and replicates infrequently and in synchrony with the host cell while utilizing the
host polymerase[61]. In our cohort, we detected EBV sequences in five seropositive donors,
and across various tissues including the lung and the bladder. The majority of hits
originate from donors 1 and 2 (TSP1 and TSP2) and from lymphoid organs such as the
spleen, where many of the susceptible host cell types reside (**Figure 4.a, SI File 4**).
Consistent with the known aspects of EBV tropism[62], we found that the majority of
sequences originate from cell types belonging to immune and epithelial compartments.

Across distributions of EBV single-cell load in different cell types, plasma cells outrank
other cell types (**Figure 4.b**). Applying BLASTx to identify protein coding sequences, we
primarily found those expressed during the lytic phase (**Figure 4.c**): membrane
glycoprotein (BILF1), DNA synthesis and genome packaging protein (BALF3)[63], and the

viral DNA polymerase (BALF5), which is used to replicate the EBV genome exclusively
during the lytic phase[61]. Notably, we observe BZLF1 expression, an important
transcriptional switch that initiates the expression of various lytic genes[64]. Although EBV
sequences are present across a variety of cell types, only plasma cells, B cells and NK
cells are positive for lytic transcripts, consistent with the fact that these cell types are
known susceptible host cell types for EBV. Considering that the latent phase of EBV life
cycle is characterized by the expression of non-coding RNAs, many of which suppress
lytic genes[56], absence of coding lytic transcripts suggests that other cell types are
experiencing latent infections. Alternatively, they may have uptaken transcripts passively
by virtue of physical proximity to sites of lytic infections, or actively through their
involvement in the clearance of lytically infected cells.
360
In addition to EBV, we detect the presence of human endogenous retroviruses (HERVs).
Introduced numerous times during the evolutionary history of primates, HERV genomes,
which are mostly defunct, occupy 8% of the human genome[65]. The HERV K (HML-2)
family, representing the latest insertions into the human genome, is the only family of
HERVs that is transcriptionally active. Differential expression of HERV K genes has been
detected across several cancers and is thought to contribute to tumor progression[66,67].
Additionally, HERV K expression and even virus-like particles have been detected during
human blastocyst formation[68]. Although this virus was tagged as a putative contaminant
by our ML decontamination model, we believe this may represent one instance of the
model's false positives. This is because we did not detect HERV K in the ~200 negative
controls obtained for the TSM dataset. We also identified it again separately through
another set of experiments described in the next section.
373
Contrary to the claim that HERV K expression is silenced in all but a few special
contexts[69], we identified HERV K (HML-2) transcripts across five donors, spanning a
diverse range of tissues (e.g. skin, fat, spleen and the salivary glands) and cell types (**SI
Table 27**). These sequences map to the three canonical retroviral genes in the HERV K
genome, namely pol, gag and env (**SI Figure 20**). HERV K expression in human donors
beyond tissues and cell types previously investigated, opens the possibility that this
enigmatic ancient relic may be more widely active and physiologically relevant than
previously appreciated.
382

383   **Tissue microbiomes through the lenses of DNA sequencing, fluorescence, and**
384   **electron microscopy**
385
To further examine our findings through other techniques beyond single-cell
transcriptomics, we developed an assay that extracts virus-like and bacterial-like particles

(VLPs and BLPs) from human tissue (see **Methods: The Extracted Human Tissue**
Microbiome dataset). We refer to the resulting dataset as the Extracted Human Tissue
Microbiome (EHTM). We used DNA sequencing, fluorescence and electron microscopy
of VLPs and BLPs extracted from 4 tissue types (pancreas, blood, lymph node, and lung)
and 8 donors (**Figure 5.a**). In this assay, we gently dissociate tissues, eliminate cell- or
capsid-free nucleic acids, and subsequently size select for VLPs and BLPs. We used
negative controls for all experimental steps to eliminate contaminants from bulk DNA
sequencing of these extracted particles. Using nucleic acid staining followed by
fluorescence microscopy of these particles from the lung and lymph nodes (**SI Figure**
21), we observed high concentrations of VLPs and BLPs (~10^7 BLPs and ~10^8 VLPs per
cm^3 of tissue). Electron microscopy of VLPs further showed familiar virus morphologies
such as a hexagonal capsid (similar to many bacteriophages) and rod-shaped capsids
(reminiscent of the tobacco mosaic virus or the rudivirus) in the pancreas (**Figure 5.b**). It
also revealed several novel morphologies, for example one with interior nodules and
fibrous protrusions residing in a lymph node. Moreover, DNA sequencing of these
extracted particles further captured numerous sequences mapping to viruses, bacteria,
and even some fungi as some have similar sizes to bacteria (**Figure 5.c**). These results
support and are consistent with the single cell transcriptomic data, thus providing
independent evidence for several of our conclusions.
407
Interestingly, EBV and HERV K were both identified in the EHTM cohort. EBV was found
in all 8 donors, which is a higher prevalence than obtained via RNA sequencing.
Considering that single-cell RNA sequencing preferentially captures transcriptionally
active organisms, one would expect to see higher EBV donor positivity rate using DNA
sequencing of VLPs. The combination of these two methodologies could perhaps be used
to decipher active versus inactive organisms to further make a distinction between
residents and passengers of the human tissue microbiome.
415
Moreover, we identified the genera that overlap between all six datasets: TSM, HMP,
Nejman *et al.*, UHGG, GEM, and the EHTM datasets (**Figure 5.d-e**). For example, we
found sequences mapping to *Rothia mucilaginosa* in the pancreas and several other
organs using single-cell transcriptomics (TSM dataset). This species, a member of the
airway microbiome (HMP dataset), appears in pancreatic, lung and breast tumors
(Nejman *et al.*). Consistent with its traces in these three datasets, we identify this species
through DNA sequencing of the extracted BLPs in the pancreas, lung, blood, and lymph
nodes. Given these additional experiments, it is perhaps fair to assume that sequences
found in single-cell transcriptomic data are not all cell-or capsid-free. Some perhaps
originate from intact organisms that escape cellular barriers to reach internal tissues and
tumors.
427
428

Discussion

The transcriptional footprints of microbial life in human organs and cells raise several questions. For example, how could microbes escape cellular barriers established by epithelial and endothelial cells? Are there any long-term residents of human tissues in a commensal relationship with our own cells? And can these microbial signatures be used as diagnostic markers for organ health? One likely route of microbial migration is from the gut to other organs, as leaky gut epithelia together with the gut-liver axis have been proposed as a route of dissemination for various microbial metabolites from the gut into the systemic circulation[70,71]. The same route is likely to harbor microbial genomes, transcripts, and perhaps whole cells and capsids. In fact, viral migration across epithelial layers has been demonstrated *in vitro*[72]. However, as we have shown there are likely other possible flow paths from other external-facing microbiomes such as the oral cavity and the urogenital tract to various internal organs.

Overall, there is growing evidence for the existence of microbial interactions with human cells at sites in the body that are typically inaccessible to sampling. While microbial sequences have already been shown as diagnostic markers for cancer detection, our understanding of their associations with human cells is largely lacking. It is possible that causal associations exist, and that the list of oncogenic microbes is much larger than previously appreciated. Our finding of significant overlaps between tumor-free tissue microbiomes and tumor microbiomes suggests that tumor microbiomes are at least partially sourced from normal tissue microbiomes. As such, mechanism-level understanding of microbe-host interactions in human tissues could further shed light on the highly variable nature of individual responses to cancer immunotherapies[73,74] and provide novel and targeted avenues for microbiome-based immune modulation, as well as lay foundational insights into the interplay between microbes, tumor cells and the host immune cells. Moreover, given that infectious agents account for ~15% of all cancers[75], and that many pathogens can manipulate the host cell cycle[76,77] and predispose host cells to cancer[78], it is highly plausible that we have only scratched the surface of causal links between microbes and various pathologies.

With the availability of high-throughput single cell sequencing technologies today, we stand at a unique time in the history of microbiology, where the study of microbe-host interactions offers an exciting new frontier. These interactions can now be studied at the level of individual cells[79–81] not just *in vitro,* but as we have shown, in the very real context of organs - in the presence of the immune system, neighboring cells, nutrient fluxes and many other important factors that shape microbial life cycles.

Conclusion

In this study we provide the first atlas of the human tissue microbiome with cell type resolution, for which we have developed new experimental and computational pipelines such as a machine learning-based decontamination model. We discovered sequences mapping to numerous species from three domains of life. We map out the possible flow routes of microbes from external-facing microbiomes to internal tissues and tumors, and show that tumor microbiomes share many common taxa with normal tissue microbiomes. We further constructed a network of associations between human cell types and microbes, shedding light on the types of organisms that cells from different tissues encounter either by way of immune clearance or physical proximity to microbial niches and sites of infections. This type of analysis can reveal the interactions between various cell types and microbial organisms. In cases of intra-cellular infectious events such as EBV, host and pathogen transcriptomic landscapes may reveal novel pathways that underlie cellular susceptibility and resistance, which would have great clinical application.

Materials and Methods

The Tabula Sapiens dataset

The Tabula Sapiens raw fastqs containing single-cell RNA sequencing data from Smart-seq2 and 10X droplet-based experimental pipelines were acquired from the Tabula Sapiens consortium[28] (https://github.com/czbiohub/tabula-sapiens). This included raw fastqs for 12 donors 1,2,4,6,7,8,9,10,11,12,14, and 16 as well as objects which contained cell type annotations for ~0.5 million cells (donor 16 cell type annotations were not included). All analysis was performed on the 10X data, which represents the vast majority of this dataset. Donor 4 tissues (mammary, myometrium, uterus) were eliminated in the TSM F7 version of the dataset due to not meeting the criteria imposed. Additionally, because we could not acquire negative control samples for tongue tissue, we do not report on this tissue. Thus we have a total of 11 donors and 19 tissues in the TSM F7 dataset.

The Extracted Human Tissue Microbiome dataset

Tissue Digestion and DNA extraction. Tissues were acquired from Donor Network West (delivered on ice) and cut into 1 cm blocks using UV and ethanol sterilized scalpels and petri dishes. Tissues were incubated for 90 minutes with collagenase IV (1 mg/mL, Worthington) at 37°C and further disrupted by pipetting. Disrupted tissue samples were then incubated for 30 minutes at 37°C with 19U/mL DNase I and 1mg/mL RNase A

(ThermoFisher) - added to eliminate as much of the cell- or capsid-free DNA and RNA
that is released during tissue digestion. The RNAase and DNAase enzymes were
deactivated using RNAsecure at 1X and EDTA at 5mM for 10 minutes at 65℃. Each
sample was then filtered through a sterile 40μm cell strainer, to eliminate the cellular
debris. The filtrate was passed through a 5μm pore size syringe filter to enrich particles
in the size range of most microbes and to eliminate host cells. To separate out as many
virus-like particles from this mixture of filtrate, we imposed another filter, this time using a
0.8μm pore size syringe filter. The filter was inverted and objects larger than 0.8μm were
collected. Although this was a protocol developed in-house, we were able to get key
insight from several sources: Qiagen TissueLyser Handbook, Worthington Tissue
Dissociation Guide and viral metagenomic studies[82]. DNA extraction was then performed
using QIAmp DNA mini kit (Qiagen) according to kit instructions. Moreover, we used
multiple negative controls for each step of the experimental workflow, including tissue
digestion, DNA extraction, and library preparation.

Sequencing and data analysis. We used SIMBA on this dataset, excluding steps used
for UMI extraction. We used the negative control samples to create a blacklist of
contaminants, which were subsequently removed. Despite significant methodological
differences between the EHTM dataset, and the TSM single-cell transcriptomic dataset,
we found members from the Actinomyces, Neisseria, Rothia, Corynebacterium,
Nesterenkonia, Streptococcus, Veillonella genera shared between the two datasets.
However, it is important to note that the level of overlap between these two datasets and
any others depends on parameters such as sampling depth and method of detection.

TEM. We plasma treated EM grids (300 mesh Carbon/Formvar coated Copper grids)
using Leica EM ACE600 to allow each sample to bind to the grid. We applied 5μL of
1:1000 diluted sample onto each EM grid and allowed the sample to bind for 3 minutes.
The grid was then exposed to 1% uranyl acetate for one minute, and subsequently dried
using a filter paper. Samples were imaged using JEOL JEM-1400.

Fluorescence imaging. Following tissue digestion and microbial extraction, filterates
were stained with SYBR™ Gold Nucleic Acid Gel Stain (ThermoFisher) at 10X for 10
minutes at room temperature and imaged using Leica CTR6000. Bright dots were
counted manually across different sections across a hemocytometer grid and average
counts were projected to estimate VLP and BLP concentrations per cubic centimeter of
tissue.

Acknowledgements

We thank Rob Phillips for insightful comments on the manuscript. We also thank Leila Sidow, Robert Jones, Roja Kolluru, Ed Marti, Elizabeth Jerrison, Mark Kowarsky, Alina Isakova, Stephanie Tzouanas Schmidt, Yuan Xue, Mira Moufarrej, Michael Swift, Ivana Cvijovic and all the other members of the Quake lab, along with the Tabula Sapiens and the Boundaries of Life consortia for their help and suggestions towards experimental and computational pipelines. Additionally, we thank the following Tabula Sapiens consortium members for their help towards tissue digestion controls: Mallory Ann Laboulange, William Kong, Siva Vijayakumar, Hua Dong, Rebeca Arroyo, Hayley Raquer, Michael Swift, Zhen Qi, Will Goodyer, Gabriel Loeb, Simon Bucher, Helen Zue Zhang, Soochi Kim, Yan Hang, Juan Irwin, Emily Trimm and Irene Fan. We are grateful to Donor Network West personnel including Ahmad Salehi and Ravi Ponnusamy, as well as the Cell Sciences Imaging Facility at Stanford University. We are especially indebted to donors and their families for their incredible contribution to science. This work was supported by the Chan Zuckerberg Biohub, the Gordon and Betty Moore Foundation and the John Templeton Foundation.

Funding

This work was supported by the Chan Zuckerberg Biohub, the Gordon and Betty Moore Foundation and the John Templeton Foundation.

Author contributions

Conceptualization: GM, SRQ
Methodology: GM, SC, SRQ
Investigation: GM, SRQ
Data Curation: GM
Formal Analysis: GM
Visualization: GM
Software: GM
Funding acquisition: GM, Tabula Sapiens Consortium, SRQ
Resources: GM, Tabula Sapiens Consortium, SRQ
Project administration: GM, Tabula Sapiens Consortium, SRQ
Writing – original draft: GM
Writing – review & editing: GM, SC, SRQ

Competing interests

570   Authors declare that they have no competing interests.
571

Data and materials availability

573   All   data   and   code   is   available   through   our   GitHub   repository:
574   https://github.com/gitamahm/human_tissue_microbiome_atlas. This repository includes
575   SIMBA, which is written in Snakemake[32], Post-SIMBA data analysis and visualization
576   organized into separate Jupyter Notebooks (written in python v3.6.10). The Jupyter
577   notebooks are listed below.
578
-        p01_databaseCreationSC_microbeDB_fungalDB.ipynb
-        p02_creating_truth_datasets_for_testing_simba.ipynb
-        p03_testing_truth_datasets.ipynb
-        p04_experimental_negative_controls.ipynb
-        p05_post_SIMBA_processing_alldonors_and_EHTM_gh.ipynb
-        p06_cell_ranger_counts_gh.ipynb
-        p07_machine_learning_modeling_contamination_and_other_filters.ipynb
-        p08_habitat_assignment_using_GEM_HMP_UHGG.ipynb
-        p09_davinci_language_model_habitat_of_origin_gh.ipynb
-        p10_impact_of_filters.ipynb
-        p11_single_cell_microbiome_stats.ipynb
-        p12_tabula_sapiens_multibiome_analysis.ipynb
-        p13_ts_intersection_with_external_datasets.ipynb
592

Figure

Figure 1. Tabula Sapiens Microbiome (TSM) atlas derived from the Tabula Sapiens single-cell transcriptomic dataset consisting of 19 tissue types from 11 donors (identified

by TSP#), 100+ cell types and 400,000+ annotated cells. Tissue microbiomes are
stratified based on taxonomy: domains (inner ring) followed by eukaryotic phyla, and
bacterial phyla, respectively. Sequences are grouped based on donor (one donor is
shown per tissue). Most tissues shown are from TSP2, except for Thymus, Liver, Salivary
gland which are from TSP14, Pancreas from TSP9, Heart from TSP12, and Skin and Fat
from TSP10. Tissues shown have at least 10 hits and are from the TSM F7 dataset.

Figure 2. Microbial flow within the body from external and internal environments (TSM F7 dataset). a) Sankey plot revealing the taxonomic classification and tissue of origin for each sequence (each line in the plot represents an individual sequence) along with additional data from the Human Microbiome Project (HMP) database, tumor microbiome dataset by Nejman *et al.*, the Unified Human Gastrointestinal Genome (UHGG) collection and the Genomes from Earth's Microbiomes (GEM). Sequences belonging to species that are not found in these additional databases, including all viral and fungal species, are labeled as "NA". Under the GEM column, "h" denotes human and "nh" denotes non-human. Only some of the largest groups in each column are denoted in the legend. The rest can be found as part of the interactive plots provided in the **SI File 1**. b) Circos plot of shared species within TSP2 (donor 2). Each link represents a unique species. Species that are not shared between tissues are not shown. c) UpSet plot for TSP2 demonstrating the number of shared species between tissues, as well as the number of species unique to each tissue. Unique sets of species are shown as filled circles. The intersection of two or more sets are shown via connected filled circles. Bars denote the size of each set or the intersection of sets.Tissues are color-coded according to panel b. bm: bone marrow, li: large intestine, si: small intestine.

a.

Figure 3. Cellular microbiomes across all donors. a) Distributions of hits per cell against
three domains of life, grouped by cell type from TSM F5. Cell types are ordered based on
the maximum value found in any of the three distributions. Y-axis corresponds to
$\log_2(h_c/g_c)$, where h_c is the number of hits per cell, normalized by the number of g_c, number
of detected genes per cell. Only cells with detected hits are included in the analysis and
only cell types with at least 10 cells. b) Network of human cell types (grouped by tissue
of origin) in association with microbes. Cell types are source nodes (circular), and

microbial species are the target nodes (square). Source nodes are weighted by $S_i/\log_2 C_i$,

where S_i represents the number of unique species associated with source node *i*, and is

normalized by the total number of cells, C, sequenced for source node *i*. The

normalization ensures that cell types with greater representation in the dataset do not

artificially appear to have more diverse microbial associations. Moreover, edges in this

network are weighted by $N_{j,i}$, the counts of sequences mapping to microbial species *j*

associated with source node *i*. For visual clarity, only larger source nodes are labeled: 1)

Thymus CD4+ T cell, 2) bone marrow CD4+ T cell, 3) thymus fibroblast, 4) lymph node

effector memory T cell, 5) lung regulatory T cell, 6) bladder intermediate epitheliocyte, 7)

pancreatic acinar cell, 8) lymph node B cell, 9) pancreas fast muscle cell, 10) bladder

macrophage. For all other nodes, please see the accompanying interactive plot (SI File

3). bm: bone marrow, li: large intestine, si: small intestine.

Figure 4. Detection of EBV across tissues and cell types (TSM F7). a) Sankey plot of
EBV sequences, demonstrating their donor, tissue, cell type, and compartment of origin.
For visual clarity, only hits from annotated cells are included (TSP1 and TSP2 hits remain)
and only large groups have been labeled for tissue and cell type rows. Please see the
accompanying interactive plot (SI File 4). b) EBV hit per cell distributions grouped by cell
type from any tissue that is EBV hit positive. Y axis denotes $\log_2(h_c/g_c)$, where h_c is the
number of hits per cell, normalized by the number of g_c, number of detected genes per
cell. Only cell types with at least 5 cells are shown. c) heatmap of protein-coding
sequences identified in TSP2 mapping to EBV, grouped by cell type. The scale bar

denotes $\log_2(h_{ct} + 1)$, where h_{ct} represents the number of hits per cell type. The functional

role of some of the EBV protein coding sequences are denoted. Note, "HP" corresponds

to "hypothetical protein".

Figure 5. Tissue microbiomes from 8 donors using DNA sequencing, fluorescence and EM imaging from the Extracted Human Tissue Microbiome dataset along with a comparison to several other datasets. **a)** study design and experimental workflow. **b)** EM images of viral fractions retrieved from the pancreas, lung and lymph nodes showing both known and novel virus-like morphologies. **c)** hits per tissue and donor stratified based on taxonomy: domains (inner ring) and phyla. **d**) Exploring the intersection of bacterial genera across various datasets using an UpSet plot: Tabula Sapiens Microbiome (TSM) F5 and F7, Extracted Human Tissue Microbiome (EHTM), Human Microbiome Project (HMP), Nejman *et al.* tumor microbiome, Unified Human Gastrointestinal Genomes (UHGG) collection, and Genomes from Earth's Microbiomes (GEM). Only sets with a minimum size of 7 are shown for visual clarity. **e**) The genera that are at the intersection of all these datasets are shown. Presence (light boxes) and absence (dark boxes) of each genus is shown. bm: bone marrow, li: large intestine, si: small intestine, panc: pancreas, GBM: glioblastoma multiforme.

SI Text

SIMBA, a computational pipeline for the identification of bacteria, viruses and fungi in single cells

With the rapidly rising number of single-cell transcriptomic datasets, there is a growing interest in detecting microbial transcripts within these datasets. Although there are many existing tools for microbial profiling in standard metagenomic datasets, tools that take single-cell transcriptomic data as input are currently more limited[83]. There are however a growing number of pipelines being developed for this purpose, including those developed by Robinson et al. (CSI-Microbes)[84], Ghaddar et al. (SAHMI)[23], and Llorens-Rico et al.[85]. CSI-Microbes supports three different existing tools for detection of microbial sequences, namely GATK PathSeq[86], CAMMiQ[87] and SRPRISM[88]. GATK PathSeq relies on BWA-MEM alignments, replacing its original use of BLAST[89] in favor of faster alignment speed. Llorens-Rico's pipeline uses Kallisto pseudo-alignments against a reference database of ~12K microbial reference genomes. SAHMI implements kmer-based aligners against a microbial and human reference dataset and implements additional filters to detect false positives.

In our pipeline, called **Si**ngle-cell **M**ultibiome **B**LAST **A**nalysis (SIMBA), we first use UMI-tools[33] to filter and extract cell barcodes and UMIs from raw fastqs using the *extract* and

whitelist commands. We used STAR[34] (v2.7) to align resulting sequences against a
reference library containing the human genome (GRCh38.p13) and ERCCs. We then took
unmapped sequences for further processing. We wrote *poly.py* to eliminate low-
complexity sequences with high fractions of consecutive bases, often representing primer
dimers and other sequences with long stretches of polyA. We used BLASTn[35] (v2.10) to
align against several custom databases followed by the nt database, which is one of
largest publicly available databases, consisting of ~70 Million sequences (11/02/2020)
pooled from GenBank, EMBL, DDBJ, PDB, and RefSeq databases. Taxonomic lineages
for each top hit against the nt database were added using TaxonKit[90]. Top hits were
filtered based on their taxonomic assignment to include only viral, bacterial and fungal
species.
710
The premise behind creating a pipeline that relies on BLAST given its high computational
cost, is based on our early observations that some of the faster tools for microbial profiling
can result in an unacceptably high false positive rate. As an example of a kmer-based
profiling method, which are among the most popular and widely used methods for
microbial profiling, we explore instances of false positives from Kraken2 against the nt
database using several truth datasets. We demonstrate why in the context of our dataset,
where the vast majority of sequences are human-derived, such tools lack the necessary
precision (see **SI: Benchmarking Kraken2 precision and recall**). Robinson et al.
similarly employ GATK PathSeq rather than faster kmer-based profiling methods[84,87]. We
further show that while Kraken2's recall is high when the input sequences are sampled
from the Bacterial Refseq dataset, it is highly variable and can be much lower when the
input sequences are sampled from other datasets. This is one example illustrating why
alignments using BLAST against large, comprehensive reference databases such as nt
are still widely considered the gold standard in precision against which new tools are often
benchmarked[91–94]. Because the study of microbiomes in low-biomass settings presents a
high burden of proof with many potential sources of false positives, in our approach we
favor precision over computational time and recall. In these settings, we and others argue
that the problem of bioinformatic false positives is critical[89,92,95,96] and on par with the
problem of experimental false positives, or contamination.
730
However, aligning large datasets against large databases using BLASTn is
computationally infeasible even with access to large compute clusters. To lower this
computational cost, we created a lean human-microbiome centric database for bacteria
and fungi which we call microbDB (see **SI: Creating custom databases to limit the**
computational burden of BLASTn against the nt database). We use microbeDB as a
first-pass filter to select sequences with the highest likelihood of being bacterial or fungal
in origin. Additionally, through subsampling, we created leaner versions of the publicly
available bacterial and fungal RefSeq databases[97] to use as second-pass filters. Viral

sequences were aligned against the Viral RefSeq database[97] (release 98) which included
9,000+ viral genomes. Surviving sequences that produced significant alignments or "hits"
(E-value < 10-5) against these intermediate databases were then aligned against the nt
database. This particular E-value threshold is commonly used[98–101]. For each sequence
the top significant hit (lowest E-value) was selected for downstream analysis. We wrote
dedup.py to create input fasta files with sequences that produced a top hit in this step,
and use the resulting file to BLASTn against the nt database. By taking this type of
screening strategy, we lower the computational burden of alignments by ~100 fold.
Without the initial screening, we would require $\sim 3 \times 10^8$ CPU hours, equivalent to ~$9
million using Amazon's compute instances (estimate made in 2020).
749
The resulting top hits that emerged from the viral, bacterial and fungal workflows against
nt were combined into one dataset consisting of ~19 million sequences for further
processing and quality control. Despite first filtering human reads using the STAR
alignment step, out of ~19 million sequences that generated significant hits against nt,
only 12% mapped to the viral, bacterial, and fungal biomes of interest, while half of them
mapped back to the human genome. This indicates the importance of implementing
multiple checkpoints to eliminate non-microbial reads, and cautions against relying solely
on smaller databases without validation against a comprehensive database such as nt.
This is especially relevant to human tissue samples where the ratio of non-microbial to
microbial sequences is high[86,89,95].
760
We applied our pipeline not only to the raw data underlying Tabula Sapiens annotated
single cell dataset but all the raw data, including those from empty droplets, which we
refer to as the Tabula Sapiens Microbiome F00 (TSM F00) dataset. The rationale for
including unannotated or empty droplets is that we suspect that microbial reads from
extracellular environments could appear in those droplets, and more generally, different
species may have different patterns of association with these two categories of droplets.
Later we incorporate this information for each species as a parameter in our machine
learning models for decontamination. Using TSM F00 as input, we analyzed reads from
150+ Million droplets, ~1.3 Million of which had more than 200 human genes and that we
treat as potentially cell-containing. By matching the droplet barcodes (along with sample
name) to those found in the Tabula Sapiens objects, we were able to acquire the cell type
annotations for a portion of the cell-containing droplets. For all other droplets, we provide
tissue and donor information. Please see **Data and Materials Availability** for the Jupyter
notebook where details of cell-containing and empty droplet counts using cell ranger can
be found.
776
The output of SIMBA once filtered to contain only bacterial, fungal or viral hits was 2.4
Million significant hits. There were ~724K significant unique hits (counted based on

Unique Molecular Identifiers or UMIs) arising from ~0.5 Million droplets - 79K cell-
containing droplets (~6% of TSM F00 cell-containing droplets) and ~380K empty droplets
(~0.3% of TSM F00 empty droplets). This dataset is what we will refer to as the TSM F0
dataset.
783

784 **Creating custom databases to limit the computational burden of BLASTn against**
785 **the nt database**

786
RefSeq databases[97] for bacterial and fungal genomes are available through NCBI,
however they are typically large databases that would create significant computational
bottlenecks. We created several custom databases to serve as lean human-microbiome
centric databases for the initial iteration of BLASTn search for bacterial and fungal
sequences. This was not done for viruses as they have much smaller genomes relative
to bacteria and fungi and thus the viral RefSeq could be used directly.
793
MicrobeDB was constructed by combining two other custom databases, bacteriaDB and
mycoDB. BacteriaDB was designed by taking the intersection of the HumGut database[102],
a comprehensive database of ~30,000 prokaryotic genomes mined from ~5,700 healthy
human metagenomes, and the bacterial RefSeq database (downloaded on 01/2021). The
HumGut dataset was obtained from https://arken.nmbu.no/~larssn/humgut/. We therefore
identified the unique set of species in HumGut and eliminated entries with unidentified
species-level taxonomy. Similarly, we identified the unique set of bacterial RefSeq entries,
and took the intersection of these two sets to arrive at several hundred bacterial species.
We used a similar strategy to bias the initial search for fungi towards species previously
identified in human mycobiome studies. We took five representative species from each
genus found in both the fungal RefSeq (downloaded on 01/2021) and a recent survey of
the healthy human mycobiome[103] to create mycoDB. While microbeDB was used for the
first search iteration, larger databases were used for the second iteration. These larger
databases, slimBacterial_RefSeq and slimFungal_RefSeq are the reduced versions of
the bacterial and fungal RefSeq databases which we subsampled to include
representative species and strains. Please see **Data and Materials Availability** for the
Jupyter notebook where details of how intermediate databases were constructed is
described.

Benchmarking SIMBA's precision and recall

We evaluated SIMBA's precision and recall against several truth datasets. For all truth
datasets, we randomly selected sequences that were 100 bases long to match the length
of sequences in the TS dataset. We will refer to each sequence from truth datasets as

"input seq" and its corresponding output hit from SIMBA as "output seq". If an input seq
goes undetected by way of not producing any corresponding output seq, we count that
as a false negative. We use taxonomic IDs on input and output seqs to measure precision
and recall at each level of taxonomic classification. For example, if an output seq does
not match an input seq in taxonomic ID or species name, we consider that as a false
positive at the species level of classification.
822
First, we created a dataset of 2400 sequences derived from human chromosomes
(GCF_000001405.39_GRCh38 assembly) including X and Y, with 100 randomly selected
sequences from each chromosome. We then employed SIMBA's bacterial and fungal
branch of the pipeline without using STAR to eliminate human-derived sequences as we
knew most of these input seqs would already be eliminated at this step. Even without this
step, the pipeline did not produce any false positives. Only one input seq produced a hit
when aligned against microbeDB and that sequence was eliminated in the subsequent
alignment step against the next larger intermediate database, thus no false positives were
detected.
832
We also used this truth dataset against the viral branch of the pipeline, again forgoing the
STAR alignment step. Out of 2400 sequences, 65 produced hits against the viral Refseq
database, and were subsequently aligned against the nt database. At this stage, all these
sequences were correctly identified as human in origin, including one human retroviral
sequence. Thus, SIMBA produced no false positives when encountering human-derived
sequences, which is an important aspect of this pipeline considering the high ratio of
human to non-human sequences in the TS dataset and in human tissues more broadly.
840
Next we tested SIMBA's performance when encountering viral sequences. To create a
truth dataset, we randomly selected 100 viral genomes and selected 10 random
sequences from each genome using the viral RefSeq database. This resulted in
sequences from 62 genera of viruses, thus representing a diverse viral truth dataset. We
found that all input seqs produced an output seq. Interestingly we noticed that some
mismatches that were encountered between the taxonomic IDs of input and output seqs
were the result of finding a prophage output seq when the input seq originated from a
phage. Because the prophage is embedded in a bacterial genome, the resulting output
seq had the taxonomic ID of the host bacterium. We excluded these special cases from
calculations of precision and recall. At the species level, we see the following precision
and recall, respectively: 0.8, 1 **(SI Table 1)**. False positives detected at the species level
correspond to sequences from highly similar viral species that typically share the same
host **(SI Table 2)**. This is expected as short sequences (e.g. ~100 bp) can be shared
identically or with very high similarity among related species. Indeed, we found that all
false positives had 100% sequence similarity across the entire length of the input seq **(SI**

Table 3). Hence, genus level classification provides a more meaningful depiction of precision as it is less prone to this type of ambiguity in the mapping of short reads. Indeed, the precision jumps to almost 1 when examined at the genus level.

Moreover, the high recall associated with our viral search is expected given that we did not use intermediate, subsampled databases to lower the computational cost for the viral search since it does not pose a computational bottleneck. The lower computational cost of searching for viruses is a result of viruses having 1) some of the smallest genomes across all domains of life, and 2) fewer representatives than bacteria in many databases, including the nt database.

Next, we characterized false positives as well as other parameters in the face of bacterial input seqs. Given that environmental bacteria can be a source of contamination in kits and reagents, we aimed to bias SIMBA to finding those bacteria that are more likely to reside in the human microbiome to 1) mask potential contaminants and 2) limit the initial search space to reduce the pipeline's computational time. To introduce this bias, we created microbeDB which serves as a first pass filter on the input data. This database is composed of bacteria and fungi commonly found in the human gut microbiome, based on the assumption that the gut likely contributes the most to human tissues due to its greater microbial biomass compared to any other site in the human body. Therefore, to evaluate the pipeline's performance, our first truth dataset contained bacteria cultured from the human gut microbiome.

We obtained taxonomic IDs from the Human Gastrointestinal Bacteria Culture Collection (HBC) - to select representative genomes from 50 random bacterial species cultured from the human gastrointestinal tract, for which we downloaded a representative genome from NCBI. We obtained the Human Gastrointestinal Bacteria Culture Collection (HBC) dataset from https://www.ebi.ac.uk/ena/browser/view/PRJEB23845?show=reads (filereport_read_run_PRJEB23845_tsv.txt). We then selected 100 random contigs, for a total of 5000 sequences. Using this truth dataset, we report the corresponding precision and recall **(SI Table 4)**. At the species level, we arrive at the following precision and recall, respectively: 0.69, 0.22. As with the viral truth dataset, when we analyzed the false positives, we found that the vast majority represent sequences from highly related bacterial species **(SI Table 5)** with nearly identical or 100% identical output seqs to input seqs. We again see a jump in precision to 0.93 going from the species level of classification to the genus level. We have a recall of 0.28 at the genus level, and thus we likely underestimate the true number of bacterial species using this pipeline. However, we wanted to limit instances of false positives even at the expense of false negatives. In future iterations of this pipeline, increasing the size and diversity of microbeDB would likely enhance the pipeline's recall.

896
To test how well SIMBA identifies members of another microbiome, we created a truth
dataset based on the expanded Human Oral Microbiome Database (HOMD), consisting
of 50 randomly selected genomes and 100 randomly selected segments from each. The
expanded Human Oral Microbiome Dataset (HOMD) was obtained from
https://www.homd.org/download#genome, specifically from HOMD_taxon_table2022-09-
15_1663245201.txt and HOMD_genome_table2022-09-15_1663245202.txt files.
Unexpectedly we found similar values for precision and recall across different
classification categories **(SI Table 6)**. This stands in contrast to the metrics we obtained
when we used a truth dataset that was not focused on bacteria from the human
microbiome.
907
We randomly selected 50 bacterial genomes from the bacterial Refseq database, and
randomly selected 100 sequences from each genome for a total of 5000 sequences.
Comparing the performance results of this truth dataset at the genus level to the gut or
oral bacteria truth datasets, we see that while the precision remains high, recall has
dropped down to only 0.05 **(SI Table 7)**. Thus, the pipeline preferentially recalls bacteria
from the human microbiome compared to any randomly selected bacterial sequence. It
can mask potential contaminant bacterial sequences. Even those input seqs that produce
output seqs have a high chance of being correctly identified due to the pipeline's high
precision afforded by the last BLASTn step against the nt database.
917
918 We created a fungal truth dataset by taking ~5000 sequences randomly sampled from a
919 study of the gut mycobiome[104]. At the genus level, we found similar values of recall and
920 precision to those we found for the bacterial truth datasets created from HOMD and
921 HBC (**SI Table 8**). Please see **Data and Materials Availability** for the Jupyter notebook
922 where our integration of datasets for benchmarking SIMBA is described in detail.
923
Benchmarking Kraken2 precision and recall
We tested the precision and recall of a popular tool in the field of microbial metagenomics,
Kraken2, both against the microbial truth datasets that we created to test SIMBA as well
as a sample of the Tabula Sapiens dataset. We chose Kraken2[105] as it is one of the fastest
and most accurate metagenomic taxonomic profiling tools available, and in general Kmer-
based approaches to fast taxonomic profiling represents a good fraction of the tools that
are available[91].
931
This was done primarily to see whether Kraken2's speed performance would justify its
use for our dataset. We assembled the nt database to use Kraken2 against. We found
that, in agreement with a previous benchmarking study[91] of Kraken2 and related tools,
Kraken2 has very high precision and recall against the bacterial RefSeq truth dataset **(SI**

Table 9). However, its recall is quite lower for the other bacterial truth two datasets,
starting from 0.53 at the genus level against the truth dataset created from HBC **(SI Table**
9).

Upon removing human-derived sequences using STAR for a bladder sample from the
Tabula Sapiens dataset, we used Kraken2 against the nt dataset. Kraken2 created
outputs marked classified and unclassified. More than 60% of reads remained
unclassified at the species level, and nearly half of the reads still remained unclassified
at the domain level **(SI Figure 2a)**. Using the provided taxonomic IDs, we obtained
taxonomic lineage information for reads that were classified.

Among the classified reads, ~285K were identified as eukaryotic, ~2.7M as bacterial and
~1.1M as viral **(SI Figure 2b)**. To test the precision and recall of these classification calls,
we subsampled each domain (except Archaea) and randomly selected 100 sequences
from those reads that had acquired full classification up to the species level. We then
used BLASTn against the nt database to measure the extent of the agreement between
Kraken2 and BLASTn taxonomic classifications. Surprisingly, we found very little
agreement and a high rate of false positives associated with Kraken2 classifications. For
bacteria and viruses, the mean precisions at the genus level are 0.06 and 0.04,
respectively **(SI Table 10)**.

Taking a closer look at instances of false positives for reads that Kraken2 classifies as
various species of bacteria, we see that many are classified as human (and other primate
species) **(SI Table 11)**. Human reads that are not sufficiently eliminated by STAR present
a major challenge for Kraken2. This issue of high false positive rate when analyzing
clinical samples has been discussed in the past[89,91]. Given a large enough number of
human reads, as is the case with Tabula Sapiens datasets, many sequences will be
misclassified as microbial or viral using this tool. On the contrary, because our pipeline
relies on BLASTn against nt, it has significantly better precision than kmer-based
approaches. Hence, we could not justify the faster speeds associated with some of the
existing metagenomic classification tools over their lack of extreme precision for this
specific application.

Measures against contamination during tissue recovery and transport

As detailed by the Tabula Sapiens consortirum[28], the tissues were procured at various
hospitals through collaboration with Donor Network West (DNW, CA) - a not-for-profit
organization that is a federally mandated organ procurement organization. Tissues were
procured consistently across all donors across different operating rooms (OR).
Contaminants from air, hair, skin, and instruments were carefully mitigated using strict

aseptic techniques as outlined by the American Association of Tissue Banks (AATB) some of which is briefly discussed in the following paragraphs.

In each OR, multiple sterile fields are created. A sterile field is created for each time a pair of sterile gloves or gowns is opened. An additional sterile field is set using sterile surgical drapes around a surgical site. Several sterile fields are created on tables where sterile surgical instruments are placed. Another sterile field is created on a table where storage and shipping supplies are placed.

To mitigate the risk of contamination from air, the OR rooms used to recover organs contain elaborate air filtration systems, and OR personnel are required to wear sterile PPE such as a face mask at all times. They also wear hair bonnets prior to putting on surgical scrubs and maintain them at all times. The OR personnel scrub their hands and arms before a procedure; using sterile surgical attire as barriers between their skin and recovery environment, they then prepare the donor for tissue recovery.

The donor is prepped for tissue recovery using rigorous aseptic protocols as outlined by AATB (cite). First, clothing is removed, and hair is removed from any recovery areas using surgical scrub soap to prevent contamination from the donor's skin. These areas are then wiped with 70% Isopropyl Alcohol to disinfect the skin while also removing any loose hair or debris not removed by the surgical soap. FDA-tested and approved surgical soaps are used. Donor draping is performed using aseptic standards of practice to isolate, form a barrier and create a sterile field. Recovered organs are protected by a triple sterile barrier and stored in enough cold, sterile preservation solution (UW solution) to sufficiently cover the organ or vessel. To capture any contamination possibly introduced through the UW solution, we performed sequencing of this solution.

It is important to note that given our study relies on post-mortem samples, thus there is a possibility of tissue barrier breakdown that would impact microbiomes. To partly address this concern, we set up a private courier service to keep the time between organ procurement and initial tissue preparation to less than one hour. This is a relatively short period considering that heart and lung procured for transplantation are viable for 4-6 hours, liver for 8-12 hours, and kidneys for 24-36 hours. As tissues were recovered in the same manner as those used for transplantation, and were rapidly prepared for single cell sequencing, the possibility of gross tissue breakdown is low. Additionally, histological staining of tissues showed viability along with certain markers used to identify rapidly dividing cells (e.g. transient amplifying cells of the intestinal epithelium)[28]. Using single-cell transcriptomes, we could differentiate between hundreds of cell types, identify differentiation trajectories, and tease apart different phases of the cell cycle[28]. Together these findings point to cellular and tissue viability at large.

Identification of contaminants based on negative controls and external datasets (Filters 1-3)

The introduction of contaminants was mitigated during tissue procurement in the OR setting by creating sterile fields and various other strict measures as these tissues were collected in the same manner as those recovered for organ transplantation. These tissues were placed in UW solution and then within sterile bags on ice and quickly shipped to different tissue expert labs to preserve tissue and cellular viability. Within different labs, organs were further dissected into smaller tissue pieces and digested to create single-cell suspensions in preparation for single-cell sequencing. A detailed description of tissue procurement and transport is provided in **SI: Measures against contamination during tissue recovery and transport.**

There are multiple potential sources of contaminants, including contamination introduced during tissue procurement, tissue transport, tissue digestion and sequencing. However, we suspect that different sources of contamination do not have equal contribution to the total list of contaminants in each sample. We posit that for dense tissue types, contaminants from instruments, reagents, or air are exposed primarily to the surface of each tissue sample, and thus surface area to volume ratio, SA:V, is an important physical constraint on each contamination source. We will illustrate the importance of SA:V using simple back-of-the envelope estimates, assuming samples are cubes for simplicity. For a cube, SA:V is $(6 \times s^2)/(s^3)$ or $6/s$, where s is the length of a side. Let us assume the organ recovered in the OR is 10 cm on each side, and that it is further dissected into 1 cm cubes in a lab and digested into 10 um cubes (or cells). In this process, the SA/V ratio starts at 0.6 cm^-1 in the OR setting. Then in the lab setting, it goes up to 6 cm^-1 at the dissection step and then to 6000 cm^-1 during the tissue digestion and single-cell suspension step. Because of the increase in SA:V, we estimate that contaminants are orders of magnitude more likely to come into contact at the tissue digestion and sequencing steps than they are at tissue procurement, transport or dissection steps.

All labs performed tissue digestion and sequencing using aseptic techniques, however because these samples were largely intended for studying the human transcriptome, no blank instrument or reagent controls were generated at the time of the original experiments. In absence of these controls, we partnered with the labs that had processed these tissues and obtained reagents and consumables from the following tissue experts for the Tabula Sapiens project: bladder, blood, bone marrow, fat, heart, intestines, kidney, liver, lung, lymph node, muscle, pancreas, salivary gland, skin, spleen, thymus, trachea, uterus, and vasculature. The reagents and consumables are listed by tissue in **SI Table 12**. The tissue dissociation protocols were performed exactly as detailed by the Tabula Sapiens consortium[28] in small scale (~250 µl final eluate) but in the absence of tissue.

1054   These protocols included incubation steps, centrifugation, sample filtration using cell
1055   strainers, enrichment using MACS columns, among other steps. Even when the same
1056   reagent was used across different labs, we obtained each lab's specific batch of that
1057   reagent to account for lab-specific contaminants.

1058

Equal volumes of supernatants obtained at the end of tissue dissociation protocols were
pooled and mRNA seq libraries were generated using the Chromium Next GEM Single
Cell 3' v3.1 chemistry (10X Genomics, Pleasanton, CA). Each pool was loaded onto a
Chromium G Chip channel for a total of 5 channels. Libraries were prepared using
manufacturers recommendations. All PCR reactions were performed using the Biorad
C1000 Touch Thermal cycler with 96-Deep Well Reaction Module. As per 10X Genomics
protocol recommendation, amplified cDNA and libraries were evaluated on a 2100
Bioanalyzer system (Agilent Technologies, Santa Clara, CA). Libraries were sequenced
on the NextSeq 500 (Illumina, San Diego, CA) to obtain a total of 86 million reads.

1068

With ~200 negative control samples, we arrived at a list of contaminant species. When
we excluded all species found in negative control samples, the number of unique hits was
reduced to ~63K, or roughly 9% of the TSM F0 unique hits. Interestingly, the number of
species dropped from ~8200 to only ~7100 and the number of genera went down from
~2500 to ~2400. Filter 1 reduced the numbers of cell-containing and empty droplets to
~13K (~17% of TSM F0 cell-containing droplets) and ~44K (~12% of TSM F0 empty
droplets), respectively.

1076

1077   We then imposed a second filter based on genera commonly found in negative controls
1078   from previous studies[36], which we will refer to as the Salter genera. The removal of these
1079   genera reduced the number of unique hits, species and genera to ~52K, ~6200, and
1080   ~2300, respectively. It reduced the numbers of cell-containing and empty droplets to ~12K
1081   and ~35K.

1082

Next we measured $R_g = \frac{c_g}{TS_g}$ where TS_g represents the number of species from genus g
found in the TSM dataset, and represents the number of species from that genus in the
negative control dataset. This simple parameter can provide insight into how confidently
species from a given genus can be flagged as true signal or contaminant. For example,
if a considerable fraction of species in a genus found in the TSM dataset are also identified
in the control dataset (high R_g), our confidence in other species from that genus would be
lower than in the reverse scenario. This is because as we showed through benchmarking
of both BLASTn and Kraken2 against several truth datasets, species-level taxonomic
assignments based on short reads (~100 bases) are less precise than genus-level
taxonomic assignments. It is also possible to experimentally undersample contaminants
and miss certain species. However, blanket removal of genera may also result in a

significant loss of true signal. Moreover, comparing the distributions of R_g for Salter
genera versus all others in the TS dataset, we observe significantly larger values (one-
sided Mann-Whitney U test, $P < 10^{-5}$), providing support that this parameter can be used
to help distinguish between genera that are known contaminants versus true signal. Thus,
rather than defining an arbitrary cutoff for R_g, we incorporated it in several machine
learning models of contamination (described in the next section) and allowed these
models to learn this parameter along with others. Indeed, the inclusion of R_g boosted the
area under the precision recall curve (AUPR) as well as the area under the receiver
operator curve (AUROC) for the machine learning models discussed in the next section.
1103

1104   We imposed a third filter to eliminate putative fungal species based on a recent study of
1105   fungi in human tissue samples[20]. This filter reduced the number of species to ~6100.
1106   Other numbers remained nearly the same (rounded numbers do not reflect a change, see
1107   exact numbers in **SI table 13**).
1108

Machine learning models of contamination (Filter 4)

Despite having collected many negative control samples representing reagents and
instruments used by the Tabula Sapiens consortium, we built 6 different types of machine
learning (ML) models to learn the shared features of the known and putative
contaminants in the TSM dataset to identify potentially missed contaminants. We used
the best performing model to identify putative contaminants in the part of the dataset that
is withheld from training and testing. As input to each model was a table where the rows
were species and the features were: 1) number of tissues and 2) number of donors
positive for that species, 3) abundance of that species, 4) median percent identity and 5)
median length of alignments corresponding to that species, 6) R_g (described in the
previous section), 7) domain-level taxonomic assignment, and 8) median number of
genes per droplet in which the species was identified. The last parameter is included to
incorporate information about whether the sequence corresponding to a given species is
coming from an empty droplet or from a cell-containing droplet. The premise behind
including the last parameter is that we suspected contaminants and true signals may
have different patterns of association with those two classes of droplets. Counts of empty
droplets were performed using Cell Ranger (v6.0.1) in addition to a custom script for
downstream analysis (see **Data and Materials availability**).

If a species was flagged as a known or putative contaminant by any of the first three
filters (see previous section), it was considered as a contaminant (labeled 1). This dataset
was split into parts A (25% of total) and B (75%). Part A dataset was split into training

(80%) and test (20%) subsets (**SI Figure 1.c**). We built k-nearest neighbor, multi-layer
perceptron, gradient boosting, random forest, logistic regression, and support vector RBF
(radial basis function) models. Because the total dataset (part A + B) contains a small
fraction of contaminating species, it represents an imbalanced dataset where the
detection of the minority positive class, or contaminant species, is of interest. Thus, to
balance the training dataset (and only the training dataset), we employed SMOTE
(Synthetic Minority Oversampling Technique)[106] to generate synthetic minority class
instances from existing ones. Next, we tested the models on the unseen test dataset and
repeated this process 10 times using different train/test splits to generate 95% confidence
intervals for test performance. We report the performance of the model on test data. The
test AUPR and AUROC performance for the models are provided in **SI Figure 1.d**,
ordered by the median value. The 95% confidence intervals for AUPR and AUROC
values are provided in **SI Table 14** and **SI Table 15**. Further details on model creation
and   hyperparameters   is   provided   in
p07_machine_learning_modeling_contamination_and_other_filters.ipynb (see **Data and**
Materials availability).

Although we provide AUROC test scores, we rely primarily on AUPR as the performance
metric of interest since AUROC may be artificially high when the dataset consists of a
minority positive class[107], though in our case the dataset is only moderately imbalanced.
The null AUPR, which corresponds to the prevalence of the positive class, is ~0.3 and
denoted by the dashed line (**SI Figure 1**). The gradient boosting classifier (GBC) had the
highest median AUPR test scores (95% confidence interval: 0.75-0.78) and highest
median AUROC (95% confidence interval: 0.82-.84).

We then fit the GBC model trained and tested in Part A to Part B. We eliminate the known
and putative contaminants in part B (based on filters 1-3) as well as the putative
contaminants identified by the GBC model (filter 4). While some of the species will be
falsely labeled as contaminants, we take a conservative approach and eliminate them
from further analysis. The application of the GBC decontamination model reduced the
number of unique hits, species and genera down to ~41K, ~4700, and ~2200. It lowered
the numbers of cell-containing and empty droplets with hits to ~10K and ~28K.

Gaussian Mixture Model of alignments (Filter 5)

BLASTn outputs with E-values < 10^{-5} against the nt database are conventionally
considered significant[98,99]. In addition to E-value, BLASTn outputs two parameters that
characterize the quality of each alignment, namely percent identity and the alignment
length. By multiplying these two parameters and dividing by 100, we arrive at the total

number of bases with a perfect match, which we refer to as n_match. Although all

significant hits had E-values < 10^{-5}, the distribution of n_match was a bimodal distribution

(**SI Figure 3**). We use a Gaussian Mixture Model (GMM) to separate these two

distributions. Because a species can have hits from both distributions, we systematically

eliminated species if they were represented by more hits from the second distribution (μ_2

= 91 bases) than the first (μ_1 = 62 bases). We posit the first distribution represents lower-

quality alignments that may be biologically interesting, as in the case of sequences from

uncharacterized microbes with only a distant relative in the current nt database. They

may also be technical artifacts. We take a conservative approach and favor precision

over recall and impose this filter. A similar combination of filters has been used by Joint

Genome Institute's microbial genome annotation pipeline (E-value ≤1e-5, ≥ 90 percent

identity, alignment ≥ 80 %)[102] and by the Kraken team when benchmarking their tool

against BLAST (E-value ≤10−5 and percent identity ≥90%)[99]. With the application of the

GMM on the alignments, we arrived at ~13K unique hits, ~2300 species, and ~1100

genera from ~1K cell-containing droplets, and ~11K empty droplets.

Identifying putative habitats of origin for 500+ bacterial genera in the Tabula

Sapiens Microbial dataset using the Genomes from Earth's Microbiomes, the

Human Microbiome Project, and the Unified Human Gastrointestinal Genome

collection (Filter 6)

1184

In addition to experimental (Filter 1) and bioinformatic decontamination (Filters 2-4) and

alignment QC (Filter 5) steps, we wanted to capture information about the likely habitats

of origin for TSM F5 taxa. We posit that some fraction of these species resides in the

human microbiome's external facing sites and have been observed in prior human

microbiome studies, while the rest may be species that are transiently introduced via

inhalation and consumption, among other possible means of transport, or may represent

less frequently observed taxa in the human microbiome. To identify and separate

members of these two groups, we leveraged the Genomes from Earth's Microbiomes[7]

(GEM) dataset to gain insight into the likely habitats from which bacterial hits in the TSM

F5 dataset originate. The GEM dataset is derived from 10,000+ metagenomes sampled

from various habitats across the earth including terrestrial, aquatic, human and animal

hosts, engineered environments, among others. This extensive dataset includes more

than 12000 species-level operational taxonomic units.

1198

We obtained the metadata for the GEM dataset from

https://portal.nersc.gov/GEM/genomes/ (genome_metadata.tsv). While most bacterial

genera were either strictly host-associated or free-living, some contained species that

belonged to both habitat categories. Thus, to identify the dominant habitat of for each

genus, we labeled species based on three categories: 1) "Human-associated" species
include organisms found in the human microbiome; 2) "other_hosts" are species isolated
from non-human vertebrates, invertebrates, fungi, and plants; 3) "free-living" are those
species isolated from human-built or natural environments. With a table where each row
represented a genus, and each column represented the fraction of its species belonging
to one of these three categories based on the GEM dataset, we performed K-nearest
neighbor clustering (number of clusters=3) and assigned each genus to a cluster (**SI**
Figure 4). We created a filter here called GEM_1 which is set to 0 (0=keep) for genera
found in the human-associated cluster, and set to 1 (1=remove) if found in the other two
clusters. GEM_1 is set to NA (NA=not applicable) if a genus is not found in the GEM
dataset. We also created another filter called GEM_2 that functions in the same way as
GEM_1 but for species. The combination of the two filters (GEM_1_2) allows for GEM_2
to override GEM_1. In other words, unless there is habitat information for a given TSM
species, we assume the filter applied on to its genus.
1217
In addition to the GEM dataset, we obtained habitat data for bacterial species from the
Unified Human Gastrointestinal Genome (UHGG) collection[5] and the Human Microbiome
Project (HMP). The UHGG dataset is a large-scale and recent dataset of gut prokaryotes,
comprising ~200K non-redundant genomes from more than 4000 gut prokaryotes. These
genomes are collected from humans living across 31 countries and 6 continents. This
comprehensive catalog includes genomes from the Gastrointestinal Bacteria Culture
Collection[29], the Culturable Genome Reference[108], as well as cultured human gut
genomes from NCBI, PATRIC[31] and IMG[109] datasets, as well as metagenome-assembled
genomes from Pasolli et al.[110], and Nyfach et al.[7] studies. We gathered the Unified Human
Gastrointestinal   Genome   (UHGG)   collection   dataset   from
http://ftp.ebi.ac.uk/pub/databases/metagenomics/mgnify_genomes/human-gut/v2.0.1/
(genomes-all_metadata.tsv). We created a filter called UHGG_species, which is set to 0
(0=keep) if a TSM species has been identified in the UHGG dataset. It is otherwise set to
NA.
1232
The   Human   Microbiome   Project   (HMP)   dataset   acquired   from
https://www.hmpdacc.org/hmp/catalog/grid.php?dataset=genomic consisted of ~3000
entries primarily from bacterial strains from various isolation sites. We chose species
found in external-facing microbiomes (oral cavity, gastrointestinal tract, urogenital tract,
airways, and the skin). When comparing the TSM dataset to these two datasets, entries
with unknown taxonomic assignments were excluded. We created a filter called
HMP_species, which is set to 0 (0=keep) if a TSM species has been identified in the HMP
dataset. It is otherwise set to NA.
1241
1242

Filter 6 combines filters based on **H**MP, **U**HGG and G**E**M datasets, or what we refer to as
the HUE datasets. Explicitly, the HUE filter (filter 6) is set to 1 (1=Remove) if any of the
GEM_1_2, UHGG_species or HMP_species filters is equal to 1. The HUE filter will be set
to 0 (0= Keep) if any of the three filters is equal to 0, effectively overriding the previous
condition. Hence, if in one dataset a given species is found to be human-associated, and
in another it has been found as not human-associated, we consider the species to be
human-associated. The HUE filter is set to NA if all three filters it comprises are set to
NA.
1251
Based on the HUE filter,159 species were determined to be human-associated and the
rest originated from other hosts or environments, or were not found in these datasets. To
obtain putative habits of origin for those species in the NA category, we used OpenAI's
Davinci large language model described in the next section. Removing bacterial species
and genera based on their habitat of origin in the HUE datasets further reduced the
number of unique hits, species, and genera down to ~10K, ~1700, and ~900, respectively,
originating from 1K cell-containing droplets and 8K empty droplets. Please see **Data and**
Materials Availability for the Jupyter notebook where further details for incorporating the
GEM, UHGG, and HMP datasets into TSM dataset can be found.
1261

Benchmarking and employing Davinci large language model for identification of
putative habitats for 1500+ microbial species (Filter 7)

The HUE datasets provide putative habitat information for many taxa, however they do
not include all taxa found in the TSM dataset. Specifically, there were 1500+ species for
which filter 6 (the HUE filter) does not provide any information. To systematically retrieve
information on where these species have previously been isolated from, we employed
openAI's Davinci Large Language Model (text-davinici-003 accessed Jan 2023). We
simply asked the following question for each species: "provide a short summary of where
species_i has been isolated from", where species_i corresponds to the species of interest.
To make queries programmatically, we employed openAI's API and set max_tokens=200.
We used the following default parameters: temperature=0.5, top_p=1, frequency_penalty,
presence_penalty=0.
1274
To better understand the model's level of self-consistency, we randomly chose 10 species
and posed the same prompt three times. A table of model outputs for all three runs is
provided in **SI table 16** and shows that the model presents a certain level of variability
across multiple outputs from which we can identify high-confidence segments based on
consensus across all three outputs. For example, here are the three outputs for
Acidaminococcus intestini: Output 1: "Acidaminococcus intestini has been isolated from
the gut of humans and pigs, as well as from soil samples and wastewater". Output 2:

"Acidaminococcus intestini has been isolated from the intestinal tracts of humans, pigs,
chickens, and horses. It has also been found in soil, water, and sewage". Output 3:
"Acidaminococcus intestini has been isolated from the intestinal tracts of humans, pigs,
and chickens. It has also been detected in the feces of horses and cows, as well as in the
environment".
1287
Next to test the validity of the model's outputs, we randomly selected 100 species from
the GEM dataset, providing us a subset of habitats that a given species has been
previously isolated from. By running three independent queries per species and
combining the model's outputs, we were able to estimate the model's precision and recall.
First, we created two categories, based on whether a given species was previously
isolated from the human microbiome or not (i.e. species isolated from natural
environments, human-built environments, or non-human hosts). Then we grouped
Davinci's responses based on the same two categories. Using GEM as the truth dataset,
we found that the model was able to identify 59 out of 63 species previously found in the
human microbiome. Out of 37 species not found in the human microbiome, the model
correctly identified 25. The model outperformed the truth dataset in some cases. For
example, among the model's false positives, we noticed four species that had been found
previously in the human body by other studies (*Serratia plymuthica*[111], *Leifsonia sp.*[112],
Rickettsia japonica[113], and *Lysobacter oculi*[114]). Correcting for these four species, we
arrive at a precision of 89% and a recall of 94%. We provide a table of these 100 test
species along with GEM habitat categories and model outputs (**SI table 17**). To increase
the precision further, we only accepted species that had been identified as having been
found in humans by all three Davinci outputs. Again, *Serratia plymuthica* and *Lysobacter*
oculi were both found in humans by all three Davinci outputs and thus were counted as
true positives. We arrived at a precision of 98% and a recall of 75% (see **SI Figure 1.e**
for the confusion table). Details of programmatic query submission along with
benchmarking can be found in a Jupyter notebook listed under **Data and Materials**
Availability.
1311
It is important to state some of the limitations of the current Davinci model. First, we found
the model's self-consistency depends on the prompt. Generally, the simpler and more
open-ended prompts provided better, less biased outputs. Additionally, while the model
performs well at classifying species into these two broad categories, we observed that it
performs poorly if the task is identifying detailed habitat information. We noticed it can
generate false results interspersed with true results which are difficult to distinguish. This
is currently a common drawback of generative models in all areas of natural language
processing, referred to as model "hallucination". Thus, although we provide the model's
combined outputs under "davinci_combined" column for each species, we suggest
readers perform independent search for obtaining more detailed habitat information.

1322

Having benchmarked the Davinci model against this prompt, we then applied the model
to the dataset of 1500+ species. To ensure high-confidence classifications, we set this
filter to 0 (0=keep) if for a given species, all three Davinci outputs independently state that
the species has been found previously in the human body. The filter is otherwise set to 1
(1=remove). We found that ~100 species met this criterion. The full dataset with Davinci
model outputs is provided as (**SI Table 18**). Imposing this filter, we arrived at ~3K unique
hits, ~300 species, and ~200 genera from the three biomes of interest. These hits
originated from ~500 cell-containing (~300 of which have cell type annotation) and ~2K
empty droplets.

1332

Impact of filters on the Tabula Sapiens Microbiome dataset

The TSM dataset consisting of significant hits (E-values <10^{-5}) from viruses, bacteria
and fungi is referred to as TSM F0. The TSM dataset that has gone through the first filter
will be referred to as the TSM F1, and so on, such that TSM F7 is the version of the
dataset that has passed through all seven filters. In addition to droplets with TSM F0 hits,
we kept track of droplets (cell-containing or empty) that did not produce any significant
hits. The combination of this group of cells and the TSM F0 group of cells is referred to
as the TSM F00.

1341

1342 F00: The total number of cell-containing and empty droplets analyzed were ~1.3M and
1343 ~156M, respectively.

1344

1345 TSM F0: There were ~724K significant unique hits (counted based on Unique Molecular
1346 Identifiers or UMIs) from three biomes spanning bacteria, fungi and viruses with E-values
1347 < 10^{-5}. These hits originated from ~79K cell-containing droplets (~6% of F00 cell-
1348 containing droplets) and ~380K empty droplets (~0.2% of F00 empty droplets).

1349

TSM F1: When we excluded all species found in negative control samples, the number
of unique hits was reduced to ~63K, or roughly 10% of the F0 hits. Interestingly, the
number of species dropped from ~8200 to only ~7100 and the number of genera went
down from ~2500 to ~2400. Filter 1 reduced the numbers of cell-containing and empty
droplets to ~13K (~16% of F0 cell-containing droplets) and ~44K (~12% of F0 empty
droplets), respectively.

1356

1357 TSM F2: The removal of Salter et al. bacterial genera, reduced the number of unique hits,
1358 species, and genera to ~52K, ~6200, and ~2300, respectively. It reduced the numbers of
1359 cell-containing and empty droplets to ~12K and ~35K.

1360

TSM F3: The removal of putative fungal contaminants previously identified by Narunsky-Haziza *et al.*, reduced the number of species to ~6100. Other numbers remained nearly the same.

TSM F4: The application of the GBC decontamination model, reduced the number of unique hits, species, and genera down to ~41K, ~4700, and ~2200. It lowered the numbers of cell-containing and empty droplets with hits to ~10K and ~28K.

TSM F5: With the application of the gaussian mixture model on the alignments, we arrived at ~13K unique hits, ~2300 species, and ~1100 genera from ~1K cell-containing droplets, and ~11K empty droplets.

TSM F6: Removing bacterial species and genera based on their habitat of origin in the HUE datasets further reduced the number of unique hits, species, and genera down to ~10K, ~1700, and ~900, respectively, originating from 1K cell-containing droplets and 8K empty droplets.

TSM F7: Lastly, removing bacterial, viral and fungal species that were not previously isolated from the human microbiome based on three independent outputs of OpenAI's Davinci model, we arrived at ~3K unique hits, ~300 species, and ~200 genera from the three biomes of interest. These hits originated from ~500 cell-containing (~300 of which have cell type annotation) and ~2K empty droplets.

The exact numbers and fractions of remaining unique hits, species, genera, cell-containing and empty droplets after each filter relative to TSM F0 dataset is shown in **SI Table 13** and **SI Figure 5**. With the imposition of all filters, we arrive at ~0.4% of the unique hits found in the TSM F0 dataset, ~3% of the species and ~5% of the genera. Similar to the percentage of unique hits remaining, the percentage of cell-containing and empty droplets is ~0.7% and ~0.4%, respectively. If we consider the percentage of *total* cell-containing and empty droplets (i.e. compare TSM F7 dataset to the TSM F00 dataset), we see that ~0.04% of cell-containing droplets and ~0.001% of empty droplets remain. Based on these numbers, we can conclude cell-containing droplets are ~40x more likely to harbor microbial reads than empty droplets.

Considering that the starting number of sequences in the TSM dataset was ~70 billion (including human reads) and the final number of unique microbial hits passing through all filters was a few thousand, we conclude that finding high-confidence microbial sequences in human cellular transcriptomes is a search roughly for one in 25 million sequences. Considering the less stringent criteria imposed on the TSM F5 dataset, it is a search for one in 2.5 million sequences. These numbers are a function of various parameters and

1401 filters affecting the recall of our search. Thus, they could be a few orders of magnitude
1402 improved with experimental assays designed specifically for capturing microbial
1403 transcripts as well as other computational improvements.
1404

1405 **Exploring the inter-and-intra donor tissue microbiomes**

1406

To explore the inter-and-intra donor tissue microbiomes, we measure the species unique
fractions. For example, for donor 2, when we sum the number of species that only appear
in one tissue and divide by the total number of species identified in that donor, we arrive
at the unique species fraction equal to 0.53. The average intra-donor unique fraction is
0.64 and the average inter-donor unique fraction, calculated for one tissue across multiple
donors, is 0.76 (**SI Figure 17**). The personal nature of human tissue microbiomes, at least
in part, likely reflects the highly individual-specific microbiomes that exist at external-
facing microbiomes[115–117]. As we also demonstrate through the overlap between species
from different isolation sites in the HMP database, external-facing microbiomes have
limited overlap at the species level **(SI Figure 22)**. Moreover, as we reduce the taxonomic
resolution from species to orders and phyla, the correlations between and within
individuals increase considerably (**SI Figures 18-19**), suggesting that high-level tissue
microbiome compositions are largely conserved.

Single-droplet statistics, outlier cells and outlier cell types

1421

In the population of hit-positive TSM F7 cells, cells with one unique hit dominate the
distribution. The median number of uhits/cell is 1, the mean is ~1.5 and the standard
deviation is 8. See **SI Table 23** for statistics on cell-containing and empty droplets from
TSM F7 dataset. We define outlier cells by those with uhits/cell > 1.5xIQR, where IQR is
the interquartile range for the uhits/cell distribution. The outlier cells are shown in **SI Table
24**. The most prevalent species in outlier cells was EBV. It appeared in outlier cells almost
exclusively from donor 2 (TSP2), with the exception of one cell from the Salivary gland of
TSP7. TSP2 outlier cells positive for EBV hits consisted of bladder B cells, Lymph node
Plasma cells and B cells, among others. In addition to EBV in TSP2, we found a bladder
macrophage containing hits from *Ureaplasma urealyticum*[118]*,* which is a commensal
organism of the urogenital tract and associated with premature birth and adverse
pregnancy outcomes. We found hits to *Aggregatibacter aphrophilus*[119]*,* an oral bacteria
associated with periodontal disease and endocarditis, in an adventitial cell of the salivary
gland in TSP7.

1436

Six out of eight cell types associated with outlier cells were immune cells, namely Plasma
cell, B cell, Natural Killer cell, Macrophage, Memory B cell, and CD34+ CD38- Multipotent

T cell. Interestingly, 72% of outlier cells had viral hits (exclusively EBV) compared to only
7% of empty droplets (**SI Table 25**). Nearly one quarter of outlier cells had bacterial hits
compared to ~83% of outlier empty droplets. Fungal hits appeared in only a small fraction
of outlier cells and droplets. These numbers support the hypothesis that bacterial and
fungal hits mostly come from extracellular spaces and are co-localized with cells or
passively uptaken, whereas in cases of intracellular pathogens such as EBV, one might
expect a higher number of positive cells and a higher number of unique hits per cell. We
found five cells that harbored unique hits from more than one species, primarily bacteria
from different orders and phyla in a single cell.
1448

We grouped cells with TSM F7 hits by their cell type, and through dividing the number of
F7 hit-positive cells by the total number of cells, we measured the fraction of hit-positive
cells for each cell type. To have sufficient numbers for statistical analysis, we chose only
cell types with a minimum of 10 cells that were sequenced and annotated. The mean and
standard deviation of the fraction of cells in each cell type positive for F7 hits were ~0.0006
and ~0.0015%, respectively. We identified outlier cell types as those that have a higher
fraction of F7 positive cells than 1.5xIQR. There were 195 cell types that were included
in the analysis, from which 36 were identified as outlier cell types (**SI table 26**). Among
the outliers, ~60% of cell types belonged to the immune compartment. Some examples
of non-immune outlier cell types are lung alveolar epithelial cells, lung ciliated cells,
intestinal transient amplifying cells, vascular and lymph endothelial cells.
1460

1461 # SI Figures

1462

SI Figure 1. Computational pipeline and post-processing steps developed for the
identification of high-confidence bacterial, viral and fungal sequences in human single-
cell data. **a)** The computational steps comprising SIMBA and post SIMBA filtration of the
Tabula Sapiens Microbiome (TSM) dataset. **b)** Databases used to mitigate and
benchmark bioinformatic false positives (blue) and experimental false positives (orange),
as well as those used for microbial flow and other analyses (green). HBC: Human

Gastrointestinal Bacteria Culture Collection, HOMD: Human Oral Microbiome Dataset, Tiew *et al.* (gut mycobiome), HMP: Human Microbiome Project, nt, Bacterial, Fungal and Viral Refseq datasets (all publically available from NCBI), TSM controls (~200 instrument and reagent negative controls), Salter *et al.* (bacterial genera previously found in prior sequencing studies), Narunsky-Haziza *et al.* (fungal contaminants seen in human tumor samples), Nejman *et al.* (bacterial taxa previously seen in human tumor samples), UHGG (Unified Human Gastrointestinal Genome collection), GEM (Genomes from Earth's Microbiomes), PATRIC (the Pathosystems Resource Integration Center), and EHTM (Extracted Human Tissue Microbiome). **c)** The process by which we used known and putative contaminants to identify additional putative contaminants in the TSM dataset using 6 different types of machine learning models. **d)** Performance metrics of each machine learning model type on withheld test data. For each model type, 10 different models using different train/test splits were created and AUPR and AUROC scores were measured. Dashed lines correspond to null model performance (0.5 for AUROC and the prevalence of the positive class for AUPR). **e)** Confusion matrix for benchmarking the Davinci large language model for microbial habitat classification against the GEM dataset using 100 randomly selected species (Precision = 98%, Recall = 75%). Species are categorized into two groups: 1) found in the human body 2) found anywhere else but the human body. TP: True Positive, FP: False Positive, FN: False Negative, TN: True Negative.

1498

1499

1500   **SI Figure 2.** Kraken2 (against nt) classification assignments for a Tabula Sapiens
1501   Microbiome sample (Bladder, TSP2). **a)** Fraction of classified input sequences by
1502   Kraken2 at different taxonomic levels. **b)** Number of classified sequences that were
1503   assigned to each domain.

1504

1505

1506

1507   **SI Figure 3.** Number of bases with exact match to a reference sequence in the nt
1508   database (n_match). The Gaussian Mixture model is superimposed on the distribution of
1509   n_match (mean or μ_1= 62 bases for first distribution, and μ_2= 91 bases for the second
1510   distribution shown in green).

1511
1512

1513

SI Figure 4. Clustering of bacterial genera based on the fraction of their species belonging to three course-grained labels: free-living (e.g. aquatic, terrestrial), human-dwelling (e.g. gi tract, skin), and other hosts (e.g. plants, insects). Each row represents a bacterial genus from the TSM dataset that also appears in the GEM dataset.

1514
1515
1516
1517

1518

Fraction of each category remaining after each filter

1519

SI Figure 5. Fraction of total microbial hits, unique hits (based on UMIs), species, genera, cell-containing droplets with hits, and empty droplets with hits remaining after each filter is applied. Filter 1: ~200 negative instrument and reagent controls, Filter 2: removal of Salter *et al.* genera, Filter 3: removal of putative fungal contaminants identified by Narunsky-Haziza *et al.*, Filter 4: applying the Gradient-boosting machine learning model for decontamination, Filter 5: applying a Gaussian Mixture Model to select species with higher quality alignments, Filter 6: putative habitat determination using HUE datasets and selecting those taxa more likely to be residents of the human microbiome, and Filter 7: putative habitat determination using the Davinci Large Language Model and selecting additional taxa that are likely residents of the human microbiome.

1530

1531

1532 **SI Figure 6.** Percentage of different types of microbial RNA discovered in the TSM F7
1533 dataset.

1534

Bacterial species (based on ribosomal RNA) colored by phyla

1535

1536   **SI Figure 7.** Ribosomal RNA from different bacterial phyla across various tissue types.
1537   Clustered heatmap of ribosomal hits (presence/yellow and absence/blue is shown) found
1538   in each donor's tissue sample. Bacterial species are color-coded based on their phylum-
1539   level classification. bm: bone marrow, li: large intestine, si: small intestine.

1540

1541

SI Figure 8. Comparisons between the Tabula Sapiens Microbiome (TSM F7) dataset
and the Human Microbiome Project (HMP). a) Number of bacterial genera shared
between TSM F7 and HMP, b) broken down by HMP isolation sites and their closest
tissue counterpart in the TSM dataset. We exclude any members of either dataset that
do not have a taxonomic assignment at the genus level in this comparison. TSM intestines
refers to both the small and the large intestine samples.

1548

1550 **SI Figure 9.** Bacterial species found in both the TSM F7 dataset and the Human
1551 Microbiome project (HMP). Presence is denoted by the light colored boxes. Boxed in red
1552 are species that were referred to in the main text. bm: bone marrow, li: large intestine, si:
1553 small intestine.

1554

1555

1556 **SI Figure 10.** Comparisons between the Tabula Sapiens Microbiome (TSM f7) and the
1557 Unified Human Gastrointestinal Genome (UHGG) collection datasets. left) Number of
1558 bacterial genera shared between TSM F7 tissue microbiomes and UHGG. right) Number
1559 of bacterial genera shared between TSM F7 intestinal tissue microbiomes (both large and
1560 small intestines) and UHGG. We treat TSM datasets as set A and UHGG as set B.

1561

1562

SI Figure 11. Exploring the intersection of bacterial genera found in the Tabula Sapiens
Microbiome (TSM) a) F7 or b) F5 datasets and tumor microbiomes from Nejman *et al.*

(only species and orders appearing in tumors that passed all of this study's contamination
filters were selected for analysis, refer to this study's Table S4). The percentage of shared
genera compared to total number of genera in each dataset are shown (set A corresponds
to TSM datasets, and set B corresponds to the Nejman *et al.* dataset). c) abundance of
bacterial genera shared between the TSM F7 stratified by tissue and donor of origin and
Nejman *et al.* datasets. Abundance of genera is based on Log_2(hits +1). Entries with
unknown genera are excluded from comparisons shown in the venn diagrams. bm: bone
marrow, li: large intestine, si: small intestine.

1573

1574

SI Figure 12. Comparison of bacterial genera in healthy tissues from TSM F7 and tumors
from Nejman *et al.* (Only genera appearing in tumors that passed all of this study's
contamination filters were selected for analysis from Nejman *et al*. Table S4, and from
their Figure 4D for tumor enriched species). a) venn-diagram representing the number of
shared genera between the two datasets. b) Number of bacterial genera shared between
each tumor type and each TSM F7 tissue type is shown. bm: bone marrow, li: large
intestine, si: small intestine, GBM: glioblastoma multiforme. c) abundance of species that
were highly enriched in tumors and also appeared in the TSM F5 dataset. Abundance of
species is based on $\log_2$ hits +1.

1584

1585

1586   **SI Figure 13.** Abundance of bacterial genera previously found in tumor samples (Nejman
1587   *et. al*, 2020) and in the Tabula Sapiens Microbiome (TSM) F5 dataset stratified by tissue
1588   of origin. Abundance is based on log$_2$(hits +1). Samples with "_T" appended correspond
1589   to tumors. bm: bone marrow, li: large intestine, si: small intestine, GBM: glioblastoma
1590   multiforme.

1591

1592

1593   **SI Figure 14.** Abundance of bacterial genera previously found in tumor samples (Nejman

1594   *et. al*, 2020) and in the Tabula Sapiens Microbiome (TSM) F7 dataset stratified by tissue

1595   of origin. Abundance is based on $\log_2$ (hits +1). Samples with "_T" appended correspond

1596   to tumors. bm: bone marrow, li: large intestine, si: small intestine, GBM: glioblastoma

1597   multiforme.

1598

TSP14 (donor 14) microbiome grouped by tissue

Tissue microbiomes grouped by donor

1599

1600   **SI Figure 15.** Circos plots of shared species within (a,b) and between donors (c,d). Each
1601   link represents a unique species. Species that are not shared are not shown. Left panels
1602   correspond to TSM F7 and the right panels correspond to TSM F5 datasets.

1603
1604

1605

SI Figure 16. UpSet plots for TSP2 (donor 2) TSM F5 species demonstrating the number
of shared taxa between tissues, as well as the number of taxa unique to each tissue.
Unique sets of species are shown as filled circles. The intersection of two or more sets
are shown via connected circles. Bars denote the size of each set or the intersection of
sets. For visual clarity, the minimum subset size has been set to 2. bm: bone marrow, li:
large intestine, si: small intestine, vasc: vasculature.

1612

1613

1614

1615

SI Figure 17. UpSet plots depicting the number of species shared and unique to each
sample. Panels a and b depict intra-donor comparisons of tissues, while c-f depict inter-
donor comparisons of the same tissue. Unique sets of species are shown as filled circles.
The intersection of two or more sets are shown via connected circles. Bars denote the
size of each set or the intersection of sets. Panel g summarizes the number and fraction
of species that appear in more than one sample within each panel a-f.

1622

1623

1624   **SI Figure 18.**   Tissue microbiomes across tissues and donors for TSM F7 bacterial
1625   dataset at the a) species, b) genera, c) orders and d) phyla-level classification. Pairwise
1626   Spearman correlation cluster maps across samples grouped by donor and tissue are
1627   shown. Only samples with more than 50 hits are shown.

1628

1629   **SI Figure 19.**   Tissue microbiomes across donor_tissue samples for TSM F5 bacterial
1630   dataset at the a) species, b) genera, c) orders and d) phyla-level classification. Pairwise
1631   Spearman correlation cluster maps across samples grouped by donor and tissue are
1632   shown. Only samples with more than 50 hits are shown.

1633

1634
1635 https://pubmed.ncbi.nlm.nih.gov/10516026/

1636 **SI Figure 20.** The genomic position of identified HERV K hits (left) with respect to the
1637 annotated HERV K genome (right) (Tönjes, 1999).

1638

1639

1640 **SI Figure 21.** Fluorescence staining of donor-derived bacterial-like particles (BLPs) from
1641 the lung (a) and Virus-like particles (VLPs) from a lung and a lymph node sample (b,c).

1642 These images were used to estimate VLP and BLP concentrations in a cubic centimeter

1643 of tissue (~10^8 VLPs/cm^3 and ~10^7 BLPs/cm^3).

1644

1646 **SI Figure 22.** UpSet plot showing the taxa shared across different external-facing

1647 microbiomes in the Human Microbiome Project (HMP) dataset. Unique sets of taxa are

1648 shown as filled circles. The intersection of two or more sets are shown via connected

1649 circles. Bars denote the size of each set or the intersection of sets.

1650

SI Tables

1652 **SI Table 1.** Precision and recall associated with SIMBA using a random selection of 100

1653 viral genomes and 10 sequences per genome as a truth dataset.

	domain	phylum	class	order	family	genus	species
precision	1.0	1.0	1.0	1.0	1.0	0.999	0.809
recall	1.0	1.0	1.0	1.0	1.0	1.000	1.000

1654

1655 **SI Table 2.** Examples of false positives associated with the viral truth dataset at the

1656 species level. "Species_truth" and "species_test" columns correspond to species found

1657 in the truth and test datasets, respectively.

	species_truth	species_test
0	Acinetobacter virus AbP2	Acinetobacter phage Bphi-R1888
1	Acinetobacter virus D2	Acinetobacter phage vB_AbaP_APK44
2	Acinetobacter virus D2	Acinetobacter phage vB_AbaP_D2M
3	Acinetobacter virus D2	Acinetobacter phage vB_AbaP_PMK34
4	Arthrobacter virus Piccoletto	Arthrobacter virus Beans
5	Campylobacter virus CP220	Campylobacter phage CP20
6	Campylobacter virus CP220	Campylobacter virus CP21
7	Campylobacter virus CP220	Campylobacter virus IBB35
8	Canna yellow mottle associated virus	Canna yellow mottle virus
9	Escherichia virus VR7	Escherichia virus VR25
10	Escherichia virus VR7	Escherichia virus VR26
11	Leuconostoc virus Lmd1	Leuconostoc virus LN6B
12	Mycobacterium phage ArcherNM	Mycobacterium phage Changeling
13	Mycobacterium phage Phaedrus	Mycobacterium phage Akoma
14	Mycobacterium phage Phaedrus	Mycobacterium phage Compostia
15	Mycobacterium phage Tiffany	Mycobacterium phage Dieselweasel
16	Mycobacterium phage Tiffany	Mycobacterium phage P28Green
17	Mycobacterium phage Tiffany	Mycobacterium phage SoilDragon
18	Mycobacterium phage Tiffany	Mycobacterium virus Bxz2
19	Mycobacterium virus Skinnyp	Mycobacterium phage ShrimpFriedEgg
20	Mycobacterium virus TA17a	Mycobacterium phage Holeinone

1658

1659 **SI Table 3.** Examples of false positives associated with the viral truth dataset and the
1660 corresponding percent identity ("pident_test" column) and the length ("length_test"
1661 column) of the alignment between the input seq and output seq. "Species_truth" and
1662 "species_test" columns correspond to species found in the truth and test datasets,
1663 respectively.

species_truth	species_test	pident_test	length_test
Pseudomonas phage C11	Pseudomonas phage vB_PaeM_SCUT-S2	100.0	100.0
Pseudomonas phage C11	Pseudomonas phage PaZq-1	100.0	100.0
Pseudomonas phage C11	Pseudomonas phage PaGz-1	100.0	100.0
Pseudomonas phage C11	Pseudomonas phage K5	100.0	100.0
Acinetobacter virus AbP2	Acinetobacter phage Bphi-R1888	100.0	100.0
Pseudomonas phage DL54	Pseudomonas phage phiPA01_EW	100.0	100.0
Pseudomonas phage DL54	Pseudomonas virus Pa222	100.0	100.0
Pseudomonas phage DL54	Pseudomonas phage otherone	100.0	100.0
Mycobacterium phage Phaedrus	Mycobacterium phage Akoma	100.0	100.0
Mycobacterium phage Phaedrus	Mycobacterium phage Compostia	100.0	100.0
Mycobacterium virus Skinnyp	Mycobacterium phage ShrimpFriedEgg	100.0	100.0
Salmonella phage SEN5	Salmonella virus SEN4	100.0	100.0
Pseudomonas virus LMA2	Pseudomonas phage PHW2	100.0	100.0
Pseudomonas virus LMA2	Pseudomonas phage willy	100.0	100.0
Pseudomonas virus LMA2	Pseudomonas phage PaGU11	100.0	100.0

1664

1665 **SI Table 4.** Precision and recall scores using a truth dataset containing 5000 sequences
1666 from 50 gut bacterial species (randomly selected sequences and species).

	domain	phylum	class	order	family	genus	species
precision	1.000	0.999	0.997	0.996	0.973	0.947	0.699
recall	0.291	0.291	0.290	0.290	0.285	0.280	0.223

1667

1668 **SI Table 5.** Species from the truth dataset and their corresponding false label at the
1669 level of species classification. First five false positives are shown. "Species_truth" and
1670 "species_test" columns correspond to species found in the truth and test datasets,
1671 respectively.

1672

species_truth	species_test
Bifidobacterium faecale	Bifidobacterium adolescentis
Enterococcus lactis	Enterococcus faecium
Lacticaseibacillus casei	Lacticaseibacillus zeae
Bacteroides faecis	Bacteroides thetaiotaomicron
Butyricimonas virosa	Butyricimonas faecalis

1673

1674 **SI Table 6.** Precision and recall scores using a truth dataset containing 5000 sequences
1675 from 50 oral bacterial species from the expanded Human Oral Microbiome Database
1676 (randomly selected sequences and species).

1677

	domain	phylum	class	order	family	genus	species
precision	1.000	1.000	0.996	0.990	0.987	0.974	0.854
recall	0.209	0.209	0.208	0.207	0.207	0.205	0.184

1678

1679 **SI Table 7.** Precision and recall scores using a truth dataset containing 5000 sequences
1680 from the bacterial RefSeq database (based on 50 randomly selected species and 100
1681 sequences from each).

1682

	superkingdom	phylum	class	order	family	genus	species
precision	1.000	1.000	1.000	0.996	0.992	0.985	0.897
recall	0.052	0.052	0.052	0.052	0.052	0.051	0.047

1683

1684 **SI Table 8.** Precision and recall scores using a truth dataset containing sequences from
1685 a fungal dataset[104] (based on ~5000 sequences).

	domain	phylum	class	order	family	genus	species
precision	1.000	0.990	0.988	0.985	0.961	0.948	0.793
recall	0.207	0.205	0.205	0.204	0.200	0.198	0.171

1686

1687 **SI Table 9.** Kraken2 precision and recall scores using 3 truth datasets each containing
1688 5000 sequences (50 randomly selected species and 100 sequences from each) from the
1689 bacterial RefSeq database, Human Gastrointestinal Bacteria Culture Collection (HBC)
1690 and the expanded Human Oral Microbiome Database (HOMD).

Bacterial Refseq

	domain	phylum	class	order	family	genus	species
precision	0.997	0.997	0.997	0.997	0.997	0.997	0.991
recall	0.984	0.983	0.981	0.980	0.979	0.977	0.863

HBC Gut bacteria

	domain	phylum	class	order	family	genus	species
precision	0.952	0.948	0.944	0.945	0.927	0.929	0.664
recall	0.619	0.606	0.603	0.603	0.586	0.534	0.417

HOMD Oral bacteria

	domain	phylum	class	order	family	genus	species
precision	0.991	0.990	0.988	0.979	0.973	0.971	0.912
recall	0.767	0.758	0.754	0.752	0.703	0.745	0.635

1691

1692 **SI Table 10.** Testing the precision of Kraken2 (against nt) classification assignments for
1693 a Tabula Sapiens Microbiome sample (Bladder, donor 2). Mean precision and standard
1694 deviation for three batches of input sequences selected from the classified reads (100
1695 randomly selected sequences per batch from each domain, excluding Archaea).

1696

1697 **Mean Precision +/- SD**

	domain	phylum	class	order	family	genus	species
Eukaryotic	0.77 ± 0.04	0.72 ± 0.07	0.61 ± 0.06	0.6 ± 0.06	0.57 ± 0.06	0.54 ± 0.07	0.54 ± 0.07
Bacterial	0.06 ± 0.01	0.06 ± 0.01	0.06 ± 0.01	0.06 ± 0.01	0.06 ± 0.01	0.06 ± 0.01	0.05 ± 0.01
Viral	0.03 ± 0.01	0.03 ± 0.01	0.03 ± 0.02	0.03 ± 0.02	0.03 ± 0.02	0.04 ± 0.03	0.03 ± 0.02

1698

1699 **SI Table 11.** Examples of the type of reads that are misclassified as bacterial by
1700 Kraken2 (against nt) and their classification based on BLASTn (against nt).

kraken2_species	species	kraken2_order	order	kraken2_superkingdom	superkingdom
Pasteurella multocida	Pan paniscus	Pasteurellales	Primates	Bacteria	Eukaryota
Paenibacillus kribbensis	Pan paniscus	Bacillales	Primates	Bacteria	Eukaryota
Kitasatospora sp. MMS16-BH015	Homo sapiens	Streptomycetales	Primates	Bacteria	Eukaryota
Coprothermobacter proteolyticus	Pan paniscus	Coprothermobacterales	Primates	Bacteria	Eukaryota
Desulfovibrio marinus	Homo sapiens	Desulfovibrionales	Primates	Bacteria	Eukaryota
Streptomyces autolyticus	Pan troglodytes	Streptomycetales	Primates	Bacteria	Eukaryota
[Clostridium] cellulosi	Homo sapiens	Clostridiales	Primates	Bacteria	Eukaryota
Citrobacter werkmanii	Homo sapiens	Enterobacterales	Primates	Bacteria	Eukaryota
Deinococcus ficus	Homo sapiens	Deinococcales	Primates	Bacteria	Eukaryota
Methylobacterium durans	Gorilla gorilla	Rhizobiales	Primates	Bacteria	Eukaryota
Pseudomonas putida	Homo sapiens	Pseudomonadales	Primates	Bacteria	Eukaryota
Ectothiorhodospiraceae bacterium BW-2	Pan paniscus	Chromatiales	Primates	Bacteria	Eukaryota
Altererythrobacter sp. TH136	Homo sapiens	Sphingomonadales	Primates	Bacteria	Eukaryota
Nocardioidaceae bacterium	Callithrix jacchus	Propionibacteriales	Primates	Bacteria	Eukaryota
Nakamurella sp. PAMC28650	Macaca mulatta	Nakamurellales	Primates	Bacteria	Eukaryota
Clostridiales bacterium	Pan paniscus	Clostridiales	Primates	Bacteria	Eukaryota
Oceanicoccus sagamiensis	Homo sapiens	Cellvibrionales	Primates	Bacteria	Eukaryota
Qipengyuania seohaensis	Pan paniscus	Sphingomonadales	Primates	Bacteria	Eukaryota
Flammeovirgaceae bacterium	Homo sapiens	Cytophagales	Primates	Bacteria	Eukaryota
Enterobacteriaceae endosymbiont of Donacia ful...	Homo sapiens	Enterobacterales	Primates	Bacteria	Eukaryota
Aeromonas encheleia	Pan paniscus	Aeromonadales	Primates	Bacteria	Eukaryota
Nostoc punctiforme	Pan paniscus	Nostocales	Primates	Bacteria	Eukaryota
Pelagovum pacificum	Homo sapiens	Rhodobacterales	Primates	Bacteria	Eukaryota
Geobacter bemidjiensis	Pan paniscus	Desulfuromonadales	Primates	Bacteria	Eukaryota

1701

SI Table 12. List of reagents and instruments per tissue used to generate negative control
samples.

SI Table 13. Absolute counts and fraction of total microbial hits, unique hits (based on
UMIs), species, genera, cell-containing droplets with hits, and empty droplets with hits
remaining after each filter is applied.

SI Table 14. 95% AUROC confidence intervals for different machine learning models
tested on unseen data. "auroc_test_95perc_ci_lower" and "auroc_test_95perc_ci_upper"
columns correspond to the lower and upper bounds for the 95% confidence interval,
respectively.

	auroc_test_95perc_ci_lower	auroc_test_95perc_ci_upper
gradient boosting classifier	0.82	0.836
random forest	0.785	0.807
multi-layer perceptron	0.765	0.797
support vector rbf	0.733	0.767
knn	0.71	0.758
logistic regression	0.703	0.727

1711

1712 **SI Table 15.** 95% AUPR confidence intervals for different machine learning models tested
1713 on unseen data. "aupr_test_95perc_ci_lower" and "aupr_test_95perc_ci_upper" columns
1714 correspond to the lower and upper bounds for the 95% confidence interval, respectively.

	aupr_test_95perc_ci_lower	aupr_test_95perc_ci_upper
gradient boosting classifier	0.748	0.776
random forest	0.715	0.753
multi-layer perceptron	0.69	0.736
support vector rbf	0.668	0.702
knn	0.636	0.69
logistic regression	0.63	0.66

1715

SI Table 16. Evaluating Davinci model's level of self-consistency to the same input
prompt. Three independent outputs are listed for each species. The temperature
parameter was set to default (0.5).

SI Table 17. Davinci model outputs for 100 test species using the GEM dataset as the
truth dataset. "davinci_0", "davinci_1", "davinci_2" columns report the three independent
outputs of the Davinci model for each species. "Davinci_human" is set to 0 if a given
species is found in humans by at least one of the three Davinci outputs, and 1 otherwise.
"Davinci_human_3x" is set to 0 if all three Davinci outputs report that a given species has
been found in humans. The "gem_human" column is the truth dataset, set to 0 if a given
species is found in humans in the GEM dataset, and 1 otherwise.

SI Table 18. Davinci model outputs for 1500+ species whose habitats were not identified
by the HUE datasets. "davinci_0" , "davinci_1", "davinci_2" columns report the three
independent outputs of the Davinci model for each species. "habitat_assignment" column
is set to 0 if all three outputs state the presence of a given species in the human body,
and set to 1 otherwise.

SI Table 19. Fraction of microbial hits in the TSM F5 and F7 datasets that have taxonomic
assignment at different levels of taxonomic categorization.

	index	f5	f7
0	species	1.00	1.00
1	genus	0.88	0.90
2	family	0.92	0.94
3	order	0.96	0.95
4	class	0.97	0.97
5	phylum	0.98	0.99
6	domain	1.00	1.00

1733

SI Table 20. The table underlying the flow paths from **Figure 2.a** (Sankey plot). Columns:
source, dest (destination), values. Source and destination columns represent the nodes
in the Sankey plot, while the values demonstrate the weight applied onto the connecting
edges. The suffixes "_phy", "_ord", "_tis", "_tum", "_cou", "_hmp", "_hab" are denoting
phyla, orders, tissues, Nejman *et al.* tumor types, UHGG countries, HMP external facing
sites, and GEM habitats, respectively. bm: bone marrow, li: large intestine, si: small
intestine.

SI Table 21. Genera found in tumor microbiomes that are not found in the TSM F7 or
TSM F5 datasets.

SI Table 22. The table underlying the flow paths from **SI File 2** (Sankey plot). Columns:
tumor_type (Nejman *et al.*), genus and tissue (TSM F7), hmp_site (HMP), Country
(UHGG), habitat2 (GEM). The suffixes are: "_phy", "_ord", "_tis", "_tum", "_cou", "_hmp",
"_hab" are denoting phyla, orders, tissues, Nejman *et al.* tumor types, UHGG countries,
HMP external facing sites, and GEM habitats, respectively. bm: bone marrow, li: large
intestine, si: small intestine.

SI Table 23. Statistics for the distribution of unique hits per cell ("cell_umi" column) and
empty droplets ("empty_umi column) in the TSM F7 dataset.

SI Table 24. Outlier cells with their associated species and domains, as well as their
tissue, donor and cell type of origin. BM: Bone Marrow.

SI Table 25. Relative fraction of outlier cells and empty droplets from TSM F7, stratified
by the domains where the hits map to. All outlier viral hits from cells and empty droplets
map to EBV.

SI Table 26. Outlier cell types in TSM F7 dataset. Fraction and total number of cells in
each cell type that had F7 hits are shown. Only cell types with a minimum of 10 cells
sequenced are included.

SI Table 27. HERV K (HML-2) transcripts across donors, spanning a diverse range of
tissues and cell types.

SI Files

1761

SI File 1. The HTML file that corresponds to **Figure 2.a** (Sankey plot). This file can be
opened using a browser for an interactive experience.

SI File 2. The HTML file of a Sankey plot showing the genera shared between several
datasets: TSM F7, GEM, UHGG, Nejman et al.). This file can be opened using a browser.
Columns from left to right are the same as those from the main Sankey plot (**Figure 2.a**).
The suffixes are: "_phy", "_ord", "_tis", "_tum", "_cou", "_hmp", "_hab" are denoting phyla,
orders, tissues, Nejman *et al.* tumor types, UHGG countries, HMP external facing sites,
and GEM habitats, respectively. bm: bone marrow, li: large intestine, si: small intestine.

SI File 3. The HTML file for the network plot shown in **Figure 3.b**. Please see **Figure 3.b**
caption for more information about node and edge weights.

SI File 4. The HTML file for EBV Sankey plot shown in **Figure 4.a**. The columns from left
to right correspond to donors, tissues, cell types and compartments. The suffixes for these
columns are "_don", "_tis", "_cel", and "_com". bm: bone marrow.

1775
1776

The Tabula Sapiens Consortium Author List

1777

Overall Project Direction and Coordination

1778

Robert C. Jones[1] , Jim Karkanias[2] , Mark Krasnow[3,4], Angela Oliveira Pisco[2] , Stephen R. Quake[1,2,5], Julia Salzman[3,6], Nir Yosef[2,7,8,9]

1779
1780

Donor Recruitment

1781

Bryan Bulthaup[10], Phillip Brown[10], William Harper[10], Marisa Hemenez[10], Ravikumar Ponnusamy[10], Ahmad Salehi[10], Bhavani A. Sanagavarapu[10], Eileen Spallino[10]

1782
1783

Surgeons

1784

Ksenia A. Aaron[11], Waldo Concepcion[10], James M. Gardner[12,13], Burnett Kelly[10,14], Nikole Neidlinger[10], Zifa Wang[10]

1785
1786

Logistical coordination

1787

Sheela Crasta[1,2], Saroja Kolluru[1,2], Maurizio Morri[2] , Angela Oliveira Pisco[2] , Serena Y. Tan[15], Kyle J. Travaglini[3] , Chenling Xu[7]

1788
1789

Organ Processing

1790

Marcela Alcántara-Hernández[16], Nicole Almanzar[17], Jane Antony[18], Benjamin Beyersdorf[19], Deviana Burhan[20], Kruti Calcuttawala[21], Matthew M. Carter[16], Charles K. F. Chan[18,22], Charles A. Chang[23], Stephen Chang[3,19], Alex Colville[21,24], Sheela Crasta[1,2], Rebecca N. Culver[25], Ivana Cvijović[1,5], Gaetano D'Amato[26], Camille Ezran[3] , Francisco X. Galdos[18], Astrid Gillich[3] , William R. Goodyer[27], Yan Hang[23,28], Alyssa Hayashi[1] , Sahar Houshdaran[29], Xianxi Huang[19,30], Juan C. Irwin[29], SoRi Jang[3] , Julia Vallve Juanico[29], Aaron M. Kershner[18], Soochi Kim[21,24], Bernhard Kiss[18], Saroja Kolluru[1,2], William Kong[18], Maya E. Kumar[17], Angera H. Kuo[18], Rebecca Leylek[16], Baoxiang Li[31], Gabriel B. Loeb[32], Wan-Jin Lu[18], Sruthi Mantri[33], Maxim Markovic[1] , Patrick L. McAlpine[11,34], Antoine de Morree[21,24], Maurizio Morri[2] , Karim Mrouj[18], Shravani Mukherjee[31], Tyler Muser[17], Patrick Neuhöfer[3,35,36], Thi D. Nguyen[37], Kimberly Perez[16], Ragini Phansalkar[26], Angela Oliveira Pisco[2] , Nazan Puluca[18] , Zhen Qi[18], Poorvi Rao[20], Hayley Raquer-McKay[16], Nicholas Schaum[18,21], Bronwyn Scott[31], Bobak Seddighzadeh[38], Joe Segal[20], Sushmita Sen[29], Shaheen Sikandar[18], Sean P. Spencer[16], Lea Steffes[17], Varun R. Subramaniam[31], Aditi Swarup[31], Michael Swift[1], Kyle J. Travaglini[3] , Will Van Treuren[16], Emily Trimm[26], Stefan Veizades[19,39], Sivakamasundari Vijayakumar[18], Kim Chi Vo[29], Sevahn K. Vorperian[1,40], Wanxin

Wang[29], Hannah N.W. Weinstein[38], Juliane Winkler[41], Timothy T.H. Wu[3] , Jamie Xie[38],
Andrea R.Yung[3] , Yue Zhang[3]

Sequencing

Angela M. Detweiler[2] , Honey Mekonen[2] , Norma F. Neff[2] , Rene V. Sit[2] , Michelle Tan[2] ,
Jia Yan[2]

Histology

Gregory R. Bean[15], Vivek Charu[15], Erna Forgó[15], Brock A. Martin[15], Michael G.
Ozawa[15], Oscar Silva[15], Serena Y. Tan[15], Angus Toland[15], Venkata N.P. Vemuri[2]

Data Analysis

Shaked Afik[7] , Kyle Awayan[2] , Rob Bierman[3,6], Olga Borisovna Botvinnik[2] , Ashley
Byrne[2] , Michelle Chen[1] , Roozbeh Dehghannasiri[3,6], Angela M. Detweiler[2] , Adam
Gayoso[7] , Alejandro A Granados[2] , Qiqing Li[2] , Gita Mahmoudabadi[1] , Aaron McGeever[2]
, Antoine de Morree[21,24], Julia Eve Olivieri[3,6,42], Madeline Park[2] , Angela Oliveira Pisco[2] ,
Neha Ravikumar[1] , Julia Salzman[3,6], Geoff Stanley[1] , Michael Swift[1] , Michelle Tan[2] ,
Weilun Tan[2] , Alexander J Tarashansky[2] , Rohan Vanheusden[2] , Sevahn K.
Vorperian[1,40], Peter Wang[3,6], Sheng Wang[2] , Galen Xing[2] , Chenling Xu[6] , Nir Yosef[2,6,7,8]

Expert Cell Type Annotation

Marcela Alcántara-Hernández[16], Jane Antony[18], Charles K. F. Chan[18,22], Charles A.
Chang[23], Alex Colville[21,24], Sheela Crasta[1,2], Rebecca Culver[25], Les Dethlefsen[43],
Camille Ezran[3] , Astrid Gillich[3] , Yan Hang[23,28], Po-Yi Ho[16], Juan C. Irwin[29], SoRi Jang[3] ,
Aaron M. Kershner[18], William Kong[18], Maya E Kumar[17], Angera H. Kuo[18], Rebecca
Leylek[16], Shixuan Liu[3,44], Gabriel B. Loeb[32], Wan-Jin Lu[18], Jonathan S Maltzman[45,46],
Ross J. Metzger[27,47], Antoine de Morree[21,24], Patrick Neuhöfer[3,35,36], Kimberly Perez[16],
Ragini Phansalkar[26], Zhen Qi[18], Poorvi Rao[20], Hayley RaquerMcKay[16], Koki
Sasagawa[19], Bronwyn Scott[31], Rahul Sinha[15,18,35], Hanbing Song[38], Sean P. Spencer[16],
Aditi Swarup[31], Michael Swift[1] , Kyle J. Travaglini[3] , Emily Trimm[26], Stefan Veizades[19,39],
Sivakamasundari Vijayakumar[18], Bruce Wang[20], Wanxin Wang[29], Juliane Winkler[41],
Jamie Xie[38], Andrea R.Yung[3]

Tissue Expert Principal Investigators

Steven E. Artandi[3,35,36], Philip A. Beachy[18,23,48], Michael F. Clarke[18], Linda C. Giudice[29],
Franklin W. Huang[38,49], Kerwyn Casey Huang[1,16], Juliana Idoyaga[16], Seung K Kim[23,28],
Mark Krasnow[3,4], Christin S. Kuo[17], Patricia Nguyen[19,39,46], Stephen R. Quake[1,2,5],

Thomas A. Rando[21,24], Kristy Red-Horse[26], Jeremy Reiter[50], David A. Relman[16,43,46],
Justin L. Sonnenburg[16], Bruce Wang[20], Albert Wu[31], Sean M. Wu[19,39], Tony Wyss-
Coray[21,24]

<u>Affiliations</u>

[1]Department of Bioengineering, Stanford University; Stanford, CA, USA.
[2]Chan Zuckerberg Biohub; San Francisco, CA, USA.
[3]Department of Biochemistry, Stanford University School of Medicine; Stanford, CA,
USA.
[4]Howard Hughes Medical Institute; USA.
[5]Department of Applied Physics, Stanford University; Stanford, CA, USA.
[6]Department of Biomedical Data Science, Stanford University; Stanford, CA, USA.
[7]Center for Computational Biology, University of California Berkeley; Berkeley, CA,
USA.
[8]Department of Electrical Engineering and Computer Sciences, University of California
Berkeley; Berkeley, CA, USA.
[9]Ragon Institute of MGH, MIT and Harvard; Cambridge, MA, USA.
[10]Donor Network West; San Ramon, CA, USA.
[11]Department of Otolaryngology-Head and Neck Surgery, Stanford University School of
Medicine; Stanford, California, USA.
[12]Department of Surgery, University of California San Francisco; San Francisco, CA,
USA. [13]Diabetes Center, University of California San Francisco; San Francisco, CA,
USA.
[14]DCI Donor Services; Sacramento, CA, USA.
[15]Department of Pathology, Stanford University School of Medicine; Stanford, CA, USA.
[16]Department of Microbiology and Immunology, Stanford University School of Medicine;
Stanford, CA, USA.
[17]Department of Pediatrics, Division of Pulmonary Medicine, Stanford University;
Stanford, CA, USA.
[18]Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School
of Medicine; Stanford, CA, USA.
[19]Department of Medicine, Division of Cardiovascular Medicine, Stanford University;
Stanford, CA, USA.
[20]Department of Medicine and Liver Center, University of California San Francisco; San
Francisco, CA, USA.
[21]Department of Neurology and Neurological Sciences, Stanford University School of
Medicine; Stanford, CA, USA.
[22]Department of Surgery - Plastic and Reconstructive Surgery, Stanford University
School of Medicine; Stanford, CA, USA.

[23]Department of Developmental Biology, Stanford University School of Medicine,
Stanford, CA, USA
[24]Paul F. Glenn Center for the Biology of Aging, Stanford University School of Medicine;
Stanford, CA, USA.
[25]Department of Genetics, Stanford University School of Medicine; Stanford, CA, USA.
[26]Department of Biology, Stanford University; Stanford, CA, USA.
[27]Department of Pediatrics, Division of Cardiology, Stanford University School of
Medicine; Stanford, CA, USA.
[28]Stanford Diabetes Research Center, Stanford University School of Medicine, Stanford,
California
[29]Center for Gynecology and Reproductive Sciences, Department of Obstetrics,
Gynecology and Reproductive Sciences, University of California San Francisco; San
Francisco, CA, USA. [30]Department of Critical Care Medicine, The First Affiliated
Hospital of Shantou University Medical College; Shantou, China.
[31]Department of Ophthalmology, Stanford University School of Medicine; Stanford, CA,
USA. [32]Division of Nephrology, Department of Medicine, University of California San
Francisco; San Francisco, CA, USA.
[33]Stanford University School of Medicine; Stanford, CA, USA.
[34]Mass Spectrometry Platform, Chan Zuckerberg Biohub; Stanford, CA, USA.
[35]Stanford Cancer Institute, Stanford University School of Medicine; Stanford, CA, USA.
[36]Department of Medicine, Division of Hematology, Stanford University School of
Medicine, Stanford, CA, USA
[37]Department of Biochemistry and Biophysics, Cardiovascular Research Institute,
University of California San Francisco; San Francisco, CA, USA.
[38]Division of Hematology and Oncology, Department of Medicine, Bakar Computational
Health Sciences Institute, Institute for Human Genetics, University of California San
Francisco; San Francisco, CA, USA.
[39]Stanford Cardiovascular Institute; Stanford CA, USA.
[40]Department of Chemical Engineering, Stanford University; Stanford, CA, USA.
[41]Department of Cell & Tissue Biology, University of California San Francisco; San
Francisco, CA, USA.
[42]Institute for Computational and Mathematical Engineering, Stanford University;
Stanford, CA, USA.
[43]Division of Infectious Diseases & Geographic Medicine, Department of Medicine,
Stanford University School of Medicine; Stanford, CA, USA.
[44]Department of Chemical and Systems Biology, Stanford University School of
Medicine, Stanford, CA, USA
[45]Division of Nephrology, Stanford University School of Medicine; Stanford, CA, USA.
[46]Veterans Affairs Palo Alto Health Care System; Palo Alto, CA, USA.

47Vera Moulton Wall Center for Pulmonary and Vascular Disease, Stanford University School of Medicine; Stanford, CA, USA.

48Department of Urology, Stanford University School of Medicine, Stanford, CA, USA

49Division of Hematology/Oncology, Department of Medicine, San Francisco Veterans Affairs Health Care System, San Francisco, CA, USA.

50Department of Biochemistry, University of California San Francisco; San Francisco, CA, USA.

Citations

1. Lynch, M. & Marinov, G. K. The bioenergetic costs of a gene. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 15690–15695 (2015).

2. Mahmoudabadi, G., Milo, R. & Phillips, R. Energetic cost of building a virus. *Proc. Natl. Acad. Sci.* **114**, E4324–E4333 (2017).

3. Gilbert, J. *et al.* Current understanding of the human microbiome. *Nat. Med.* **24**, 392–400 (2018).

4. Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human Microbiome Project. *Nature* **550**, 61–66 (2017).

5. Almeida, A. *et al.* A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nat. Biotechnol.* **39**, 105–114 (2021).

6. Mahmoudabadi, G. & Phillips, R. A comprehensive and quantitative exploration of thousands of viral genomes. *eLife* **7**, e31955 (2018).

7. Nayfach, S. *et al.* A genomic catalog of Earth's microbiomes. *Nat. Biotechnol.* **39**, 499–509 (2021).

8. Kowarsky, M. *et al.* Numerous uncharacterized and highly divergent microbes which colonize humans are revealed by circulating cell-free DNA. *Proc. Natl. Acad. Sci.* **114**, 9623–9628 (2017).

9. Dickson, R. P., Erb-Downward, J. R., Martinez, F. J. & Huffnagle, G. B. The Microbiome and the Respiratory Tract. *Annu. Rev. Physiol.* **78**, 481–504 (2016).

10. Turnbaugh, P. J. *et al.* A core gut microbiome in obese and lean twins. *Nature* **457**, 480–

484 (2009).

11. Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes.

Nature **490**, 55–60 (2012).

12. Cryan, J. F., O'Riordan, K. J., Sandhu, K., Peterson, V. & Dinan, T. G. The gut microbiome

in neurological disorders. *Lancet Neurol.* **19**, 179–194 (2020).

13. Sharon, G. *et al.* Human Gut Microbiota from Autism Spectrum Disorder Promote

Behavioral Symptoms in Mice. *Cell* **177**, 1600-1618.e17 (2019).

14. Asnicar, F. *et al.* Microbiome connections with host metabolism and habitual diet from 1,098

deeply phenotyped individuals. *Nat. Med.* **27**, 321–332 (2021).

15. Jie, Z. *et al.* The gut microbiome in atherosclerotic cardiovascular disease. *Nat. Commun.* **8**,

845 (2017).

16. Fujimura, K. E. *et al.* Neonatal gut microbiota associates with childhood multisensitized

atopy and T cell differentiation. *Nat. Med.* **22**, 1187–1191 (2016).

17. Wolter, M. *et al.* Leveraging diet to engineer the gut microbiome. *Nat. Rev. Gastroenterol.*

Hepatol. 1–18 (2021) doi:10.1038/s41575-021-00512-7.

18. Poore, G. D. *et al.* Microbiome analyses of blood and tissues suggest cancer diagnostic

approach. *Nature* **579**, 567–574 (2020).

19. Dohlman, A. B. *et al.* The cancer microbiome atlas: a pan-cancer comparative analysis to

distinguish tissue-resident microbiota from contaminants. *Cell Host Microbe* **29**, 281-298.e5

(2021).

20. Narunsky-Haziza, L. *et al.* Pan-cancer analyses reveal cancer-type-specific fungal ecologies

and bacteriome interactions. *Cell* **185**, 3789-3806.e17 (2022).

21. Nejman, D. *et al.* The human tumor microbiome is composed of tumor type-specific

intracellular bacteria. *Science* **368**, 973–980 (2020).

22. Riquelme, E. *et al.* Tumor Microbiome Diversity and Composition Influence Pancreatic

Cancer Outcomes. *Cell* **178**, 795-806.e12 (2019).

23. Ghaddar, B. *et al.* Tumor microbiome links cellular programs and immunity in pancreatic

cancer. *Cancer Cell* **40**, 1240-1253.e5 (2022).

24. Aykut, B. *et al.* The fungal mycobiome promotes pancreatic oncogenesis via activation of

MBL. *Nature* **574**, 264–267 (2019).

25. Tzeng, A. *et al.* Human breast microbiome correlates with prognostic features and

immunological signatures in breast cancer. *Genome Med.* **13**, 60 (2021).

26. Hieken, T. J. *et al.* The Microbiome of Aseptically Collected Human Breast Tissue in Benign

and Malignant Disease. *Sci. Rep.* **6**, 30751 (2016).

27. Shao, D. *et al.* Microbial characterization of esophageal squamous cell carcinoma and

gastric cardia adenocarcinoma from a high-risk region of China. *Cancer* **125**, 3993–4002

(2019).

28. THE TABULA SAPIENS CONSORTIUM. The Tabula Sapiens: A multiple-organ, single-cell

transcriptomic atlas of humans. *Science* **376**, eabl4896 (2022).

29. Forster, S. C. *et al.* A human gut bacterial genome and culture collection for improved

metagenomic analyses. *Nat. Biotechnol.* **37**, 186–192 (2019).

30. HOMD :: Human Oral Microbiome Database. https://www.homd.org/.

31. Davis, J. J. *et al.* The PATRIC Bioinformatics Resource Center: expanding data and

analysis capabilities. *Nucleic Acids Res.* **48**, D606–D612 (2020).

32. Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Research* **10**, 33 (2021).

33. Smith, T. S., Heger, A. & Sudbery, I. UMI-tools: Modelling sequencing errors in Unique

Molecular Identifiers to improve quantification accuracy. *Genome Res.* gr.209601.116

(2017) doi:10.1101/gr.209601.116.

34. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

35. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment

search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

36. Salter, S. J. *et al.* Reagent and laboratory contamination can critically impact sequence-
based microbiome analyses. *BMC Biol.* **12**, 87 (2014).

37. Woellmer, A. & Hammerschmidt, W. Epstein-Barr virus and host cell methylation:
Regulation of latency, replication and virus reactivation. *Curr. Opin. Virol.* **3**, 260–265
(2013).

38. Tierney, R. J., Shannon-Lowe, C. D., Fitzsimmons, L., Bell, A. I. & Rowe, M. Unexpected
patterns of Epstein–Barr virus transcription revealed by a High throughput PCR array for
absolute quantification of viral mRNA. *Virology* **474**, 117–130 (2015).

39. Forbes, J. D., Bernstein, C. N., Tremlett, H., Van Domselaar, G. & Knox, N. C. A Fungal
World: Could the Gut Mycobiome Be Involved in Neurological Disease? *Front. Microbiol.* **9**,
3249 (2019).

40. Qin, J. *et al.* A human gut microbial gene catalogue established by metagenomic
sequencing. *Nature* **464**, 59–65 (2010).

41. Al Bataineh, M. T. *et al.* Revealing links between gut microbiome and its fungal community
in Type 2 Diabetes Mellitus among Emirati subjects: A pilot study. *Sci. Rep.* **10**, 9624
(2020).

42. Milo, R. & Phillips, R. *Cell Biology by the numbers*. (Garland Science, 2015).
doi:10.1201/9780429258770.

43. Schaechter, M., MaalØe, O. & Kjeldgaard, N. O. Y. 1958. Dependency on Medium and
Temperature of Cell Size and Chemical Composition during Balanced Growth of Salmonella
typhimurium. *Microbiology* **19**, 592–606.

44. Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T. Interdependence of cell
growth and gene expression: origins and consequences. *Science* **330**, 1099–1102 (2010).

45. Milo, R., Jorgensen, P., Moran, U., Weber, G. & Springer, M. BioNumbers--the database of
key numbers in molecular and cell biology. *Nucleic Acids Res.* **38**, D750-753 (2010).

46. Grimm, C., Bartuli, J. & Fischer, U. Cytoplasmic gene expression: lessons from poxviruses.

Trends Biochem. Sci. (2022) doi:10.1016/j.tibs.2022.04.010.

47. Wylie, S. J., Luo, H., Li, H. & Jones, M. G. K. Multiple polyadenylated RNA viruses detected

in pooled cultivated and wild plant samples. *Arch. Virol.* **157**, 271–284 (2012).

48. Hajnsdorf, E. & Kaberdin, V. R. RNA polyadenylation and its consequences in prokaryotes.

Philos. Trans. R. Soc. B Biol. Sci. **373**, 20180166 (2018).

49. Maes, A. *et al.* Landscape of RNA polyadenylation in E. coli. *Nucleic Acids Res.* **45**, 2746–

2756 (2017).

50. Ozsolak, F. *et al.* Comprehensive Polyadenylation Site Maps in Yeast and Human Reveal

Pervasive Alternative Polyadenylation. *Cell* **143**, 1018–1029 (2010).

51. Mayer, A. *et al.* Native Elongating Transcript Sequencing Reveals Human Transcriptional

Activity at Nucleotide Resolution. *Cell* **161**, 541–554 (2015).

52. Chukkapalli, S. S. *et al.* Invasion of Oral and Aortic Tissues by Oral Spirochete Treponema

denticola in ApoE−/− Mice Causally Links Periodontal Disease and Atherosclerosis. *Infect.*

Immun. **82**, 1959–1967 (2014).

53. Santos, P. B. do R. E. dos *et al.* Review- The periodontal pathogen Treponema denticola:

an atherosclerosis risk factor. *Res. Soc. Dev.* **10**, e25810111637–e25810111637 (2021).

54. Okuda, K. *et al.* Detection of Treponema denticola in Atherosclerotic Lesions. *J. Clin.*

Microbiol. (2001) doi:10.1128/JCM.39.3.1114-1117.2001.

55. Schenkein, H. A., Papapanou, P. N., Genco, R. & Sanz, M. Mechanisms underlying the

association between periodontitis and atherosclerotic disease. *Periodontol. 2000* **83**, 90–

106 (2020).

56. Young, L. S., Yap, L. F. & Murray, P. G. Epstein–Barr virus: more than 50 years old and still

providing surprises. *Nat. Rev. Cancer* **16**, 789–802 (2016).

57. Dunmire, S. K., Verghese, P. S. & Balfour, H. H. Primary Epstein-Barr virus infection. *J.*

Clin. Virol. **102**, 84–92 (2018).

58. Thorley-Lawson, D. A. EBV Persistence—Introducing the Virus. in *Epstein Barr Virus*

Volume 1: One Herpes Virus: Many Diseases (ed. Münz, C.) 151–209 (Springer

International Publishing, 2015). doi:10.1007/978-3-319-22822-8_8.

59. Smatti, M. K. *et al.* Epstein–Barr Virus Epidemiology, Serology, and Genetic Variability of

LMP-1 Oncogene Among Healthy Population: An Update. *Front. Oncol.* **8**, 211 (2018).

60. Bjornevik, K. *et al.* Longitudinal analysis reveals high prevalence of Epstein-Barr virus

associated with multiple sclerosis. *Science* (2022) doi:10.1126/science.abj8222.

61. Narita, Y. *et al.* Pin1 Interacts with the Epstein-Barr Virus DNA Polymerase Catalytic

Subunit and Regulates Viral DNA Replication. *J. Virol.* **87**, 2120–2127 (2013).

62. van Zyl, D. G., Mautner, J. & Delecluse, H.-J. Progress in EBV Vaccines. *Front. Oncol.* **9**,

104 (2019).

63. Chiu, S.-H. *et al.* Epstein-Barr Virus BALF3 Has Nuclease Activity and Mediates Mature

Virion Production during the Lytic Cycle. *J. Virol.* **88**, 4962–4975 (2014).

64. Wen, W. *et al.* Epstein-Barr Virus BZLF1 Gene, a Switch from Latency to Lytic Infection, Is

Expressed as an Immediate-Early Gene after Primary Infection of B Lymphocytes. *J. Virol.*

81, 1037–1042 (2007).

65. Dervan, E., Bhattacharyya, D. D., McAuliffe, J. D., Khan, F. H. & Glynn, S. A. Ancient

Adversary – HERV-K (HML-2) in Cancer. *Front. Oncol.* **11**, 1707 (2021).

66. Agoni, L., Guha, C. & Lenz, J. Detection of Human Endogenous Retrovirus K (HERV-K)

Transcripts in Human Prostate Cancer Cell Lines. *Front. Oncol.* **3**, 180 (2013).

67. Chan, S. M. *et al.* The HERV-K accessory protein Np9 controls viability and migration of

teratocarcinoma cells. *PLOS ONE* **14**, e0212970 (2019).

68. Grow, E. J. *et al.* Intrinsic retroviral reactivation in human preimplantation embryos and

pluripotent cells. *Nature* **522**, 221–225 (2015).

69. Wang, T. *et al.* Regulation of stem cell function and neuronal differentiation by HERV-K via

mTOR pathway. *Proc. Natl. Acad. Sci.* **117**, 17842–17853 (2020).

70. Adawi, D., Molin, G. & Jeppsson, B. Gut–liver axis. *HPB* **1**, 173–186 (1999).

71. Tripathi, A. *et al.* The gut–liver axis and the intersection with the microbiome. *Nat. Rev.*
Gastroenterol. Hepatol. **15**, 397–411 (2018).

72. Nguyen, S. *et al.* Bacteriophage Transcytosis Provides a Mechanism To Cross Epithelial
Cell Layers. *mBio* **8**, e01874-17.

73. Zhou, C.-B., Zhou, Y.-L. & Fang, J.-Y. Gut Microbiota in Cancer Immune Response and
Immunotherapy. *Trends Cancer* **7**, 647–660 (2021).

74. York, A. Gut microbiota sways response to cancer immunotherapy. *Nat. Rev. Microbiol.* **16**,
121–121 (2018).

75. Martel, C. de, Georges, D., Bray, F., Ferlay, J. & Clifford, G. M. Global burden of cancer
attributable to infections in 2018: a worldwide incidence analysis. *Lancet Glob. Health* **8**,
e180–e190 (2020).

76. Evans, E. L., Becker, J. T., Fricke, S. L., Patel, K. & Sherer, N. M. HIV-1 Vif's Capacity To
Manipulate the Cell Cycle Is Species Specific. *J. Virol.* **92**, e02102-17.

77. Fan, Y., Sanyal, S. & Bruzzone, R. Breaking Bad: How Viruses Subvert the Cell Cycle.
Front. Cell. Infect. Microbiol. **8**, 396 (2018).

78. Gagnaire, A., Nadel, B., Raoult, D., Neefjes, J. & Gorvel, J.-P. Collateral damage: insights
into bacterial mechanisms that predispose host cells to cancer. *Nat. Rev. Microbiol.* **15**,
109–128 (2017).

79. Russell, A. B., Trapnell, C. & Bloom, J. D. Extreme heterogeneity of influenza virus infection
in single cells. *eLife* **7**, e32303 (2018).

80. Zanini, F., Pu, S.-Y., Bekerman, E., Einav, S. & Quake, S. R. Single-cell transcriptional
dynamics of flavivirus infection. *eLife* **7**, e32942 (2018).

81. Xue, Y. *et al.* A single-parasite transcriptional atlas of Toxoplasma Gondii reveals novel
control of antigen expression. *eLife* **9**, e54129 (2020).

82. Weynberg, K. D., Wood-Charlson, E. M., Suttle, C. A. & van Oppen, M. J. H. Generating
viral metagenomes from the coral holobiont. *Front. Microbiol.* **5**, 206 (2014).

83. Wang, Q. *et al.* Computational methods and challenges in analyzing intratumoral

microbiome data. *Trends Microbiol.* S0966842X23000306 (2023)

doi:10.1016/j.tim.2023.01.011.

84. Robinson, W. *et al.* scRNA-seq analysis of colon and esophageal tumors uncovers

abundant microbial reads in myeloid cells undergoing proinflammatory transcriptional

alterations. 2020.05.14.096230 Preprint at https://doi.org/10.1101/2020.05.14.096230

(2023).

85. Lloréns-Rico, V. *et al.* Clinical practices underlie COVID-19 patient respiratory microbiome

composition and its interactions with the host. *Nat. Commun.* **12**, 6243 (2021).

86. Walker, M. A. *et al.* GATK PathSeq: a customizable computational tool for the discovery and

identification of microbial sequences in libraries from eukaryotic hosts. *Bioinformatics* **34**,

4287–4289 (2018).

87. Zhu, K. *et al.* Strain level microbial detection and quantification with applications to single

cell metagenomics. *Nat. Commun.* **13**, 6430 (2022).

88. Morgulis, A. & Agarwala, R. SRPRISM (Single Read Paired Read Indel Substitution

Minimizer): an efficient aligner for assemblies with explicit guarantees. *GigaScience* **9**,

giaa023 (2020).

89. Kostic, A. D. *et al.* PathSeq: software to identify or discover microbes by deep sequencing of

human tissue. *Nat. Biotechnol.* **29**, 393–396 (2011).

90. Shen, W. & Ren, H. TaxonKit: A practical and efficient NCBI taxonomy toolkit. *J. Genet.*

Genomics **48**, 844–850 (2021).

91. Ye, S. H., Siddle, K. J., Park, D. J. & Sabeti, P. C. Benchmarking Metagenomics Tools for

Taxonomic Classification. *Cell* **178**, 779–794 (2019).

92. McIntyre, A. B. R. *et al.* Comprehensive benchmarking and ensemble approaches for

metagenomic classifiers. *Genome Biol.* **18**, 182 (2017).

93. Bazinet, A. L., Ondov, B. D., Sommer, D. D. & Ratnayake, S. BLAST-based validation of

metagenomic sequence assignments. *PeerJ* **6**, e4892 (2018).

94. Wood, D. E. & Salzberg, S. L. Kraken: ultrafast metagenomic sequence classification using

exact alignments. *Genome Biol.* **15**, R46 (2014).

95. Liebhoff, A.-M. *et al.* Pathogen detection in RNA-seq data with Pathonoia. *BMC*

Bioinformatics **24**, 53 (2023).

96. Jurado-Rueda, F. *et al.* Benchmarking of microbiome detection tools on RNA-seq synthetic

databases according to diverse conditions. *Bioinforma. Adv.* **3**, vbad014 (2023).

97. O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: current status,

taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44**, D733-745 (2016).

98. Huntemann, M. *et al.* The standard operating procedure of the DOE-JGI Microbial Genome

Annotation Pipeline (MGAP v.4). *Stand. Genomic Sci.* **10**, 86 (2015).

99. Schulz, F. *et al.* Hidden diversity of soil giant viruses. *Nat. Commun.* **9**, 4881 (2018).

100.   González-Pech, R. A., Stephens, T. G. & Chan, C. X. Commonly misunderstood

parameters of NCBI BLAST and important considerations for users. *Bioinformatics* **35**,

2697–2698 (2019).

101.   Hung, J.-H. & Weng, Z. Sequence Alignment and Homology Search with BLAST and

ClustalW. *Cold Spring Harb. Protoc.* **2016**, pdb.prot093088 (2016).

102.   Hiseni, P., Rudi, K., Wilson, R. C., Hegge, F. T. & Snipen, L. HumGut: a comprehensive

human gut prokaryotic genomes collection filtered by metagenome data. *Microbiome* **9**, 165

(2021).

103.   Tiew, P. Y. *et al.* The Mycobiome in Health and Disease: Emerging Concepts,

Methodologies and Challenges. *Mycopathologia* **185**, 207–231 (2020).

104.   Nash, A. K. *et al.* The gut mycobiome of the Human Microbiome Project healthy cohort.

Microbiome **5**, 153 (2017).

105.   Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol. **20**, 257 (2019).

106.   Chawla, N. V., Bowyer, K. W., Hall, L. O. & Kegelmeyer, W. P. SMOTE: synthetic

2154   minority over-sampling technique. *J. Artif. Intell. Res.* **16**, 321–357 (2002).

2155   107.   Lopes, M. & Bontempi, G. *On the Null Distribution of the Precision and Recall Curve*.

2156   337 (2014). doi:10.1007/978-3-662-44851-9_21.

2157   108.   Zou, Y. *et al.* 1,520 reference genomes from cultivated human gut bacteria enable

2158   functional microbiome analyses. *Nat. Biotechnol.* **37**, 179–185 (2019).

2159   109.   Chen, I.-M. A. *et al.* IMG/M v.5.0: an integrated data management and comparative

2160   analysis system for microbial genomes and microbiomes. *Nucleic Acids Res.* **47**, D666–

2161   D677 (2019).

2162   110.   Pasolli, E. *et al.* Extensive Unexplored Human Microbiome Diversity Revealed by Over

2163   150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell* **176**,

2164   649-662.e20 (2019).

2165   111.   Carrero, P. *et al.* Report of six cases of human infection by Serratia plymuthica. *J. Clin.*

2166   *Microbiol.* **33**, 275–276 (1995).

2167   112.   Al-Sardi, M., Radwan, H., Itbaileh, A. B. & AlMusa, Z. Leifsonia Species Bacteremia in a

2168   Hemodialysis Patient: A Difficult-to-Identify Organism. *Cureus* **13**, e17994.

2169   113.   Rickettsia Japonica - an overview | ScienceDirect Topics.

2170   https://www.sciencedirect.com/topics/medicine-and-dentistry/rickettsia-japonica.

2171   114.   Bai, H. *et al.* Lysobacter oculi sp. nov., isolated from human Meibomian gland

2172   secretions. *Antonie Van Leeuwenhoek* **113**, 13–20 (2020).

2173   115.   Shkoporov, A. N. *et al.* The Human Gut Virome Is Highly Diverse, Stable, and Individual

2174   Specific. *Cell Host Microbe* **26**, 527-541.e5 (2019).

2175   116.   Franzosa, E. A. *et al.* Identifying personal microbiomes using metagenomic codes. *Proc.*

2176   *Natl. Acad. Sci.* (2015) doi:10.1073/pnas.1423854112.

2177   117.   Integrative HMP (iHMP) Research Network Consortium. The Integrative Human

2178   Microbiome Project: dynamic analysis of microbiome-host omics profiles during periods of

human health and disease. *Cell Host Microbe* **16**, 276–289 (2014).

118.   Glass, J. I. *et al.* The complete sequence of the mucosal pathogen Ureaplasma

urealyticum. *Nature* **407**, 757–762 (2000).

119.   Lindholm, M., Min Aung, K., Nyunt Wai, S. & Oscarsson, J. Role of OmpA1 and OmpA2

in Aggregatibacter actinomycetemcomitans and Aggregatibacter aphrophilus serum

resistance. *J. Oral Microbiol.* **11**, 1536192 (2019).