Title:

The integrity of dopaminergic and noradrenergic brain regions is associated with different aspects of late-
life memory performance

Authors:

Martin J. Dahl [1,2,*], Shelby L. Bachman [2], Shubir Dutt [2,3], Sandra Düzel [1], Nils C. Bodammer [1], Ulman
Lindenberger [1,4], Simone Kühn [4,5,6], Markus Werkle-Bergner [1], & Mara Mather [2]

Affiliations:

[1] Center for Lifespan Psychology, Max Planck Institute for Human Development, 14195 Berlin, Germany

[2] Leonard Davis School of Gerontology, University of Southern California, 90089 Los Angeles, CA, USA

[3] Department of Psychology, University of Southern California, 90089, Los Angeles, CA, USA

[4] Max Planck UCL Centre for Computational Psychiatry and Ageing Research, London, England, and
Berlin, Germany

[5] Lise Meitner Group for Environmental Neuroscience, Max Planck Institute for Human Development,
14195 Berlin, Germany

[6] Department of Psychiatry and Psychotherapy, University Clinic Hamburg-Eppendorf, Hamburg,
Germany

* Corresponding author:

MJD (dahl@mpib-berlin.mpg.de)

Abstract:

Researchers have identified changes in dopaminergic neuromodulation as playing a key role in adult memory decline. Facilitated by technical advancements, recent research has also implicated noradrenergic neuromodulation in shaping late-life memory development. However, it is not yet clear whether these two neuromodulators have distinct roles in age-related cognitive changes. Combining longitudinal high-resolution magnetic resonance imaging of the dopaminergic substantia nigra–ventral tegmental area (SN–VTA) and the noradrenergic locus coeruleus (LC), we found that dopaminergic and noradrenergic integrity are differentially associated with individual differences in younger (n = 69) and older adults' (n = 251) memory performance. While LC integrity was related to better episodic memory across several memory tasks, SN–VTA integrity was linked to working memory. Moreover, consistent with their dense interconnection and a largely shared biosynthesis, dopaminergic and noradrenergic brain regions' integrity were positively related, and correlated with medial temporal lobe volumes. Longitudinally, we found that older age was associated with more-negative change in SN–VTA and LC integrity (time point 1–time point 2; mean delay ~1.9 years). Importantly, changes in LC integrity reliably predicted future episodic memory performance (at time point 3). These findings support the feasibility of in-vivo indices for catecholaminergic integrity with potential clinical utility, given the degeneration of both neuromodulatory systems in several age-associated diseases. Moreover, they highlight differential roles of dopaminergic and noradrenergic neuromodulatory nuclei in late-life cognitive decline.

Introduction:

Our memory fades as we age [1]. On average, old age is characterized by impaired abilities to retain and manipulate information over shorter time scales—termed working memory [2–4]—and to recollect past experiences within their temporal and spatial context—called episodic memory [1,5,6]. On a neural level, senescent declines in memory have been linked to dopaminergic neuromodulation [7–9] and, more recently, also to noradrenergic neuromodulation [10–12]. The degeneration of catecholaminergic (i.e. dopaminergic and noradrenergic) systems also is a core feature of age-related pathologies, such as Alzheimer's and Parkinson's disease [11,13–17] that are characterized by amnestic impairments [18,19]. However, investigations disentangling the contribution of the two neuromodulators to human memory in aging and disease are scarce.

Neuromodulators are a group of neurochemicals synthesized in circumscribed subcortical nuclei from which they are released throughout the brain via widely branching axonal projections [20]. Dopaminergic neurons are based mainly in the midbrain substantia nigra–ventral tegmental area (SN–VTA) [21], whereas noradrenergic neurons are primarily found in the brainstem locus coeruleus (LC) [22].

There are several mechanistic accounts linking dopaminergic and noradrenergic neuromodulation to aging memory. Computational models posit that catecholamines modulate the input–output relation of neurons (i.e., gain change) which increases the signal-to-noise ratio in neural processing [23] and influences cognition [24–26]. Age-related neurodegeneration of dopaminergic and noradrenergic nuclei thus results in noisier neural information processing (i.e., gain reduction) [8]. Specifically, declining catecholaminergic drive with increasing age is hypothesized to lead to less distinctive cortical representations and senescent memory decline [8,27].

A second mechanism linking dopaminergic and noradrenergic neuromodulation to aging memory is their modulation of prefrontal processing [28]. Lateral prefrontal circuits can represent external stimuli in the absence of sensory stimulation, even in the face of distractors, by means of persistently firing delay cells [29]. Catecholaminergic inputs orchestrate recurrent activity in delay cell circuits that is essential for higher-order cognitive functions, such as working memory [30]. Specifically, the stimulation of dopaminergic D1-receptors and noradrenergic α_{2a}-receptors boosts prefrontal delay activity with an inverted-u dose–response curve [12]. Age-related memory deficits, in turn, have been associated with reduced delay cell firing, which could be partially restored by catecholaminergic drugs [31–34].

Finally, dopamine and noradrenaline modulate hippocampal long-term potentiation (LTP) and long-term depression (LTD) [35–39], which are critical for synaptic plasticity and memory. Initial accounts proposed a ventral tegmental area–hippocampal circuit by which neuromodulatory inputs facilitate the consolidation of salient experiences [35,38,40]. Interestingly, more recent investigations indicate that while the SN–VTA and LC both project to the dorsal hippocampus, the latter sends denser inputs [41–43]. LC neurons also produce dopamine as biosynthetic precursor of noradrenaline [44], and can co-release both catecholamines to modulate hippocampal synaptic plasticity and memory [41,42,45–47]. Older age is characterized by impaired hippocampal plasticity [48,49], likely exacerbated by deficient catecholaminergic innervation from the LC and SN–VTA [50].

Taken together, dopaminergic and noradrenergic neuromodulation mechanistically sculpts senescent memory via several pathways, including gain modulation [8,10], frontal delay activity[29,31], and hippocampal synaptic plasticity [35,46]. Notably, these mechanisms are specified for dopaminergic *and* noradrenergic neuromodulation (gain modulation [23]; delay activity [12]; synaptic plasticity [46]). However, research sampling a broad array of cognitive tasks to identify *unique* associations with late-life memory is lacking. That is, while animal research has demonstrated considerable overlap of

catecholamines at the neural level, the question remains how much dopaminergic and noradrenergic

nuclei overlap in their association with behavior.

Comparative studies of catecholaminergic systems in humans were long hampered by technical

challenges in reliable non-invasive assessments of the small subcortical nuclei [51,52]. However, recent

advances in high-resolution magnetic resonance imaging (MRI) reveal the SN–VTA [17,53] and LC

[54,55] as focal hyperintensity on MRI scans. Multimodal post-mortem validation studies suggest this

hyperintensity as a marker for catecholaminergic neurons [56–58]. Neuromelanin, a catecholamine-

derived paramagnetic pigment accumulating in the LC and SN–VTA [59,60], presumably contributes to

the MRI contrast of the nuclei [55,57,58,61,62]. However, also other factors likely play a role, such as the

large cellular bodies of catecholaminergic neurons [63,64] that results in a high abundance of ions and

water protons [64] as well as a lower macromolecular fraction [65,66]. Importantly, first in-vivo studies

suggest an association between the MRI intensity of catecholaminergic nuclei and their functionality [57].

Furthermore, investigations in clinical populations confirm the validity of the imaging approach [17,67–

71].

Catecholaminergic nuclei are among the first brain structures to accumulate pathologies in age-associative

diseases like Parkinson's and Alzheimer's [13,72,73] and show severe degeneration over the course of

these diseases [11,14,16]. In line with this, LC [67–69] and SN–VTA [17,70,71] imaging using dedicated

MRI sequences (i.e., Magnetization Transfer [MT], Fast Spin Echo [FSE]) reveals reduced

hyperintensities in patients relative to controls. In healthy lifespan samples, initial cross-sectional

evidence reveals a negative quadratic relationship between age and catecholaminergic hyperintensity [74–

76], whereby lower contrast with advancing age might be linked to impending pathology [67,77,78].

Taken together, recent advances in imaging techniques open the door for comparative non-invasive

assessments of catecholaminergic nuclei integrity, which are sensitive for age and disease-related changes

[17,53,55,70,71].

In the present study, we took advantage of these new imaging approaches to attempt to disentangle the

relative contribution of declining dopaminergic and noradrenergic neuromodulation to aging cognition.

We repeatedly assessed cognitive performance and high-resolution MRI in large samples of younger and

older adults across multiple time points [79–82]. Furthermore, we leveraged latent-variable modeling of

multimodal imaging [83,84] and comprehensive cognitive assessments [84,85] to decrease measurement

error and increase generalizability [86]. In sum, the goal of this study was to extend our knowledge about

the respective roles of dopaminergic and noradrenergic neuromodulation in late-life memory decline.

Results:

Locus coeruleus and substantia nigra–ventral tegmental area intensity shows high agreement across imaging modalities

We applied a validated semi-automatic procedure [87–91] to extract intensity information of catecholaminergic nuclei from different imaging sequences (FSE, MT+, and MT–; see Methods)—by standardizing MRI intensity in the LC and SN–VTA based on the intensity in an adjacent white matter reference region [54,55,57] (see Figures 1 and 2). Based on earlier post-mortem validations [56–58,68], we interpret individual differences in standardized MRI intensity as proxy for the integrity of catecholaminergic nuclei. Next, we leveraged an established factor structure [87] to integrate intensity information across hemispheres for each imaging sequence and age group (see Figure S1).

Previous in-vivo studies of catecholaminergic nuclei relied on different imaging approaches (mostly MT+ and FSE; [17,53–55]), but there are few comparisons between these MRI sequences, limiting cross-study comparability. Contrasting LC and SN–VTA estimates, we found strong differences across MRI sequences in their average intensity, $\Delta\chi^2(df = 2) = 693.55$; $p < 0.001$ for older adults' LC; $\Delta\chi^2(df = 1) = 657.37$; $p < 0.001$ for older adults' SN–VTA. That is, standardized to a reference region, catecholaminergic nuclei appeared brightest in the MT+, followed by the FSE, and finally the MT– sequences, for older adults' LC (mean [standard error]): MT+, 25.816 (0.304); FSE, 20.144 (0.37); MT–, 6.425 (0.194); for older adults' SN–VTA: MT+, 19.523 (0.235); MT–, 2.979 (0.135). Crucially, despite these mean differences, intensities were highly correlated across imaging modalities, $r = 0.43–0.621$; $p < 0.001$ for older adults' LC; $r = 0.503$; $p < 0.001$ for older adults' SN–VTA (see Figure 3 and S6), indicating that these sequences provide convergent information about the same underlying construct (i.e., catecholaminergic nuclei).

We thus aggregated the information shared across imaging modalities by estimating multimodal latent factors expressing LC integrity and SN–VTA integrity (see Figure S2; for comparable approaches and a discussion, see [84] and [83,92]). Such latent variables capture the commonalities across scan modalities while removing the modality-specific measurement error and hence increase statistical power to detect true effects [86]. Model visualizations, model fit, and younger adults' findings are reported in the supplementary information (see Figures S1–2, S4–5, and [93]).

Taken together, we extracted the intensity of the LC and SN–VTA from different MRI sequences sensitive for catecholaminergic nuclei. We found a high agreement in intensities across imaging modalities and thus aggregated across sequences to obtain individual integrity estimates for the two catecholaminergic nuclei.

Figure 1. LC and SN–VTA-sensitive magnetic resonance imaging sequences. Hyperintensities corresponding to the (**a**) LC and (**b**) SN–VTA are indicated by red arrows on axial slices of group templates based on a Fast Spin Echo (FSE) sequence, and a Magnetization Transfer sequence, acquired once with a dedicated magnetic saturation pulse (MT+) and once without, resulting in a proton density image (MT−). A Magnetization Prepared Gradient-Echo (MPRAGE) sequence template shows the location of a previously-established LC [67] and SN–VTA [57] volume of interest (red overlays). Note that our FSE sequence only covered the brainstem. All templates were registered to MNI152 0.5mm linear space and are available at [93]. LC, locus coeruleus; SN–VTA, substantia nigra–ventral tegmental area.

162
Figure 2. Normalized intensity in the LC, SN–VTA, and corresponding white-matter reference regions.
Hyperintensities corresponding to the (**a**) LC and (**b**) SN–VTA are evident in group templates based on a Fast Spin
Echo (FSE) sequence, and a Magnetization Transfer sequence, acquired once with a dedicated magnetic saturation
pulse (MT+) and once without, resulting in a proton density image (MT–). LC and reference volumes of interest are
taken from [67], SN–VTA and reference volumes of interest are based on [57]. The reference regions are located
anterior of the LC and SN–VTA, respectively. In the visualization, they can be seen as rectangular shape to the right
of the LC (i.e., the two red columns) as well as in front of the SN–VTA (i.e., the red curved shape). Note that our
FSE sequence only covered the brainstem. LC, locus coeruleus; SN–VTA, substantia nigra–ventral tegmental area.
171
172

173
Figure 3. Younger and older adults' LC intensities are correlated across imaging modalities—a marker for their
agreement—and across time points—a marker for their stability. Visualized data are based on the statistical model
depicted in Figure S12. For the same analyses using SN–VTA data, see Figures S16 and S18. Note, the diagonal
shows LC intensity, standardized across all sequences and time points, to facilitate comparing intensity distributions.
Imaging sequences included a Fast Spin Echo (FSE) sequence, and a Magnetization Transfer sequence, acquired
once with a dedicated magnetic saturation pulse (MT+) and once without, yielding a proton density image (MT−).
LC, locus coeruleus; TP, time point; SN–VTA, substantia nigra–ventral tegmental area.
181

Multimodal locus coeruleus and substantia nigra–ventral tegmental area integrity factors show high stability over time

Making use of the longitudinal nature of this dataset, we next explored the stability of our integrity estimates over time (TP1–TP2; mean delay ~1.9 (SD 0.7) years), as a proxy for their reliability [94]. Longitudinal studies investigating the reliability of imaging sequences for catecholaminergic nuclei are sparse. Thus, as a reference, we first assessed the stability of the modality-specific intensity factors and found evidence for an intermediate stability, for younger and older adults' LC: MT+, $r = 0.6$; $p < 0.001$; MT–, $r = 0.63$; $p < 0.001$; for younger and older adults' SN–VTA: MT+, $r = 0.66$; $p < 0.001$; MT–, $r = 0.18$; $p = 0.17$ (see Figures 3 and S12, S16, S18; FSE sequence only available for TP2 [see Methods]). If our multimodal integrity factors remove modality-specific measurement error [84], we should observe a higher stability across time points for the multimodal as compared to the modality-specific factors. Importantly, our multimodal LC and SN–VTA factors indeed evinced a higher stability, pointing to the benefits of the multimodal imaging approach, $r = 0.88$; $p < 0.001$ for younger and older adults' LC (test against mean modality-specific stability, i.e., $r = 0.615$, $z = 11.713$, $p < 0.001$); $r = 0.67$; $p < 0.001$ for younger and older adults' SN–VTA (test against mean modality-specific stability, i.e., $r = 0.42$, $z = 6.452$; $p < 0.001$) [95].

Figure 4. Multimodal LC and SN–VTA estimates show high stability over time points (mean delay ~ 1.9 years), $r = 0.88$; $p < 0.001$ for younger and older adults' LC (test against mean modality-specific stability, i.e., $r = 0.615$, $z = 11.713$; $p < 0.001$); $r = 0.67$; $p < 0.001$ for younger and older adults' SN–VTA (test against mean modality-specific stability, i.e., $r = 0.42$, $z = 6.452$; $p < 0.001$). LC, locus coeruleus; SN–VTA, substantia nigra–ventral tegmental area.

Locus coeruleus and substantia nigra–ventral tegmental area are associated with different aspects of late-life memory performance

Next, we cross-sectionally probed the behavioral implications of interindividual differences in LC and SN–VTA integrity, using data of TP2. For this, we leveraged a comprehensive cognitive battery and a previously established factor structure [85] to aggregate performance across several working memory, episodic memory, and fluid intelligence tasks and capture their shared variance on a latent level (see Figure S7). We observed strong age differences in each of the cognitive domains (see Figure 5), older relative to younger adults (mean age difference [standard error]): working memory, –2.265 (0.309); episodic memory, –2.287 (0.309); fluid intelligence, –2.073 (0.295); all $p < 0.001$. Importantly, however, when we added estimates for the catecholaminergic nuclei to the model, we found that higher LC and SN–VTA integrity were related to better performance (i.e., less age-related cognitive impairments) across domains, $\Delta\chi^2(df = 3) = 25.11$; $p < 0.001$ for older adults' LC; $\Delta\chi^2(df = 3) = 7.86$; $p = 0.049$ for older adults' SN–VTA (see S8).

Dopaminergic and noradrenergic neuromodulatory centers are densely interconnected [37,96]. In addition, dopamine is the biosynthetic precursor of noradrenaline [44], and indeed we detected a positive association between the structural metrics of the two neuromodulatory nuclei, $r = 0.25$; $\Delta\chi^2(df = 1) = 5.75$; $p = 0.017$ for older adults (cf. Figure 6 and see S8). However, while the LC and SN–VTA were positively coupled, they, differed in their association with late-life cognition, $\Delta\chi^2(df = 3) = 15.66$; $p = 0.001$ for older adults.

Follow-up analyses showed that higher LC integrity was related to better episodic memory performance, $r = 0.49$; $\Delta\chi^2(df = 1) = 21.44$; $p < 0.001$ for older adults (cf. Figure 6 and see S8) and that this association was specific: That is, LC's relation to cognition differed across domains (episodic memory, working memory, and fluid intelligence), $\Delta\chi^2(df = 2) = 10.64$; $p = 0.005$ for older adults. Moreover, the LC–episodic memory association was stronger than that of the SN–VTA and episodic memory, $\Delta\chi^2(df = 1) = 6.63$; $p = 0.01$ for older adults. Taken together, our findings show a general (i.e., task and imaging sequence independent), yet specific relation of LC and late-life episodic memory performance, corroborating and extending earlier work [67,68,87,97,98].

In contrast to the LC, higher SN–VTA integrity correlated with better late-life working memory, $r = 0.28$; $\Delta\chi^2(df = 1) = 6.76$; $p = 0.009$ for older adults (cf. Figure 6 and see S8). There was a numerical tendency for differential associations of SN–VTA integrity with performance in each of the cognitive domains (working memory, episodic memory, and fluid intelligence), $\Delta\chi^2(df = 2) = 5.73$; $p = 0.057$ for older adults. However, we did not observe reliable differences in the relation of the two neuromodulatory nuclei to working memory, $\Delta\chi^2(df = 1) = 2.01$; $p = 0.156$ for older adults. In sum, our findings suggest a differential association of the two neuromodulatory systems with late-life memory performance. While episodic memory was associated with LC integrity, SN–VTA integrity was related to working memory. Note that our model (see Figure S8) specified correlations between the latent catecholaminergic and cognitive factors. That is, it probed associations of one neuromodulatory system with cognition irrespective of the other neuromodulatory system. Thus, we additionally specified a second, statistically equivalent model in which we searched for unique associations with cognition for each catecholaminergic system while controlling for the respective other system, using a multiple regression approach (see [84] for a comparable approach; see Figure 6 and S9). Crucially, we again detected reliable LC–episodic memory, $r = 0.5$; $\Delta\chi^2(df = 1) = 19.55$; $p < 0.001$, and SN–VTA–working memory associations, $r = 0.28$; $\Delta\chi^2(df = 1) = 6.05$; $p = 0.014$, for older adults, underlining differential relations to cognition of the two

247 neuromodulatory nuclei. Model visualizations, model fit, and younger adults' findings are reported in the
248 supplementary information (see S8 and S9 and [93]).
249

250
251 ***Figure 5.*** Cross-sectional age differences in working memory, episodic memory, and fluid intelligence factors at
252 TP2 (for the statistical model, see Figure S7). Older adults show lower performance relative to younger adults across
253 cognitive domains (mean age difference [standard error]): working memory, –2.265 (0.309); episodic memory, –
254 2.287 (0.309); fluid intelligence, –2.073 (0.295); all *p* < 0.001. Raincloud plots based on [99].

Figure 6. Schematic pictorial rendition of a structural equation model probing the interrelation of catecholaminergic nuclei and medial temporal lobe volume (paths **a** and **b**) and their unique associations with late-life cognition (paths **c–e**) in older adults at TP2. Note that covariances among cognitive factors, intercepts, and error terms are omitted for clarity. For the full statistical model, see Figure S11. Task*, cognitive paradigm (indicator; [square]) for the respective cognitive domain (latent factor; [ellipse]); One-headed arrows, regression; Double-headed arrow, correlation. Note that the brightness of paths indicates their significance. Medial temporal lobe volumes only demonstrate a reliable association with episodic memory in a correlational model (see grey line in scatter plot **e**; cf. Figure S10), but not when controlling for catecholaminergic neuromodulation (see black line in scatter plot **e**). Locus coeruleus (LC) and substantia nigra–ventral tegmental area (SN–VTA) factors are derived from a Fast Spin Echo (FSE) sequence, and a Magnetization Transfer sequence, acquired once with a dedicated magnetic saturation pulse (MT+) and once without, resulting in a proton density image (MT–). Hipp, hippocampus volume; Parahipp, parahippocampus volume.

Locus coeruleus and substantia nigra–ventral tegmental area are associated with memory performance over and above medial temporal lobe volumes

Via direct projections, the LC and SN–VTA release catecholamines in memory-relevant brain regions, such as the medial temporal lobe [41,43,46], which improves retention performance [35,37,41,43,45,46]. Abnormally phosphorylated tau, an indicator of neurodegenerative diseases such as Alzheimer's, starts to accumulate early in life in catecholaminergic nuclei [13,100–102]. With advancing age, abnormal tau also appears in projection targets of the neuromodulatory nuclei, like the medial temporal lobe [13,101,103,104], which may facilitate degeneration [67,68,104,105]. Thus, as a control analysis, we also incorporated medial temporal lobe volumes in our models linking catecholaminergic nuclei and cognition (see Figure S10 and S11).

In a correlational model, we observed that the integrity of both catecholaminergic nuclei was positively associated with medial temporal lobe volumes, $r = 0.41$; $\Delta\chi^2(df = 1) = 27.45$; $p < 0.001$ for older adults' LC; $r = 0.23$; $\Delta\chi^2(df = 1) = 6.29$; $p = 0.012$ for older adults' SN–VTA (cf. Figure 6 and see S10), potentially indicating neuroprotective catecholaminergic effects [106,107]. In addition, higher medial temporal lobe volumes were related to better late-life episodic memory performance, $r = 0.33$; $\Delta\chi^2(df = 1) = 14.22$; $p < 0.001$ for older adults, in line with its role in memory processing [108].

Importantly, when we specified a second, multiple-regression model that searches for unique effects, we found that the LC was still reliably associated with episodic memory performance, $r = 0.43$; $\Delta\chi^2(df = 1) = 11.96$; $p < 0.001$, while medial temporal lobe volumes were not, $r = 0.16$; $\Delta\chi^2(df = 1) = 2.46$; $p = 0.117$ (see Figure 6 and S11), when accounting for the respective other regions (e.g., controlling for SN–VTA and medial temporal lobe volume when evaluating the association between LC and episodic memory). Similarly, the SN–VTA was related to working memory after controlling for medial temporal lobe volumes and LC integrity, $r = 0.28$; $\Delta\chi^2(df = 1) = 5.8$; $p = 0.016$. Taken together, our results suggest a robust association of catecholaminergic nuclei and memory that cannot be fully accounted for by medial temporal lobe volume.

Longitudinal changes in locus coeruleus integrity predict future episodic memory performance

Cross-sectional studies point to late-life differences in catecholaminergic nuclei [68,74–76,87,109], however, longitudinal data showing within-person changes are scarce. Thus, we combined imaging data of the two time points (TP1–TP2; mean delay ~1.9 years) to test for individual changes in LC and SN–VTA integrity estimates in later life [86,110].

First, we observed that change in the catecholaminergic nuclei was correlated across imaging modalities (MT+, MT−; no longitudinal FSE data available), suggesting that the different MR sequences pick up a common underlying construct (i.e., change in catecholaminergic nuclei), $r = 0.16$; $\Delta\chi^2(df = 1) = 6.09$; $p = 0.014$ for older adults' LC; $r = 0.13$; $\Delta\chi^2(df = 1) = 5.91$; $p = 0.015$ for older adults' SN–VTA (see S14 and S19). Thus, we again integrated across sequences to retrieve multimodal latent change factors for LC and SN–VTA integrity (see S15 and S20). For both catecholaminergic systems, we found reliable individual differences in change, $\Delta\chi^2(df = 1) = 6.09$; $p = 0.014$ for older adults' LC; $\Delta\chi^2(df = 1) = 5.91$; $p = 0.015$ for older adults' SN–VTA. However, there was no reliable mean change at the group level in either nucleus, ps > 0.1 in older adults. That is, we observed that older adults differed from one another in how their LC and SN–VTA changed over time—while some older adults showed increases in intensity ratios, others showed decreases (see Figure 7; cf. [68,74]).

Finally, to test the behavioral implications of these late-life changes in catecholaminergic nuclei, we
probed whether our neural change model (TP1–TP2) could be used to predict future cognition (at TP3;
using the previously established cognitive factor structure, cf. Figure S7). We additionally included
chronological age (at TP2) as a predictor, to test and account for potential age effects in rate of decline
(see S23 and S24).
In line with a late-life degeneration of neuromodulatory centers [7,10,11,68], we found more negative
change in the LC and SN–VTA with increasing age, $r = -0.18$; $\Delta\chi^2(df = 1) = 4.81$; $p = 0.028$ for older
adults' LC; $r = -0.29$; $\Delta\chi^2(df = 1) = 3.95$; $p = 0.047$ for older adults' SN–VTA. Moreover, we observed
that changes in LC integrity predicted subsequent episodic memory performance over and above
chronological age, $r = 0.23$; $\Delta\chi^2(df = 1) = 4.73$; $p = 0.03$ for older adults' LC (see Figure 7); $r = 0.27$;
$\Delta\chi^2(df = 1) = 1.55$; $p = 0.213$ for older adults' SN–VTA (see Figure S25). Taken together, our results are
in line with the proposition that a decline of the LC in later life is associated with diminished episodic
memory performance [10,68,87,111].
323

324
Figure 7. Longitudinal changes in LC intensity ratios and their association with age and future memory
performance. **a**, Numerically, older adults show more negative average change in LC intensity across time points as
compared to younger adults. MRI sequences include a Magnetization Transfer sequence, acquired once with a
dedicated magnetic saturation pulse (MT+) and once without, resulting in a proton density image (MT–). For the
Fast Spin Echo-sequence, only cross-sectional data are available. **b,** Schematic depiction of the structural equation
model probing the association of longitudinal change in multimodal LC integrity with future episodic memory
performance, accounting for chronological age. For the full model, see Figure S23. **c**, Scatter plots showing (1) more
negative LC change in older adults of higher age and (2) a prediction of future memory performance based on LC
change (controlling for chronological age). For comparable analyses using SN–VTA and working memory data, see

Figures S24 and S25. Raincloud plots based on [99]. LC, locus coeruleus; SN–VTA, substantia nigra–ventral
tegmental area.

Discussion:

This study sought to disentangle the effects of declining dopaminergic and noradrenergic neuromodulation on late-life memory. We took advantage of a multimodal imaging protocol and extensive cognitive assessments across several time points to contrast the behavioral implications of LC and SN–VTA integrity.

Our findings indicate that different imaging approaches for catecholaminergic nuclei (FSE, MT+, MT–) show a high agreement. Thus, we used latent-variable modeling to integrate across MRI modalities and retrieve multimodal LC and SN–VTA integrity factors that demonstrated a markedly increased reliability compared with their individual components. After establishing reliable in-vivo integrity proxies, we probed their associations with late-life cognition. We used an extensive neuropsychological test battery and a previously identified factor structure to demonstrate that these two catecholaminergic systems, although positively coupled, differ in their relationship three domains of aging cognition: episodic memory, working memory, and fluid intelligence.

We observed a general (i.e., task and imaging sequence independent), yet specific association of LC integrity and late-life episodic memory performance (i.e., stronger for episodic memory as compared to working memory and fluid intelligence). By contrast, SN–VTA integrity was linked to better working memory. Remarkably, both associations remained reliable even after accounting for the respective other neuromodulatory system and a key node in the memory network, the medial temporal lobe, suggesting robust effects.

Leveraging the longitudinal nature of this dataset, we also investigated late-life changes in the LC and SN–VTA over a delay of approximately two years. A principal finding was that older age was associated with more negative change in each catecholaminergic system, in line with the late-life degeneration of neuromodulatory centers. Moreover, we showed that changes in the LC predicted future episodic memory performance over and above chronological age. Taken together, this study suggests dopaminergic and noradrenergic neuromodulation play domain-specific roles in determining the trajectory of cognition in later life and provides novel insights into the neural basis of human senescent memory decline.

The loss of dopaminergic neuromodulation has long been recognized as crucial determinant of late-life cognitive deficits [7–9]. More recently, technical advances have also facilitated studies of the noradrenergic LC [55] that long seemed unattainable [51,52]. We used two types of imaging sequences (FSE, MT+) validated on postmortem specimen of the LC [56] and SN–VTA [57,58]. Neuromelanin, an insoluble catecholamine-derived pigment that traps metals and exhibits paramagnetic properties, is thought to contribute to the hyperintensity of catecholaminergic nuclei [17,55,57,58,61,62]. Moreover, the high density of water protons and paramagnetic ions in large catecholaminergic neurons has also been proposed as source of their MRI contrast [64,65]. In line with this, we provide the first quantification of the LC and SN–VTA based on a proton density-weighted sequence (i.e., without dedicated magnetization transfer preparation pulse; MT–). Magnetization transfer imaging studies frequently estimate a ratio score based on sequences with and without dedicated preparation pulse (i.e., MT+ and MT–) [112]. However, the sensitivity of our MT– sequence for the LC and SN–VTA suggests that this ratio would reduce the detectability of these nuclei. Notably, we observed that despite differences in mean contrast, LC and SN–VTA hyperintensities were correlated across all imaging modalities (FSE, MT+, MT–), suggesting that they provide convergent information about the same underlying construct (i.e., catecholaminergic nuclei). We thus leveraged our multimodal approach to estimate latent factors for catecholaminergic nuclei integrity based on the commonalities across imaging sequences while removing modality-specific measurement error [83,113]. Moreover, using data from both imaging time points, we show that

semiautomatic analyses of the intensity of catecholaminergic nuclei have high reliability, especially when
multimodal assessments are available [114].
The retention of salient experiences is enhanced by catecholamine release in the hippocampus which
facilitates synaptic plasticity and memory [35,37,46]. While the SN–VTA has long been attributed as
source of the memory-enhancing dopaminergic inputs [35,38,43], recent findings point to a denser
innervation by the LC that can provide noradrenergic as well as dopaminergic signals [41,45,46]. Here we
compared the association of the two catecholaminergic centers with an extensive set of tasks that are
thought to depend more (episodic memory) or less (working memory) on hippocampal processing
[108,115]. We observed that LC integrity was specifically related to late-life episodic memory
performance (as compared to working memory or fluid intelligence) and that this association was stronger
than the SN–VTA–episodic memory relationship. This finding might be explained by a LC-related
modulation of hippocampal synaptic plasticity [41,45,46], but our data cannot rule out other memory-
related mechanisms, such as gain modulation [8,23–26] and prefrontal delay activity [12,29]. Our
observations are supported by a series of large-scale in-vivo imaging studies that showed reliable LC–
cognition associations in aging [68,87,97,98], and particularly with episodic memory [68,87,116]. In
addition, they concord with a recent report linking anteromedial and superior substantia nigra intensity to
attentional performance [117], a cognitive concept overlapping with working memory [24].
Cross-sectional studies point to late-life differences in catecholaminergic nuclei [74–76,87,109]. Here we
provide the first characterization of late-life longitudinal changes in the LC and SN–VTA. In general
agreement with extrapolations from cross-sectional data, we found more negative change in both
catecholaminergic systems with increasing age. Subcortical neuromodulatory centers like the LC and the
SN–VTA are among the first sites to accumulate pathology in age-associated diseases like Alzheimer's
[13] and Parkinson's [72], and show severe degeneration with disease progression [16,118]. In
combination with earlier work suggesting lower catecholaminergic contrast with advancing age might be
linked to impending pathology [67,70,77,78,117], our findings may indicate subthreshold pathological
processes in a subset of our older participants. This interpretation is supported by our prediction of poorer
future memory performance in individuals with more negative LC change [111]. Accordingly, recent
meta-analyses demonstrate the efficacy of noradrenergic treatments in improving cognitive symptoms in
Alzheimer's disease [119]. Notably, MRI-indexed catecholaminergic integrity has been suggested as
useful tool for stratifying patients suffering for neurodegenerative diseases in clinical trials that include
noradrenergic treatments [55,120].
The present study highlights the utility of multimodal longitudinal assessments of
catecholaminergic nuclei to elucidate the neurobiological basis of senescent memory decline. We
dissociated the roles of the noradrenergic LC and dopaminergic SN–VTA in late life cognition. While the
former showed robust associations with current and future episodic memory performance, the latter
showed a relationship to better working memory performance. These findings point to differential roles of
dopaminergic and noradrenergic neuromodulation in late-life cognition with potential implications for
age-associated diseases like Alzheimer's and Parkinson's [13,68,72]. Accurate longitudinal assessments
of catecholaminergic nuclei may constitute promising early markers for predicting cognitive decline in
later life.
420

Methods:

Study design and participants:

Data were collected as part of the Berlin Aging Study-II (BASE-II), an ongoing longitudinal study that investigates neural, cognitive, physical, and social conditions related to successful aging (for more information, see https://www.base2.mpg.de/en and references [79–82].

Cognitive performance was assessed in three time periods (TP1–TP3) between 2013 and 2020 (TP1: 2013–2015; TP2: 2015–2016; TP3: 2018–2020) with a mean duration between cognitive assessments of 2.246 years (TP1–2; SD: 0.603) and 2.917 years (TP2–TP3; SD: 0.438).

A subset of BASE-II participants also underwent magnetic resonance imaging (MRI). Eligible participants had no history of neurological or psychiatric disorders, or head injuries, and did not take medication that may affect memory function. Imaging data were collected in two time periods (TP1, TP2) in temporal proximity to the cognitive assessments (mean delay between MRI waves 1.894 years; SD: 0.656). Participants were only considered for further analyses if at least one type of imaging sequence sensitive for dopaminergic or noradrenergic neuromodulatory centers was available (see below). For TP1, this corresponds to 288 participants out of a total of 488 participants with imaging data, whereas for TP2 this corresponds to 320 out of 323 participants with imaging data. Thus, our analyses included a total of 320 individual participants. While not all imaging sequences were available for all participants (see Table 1), 203 participants have relevant MRI data for both TP1 and TP2.

The final sample (n = 320) included 69 younger adults (22 female; mean age (SD): 32.705 (3.884) years [at TP2]) and 251 older adults (91 female; mean age (SD): 72.414 (4.045) years [at TP2]). Sample descriptives are reported in Table 2.

The cognitive and imaging assessments were approved by the institutional review boards of the Max Planck Institute for Human Development and the German Psychological Society (DGPS), respectively. Participants provided written informed consent and were reimbursed for their participation.

447   **Table 1:** Overview of participants with available imaging data

Magnetic resonance imaging sequence	Time period	Number of participants with relevant imaging data	Total number of participants with imaging data
Magnetization Transfer (MT+) *and* Proton Density (MT–)	1	288	448
Magnetization Transfer (MT+) *and* Proton Density (MT–)	2	260	323
Fast Spin Echo (FSE)	1	0	448
Fast Spin Echo (FSE)	2	316	323
Fast Spin Echo (FSE) *and* Magnetization Transfer (MT+) *and* Proton Density (MT–)	1	0	448
Fast Spin Echo (FSE) *or* Magnetization Transfer (MT+) *and* Proton Density (MT–)	1	288	448
Fast Spin Echo (FSE) *and* Magnetization Transfer (MT+) *and* Proton Density (MT–)	2	256	323
Fast Spin Echo (FSE) *or* Magnetization Transfer (MT+) *and* Proton Density (MT–)	2	320	323

Note: Relevant imaging data refers to participants with a whole-brain T_1-weighted (MPRAGE) sequence
as well as a sequence sensitive for dopaminergic or noradrenergic neuromodulatory centers
(Magnetization Transfer (MT+), Proton Density (MT–) or Fast Spin Echo (FSE)). By contrast, the total
number of participants with imaging data reflects all participants that underwent MRI, irrespective of
sequence type. The Magnetization Transfer and Proton Density sequences were acquired in succession in
each scan session (i.e., identical sequence, acquired once with and one without dedicated MT preparation
pulse). Thus, they are grouped in the table.

Table 2: Summary of sample descriptives for younger and older adults

	Younger adults (n = 69; 22 female)		Older adults (n = 251; 91 female)	
	Mean (SD)	Range	Mean (SD)	Range
Age (years; at time period 2)	32.705 (3.884)	25.414–40.203	72.414 (4.045)	62.534–83.162
Education (years) (available for n = 285)	14.242 (2.519)	10–18	14.08 (2.924)	7–18
Mini Mental State Examination (available for n = 204)	–	–	28.613 (1.336)	24–30

Note: Mini Mental State data was assessed around TP3, during a medical examination at a different visit
as part of the longitudinal Berlin Aging Study-II. *SD,* Standard Deviation.

Cognitive data assessment:

At TP1–TP3, cognition was tested using a comprehensive computerized battery probing key cognitive
functions. Performance was assessed in small groups of four to six participants. Cognitive test sessions
lasted approximately 3.7 h and included 16 tasks (at TP2), of these, three measured working memory,
four episodic memory, and three fluid intelligence [121]. While the exact composition of the cognitive
assessments changed across waves, the same tasks were used to measure working memory, episodic
memory and fluid intelligence at TP1–TP3. Only older adults were tested at TP3. For a detailed task
description, please refer to [121–123], below we provide a brief overview of the measures relevant to the
current analyses.

Working memory assessment:

Spatial Updating (abbreviated as *spatial-u*): Participants were shown a display of two to three 3 × 3 grids,
in each of which a blue dot was presented in one of the nine tiles. Participants were asked to memorize
the locations of the blue dots and mentally update them according to shifting operations that were
indicated by arrows appearing below the dots. Six updating operations were required before the 3 × 3
grids reappeared and participants indicated the end position of the blue dots (by mentally combining their
start position and the six shifting operations). We used the number of correct placements as an indicator
of working memory performance [121,123].
476
477   *Letter Updating* (abbreviated as *letter-u*): Participants were shown a sequence of 7–13 letters. Once the
478   presentation ended, they were asked to report, in correct order, the last three letters that were shown. The
479   number of correctly reported letters was used as a measure of working memory performance [121].
480
Number N-back (abbreviated as *number*): Three digits (1–9) were presented consecutively in three
adjacent cells, followed by the next sequence of three digits. Participants indicated by button press
whether the currently presented digit matched the digit shown three steps before [121,123]. We took
participants' accuracy as indicator of their working memory performance.

Episodic memory assessment:

Scene encoding (abbreviated as *scene*): Participants incidentally encoded 88 scene images by performing
indoor/outdoor judgments on each image. The encoding phase was followed by an old/new recognition
memory test, which included confidence judgments. Recognition memory was tested after a delay of
approximately 2.5 hours and served as episodic memory performance index (hits – false alarms)
[84,87,121].
491
Verbal learning and memory task (abbreviated as *list*): Participants first learned a 15-word list that was
presented via headphones. The task was composed of five learning trials each followed by a free recall
period in which participants entered the words they remembered via keyboard (trial 1–5; recall of learning
list). After these initial learning–recall cycles, participants were presented an interference list and their
delayed recall and recognition memory was assessed. The sum of correctly recalled words during the
learning–recall cycles (trial 1–5) served as episodic memory measure [84,87,121].
498

499 *Face–profession task* (abbreviated as *face*): Participants studied 45 pairs of face images and profession
500 words. The tasks consisted of an incidental encoding phase, a 2-minute distraction phase, and finally a
501 recognition memory task including old, new as well as rearranged face–profession pairs. Corrected
502 recognition memory scores for rearranged pairs was used as the performance index.
503
Object–location task (abbreviated as *object*): Participants encoded the location of 12 digital photographs
of real life objects on a 6 × 6 grid. After encoding, the objects reappeared next to the grid and participants
were instructed to reproduce their correct location by placing the items in the grid. The sum of correct
placements served as index of episodic memory.

Fluid intelligence assessment:
Practical problems (abbreviated as *problem*): Participants were sequentially presented eighteen items
depicting everyday problems (e.g., the hours of a bus timetable, a subway map etc.), in order of ascending
difficulty. For each of these problems, five response alternatives were provided and participants selected
the correct option by clicking on it. We took the sum of correctly solved problems as measure of fluid
intelligence [121,124].
514
515 *Figural analogies* (abbreviated as *analog*): Participants were instructed to draw analogies. They were
516 presented with twenty-two items in ascending difficulty that followed the format *A* is to *B* as *C* is to *X*?
517 Below each item, five response alternatives were presented and participants selected the correct option by
518 clicking on it. The sum of correctly given answers served as index of fluid intelligence [121,124].
519
520 *Letter series* (abbreviated as *letter*): Participants were shown twenty-two series of five letters, each ending
521 with a question mark (e.g., c-e-g-i-k-?). Each series followed a simple rule (e.g., +1, –1, +2; or +2, –1),
522 with increasing difficulty. Below each letter series, five response alternatives were presented and
523 participants selected the correct option by clicking on it. The sum of correct responses served as fluid
524 intelligence measure [121,124].
525
526 All tasks included practice blocks to familiarize participants with the instructions. Note that these tasks
527 have previously been used to estimate latent factors of working memory, episodic memory, and fluid
528 intelligence [121] (also see [84,87,125]).
529

Magnetic resonance imaging data assessment:

To investigate the associations of dopaminergic and noradrenergic integrity with late-life cognition,
younger and older participants underwent 3T-MRI at TP1 and TP2 (MAGNETOM TIM Trio, Siemens
Healthcare). Only those sequences used in the current analyses are described below. The imaging protocol
included three scans sensitive for the SN–VTA and LC—a Fast Spin Echo sequence (FSE; sometimes
also called Turbo Spin Echo [TSE]), and a Magnetization Transfer sequence, acquired once with a
dedicated magnetic saturation pulse (MT+) and once without, resulting in a proton density image (MT–).
Moreover, a Magnetization Prepared Gradient-Echo (MPRAGE) sequence, comparable to the ADNI
protocol (www.adni-info.org), was collected to facilitate coregistration to standard space and to estimate
volumes for regions of interest. Moreover, the MPRAGE sequence was used during acquisition to align
the FSE sequence perpendicularly to the plane of a participant's brainstem. Note that for some
participants specific absorption rate (SAR) limits were exceeded during the FSE acquisition, as reported
previously [87] (also see [54,126]). Sequence parameters are reported in Table 3.
543
544   **Table 3:** Overview of magnetic resonance imaging sequences

Magnetic resonance imaging sequence	Acquisition matrix; Slices (orientation; distance factor)	Voxel size (x, y, z; mm)	Repetition time (TR; ms)	Echo time (TE; ms)	Flip angle (°)
Magnetization Transfer (MT+)[1]	192 × 256 × 48 (axial; –)	1 × 1 × 3	38	5.5	10
Proton Density (MT–)[2]	192 × 256 × 48 (axial; –)	1 × 1 × 3	38	5.5	10
Fast Spin Echo (FSE)[3]	440 × 512 10 (axial; 20 %)	0.5 × 0.5 × 2.5	600	11	120
Magnetization Prepared Gradient-Echo (MPRAGE)[4]	256 × 256 × 192 (sagittal; –)	1 × 1 × 1	2500	4.77	7

Note: All sequences were acquired with a standard 32-channel head coil.
[1]: Magnetization Transfer Contrast (MTC) option enabled (MT pulse: 1200 Hz off-resonance, 16 ms [109]); 3-D
sequence
[2]: Magnetization Transfer Contrast (MTC) option disabled; 3-D sequence
[3]: Each FSE acquisition included four online averages and yielded two brainstem images. The values extracted from
these images were aggregated offline. SAR limits resulted in a slight variation in the number of slices [87];
2-D sequence; sometimes also called Turbo Spin Echo (TSE).
[4]: Inversion time (TI; ms): 1100; 3-D sequence
553

Magnetic resonance imaging data analysis:

We applied a previously established semi-automatic analysis procedure to extract individual LC and SN–
VTA intensity values from structural imaging data (for a detailed description and validation, see [87], for
applications, see [88,89,127]; for an independent validation see [91]). The following procedure was
performed separately for TP1 and TP2 imaging data.

Template construction and standardization:

First (step 1), scans of each scan modality (MPRAGE, FSE, MT+ and MT–) were iteratively aligned
across participants using a template-based procedure implemented in Advanced Normalization Tools
[ANTs] (v. 2.3 [128,129]; *antsMultivariateTemplateConstruction,* 6 iterations, including
N4BiasFieldCorrection). Before template construction, MPRAGE and MT– scans were resampled to 0.5
mm isometric resolution (ANTs' *ResampleImage*). Moreover, to facilitate template construction,
participants' native space FSE scans were aligned to their template-space MPRAGE scans
(*antsRegistrationSyNQuick*) [87]. Native space MT+ scans were aligned to resampled MT– scans to
account for potential movement effects between scan acquisitions (*antsRegistrationSyNQuick*). After their
alignment, MT– and MT+ scans were submitted to a common multimodal template construction, while
FSE and MPRAGE scans each were used to generate a brainstem and whole-brain template, respectively.
Next (step 2), modality-specific group templates (MPRAGE, FSE, MT+ and MT–) were linearly and non-
linearly coregistered (*antsRegistration*) to standard space (MNI-ICBM 152 linear, 0.5 mm [130]).
Specifically, templates with a sensitivity for catecholaminergic nuclei (FSE, MT+, MT–) were first
standardized to the whole-brain MPRAGE template (using a coregistration mask). Next, the MPRAGE
template was coregistered to MNI space and the transformations were applied to the other templates
(FSE, MT+, MT–; *antsApplyTransforms*). To improve coregistration accuracy, whole-brain templates
(MPRGAE, MT–, MNI) were skull stripped before alignment using the FMRIB Software Library [131].
Finally (step 3), all transformation matrices were concatenated and applied to individual participants'
scans to bring them from native to MNI space in a single step (*antsApplyTransforms*).

Semi-automatic intensity assessment:

To extract the intensity values of catecholaminergic nuclei, in standard space, individual scans were
masked with binary volumes of interest using Statistical Parametric Mapping toolbox version 12
(SPM12) [132] in Matlab (MathWorks). In particular, for the LC, we applied a previously established
high-confidence consensus mask [67]. For the SN–VTA, we relied on a previously established mask that
was based on manual tracings in template space [57]. Inter-participant comparisons of arbitrality scaled
MRI intensity values require that intensity values are normalized within participants [55]. Thus, we also
masked scans (FSE, MT+, MT–) with volumes of interest in potine [67] and midbrain [57] reference
areas, in line with earlier research [54,57,87]. Note that the fourth ventricle, which is in close proximity to
the LC, appears hyperintense in our MT+ scans. Thus, to rule out that the hyperintensity of the ventricle
could interfer with automatized LC assessments, we generated a sample-specific ventricle volume-of-no-
interest (based on the MT– group template), which we removed from MT+ and MT– scans before value
extraction. Within the masked scans (FSE, MT+, MT–), we then automatically searched for the voxel of
highest intensity in the LC, SN–VTA, and reference regions. Next, for each participant, spatially-resolved
intensity ratios for the LC and SN–VTA were computed per hemisphere (left, right) on a slice-by-slice
basis using the following formula [54,87]:

$$LC_{Ratio} = \frac{max(ROI) - max(Ref)}{max(Ref)}$$

Where max(*ROI*) denotes the peak intensity for a given slice in the LC or SN–VTA regions of interest
and max(*Ref*) indicates the peak intensity in the respective reference region. For the FSE modality, two
scans were available per participant (see *Magnetic resonance imaging data assessment*) and we averaged
the extracted intensity ratios within participants to obtain more stable estimates. For further analyses, for
all modalities (FSE, MT+, MT–) the peak intensity ratio across the regions of interest (LC, SN–VTA) was
calculated as an overall integrity measure [52,67,87]. Outlier values exceeding ± 3 standard deviations
were dropped, while all other values were linearly scaled (× 100) to facilitate subsequent model
estimation. Note that all analyses including LC or SN–VTA data were based on peak intensity ratios. That
is, the peak intensity of catecholaminergic nuclei standardized using nearby white matter regions (not
their raw intensity values). To facilitate readability, we will nonetheless use the term *intensity* in our
description.
At acquisition, the FSE sequence was centered on the pons and contained fewer slices compared to the
MT+ and MT– sequences. As evident in Figure 2 (a), the most rostral slices of our volume of interest (LC
meta mask and reference mask) reach the edges of the brainstem template and include high intensity
artifacts, which, however, are reliably excluded from analyses using the semiautomatic procedure (cf.
peak detection above).

Structural equation modeling:

We used structural equation modeling (SEM) to evaluate inter- and intra-individual differences in
catecholaminergic nuclei and their association with cognition using the Ωnyx software environment [133]
and the lavaan R package [134]. All models used full information maximum likelihood (FIML)
estimation to account for missing values. The adequacy of the reported models was evaluated using χ^2-
tests (i.e., an absolute fit index), as well as two frequently reported approximate fit indices: the root mean
square error of approximation (RMSEA) and the comparative fit index (CFI). RMSEA values close to or
below 0.06 and CFI values close to 0.95 or greater indicate good model fit [135,136]. Unless otherwise
noted, multi-group models were fit, comprising submodels for younger and older adults. For this,
invariance across age groups was evaluated by a hierarchical series of likelihood-ratio tests, probing
group differences in (1) factor loadings (weak invariance), (2) indicator intercepts (strong invariance), and
(3) residual variances (strict invariance) [137]. In case of longitudinal models, the same criteria were
applied to test invariance across time [110]. After establishing adequate model fit and invariance, the
significance of parameters of interest was evaluated using likelihood-ratio tests. That is, we created two
nested models—in one, the parameter of interest was freely estimated, whereas in the other model it was
fixed to zero. If a fixing the parameter of interest to zero resulted in a drop in model fit, as evaluated using
a likelihood-ratio test comparing the two nested models, this indicated the significance of the parameter
[138]. We used an alpha level of .05 for all statistical tests. Statistical results with *p*-values between 0.05
and 0.1 are described as statistical trend. In the following, cross-sectional models refer to models that
include TP2 data only, while longitudinal models evaluate the change in parameters of interest over time
(TP1–TP2 or TP1–TP3). Model code, visualizations, and output are available from [93].

Cross-sectional cognitive models:

We made use of a previously established factor structure [121] to integrate performance across several
working memory, episodic memory, and fluid intelligence tasks (see *Cognitive data assessment*) and
capture their shared variance on a latent level. Latent variables account for measurement error in the
observed scores (cognitive tasks) and thus increase statistical power to detect true effects [86].
We added covariances between the latent working memory, episodic memory, and fluid intelligence
factors, as performance across these cognitive domains had been shown to be correlated [121] (see Figure
S7).

Cross-sectional neural models:

We also adapted a previously established factor structure [87] to capture LC and SN–VTA intensity on a
latent level. Specifically, for each region (LC, SN–VTA) and scan modality (FSE, MT+, MT–), we used
the left and right hemispheric peak intensity as observed scores to estimate a modality-specific integrity
factor on a latent level. Note that our FSE sequence only covered the brainstem. Thus, we cannot obtain
SN–VTA estimates for this scan modality. To test the agreement in integrity estimates across modalities
(FSE, MT+, MT–), we added covariances among the modality-specific latent factors for each brain region
(see Figure S1).
Using the model describe above, we found a high correspondence in the integrity estimates for each
nucleus across modalities (see *Results*). Thus, in a second model, we introduced a multimodal integrity
factor for the LC and SN–VTA that captures the commonalities across scan modalities while removing
the modality-specific measurement error (for similar approaches and a detailed discussion, see [83,84];
see Figure S2). Finally, as dopamine and noradrenaline are products of the same biosynthesis pathway,
with dopamine a direct precursor to noradrenaline [44], we evaluated the association of the multimodal
LC and SN–VTA integrity factors by estimating their covariance.

Cross-sectional neuro–cognitive models:

After separately establishing models for our cognitive and neural measures that showed acceptable fit and
invariance across age groups, we unified these models to probe associations between catecholaminergic
nuclei and cognition [87]. In the unified neuro-cognitive model, we estimated covariances between the
multimodal LC and SN–VTA integrity factors and working memory, episodic memory, and fluid
intelligence (correlation model; see Figure S8 and S9). In addition, we specified a second neuro-cognitive
model, in which regression paths were drawn from the latent neural to the cognitive factors (regression
model; see Figures S10 and S11). The correlation model evaluates associations between the LC and
cognition irrespective of those between the SN–VTA and cognition. By contrast, the regression model
tests whether one region explains variance in the cognitive factors over and above the other region, thus
providing complementary information [84].

Longitudinal cognitive models:

Making use of the repeated assessments of cognitive performance, we tested for late-life changes in
working memory and episodic memory tasks over time (TP1–TP3). Note that cognitive data at TP3 was
only available for older adults (see *Cognitive data assessment*). Thus, here we relied on single-group
models that excluded younger adults. In particular, we specified a latent change score model [86,139] for

each cognitive task. These latent change score models yield a latent slope factor for each task that
expresses participants' performance difference between TP1 and TP3 (see Figures S21 and S22).
Next, we tested whether there is a common latent factor of working memory or episodic memory change.
If changes in performance were shared across tasks (i.e., task independent) of one cognitive domain, they
could be captured on a higher order latent level (cf. the latent working memory and episodic memory
factors in the cross-sectional cognitive model; Figure S7). To evaluate potential associations in the
changes across working memory and episodic memory tasks, we added covariances among the task-
specific slope terms (see Figures S21 and S22).
Most covariances across task-specific slope terms did not reach significance (see *Supplementary results*).
Thus, we did not further attempt to capture task-independent changes in working memory or episodic
memory on a latent level.

Longitudinal neural models:

Assessment of across-time stability:

Our cross-sectional analyses of different MRI sequences sensitive for catecholaminergic nuclei (FSE,
MT+, MT−) demonstrated a high agreement in integrity estimates across imaging modalities (see *Cross-*
sectional neural models and *Results* Figure 3). Leveraging the longitudinal nature of this study, we
additionally explored the stability of the modality-specific integrity estimates over time (TP1, TP2), as a
proxy for their reliability [94]. Specifically, we started with a modality-specific SN–VTA and LC models
for TP2 (see *Cross-sectional neural models* and Figure S1) and then appended the same variables for
TP1. We introduced covariances linking modality-specific factors of TP1 and TP2 to evaluate the stability
in integrity estimates over time (see Figures S12 and S16). Moreover, we allowed for correlated residuals
over time, as suggested for longitudinal models [110].
Using the model describe above, we found a high stability of the modality-specific integrity estimates for
each region across time (see *Results* Figure 3 and S18). Comparable to our cross-sectional analyses, we
thus again added multimodal integrity factors for the LC and SN–VTA for each time point (TP1, TP2). If
the multimodal integrity factors indeed remove measurement error [84], we should observe a higher
stability across time points of the multimodal as compared to the modality-specific integrity factors. To
test this hypothesis, we computed the covariance between the multimodal factors of TP1 and TP2 (see
Figures S13 and S17). In addition, we directly compared the stability estimates, our reliability proxy, for
the modality-specific and multimodal SN–VTA and LC factors. Note that the modality-specific and
multimodal integrity models including TP1 and TP2 were fit across younger and older adults to obtain a
single reliability proxy for the complete sample.

Assessment of within-person changes:

Previous cross-sectional research observed between-person age differences in catecholaminergic nuclei
integrity [74,75], however, longitudinal studies that evaluate within-person changes in the SN–VTA and
LC are scarce. Making use of the imaging data of both time points (TP1, TP2), we thus estimated changes
in catecholaminergic nuclei using latent change score models [86,139] for each imaging modality (MT+,
MT−). Note that the FSE sequence was only acquired at TP2; precluding change analyses (see Table 1).
To reduce model complexity, we averaged intensity ratios across hemispheres for these models.
Comparable to our longitudinal cognitive analyses, we first evaluated whether the change in each region

was shared across imaging modalities (i.e., sequence independent) by computing the covariance of the
modality-specific slope terms (see Figures S14 and S19).
For the SN–VTA as well as the LC we observed that the changes were indeed correlated across imaging
modalities (MT+, MT–). Thus, in a second set of models, we introduced a higher order multimodal slope
factor to capture the shared variance across modality-specific slope factors (see Figures S15 and S20).

Longitudinal neuro–cognitive models:

In older adults, we found that SN–VTA and LC integrity changed across time (TP1–TP2; see
Longitudinal neural models). To evaluate the behavioral implications of these changes, we leveraged a
previously established cognitive factor structure [121,125] (see *Cross-sectional cognitive models*).
Specifically, we were interested in testing whether changes in catecholaminergic nuclei (TP1–TP2) could
be used to predict future cognition (at TP3). For this, we unified our neural change models (see Figures
S15 and S20) with models of working memory and episodic memory (at TP3; cf. Figure S7). In the
unified model, we specified regression paths from the neural change factors to the cognitive factors (SN–
VTA to working memory; LC to episodic memory). Finally, we added chronological age (at TP2) as an
additional predictor to test (1) whether catecholaminergic nuclei explains future cognition over and above
age and (2) how changes in catecholaminergic nuclei differ in old age (see Figures S23 and S24). Note
that cognition at TP3 was only assessed for older adults. Thus, we specified single-group models
excluding younger adults.

Control analyses:

Catecholaminergic neuromodulation influences neural processing in the medial temporal lobe, a key node
of the memory network [35,37,41,43,45,46]. Moreover, catecholaminergic nuclei have direct projections
to the medial temporal lobe [41,43,140,141] and their integrity has been linked to tau pathology in these
areas [67–69]. Thus, as a control analysis, we additionally evaluated medial temporal lobe volumes (at
TP2). This allowed us to compare our measures of catecholaminergic nuclei and their association with
memory performance to those of a well-established player in the memory network [108].

Voxel-based morphometry assessment:

Whole-brain MPRAGE images were processed using the voxel-based morphometry pipeline in SPM12 in
MATLAB [132,142]. First, images were segmented into distinct tissue classes (e.g., grey matter, white
matter, cerebrospinal fluid) using SPM12's Unified Segmentation procedure [143]. Next, a study-specific
DARTEL template was created and segmented images were aligned to the template, followed by spatial
normalization, modulation, and smoothing with a 2 mm full-width at half-maximum isotropic Gaussian
kernel [144,145]. The resulting normalized, modulated, and smoothed grey matter images were used to
derive region-of-interest volumes. Region-of-interest masks from the AAL3 atlas were applied to
processed grey matter images using native SPM functions and the get_totals script
(http://www0.cs.ucl.ac.uk/staff/g.ridgway/vbm/get_totals.m) to calculate volumes for the
parahippocampal and hippocampal regions [146]. Outlier values exceeding ± 3 standard deviations were
dropped, while all other values were linearly scaled (× 10,000) to facilitate subsequent model estimation.

Cross-sectional neuro–cognitive models including voxel-based morphometry factor:

Mimicking our integrity factors of catecholaminergic nuclei (see Figure S1), we aggregated across left
and right hemispheric volumes to estimate latent parahippocampal and hippocampal factors for older
adults. We specified a covariance between the two regions to evaluate their association (see Figure S3).
We observed that parahippocampal and hippocampal factors were highly correlated (see *Supplementary*
results). Thus, we introduced a higher-order medial temporal lobe factor to capture their shared variance,
which we then included in our neuro-cognitive models (see *Cross-sectional neuro–cognitive models*).
Specifically, for older adults we estimated one correlation model and one regression model [84]. In each
model, we compared the associations of catecholaminergic nuclei and cognition with the medial temporal
lobe–cognition association (see Figures 6 and S10, S11). In addition, we quantified the interrelations of
SN–VTA and LC integrity and medial temporal lobe volume by estimating their covariance.

Data availability

The data that our results are based on are available from the BASE-II steering committee upon approved
research proposal (see https://www.base2.mpg.de/en). For inquiries, please contact Dr. Ludmilla Müller,
BASE-II project coordinator (lmueller@mpib-berlin.mpg.de).
To facilitate comparability of study results, we share the group templates with sensitivity for
catecholaminergic nuclei (FSE, MT+, MT–) in MNI 0.5 mm linear space (https://osf.io/eph9a/) [93].
The LC consensus volume of interest (LC meta mask and pontine reference mask) is available via
https://osf.io/sf2ky/ [89].
We provide two synthetic datasets of simulated cases (n = 250) that follow the population described in
our models, with the parameter values displayed in the model visualizations (generated using [133]). In
combination with the model code (see below), these data allow reproduction of key results.

Code availability

All statistical models that our inferences are based on and their outputs are available via:
https://osf.io/eph9a [93]
Vector-graphics visualizations for all models are also available via OSF.
Additional custom code used for processing behavioral and neural data are available from the
corresponding authors upon request.

Competing interests:

The authors declare no competing interests.

Acknowledgements:

This article uses data from the Berlin Aging Study II (BASE-II), which was supported by the German Federal Ministry of Education and Research (Bundesministerium für Bildung und Forschung, BMBF) under grant numbers #16SV5536 K, #16SV5537, #16SV5538, and #16SV5837, #01UW070 and #01UW0808. Additional contributions (e.g., financial, equipment, logistics, personnel) are made from each of the other participating sites, i.e., the Max Planck Institute for Human Development (MPIB), Max Planck Institute for Molecular Genetics (MPIMG), Charite-Universiätsmedizin, University Medicine, German Institute for Economic Research (DIW), Humboldt-Universität zu Berlin, all located in Berlin, Germany, and University of Lübeck, and University of Tübingen, Germany. For further information about the BASE-II project, see https://www.base2.mpg.de/en

MW-B received support from the German Research Foundation (DFG, WE 4269/5-1) and the Jacobs Foundation (Early Career Research Fellowship 2017–2019).

MM's work was supported by an Alexander von Humboldt fellowship and by National Institutes of Health grant R01AG025340.

SB was supported by a National Institutes of Health grant T32AG000037 and a National Science Foundation Grant DGE-1842487.

SD received funding from the National Science Foundation (DGE1418060).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

We thank Dr. Clifford Cassidy for providing the substantia nigra and crus cerebri masks [57]. We are grateful to Dr. Ylva Köhncke and Sarah Polk for preprocessing the cognitive data.

We thank Sumedha Attanti for visually inspecting the VBM-processed segmentations.

References:

Nyberg, L. *et al.* (2012) Memory aging and brain maintenance. *Trends Cogn. Sci.* 16, 292–305

Sander, M.C. *et al.* (2012) Lifespan age differences in working memory: A two-component framework. *Neurosci. Biobehav. Rev.* 36, 2007–2033

Gazzaley, A. and Nobre, A.C. Top-down modulation: Bridging selective attention and working memory. , *Trends in Cognitive Sciences*, 16. 01-Feb-(2012) , Elsevier Current Trends, 129–135

Hasher, L. and Zacks, R.T. (1988) Working memory, comprehension, and aging: A review and a new view. *Psychol. Learn. Motiv. - Adv. Res. Theory* 22, 193–225

Shing, Y.L. *et al.* (2010) Episodic memory across the lifespan: The contributions of associative and strategic components. *Neurosci. Biobehav. Rev.* 34, 1080–1091

Tulving, E. (1983) *Elements of episodic memory*, Clarendon Press.

Bäckman, L. *et al.* (2006) The correlative triad among aging, dopamine, and cognition: Current status and future prospects. *Neurosci. Biobehav. Rev.* 30, 791–807

Li, S.-C. *et al.* (2001) Aging cognition: From neuromodulation to representation. *Trends Cogn. Sci.* 5, 479–486

Braver, T.S. and Barch, D.M. (2002) A theory of cognitive control, aging cognition, and neuromodulation. *Neurosci. Biobehav. Rev.* 26, 809–817

Mather, M. and Harley, C.W. (2016) The locus coeruleus: Essential for maintaining cognitive function and the aging brain. *Trends Cogn. Sci.* 20, 214–226

Weinshenker, D. (2018) Long road to ruin: Noradrenergic dysfunction in neurodegenerative disease. *Trends Neurosci.* 41, 211–223

Arnsten, A.F.T. *et al.* (2012) Neuromodulation of thought: flexibilities and vulnerabilities in prefrontal cortical network synapses. *Neuron* 76, 223

Braak, H. *et al.* (2011) Stages of the pathologic process in Alzheimer disease: Age categories from 1 to 100 years. *J. Neuropathol. Exp. Neurol.* 70, 960–969

Marien, M.R. *et al.* (2004) Noradrenergic mechanisms in neurodegenerative diseases: A theory. *Brain Res. Rev.* 45, 38–78

Matchett, B.J. *et al.* (2021) The mechanistic link between selective vulnerability of the locus coeruleus and neurodegeneration in Alzheimer's disease. *Acta Neuropathol.* 1, 3

Lyness, S.A. *et al.* (2003) Neuron loss in key cholinergic and aminergic nuclei in Alzheimer disease: A meta-analysis. *Neurobiol. Aging* 24, 1–23

Lee, J.-Y. *et al.* (2022) Multimodal brain and retinal imaging of dopaminergic degeneration in Parkinson disease. *Nat. Rev. Neurol. 2022* 0123456789, 1–18

Aarsland, D. *et al.* (2021) Parkinson disease-associated cognitive impairment. *Nat. Rev. Dis. Prim. 2021 71* 7, 1–21

Knopman, D.S. *et al.* (2021) Alzheimer disease. *Nat. Rev. Dis. Prim.* 7, 33

Lőrincz, M.L. and Adamantidis, A.R. (2017) Monoaminergic control of brain states and sensory processing: Existing knowledge and recent insights obtained with optogenetics. *Prog. Neurobiol.* 151, 237–253

Düzel, E. *et al.* (2009) Functional imaging of the human dopaminergic midbrain. *Trends Neurosci.* 32, 321–328

Poe, G.R. *et al.* (2020) Locus coeruleus: a new look at the blue spot. *Nat. Rev. Neurosci.* 21, 644–659

Servan-Schreiber, D. *et al.* (1990) A network model of catecholamine effects: Gain, signal-to-noise ratio, and behavior. *Science (80-.).* 249, 892–5

Thiele, A. and Bellgrove, M.A. (2018) Neuromodulation of attention. *Neuron* 97, 769–785

Aston-Jones, G. and Cohen, J.D. (2005) An integrative theory of locus coeruleus-norepinephrine function: Adaptive gain and optimal performance. *Annu. Rev. Neurosci.* 28, 403–450

Mather, M. *et al.* (2016) Norepinephrine ignites local hotspots of neuronal excitation: How arousal amplifies selectivity in perception and memory. *Behav. Brain Sci.* 39, e200

853	27	Koen, J.D. and Rugg, M.D. (2019) Neural dedifferentiation in the aging brain. *Trends Cogn. Sci.*
854		23, 547–559
855	28	Robbins, T.W. and Arnsten, A.F.T. (2009) The neuropsychopharmacology of fronto-executive
856		function: Monoaminergic modulation. *Annu. Rev. Neurosci.* 32, 267–287
857	29	Cools, R. and Arnsten, A.F.T. (2021) Neuromodulation of prefrontal cortex cognitive function in
858		primates: the powerful roles of monoamines and acetylcholine. *Neuropsychopharmacology* DOI:
859		10.1038/s41386-021-01100-8
860	30	Goldman-Rakic, P.S. (1995) Cellular basis of working memory. *Neuron* 14, 477–485
861	31	Wang, M. *et al.* (2011) Neuronal basis of age-related working memory decline. *Nature* 476, 210–
862		213
863	32	Arnsten, A.F.T. and Goldman-Rakic, P.S. (1985) Alpha 2-adrenergic mechanisms in prefrontal
864		cortex associated with cognitive decline in aged nonhuman primates. *Science (80-.).* 230, 1273–
865		1276
866	33	Arnsten, A.F.T. and Contant, T.A. (1992) Alpha-2 adrenergic agonists decrease distractibility in
867		aged monkeys performing the delayed response task. *Psychopharmacology (Berl).* 108, 159–69
868	34	Cai, J.X. and Arnsten, A.F.T. (1997) Dose-dependent effects of the dopamine D1 receptor agonists
869		A77636 or SKF81297 on Spatial working memory in aged monkeys. *J. Pharmacol. Exp. Ther.*
870		283, 183 LP – 189
871	35	Lisman, J.E. and Grace, A.A. The hippocampal-VTA loop: Controlling the entry of information
872		into long-term memory. , *Neuron*, 46. 02-Jun-(2005) , Neuron, 703–713
873	36	O'Dell, T.J. *et al.* (2015) β-Adrenergic receptor signaling and modulation of long-term
874		potentiation in the mammalian hippocampus. *Learn. Mem.* 22, 461–471
875	37	Sara, S.J. (2009) The locus coeruleus and noradrenergic modulation of cognition. *Nat. Rev.*
876		*Neurosci.* 10, 211–223
877	38	Lisman, J. *et al.* (2011) A neoHebbian framework for episodic memory; role of dopamine-
878		dependent late LTP. *Trends Neurosci.* 34, 536–547
879	39	Hagena, H. *et al.* (2016) β-adrenergic control of pippocampal function: Subserving the
880		choreography of synaptic information storage and memory. *Cereb. Cortex* 26, 1349–1364
881	40	Gruber, M.J. and Ranganath, C. (2019) How curiosity enhances hippocampus-dependent memory:
882		The prediction, appraisal, curiosity, and exploration (PACE) Framework. *Trends Cogn. Sci.* 23,
883		1014–1025
884	41	Takeuchi, T. *et al.* (2016) Locus coeruleus and dopaminergic consolidation of everyday memory.
885		*Nature* 537, 357–362
886	42	Kempadoo, K.A. *et al.* (2016) Dopamine release from the locus coeruleus to the dorsal
887		hippocampus promotes spatial learning and memory. *Proc. Natl. Acad. Sci. U. S. A.* 113, 14835–
888		14840
889	43	Tsetsenis, T. *et al.* (2021) Midbrain dopaminergic innervation of the hippocampus is sufficient to
890		modulate formation of aversive memories. *Proc. Natl. Acad. Sci.* 118, e2111069118
891	44	Eisenhofer, G. (2004) Catecholamine metabolism: A contemporary view with implications for
892		physiology and medicine. *Pharmacol. Rev.* 56, 331–349
893	45	Wagatsuma, A. *et al.* (2018) Locus coeruleus input to hippocampal CA3 drives single-trial
894		learning of a novel context. *Proc. Natl. Acad. Sci. U. S. A.* 115, E310–E316
895	46	Duszkiewicz, A.J. *et al.* (2019) Novelty and dopaminergic modulation of memory persistence: A
896		tale of two systems. *Trends Neurosci.* 42, 102–114
897	47	Chowdhury, A. *et al.* (2022) A locus coeruleus-dorsal CA1 dopaminergic circuit modulates
898		memory linking. *Neuron* DOI: 10.1016/J.NEURON.2022.08.001
899	48	Burke, S.N. and Barnes, C.A. (2006) Neural plasticity in the ageing brain. *Nat. Rev. Neurosci.*
900		*2006 71* 7, 30–40
901	49	Barnes, C.A. (2003) Long-term potentiation and the ageing brain. *Philos. Trans. R. Soc. B Biol.*
902		*Sci.* 358, 765
903	50	Lubec, J. *et al.* (2021) Reinstatement of synaptic plasticity in the aging brain through specific

dopamine transporter inhibition. *Mol. Psychiatry* 26, 7076–7090

51 Astafiev, S. V. *et al.* (2010) Comment on "Modafinil shifts human locus coeruleus to low-tonic,
high-phasic activity during functional MRI" and "Homeostatic sleep pressure and responses to
sustained attention in the suprachiasmatic area." *Science (80-.).* 328, 309

52 Keren, N.I. *et al.* (2009) In vivo mapping of the human locus coeruleus. *Neuroimage* 47, 1261–
1267

53 Sulzer, D. *et al.* (2018) Neuromelanin detection by magnetic resonance imaging (MRI) and its
promise as a biomarker for Parkinson's disease. *npj Park. Dis.* 4, 11

54 Liu, K.Y. *et al.* (2017) Magnetic resonance imaging of the human locus coeruleus: A systematic
review. *Neurosci. Biobehav. Rev.* 83, 325–355

55 Betts, M.J. *et al.* (2019) Locus coeruleus imaging as a biomarker for noradrenergic dysfunction in
neurodegenerative diseases. *Brain* 142, 2558–2571

56 Keren, N.I. *et al.* (2015) Histologic validation of locus coeruleus MRI contrast in post-mortem
tissue. *Neuroimage* 113, 235–245

57 Cassidy, C.M. *et al.* (2019) Neuromelanin-sensitive MRI as a noninvasive proxy measure of
dopamine function in the human brain. *Proc. Natl. Acad. Sci. U. S. A.* 116, 5108–5117

58 Kitao, S. *et al.* (2013) Correlation between pathology and neuromelanin MR imaging in
Parkinson's disease and dementia with Lewy bodies. *Neuroradiology* 55, 947–953

59 Zecca, L. *et al.* (2004) The role of iron and copper molecules in the neuronal vulnerability of locus
coeruleus and substantia nigra during aging. *Proc. Natl. Acad. Sci. U. S. A.* 101, 9843–9848

60 Zucca, F.A. *et al.* (2017) Interactions of iron, dopamine and neuromelanin pathways in brain aging
and Parkinson's disease. *Prog. Neurobiol.* 155, 96–119

61 Carballo-Carbajal, I. *et al.* (2019) Brain tyrosinase overexpression implicates age-dependent
neuromelanin production in Parkinson's disease pathogenesis. *Nat. Commun.* 10, 973

62 Trujillo, P. *et al.* (2017) Contrast mechanisms associated with neuromelanin-MRI. *Magn. Reson.*
Med. 78, 1790–1800

63 Simpson, K.L. and Lin, R.C.S. (2007) Neuroanatomical and chemical organization of the locus
coeruleus. In *Brain Norepinephrine* (Ordway, G. A. et al., eds), pp. 9–52, Cambridge University
Press

64 Watanabe, T. *et al.* (2019) Magnetic resonance imaging of noradrenergic neurons. *Brain Struct.*
Funct. 224, 1609–1625

65 Priovoulos, N. *et al.* (2020) Unraveling the contributions to the neuromelanin-MRI contrast. *Brain*
Struct. Funct. 1, 3

66 Trujillo, P. *et al.* (2019) Quantitative magnetization transfer imaging of the human locus
coeruleus. *Neuroimage* 200, 191–198

67 Dahl, M.J. *et al.* (2022) Locus coeruleus integrity is related to tau burden and memory loss in
autosomal-dominant Alzheimer's disease. *Neurobiol. Aging* 112, 39–54

68 Jacobs, H.I.L. *et al.* (2021) In vivo and neuropathology data support locus coeruleus integrity as
indicator of Alzheimer's disease pathology and cognitive decline. *Sci. Transl. Med.* 13, eabj2511

69 Cassidy, C.M. *et al.* (2022) Association of locus coeruleus integrity with Braak stage and
neuropsychiatric symptom severity in Alzheimer's disease. *Neuropsychopharmacol. 2022* DOI:
10.1038/s41386-022-01293-6

70 Gaurav, R. *et al.* (2021) Longitudinal Changes in Neuromelanin MRI Signal in Parkinson's
Disease: A Progression Marker. *Mov. Disord.* 36, 1592–1602

71 Matsuura, K. *et al.* (2016) A longitudinal study of neuromelanin-sensitive magnetic resonance
imaging in Parkinson's disease. *Neurosci. Lett.* 633, 112–117

72 Braak, H. *et al.* (2003) Staging of brain pathology related to sporadic Parkinson's disease.
Neurobiol. Aging 24, 197–211

73 Brettschneider, J. *et al.* Spreading of pathology in neurodegenerative diseases: A focus on human
studies. , *Nature Reviews Neuroscience*, 16. 01-Jan-(2015) , Nature Publishing Group, 109–120

74 Liu, K.Y. *et al.* (2019) In vivo visualization of age-related differences in the locus coeruleus.

955		*Neurobiol. Aging* 74, 101–111
956	75	Shibata, E. *et al.* (2006) Age-related changes in locus coeruleus on neuromelanin magnetic
957		resonance imaging at 3 Tesla. *Magn. Reson. Med. Sci.* 5, 197–200
958	76	Xing, Y. *et al.* (2018) Life span pigmentation changes of the substantia nigra detected by
959		neuromelanin-sensitive MRI. *Mov. Disord.* 33, 1792–1799
960	77	Jacobs, H.I.L. *et al.* (2021) Alzheimer's disease pathology: pathways between central
961		norepinephrine activity, memory, and neuropsychiatric symptoms. *Mol. Psychiatry* 26, 897–906
962	78	Jacobs, H.I.L. *et al.* (2022) Waning locus coeruleus integrity precedes cortical tau accrual in
963		preclinical autosomal dominant Alzheimer's disease. *Alzheimers. Dement.* DOI:
964		10.1002/ALZ.12656
965	79	Delius, J.A.M. *et al.* (2015) Berlin Aging Studies (BASE and BASE-II). In *Encyclopedia of*
966		*geropsychology* (Pachana, N. A., ed), pp. 386–395, Springer
967	80	Gerstorf, D. *et al.* (2016) The Berlin Aging Study II: An overview [Editorial]. *Gerontology* 62,
968		311–315
969	81	Bertram, L. *et al.* (2014) Cohort profile: The Berlin Aging Study II (BASE-II). *Int. J. Epidemiol.*
970		43, 703–712
971	82	Demuth, I. *et al.* (2021) Cohort profile: follow-up of a Berlin Aging Study II (BASE-II) subsample
972		as part of the GendAge study. *BMJ Open* 11, e045576
973	83	Kühn, S. *et al.* (2017) In search of features that constitute an "enriched environment" in humans:
974		Associations between geographical properties and brain structure. *Sci. Rep.* 7, 1–8
975	84	Köhncke, Y. *et al.* (2021) Hippocampal and Parahippocampal Gray Matter Structural Integrity
976		Assessed by Multimodal Imaging Is Associated with Episodic Memory in Old Age. *Cereb. Cortex*
977		31, 1464–1477
978	85	Düzel, S. *et al.* (2016) The Subjective Health Horizon Questionnaire (SHH-Q): Assessing Future
979		Time Perspectives for Facets of an Active Lifestyle. *Gerontology* 62, 345–353
980	86	Kievit, R.A. *et al.* (2018) Developmental cognitive neuroscience using latent change score models:
981		A tutorial and applications. *Dev. Cogn. Neurosci.* 33, 99–117
982	87	Dahl, M.J. *et al.* (2019) Rostral locus coeruleus integrity is associated with better memory
983		performance in older adults. *Nat. Hum. Behav.* 3, 1203–1214
984	88	Gallant, S.N. *et al.* (2022) Behavioral and fMRI evidence that arousal enhances bottom-up
985		selectivity in young but not older adults. *Neurobiol. Aging* DOI:
986		10.1016/J.NEUROBIOLAGING.2022.08.006
987	89	Dahl, M.J. *et al.* (2022) Locus coeruleus integrity is related to tau burden and memory loss in
988		autosomal-dominant Alzheimer's disease. *Neurobiol. Aging* 112, 39–54
989	90	Bachman, S.L. *et al.* (2022) Daily heart rate variability biofeedback training decreases locus
990		coeruleus MRI contrast in younger adults. *medRxiv* DOI: 10.1101/2022.02.04.22270468
991	91	Giorgi, F.S. *et al.* (2022) Locus Coeruleus magnetic resonance imaging: a comparison between
992		native-space and template-space approach. *J. Neural Transm. 2022* 1, 1–8
993	92	Polk, S.E. *et al.* (2022) Change in latent gray-matter structural integrity is associated with change
994		in cardiovascular fitness in older Adults who engage in at-home aerobic exercise. *Front. Hum.*
995		*Neurosci.* 16,
996	93	Dahl, M.J. *et al.* (2022) The integrity of dopaminergic and noradrenergic brain regions is
997		associated with different aspects of late-life memory performance. *OSF* DOI:
998		https://doi.org/10.17605/OSF.IO/EPH9A
999	94	Brandmaier, A.M. *et al.* (2018) Precision, reliability, and effect size of slope variance in latent
1000		growth curve models: Implications for statistical power analysis. *Front. Psychol.* 9, 294
1001	95	Eid, M. *et al.* (2015) *Statistik und Forschungsmethoden: Lehrbuch*, Beltz.
1002	96	Briand, L.A. *et al.* (2007) Modulators in concert for cognition: Modulator interactions in the
1003		prefrontal cortex. *Prog. Neurobiol.* 83, 69–91
1004	97	Elman, J.A. *et al.* (2021) MRI-assessed locus coeruleus integrity is heritable and associated with
1005		cognition, Alzheimer's risk, and sleep-wake disturbance. *Alzheimer's Dement.* 17, 1017– 1025

98   Liu, K.Y. *et al.* (2020) Noradrenergic-dependent functions are associated with age-related locus
coeruleus signal intensity differences. *Nat. Commun.* 11, 1712
99   Allen, M. *et al.* (2021) Raincloud plots: a multi-platform tool for robust data visualization.
Wellcome Open Res. 2021 463 4, 63
100   Ehrenberg, A.J. *et al.* (2017) Quantifying the accretion of hyperphosphorylated tau in the locus
coeruleus and dorsal raphe nucleus: The pathological building blocks of early Alzheimer's disease.
Neuropathol. Appl. Neurobiol. 43, 393–408
101   Stratmann, K. *et al.* (2016) Precortical phase of Alzheimer's disease (AD)-related tau cytoskeletal
pathology. *Brain Pathol.* 26, 371–386
102   Theofilas, P. *et al.* (2015) Turning on the light within: Subcortical nuclei of the isodentric core
and their role in Alzheimer's disease pathogenesis. *J. Alzheimer's Dis.* 46, 17–34
103   Chalermpalanupap, T. *et al.* (2017) Down but not out: The consequences of pretangle tau in the
locus coeruleus. *Neural Plast.* 2017, 7829507
104   Ghosh, A. *et al.* (2019) An experimental model of Braak's pretangle proposal for the origin of
Alzheimer's disease: The role of locus coeruleus in early symptom development. *Alzheimer's Res.*
Ther. 11, 59
105   Rorabaugh, J.M. *et al.* (2017) Chemogenetic locus coeruleus activation restores reversal learning
in a rat model of Alzheimer's disease. *Brain* 140, 3023–3038
106   Feinstein, D.L. *et al.* (2002) Noradrenergic regulation of inflammatory gene expression in brain.
Neurochem. Int. 41, 357–365
107   Rommelfanger, K.S. and Weinshenker, D. (2007) Norepinephrine: The redheaded stepchild of
Parkinson's disease. *Biochem. Pharmacol.* 74, 177–190
108   Moscovitch, M. *et al.* (2016) Episodic memory and beyond: The hippocampus and neocortex in
transformation. *Annu. Rev. Psychol.* 67, 105–134
109   Bunzeck, N. *et al.* (2007) Mesolimbic novelty processing in older adults. *Cereb. Cortex* 17, 2940–
2948
110   Little, T.D. (2013) *Longitudinal structural equation modeling.*, Guilford Press.
111   Wilson, R.S. *et al.* (2013) Neural reserve, neuronal density in the locus ceruleus, and cognitive
decline. *Neurology* 80, 1202–1208
112   Seiler, S. *et al.* (2014) Magnetization Transfer Imaging for in vivo Detection of Microstructural
Tissue Changes in Aging and Dementia: A Short Literature Review. *J. Alzheimer's Dis.* 42, S229–
S237
113   Köhncke, Y. *et al.* (2021) Hippocampal and Parahippocampal Gray Matter Structural Integrity
Assessed by Multimodal Imaging Is Associated with Episodic Memory in Old Age. *Cereb. Cortex*
31, 1464–1477
114   Tona, K.D. *et al.* (2017) In vivo visualization of the locus coeruleus in humans: Quantifying the
test–retest reliability. *Brain Struct. Funct.* 222, 4203–4217
115   Allen, R.J. *et al.* (2022) Intact high-resolution working memory binding in a patient with
developmental amnesia and selective hippocampal damage. *Hippocampus* 32, 597–609
116   Ciampa, C.J. *et al.* (2022) Associations among locus coeruleus catecholamines, tau pathology, and
memory in aging. *Neuropsychopharmacol. 2022* DOI: 10.1038/s41386-022-01269-6
117   Biondetti, E. *et al.* (2020) Spatiotemporal changes in substantia nigra neuromelanin content in
Parkinson's disease. *Brain* 143, 2757–2770
118   Theofilas, P. *et al.* (2017) Locus coeruleus volume and cell population changes during
Alzheimer's disease progression: A stereological study in human postmortem brains with potential
implication for early-stage biomarker discovery. *Alzheimer's Dement.* 13, 236–246
119   David, M.C.B. *et al.* (2022) Cognitive and neuropsychiatric effects of noradrenergic treatment in
Alzheimer's disease: systematic review and meta-analysis. *J. Neurol. Neurosurg. Psychiatry* 0,
jnnp-2022-329136
120   O'Callaghan, C. *et al.* (2021) Locus coeruleus integrity and the effect of atomoxetine on response
inhibition in Parkinson's disease. *Brain* 144, 2513–2526

121   Düzel, S. *et al.* (2016) The Subjective Health Horizon Questionnaire (SHH-Q): Assessing Future
1058   Time Perspectives for Facets of an Active Lifestyle. *Gerontology* 62, 345–353
1059   122   Lindenberger, U. *et al.* (1993) Speed and intelligence in old age. *Psychol. Aging* 8, 207–220
1060   123   Schmiedek, F. *et al.* (2010) Hundred days of cognitive training enhance broad cognitive abilities in
1061   adulthood: Findings from the COGITO study. *Front. Aging Neurosci.* 2,
1062   124   Lindenberger, U. *et al.* (1993) Speed and intelligence in old age. *Psychol. Aging* 8, 207–220
1063   125   Kühn, S. *et al.* (2019) Food for thought: association between dietary tyrosine and cognitive
1064   performance in younger and older adults. *Psychol. Res.* 83, 1097–1106
1065   126   Chen, X. *et al.* (2014) Simultaneous imaging of locus coeruleus and substantia nigra with a
1066   quantitative neuromelanin MRI approach. *Magn. Reson. Imaging* 32, 1301–1306
1067   127   Bachman, S.L. *et al.* (2022) Daily heart rate variability biofeedback training decreases locus
1068   coeruleus MRI contrast in younger adults. *medRxiv* DOI: 10.1101/2022.02.04.22270468
1069   128   Avants, B.B. *et al.* (2009) Advanced Normalization Tools: V1.0. *Insight J.* 2,
1070   129   Avants, B.B. *et al.* (2011) A reproducible evaluation of ANTs similarity metric performance in
1071   brain image registration. *Neuroimage* 54, 2033–2044
1072   130   Ciric, R. *et al.* (2021) TemplateFlow: a community archive of imaging templates and atlases for
1073   improved consistency in neuroimaging. *bioRxiv* DOI: 10.1101/2021.02.10.430678
1074   131   Smith, S.M. (2002) Fast robust automated brain extraction. *Hum. Brain Mapp.* 17, 143–155
1075   132   Friston, K. *et al.* (2007) *Statistical Parametric Mapping: The Analysis of Functional Brain
1076   Images*, 1st edn.Academic Press.
1077   133   von Oertzen, T. *et al.* (2015) Structural equation modeling with Ωnyx. *Struct. Equ. Model.* 22,
1078   148–161
1079   134   Rosseel, Y. (2012) Lavaan: An R package for structural equation modeling. *J. Stat. Softw.* 48, 1–
1080   36
1081   135   Brown, T.A. (2006) *Confirmatory factor analysis for applied research*, Guilford Press.
1082   136   Hu, L.T. and Bentler, P.M. (1999) Cutoff criteria for fit indexes in covariance structure analysis:
1083   Conventional criteria versus new alternatives. *Struct. Equ. Model.* 6, 1–55
1084   137   Meredith, W. and Teresi, J.A. (2006) An essay on measurement and factorial invariance. *Med.
1085   Care* 44, S69–S77
1086   138   Kline, R.B. (2016) *Principles and practice of structural equation modeling, 4th ed.*, Guilford
1087   Press.
1088   139   McArdle, J.J. (2009) Latent variable modeling of differences and changes with longitudinal data.
1089   *Annu. Rev. Psychol.* 60, 577–605
1090   140   Solders, S.K. *et al.* (2021) Diffusion MRI tractography of the locus coeruleus-transentorhinal
1091   cortex connections using GO-ESP. *Magn. Reson. Med.* 00, 1–16
1092   141   Porat, S. *et al.* (2022) Age differences in diffusivity in the locus coeruleus and its ascending
1093   noradrenergic tract. *Neuroimage* 251, 119022
1094   142   Ashburner, J. and Friston, K.J. (2000) Voxel-based morphometry - The methods. *Neuroimage* 11,
1095   805–821
1096   143   Ashburner, J. and Friston, K.J. (2005) Unified segmentation. *Neuroimage* 26, 839–851
1097   144   Ashburner, J. (2007) A fast diffeomorphic image registration algorithm. *Neuroimage* 38, 95–113
1098   145   Ashburner, J. and Friston, K.J. (2009) Computing average shaped tissue probability templates.
1099   *Neuroimage* 45, 333–341
1100   146   Rolls, E.T. *et al.* (2020) Automated anatomical labelling atlas 3. *Neuroimage* 206, 116189
1101   147   Hu, L.T. and Bentler, P.M. (1999) Cutoff criteria for fit indexes in covariance structure analysis:
1102   Conventional criteria versus new alternatives. *Struct. Equ. Model.* 6, 1–55

1103
1104