[bioRxiv, revised version]

Environment-mediated vertical transmission fostered uncoupled phylogenetic

relationships between longicorn beetles and their symbionts

Yasunori Sasakura[1], Nobuhisa Yuzawa[2], Junsuke Yamasako[3], Kazuki Mori[2], Takeo

Horie[4], Masaru Nonaka[5]

1. Shimoda Marine Research Center, University of Tsukuba, Shimoda, Shizuoka 415-

0025, Japan

2. Independent Researcher

3. Institute for Plant Protection, National Agriculture and Research Organization,

Tsukuba, Ibaraki 305-8517, Japan

4. Laboratory for Single-cell Neurobiology, Graduate School of Frontier Biosciences,

Osaka University, Suita, Osaka, 565-0871, Japan

5. Department of Biological Sciences, Graduate School of Science, Tokyo

Metropolitan University, Hachioju, Tokyo 192-0397, Japan

This manuscript consists of 33 pages of text including References and Figure legends;

nine Figures; one Table; five Supplementary Tables.

Running Head: Fungal symbionts of longicorn beetles

Corresponding to: Yasunori Sasakura

Phone: +81-558-22-6605; E-mail: sasakura@shimoda.tsukuba.ac.jp

ABSTRACT

The Coleoptera Cerambycidae (longicorn beetles) use wood under different states

(living healthy, freshly snapped, completely rot, etc.) in a species-specific manner for

their larval diet. Larvae of some Cerambycidae groups have mycetomes, accessory

organs associated with the midgut that harbor fungal symbiont cells. The symbionts are

thought to improve nutrient conditions; however, this has yet to be shown

experimentally. To deduce the evolutionary history of this symbiosis, we investigated

the characteristics of the mycetomes in the larvae of longicorn beetles collected in

Japan. Lepturinae, Necydalinae, and Spondylidinae are the only groups that possess

mycetomes, and these three groups' mycetomes and corresponding fungal cells exhibit

different characteristics between the groups. However, the phylogenetic relationship of

symbiont yeasts does not coincide with that of the corresponding longicorn beetle

species, suggesting they have not co-speciated. The imperfect vertical transmission of

symbiont yeasts from female to offspring is a mechanism that could accommodate the

host-symbiont phylogenetic incongruence. Some Lepturinae species secondarily lost

mycetomes. The loss is associated with their diet choice, suggesting that different

conditions between feeding habits could have allowed species to discard this organ. We

found that symbiont fungi encapsulated in the mycetomes are dispensable for larval

growth if sufficient nutrients are given, suggesting that the role of symbiotic fungi could

be compensated by the food larvae take. *Aegosoma sinicum* is a longicorn beetle

classified to the subfamily Prioninae, which does not possess mycetomes. However, this

species contains a restricted selection of yeast species in the larval gut, suggesting that

the symbiosis between longicorn beetles and yeasts emerged before acquiring the

mycetomes.

KEY WORDS

Longicorn beetle, fungi, yeast, symbiosis, mycetome, Lepturinae

INTRODUCTION

Heterotrophic animals use symbiotic microorganisms to improve their uptake of nutrients. This kind of symbiosis is generally mutualism because host animals provide a space for microorganisms to proliferate comfortably with essential nutrients, while microorganisms support food digestion and supply nutrients (Lindsay et al., 2020; Modolon et al., 2020). Through symbiosis, animals have found ways to adjust to variable environments that are not always conducive to survival. Therefore, symbiotic systems are crucial for understanding the flourishing of animals on Earth.

The wood-eating groups in the coleopteran insects, which comprise the most prominent animal order, have significant relationships with fungi, particularly yeasts (Vega and Blackwell 2005; Gibson and Hunter 2010; Hulcr et al., 2017; Biedermann and Vega 2020). Many yeasts have been isolated from the frass, tunnel, and digestive tubes of coleopteran larvae (Suh et al., 2005; Endoh et al., 2011; Davis et al., 2015). In some groups, adults have a specific structure (the mycangium) that stores fungal cells and passes them to the next generation during oviposition (Tanahashi et al., 2010; Joseph et al., 2021). The fungi proliferate in the tunnel where larvae dwell and are incorporated into the larval digestive tube. These fungi are consumed not only as part of the larval diet (Toki et al., 2012) but are also thought to contribute to creating a suitable condition for the larvae and assisting in the digestion of wood. Indeed, many yeasts isolated from coleopteran larvae have cellulose-, cellobiose- and xylose-digesting activities (Davis et al., 2015).

The Cerambycidae family (longicorn beetles) is a widespread coleopteran group of wood-eating beetles. Larvae of longicorn beetles have a unique fungi-storing

organ, the mycetome, near the anterior end of the midgut (Schomann, 1937; Nardon and

Grenier, 1989; Grünwald et al., 2010). The mycetome consists of a mass of knob-shaped

sacs that attach to the outer membrane (serosa) of the gut. Mycetome sacs are filled with

nutritional particles, and fungal symbiont cells consisting primarily of yeasts dwell in

the lumen or mycetocytes (Jurzitza, 1960). The fungal cells are constantly transferred to

the gut lumen (Grünwald et al., 2010) and are thought to assist in the digestion of wood.

Although the mycetome seems essential for assisting the symbiosis with fungi,

its function is still obscure. Indeed, the mycetome is not conserved among all longicorn

beetle groups (Schomann, 1937). Therefore, this structure may have been acquired in

some groups that require an additional benefit from symbiont fungi to increase their

survival fitness, such as developing efficient digestion assisted by fungi. However, the

advantages that symbiont fungi confer to longicorn beetles still need to be resolved. To

deduce the evolutionary history of the symbiosis facilitated by the mycetome, we

investigated the distribution of this structure in various species from different

subfamilies of longicorn beetles in Japan. We inspected larvae from two families, six

subfamilies, 61 genera, 73 species, and 289 individuals of longicorn beetles. Species of

the subfamily Lepturinae are dominant among the species containing a mycetome;

however, some Lepturinae species do not have this structure. We found that the loss of a

mycetome is correlated with diet choice. The Japanese longicorn beetle groups/species

that had or that lacked a mycetome in our analysis were in good accordance with those

in a previous report of the presence/absence of symbiont yeasts in European species

(Schomann, 1937). *Aegosoma sinicum* belongs to the subfamily Prioninae, whose

members do not have a mycetome. We found that its larvae exhibit symbiosis with

limited yeast species classified into the genus *Scheffersomyces*, suggesting that

symbiosis with yeasts preceded the acquisition of mycetomes during the evolution of

Cerambicydae.

MATERIALS AND METHODS

Animals

Most of the longicorn beetle larvae were collected from dead trees. We used

cultured larvae to analyze the species whose larvae are difficult to capture from the

wild. Their species names were identified by the cuticle structure on the thorax

(Cherepanov, 1988; Koiwaya, 2017). When the thoracic cuticle structure could not

identify species, genomic DNA was isolated from larval somatic tissues with a Wizard

Genomic DNA Purification kit (Promega), followed by PCR amplification of 28S rDNA

with ExTaq DNA polymerase hot start version (Takara-bio) and with the primers 5'-

GAACTTTGAAGAGAGAGTT-3' and 5'-CAGGCATAGTTCACCATCTT-3'. After

the PCR fragments were purified with the Qiaquick Gel Extraction kit (Qiagen), the

sequence was determined by the conventional Sanger method, which was used to

identify species with the aid of 28S rDNA sequence isolated from adults (Accession

number PP375410-PP375497). The larval midgut was surgically isolated with fine

tweezers in phosphate-buffered saline (PBS) to observe mycetomes and collect gut

materials. Frass was collected from the tunnel on wood with cotton swabs. Larvae in the

tunnel were isolated together with frass to identify species.

Identification of fungi

Mycetome contents, gut materials, frass or fluid from adult genitalia were

spread on potato dextrose agar (PDA) plates (4-1126-05 Sani-Spec, Japan or 05709

Shimazu Diagnostics, Japan), including 15 μg/ml chloramphenicol and 50 μg/ml

ampicillin. 1L PDA agar broth contains 4g potato extract, 20g glucose, and 15g agar.

Eggs were immersed in PBS for a few minutes, and the supernatant was spread onto

PDA plates. The plates were incubated for a few days at 25ºC or longer at 4ºC. DNA

was extracted from yeast or yeast-like colonies by a microwave-based method (Izumitsu

et al., 2012) and was subjected to amplification of the internal transcribed spacer (ITS)

rDNA with ExTaq DNA polymerase hot start version (Takara-bio) and with the primers

5'-CTTGGTCATTTAGAGGAAGTAA-3' and 5'-TCCGTAGGTGAACCTGCGG-3'

according to the previous method (White et al., 1990; Gardes and Bruns 1993). In this

analysis, up to four colonies on a single plate were soaked together in 100 μl Tris-Cl

buffer if the colonies were available, before DNA extraction. Sequences were

determined from the amplified bands after purification with a Qiaquick Gel Extraction

kit (Qiagen). When sequences could not be determined due to the low quality of the

chromatographs, each colony was subjected to sequencing analysis separately. When the

mycetome contents did not yield colonies because the symbiont fungi were

unculturable, we isolated DNA from mycetomes with Isoplant II (Nippon-Gene)

according to the manufacturer's instructions, followed by PCR amplification and

sequencing of rDNA of fungal cells.

The identified rDNA sequences of fungal cells were compared with the

reported DNA sequences in the standard nucleotide collection (nr/nt) database using the

blastn program (Altschul et al., 1997). The species names of the source organisms of the

DNA sequence that exhibited the highest similarity in the blastn analysis are listed in

the "Blastn Best Hit" column in Table S1. We used the source organism name to refer to

the isolated fungal cells for further analyses. When an undescribed species (for example,

Scheffersomyces sp.) exhibited the best-hit, its strain identifier (such as strain Y13) is

referred. We have to note that it does not always guarantee the actual yeast species

because in many cases, our rDNA sequences did not exhibit a perfect match to a known

rDNA sequence (Table S1, "Blastn score" column).

The internal transcribed spacer (ITS) of *Scheffersomyces* sp. Y1052_SICYLG3

has one *Eco*RI restriction site. The yeasts isolated from adult genitalia of *Leptura*

ochraceofasciata females were identified by the presence of the *Eco*RI site. After PCR

amplification, the PCR fragments were digested by *Eco*RI, followed by agarose gel

electrophoresis.

Mycetome-possessing Lepturinae and mycetome-lacking longicorn beetle

groups are compared according to the number of larvae possessing symbiont yeasts, and

the number of yeast species characterized in these groups. We performed two-tailed

Chi-square test using R x64.4.1.2 and RStudio softwares (R Core Team 2021; RStudio

Team 2020). In the latter comparison, we counted the "Blastn Best Hit" names of yeasts

that formed colonies on PDA plates. *Encyclops olivaceus* and species in Necydalinae

and Spondylidinae have unculturable symbiont fungi in their mycetomes. On the other

hand, we collected only culturable yeasts from larvae in a mycetome-lacking group by

streaking gut contents or frass onto the PDA plates. For unbiased comparisons, the

identified unculturable fungi were excluded from these counts.

Phylogenetic analyses

The sequences were aligned using the M-COFFEE program

(https://tcoffee.crg.eu/apps/tcoffee/do:mcoffee; Notredame et al., 2000; Wallace et al.,

2006). After removing the residues at the ends that seemed not to align properly due to

the differences in the lengths between the sequences, maximum likelihood trees were

constructed with the MEGA software version 11 (Tamura et al., 2021). Trees were

tested with 1,000 bootstrap pseudoreplicates. The aligned nucleotides used for

phylogenetic constructions are shown in Table S2.

The phylogenetic tree for Lepturinae beetles was constructed using 28S rDNA

sequence. The phylogenetic trees of the yeasts isolated from longicorn beetle larvae

were constructed as follows. Referring to the results of the blastn search, the ITS

sequences of the yeast from the Lepturinae, Spondylidinae, and *Aegosoma sinicum*

larvae were used for the phylogenetic tree construction. When a yeast species was

detected from more than one larvae of the same longicorn beetle species, only one

sequence was used for the analysis. The ITS sequences from *Hyphopichia rhagii*

(isolated from *Rhagium femorale* and *R. japonicum*), *Candida wanchemiae* (isolated

from *Anastrangalia scotodes*), and uncultured fungus clone OUT_F389 (isolated from

Megasemum quadricostulatum) are considerably shorter than those from other yeasts

and were removed from the analysis in view of their potential influence on alignment.

Culture of Cerambycidae larvae

Our artificial diet was developed based on the work of Gardiner (1970). The

artificial diet Insecta FII (Nosan Corporation, Japan) prevents the growth of fungi with

the aid of anti-fungal chemicals that are not open by the company. Nine grams of

Insecta FII, 6 g of dead *Symplocos coreana* powder, and 15 g of cellulose (ISC powder)

were kneaded in the following liquid to create ISC paste: 30 or 45 ml of water, 30 or 45

ml of PDA dissolved in water, or 30 or 45 ml of 1x PDA suspended with one of two

yeasts, *Candida vrieseae* or *Scheffersomyces* sp. Y1052_SICYLG3. These yeasts were

cultured on PDA plates until colonies grew to cover the surface of the broth. The

colonies were recovered with a cotton swab, and suspended in PDA liquid so that the

liquid becomes cloudy. We did not quantify the suspended yeast cells. The yeasts were

expected to improve larval growth; however, the effect was questionable and was not

analyzed in detail. The ISC paste was spread in a polypropylene box and was

autoclaved (20 min at 105°C) for sterilization. Eggs, hatched larvae or larvae collected

from dead woods were placed on the autoclaved diet and were kept at room temperature

until observation. For the introduction of selected yeast to larvae, eggs were sterilized

for more than one day by placing them on sterilized ISC paste. The powder of PDA

broth was suspended in distilled water and kept until agar powder sank. Then the

supernatant was collected to obtain 1 x PDA liquid. A 70 mm x 30 mm x 2 mm

rectangular wood plate of *Symplocos coreana* was sterilized by autoclaving, then

applied with 1 x PDA liquid in which yeast cells were suspended by the same method as

ISC paste. Sterilized eggs were cultured on the yeast-applied wood plate until hatching.

Hatched larvae were collected and cultured in sterilized ISC paste for about one month

until they became sufficiently large for the surgical isolation of mycetomes.

RESULTS

Distribution of mycetomes in Japanese Cerambycidae

To deduce the evolutionary history of the mycetome in longicorn beetles, we

investigated the distribution of mycetomes in the Japanese species (Fig. 1). The majority

of the cerambycid species we inspected representing the subfamilies Lepturinae,

Necydalinae, and Spondylidinae, had mycetomes near the anterior end of the larval

midgut (Fig. 1 and Table 1). There are group-specific characteristics in the shapes of

mycetomes. The mycetomes of most Lepturinae species (Fig. 1B-C) and all

Spondylidinae (Fig. 1D) adhered tightly to the serosa of the midgut. By contrast, the

large mycetomes of Necydalinae species were well separated from the serosa with a

connection through several thin tubes (Fig. 1E). The mycetomes of Lepturinae form a

single circular ring surrounding the serosa (Fig. 1B). In contrast, those of Spondylidinae

larvae form two narrower rings (Fig. 1D). The characteristics of the appearance of

mycetomes in Lepturinae and Spondylidinae agree with those in the previous report

(Buchner 1965).

We found mycetomes in 22 out of 31 species of Lepturinae and all 3 of the

examined Necydalinae species (Table 1). *Paragaurotes doris*, *Japanocorus*

caeruleipennis, *Brachyta bifasciata,* and all *Pidonia* lacked mycetomes (Fig. 1F and

Fig. 2). In these cases, the mycetomes were probably lost secondarily during speciation.

Among the mycetome-lacking Lepturinae, larvae of *B. bifasciata* eat the roots of living

Paeonia (Table S1, "Feeding habit" column). All of the other mycetome-lacking

Lepturinae are inner bark-eaters at the larval stage. Inner bark-eating Lepturinae do not

necessarily lack mycetomes; two *Rhagium* species are inner bark-eaters but have

mycetomes. We found mycetomes in all examined Spondylidinae species (n=5; Table

1).

Characteristics of yeasts or yeast-like fungi in mycetomes

We found that mycetomes and fungi dwelling in them had group-specific

characteristics. Each lobe of the Lepturinae mycetome contains tear-shaped fungal cells,

a membranous structure, and particles much smaller than yeast cells (Fig. 3A and B,

B'). The density of yeast cells in a mycetome is usually not high, and the cells do not

fully occupy the lumen. When cultured on potato dextrose agar (PDA) plates, almost all

of the fungi from the mycetomes of Lepturinae larvae formed colonies (Fig. 3C), and

their rDNA sequences indicated that they were yeasts.

We classified yeasts isolated from the Lepturinae mycetome or gut by

determining the sequence of the ITS site of mitochondria rDNA (Table S1). These

rDNA sequences were used as queries of blastn searches to estimate the species of

yeasts (Altschul et al., 1997). In the following section, we refer to the characterized

yeasts as the species exhibiting the highest similarity to the rDNA sequences in the

blastn analyses for simplicity.

In almost all cases, only a single yeast species was detected from the mycetome

of a Lepturinae larva (87%, n=109/125; Table 1). However, seven larvae had more than

one yeast species in the mycetome (rows 6, 8, 73-76, 107 in Table S1). Moreover, the

rDNA sequence derived from the mycetomes of single larvae exhibited nucleotide

variations in three cases (rows 10, 23, and 119 in Table S1). These results suggest that

the yeast cells in mycetomes are not always clones. There are species-specificities

between longicorn beetles and yeasts. For example, two Japanese *Rhagium* species had

a yeast whose rDNA exhibited the best nucleotide sequence similarity to that of

Hyphopichia (or *Candida*) *rhagii*, also found in a *Rhagium* species in Germany

(Grünwald et al., 2010).

The phylogenetic analysis (Fig. 4) classified the yeasts isolated from

Lepturinae mycetomes into eight species or unidentified species (*Scheffersomyces*

ergatensis, *Scheffersomyces paraergatensis*, *Scheffersomyces coipomoensis*,

Scheffersomyces sp. Y1052_SICYLG3, *Scheffersomyces* sp. Y6, *Scheffersomyces* sp.

Y13, *Candida tenuis*, *Candida vrieseae*). Two indentified species (*Scheffersomyces* sp.

Y6 and Y13) and *S. paraergatensis* are included into the branch of *S. ergatensis*. We did

not include two species (*Hyphopicha rhagii* and *Candida wanchemiae*) in this analysis

due to their short ITS sequences. In total, ten culturable yeast species were isolated from

22 mycetome-possessing Lepturinae species, suggesting that some yeast species were

shared among different Lepturinae species (Table 1, Fig. 2, and Fig. 4). However, the

Lepturinae species that share the same symbiont yeast are not always phylogenetically

close to each other (Fig. 2). For example, the larvae of *Leptura ochraceofasciata*

isolated from both hardwoods and conifers almost always had *Scheffersomyces* sp.

Y1052_SICYLG3 (Table S1). This yeast was also found in the larvae of *Caraphia*

lepturoides, although *L. ochracerofasciata* and *C. lepturoides* are phylogenetically

distant. Larvae of *Pyrrhona laeticolor*, *Ohbayashia nigromarginata*, *Ephies japonicas*,

Pigostrangalia dulcis, and *Mimostrangalia kurosonensis* had the yeast related to

Candida vrieseae in their mycetomes. All of these Lepturinae species use the hardwood

group Symplocaceae as their larval hosts. This was not due to the ubiquity of *C.*

vrieseae in the dead Symplocaceae tree since larvae of the stag beetle *Aesalus asiaticus*

isolated from a dead *Symplocos coreana* together with *Pyrrhona laeticolor* had a

different yeast (*Scheffersomyces ergatensis*) in their digestive tube (Table S1, stag

beetle). However, the relationships between yeast species and host plants of longicorn

beetles are not expected to be rigid since *Anastrangalia scotodes* and *Strangalia*

koyaensis use several conifers as hosts possessed *Candida vrieseae* (Table S1 and Fig.

4).

The characteristics of the mycetome contents of the Lepturinae species

Encyclops olivaceus and the three examined Necydalinae species were quite different

from those of the other Lepturinae species. *Encyclops* and Necydalinae mycetomes were

fully occupied by fungal cells, and we could not find a membranous structure (Fig. 3).

The elongated fungal cells in these Necydalinae mycetomes were obviously larger than

the yeast cells of Lepturinae, and the Necydalinae symbionts did not form a colony on

PDA plates. The unculturable symbionts of *Encyclops* are similar in shape but smaller

than those of Necydalinae species (Fig. 3E). All analyzed larvae of Necydalinae species

are dependent on dead hardwood trees (Table S1). To further determine whether the

symbiont characteristics are shared among Necydalinae species, we examined the

symbiont of *N. sachalinensis*, which is dependent on the dead part of living conifers.

Because its larvae were unavailable, we isolated fungal cells from the oviducts of *N.*

sachalinensis females. The fungal cells had the same characteristics as those of the

fungi isolated from the other Necydalinae larvae (Fig. 3H' and Table S3).

We attempted to characterize the fungi from *Encyclops olivaceus* and

Necydalinae mycetomes by directly isolating DNA from their mycetomes. The DNA

fragments amplified from their mycetomes did not show sufficient similarity to those of

any known fungi (Table S1). The DNA fragments from *Encyclops olivaceus* and

Necydalis sachalinensis exhibited homology to the genome sequences of the other

Coleopteran species (Table S1 and S3), remaining the possibility that these DNAs were

not derived from the symbionts. Because Lepturinae symbiont yeasts yielded DNA

bands specific to them by the same methods, these results confirm that the symbionts of

the *Encyclops olivaceus* and Necydalinae species are distinct from yeasts.

The contents of Spondylidinae mycetomes were similar to those of typical

Lepturinae in that it contained membranous structures, and the density of fungal cells

was not high (Fig. 3J). The shape of Spondylidinae fungal cells was rod-like (Fig. 3K-

K'). None of these fungi made a colony on PDA plates (Fig. 3L). Unlike Necydalinae,

Spondylidinae fungal cells are classified into the yeast group

Debaryomyces/*Schwanniomyces* (Table S1 and Fig. 4). These yeasts derived from

Spondylidinae form a single clade in the phylogenetic tree of the yeasts isolated from

mycetome-positive Lepturinae/Spondylidinae (Fig. 4). Therefore, there is broad group-

specificity in the choice of symbiont fungi when Lepturinae, Necydalinae, and

Spondylidinae are compared. However, when viewed from the species level, Lepturinae

longicorn beetles' phylogenetic relationships do not reflect in those of the symbiont

yeasts (Fig. 2). This incongruence is our next target to address.

Imperfect delivery of yeasts across generations

It has been suggested that the maternal parent provides the symbiont fungi in

the larval mycetome (Schomann, 1937; Douglas, 1989). We characterized yeasts

isolated from eight species of Lepturinae adult females. Although not perfectly matched,

the adult females had yeast species isolated from the same species of larvae (Fig. 5;

Table S3), confirming the vertical transfer of symbionts. We found that yeasts in the

female genitalia could be easily excreted by gently pushing the oviduct. This suggests

that yeasts are smeared around the eggs during oviposition, as previous studies

described (Schomann, 1937; Douglas, 1989). Hatched larvae may incorporate yeasts

with egg shells and nearby wood into their gut. Indeed, 1st instar larvae of *Ohbayashia*

nigromarginata and *Necydalis formosana* a few days after hatching already had

mycetome-like knobs at the anterior midgut and fungal cells in the structures (Fig. 6).

This process of yeast delivery does not guarantee the accurate transmission of

maternal yeasts to their descendants because larvae that accidentally move from the

oviposited area before yeast uptake may have a chance to incorporate a different yeast

species from the environment. For example, larvae of *L. ochraceofasciata* almost

always have *Scheffersomyces* sp. Y1052_SICYLG3; however, we found a single larva

with the yeast *Candida vrieseae* (Table S1). This individual was found in a dead tree of

Symplocaceae. As we stated above, Lepturinae species that depend on this tree group

usually have *Candida vrieseae*, and the larva of *L. ochraceofasciata* is suspected of

having accidentally incorporated *Candida vrieseae* supplied by another Lepturinae

species. We also found two larvae of *L. ochraceofasciata* and *P. laeticolor*, and five

larvae of *Ohbayashia nigromarginata* that had the yeast *Scheffersomyces coipomoensis*

(Table S1). These larvae were isolated from one dead tree of *Symplocos coreana* at Mt.

Amagi. We isolated *Scheffersomyces coipomoensis* from yellowish dead trees of

Symplocos coreana on the same mountain, suggesting that this yeast dwells in the

environment and could have been accidentally incorporated into the larvae.

If Lepturinae larvae could incorporate yeasts from environment, it would be

possible to generate yeast-free larvae or larvae having a yeast species different from that

dominantly possessed by larvae in the wild populations, by removing maternally

supplied yeasts. To show this, we developed a way to culture larvae under a condition

chemically suppressing fungal growth. After oviposition, we placed eggs on an artificial

diet containing anti-fungal chemicals for sterilization, so that the hatched larvae did not

come into contact with external yeasts during and after birth. Larvae of *Leptura*

ochraceofasciata and *Stictoleptura dichroa* grew well under this condition (Fig. 7A).

They had brownish mycetomes near the anterior end of the midgut (Fig. 7B and D). The

position of their mycetomes was identical to those of larvae from the wild. Microscopic

observations showed that these brown mycetomes did not include yeasts (Fig. 7C and

E), which was confirmed by culturing mycetome contents on PDA plates (Fig. 7C, inset), suggesting that the yeasts needed to be supplied externally for larvae to acquire yeasts in their mycetomes. This experiment also indicates that mycetomes can develop without yeasts.

In the above culture experiments, some of the larvae had yeasts not found in the mycetomes of larvae isolated from wild populations. In the experiment with *L. ochraceofasciata*, we discovered that a few larvae (12.5%, n=2/16) had *Meyerozyma caribbica*. *Meyerozyma* is an indigenous yeast group arising in our laboratory. In the experiment using *S. dichroa*, a portion of eggs hatched on the wood onto which their mothers had laid eggs before sterilization. We cultured the pre-hatched eggs and 1st instar larvae in the same container. After one month, we observed that the larvae were divided into three groups: larvae with yeast-free mycetomes (n=5; Fig. 7D and E), larvae with mycetomes that included yeast cells with a low density (n=4; Fig. 7F and G), and larvae with relatively yeast-rich mycetomes (n=9; Fig. 7H and I). The yeast species isolated from the yeast-poor mycetomes differed from those isolated from yeast-rich mycetomes: the former was *Meyerozyma caribbica,* while the latter was *Scheffersomyces ergatensis*. *S. ergatensis* is the default species of this longicorn beetle in the wild (Table S1). We suspected that the larvae with *S. ergatensis* hatched on the egg-laid wood and had a chance to incorporate maternally gifted yeast.

To investigate the above hypothesis about environmental yeast acquisition, we next tested whether we could deliver a specific yeast species into mycetomes. In this experiment, sterilized eggs were placed on a wood plate to which a yeast species was applied. After birth, we transferred the larvae to a plate with the artificial diet to avoid further yeast incorporation during growth until the larvae reached a sufficient size to

isolate mycetomes. We examined 33 larvae, and 22 had the given yeast species (Table

S4). These yeasts were never detected in the control groups to which we did not provide

the yeasts (0%, n=50), suggesting that this experiment successfully replaced the

symbiont yeast with the arbitrary yeast. We concluded from these results that Lepturinae

longicorn beetles acquire yeasts externally after hatching. Although the yeasts are

usually maternally supplied, a different yeast could be obtained from the environment.

Lepturinae mycetomes protect symbionts from anti-fungal chemicals in food

397   During the culture of longicorn beetle larvae with the artificial diet, we found

that the mycetomes of Lepturinae species protect the symbionts from the anti-fungal

chemicals in the artificial diet. A larva of *Etorofus vicaria* and *Leptura ochraceofasciata*

maintained their symbiont yeasts in the mycetomes even after culturing for about one

year (Fig. 8A and Table S5). By contrast, symbionts of *Necydalis formosana* larvae

were eliminated from their mycetomes (Fig. 8B) even though they increased their

weight during the culture (Table S5). These results suggest that the mycetome of this

species is weaker in protecting symbionts from chemicals taken with the food.

Yeast transmission between generations in the longicorn beetles without mycetomes

407   There is an association between mycetome-possessing Lepturinae longicorn

beetles and yeast species. Our next question is how their symbiosis started, and notably,

how mycetomes contributed to the evolution of the symbiosis. To address these

questions, we investigated yeasts in the digestive tubes and frass of mycetome-lacking

longicorn beetles. We observed 135 larvae of mycetome-lacking longicorn beetles

(including Lepturinae) corresponding to 46 species from five families/subfamilies

(Table 1). Yeasts were detected in 46% (n=63/135) of larvae. This score is significantly

low compared to that of mycetome having Lepturinae (92%, n=116/125, $p<0.001$),

suggesting that mycetomes have a role in stabilizing symbiosis with yeasts. We isolated

in total 45 culturable yeast species from 46 mycetome-lacking longicorn beetle species

(Table 1). This score is two times more than mycetome-having Lepturinae species (10

out of 21 excluding *Encyclops olivaceus* whose symbiont is unculturable; $p<0.001$),

suggesting that yeasts habiting in the mycetome-lacking species are more variable than

mycetome-possessing species. These wobbles could be explained if mycetome-lacking

longicorn beetles do not usually possess a way to limit the species of symbiont yeasts,

such as a delivery system between generations.

We found an exceptional mycetome-lacking species that could transmit specific

yeasts between generations. *Aegosoma sinicum* is a kind of Prioninae that does not have

mycetomes (Fig. 1G). Two yeast groups belonging to the genus *Scheffersomyces* were

repeatedly isolated from larvae of *A. sinicum* collected in different locations and from

different host trees (Table S1 and Fig. 9), suggesting a strong correlation between *A.*

sinicum and the yeasts. Individual *A. sinicum* larvae exclusively possessed either of the

two *Scheffersomyces* groups.

Two stand out among the possible explanations for the relationship between *A.*

sinicum and *Scheffersomyces*. The first is that *A. sinicum* females might lay eggs on the

host trees in which the yeasts habit. The second possibility is that the yeasts were

transferred from adult females to their progeny. We isolated the colonies of the same

Scheffersomyces yeasts from the genitalia of females and the egg surfaces of *A. sinicum*

(Table S2). Therefore, it is likely that *A. sinicum* vertically transfers yeasts between

generations without mycetomes, although we cannot rule out the first possibility.

DISCUSSION

Many dead wood-eating beetles depend on yeasts or related fungi at their larval stage (Davis, 2015). Although the role of this symbiosis has yet to be fully understood, its suspected function is to increase efficiency in the nutrient absorption from dead wood that is less nutritious than living plants. Larvae of longicorn beetles, which form one of the most diversified groups in the Coleoptera, evolved a characteristic fungi-storing organ, the mycetome, at the midgut (Schomann, 1936; Buchner, 1965). Among six subfamilies of Cerambycidae in Japan, the species in Lepturinae, Necydalinae, and Spondylidinae have mycetomes. The Japanese species in Disteniidae, which is a sister group of Cerambycidae, does not have the mycetome, confirming the previous report (Svacha and Lawrence 2014). Svacha and Lawrence (2014) also reported that the larvae in the families Vesperidae and Oxypeltidae do not have mycetomes. Because mycetome-absent families are dominant in longicorn beetles, mycetome was likely to have evolved in the subfamilies of Cerambycidae. If this is true, the mycetome emerged at least twice in Spondylidinae and Lepturinae/Necydalinae according to the phylogenetic tree of the Cerambycidae subfamilies (Fig. 1; Haddad et al., 2018; Nie et al., 2021). The emergence of the mycetome in Lepturinae and Necydalinae is likely to have been the single event at the root of these groups. This is supported by the recent phylogenetic study suggesting that Necydalinae is regarded to be included in Lepturinae (Nie et al., 2021). However, we cannot rule out the possibility that these mycetomes were evolved independently because Necydalinae mycetomes have characteristics that are not seen in Lepturinae mycetomes.

The symbionts of Necydalinae species and that of *Encyclops olivaceus* are

unculturable and fully occupy the lumens of the mycetomes. The shared characteristics

of the mycetomes and symbionts among *Encyclops* and Necydalinae species suggest

that they are phylogenetically related. The previous study showed that the Lepturinae

species *Oxymirus cursor* has gigantic mycetomes fully occupied with slender tear-

shaped and unculturable symbiont fungal cells (Buchner 1965). These described

characteristics are also similar to those in Necydalinae. The phylogenetic relationships

between Necydalis and these Lepturinae species should be considered in future studies.

The emergence of mycetomes in Spondylidinae and Lepturinae/Necydalinae

suggests that the acquisition of this organ is beneficial for these groups. The benefits of

this symbiosis need to be considered from the respective perspectives of the symbionts

and longicorn beetles. Establishing symbiotic relationships between Cerambycidae

through mycetomes would have apparent advantages for fungi. Our study and previous

studies have shown that the mycetomes of some species include fungi that cannot

proliferate on a PDA plate, although such a plate is sufficient for many yeasts to grow

(Müller, 1934). From this result, one of the functions of the mycetome is to provide

fungi with nutrients necessary for their proliferation.

Moreover, we here showed that the mycetomes of larval Lepturinae possessed

yeast cells even when the larvae ate an artificial diet containing anti-fungal chemicals,

which can inactivate yeast cells by overnight incubation, suggesting that another role of

the mycetome is to protect fungi from the chemicals that could be included in plants or

secreted from competitive fungi. Unlike larvae of the species in Lepturinae, *Necydalis*

formosana larvae lost symbionts from their mycetomes when cultured with a diet

containing anti-fungal chemicals, suggesting that the mycetomes of Lepturinae and

Necydalinae are functionally and structurally different. In most cases, a single yeast

species occupies the mycetome of a larva, even though the larvae have a chance to

incorporate external yeasts or yeast-like fungi that are spontaneously present in dead

trees. Therefore, the mycetome can protect symbiotic fungi from competition with other

fungi.

In contrast to the advantages for fungi, the benefits conferred to the longicorn

beetles remain obscure. Because the mycetome is a longicorn beetle-derived organ and

fungi are not necessary for its formation, symbiosis with fungi is advantageous for

longicorn beetles. It has been proposed that symbiont fungi are nutritionally beneficial

for host beetles, as in the case of many wood-eating insects and their symbiotic gut

microorganisms (Gibson and Hunter, 2010). Our study provided evidence suggesting

that symbiosis is related to nutrition and/or feeding habits. All species in Spondylidinae

examined in this study and most of the mycetome-possessing species in Lepturinae eat

brown rot supwood. This shared food choice would have been made possible by the

acquisition of mycetomes. Some Lepturinae species do not have mycetomes, and the

loss of this organ occurred over several rounds of speciation (Fig. 2). The loss of the

mycetome is also strongly associated with the choice of diet. All mycetome-lacking

Lepturinae species eat inner bark or living roots. Inner bark and living roots have

different nutritional or organismal conditions from those in rot supwood. For example,

inner bark and living plant tissue contain more nutrients than supwood (Schowalter

1992). The change in diet preference may have enabled the Lepturinae species to be free

from mycetome-dwelling yeasts due to the improved nutrient uptake. When fed an

artificial diet, we also showed that mycetome-possessing Lepturinae larvae can grow

without symbiont yeasts. For this experiment, we used an artificial diet based on the

silkworm diet, which is much richer in nutrients than dead wood. Therefore, the species

having mycetomes do not require symbiont yeasts under such improved nutritional

conditions, which paradoxically supports the idea that the benefits from symbiont fungi

are associated with nutrients. To confirm more conclusively, we need to compare the

growth rate between larvae having mycetomes with or those having mycetomes without

yeast when fed a diet having low nutritional value. Our method of obtaining larvae with

yeast-free mycetomes will assist in future experiments.

Among the Cerambycidae group, many Lamiinae and Cerambycinae species

are known to eat inner bark of fresh dead trees or live trees (Svacha and Lawrence

2014). The species in Lamiinae and Cerambycinae do not have mycetomes, probably

because their diets are more nutritive than those of mycetome-possessing groups.

Lamiinae is the sister group of Spondylidinae, which is dominant in mycetome-

possessing species. Unlike Lamiinae, Spondylidinae longicorn beetles in Japan depend

on brown rot supwoods (Table S1). The symbiosis strategy with fungi in these two

groups may have differed in their diet choices after the split. We have to note that the

tendency of diet choice is not always supported since several mycetome-lacking

Prioninae (such as *Psephactus remiger*), Cerambycinae (such as *Thranius variegatus*,

Paraclytus excultus, *Xylotrechus cuneipennis*) and Lamiinae (such as *Nanohammus*

rufescens) species eat strongly rot wood or outer bark. These species may have another

way to strengthen their nutritional condition, including establishing symbiosis with

another microorganism group, such as bacteria, as reported previously (Grünwald et al.,

2010; Reid et al., 2011).

In our experiments, mycetome-lacking larvae of longicorn beetles tended not to

have yeast in their gut compared to those in mycetome-possessing larvae (Table 1),

suggesting that many of mycetome-lacking species do not transfer yeasts between

generations. When yeasts were found in these larvae, the assortment of yeast species

usually varied among individuals. A possible explanation is that mycetome-lacking

larvae uptake yeasts from their environment, and yeast species that can withstand larval

digestive systems proliferate in the gut. Such environment-dependent acquisition of

yeast is likely to be the ancestral form of the symbiosis between longicorn beetles and

yeasts, as was suggested previously (Douglas, 1989), and the external delivery of yeasts

observed in mycetome-possessing species is probably the remnant of this ancestral

symbiosis. Because the mycetome-lacking longicorn beetle *A. sinicum* can share a

limited number of yeast species between generations, the emergence of mycetomes is

not requisite for longicorn beetles to initiate symbiosis with specific yeasts.

Because mycetome-possessing longicorn beetles tend to have a single preferred

yeast or fungal species/group as their symbiosis partner, mycetomes could limit the

symbiont to a specific species. This role of mycetomes should be in concert with the

development of the sac in the female genitalia that stores symbiont yeasts (Schomann,

1937; Buchner, 1965). With the aid of the mechanism of vertical transmission, the

mycetomes and symbionts have group-specific features. Larvae of the species in

Lepturinae tend to have yeasts culturable on the PDA plate and the yeast cells do not

fully occupy the lumen of the lobes of mycetomes. Larvae of the species in

Spondylidinae have elongated yeasts unculturable on the PDA plate. Larvae of the

species in Necydalinae have unculturable fungal cells that are larger than the yeast cells

in Lepturinae. The Necydalinae mycetomes are well-developed whose lumen is fully

occupied by the fungal cells. The presence of these group-specific characteristics

suggests that some limitations worked to prevent the loss of these characteristics even as

speciation occurred within these groups. It is possible that the constraint is associated

with group-specific diet choices. These group-specific diet choices include the

hardness/softness and degree of moisture of rotten wood, and dependency on specific

fungal infestation (Svacha and Lawrence, 2014).

However, our data showed that the vertical transmission of symbionts between

generations is imperfect, and larvae sometimes have a different yeast from the symbiont

species. These exceptions could be explained by somewhat imperfect manner of yeast

transmission between generations. Moreover, this accidental "mismatch" could disturb

the phylogenetic relationship between symbiont yeasts and longicorn beetle species.

During the speciation of longicorn beetles, the acquisition of symbiont fungi from the

environment allowed the diversification of symbiont yeast species that is not always

associated with longicorn beetle speciation. We found that the Lepturinae species using

Symplocaceae as their host predominantly have the symbiont yeast related to *Candida*

vrieseae, even though these Lepturinae species are not monophyletic. Probably the use

of the same dead tree promoted the sharing of the yeast species between the beetle

species. The mismatch between the phylogenies of host beetle species and symbiont

yeasts has also been reported in stag beetles (Kubota et al., 2020), suggesting that this

incongruent evolution is shared among Coleopteran groups. Such incongruent evolution

was probably permitted because symbiont yeasts have similar physiological

characteristics and because the function of yeasts may be to provide inessential dietary

supplementation, which may have been dispensable when longicorn beetles switched

their food from dead wood to inner bark.

ACKNOWLEDGMENTS

We are grateful to Drs. Ryutaro Iwata, Rikiya Endo, Takahito Watanabe, Yosuke

Degawa and Benjamin Harvey for helpful discussions. We also thank Dr. Ryosuke

Yamamoto, Tomoji Mikage, Mitsuru Nonaka, Naho Nakao, Haruto Nakao, Keiichi

Sasaki and Sigeo Tsuyuki for helping us with the sample collection.

COMPETING INTERESTS

The authors have no competing interest to declare.

AUTHOR CONTRIBUTIONS

YS, NY, KM, and MN conceived and designed the study. YS and MN performed most

experiments. NY, and KM performed sampling. JY provided the phylogenetic analyses.

TH performed molecular experiments. YS wrote the manuscript with the aid of the other

authors.

SUPPLEMENTARY MATERIALS

Supplementary materials for this article are available online.

Table S1.

Symbiont fungi isolated from larvae of longicorn beetles in Japan. Each row

corresponds to the results of a single larva. "Culture" indicates the cultured larvae used

for the analyses. The digits at the columns of "*Aegosoma sinicum*" correspond to those

in Figure 9. The numbers of colonies on the plates are indicated when a few colonies

appeared.

Table S2.

Alignment of nucleotide sequences used for phylogenetic tree construction.

Table S3.

Symbiont fungi isolated from female adults of longicorn beetles in Japan. Each row

corresponds to the results of a single female.

Table S4.

Artificial transfer of yeasts to larvae. The same Exp number indicates the experiment

using the eggs from the same female individual.

Table S5.

Possession of symbiont cells in the mycetome after long culture in the artificial diet

with anti-fungal chemicals.

REFERENCES

Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997)
Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs. Nucleic Acids Res 25: 3389-3402.
Biedermann PHW, Vega FE (2020) Ecology and Evolution of Insect-Fungus
Mutualisms. Annu Rev Entomol 65: 431-455.
Buchner P (1965) Symbiosis in insects feeding on cellulose and similar substances. In:
Endosymbiosis of animals with plant microorganisms. Interscience Publishers, NY,
London, Sydney.
Cherepanov AI (1988) Cerambycidae of Northern Asia. Brill, Leiden, Netherlands.
Danilevsky M (2020) Catalogue of Palaearctic Coleoptera Volume 6/1 (Revised and
updated second edition) Chrysomeloidea I (Vesperidae, Disteniidae, Cerambycidae)

Brill, Leiden, Netherlands.

Davis TS (2015) The ecology of yeasts in the bark beetle holobiont: a century of

research revisited. Microb Ecol 69: 723-732.

Douglas AE (1989) Mycetocyte symbiosis in insects. Biol Rev 64: 409-434.

Endoh R, Suzuki M, Okada G, Takeuchi Y, Futai K (2011) Fungus symbionts

colonizing the galleries of the ambrosia beetle Platypus quercivorus. Microb Ecol 62:

106-120.

Gardes M, Bruns TD (1993) ITS primers with enhanced specifity for Basidiomycetes:

application to identification of mycorrhizae and rusts. Mol Ecol 2: 113-118.

Gardiner LM (1970) Rearing wood-boring beetles (Cerambycidae) on artificial diet.

Can Entomol 102: 113-117.

Gibson CM, Hunter MS (2010) Extraordinarily widespread and fantastically complex:

comparative biology of endosymbiotic bacterial and fungal mutualists of insects. Ecol

Lett 13: 223-34.

Haddad S, Shin S, Lemmon AR, Lemmon EM, Svacha P, Farrell B, Slipinski A,

Windsor D, McKenna DD (2018) Anchored hybrid enrichment provides new insights

into the phylogeny and evolution of longhorned beetles (Cerambycidae). Syst

Entomol 43: 68-89.

Hulcr J, Stelinski LL (2017) The Ambrosia Symbiosis: From Evolutionary Ecology to

Practical Management. Annu Rev Entomol 62: 285-303.

Izumitsu K, Hatoh K, Sumita T, Kitade Y, Morita A, Gafur A, Ohta A, Kawai M,

Yamanaka T, Neda H, Ota Y, Tanaka C (2012) Rapid and simple preparation of

mushroom DNA directly from colonies and fruiting bodies for PCR. Mycosci 53:

396-401.

Joseph R, Keyhani NO (2021) Fungal mutualisms and pathosystems: life and death in

the ambrosia beetle mycangia. Appl Microbiol Biotechnol 105: 3393-3410.

Jurzitza G (1960) Zur systematik einiger Cerambycidensymbionten. Arch Mikrobiol 36:

229-243.

Koiwaya S (2017) Kakedashi tengyuya no hisokana yokubou (1) [The secret desires of

a longicorn beetle collector] Gekkan-Mushi 560: 28-36 (in Japanese)

Lindsay EC, Metcalfe NB, Llewellyn MS (2020) The potential role of the gut

microbiota in shaping host energetics and metabolic rate. J Anim Ecol 89: 2415-

2426.

Modolon F, Barno AR, Villela HDM, Peixoto RS (2020) Ecological and

biotechnological importance of secondary metabolites produced by coral-associated

bacteria. J Appl Microbiol 129: 1441-1457.

Müller W (1934) Untersuchungen über die Symbiose von Tieren mit Pilzen und

Bakterien III Mitteilung: Über die Pilzsymbiose holzfressender Insektenlarven.

Archiv für Mikrobiologie 5: 84-147.

Nardon P, Grenier AM (1989) Endocytobiosis in Coleoptera: Biological, biochemical

and genetic aspects. Ins Endocymbiosis: Morphology, Physiology, Genetics

Evolution (ed. by S. W. & G. G.), pp. 175-216. CRC Press, Boca Raton, FL, USA.

Nie R, Vogler AP, Yang X, Lin M (2020) Higher-level phylogeny of longhorn beetles

(Coleoptera: Chrysomeloidea) inferred from mitochondrial genomes. Sys Entomol

46: 56-70.

Notredame C, Higgins DG, Heringa J (2000) T-Coffee: A novel method for fast and

accurate multiple sequence alignment. J Mol Biol 302: 205-217.

R Core Team (2021) R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/.

RStudio Team (2020) RStudio: Integrated Development for R. RStudio, PBC, Boston,

MA URL http://www.rstudio.com/.

Schomann H (1937) Die Symbiose der Bockkäfer. Zeitschrift für Morphologie und

Ökologie der Tiere 32: 542-612.

Suh SO, McHugh JV, Pollock DD, Blackwell M (2005) The beetle gut: a hyperdiverse

source of novel yeasts. Mycol Res 109: 261-265.

Svacha P, Lawrence, JF (2014) In: Handbook of Zoology, Arthropoda. Insecta.

Coleoptera. Volume 3: Coleoptera, Beetles. Morphology and Systematics

(Phytophaga). Ed by Kristensen NP, Beutel RG, pp 16-177. De Gruyter, Berlin,

Germany.

Tanahashi M, Kubota K, Matsushita N, Togashi K (2010) Discovery of mycangia and

the associated xylose-fermenting yeasts in stag beetles (Coleoptera: Lucanidae).

Naturwissenschaften 97: 311-317.

Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics

Analysis Version 11. Mol Biol Evol 38: 3022-3027.

Toki W, Takahashi Y, Togashi K (2013) Fungal garden making inside bamboos by a

non-social fungus-growing beetle. PLoS One 8: e79515.

Vega FE, Blackwell M (2005) Insect-fungal associations: ecology and evolution.

Oxford University Press, Oxford.

Wallace IM, O'Sullivan O, Higgins DG, Notredame C (2006) M-Coffee: combining

multiple sequence alignment methods with T-Coffee. Nucleic Acids Res 34: 1692-

1699.

White T, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics. In: PCR protocols, a guide to methods and
applications. Ed by M Innis, D Gelfand, J Sninsky, T White, pp. 315-322. Academic
Press, CA, USA.

FIGURE LEGENDS AND TABLE CAPTIONS

Fig. 1.

Detection of mycetomes in Japanese cerambycid larvae. **(A)** The phylogenetic

relationships of Japanese longicorn beetle families/subfamilies. Red circles indicate

groups including mycetome-possessing species. The number of species examined is

shown in parentheses. This phylogenetic tree follows that of Hadded et al. (2018),

although the branch length does not reflect phylogenetic distances. **(B)** Mycetomes

(arrow) on the anterior midgut serosa of the Lepturinae *Caraphia lepturoides*. The

anterior is toward the top. Scale bar, 500 μm. **(C)** Mycetomes of *Encyclops olivaceus*.

Scale bar, 500 μm. **(D)** Mycetomes of the Spondylidinae *Asemum striatum*. Scale bar,

500 μm. The mycetomes of this group form two rings surrounding the serosa, which are

shown by two arrows. **(E)** Giant mycetomes (My) of *Necydalis formosana*. Scale bar,

500 μm. MG, midgut; Th, thorax. The arrowhead indicates a thin tube that connects

mycetomes with the serosa. **(F)** Mycetome-lacking midgut of the Lepturinae

Paragaurotes doris. Scale bar, 500 μm. **(G)** Anterior midgut of the Prioninae *Aegosoma*

sinicum. Scale bar, 500 μm. The mycetome is absent.

Fig. 2.

Distribution of mycetomes in the subfamilies Lepturinae and Necydalinae. **(A)** This

phylogenetic tree was generated by the maximum likelihood method based on the

alignment of 28S rDNA sequences. Crossed circles indicate mycetome-lacking species.

Open circles indicate Lepturinae species that preferentially include the yeast

Scheffersomyces sp. Y1052_SICYLG3 in their mycetomes. Black circles indicate

Lepturinae species that preferentially include the yeast *Scheffersomyces ergatensis*.

Squares indicate Lepturinae species that preferentially include the yeast *Candida*

vrieseae with the information of their host trees. These three yeasts were frequently

isolated from various species. Note that this tree does not include all species examined

in this study for simplicity. The numbers beside branches indicate the percentage of

times that a node was supported in 1,000 bootstrap pseudoreplication. The scores that

are equal to or exceed 50% are shown. Species names are based on Danilevsky (2020).

The accession numbers of the DNA sequences used for the phylogenetic tree

construction are shown at the right side of the species names. **(B)** A simplified image of

(A), highlighting the distribution of the symbionts and the loss of mycetomes.

Fig. 3.

Fungal cells dwelling in the mycetomes. **(A-C)** Results for typical Lepturinae species.

The names of the examined longicorn beetles are shown at the bottom of each panel.

(A) Mycetome contents and yeast cells. Membranous structures are shown by arrows.

An arrowhead indicates an example of symbiont yeast cells. Yeast cells are shown in

higher-magnified images in **(B)** and **(B')**. Scale bar, 50 μm in **(A)** and 10 μm in **(B)**. **(C)**

Yeast colonies on a PDA plate. **(D-F)** The results for Lepturinae *Encyclops olivaceus*. In

(F), no colony was detected. **(G-I)** The results for Necydalinae species. The fungal cell

in **(H')** was isolated from a female oviduct. **(J-L)** The results for Spondylidinae species.

Arrows indicate the membranous structures.

Fig. 4.

Phylogenetic relationships of yeasts isolated from the mycetome-possessing longicorn

beetle larvae in this study. This phylogenetic tree was generated by the maximum

likelihood method based on the alignment of ITS rDNA sequences. *Hyphopichia rhagii*

(isolated from *Rhagium femorale* and *R. japonicum*), *Candida wanchemiae* (isolated

from *Anastrangalia scotodes*), and uncultured fungus clone OUT_F389 (isolated from

Megasemum quadricostulatum) were excluded from this tree because the yeast has

much shorter ITS rDNA sequence than those of the other yeasts. The species names of

yeasts are abbreviated. For example, "*Judolidia bangi* Se OP537542" indicates

"*Scheffersomyces ergatensis* isolated from *Judolidia bangi*, and the accession number of

the sequence used for the phylogenetic tree construction". The number noted at a given

branch indicates the percentage of times that a node was supported in 1,000 bootstrap

pseudoreplications. The scores that are equal to or exceed 50% are shown.

Fig. 5.

Phylogenetic relationships of yeasts isolated from larvae and adults of the Lepturinae

beetles. This phylogenetic tree was generated by the maximum likelihood method based

on the alignment of ITS rDNA sequences. The sequences of the yeasts isolated from the

larvae are the same as the ones used in Fig. 4. Ad, yeast isolated from adult; Lv, yeast

isolated from larva. The yeast species names are shown on the right side of the tree. For

example, "*Pachytodes cometes* Lv OP537597" indicates "*Scheffersomyces ergatensis*

isolated from a larva of *Pachytodes cometes*, and the accession number of the sequence

used for the phylogenetic tree construction". The clades that include the yeasts from

both adults and larvae of the same Lepturinae species are marked with black dots. The

number noted at a given branch indicates the percentage of times that a node was

supported in 1,000 bootstrap pseudoreplications. The scores that are equal to or exceed

50% are shown.

Fig. 6.

Fungal cells in the 1st instar larvae of Lepturinae/Necydalinae beetles. The observation

was carried out over the first few days after hatching. **(A-A')** The anterior midgut of

Ohbayashia nigromarginata and yeast cells isolated from the gut. The bracket indicates

mycetomes. Bar, 50 μm for A and 25 μm for A'. **(B-B')** The anterior midgut of

Necydalis formosana and a fungal cell isolated from the gut. Mycetome-like knobbed

structures are shown by the bracket. Bar, 50 μm for B and 25 μm for B'.

Fig. 7.

Yeast-free mycetomes in Lepturinae larvae. **(A)** *Leptura ochraceofasciata* larvae

cultured with an artificial diet. The images were obtained approximately one month

after hatching. **(B)** Yeast-free mycetomes (knobs shown by the bracket). **(C)** A

magnified image of the contents of yeast-free mycetomes. The inset shows the results of

culturing of the contents on a PDA plate. No colony appeared. **(D-I)** The results for

Stictoleptura dichroa. **(D-E)** Yeast-free mycetomes. **(F-G)** Yeast-poor mycetomes. Note

that the color of the mycetomes is somewhat whitish. Four examples of yeast cells are

shown by arrowheads. The yeast *Meyerozyma caribbica* was identified from the

mycetomes. **(H-I)** Yeast-rich mycetomes. The yeast *Scheffersomyces ergatensis* was

identified from these mycetomes.

Fig. 8.

The presence of symbionts in the mycetomes after long-time culture on the artificial diet

with anti-fungal chemicals. **(A)** *Etorofus vicarius* mycetome. Arrowheads are examples

of symbiont yeasts. Bar, 100 μm. **(B)** *Necydalis formosana* mycetome. Bar, 100 μm.

Fig. 9.

Aegosoma sinicum larvae have a symbiotic relationship with the two yeast groups. This

phylogenetic tree was generated by the maximum likelihood method based on the

alignment of ITS rDNA sequences. The yeasts from mycetome-possessing longicorn

beetles used in Fig. 4 are also shown in this phylogenetic tree to investigate the

phylogenetic relationships between mycetome-dependent and *Aegosoma* yeasts. The

yeasts isolated from larvae of *Aegosoma sinicum* are indicated by the sampling location

and identification numbers corresponding to those in Table S1. The number noted at a

given branch indicates the percentage of times that a node was supported in 1,000

bootstrap pseudoreplications. The scores that are equal to or exceed 50% are shown.

Table 1.

Summary of symbiont characterization among longicorn beetle groups collected in

Japan. The sum of yeast species isolated from Lepturinae having mycetome,

Necydalinae and Spondylidinae, excluding duplicates between groups. The sum of yeast

species isolated from Lepturinae without mycetome, Prioninae, Cerambycinae,

Lamiinae, and Disteniidae, excluding duplicates between groups.

Sasakura et al., Figure 1.

A

Cerambycinae (0/17)
Prioninae (0/4)
Necydalinae (3/3) ●
Lepturinae (22/31) ●
Spondylidinae (5/5) ●
Lamiinae (0/14)
Disteniidae (0/1)

Cerambycidae

● Groups that include larvae having mycetomes

anterior
posterior

B
Lepturinae
(*Caraphia lepturoides*)

C
Lepturinae
(*Encyclops olivacea*)

D
Spondylidinae
(*Asemum striatum*)

E
Th
My   MG   My
Necydalinae
(*Necydalis formosana*)

F
Lepturinae
(*Paragaurotes doris*)

G
Prioninae
(*Aegosoma sinicum*)

A

Ephies japonicus (OP562690) — 93
Pigostrangalia dulcis (OP562693) ■ Symplocos
Mimostrangalia kurosonensi (OP562691)
Strangalia koyaensis (OP562692) ■ Conifer
Judolidia bangi (OP562676) ●
Munamizoa maculata (OP5626977) ●
Leptura ochraceofasciata (OP562683) — 100
Leptura annularis (OP562684) ○
Pachytodes cometes (OP562678) ●
Stictoleptura dichroa (OP562679) ● — 98
Anastrangalia scotodes (OP562686) ■ Conifer
Etorofus vicarius (OP562681) ●
Eustrangalis anticereducta (OP562687) ○ — 54
Rhagium femorale (OP562689) — 99
Rhagium japonicum (OP562696)
Pidonia aegrota (OP562697) — 87
Pidonia oyamae (OP562698) ⊗
Strangalomorpha tenuis (OP562685) ○
Brachyta bifasciata (OP562703) ⊗
Pyrrhona laeticolor (OP562680)
Ohbayashia nigromarginata (OP562682) ■ Symplocos — 100
Parastrangalis lesnei (OP562694) ■ Non-Symplocos Hardwood
Encyclops olivaceus (OP562700)
Paragaurotes doris (OP562699) ⊗
Caraphia lepturoides (OP562688) ○
Japanocorus caeruleipennis (OP562695) ⊗
Necydalis gigantea (OP562701) — Necydalinae
Necydalis formosana (OP562702) — 88

B
A simplified image of (A)

Rhagium

Encyclops olivaceus

Necydalinae

⊗ Mycetome-lacking species/groups

○ Species or group that possessed *Scheffersomyces* sp. Y1052_SICYLG3

● Species or group that possessed *Scheffersomyces ergatensis*

■ Species or group that possessed *Candida vrieseae*

Mycetome Fungal cells Colonies on PDA plates

Typical Lepturinae

A — *L. ochraceofasciata*

B — *L. ochraceo-fasciata*

B' — *S. dichroa*

C — *P. laeticolor*

Encyclops olivaceus

D — *E. olivacea*

E — *E. olivaceus*

F — *E. olivaceus*

Necydalinae

G — *N. gigantea*

H — *N. formosana*

H' — *N. sachalinensis*

I — *N. formosana*

Spondylidinae

J — *M. quadricostulatum*

K — *M. quadricostula*

K' — *T. castaneum*

L — *T. castaneum*

1st instar larva of *O. nigromarginata*   1st instar larva of *N. formosana*

L. ochraceofasciata

S. dichroa

A
Etorofus vicarius

B
Necydalis formosana

Table 1. Summary of symbiont characterization among Longicorn beetle groups in Japan

Groups	Number of observed species	Number of observed larvae	N of yeast colony yielded larvae	N of larvae which had more than one yeast species	N of yeast species
Lepturinae having mycetome	22	125	116	7	10
Necydalinae	3	12	0	0	0
Spondylidinae	5	19	3	1	8
Lepturinae without mycetome	9	38	13	5	15
Prioninae	4	25	20	2	11
Cerambycinae	17	38	15	3	13
Lamiinae	14	29	11	3	12
Disteniidae	1	5	4	0	4
Mycetome present groups	30	156	119	8	18
Mycetome absent groups	46	135	63	13	46

Table S1. Symbiont fungi isolated from larvae of Longicorn beetles in Japan

Group	Species	Collected location*	host	mycetome	Analyzed region	Colonies on plate**	Blastn Best Hit	Blastn Score	Accession Number
Lepturinae with mycetome	*Anastrangalia scotodes*	Shizuoka	*Pinus thunbergii*	y	mycetome	y 10	[*Candida*] *vrieseae* CBS 10829	592/606	OP537521
Lepturinae with mycetome	*Anastrangalia scotodes*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	[*Candida*] *vrieseae* CBS 10829	578/593	OP537522
Lepturinae with mycetome	*Anastrangalia scotodes*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	not analyzed		
Lepturinae with mycetome	*Anastrangalia scotodes*	Nara	*Pinus densiflora*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829 *Candida wanchemiae* CBS 12090	612/630 371/420	OP537523 OP537524
Lepturinae with mycetome	*Anastrangalia scotodes*	Gifu	*Betula platyphylla*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537525
Lepturinae with mycetome	*Anastrangalia scotodes*	Gifu	*Abies veitchii*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829 *Schefferomyces ergatensis* strain ATCC MYA-4655	614/629 614/615	OP537526 OP537527
Lepturinae with mycetome	*Anastrangalia scotodes*	Gifu	*Abies veitchii*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537528
Lepturinae with mycetome	*Stictoleptura dichroa*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537529
Lepturinae with mycetome	*Stictoleptura dichroa*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537530
Lepturinae with mycetome	*Stictoleptura dichroa*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537531
Lepturinae with mycetome	*Stictoleptura dichroa*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537532
Lepturinae with mycetome	*Stictoleptura dichroa*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	614/616	OP537533
Lepturinae with mycetome	*Stictoleptura dichroa*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	614/615	OP537534
Lepturinae with mycetome	*Caraphia lepturoides*	Aichi	*Clerodendrum trichotomum*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537535
Lepturinae with mycetome	*Caraphia lepturoides*	Aichi	*Clerodendrum trichotomum*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537536
Lepturinae with mycetome	*Ephies japonicus*	Yamaguchi	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	584/597	OP537537
Lepturinae with mycetome	*Ephies japonicus*	Yamaguchi	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	584/597	OP537538
Lepturinae with mycetome	*Eustrangalis anticereducta*	Gifu	*Robinia pseudoacacia*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	617/619	OP537539
Lepturinae with mycetome	*Eustrangalis anticereducta*	Gifu	*Robinia pseudoacacia*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	587/587	OP537540
Lepturinae with mycetome	*Eustrangalis anticereducta*	Gifu	*Robinia pseudoacacia*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	617/619	OP537541
Lepturinae with mycetome	*Judolidia bangi*	Yamanashi	*Larix kaempferi*	y	mycetome/gut	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	614/615	OP537542
Lepturinae with mycetome	*Leptura annulalis*	Ibaraki	*Salix* sp.	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	593/593	OP537543
Lepturinae with mycetome	*Leptura annulalis*	Ibaraki	*Salix* sp.	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537544
Lepturinae with mycetome	*Leptura annulalis*	Kagoshima	unknown	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537545
Lepturinae with mycetome	*Leptura annulalis*	Yamanashi	*Sorbus commixta*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537546
Lepturinae with mycetome	*Leptura annulalis*	Gifu	*Betula platyphylla*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537547
Lepturinae with mycetome	*Leptura ochraceofasciata*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537548
Lepturinae with mycetome	*Leptura ochraceofasciata*	Nara	*Carpinus tschonoskii*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537549
Lepturinae with mycetome	*Leptura ochraceofasciata*	Nara	*Carpinus tschonoskii*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537550
Lepturinae with mycetome	*Leptura ochraceofasciata*	Nara	*Carpinus tschonoskii*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	586/587	OP537551
Lepturinae with mycetome	*Leptura ochraceofasciata*	Nara	*Carpinus tschonoskii*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	595/596	OP537552
Lepturinae with mycetome	*Leptura ochraceofasciata*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	586/587	OP537553
Lepturinae with mycetome	*Leptura ochraceofasciata*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537554
Lepturinae with mycetome	*Leptura ochraceofasciata*	Aichi	*Cryptomeria japonica*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537555
Lepturinae with mycetome	*Leptura ochraceofasciata*	Aichi	*Cryptomeria japonica*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537556
Lepturinae with mycetome	*Leptura ochraceofasciata*	Kagoshima Yakushima Island	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	597/612	OP537557
Lepturinae with mycetome	*Leptura ochraceofasciata*	Aichi	*Lindera triloba*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	592/593	OP537558
Lepturinae with mycetome	*Leptura ochraceofasciata*	Mie	*Cryptomeria japonica*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	616/621	OP537559
Lepturinae with mycetome	*Leptura ochraceofasciata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537560
Lepturinae with mycetome	*Leptura ochraceofasciata*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537561
Lepturinae with mycetome	*Leptura ochraceofasciata*	Kagoshima	unknown	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537562
Lepturinae with mycetome	*Leptura ochraceofasciata*	Gifu	*Hamamelis japonica*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537563
Lepturinae with mycetome	*Leptura ochraceofasciata*	Gifu	*Fagus crenata*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	618/619	OP537564
Lepturinae with mycetome	*Leptura subtilis*	Yamanashi	*Betula platyphylla*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537565

Lepturinae with mycetome	*Leptura subtilis*	Yamanashi	*Betula platyphylla*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537566
Lepturinae with mycetome	*Pigostrangalia dulcis*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	607/623	OP537567
Lepturinae with mycetome	*Pigostrangalia dulcis*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537568
Lepturinae with mycetome	*Mimostrangalia kurosonensis*	Kagoshima Yakushima Island	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	627/641	OP537569
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	Failed to observe	gut	y	*Scheffersomyces* sp. strain Y13	588/589	OP537570
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	Failed to observe	gut	y	*Scheffersomyces* sp. strain Y6	583/583	OP537571
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537572
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/616	OP537573
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537574
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	615/615	OP537575
Lepturinae with mycetome	*Munamizoa maculata*	Yamanashi	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces ergatensis* strain ATCC 22589	604/611	OP537576
Lepturinae with mycetome	*Etorofus vicarius*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	595/596	OP537577
Lepturinae with mycetome	*Etorofus vicarius*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. strain Y13	589/593	OP537578
Lepturinae with mycetome	*Etorofus vicarius*	Yamanashi	*Abies* sp.	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	613/616	OP537579
Lepturinae with mycetome	*Etorofus vicarius*	Yamanashi	*Abies* sp.	y	mycetome	y	*Scheffersomyces* sp. strain Y13	586/589	OP537580
Lepturinae with mycetome	*Etorofus vicarius*	Yamanashi	*Abies* sp.	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	613/616	OP537581
Lepturinae with mycetome	*Etorofus vicarius*	Yamanashi	*Larix kaempferi*	Failed to observe	gut	y 1	*Schefferomyces ergatensis* strain ATCC MYA-4655	613/616	OP537582
Lepturinae with mycetome	*Etorofus vicarius*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Schefferomyces ergatensis* strain ATCC MYA-4655	612/616	OP537583
Lepturinae with mycetome	*Etorofus vicarius*	Shizuoka	*Chamaecyparis obtusa*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	619/619	OP537584
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	not observed	Frass	no yeast colony			
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537585
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	581/595	OP537586
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537587
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	505/519	OP537588
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537589
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	588/602	OP537590
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537591
							[*Candida*] *vrieseae* CBS 10829	615/629	OP537592
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Wakamaya	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	614/629	OP537593
							Scheffersomyces coipomoensis strain Y3	615/615	OP537594
Lepturinae with mycetome	*Ohbayashia nigromarginata*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	612/634	OP537595
							Scheffersomyces coipomoensis strain Y3	615/615	OP537596
Lepturinae with mycetome	*Pachytodes cometes*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	615/615	OP537597
							Scheffersomyces sp. strain Y13	592/592	OP537598
Lepturinae with mycetome	*Pachytodes cometes*	Aichi	*Cryptomeria japonica*	y	mycetome	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	615/615	OP537599
Lepturinae with mycetome	*Pachytodes cometes*	Yamanashi	*Larix kaempferi*	y	mycetome	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	615/615	OP537600
Lepturinae with mycetome	*Parastrangalia lesnei*	Gifu	*Hamamelis japonica*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	582/596	OP537601
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	not observed	frass	no yeast colony			
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	not observed	frass	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537602
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	not observed	frass	y	[*Candida*] *vrieseae* CBS 10829	592/606	OP537603
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	not observed	frass	no yeast colony			
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537604
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537605
Lepturinae with mycetome	*Pyrrhona laeticolor*	Yakushima Island	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	627/641	OP537606
Lepturinae with mycetome	*Pyrrhona laeticolor*	Yakushima Island	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537607
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537608
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537609
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537610

Group	Species	Location	Host plant	Mycetome	Tissue	Yeast colony	Yeast identification	bp	Accession
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537611
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537612
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	613/631	OP537613
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537614
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537615
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537616
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537617
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537618
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537619
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537620
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537621
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	605/619	OP537622
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537623
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	613/631	OP537624
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537625
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537626
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome, gut	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537627
							Scheffersomyces ergatensis strain ATCC MYA-4655	609/615	OP537628
							Scheffersomyces paraergatensis strain NYNU 16782	586/587	OP537629
							[*Candida*] *vrieseae* CBS 10829	615/629	OP537630
Lepturinae with mycetome	*Pyrrhona laeticolor*	Wakamaya	*Symplocos myrtacea*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	583/597	OP537631
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	615/629	OP537632
Lepturinae with mycetome	*Pyrrhona laeticolor*	Shizuoka	*Symplocos coreana*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	613/632	OP537633
Lepturinae with mycetome	*Rhagium femorale*	Nagano	*Abies veitchii*	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	369/369	OP537634
Lepturinae with mycetome	*Rhagium femorale*	Nagano	*Abies veitchii*	y	mycetome	y	*Candida tenuis*	566/566	OP537635
Lepturinae with mycetome	*Rhagium femorale*	Tochigi	unknown	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	369/369	OP537636
Lepturinae with mycetome	*Rhagium femorale*	Yamanashi	*Abies veitchii*	y	mycetome	y	*Candida tenuis*	581/581	OP537637
Lepturinae with mycetome	*Rhagium femorale*	Yamanashi	*Abies veitchii*	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	405/405	OP537638
Lepturinae with mycetome	*Rhagium femorale*	Yamanashi	*Abies veitchii*	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	377/379	OP537639
Lepturinae with mycetome	*Rhagium femorale*	Yamanashi	*Abies veitchii*	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	405/405	OP537640
Lepturinae with mycetome	*Rhagium japonicum*	Tochigi	unknown	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	382/382	OP537641
Lepturinae with mycetome	*Rhagium japonicum*	Tochigi	unknown	y	mycetome	y	*Hyphopichia rhagii* culture CBS:618	404/405	OP537642
Lepturinae with mycetome	*Strangalia koyaensis*	Aichi	*Cryptomeria japonica*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	613/629	OP537643
Lepturinae with mycetome	*Strangalia koyaensis*	Aichi	*Cryptomeria japonica*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	612/629	OP537644
Lepturinae with mycetome	*Strangalia koyaensis*	Aichi	*Cryptomeria japonica*	y	mycetome	y	[*Candida*] *vrieseae* CBS 10829	612/630	OP537645
Lepturinae with mycetome	*Strangalomorpha tenuis*	Yamanashi	*Sorbus commixta*	y	mycetome	y	*Scheffersomyces* sp. Y1052_SICYLG3	586/586	OP537646
Lepturinae with mycetome	*Encyclops olivaceus*	Aichi	*Quercus crispula*	y	mycetome	no yeast colony	*Chrysolina oricalcia* genome assembly, chromosome: 12	186/190	OP537662
Lepturinae with mycetome	*Encyclops olivaceus*	Aichi	*Quercus crispula*	y	mycetome	no yeast colony	*Acalymma blandulum*	149/153	OP537663
Lepturinae with mycetome	*Encyclops olivaceus*	Gifu	*Quercus crispula*	y	mycetome	no yeast colony	not analyzed		
Lepturinae with mycetome	*Encyclops olivaceus*	Gifu	*Quercus crispula*	y	mycetome	no yeast colony	*Chrysolina oricalcia* genome assembly, chromosome: 12	185/190	OP537664
Lepturinae w/o mycetome	*Pargaurotes doris*	Nagano		n	gut	y	[*Candida*] *vrieseae* CBS 10829	607/636	OP537647
Lepturinae w/o mycetome	*Pargaurotes doris*	Nagano		n	gut	y	*Yamadazyma mexicana* culture CBS:2089	606/618	OP537648
Lepturinae w/o mycetome	*Pargaurotes doris*	Nagano		n	gut	y	[*Candida*] *montana* isolate yHRVM49	396/408	OP537649
							Wickerhamomyces canadensis culture CBS:5676	522/586	OP537650
Lepturinae w/o mycetome	*Pargaurotes doris*	Yamanashi		n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pargaurotes doris*	Yamanashi		n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pargaurotes doris*	Tochigi	*Fagus crenata*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pargaurotes doris*	Gifu	*Abies veitchii*	n	gut	y	[*Candida*] *rhagii* culture CBS:618	373/373	OP537651
Lepturinae w/o mycetome	*Pargaurotes doris*	Gifu	*Abies veitchii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pargaurotes doris*	Gifu	*Abies veitchii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia aegrota*	Nara	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia aegrota*	Nara	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia aegrota*	Nara	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia aegrota*	Mie	*Pieris japonica*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia aegrota*	Shizuoka	*Carpinus* sp.	n	gut	y 3	*Candida boleticola* culture CBS:6420	612/613	OP537652

Subfamily	Species	Location	Host plant		Tissue	Yeast	Closest match	Identity	Accession
Lepturinae w/o mycetome	*Pidonia amentata?*	Nara	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia amentata?*	Nara	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia discoidalis*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Yamanashi	unknown	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Yamanashi	unknown	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Yamanashi	unknown	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Mie	*Styrax japonica*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Mie	*Styrax japonica*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Mie	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Mie	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Mie	*Carpinus tschonoskii*	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia oyamae/masakii*	Shizuoka	*Carpinus* sp.	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia sp.*	Yamanashi	unknown	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Pidonia sylvicola/himehana/maculitho*	Aichi	*Kalopanax septemlobus*	n	gut	y	[*Candida*] *montana* isolate yHRVM49	396/408	OP537653
Lepturinae w/o mycetome	*Pidonia sylvicola/himehana/maculitho*	Aichi	*Kalopanax septemlobus*	n	gut	y	*Wickerhamomyces canadensis* culture CBS:5676	522/586	OP537654
							[*Candida*] *montana* isolate yHRVM49	396/408	OP589179
Lepturinae w/o mycetome	*Pidonia sylvicola/himehana/maculitho*	Aichi	*Kalopanax septemlobus*	n	gut	y	[*Candida*] *montana* isolate yHRVM49	396/408	OP537655
Lepturinae w/o mycetome	*Pidonia sylvicola/himehana/maculitho*	Yamanashi	unknown	n	gut	no yeast colony			
Lepturinae w/o mycetome	*Japanocorus caeruleipennis*	Culture		n	gut	y	[*Candida*] *gorgasii* ATCC MYA-4334	584/621	OP537656
							[*Candida*] *michaelii* culture CBS:9502	613/659	OP589180
							Sugiyamaella smithiae strain NS1	549/549	OP589181
Lepturinae w/o mycetome	*Japanocorus caeruleipennis*	Culture		n	gut	y	[*Candida*] *insectorum*	625/625	OP537657
							Nakazawaea sp. isolate UFMG-CM-Y6909	609/611	OP537658
Lepturinae w/o mycetome	*Japanocorus caeruleipennis*	Culture		n	gut	y	[*Candida*] *vrieseae* CBS 10829	587/613	OP537659
Lepturinae w/o mycetome	*Japanocorus caeruleipennis*	Culture		n	gut	no yeast colony			
Lepturinae w/o mycetome	*Japanocorus caeruleipennis*	Culture		n	gut	y	*Candida* sp. Af110-2-1	591/591	OP537660
							Pichia hampshirensis strain ATCC 62896	551/566	OP589182
							Pichia veronae strain CECT 11800	594/596	OP589183
Lepturinae w/o mycetome	*Brachyta bifasciata*	Culture		n	gut	no yeast colony			
Lepturinae w/o mycetome	*Brachyta bifasciata*	Culture		n	gut	y l	*Zygoascus meyerae* culture CBS:6360	571/572	OP537661
Necydalinae	*Necydalis formosana*	Shizuoka	*Symplocos coreana*	not observed	frass	no yeast colony			
Necydalinae	*Necydalis formosana*	Shizuoka	*Symplocos coreana*	y	mycetome	no yeast colony	Uncultured fungus clone Oc_O_AugC04	467/552	OP537665
Necydalinae	*Necydalis formosana*	Shizuoka	*Symplocos coreana*	y	mycetome	no yeast colony			
Necydalinae	*Necydalis formosana*	Shizuoka	*Symplocos coreana*	y	mycetome	no yeast colony			
Necydalinae	*Necydalis formosana*	Shizuoka	*Symplocos coreana*	y	mycetome	not done	*Phacidiales* sp. strain 19GCAS019	359/410	OP537666
Necydalinae	*Necydalis gigantea*	Nara	*Neolitsea aciculata*	not observed	frass	no yeast colony			
Necydalinae	*Necydalis gigantea*	Nara	*Neolitsea aciculata*	y		no yeast colony	*Aotearoamyces nothofagi* voucher PDD:55517	493/555	OP537667
Necydalinae	*Necydalis harmandi*	Yamanashi	*Sorbus commixta*	y		no yeast colony	*Variabilispora flava* CBS 144845	374/421	OP537668
Necydalinae	*Necydalis harmandi*	Yamanashi	*Sorbus commixta*	y	mycetome	no yeast colony	Uncultured fungus clone 3306E21	477/544	OP537669
Necydalinae	*Necydalis harmandi*	Yamanashi	*Sorbus commixta*	y	mycetome	no yeast colony	Uncultured fungus clone 3306E21	476/545	OP537670
Necydalinae	*Necydalis harmandi*	Yamanashi	*Sorbus commixta*	y	mycetome gut	no yeast colony	Uncultured fungus clone 3306E21	477/544	OP537671
Necydalinae	*Necydalis harmandi*	Yamanashi	*Sorbus commixta*	y	mycetome gut	no yeast colony	*Variabilispora flava* CBS 144845	445/509	OP537672
Spondylidinae	*Arhopalus coreanus*	Aichi	*Pinus densiflora*	y	mycetome	no yeast colony			
Spondylidinae	*Arhopalus coreanus*	Aichi	*Pinus densiflora*	y	mycetome	no yeast colony			
Spondylidinae	*Asemum striatum*	Nagano	*Larix kaempferi*	y	mycetome	y	*Candida oregonensis* CBS 5036	386/386	OP537673
Spondylidinae	*Asemum striatum*	Nagano	*Larix kaempferi*	y	mycetome	y l	uncultured fungus/*Schwanniomyces capriottii* isolate NG_p11	574/611	OP537674
Spondylidinae	*Asemum striatum*	Nagano	*Larix kaempferi*	y	mycetome	no yeast colony	*Debaryomyces castellii* culture CBS:2923	562/599	OP537675
Spondylidinae	*Cephalallus unicolor*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	*Debaryomyces* sp. UFMG-CM-Y343	573/628	OP537676
Spondylidinae	*Cephalallus unicolor*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	*Debaryomyces* sp. UFMG-CM-Y343	572/627	OP537677
Spondylidinae	*Cephalallus unicolor*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	*Schwanniomyces vanrijiae* var. yarrowii culture CBS:6246	584/638	OP537678
Spondylidinae	*Cephalallus unicolor*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	*Schwanniomyces vanrijiae* var. yarrowii culture CBS:6246	584/638	OP537679
Spondylidinae	*Cephalallus unicolor*	Shizuoka	*Pinus thunbergii*	y	mycetome	no yeast colony	*Schwanniomyces vanrijiae* var. yarrowii culture CBS:6246	584/638	OP537680
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Abies veitchii*	not observed	gut	no yeast colony			

Subfamily	Species	Location	Host		Tissue	Yeast colony	Identification	Match	Accession
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Abies veitchii*	y	mycetome	no yeast colony			
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Abies veitchii*	y	mycetome	no yeast colony	*Debaryomyces castellii* culture CBS:2923	589/627	OP537681
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Abies veitchii*	y	mycetome	y 6	*Debaryomyces castellii* culture CBS:2923 (mycetome)	589/627	OP537682
							Scheffersomyces sp. strain Y13 (colony)	577/578	OP537683
							Candida ergatensis strain ATCC MYA-4655 (colony)	613/616	OP537684
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Abies veitchii*	y	mycetome	no yeast colony	*Debaryomyces castellii* culture CBS:2923	589/627	OP537685
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Larix kaempferi*	y	mycetome	no yeast colony	*Debaryomyces castellii* culture CBS:2923	589/627	OP537686
Spondylidinae	*Megasemum quadricostulatum*	Yamanashi	*Larix kaempferi*	y	mycetome	no yeast colony	Uncultured fungus clone OTU_F389	43/43	OP537687
							Debaryomyces castellii culture CBS:2923	589/627	OP537688
Spondylidinae	*Tetropium castaneum*	Nagano	*Picea jezoensis*	y	mycetome	no yeast colony	*Debaryomyces* sp. UFMG-CM-Y343	618/660	OP537689
Spondylidinae	*Tetropium castaneum*	Nagano	*Picea jezoensis*	y	mycetome	no yeast colony			
Prioninae	*Psephactus remiger*	Shizuoka	*Fagus crenata*	n	gut	y	*Scheffersomyces* sp. strain Y13	601/601	OP537694
Prioninae	*Psephactus remiger*	Shizuoka	*Fagus crenata*	n	gut	y	*Candida boleticola*	590/590	OP537695
Prioninae	*Psephactus remiger*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Prioninae	*Psephactus remiger*	Shizuoka	*Fagus crenata*	n	gut	y 2	*Sugiyamaella smithiae* strain NS1	549/550	OP537696
Prioninae	*Psephactus remiger*	Yamanashi	*Abies* sp.	n	gut	y	*Scheffersomyces coipomoensis* strain Y3	614/615	OP537697
							Scheffersomyces sp. Y1052_SICYLG	619/619	OP537698
Prioninae	*Psephactus remiger*	Mie	*Mallotus japonicus*	n	gut	y	[*Candida*] *boleticola* strain UCDFST:80-62#	590/590	OP537699
Prioninae	*Prionus insularis*	Cultured		n	gut	no yeast colony			
Prioninae	*Prionus insularis*	Cultured		n	gut	no yeast colony			
Prioninae	*Aegosoma sinicum*	Shizuoka1	*Mallotus japonicus*	n	gut	y	*Scheffersomyces anoplophorae*	591/591	OP537700
Prioninae	*Aegosoma sinicum*	Mie1	*Mallotus japonicus*	n	gut	y	*Scheffersomyces anoplophorae*	579/579	OP537701
							Rhodotorula mucilaginosa strain 2018A0762M12	576/577	OP537702
Prioninae	*Aegosoma sinicum*	Mie2	*Mallotus japonicus*	n	gut	y	*Scheffersomyces anoplophorae*	579/579	OP537703
Prioninae	*Aegosoma sinicum*	Mie3	*Mallotus japonicus*	n	gut	y	*Scheffersomyces* sp. YW28.1	613/613	OP537704
Prioninae	*Aegosoma sinicum*	Shiga	*Fagus crenata*	n	gut	y 7	*Scheffersomyces anoplophorae*	591/591	OP589184
Prioninae	*Aegosoma sinicum*	Kyoto1	*Populus* sp.	not observed	gut	no yeast colony			
Prioninae	*Aegosoma sinicum*	Kyoto2	*Populus* sp.	not observed	gut	y	*Scheffersomyces anoplophorae*	587/591	OP537705
Prioninae	*Aegosoma sinicum*	Kyoto3	*Populus* sp.	not observed	gut	y	*Scheffersomyces anoplophorae*	577/581	OP537706
Prioninae	*Aegosoma sinicum*	Ibaraki1	*Salix* sp.	n	gut	y	Uncultured Scheffersomyces clone 15y-3	582/584	OP537707
							Scheffersomyces sp. YW01.2	577/577	OP537708
Prioninae	*Aegosoma sinicum*	Ibaraki2	*Salix* sp.	n	gut	y 4	*Scheffersomyces* sp. YW01.2	612/612	OP537709
Prioninae	*Aegosoma sinicum*	Ibaraki3	*Salix* sp.	n	gut	y	*Scheffersomyces anoplophorae*	591/591	OP537710
Prioninae	*Aegosoma sinicum*	Ibaraki4	*Salix* sp.	n	gut	y 3	*Scheffersomyces anoplophorae*	582/582	OP537711
Prioninae	*Aegosoma sinicum*	Ibaraki5	*Salix* sp.	n	gut	y	*Scheffersomyces anoplophorae*	590/591	OP537712
Prioninae	*Aegosoma sinicum*	Kagoshima Yakushima Island	unknown	n	frass gut	y	*Scheffersomyces* sp. YW01.2	612/612	OP537713
Prioninae	*Aegosoma sinicum*	Shizuoka2	*Fagus crenata*	n	gut	y	*Scheffersomyces* sp. YW27.1	612/612	OP537714
Prioninae	*Eurypoda batesi*	Kagoshima Yakushima Island	unknown	n	gut	y	*Scheffersomyces coipomoensis* strain Y3	615/615	OP537715
Prioninae	*Eurypoda batesi*	Kagoshima Yakushima Island	unknown	n	gut	no yeast colony			
Cerambycinae	*Xystrocera globosa*	Aichi	*Albizia julibrissin*	n	gut	y	*Ogataea naganishii* culture CBS:7259	711/712	OP537716
							Kluyveromyces sp. UFMG-CM-Y3682	657/676	OP537717
Cerambycinae	*Xystrocera globosa*	Aichi	*Albizia julibrissin*	n	gut	y	*Ogataea naganishii* culture CBS:7259	711/711	OP537718
Cerambycinae	*Xylotrechus emaciatus*	Kyoto	*Celtis jessoensis*	n	gut	no yeast colony			
Cerambycinae	*Xylotrechus emaciatus*	Kyoto	*Celtis jessoensis*	n	gut	no yeast colony			
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Acer* sp.	not observed	frass	y	*Scheffersomyces* sp. strain Y13	603/603	OP537719
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Carpinus* sp.	n	frass	y	*Scheffersomyces* sp. strain Y13	590/590	OP537720
							Candida khao-*thaluensis* strain CBS 9930	608/622	OP537721
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Carpinus* sp.	n	frass	no yeast colony			
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Cerambycinae	*Xylotrechus cuneipennis*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Cerambycinae	*Xylotrechus clarinus*	Nagano	*Betula platyphylla*	failed to observe	gut	no yeast colony			
Cerambycinae	*Xylotrechus clarinus*	Nagano	*Betula platyphylla*	n	gut	no yeast colony			
Cerambycinae	*Thranius variegatus*	Aichi	*Mallotus japonicus*	n	gut	y	*C. schatavii*	586/593	OP537722

Subfamily	Species	Location	Host	Obs	Source	Yeast	Identity	Match	Accession
Cerambycinae	*Thranius variegatus*	Aichi	*Mallotus japonicus*	n	gut	y	*C. schatavii*	586/593	OP537723
Cerambycinae	*Purpuricenus temminckii*	Aichi	*Phyllostachys edulis*	n	gut	y	*Candida blattae*	350/351	OP537724
Cerambycinae	*Purpuricenus temminckii*	Aichi	*Phyllostachys edulis*	n	gut	y	*Hyphopichia* sp. strain NYNU 20899	392/395	OP537725
Cerambycinae	*Phymatodes testaceus*	Nagano	*Quercus serrata*	n	gut	no yeast colony			
Cerambycinae	*Paraclytus excultus*	Yamanashi	Hardwood	n	gut	no yeast colony			
Cerambycinae	*Paraclytus excultus*	Yamanashi	Hardwood	n	gut	no yeast colony			
Cerambycinae	*Paraclytus excultus*	Yamanashi	Hardwood	n	gut	no yeast colony			
Cerambycinae	*Paraclytus excultus*	Yamanashi	Hardwood	failed to observe	gut	no yeast colony			
Cerambycinae	*Trichoferus campestris*	Yamanashi	*Prunus persica*	n	gut	y	*Hyphopichia homilentoma* culture CBS:6099	376/379	OP537726
Cerambycinae	*Trichoferus campestris*	Yamanashi	*Prunus persica*	n	gut	no yeast colony			
Cerambycinae	*Grammographus notabilis*	Yamanashi	*Pterocarya rhoifolia*	n	gut	y 4	*Hyphopichia homilentoma* culture CBS:6099	385/386	OP537727
							Yamadazyma akitaensis CBS 6701	604/611	OP537728
Cerambycinae	*Epania dilaticornis*	Kagoshima	*Diospyros lotus*	failed to observe	frass	no yeast colony			
Cerambycinae	*Epania dilaticornis*	Kagoshima	*Diospyros lotus*	failed to observe	frass	no yeast colony			
Cerambycinae	*Epania dilaticornis*	Kagoshima	*Diospyros lotus*	failed to observe	frass	no yeast colony			
Cerambycinae	*Cyrtoclytus caproides*	Ibaraki	*Salix* sp.	n	gut	no yeast colony	not analyzed		
Cerambycinae	*Clytus auripilis*	Kyoto	*Celtis jessoensis*	n	gut	y 3	Uncultured *Candida*	608/615	OP537729
Cerambycinae	*Ceresium holophaeum*	Aichi	*Euonymus japonicus*	n	gut	y	*Hyphopichia pseudoburtonii*	400/400	OP537730
Cerambycinae	*Ceresium holophaeum*	Aichi	*Euonymus japonicus*	n	gut	y	*Meyerozyma carpophila*	568/568	OP537731
Cerambycinae	*Callidiellum rufipenne*	Nagano	*Chamaecyparis obtusa*	failed to observe	gut	no yeast colony			
Cerambycinae	*Callidiellum rufipenne*	Nagano	*Chamaecyparis obtusa*	n	gut	no yeast colony			
Cerambycinae	*Callidiellum rufipenne*	Nagano	*Chamaecyparis obtusa*	n	gut	no yeast colony			
Cerambycinae	*Aromia bungii*	Oosaka	*Cerasus* sp.	not observed	frass	y	[*Candida*] *succiphila* culture CBS:7297	623/623	OP537732
Cerambycinae	*Aromia bungii*	Oosaka	*Cerasus* sp.	not observed	frass	no yeast colony			
Cerambycinae	*Anaglyptus niponensis*	Yamanashi	*Pterocarya rhoifolia*	n	gut	y 2	*Hyphopichia homilentoma* culture CBS:6099	409/412	OP537733
Lamiinae	*Acalolepta sejuncta*	Shizuoka	*Acer* sp.	not observed	frass	no yeast colony			
Lamiinae	*Acalolepta sejuncta*	Shizuoka	*Carpinus* sp.	n	gut	no yeast colony			
Lamiinae	*Eumecocera argyrosticta*	Shizuoka	*Acer* sp.	not observed	frass	no yeast colony			
Lamiinae	*Eumecocera gleneoides*	Shizuoka	*Cornus controversa*	not observed	frass	no yeast colony			
Lamiinae	*Eutetrapha chrysochloris*	Shizuoka	*Tilia japonica*	not observed	frass	y	[*Candida*] *gorgasii* ATCC MYA-4334	607/644	OP537734
Lamiinae	*Eutetrapha chrysochloris*	Shizuoka	*Acer* sp.	not observed	frass	no yeast colony			
Lamiinae	*Eutetrapha chrysochloris*	Nagano	*Quercus serrata*	n	gut	y	*Candida* sp. (in: *Saccharomycetales*) strain NYUN 211183	435/436	OP537735
Lamiinae	*Eutetrapha chrysochloris*	Nagano	*Quercus serrata*	n	gut	y	*Yamadazyma mexicana* culture CBS:2089	606/618	OP537736
							Pichia hampshirensis strain ATCC 62896	572/587	OP537737
							Candida sp. (in: *Saccharomycetales*) strain NYUN 211183	435/436	OP537738
Lamiinae	*Leiopus stillatus*	Shizuoka	*Carpinus sp.*	not observed	frass	no yeast colony			
Lamiinae	*Leiopus stillatus*	Shizuoka	*Acer* sp.	not observed	frass	y	*Candida* sp. (in: *Saccharomycetales*) isolate P10-3-5	568/570	OP537739
Lamiinae	*Mecynippus pubicornis*	Nagano	*Salix* sp.	n	gut	no yeast colony			
Lamiinae	*Mecynippus pubicornis*	Nagano	*Salix* sp.	n	gut	no yeast colony			
Lamiinae	*Monochamus alternatus*	Shizuoka	*Pinus thunbergii*	n	gut	y	*Hyphopichia heimii* strain SSA1530	364/366	OP537740
Lamiinae	*Monochamus alternatus*	Shizuoka	*Pinus thunbergii*	n	gut	y	*Hyphopichia heimii* strain SSA1530	364/366	OP537741
Lamiinae	*Nanohammus rufescens*	Aichi	*Quercus crispula*	n	gut	no yeast colony			
Lamiinae	*Nanohammus rufescens*	Aichi	*Quercus crispula*	n	gut	no yeast colony			
Lamiinae	*Niphona furcata*	Aichi	*Bambuseae*	n	gut	y	*Hyphopichia burtonii* strain XWB32-4	404/404	OP537742
							Hyphopichia sp. isolate SFC101142	385/390	OP537743
Lamiinae	*Niphona furcata*	Aichi	*Bambuseae*	n	gut	y	*Hyphopichia pseudoburtonii* isolate makgeolli	377/377	OP537744
Lamiinae	*Niphona furcata*	Aichi	*Bambuseae*	n	gut	y	*Hyphopichia burtonii* culture CBS:2452	425/428	OP537745
Lamiinae	*Oplosia suvorovi*	Aichi	*Tilia japonica*	n	gut	no yeast colony			
Lamiinae	*Palimna liturata*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Lamiinae	*Palimna liturata*	Shizuoka	*Fagus crenata*	n	gut	no yeast colony			
Lamiinae	*Palimna liturata*	Kagoshima	unknonw	n	gut	no yeast colony			
Lamiinae	*Parechthistatus gibber*	Shizuoka	*Fagus crenata*	n	frass	no yeast colony			
Lamiinae	*Parechthistatus gibber*	Shizuoka	*Fagus crenata*	n	frass	y	*Candida* sp. W370	539/541	OP537746
Lamiinae	*Parechthistatus gibber*	Shizuoka	*Fagus crenata*	n	gut	y	*Scheffersomyces* sp. Y1052_SICYLG3	596/596	OP537747
							Sugiyamaella boreocaroliniensis CBS 10344	555/555	OP537748

Lamiinae	*Pseudocalamobius montanus*	Gifu	*Ribes japonicum*	n	gut	no yeast colony			
Lamiinae	*Pseudocalamobius montanus*	Gifu	*Ribes japonicum*	n	gut	no yeast colony			
Lamiinae	*Saperdini sp.*	Aichi	*Tilia japonica*	n	gut	no yeast colony			
Disteniidae	*Distenia gracilis*	Mie	*Carpinus tschonoskii*	n	gut	y	[*Candida*] *boleticola* strain UCDFST:80-62#	613/613	OP537690
Disteniidae	*Distenia gracilis*	Ibaraki	*Salix* sp.	n	gut	y 8	[*Candida*] *intermedia* strain DL6Y	374/374	OP537691
Disteniidae	*Distenia gracilis*	Shizuoka	*Fagus crenata*	n	gut	no colony			
Disteniidae	*Distenia gracilis*	Shizuoka	*Carpinus* sp.	n	gut	y 2	*Candida rhagii* culture CBS:618	376/415	OP537692
Disteniidae	*Distenia gracilis*	Shizuoka	*Carpinus* sp.	n	gut	y 2	*Candida* sp. (in: *Saccharomycetales*) strain NYUN 211183	435/437	OP537693
Stag beetle	*Aesalus asiaticus*	Shizuoka	*Symplocos coreana*	n	gut	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	609/615	OP537749
Stag beetle	*Aesalus asiaticus*	Shizuoka	*Symplocos coreana*	n	gut	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	609/615	OP537750
Stag beetle	*Aesalus asiaticus*	Shizuoka	*Symplocos coreana*	n	gut	y 1	not analyzed		
Stag beetle	*Aesalus asiaticus*	Shizuoka	*Symplocos coreana*	n	gut	y	*Scheffersomyces ergatensis* strain ATCC MYA-4655	609/615	OP537751

*The digits at the columns of "*Aegosoma sinicum*" correspond to those in Figure 9.

**The numbers of colonies on the plates are indicated when a few colonies appeared.

Table S2. Alignment of nucleotide sequences used for phylogenetic tree construction

Corresponding Figure	Indications in the tree	DNA sequence with gaps
	Judolidia bangi	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Munamizoa maculata	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Pachytodes cometes	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGAC--CG--TCGGCGTCGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Aredolpona succedanea	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGGAAATTCATTCGCGTTTCGGCA--G--TCGGCGTCGGATGGATAGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Pyrrhona laeticolor	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--G--TCGGCGTTGGATGG-TCGAGTGACGGACGGCG-T-A-CTCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Nakanea vicaria	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCGGT-C-TACGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Ohbayashia nigromarginata	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--G--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CTCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Leptura ochraceofasciata	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCC--G--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCTTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Leptura modicenotata	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCC--G--TTGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCTTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTGGGACGTC
	Strangalomorpha tenuis	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCTCTG--TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCTCGTCCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Anastrangalia scotodes	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCA--G--TCGGCGTTGGATG-GTCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCGCCTAGTAGGACGTC
	Eustrangalis anticereducta	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTCGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGGACGTC
	Caraphia lepturoides	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTAGGATGG-TCGTGCGACGGACGGCG-TTCGCGCGTCG-TCCGCGTCCTCTT--TTCTTCGAACCGTCGACGCGTGCACTTTTCCCCTAGTAGGACGTC
Fig. 2	*Rhagium femorale*	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCTTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Ephies japonicus	AGTACGTGAAACCGTTTAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGTCT--G--TCGGCGTTGGACGGGTCGACTGACGGACGACG-T-A-CGCGTCG-TCCGCTTCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Mimostrangalia kurosonensi	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCTTTG--TTGGCGTCGAATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Strangalia koyaensis	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCTTCG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Mimostrangalia dulcis	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGCTTTGTGTCGGCGTCGAATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TTCTTCGAACCGACGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Parastrangalia lesnei	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTCGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Stenocorus caeruleipennis	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTGGGATGG-TGGGAGTGACGGATGGCG-TTC-TGCGTCG-TCCGCGCCTTCTT--TCCTTCGTACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Rhagium japonicum	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Pidonia aegrota	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--TG--CCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-ACCGCGCCTTCCT--TTCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Pidonia oyamae	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--TG--CCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCTT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Gaurotes doris	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTTGGATGG-TCGAGTGACGGATGGCG-T-A-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGATCCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Encyclops olivacea	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGGC--CG--TCGGCGTCGGATGG-TCGAGTGACGGATGGCG-T-T-CGCGTCG-TCCGCGCCTTCCT--TCCTTCGAACCGGCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Necydalis gigantea	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTTTCGAT--CG--CTGGCGTGGGACGGGTCGTGCGACGGACGACG-T-A-CGCGTCG-TTCGCGCCTTCCT--TCTTTCGAACCGCGATCGAACGCGTGCACTTTTCCCCTAGTGAGGACGTC
	Necydalis formosana	AGTACGTGAAACCGTTCAGGGGTAAACCTGAGAAACCCGAAAGGTCGAAAGGGGAAATTCATTCGCGTGTCGAC--GC--GCGGCGAGGGTCGGATCGGGCGACGGTCGGCG-T-A-CGCGCCGTTCCGCGCCTTCCT--TTCTTCGAACCGTCGTCGAACGCGTGCACTTTTCCCCTAGTAGGACGTC
	Anastrangalia scotodes Cv OP537521	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------AATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Anastrangalia scotodes Se OP537527	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Stictoleptura dichroa Se OP537530	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Caraphia lepturoides Ssp OP537535	TTGCCAGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Ephies japonicus Cv OP537537	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------TATTGTCGGGCCAGAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTATTTGTC----AATTTGTTTGATTAATTTC
	Eustrangalis anticereducta Ssp OP537539	TTGCCGGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Judolidia bangi Se OP537542	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Leptura annulalis Ssp OP537543	TTGCCAGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Leptura ochraceofasciata Ssp OP537548	TTGCCAGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Leptura ochraceofasciata Cv OP537557	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------AATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Leptura ochraceofasciata Sc OP537560	TTACCTGCGCTTAATTGCGCGGTAAACA------AATCT-TACACACAGTGTTTTTCTTTAT-T--AGAAACTTTTGCTTTGGTTTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAT---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAATTTC
	Pigostrangalia dulcis Cv OP537567	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------AATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Mimostrangalia kurosonensis Cv OP537569	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTATGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------TATTGTCGGGCCAGAGACTACACTAAACTTCAGATTTTTATTTGAATTGTT---TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Munamizoa maculata Se OP537572	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Munamizoa maculata Ssp OP537570	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Munamizoa maculata Ssp OP537571	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
Fig. 4	*Etorofus vicarius* Se OP537580	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Etorofus vicarius Ssp OP537584	TTGCCAGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Etorofus vicarius Se OP537581	TTACCTGCGCTTAATTGCGCGGTAAACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Ohbayashia nigromarginata Sc OP537585	TTACCTGCGCTTAATTGCGCGGTAAACA------AATCT-TACACACAGTGTTTTTCTTTAT-T--AGAAACTTTTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAT---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAATTTC
	Ohbayashia nigromarginata Cv OP537587	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------AATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Pachytodes cometes Se OP537597	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Pachytodes cometes Ssp OP537598	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Parastrangalia lesnei Cv OP537601	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------AATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Pyrrhona laeticolor Cv OP537602	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------AATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Pyrrhona laeticolor Sc OP537615	TTACCTGCGCTTAATTGCGCGGTAAACA------AATCT-TACACACAGTGTTTTTCTTTAT-T--AGAAACTTTTGCTTTGGTTTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAT---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAATTTC
	Pyrrhona laeticolor Se OP537628	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Pyrrhona laeticolor Sp OP537629	TTACCTGCGCTTAATTGCGCGGTAGACA------AA-CT-TACACACAGTGTTTT-CTTTAT-T---GAAACTATTGCTTTGGTCTGATTT-AGAAATA-AGTTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Rhagium femorale Ct OP537635	TTGCCAGCGCTTAACTGCGCGGCGAAAAC----TAATCCTTACACACTGTGTTTTTTGTTAT-TACAAGAACCATTGCTTTGGCTTGGCCG------CAAGGTTGGGCCAGAGGACACAC-AACTTCAATTTT--ATT-GAATTGTTAATTTTAATTATTTTGTC----AATT-GTTTGATTAATTTC
	Strangalia koyaensis Cv OP537643	CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTT-TT---GAAACATTGCTTTGGCTTGGCTT------CATTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTT--TTAATTTGTTTGTC----AATTTGTTTGATTAATTTC
	Strangalomorpha tenuis Ssp OP537646	TTGCCAGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Leptura subtilis Ssp OP537566	TTGCCAGCGCTTAACTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTT-CTTTATTA---GAAACTATTGCTTTGGTCTGGCTT-AGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTATTATTGAATTGTTATTTATTTAATTTGTC----AATTG--TTGATTAAATTC
	Megasemum quatricostulatum Dc OP537681	TTGCCAGCGCTTAATTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTTTCTTATTA---GAAACTATTGCTTTGGTCTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAATTCAATTTTT-TAATTGAATTGTTTATT-TAATTATAA----TGTCAATTTG--TTGATTAAATTC
	Megasemum quatricostulatum Ssp OP537683	TTACCTGCGCTTAATTGCGCGGTAAACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Megasemum quatricostulatum Se OP537684	TTACCTGCGCTTAATTGCGCGGTAACA------AA-CT-TACACACAGTGTTTTTCTTTAT-T---GAAACTATTGCTTTGGTCTGACTT-AGAAATA-AGCTGGGCCAGAGGTTTAA---AACTTCAATTTT-TAATTGAATTGTTTATTTATTTAATT-GTC----AATTG--TTGATTAAATTC
	Asemum striatum Sc OP537674	TTGCCAGCGCTTAATTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTTTCTTATT-A---AAACTATTGCTTTGGTCTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAATTTCAATTTT-TAATTGAATTGTTATT-TAATTA-ATTGTC----AATTG--TTGATTAAATTC
	Asemum striatum Dc OP537675	TTGCCAGCGCTTAATTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTTTCTTATT-A---AAACTATTGCTTTGGTCTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAATTTCAATTTT-TAATTGAATTGTTATT-TAATTA-ATTGTC----AATTG--TTGATTAAATTC
	Cephalallus unicolor Sv OP537677	TTGCCAGCGCTTAATTGCGCGGTAAAAA------AACCTTTACACACAGTGTTTTTCTTATTTA---GAAACTATTGCTTTGGTTTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT-TAATTATATAATGTATCAACTT----TGTTTAATTC

Fig. 5

```
Cephalallus unicolor  Dsp OP537676          TTGCCAGCGCTTAATTGCGCGGTAAAAA------AACCTTTACACACAGTGTTTTTCTTATTTA---GAAACTATTGCTTTGGTTTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT-TAATTATATAATGTATCAACTT----TGTTTAAATTC
Tetropium castaneum  Dsp OP537689           TTGCCAGCGCTTAGTTGCGCGGCGAAAA------AACCT-TACACACAGTGTTTTTCTTATT-A---AAAACTATTGCTTTGGTCTGGCTCTAGAAGTAGAGTTGGGTCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT-TAATTATAATGTC----AATTTG--TTGATTAAATTC
Strangalomorpha tenuis  Ad OP537752         ACA----G-TATTCT----TTTTGCCAGCGCTTAACTGCGCGGCG-------AAAAAACCTTACACACAGTG-TTTTCTTTATTAGAAACTATTGCTTTGGTCTGGCTTAGAAATAAGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATTTTATTTAATTTGTCAATTT-G-
Strangalomorpha tenuis  Ad OP537753         ACA----G-TATTCT----TTTTGCCAGCGCTTAACTGCGCGGCG-------AAAAAACCTTACACACAGTG-TTTTCTTTATTAGAAACTATTGCTTTGGTCTGGCTTAGAAATAAGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATTTTATTTAATTTGTCAATTT-G-
Strangalomorpha tenuis  Ad OP537754         ACA----G-TATTCT----TTTTGCCAGCGCTTAACTGCGCGGCG-------AAAAAACCTTACACACAGTG-TTTTCTTTATTAGAAACTATTGCTTTGGTCTGGCTTAGAAATAAGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATTTTATTTAATTTGTCAATTT-G-
Pyrrhona laeticolor  Ad OP537758            ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAATCTTACACACAGTGTTTTTCTTTATTAGAAACTTTTGCTTTGGTTTGGCTCTAGAAATAAGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pyrrhona laeticolor  Ad OP537759            ACAGTTAG-TTTATGTTCTCTCAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTTTTGAAAC-ATTGCTTTGGCTTGGCTTAA----TTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTTTTAATTTGTT-TGTCAATTTTGT
Pachytodes cometes  Ad OP537760             ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pachytodes cometes  Ad OP537761             ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pachytodes cometes  Ad OP537762             ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pachytodes cometes  Ad OP537763             ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pachytodes cometes  Ad OP537764             ACA----G-TATTCT----TTTTGCCAGCGCTTAACTGCGCGGCG-------AAAAAACCTTACACACAGTG-TTTTCTTTATTAGAAACTATTGCTTTGGTCTGGCTTAGAAATAAGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Ohbayashia nigromarginata  Ad OP537766      ACA----G-TATTC-----TTTTGCCAGCGCTTAACTGCGCGGCG-------AA-AAACCTTACACACAGTG-TCTTTTTGATACAGAACTCTTGCTTTGGTTTGGCTAGAGATAGGTTGGGCCAGAGGTTTA---AC-----AAAACA-CAATTTAATTATTTTTACAGTTA----GTCAAATT-T-
Ohbayashia nigromarginata  Ad OP589185      ACAGTTAG-TTTATGTTCTCTCAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTTTTGAAAC-ATTGCTTTGGCTTGGCTTAA----TTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTTTTAATTTGTT-TGTCAATTTTGT
Ohbayashia nigromarginata  Ad OP537767      ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAATCTTACACACAGTGTTTTTCTTTATTAGAAACTTTTGCTTTGGTTTGGCTCTAGAAATAAGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Ohbayashia nigromarginata  Ad OP537768      ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAATCTTACACACAGTGTTTTTCTTTATTAGAAACTTTTGCTTTGGTTTGGCTCTAGAAATAAGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Etorofus vicarius  Ad OP537769              AAT----GTTATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Etorofus vicarius  Ad OP537770              AAT----GTTATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Etorofus vicarius  Ad OP537771              ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Strangalomorpha tenuis  Lv OP537646         ACA----G-TATTCT----TTTTGCCAGCGCTTAACTGCGCGGCG-------AAAAAACCTTACACACAGTG-TTTTCTTTATTAGAAACTATTGCTTTGGTCTGGCTTAGAAATAAGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATTTTATTTAATTTGTCAATTT-G-
Pyrrhona laeticolor  Lv OP537602            ACAGTTAG-TTTATGTTCTCTCAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTTTTGAAAC-ATTGCTTTGGCTTGGCTTAA----TTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTTTTAATTTGTT-TGTCAATTTTGT
Pyrrhona laeticolor  Lv OP537615            ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAATCTTACACACAGTGTTTTTCTTTATTAGAAACTTTTGCTTTGGTTTGGCTCTAGAAATAAGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pyrrhona laeticolor  Lv OP537628            ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTG-TTTTCTTTATT-GAAACTATTGCTTTGGTCTGATTTAGAAATAAGTTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pyrrhona laeticolor  Lv OP537629            ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTG-TTTTCTTTATT-GAAACTATTGCTTTGGTCTGATTTAGAAATAAGTTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pachytodes cometes  Lv OP537597             ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Pachytodes cometes  Lv OP537598             ATT----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------GACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Ohbayashia nigromarginata  Lv OP537585      ACA----G-TATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAATCTTACACACAGTGTTTTTCTTTATTAGAAACTTTTGCTTTGGTTTGGCTCTAGAAATAAGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Ohbayashia nigromarginata  Lv OP537587      ACAGTTAG-TTTATGTTCTCTCAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTTACACACTGTGTTTTTGTTTTTTTGAAAC-ATTGCTTTGGCTTGGCTTAA----TTGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTTTTTAATTTGTT-TGTCAATTTTGT
Etorofus vicarius  Lv OP537580              AAT----GTTATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Etorofus vicarius  Lv OP537581              AAT----GTTATTAG----TTTTACCTGCGCTTAATTGCGCGGTA-------AACAAA-CTTACACACAGTGTTTTTCTTTATT-GAAACTATTGCTTTGGTCTGACTTAGAAATAAGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTATTTATTTAAT-TGTCAATTT-G-
Etorofus vicarius  Lv OP537584              ACA----G-TATTCT----TTTTGCCAGCGCTTAACTGCGCGGCG-------AAAAAACCTTACACACAGTG-TTTTCTTTATTAGAAACTATTGCTTTGGTCTGGCTTAGAAATAAGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATTTTATTTAATTTGTCAATTT-G-
```

Fig. 9

```
Anastrangalia scotodes  Cv OP537521          CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TAA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Anastrangalia scotodes  Se OP537527          TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Stictoleptura dichroa  Se OP537530           TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Caraphia lepturoides  Ssp OP537535           TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Ephies japonicus  Cv OP537537                CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTTGTTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TTA---T--TGTCGGGCCAGAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTATTTGTCAATTTGTTTGATTAAATTCAA
Eustrangalis anticereducta  Ssp OP537539     TTGCCGGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Judolidia bangi  Se OP537542                 TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Leptura annulalis  Ssp OP537543              TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Leptura ochraceofasciata  Ssp OP537548       TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Leptura ochraceofasciata  Cv OP537557        CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TAA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Leptura ochraceofasciata  Sc OP537560        TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Pigostrangalia dulcis  Cv OP537567           CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TAA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Mimostrangalia kurosonensis  Cv OP537569     CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTATGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TTA---T--TGTCGGGCCAGAGACTACACTAAACTTCAGATTTTTATTTGAATTGTT-----TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Munamizoa maculata  Se OP537572              TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Munamizoa maculata  Ssp OP537570             TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Munamizoa maculata  Ssp OP537571             TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Etorofus vicarius  Ssp OP537580              TTACCTGCGCTTAATTGCGCGGTAAACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Etorofus vicarius  Ssp OP537584              TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Etorofus vicarius  Se OP537581               TTACCTGCGCTTAATTGCGCGGTAAACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Ohbayashia nigromarginata  Sc OP537585       TTACCTGCGCTTAATTGCGCGGTAAACAA------A-TCT-T-ACACACAGTGTTTTTC--TTTATTAGAAACTTTTGCTTTGGTTTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Ohbayashia nigromarginata  Cv OP537587       CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TAA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Pachytodes cometes  Se OP537597              TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Pachytodes cometes  Ssp OP537598             TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Parastrangalia lesnei  Cv OP537601           CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TAA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Pyrrhona laeticolor  Cv OP537602             CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TAA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Pyrrhona laeticolor  Sc OP537615             TTACCTGCGCTTAATTGCGCGGTAAACAA------A-TCT-T-ACACACAGTGTTTTTC--TTTATTAGAAACTTTTGCTTTGGTTTGGTCT-TAGAAATA-AGTTGGGCCAGAGGTTTA---TAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Pyrrhona laeticolor  Se OP537628             TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTT-C--TTTATT-GAAACTATTGCTTTGGTCTGATT-TAGAAATA-AGTTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Pyrrhona laeticolor  Sp OP537629             TTACCTGCGCTTAATTGCGCGGTAGACAA------A-CT-T-ACACACAGTGTTTT-C--TTTATT-GAAACTATTGCTTTGGTCTGATT-TAGAAATA-AGTTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Rhagium femorale  Ct OP537635                TTGCCAGCGCTTAACTGCGCGGCGAAAAC----TAATCCTT-ACACACTGTGTTTTTTGTTATTACAAGAACCATTGCTTTGGCTTGGCT-TGGCC-GCA---A--GGTTGGGCCAGAGGACACAC-AAACTTCAATTT--TATT-GAATTGTTAAT--TTTAATTATTTGTCAATTT--GTTTGATTAAATTCAA
Strangalia koyaensis  Cv OP537643            CAGCCTGCGCTTAATTGCGCGGCGAGGAACAGAC-AACCTT-ACACACTGTGTTTTTGTTTTTT---TGAAACATTGCTTTGGCTTGGCT-TCA---T--TGTCGGGCCAAAGACTACACTAAACTTCAGATTTTTATTTGAATTGTTT-T--TTAATTTGTTTGTCAATTTGTTTGATTAAATTCAA
Strangalomorpha tenuis  Ssp OP537646         TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Leptura subtilis  Ssp OP537566               TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-TAGAAATA-AGTTGGGCCAGAGGTTTAACTAAACTTCAATTTTTATTATTGAATTGTTATT--TATTTAATTTGTCAATTT---GTTGATTAAATTCAA
Megasemum quatricostulatum  Dc OP537681      TTGCCAGCGCTTAATTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTTC--TTATTTAGAAACTATTGCTTTGGTCTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT---TAATTATAATGTCAATTG--TTGATTAAATTCAA
Megasemum quatricostulatum  Ssp OP537683     TTACCTGCGCTTAATTGCGCGGTAAACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Megasemum quatricostulatum  Se OP537684      TTACCTGCGCTTAATTGCGCGGTAAACAA------A-CT-T-ACACACAGTGTTTTTC--TTTATT-GAAACTATTGCTTTGGTCTGACT-TAGAAATA-AGCTGGGCCAGAGGTTTA---AAACTTCAATTTT-TAATTGAATTGTTTAT--TTATTTAATT-GTCAATTT--GTTGATTAAATTCAA
Asemum striatum  Sc OP537674                 TTGCCAGCGCTTAATTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTTC--TTATT-AAAAACTATTGCTTTGGTCTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAATTCAATTTT-TAATTGAATTGTTATT---TAATTAA-TTGTCAATTTG--TTGATTAAATTCAA
```

Asemum striatum Dc OP537675	TTGCCAGCGCTTAATTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTTC--TTATT-AAAAACTATTGCTTTGGTCTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAATTTCAATTTT-TAATTGAATTGTTATT---TAATTAA-TTGTCAATTTG--TTGATTAAATTCAA
Cephalallus unicolor Sv OP537677	TTGCCAGCGCTTAATTGCGCGGTAAAAA------ACCT-TTACACACAGTGTTTTTC--TTATTTAGAAACTATTGCTTTGGTTTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT---TAATTATATAATGTATCAACTTTGTTTAAATTCAA
Cephalallus unicolor Dsp OP537676	TTGCCAGCGCTTAATTGCGCGGTAAAAA------ACCT-TTACACACAGTGTTTTTC--TTATTTAGAAACTATTGCTTTGGTTTGGCTCTAGAAATAGAGTTGGGCCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT---TAATTATATAATGTATCAACTTTGTTTAAATTCAA
Tetropium castaneum Dsp OP537689	TTGCCAGCGCTTAGTTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTTC--TTATT-AAAAACTATTGCTTTGGTCTGGCTCTAGAAGTAGAGTTGGGTCAGAGGTCAAACCAAACTTCAATTTT-TAATTGAATTGTTATT---TAATTATAATGTCAATTTG--TTGATTAAATTCAA
As Shizuoka1 OP537700	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Mie1 OP537701	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Mie2 OP537703	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Shiga1 OP589184	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Mie3 OP537704	TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-CAGAAATG-AGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTTA---TTGAATTGTTATT--TTATT-ATTTGTCAATTT--GTTGATTAAATTCAA
As Kyoto2 OP537705	TTGCCTGCGCTTAATTGCGCGGCGAAAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTT-TACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Kyoto3 OP537706	TTGCCTGCGCTTAATTGCGCGGCGAAAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTT-TACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Ibaraki1 OP537708	TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-CAGAAATG-AGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTTA---TTGAATTGTTATT--TTATT-AATTTGTCAATTT--GTTGATTAAATTCAA
As Ibaraki2 OP537709	TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-CAGAAATG-AGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTTA---TTGAATTGTTATT--TTATT-AATTTGTCAATTT--GTTGATTAAATTCAA
As Ibaraki3 OP537710	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Ibaraki4 OP537711	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Ibaraki5 OP537712	TTGCCTGCGCTTAATTGCGCGGCGAGAAA------ACCT-T-ACACACAATGTTTTTGTTTTTATTAAGAAACTTTGCTTTGGTTTGGCG-TAGAGATA-CGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTT-TAATTGAATTGTTTTTAACTATTTTATTTGTCAATTGT-GATTTTATAATCAAT
As Yakushima OP537713	TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTT-C--TTTATTAGAAACTATTGCTTTGGTCTGGCT-CAGAAATG-AGTTGGGCCAGAGGTTTA-CCAAACTTCAATTTTA---TTGAATTGTTATT--TTATT-AATTTGTCAATTT--GTTGATTAAATTCAA
As Shizuoka2 OP537714	TTGCCAGCGCTTAACTGCGCGGCGAAAAA------ACCT-T-ACACACAGTGTTTT-C--TTTATTAGAAACTATTGCTTTGGTTTGGCT-CAGAAATG-AGTTGGGCCAGAGGTTTA-CTAAACTTCAATTTTA---TTGAATTGTTATT--TTATT-AATTTGTCAATTT--GTTGATTAAATTCAA

Table S2. Symbiont fungi isolated from female adults of Longicorn beetles in Japan

Group	Species	Collected location	Analyzed region	Colonies on plate	Blastn Best Hit	Accession Number
Lepturinae	*Strangalomorpha tenuis*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	OP537752
Lepturinae	*Strangalomorpha tenuis*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	OP537753
Lepturinae	*Strangalomorpha tenuis*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	OP537754
Lepturinae	*Rhagium femorale*	Yamanashi	Female oviduct	no yeast colony		
Lepturinae	*Rhagium femorale*	Yamanashi	Female oviduct	y	[*Candida*] *rhagii* culture CBS:618	OP537755
Lepturinae	*Rhagium femorale*	Yamanashi	Female oviduct	y	[*Candida*] *rhagii* culture CBS:618	OP537756
Lepturinae	*Rhagium femorale*	Yamanashi	Female oviduct	y	[*Candida*] *rhagii* culture CBS:618	OP537757
Lepturinae	*Pyrrhona laeticolor*	Shizuoka	Female oviduct	y	*Scheffersomyces coipomoensis* strain Y3	OP537758
Lepturinae	*Pyrrhona laeticolor*	Shizuoka	Female oviduct	y	[*Candida*] *vrieseae* CBS 10829	OP537759
Lepturinae	*Pachytodes cometes*	Gifu	Female oviduct	y	*Candida ergatensis* strain ATCC MYA-4655	OP537760
Lepturinae	*Pachytodes cometes*	Gifu	Female oviduct	y	*Candida ergatensis* strain ATCC MYA-4655	OP537761
Lepturinae	*Pachytodes cometes*	Gifu	Female oviduct	y	*Candida ergatensis* strain ATCC MYA-4655	OP537762
Lepturinae	*Pachytodes cometes*	Gifu	Female oviduct	y	*Scheffersomyces* sp. strain Y13	OP537763
Lepturinae	*Pachytodes cometes*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	OP537764
Lepturinae	*Pachypidonia bodemeyeri*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	OP537765
Lepturinae	*Ohbayashia nigromarginata*	Wakamaya	Female oviduct	y	*Meyerozyma guilliermondii* isolate NTOU5003	OP537766
					[*Candida*] *vrieseae* CBS 10829	OP589185
Lepturinae	*Ohbayashia nigromarginata*	Shizuoka	Female oviduct	y	*Scheffersomyces coipomoensis* strain Y3	OP537767
Lepturinae	*Ohbayashia nigromarginata*	Shizuoka	Female oviduct	y	*Scheffersomyces coipomoensis* strain Y3	OP537768
Lepturinae	*Etorofus vicarius*	Nagano	Female oviduct	no yeast colony		
Lepturinae	*Etorofus vicarius*	Nagano	Female oviduct	y	*Candida ergatensis* strain ATCC MYA-4655	OP537769
Lepturinae	*Etorofus vicarius*	Nagano	Female oviduct	y	*Candida ergatensis* strain ATCC MYA-4655	OP537770
Lepturinae	*Etorofus vicarius*	Yamanashi	Female oviduct	y	*Candida ergatensis* strain ATCC MYA-4655	OP537771
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Yamanashi	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Nagano	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Nagano	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Nagano	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Nagano	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Leptura ochraceofasciata*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. Y1052_SICYLG3	
Lepturinae	*Encyclops olivaceus*	Yamanashi	Female oviduct	no yeast colony	*Chrysolina oricalcia* genome assembly, chromosome: 12	OP537772
Lepturinae	*Encyclops olivaceus*	Yamanashi	Female oviduct	no yeast colony	*Chrysolina oricalcia* genome assembly, chromosome: 12	OP537773
Necydalinae	*Necydalis sacharinensis*	Nagano	Female oviduct	no yeast colony	*Chrysolina oricalcia* genome assembly, chromosome: 12	OP537774
					Uncultured fungus isolate FICUS 144	OP537775
Necydalinae	*Necydalis sacharinensis*	Nagano	Female oviduct	no yeast colony	*Chrysolina oricalcia* genome assembly, chromosome: 12	OP537776
					Uncultured fungus isolate FICUS 144	OP537777
Prioninae	*Aegosoma sinicum*	Shizuoka	Female oviduct	y	*Scheffersomyces anoplophorae*	OP537778
Prioninae	*Aegosoma sinicum*	Shizuoka	Female oviduct	y	*Scheffersomyces* sp. YW27.1	OP537779
Prioninae	*Aegosoma sinicum*	Shizuoka	Female oviduct	y	*Scheffersomyces anoplophorae*	OP537780
					Scheffersomyces anoplophorae	OP537781
Prioninae	*Aegosoma sinicum*	Shizuoka	Egg surface	y	*Meyerozyma guilliermondii* strain CBS 2030	OP537782
					Meyerozyma carpophila strain CBS 5256	OP537783
					Scheffersomyces anoplophorae	OP537784
					Kodamaea ohmeri strain DM9Y	OP537785
					Kodamaea ohmeri strain SM14UFAM	OP537786
					Debaryomyces hansenii IFM 55872	OP537787
Prioninae	*Aegosoma sinicum*	Osaka	Egg surface	y	[*Candida*] *railenensis* isolate R97308_ITS	OP537788
					Meyerozyma caribbica strain CBS 9966	OP537789
					Meyerozyma guilliermondii isolate NTOU5003	OP537790
					[*Candida*] *oleophila* isolate B2-15b	OP537791
					Wickerhamomyces sydowiorum culture CBS:7276	OP537792

Table S4. Artificial transfer of yeasts to larvae

Species and biological replicate	Sterilization	Given yeast	Detected yeast	Number of larvae	Total
Pyrrhona laeticolor	No	Not given	Other yeast	5	10
Pyrrhona laeticolor	No	Not given	Symbiont yeast	2	
Pyrrhona laeticolor	No	Not given	No yeast detected	3	
Pyrrhona laeticolor	Yes	Not given	No yeast detected	13	28
Pyrrhona laeticolor	Yes	*Scheffersomyces* sp. Y1052_SICYLG3	Given yeast	12	
Pyrrhona laeticolor	Yes	*Scheffersomyces* sp. Y1052_SICYLG3	No yeast detected	2	
Pyrrhona laeticolor	Yes	Not given	Other yeast	1	
Ohbayashia nigromarginata	No	Not given	No yeast detected	1	3
Ohbayashia nigromarginata	No	Not given	Symbiont yeast	2	
Ohbayashia nigromarginata	Yes	Not given	No yeast detected	3	9
Ohbayashia nigromarginata	Yes	*Scheffersomyces* sp. Y1052_SICYLG3	No yeast detected	5	
Ohbayashia nigromarginata	Yes	*Scheffersomyces coipomoensis* strain Y3	Given yeast	1	
Leptura ochraceofasciata Exp 1	No	Not given	Symbiont yeast	15	15
Leptura ochraceofasciata Exp 1	Yes	Not given	No yeast detected	6	19
Leptura ochraceofasciata Exp 1	Yes	Not given	Other yeast	1	
Leptura ochraceofasciata Exp 1	Yes	*Scheffersomyces coipomoensis* strain Y3	Given yeast	8	
Leptura ochraceofasciata Exp 2	Yes	*Scheffersomyces coipomoensis* strain Y3	No yeast detected	3	
Leptura ochraceofasciata Exp 2	Yes	*Candida vrieseae*	Other yeast	1	

Table S5. Possession of symbiont cells in the mycetome after long culture

Species name	Number of exaimined larva	Number of larva possessing symbiont cells in the mycetomes	Weight at collection (mg)	Weight at observation (mg)	Date of collection	Date of observation
Etorofus vicarius	1	1	356	420	20-Aug-2020	03-July-2021
Leptura ochraceofasciata	1	1	40	199	20-Aug-2020	03-July-2021
Necydalis formosana	2	0	68 129	267 352	20-Aug-2020	03-July-2021