Diving Behavior Reveals Humidity Sensing Ability of Water Deprived Planarians

Yu Pei[1], Renzhi Qian[1], Yuan yan[1], Yixuan Zhang[1], Liyuan Tan[1], Xinran Li[1], Chenxu Lu[1], Yuxuan Chen[1], Yuanwei Chi[1], Kun Hao[1], Zhen Xu[1], Guang Yang[1], Zilun Shao[1], Yuhao Wang[1] and Kaiyuan Huang[1,2,3]

[1]College of Biological Science, China Agricultural University; Beijing, 100193, China.

[2]Tsinghua Institute of Multidisciplinary Biomedical Research (TIMBR), Tsinghua University;

17 Beijing, 100084, China

[3]National Institute of Biological Sciences (NIBS); Beijing, 102206, China.

* Kaiyuan Huang

Email: huangkaiyuan@nibs.ac.cn

Author Contributions: Yu Pei, Renzhi Qian and Yuan yan contributed equally to this work. Kaiyuan Huang, Yu Pei, Renzhi Qian and Yuan yan designed research. Yu Pei, Renzhi Qian, Yuan yan, Yixuan Zhang, Liyuan Tan, Xinran Li, Chenxu Lu, Yuxuan Chen, Yuanwei Chi and Kun Hao performed research. Zhen Xu, Guang Yang, Zilun Shao and Yuhao Wang analyzed data. Kaiyuan Huang and Yu Pei wrote the manuscript.

Competing Interest Statement: The authors declare that there is no competing interest in the study.

Classification: Biological Sciences

Keywords: planarians, humidity sensing, aquatic animals, decision making, survival seeking.

Abstract

Humidity sensing ability is crucial to terrestrial animals for fitting the environment. Researchers
made great progress in recent study about humidity sensing mechanisms of terrestrial animals.
However, it is poorly understood whether humidity sensing exists in aquatic animals. Here, we
demonstrate that the aquatic planarians, one of the primitive forerunners of later animals, has the
ability of humidity sensing and is capable of using the ability to perceive the direction of water
from a drought place to seek survival. The behavior we discovered is described as diving
because the worms twist its body to break away from the mucus that make them adhere to the
drought place and drop into the water. The behavior is triggered by rapidly increasing humidity.
This finding suggests that humidity sensing ability exists in the lower aquatic animals, and the
ability might be used to seek for water when aquatic animals are facing desiccation. The finding
also suggests that survival-seeking and decision-making behavior have appeared in the primitive
planarian worms.

Main Text
16
17   **Introduction**
18
As a universal medium for biochemical events, water is an indispensable resource for all
animals. For terrestrial animals, they are at constant risk for desiccation due to unpredictable
climate change. Therefore, humidity sensing has been widely investigated in terrestrial animals
for their need for a comfortable environment. Recent studies revealed detailed mechanisms of
how terrestrial animals sense humidity(1-3).
In contrast to terrestrial animals, aquatic animals have much less possibility to face a situation
of desiccation. However, dehydration is usually fatal to aquatic animals for they need water to
respire. So, it might also be crucial for some aquatic animals to perceive the direction of water
when facing an emergent situation of water depletion. Nevertheless, it is poorly understood
whether this ability exists in aquatic animals.
Planarian is a kind of aquatic free-living flatworm to have first evolved a centralized brain. As a
primitive forerunner of later animals, the planarians can be evolutionarily instructive for the
investigation of later animals. For freely living in the natural environment, planarians have evolved
various sensory abilities, including sensitivity to light(4), temperature(5), water currents(6),
chemical gradients(7), vibration(8), magnetic fields(9) and electric fields(10), but its humidity
sensing ability is not yet identified.
Unlike most aquatic animals who live freely in the water, planarians usually live and stick under
rocks, debris and water plants in streams, ponds, and springs(11). Therefore, they are confronted
with frequently falling water levels and might be lifted out of the water. So, it might be important
for planarians to perceive the direction of water to seek survival under such emergent situations.
Thus, we speculate that planarians have the ability of humidity sensing to carry out such tasks.
To prove this hypothesis, we established a behavioral paradigm of planarians called 'diving',
which will be explained in detail in the result section. And then we demonstrate that the worms
can perceive humidity and its increasing speed to judge the direction of the water. Our finding
identified the humidity sensing ability of a kind of aquatic animal and explained what this ability is
used for, which is yet not discovered in this field. This finding also suggests that survival-seeking
and decision-making behavior have appeared in the primitive planarian worms and might shed
light on how these abilities evolved. The finding also provides a 'diving' behavioral paradigm for
future study.
48
49   **Results**
50
1. Rapidly increasing humidity induces diving behavior of planarians

We established a behavioral paradigm of planarians called 'diving' (Fig. 1A, SI movie 1). A
planarian worm is put in a petri dish and its surrounding water is wiped out. Then the petri dish is
inverted onto a 250 mL beaker containing 200 mL of water. The worm will first struggle to find
water and then uplift its head. When the worm decides to dive, it will twist its body to break away
from the mucus and drop into the water.
We totally tested 20 worms through the diving paradigm and most of the worms started the
diving behavior in 60 seconds, showing that they might be able to perceive the water under them.
Then we tested 20 worms with a dry 250 mL beaker, all of the worms did not perform the diving
behavior and finally stopped moving (Fig. 1B).
We speculate that this behavior is related to the increase of humidity. Thus, to control the
humidity conditions, all of the diving experiment was carried out in a 38%±1% relative humidity
(RH) environment if not otherwise stated. We measured the RH variation in the two processes
above. To simulate the situation of a worm in the experiment while measuring the RH variation,
we embedded the humidity meter probe in the middle of a foam plastic board and then put the
board on the beaker (Fig. 1E). The result reveals a rapid increasing RH in the 250 mL beaker
containing 200 mL water, which increases from 38% RH to more than 60% RH in 30 seconds.
(Fig. 2A). In contrast, the RH of the dry 250 mL beaker maintained relatively constant at 38%±1%.
The time that a worm starts the diving behavior is counted and synchronized with the RH
variation. (Fig. 2A, Fig. 2B).
The worms dropped at the mean time of 13 seconds and the corresponding RH is 54.1%. We
argued that the diving behavior might be induced by high constant humidity rather than rapidly
increasing humidity. Therefore, we tested 20 worms with dry 250 mL beaker at an at a constant
RH of 65%±5%. Although the worms can struggle to crawl for a long time due to low evaporation
rate, none of the worms performed diving behavior. Hence, we conclude that the diving behavior
of the worm is induced by rapidly increasing humidity rather than constant high humidity.
77
2. Slower increasing humidity hinders planarians' decision to dive
To investigate whether diving behavior can be induced by slower increasing humidity, we
tested 20 worms with a 250 mL beaker containing 50 mL and measured its humidity variation.
(Fig. 1C, Fig. 2B). The humidity increase rate in this case is about one half of 250 mL beaker
containing 200 mL of water. Surprisingly, some of the worms didn't drop and some of the worms
took about minutes to drop. We reasoned that whether to execute the diving behavior involves
the worm's decision. Slower increasing humidity makes some of the worms hesitate to drop,
which further proves that only rapidly increasing humidity can solidly induce the diving behavior of
planarians.
87
3. Rapidly increasing humidity can mislead planarians drop into a dry place
To further demonstrate the diving behavior is induced by rapidly increasing humidity, we
simulated a situation of rapidly increasing humidity, yet no water is provided if the worm drops.
Instead of using a large quantity of water, we sprayed water droplets on the wall of the beaker
and put a piece of dry plastic to cover the bottom of the beaker (Fig. 1D). Then we tested 20
worms in this beaker and measured the humidity variation, all of the worms started to drop on the
dry plastic in 60 seconds (Fig. 2D). This result confirmed the conclusion that rapidly increasing
humidity induces the diving behavior of planarians.
96
Discussion
The investigation of mechanisms of humidity sensing had been focused on terrestrial animals.
Including how hygroreceptor works in insects like P. americana(12) and D. melanogaster(13), and
the integration of mechano and thermo inputs of C. elegans(2) and humans(3). However, the
humidity sensing of aquatic animals was hardly ever considered in previous studies. In the
present study, we unveiled the ability of humidity sensing of aquatic planarians by establishing
the diving behavioral paradigm, which they use to seek survival under dehydration conditions.

Our work reveals that in the diving behavioral paradigm, a worm facing dehydration has to
make a quick decision whether or not to secede from the attached surface before it cannot move
anymore. In this process, the worm continues to raise its head probably to sense the increasing
humidity, which would accelerate the rate of evaporation. So, the judgment of the worm must be
accurate to deal with such an emergent situation. As our result shows increasing rate of humidity
have become a crucial indicator for the worm's decision.
As illustrated above, the diving behavior of planarians can be classified as a decision-making
and survival-seeking behavior. Being one of the first kinds of animals to have evolved a
centralized brain, planarians' behavior can provide instructions from the evolutionary perspective
for investigating the behaviors of later animals. Our results demonstrate that the decision-making
and survival-seeking behavior had already developed in the primitive planarian worms, which
might provide a new evolutionary perspective for investigating such behaviors.
116
117 **Materials and Methods**
118
119 **Planarians** A laboratory strain of *D. japonica*, originating from wild collected *D. japonica*
120 (identified by cytochrome c oxidase subunit 1 gene) from the Cherry-Valley in Beijing Botanical
121 Garden, Haidian district, Beijing, China in 2019. Worms are maintained in Montjuic Water(14) in
122 the dark and fed with chicken liver twice a week. Worms are fed 2 days before experiment. The
123 length of planarians used in the experiment varies from 1.5 cm to 2.5 cm.
124
125 **Experimental Setup** A 250mL glass beaker and a plastic petri dish is used in the experiment.
126 Kimwipes paper towel is used to wipe water. A UT331+ humidity meter (Uni-Trend Technology
127 (China) Co., Ltd.) is used to measure and record the RH.
128
129 **Test Procedure** A planarian is transferred to the petri dish containing water from home well by a
130 transfer pipette. Wait until the worm sink and attach to the bottom of the petri dish. Slowly pour
131 the water out while maintaining the worm attached to the bottom. Wipe out the water in the petri
132 dish but do not touch the worm. The petri dish is washed by water and wiped between each
133 worm's test.
134
135 **Humidity Measure Procedure** The primer of the humidity meter is embedded into a foam plastic
136 board then cover the beaker and immediately start measuring for 90 second.
137
138 **Statistical Analysis** All data were analyzed using PRISM (GraphPad Prism 9.0.0(121)).
139 Nonlinear regression (curve fit): polynomial (fourth order) is used to generate fitting results of RH
140 variation.
141
142 **Acknowledgments**
143

144 We wish to thank Prof. Baoqing Wang, Prof. Zhengxin Ying and Dr. Wei Wu for suggestions and
145 financial support. Figure 1 is created with BioRender.com

146
References

1. Enjin A*, et al.* (2016) Humidity Sensing in Drosophila. *Current biology : CB* 26(10):1352-
1358.
2. Russell J, Vidal-Gadea A, Makay A, Lanam C, & Pierce-Shimomura J (2014) Humidity
sensation requires both mechanosensory and thermosensory pathways in
Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United*
States of America 111.

3.   Filingeri D, Fournet D, Hodder S, & Havenith G (2014) Why wet feels wet? A
155   neurophysiological model of human cutaneous wetness sensitivity. *Journal of*
156   *Neurophysiology* 112:1457.
157   4.   Shettigar N*, et al.* (2017) Hierarchies in light sensing and dynamic interactions between
158   ocular and extraocular sensory networks in a flatworm. *Science Advances* 3:e1603025.
159   5.   Inoue T, Hoshino H, Yamashita T, Shimoyama S, & Agata K (2015) Planarian shows
160   decision-making behavior in response to multiple stimuli by integrative brain function.
161   *Zoological Letters* 1(1):7.
162   6.   Allen GD (1915) Reversibility of the Reactions of Planaria Dorotocephala to a Current of
163   Water. *Biological Bulletin* 29(2):111-128.
164   7.   Mason P (1975) Chemo-klino-kinesis in planarian food location. *Animal behaviour*
165   23:460-469.
166   8.   Dessì-Fulgheri F & Messeri P (1973) [Use of 2 different negative reinforcements in light-
167   darkness discrimination of planarians]. *Bollettino della Società italiana di biologia*
168   *sperimentale* 49:1141-1145.
169   9.   Brown F & Chow C (1975) Differentiation between Clockwise and Counterclockwise
170   Magnetic Rotation by the Planarian, Dugesia dorotacephala. *Physiological Zoology*
171   48:168-176.
172   10.   Brown H & Ogden T (1968) The Electrical Response of the Planarian Ocellus. *The Journal*
173   *of general physiology* 51:237-253.
174   11.   Vila-Farré M & Rink J (2018) The Ecology of Freshwater Planarians.), Vol 1774, pp 173-
175   205.
176   12.   Tichy H & Kallina W (2010) Insect Hygroreceptor Responses to Continuous Changes in
177   Humidity and Air Pressure. *Journal of neurophysiology* 103:3274-3286.
178   13.   Liu L*, et al.* (2007) Drosophila hygrosensation requires the TRP channels water witch and
179   nanchung. *Nature* 450:294-298.
180   14.   Merryman S, Sánchez Alvarado A, & Jenkin J (2018) Culturing Planarians in the
181   Laboratory.), Vol 1774, pp 241-258.

182
183

184
185 **Figures**

186
187 **Figure 1.** Illustration of the experiment process. (A-D) The diving experiment. (E) The relative
188 humidity measurement.
189
190 <insert page break here>
191
192

193
Figure 2. Diving behavior are induced by rapidly increasing humidity. 3 sets of RH data are used
for nonlinear curve fitting in (A-C). The lower panel of (A-C) shows that the time a worm starts the
diving behavior synchronized with the RH variation, time data is presented as mean ± SEM. (A)
Rapidly increasing humidity induces diving behavior of planarians (n=20). (B) Slower increasing
humidity hinders planarians' decision to dive (n=20, 6 worms did not dive, 2 worms used more
than 90 seconds). (C) Rapidly increasing humidity can mislead planarians drop into a dry place
(n=20). (D) The percentage of worms dived in each group.
201
202
203