Diving Behavior Reveals Humidity Sensing Ability of Water Deprived Planarians

Yu Pei[1], Renzhi Qian[1], Yuan yan[1], Yixuan Zhang[1], Liyuan Tan[1], Xinran Li[1], Chenxu Lu[1], Yuxuan Chen[1], Yuanwei Chi[1], Kun Hao[1], Zhen Xu[1], Guang Yang[1], Zilun Shao[1], Yuhao Wang[1] and Kaiyuan Huang[1]

[1]College of Biological Science, China Agricultural University; Beijing, 100193, China.

* Kaiyuan Huang

Email: cfhuangkaiyuan@gmail.com

Author Contributions: Yu Pei, Renzhi Qian and Yuan yan contributed equally to this work. Kaiyuan Huang, Yu Pei, Renzhi Qian and Yuan yan designed research. Yu Pei, Renzhi Qian, Yuan yan, Yixuan Zhang, Liyuan Tan, Xinran Li, Chenxu Lu, Yuxuan Chen, Yuanwei Chi and Kun Hao performed research. Zhen Xu, Guang Yang, Zilun Shao and Yuhao Wang analyzed data. Kaiyuan Huang and Yu Pei wrote the manuscript.

Competing Interest Statement: The authors declare that there is no competing interest in the study.

Classification: Biological Sciences

Keywords: planarians, humidity sensing, aquatic animals, decision making, survival seeking.

Abstract

Humidity sensing ability is crucial to terrestrial animals for fitting the environment. Researchers
made great progress in recent study about humidity sensing mechanisms of terrestrial animals.
However, it is poorly understood whether humidity sensing exists in aquatic animals. Here, we
demonstrate that the aquatic planarians, one of the primitive forerunners of later animals, has the
ability of humidity sensing and is capable of using the ability to perceive the water beneath itself
from a drought place to seek survival. The behavior we discovered is described as diving
because the worms twist its body to break away from the mucus that make them adhere to the
drought place and drop into the water. The behavior is triggered by rapidly increasing humidity.
This finding suggests that humidity sensing ability exists in the lower aquatic animals, and the
ability might be used to seek for water when aquatic animals are facing desiccation. The finding
also suggests that survival-seeking and decision-making behavior have appeared in the primitive
planarian worms.

Main Text

16
17 **Introduction**

18
As a universal medium for biochemical events, water is an indispensable resource for all
animals. For terrestrial animals, they are at constant risk for desiccation due to unpredictable
climate change. Therefore, humidity sensing has been widely investigated in terrestrial animals
for their need for a comfortable environment. Recent studies revealed detailed mechanisms of
how terrestrial animals sense humidity[1-3].
In contrast to terrestrial animals, aquatic animals have much less possibility to face a situation
of desiccation. However, dehydration is usually fatal to aquatic animals for they need water to
respire. So, it might also be crucial for some aquatic animals to perceive the direction of water
when facing an emergent situation of water depletion. Nevertheless, it is poorly understood
whether this ability exists in aquatic animals.
Planarian is a kind of aquatic free-living flatworm to have first evolved a centralized brain. As a
primitive forerunner of later animals, the planarians can be evolutionarily instructive for the
investigation of later animals. For freely living in the natural environment, planarians have evolved
various sensory abilities, including sensitivity to light[4], temperature[5], water currents[6],
chemical gradients[7], vibration[8], magnetic fields[9] and electric fields[10], but its humidity
sensing ability is not yet identified.
Unlike most aquatic animals who live freely in the water, planarians usually live and stick under
rocks, debris and water plants in streams, ponds, and springs[11]. Therefore, they are confronted
with frequently falling water levels and might be lifted out of the water. So, it might be important
for planarians to perceive the direction of water to seek survival under such emergent situations.
Thus, we speculate that planarians have the ability of humidity sensing to carry out such tasks.
To prove this hypothesis, we established a behavioral paradigm of planarians called 'diving',
which will be explained in detail in the result section. And then we demonstrate that the worms
can perceive humidity and its increasing speed to judge the direction of the water. Our finding
identified the humidity sensing ability of a kind of aquatic animal and explained what this ability is
used for, which is yet not discovered in this field. This finding also suggests that survival-seeking
and decision-making behavior have appeared in the primitive planarian worms and might shed

light on how these abilities evolved. The finding also provides a 'diving' behavioral paradigm for
future study.
48
49 **Results**
50
1. Rapidly increasing humidity induces diving behavior of planarians
We established a behavioral paradigm of planarians called 'diving' (Fig. 1A, SI movie 1). A
planarian worm is put in a petri dish and its surrounding water is wiped out. Then the petri dish is
inverted onto a 250 mL beaker containing 200 mL of water. The worm will first attempt to explore
around and then uplift its head. Finally, it will crawl and twist its body to break away from the
mucus and drop into the water. This is more likely to be a behavior rather than a physical
phenomenon.
We totally tested 20 worms through the diving paradigm and most of the worms started the
diving behavior in 60 seconds, showing that they might be able to perceive the water under them.
Then we tested 20 worms with a dry 250 mL beaker, all of the worms did not perform the diving
behavior and finally stopped moving (Fig. 1B, SI movie 2). A worm will retract its extending head
and reattach to the petri dish several times without diving. We speculate that this behavior is
related to the increase in humidity. We measured the RH variation in the two processes above.
To simulate the situation of a worm in the experiment while measuring the RH variation, we
embedded the humidity meter probe in the middle of a foam plastic board and then put the board
on the beaker (Fig. 1E). The result reveals a rapidly increasing RH in the 250 mL beaker
containing 200 mL water, which increases from 38% RH to more than 60% RH in 30 seconds.
(Fig. 2A). In contrast, the RH of the dry 250 mL beaker was maintained relatively constant at 38%
±1% (Fig. 2B). The time that a worm starts the diving behavior is counted and synchronized with
the RH variation. (Fig. 2A, Fig. 2F).
We argued that the diving behavior might be induced by other factors such as temperature,
density of mucus, or gravity. To control the humidity conditions, all of the diving experiment was
carried out in a 38%±1% relative humidity (RH) environment if not otherwise stated. All
experiments were carried out at $25\pm1℃$ to eliminate the influence of temperature. In the 250 mL
group, the worms dropped at the mean time of 13 seconds and the corresponding RH is 54.1%.
To exclude other factors that might interpret the diving behavior as a physical phenomenon rather
than behavior, we tested 20 worms with dry 250 mL beaker at an constant RH of 65%±5% (Fig.
2C). Although the worms could struggle to crawl for a long time due to high humidity, none of the
worms performed diving behavior. Hence, we conclude that the diving behavior of the worm is
induced by rapidly increasing humidity rather than constant high humidity.
81
2. Slower increasing humidity hinders planarians' decision to dive
To investigate whether diving behavior can be induced by slower increasing humidity, we
tested 20 worms with a 250 mL beaker containing 50 mL and measured its humidity variation.
(Fig. 1C, Fig. 2D). The humidity increase rate in this case is about one half of 250 mL beaker
containing 200 mL of water. Surprisingly, some of the worms didn't drop and some of the worms
took about minutes to drop (Fig. 2D, Fig. 2F). We reasoned that whether to execute the diving
behavior involves the worm's decision. Slower increasing humidity makes some of the worms
hesitate to drop, which further proves that only rapidly increasing humidity can solidly induce the
diving behavior of planarians.
91
3. Rapidly increasing humidity can mislead planarians drop into a dry place
To further demonstrate the diving behavior is induced by rapidly increasing humidity, we
simulated a situation of rapidly increasing humidity, yet no water is provided if the worm drops.
Instead of using a large quantity of water, we sprayed water droplets on the wall of the beaker
and put a piece of dry plastic to cover the bottom of the beaker (Fig. 1D). Then we tested 20
worms in this beaker and measured the humidity variation, all of the worms started to drop on the

dry plastic in 60 seconds (Fig. 2E, Fig. 2F). This result confirmed the conclusion that rapidly
increasing humidity induces the diving behavior of planarians.
100
Discussion
We noticed that the diving of the planarian is quite different from a simple drop. A planarian will
first attempt to crawl around, but soon stop crawling and extend its head. If a worm would dive, as
shown by the 250 mL group (Fig. 1A, SI movie 1), it would first uplift its head (extending more
than 1/2 of its body), then twist and crawl to make the tail (attached to the petri dish) move
forward. As the attachment gradually decreased to a point, the planarian severely twisted its body
to break away from mucus and dropped into the water eventually. If a worm would not dive, as
shown by the two constant humidity groups (Fig. 1B, SI movie 2), it would also uplift its head but
soon retracted its head (usually extending more than 1/2 of its body) and reattach to the petri
dish. The worms might attempt to uplift its head for several times, but eventually, it will dehydrate
and die on the petri dish. It seems that the planarian sensed no water below and decided not to
dive. In addition, because high constant RH of $65\% \pm 5\%$ did not trigger the diving behavior of
planarians (Fig. 2C, Fig 2F), this diving behavior cannot be concluded as it is the lower density of
mucus that makes worms drop and high density mucus lost moisture which prevented the drop.
Above illustrates that diving is an innate behavior induced by the rapidly increasing humidity
rather than a physical phenomenon.
The investigation of mechanisms of humidity sensing had been focused on terrestrial animals.
Including how hygroreceptor works in insects like P. americana[12] and D. melanogaster[13], and
the integration of mechano and thermo inputs of C. elegans[2] and humans[3]. However, the
humidity sensing of aquatic animals was hardly ever considered in previous studies. In the
present study, we unveiled the ability of humidity sensing of aquatic planarians by establishing
the diving behavioral paradigm, which they use to seek survival under dehydration conditions. As
a kind of aquatic animal, planarians might not have hygroreceptors to directly sense water, and
there is an obvious chemical barrier that would limit a planarian covered in mucus to sense the
external air. Therefore, we speculate that planarians can indirectly perceive humidity through the
change of mucus properties.
Our work reveals that in the diving behavioral paradigm, a worm facing dehydration has to
make a quick decision whether or not to secede from the attached surface before it cannot move
anymore. In this process, the worm continues to raise its head probably to sense the increasing
humidity, which would accelerate the rate of evaporation. So, the judgment of the worm must be
accurate to deal with such an emergent situation. As our result shows increasing rate of humidity
has become a crucial indicator for the worm's decision.
As illustrated above, the diving behavior of planarians can be classified as a decision-making
and survival-seeking behavior. Being one of the first kinds of animals to have evolved a
centralized brain, planarians' behavior can provide instructions from the evolutionary perspective
for investigating the behaviors of later animals. Our results demonstrate that the decision-making
and survival-seeking behavior had already developed in the primitive planarian worms, which
might provide a new evolutionary perspective for investigating such behaviors.
139
140 **Materials and Methods**
141
142 **Planarians** A laboratory strain of *D. japonica*, originating from wild collected *D. japonica*
143 (identified by cytochrome c oxidase subunit 1 gene) from the Cherry-Valley in Beijing Botanical
144 Garden, Haidian district, Beijing, China in 2019. Worms are maintained in Montjuic Water[14] in
145 the dark and fed with chicken liver twice a week. Worms are fed 2 days before experiment. The
146 length of planarians used in the experiment varies from 1.5 cm to 2.5 cm.
147
148 **Experimental Setup** A 250mL glass beaker and a plastic petri dish is used in the experiment.
149 The internal diameter of 250 mL beaker is 66.32 mm, the external diameter is 70.38 mm, the
150 internal height is 94.24 mm, and the external height is 96.72 mm. Kimwipes paper towel is used

151 to wipe water. A UT331+ humidity meter (Uni-Trend Technology (China) Co., Ltd.) is used to
152 measure and record the RH.
153
Test Procedure A planarian is transferred to the petri dish containing water from home well by a
transfer pipette. Wait until the worm sink and attach to the bottom of the petri dish. Slowly pour
the water out while maintaining the worm attached to the bottom. Wipe out the water in the petri
dish but not touch the worm. Then absorb water on the worm from its caudal direction until there
is no water film. At this time, there is little water on the worm and no water in the petri dish. The
petri dish is washed by water and wiped between each worm's test. After the test, the animals
were saved and well maintained.
161
162 **Humidity Measure Procedure** The primer of the humidity meter is embedded into a foam plastic
163 board then cover the beaker and immediately start measuring for 90 second.
164
165 **Statistical Analysis** All data were analyzed using PRISM (GraphPad Prism 9.0.0(121)).
166 Nonlinear regression (curve fit): polynomial (fourth order) is used to generate fitting results of RH
167 variation.
168
169 **Acknowledgments**
170

171 We wish to thank Prof. Baoqing Wang, Prof. Zhengxin Ying and associate Prof. Wei Wu for
172 suggestions and financial support. Figure 1 is created with BioRender.com

173
References

1. Enjin A*, et al.* (2016) Humidity Sensing in Drosophila. *Current biology : CB* 26(10):1352-
1358.
2. Russell J, Vidal-Gadea A, Makay A, Lanam C, & Pierce-Shimomura J (2014) Humidity
sensation requires both mechanosensory and thermosensory pathways in
Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United*
States of America 111.
3. Filingeri D, Fournet D, Hodder S, & Havenith G (2014) Why wet feels wet? A
neurophysiological model of human cutaneous wetness sensitivity. *Journal of*
Neurophysiology 112:1457.
4. Shettigar N*, et al.* (2017) Hierarchies in light sensing and dynamic interactions between
ocular and extraocular sensory networks in a flatworm. *Science Advances* 3:e1603025.
5. Inoue T, Hoshino H, Yamashita T, Shimoyama S, & Agata K (2015) Planarian shows
decision-making behavior in response to multiple stimuli by integrative brain function.
Zoological Letters 1(1):7.
6. Allen GD (1915) Reversibility of the Reactions of Planaria Dorotocephala to a Current of
Water. *Biological Bulletin* 29(2):111-128.
7. Mason P (1975) Chemo-klino-kinesis in planarian food location. *Animal behaviour*
23:460-469.
8. Dessì-Fulgheri F & Messeri P (1973) [Use of 2 different negative reinforcements in light-
darkness discrimination of planarians]. *Bollettino della Società italiana di biologia*
sperimentale 49:1141-1145.
9. Brown F & Chow C (1975) Differentiation between Clockwise and Counterclockwise
Magnetic Rotation by the Planarian, Dugesia dorotacephala. *Physiological Zoology*
48:168-176.

10.   Brown H & Ogden T (1968) The Electrical Response of the Planarian Ocellus. *The Journal*
200   *of general physiology* 51:237-253.
201   11.   Vila-Farré M & Rink J (2018) The Ecology of Freshwater Planarians.), Vol 1774, pp 173-
202   205.
203   12.   Tichy H & Kallina W (2010) Insect Hygroreceptor Responses to Continuous Changes in
204   Humidity and Air Pressure. *Journal of neurophysiology* 103:3274-3286.
205   13.   Liu L*, et al.* (2007) Drosophila hygrosensation requires the TRP channels water witch and
206   nanchung. *Nature* 450:294-298.
207   14.   Merryman S, Sánchez Alvarado A, & Jenkin J (2018) Culturing Planarians in the
208   Laboratory.), Vol 1774, pp 241-258.

209
210

211
212 **Figures**

213
214 **Figure 1.** Illustration of the experiment process. (A-D) The diving experiment. (E) The relative
215 humidity measurement.
216
217 <insert page break here>
218
219

220
Figure 2. Diving behavior are induced by rapidly increasing humidity. 3 sets of RH data are used
for nonlinear curve fitting in (A-E) Nonlinear regression (curve fit): polynomial (fourth order) is
used to generate fitting results of RH variation. The lower panel of (A-E) shows that the time a
worm starts the diving behavior synchronized with the RH variation, time data is presented as
mean ± SEM. (A) Rapidly increasing humidity induces diving behavior of planarians (n=20). (B)

226 Constant RH at around 38% cannot induce diving behavior of planarians(n=20). (C) Constant RH
227 at around 65% cannot induce diving behavior of planarians(n=20). (D)Slower increasing humidity
228 hinders planarians' decision to dive (n=20, 6 worms did not dive, 2 worms used more than 90
229 seconds). (E) Rapidly increasing humidity can mislead planarians drop into a dry place (n=20).
230 (F) The percentage of worms dived in each group.
231