A meta-analysis of neural systems underlying delay discounting: implications for transdiagnostic

research

Min Kang Souther[1]*

Bema Boateng[1]

Joseph W. Kable[1]

[1] Department of Psychology, University of Pennsylvania

*Corresponding author:
Min Kang Souther, MA
Department of Psychology
University of Pennsylvania
3700 Hamilton Walk
Philadelphia, PA 19104
minskang@sas.upenn.edu

ABSTRACT

The National Institute of Mental Health launched the Research Domain Criteria (RDoC) initiative to identify common biological mechanisms underlying mental disorders. A central challenge for this approach is to identify reliable biobehavioral paradigms that engage specific neural circuits implicated across psychiatric disorders. Delay discounting paradigms have been proposed as a promising candidate because both excessive and insufficient tendency to discount future rewards have been reported across a wide range of diagnoses. Because delay discounting involves multiple facets of neurocognitive functions, identifying specific neural systems engaged during the task can inform ongoing efforts in clinical neuroscience to link neural patterns to a core symptom dimension. To this end, we conducted a meta-analysis of 72 fMRI studies of delay discounting by common analytic approaches (contrasts) to test the predictions of commonly proposed mechanisms underlying the decision-making process. Despite being widely used in previous studies, comparing impulsive and patient choices did not reveal any reliable clustering of effects. Our results instead revealed that the subjective value, magnitude, and delay contrasts reliably engage the valuation network, with the subjective value contrast eliciting more widespread activation. Furthermore, both the difficulty and task contrasts engage the frontoparietal network. Taken together, our findings suggest that both reward valuation and action selection are important but do not fully explain impulsive versus patient choices. Taken together, these results provide further support that delay discounting task is a promising measure of reward valuation that reliably activates the valuation network. In addition, we discuss implications for analytic strategies for future studies in clinical neuroscience.

In 2009, the National Institute of Mental Health launched the Research Domain Criteria (RDoC) initiative to identify common biological mechanisms for understanding mental disorders[1]. After over a decade of research, reliability remains a central challenge for identifying biobehavioral measures that provide novel insights into the core neurocognitive features exhibited across disorders[2]. Delay discounting, or the tendency to devalue rewards that are distant in the future, has been proposed as a promising candidate paradigm for shared reward-related changes across diagnostic populations[3]. Currently categorized as a subconstruct of reward valuation under positive valence systems, delay discounting has been implicated in a wide range of psychiatric disorders, as both excessive discounting (e.g., substance use disorder, depression, schizophrenia, binge eating disorder) and insufficient discounting (e.g., anorexia) can be highly maladaptive[4,5]. In addition to this dimensional feature that cuts across diagnoses, delay discounting has been shown to be a strong prognostic indicator of treatment response and relapse[6,7], and can be modified using pharmacological[8–10], neural[11], and behavioral[12,13] interventions, which demonstrates its potential for clinical utility. Coupled with its high test-retest reliability[14] and evidence for genetic basis[15,16], delay discounting indeed has many appealing features as a transdiagnostic construct. Moreover, delay discounting is thought to engage multiple key neurocognitive processes that are relevant to many psychiatric disorders, such as impulsivity, self-regulation, and valuation[17]. Given the multifaceted nature of neurocognitive processes likely engaged in this task, previous research has implemented a wide range of analytic approaches to neuroimaging analysis. Therefore, systematically evaluating the reliability of neural systems hypothesized to underlie these facets can provide important insights for future transdiagnostic research.

A prominent hypothesis for neural mechanisms underlying maladaptive decision-making across disorders is the Competing Neurobehavioral Decision Systems (CNDS) model[17]. This account draws upon one of the first neuroimaging studies of delay discounting that identified two separate neural systems in the healthy brain, one for impulsive, emotional choices that biases toward immediate reward and another that rationally weighs both options[18]. Therefore, the CNDS model posits that impulsive choices are driven by heightened sensitivity to bottom-up reward signals in the limbic areas in response to rewards that are available sooner. On the other hand, hyperactivation of the top-down inhibitory control signals in the frontal cognitive regions can override this sensitivity, leading to more patient choices. Taken together, both excessive and insufficient discounting can arise when competition between the bottom-up reward signals and top-down inhibitory signals is not balanced. Indeed, evidence consistent with this model has been reported across disorders. For one, patients with anorexia nervosa showed hyperactivations in the dorsal cognitive regions, whereas those with bulimia nervosa showed reduced activations during reward processing[19]. This framework has become instrumental for subsequent clinical neuroscience research, as numerous studies have compared neural activity by choice to dissociate the systems underlying impulsivity and self-regulation as a feature of various psychiatric disorders.

Another prominent framework for the neural systems underlying discounting behavior is the single valuation model[20]. Contrary to the CNDS, this hypothesis posits that, instead of two competing neural systems, there is a single system that tracks the subjective value of both options and people make choices that maximize subjective value[20,21]. Subjective value is an estimated value of the future reward after depreciation, which accounts for the objective amount (magnitude), devaluation from the waiting time (delay), and individual differences in one's

preference for the sooner option (discount rate). The rationale is that, within the same neural system, activation increases in response to greater reward amount and simultaneously decreases as the reward becomes more distant in the future. Under this framework, excessive discounting can be driven by exaggerated response to reward amount, heightened aversion to delay, or both. Conversely, insufficient discounting can be driven by a combination of dampened response to reward and reduced sensitivity to delay. This account is consistent with previous meta-analytic work which found that the same regions within the valuation network, consisting of ventral striatum and ventromedial prefrontal cortex (vmPFC), encode both positive and negative effects of subjective value[22]. Disruptions in value signals in either direction can be maladaptive. Reduced value response has been reported across disorders characterized by anhedonia or reduced motivation[23,24], whereas pathological gambling has been linked to exaggerated value signals[25]. Although the single valuation model has been widely examined in decision neuroscience, relatively fewer clinical neuroimaging studies of delay discounting have taken this approach[25,26].

Lastly, an emerging line of research in computational neuroscience calls for a third hypothesis, termed the evidence accumulation model[27]. This model draws upon previous findings of frontoparietal control activations during decision-making that do not require impulse control, such as perceptual judgment tasks[28,29]. This framework is complementary to the single valuation model but speaks to the post-valuation process, in which subjective values are evaluated and translated into decisions. This value accumulation and response selection process is believed to engage activations in the frontoparietal control areas including the dorsolateral prefrontal cortex (dlPFC), with increased activity when two options are highly conflicting. Applying this logic to delay discounting specifically, rewards that are similar in subjective value

are considered to be more conflicting and difficult, and thus would elicit greater activations in these regions. Indeed, increased frontoparietal activations as options become closer in value have been reported in delay discounting[30] and attenuated response to difficult decisions have been reported in patients with methamphetamine dependence[31].

In this study, we performed a series of meta-analyses of fMRI findings derived from analytic strategies grounded in the three prominent neurocognitive models of delay discounting with the goal of identifying the neural systems that are reliable and robust to individual study designs, including the presence of psychiatric diagnoses. Each model makes different predictions about the expected neural patterns. The CNDS model would predict greater fronto-striatal activations when presented with an immediate option ("immediacy bias") and greater frontoparietal activations for difficult decisions. Furthermore, the former regions would be more active for impulsive choices and the latter regions would be more active for patient choices. On the other hand, the single valuation model would predict activations in the valuation network as a function of subjective value, magnitude, and delay, and that these activations would occur in overlapping regions. An extension of this account by the value accumulation model would predict increased frontoparietal activations as a function of choice difficulty, which is consistent with the prediction made by the CNDS model. For each of these analytic contrasts, we conducted a multi-level kernel density analysis to quantify the proportion of studies that report consistent effects in a given anatomical location. We further tested the predictions of the three neurocognitive models by identifying unique and overlapping regions across analytic contrasts. We hope this work will shed light on the core neurocognitive processes underlying delay discounting behavior and contribute to building a strong foundation for future transdiagnostic research in clinical neuroscience.

Methods

Search strategy. Preregistration of the present meta-analysis is available on PROSPERO under protocol CRD42020150691. We conducted a systematic literature search on PubMed and PsycInfo (final search on February 2, 2021). Based on previous meta-analyses of delay discounting[4,32], we used the following search terms, which yielded 392 unique results:

> ((Delay discounting) OR (Temporal discounting) OR (Delay of gratification) OR
> (Intertemporal choice) OR (time discounting)) AND (fMRI OR functional magnetic
> resonance imaging)

We also reviewed the reference lists of previously published review articles and searches by frequently occurring authors (i.e., 10 or more unique articles in the initial search results), which identified 2 additional articles.

Study selection. All articles identified through the above search strategy were evaluated according to the preregistered inclusion/exclusion criteria. For inclusion, the study had to meet the following criteria: (1) English or an English translation is available; (2) human participants with no brain lesions; (3) fMRI data collected while the participants were completing a delay discounting task in the scanner; (4) the task included a delay of at least 1 day and used monetary rewards, regardless of payment contingency; (5) conducted a parametric or categorical contrast analysis across the whole brain; (6) spatial coordinates are available in Talairach or Montreal Neurological Institute (MNI) space, either reported in the manuscript or provided by the corresponding author upon request; and (7) diagnosis of psychiatric disorders can be present. Although existing meta-analyses of self-regulatory decision-making tend to focus on healthy adults or group comparisons[33–35], inclusion of individuals with psychiatric disorders achieves two goals. First, it allows greater variability of the clinical or cognitive constructs in question

(impulsivity, self-regulation, valuation, conflict monitoring, and action planning) and others that may be related to delay discounting. Second, given our ultimate aim to understand specific neural systems engaged during the task to inform its implications for transdiagnostic processes, inclusion of psychopathology without regard to specific diagnoses seemed critical. We excluded studies that (1) were not a primary source; (2) used non-monetary rewards; (3) had pharmacological manipulation unless they report effects in the placebo group; (4) resting-state or volumetric analyses; (5) connectivity and multi-voxel pattern analyses; (6) overlapping samples with another included study. In the event of overlapping samples, we included the largest sample. To ensure inclusion criteria are standard and to limit bias, all studies were reviewed using Rayyan (https://rayyan.qcri.org) by at least two independent reviewers (MSK and JWK, MSK and BB, or all three). The inter-rater reliability was high ($\kappa = 0.86$; 93.9% agreement) and all disagreements were discussed and resolved by a consensus decision. A total of 74 studies (1262 foci; 2911 participants) across were included in the quantitative meta-analysis (see Figure 1 for the PRISMA flow diagram; Table 1 for the list of studies and contrasts).

Coding contrasts. Following study selection, we coded the studies into the following contrast categories: general task-related activity (*Task*), choices involving an immediate reward (*Now*), choosing the smaller-sooner reward (*SSR*), choosing the larger-later reward (*LLR*), difficult decisions (*Hard*), easy decisions (*Easy*), subjective value (*SV*), objective amount (*Magnitude*), and length of delay (*Delay*). For *Task*, we included positive effects of comparing the delay discounting task to a fixation or implicit baseline or a control task (e.g., choose left/right, choose larger). We also categorized comparing a subset of the task to baseline (e.g., Hard > baseline) as *Task*. For *Now*, we included comparing trials involving an immediate reward to either all trials or trials involving delayed options only. *SSR* included direct comparisons

between the two choices (positive effects of SSR > LLR or negative effects of LLR > SSR) and *LLR* included the converse. As described earlier, trials in which the two options are closer in subjective value or near the indifference point are considered to be difficult, and studies categorize trials as hard or easy based on the value distance or distance from the indifference point. Therefore, *Hard* included direct comparisons between hard and easy trials (positive effects of Hard > Easy or negative effects of Easy > Hard) whereas *Easy* included the opposite effects. We also coded negative correlations with subjective value difference as *Hard* and positive correlations as *Easy*. For *SV*, we included brain regions showing positive correlation with the discounted value of the delayed reward. Although SV is frequently calculated using hyperbolic discounting[36], we deferred to the judgment of the original authors and reviewers and included all discounted value regressors regardless of how SV was calculated. For *Magnitude*, we included positive correlations with the objective amount of either the LLR or SSR, although all studies in our meta-analytic sample examined the amount of the delayed option except for one study[37], which examined varying amounts of the immediate option while keeping the amount of delayed reward constant throughout the task. For *Delay*, we included brain regions showing reduced activations as the delay increases (i.e., the negative effects of delay or positive effects of inverse delay). For studies that reported the peaks by comparison groups (e.g., young and older adults, healthy and psychopathology), we included them as separate studies rather than selecting a single group. After coding, there were 25 studies (250 foci) in *Task*, 8 (78 foci) in *Now*, 15 (95 foci) in *SSR*, 22 (171 foci) in *LLR*, 16 (112 foci) in *Hard*, 12 (95 foci) in *Easy*, 22 (263 foci) in *SV*, 9 (82 foci) in *Magnitude* and 9 (116 foci) in *Delay*.

Multilevel kernel density analysis. Multilevel kernel density analysis (MKDA) is a widely used technique for coordinate-based meta-analysis to quantify the proportion of studies that

report activation foci within a given radius, with the null hypothesis that foci are located randomly throughout the brain[22,38]. Unlike the traditional meta-analysis that examines the magnitude of effect sizes, here we aimed to identify anatomical locations where an effect is observed consistently across studies that used the same contrast. A previous meta-analysis from our group describes in detail the rationale for refining the MKDA procedure used in the present study[22]. We first converted all peak coordinates from each study into the MNI space using mni2tal to create a binary indicator map (IM)—for interpretability, we assigned 100 for voxels within 10mm of the peaks and 0 for others. We then averaged these maps across studies, resulting in the mean IM ranging from 0-100 that can be interpreted as the percentage of studies reporting a peak within a 10mm radius. For significance testing, we implemented permutation testing in Matlab by simulating a null-hypothesis that foci are randomly distributed, while maintaining the number of studies and number of foci per study. We also applied a cluster-forming threshold for each contrast to improve sensitivity. For each study, we first accounted for the probability of gray matter, where there is a higher probability of observing a positive IM value, by calculating the proportion of voxels with a positive IM value in the top quintile, as indicated by the ICBM tissue probability map (www.loni.ucla.edu/ICBM). These proportions were then treated as independent probabilities to derive the mean IM value that would be exceeded with a probability of less than 0.01 by chance alone. We applied this threshold to the results of each random permutation and recorded the largest cluster mass (sum of all IM values in a given cluster) in the null distribution. We also applied the same threshold to the real data, and the resulting clusters were assigned corrected p-values based on the frequency of equal or larger clusters in the null distribution. All reported results are based on 200 iterations of

permutations (one of which was the observed data) and a threshold of corrected p = 0.05 (one-tailed).

Spatial overlap. Lastly, a series of overlap analyses were performed using the maps obtained from the density analyses described above to reveal activity in unique and shared brain areas. Suprathreshold cluster mass were binarized and summed to create a conjunction map, which was generated and visualized in FSL. The centroid locations for overlapping clusters were identified and the number of contrasts that found significant clustering in a given voxel was examined.

Results

Individual contrasts. Significant clusters for each contrast are presented in Figure 2 and Table 2. We first examined general task-related activations (*Task*), which were significantly clustered in the superior frontal gyrus, inferior frontal gyrus, frontal pole, intraparietal sulcus, and insula. We also observed a cluster in the occipital pole, likely an effect of visual input from the task. We then examined specific contrasts, guided by different neurocognitive models of delay discounting. As previously described, the CNDS model predicts greater activations in the reward regions for choices involving an immediate option (*Now*) and choosing the smaller-sooner reward (*SSR*), as well as distinct, greater frontoparietal activations for difficult choices (*Hard*) and choosing the larger-later reward (*LLR*). Contrary to these predictions, we also observed no areas of significant clustering for *Now* and *SSR*, although the former null finding should be interpreted with caution as there were only 8 studies (78 foci) in this meta-analysis. For *LLR*, significant clustering occurred in the left precentral gyrus. Given this localization in motor cortex, we examined the studies that contributed to this cluster for motor confounds and found that they all instructed participants to choose the LLR using their right middle finger (and

SSR with the right index finger)[39–43]. Both the CNDS and anDDM models predict greater frontoparietal activation for difficult decisions (*Hard*). Significant clusters were found in the dorsomedial prefrontal cortex (dmPFC) and supramarginal gyrus for *Hard*, and in the posterior cingulate cortex (PCC) for *Easy* (i.e., negative effects of choice difficulty). The unitary valuation model predicts activations in the valuation network across *SV*, *Magnitude*, and *Delay* in overlapping regions. The highest meta-analytic statistics (i.e., percentage of studies reporting a peak within 10mm from a given voxel) were found in the striatum for *SV* (63.6%). Additionally, we observed significant clusters in the vmPFC, PCC, and middle temporal gyrus. Foci *Magnitude* were also clustered in the striatum. For negative effects of *Delay*, significant clustering was similarly found in the striatum, as well as medial prefrontal cortex, inferior frontal gyrus, and middle frontal gyrus.

Spatial overlap. We conducted an overlap analysis to identify unique and overlapping neural systems according to different neurocognitive models of delay discounting (Figure 3). Although the CNDS model would predict an overlap between *Now* and *SSR*, and between *Hard* and *LLR*, we did not conduct this follow up analysis because there were no significant clusters in the individual contrasts for *Now*, *SSR*, and *LLR* (other than motor activity). The single valuation model would predict an overlap among *SV*, *Magnitude*, and *Delay*. Consistent with this account, there was a considerable overlap among the clusters in the striatum, although there were no overlapping voxels shared by all three contrasts. The anDDM model would predict activations in the frontoparietal regions distinct from the valuation system. Contrary to this prediction, we found an overlap in the PCC for *SV* and *Easy*. We also explored overlapping clusters between task-related activations and individual contrasts, and found overlapping voxels between *Task* and

Hard in the medial prefrontal cortex, and between *Task* and *Delay* in the dmPFC and the right inferior frontal gyrus.

Discussion

In this paper, we report a series of meta-analyses examining neural effects predicted by existing models of delay discounting. Our first step was to identify general task-related activations, which revealed reliable clusters corresponding with the frontoparietal task control network[44]. We then tested the predictions of the CNDS model, single valuation model, and anDDM by examining individual contrasts. Contrary to the CNDS model, our meta-analysis did not reveal reliable effects of choice (impulsive or patient) at the neural level. Similarly, our results did not find neural effects of an immediacy bias, although this may be in part due to the small number of studies included in our analysis (n=8). Consistent with both the CNDS model and anDDM, our findings revealed significant effects of choice difficulty in the dmPFC, although we did not find clusters in the dlPFC as both models predict. Conversely, easy choices engaged the PCC, a node in the task-negative or default mode network[44], suggesting reduced cognitive engagement when the options have highly distinct values. In line with the unitary valuation model, the most reliable clusters were found throughout the valuation network as a function of subjective value, magnitude, and delay. Although we saw a more widespread engagement of the valuation network for subjective value than magnitude or delay alone, this comparison should be interpreted with caution given the relatively small sample sizes for magnitude and delay. Taken together, these results offer meta-analytic support for the unitary valuation model and anDDM, suggesting that both valuation and value accumulation processes are important for delay discounting. Although our findings may seem to contradict the CNDS model, the initial fMRI study that provided a foundation for this model in fact agrees that regions

activated by the task and value are distinct[18]. However, the interpretation that these networks are differentially associated with impulsive or patient choices is not supported by our work.

Similarly to the results of individual contrasts, our spatial overlap analysis suggests strong support for the unitary valuation model, as subjective value, magnitude, and delay were encoded in overlapping clusters of the striatum. Although not specifically predicted by any of the models of delay discounting, as one might expect, both the task and hard decisions engage the dmPFC, suggesting that difficult decisions require more task engagement. Our exploratory results also revealed overlaps between subjective value and easy choices in the PCC, and between task and delay in the dorsomedial prefrontal cortex. Although this finding was unexpected, we speculate that choices involving either extremely high values or extreme delays are easy and require less task engagement, indicated by deactivation of the task-positive network. Previous a meta-analysis which found similarities in the valuation network and default mode network also speculates a link between valuation and deactivation of task-related activity because value computation inherently involves episodic memory and mental simulations[45].

Since we began this work, a similar meta-analysis comparing a subset of these contrasts in healthy adults was published[33]. Because we included studies across the lifespan and psychiatric diagnoses, we had a much larger meta-analytic sample and therefore increased power to identify more clusters. Despite the differences in both study selection and meta-analytic approach (MKDA vs. activation likelihood estimation[46]), results are largely consistent for the valuation system (*SV*, *Magnitude*), frontoparietal control system (*Hard*), as well as task-related activations. However, our results did not replicate significant clusters previously reported in the

vmPFC for the immediacy bias (*Now*), and inferior frontal gyrus and insula for patient choices (*LLR*). They did not examine the *Delay*, *Easy*, and *SSR* contrasts.

Our ultimate goal for examining reliable neural effects by specific contrasts was to inform future transdiagnostic research. A sizable number of studies included in our quantitative analysis used delay discounting to investigate a wide range of psychiatric disorders and related symptoms – such as substance use[31,41,47–50], pathological gambling[25,26,40,51], obsessive compulsive disorder[39,52], ADHD[39], schizophrenia[53], major depressive disorder[54], obesity[55,56], prediabetics[57], and poor sleepers[58]. Although there were not enough studies for a meaningful quantitative meta-analysis for any given contrast, our work identified neural systems that are reliable across studies and robust to experimental variations, including psychiatric diagnoses. Among the studies with a psychiatric sample included in our analyses, *Task* and *SSR* were the most frequently used contrasts (n=8 for both), whereas only two studies[25,26] examined the *SV* contrast and only one study[54] examined the effects of *Magnitude* and *Delay*. While comparing choices is a logical approach to target neural systems underlying impulsivity, the hypothesis that there are distinct systems associated with impulsive and patient choices is not supported by our work. Based on our findings that the unitary valuation model yielded the most reliable effects, it would be worthwhile for future clinical neuroimaging studies to examine differential effects of magnitude and delay to characterize excessive or insufficient discounting, rather than choices per se.

There are several limitations to consider when interpreting our findings. Although our results suggest neural evidence for value computation and accumulation, neither account can yet fully explain how they ultimately translate to individual choices. It would be important for future research to focus on how valuation and value accumulation jointly shape discounting. Also, we

did not have enough studies for a categorical comparison between healthy volunteers and individuals with psychiatric disorders. However, we determined that inclusion of both healthy and psychiatric populations would be more in line with the goals of RDoC and relevant for future studies adopting this framework. Relatedly, because of the emphasis on including psychiatric populations, we coded studies that reported the effects of contrasts separated by comparison groups as independent studies, rather than selecting a single set of results to be included. Because these studies are not independent, our results may contain a bias for such studies that may not be present in other meta-analyses. It is also important to note that most studies contributed more than a single contrast, and these contrasts tended to go hand in hand (e.g., SSR and LLR, magnitude and delay, hard and easy), which could pose a confound for the spatial overlap results. Our work exclusively focused on monetary rewards, but there are a number of behavioral studies with paradigms using clinically relevant rewards (e.g., cigarettes for smokers[59], food for obese individuals[60,61]). Although we are cautiously optimistic about the generalizability of the present study given the evidence supporting a common neural valuation system across commodities[62], it is likely that there are commodity-specific activations that are not captured in our results. Lastly, because we were estimating the location of effects rather than the size of effects, there was no statical penalty for reported null reports. To this end, it is worth nothing that a large number of studies that used the choice contrast (LLR, SSR) reported no significant clusters for this contrast[20,31,43,48,50,56,58,63–70], whereas only two studies reported no significant effects of SV[71,72]. This comparison further highlights the importance for identifying and implementing more reliable analytic strategies for future research.

The present meta-analysis offers several exciting directions for future research. As data archiving becomes more common in the field, future meta-analysis can move beyond coordinate-

based meta-analysis and conduct image-based meta-analysis[73]. Unlike the coordinate-based methods, which only include the peak voxels, image-based methods can include the whole statistical map, allowing researchers to detect consistent but relatively weaker effects and to calculate effect sizes, which is a limitation of coordinate-based methods that only examines effect locations. In order to make image-based approach possible, it would be crucial for researchers to make whole-brain statistical maps publicly available in established archives, such as NeuroVault (https://neurovault.org/). Future meta-analytic work in delay discounting might also aim to examine the link between these neural effects and individual differences, such as discount rates, cognitive functioning, and psychopathology. As more neuroimaging studies of delay discounting in clinical populations become available, it would be worthwhile to conduct a future meta-analysis, both within specific disorders and across psychopathology in line with the RDoC framework. A practical goal of our work was to identify an analytic strategy that is reliable and robust to experimental variations. Given the high reliability and interpretability of effects produced by the subjective value contrast, our recommendation for future work is to use this contrast in the absence of specific hypotheses that call for a different approach. Another, also practical, outcome of our work is the generation of cluster maps that can be used as *a priori* ROIs for future research. This will be especially valuable for clinical neuroimaging studies of delay discounting that might be underpowered to detect modest effects using whole-brain approaches. All results will be made available in nifti format for this purpose.

References

1. Insel TR. Translating Scientific Opportunity Into Public Health Impact: A Strategic Plan for Research on Mental Illness. *Arch Gen Psychiatry*. 2009;66(2):128-133. doi:10.1001/archgenpsychiatry.2008.540

2. MacNamara A, Phan KL. Psychobiological operationalization of RDoC constructs: Methodological and conceptual opportunities and challenges. *Psychophysiology*. 2016;53(3):406-409. doi:https://doi.org/10.1111/psyp.12587

3. Lempert KM, Steinglass JE, Pinto A, Kable JW. Can delay discounting deliver on the promise of RDoC? *Psychol Med*. 2019;49(2):190-199. doi:10.1017/S0033291718001770

4. Amlung M, Marsden E, Holshausen K, et al. Delay Discounting as a Transdiagnostic Process in Psychiatric Disorders: A Meta-analysis. *JAMA Psychiatry*. Published online August 28, 2019. doi:10.1001/jamapsychiatry.2019.2102

5. Amlung M, Vedelago L, Acker J, Balodis I, MacKillop J. Steep delay discounting and addictive behavior: a meta-analysis of continuous associations. *Addict Abingdon Engl*. 2017;112(1):51-62. doi:10.1111/add.13535

6. Washio Y, Higgins ST, Heil SH, et al. Delay discounting is associated with treatment response among cocaine-dependent outpatients. *Exp Clin Psychopharmacol*. 2011;19(3):243-248. doi:10.1037/a0023617

7. Sheffer CE, Christensen DR, Landes R, Carter LP, Jackson L, Bickel WK. Delay discounting rates: A strong prognostic indicator of smoking relapse. *Addict Behav*. 2014;39(11):1682-1689. doi:10.1016/j.addbeh.2014.04.019

8. Orsini CA, Mitchell MR, Heshmati SC, et al. Effects of nucleus accumbens amphetamine administration on performance in a delay discounting task. *Behav Brain Res*. 2017;321:130-136. doi:10.1016/j.bbr.2017.01.001

9. de Wit H, Mitchell SH. Drug effects on delay discounting. In: *Impulsivity: The Behavioral and Neurological Science of Discounting*. American Psychological Association; 2010:213-241. doi:10.1037/12069-008

10. Demoto Y, Okada G, Okamoto Y, et al. Neural and personality correlates of individual differences related to the effects of acute tryptophan depletion on future reward evaluation. *Neuropsychobiology*. 2012;65(2):55-64. doi:10.1159/000328990

11. Hecht D, Walsh V, Lavidor M. Bi-frontal direct current stimulation affects delay discounting choices. *Cogn Neurosci*. 2013;4(1):7-11. doi:10.1080/17588928.2011.638139

12. Ciaramelli E, Sellitto M, Tosarelli G, di Pellegrino G. Imagining Events Alternative to the Present Can Attenuate Delay Discounting. *Front Behav Neurosci*. 2019;13. doi:10.3389/fnbeh.2019.00269

13. Lempert KM, Speer ME, Delgado MR, Phelps EA. Positive autobiographical memory retrieval reduces temporal discounting. *Soc Cogn Affect Neurosci*. 2017;12(10):1584-1593. doi:10.1093/scan/nsx086

14. Enkavi AZ, Eisenberg IW, Bissett PG, et al. Large-scale analysis of test–retest reliabilities of self-regulation measures. *Proc Natl Acad Sci*. 2019;116(12):5472-5477. doi:10.1073/pnas.1818430116

15. Gray JC, Sanchez-Roige S, de Wit H, MacKillop J, Palmer AA. Genomic basis of delayed reward discounting. *Behav Processes*. 2019;162:157-161. doi:10.1016/j.beproc.2019.03.006

16. Anokhin AP, Grant JD, Mulligan RC, Heath AC. The Genetics of Impulsivity: Evidence for the Heritability of Delay Discounting. *Biol Psychiatry*. 2015;77(10):887-894. doi:10.1016/j.biopsych.2014.10.022

17. Bickel WK, Athamneh LN, Basso JC, et al. Excessive discounting of delayed reinforcers as a trans-disease process: Update on the state of the science. *Curr Opin Psychol*. 2019;30:59-64. doi:10.1016/j.copsyc.2019.01.005

18. McClure SM, Laibson DI, Loewenstein G, Cohen JD. Separate Neural Systems Value Immediate and Delayed Monetary Rewards. *Science*. 2004;306(5695):503-507. doi:10.1126/science.1100907

19. Kaye WH, Wagner A, Fudge JL, Paulus M. Neurocircuity of Eating Disorders. In: Adan RAH, Kaye WH, eds. *Behavioral Neurobiology of Eating Disorders*. Current Topics in Behavioral Neurosciences. Springer; 2011:37-57. doi:10.1007/7854_2010_85

20. Kable JW, Glimcher PW. An "as soon as possible" effect in human intertemporal decision making: behavioral evidence and neural mechanisms. *J Neurophysiol*. 2010;103(5):2513-2531. doi:10.1152/jn.00177.2009

21. Kable JW, Glimcher PW. The neural correlates of subjective value during intertemporal choice. *Nat Neurosci*. 2007;10(12):1625-1633. doi:10.1038/nn2007

22. Bartra O, McGuire JT, Kable JW. The valuation system: A coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *NeuroImage*. 2013;76:412-427. doi:10.1016/j.neuroimage.2013.02.063

23. Kang MS, Wolf D, Kazinka R, et al. Decision value signals in the ventromedial prefrontal cortex and anhedonia across mood and psychotic disorders. *bioRxiv*. Published online December 2, 2020:2020.12.01.407197. doi:10.1101/2020.12.01.407197

24. Brown VM, Wilson J, Hallquist MN, Szanto K, Dombrovski AY. Ventromedial prefrontal value signals and functional connectivity during decision-making in suicidal behavior and impulsivity. *Neuropsychopharmacology*. 2020;45(6):1034-1041. doi:10.1038/s41386-020-0632-0

25. Miedl SF, Peters J, Büchel C. Altered neural reward representations in pathological gamblers revealed by delay and probability discounting. *Arch Gen Psychiatry*. 2012;69(2):177-186.

26. Wiehler A, Petzschner FH, Stephan KE, Peters J. Episodic Tags Enhance Striatal Valuation Signals during Temporal Discounting in pathological Gamblers. *eNeuro*. 2017;4(3). Accessed January 1, 5AD. https://pubmed.ncbi.nlm.nih.gov/28612049/

27. Hutcherson CA, Tusche A. Evidence accumulation, not "self-control," explains dorsolateral prefrontal activation during normative choice. *bioRxiv*. Published online January 20, 2021:2020.10.06.328476. doi:10.1101/2020.10.06.328476

28. Pedersen ML, Endestad T, Biele G. Evidence Accumulation and Choice Maintenance Are Dissociated in Human Perceptual Decision Making. *PLOS ONE*. 2015;10(10):e0140361. doi:10.1371/journal.pone.0140361

29. Noppeney U, Ostwald D, Werner S. Perceptual decisions formed by accumulation of audiovisual evidence in prefrontal cortex. *J Neurosci Off J Soc Neurosci*. 2010;30(21):7434-7446. doi:10.1523/JNEUROSCI.0455-10.2010

30. Rodriguez CA, Turner BM, Van Zandt T, McClure SM. The neural basis of value accumulation in intertemporal choice. *Eur J Neurosci*. 2015;42(5):2179-2189.

31. Monterosso JR, Ainslie G, Xu J, Cordova X, Domier CP, London ED. Frontoparietal cortical activity of methamphetamine-dependent and comparison subjects performing a delay discounting task. *Hum Brain Mapp*. 2007;28(5):383-393.

32. Wesley MJ, Bickel WK. Remember the future II: Meta-analyses and functional overlap of working memory and delay discounting. *Biol Psychiatry*. 2014;75(6):435-448. doi:10.1016/j.biopsych.2013.08.008

33. Schüller CB, Kuhn J, Jessen F, Hu X. Neuronal correlates of delay discounting in healthy subjects and its implication for addiction: an ALE meta-analysis study. *Am J Drug Alcohol Abuse*. 2019;45(1):51-66. doi:10.1080/00952990.2018.1557675

34. Han JE, Boachie N, Garcia-Garcia I, Michaud A, Dagher A. Neural correlates of dietary self-control in healthy adults: A meta-analysis of functional brain imaging studies. *Physiol Behav*. 2018;192:98-108. doi:10.1016/j.physbeh.2018.02.037

35. Yuan P, Raz N. Prefrontal cortex and executive functions in healthy adults: A meta-analysis of structural neuroimaging studies. *Neurosci Biobehav Rev*. 2014;42:180-192. doi:10.1016/j.neubiorev.2014.02.005

36. Commons ML, Mazur JE, Nevin JA, Rachlin H. *The Effect of Delay and of Intervening Events on Reinforcement Value: Quantitative Analyses of Behavior*. Psychology Press; 2013.

37. Albert WD. Neurodevelopmental Substrates of Peer Influences on Adolescents' Choice Evaluation and Decision Making. Published online 2011. Accessed May 5, 2021. https://scholarshare.temple.edu/handle/20.500.12613/660

38. Wager TD, Lindquist MA, Nichols TE, Kober H, Van Snellenberg JX. Evaluating the consistency and specificity of neuroimaging data using meta-analysis. *Neuroimage*. 2009;45(1 Suppl):S210-S221. doi:10.1016/j.neuroimage.2008.10.061

39. Norman LJ, Carlisi CO, Christakou A, et al. Neural dysfunction during temporal discounting in paediatric Attention- Deficit/Hyperactivity Disorder and Obsessive-Compulsive Disorder. *Psychiatry Res Neuroimaging*. 2017;269:97-105. doi:10.1016/j.pscychresns.2017.09.008

40. Miedl SF, Wiswede D, Marco-Pallarés J, et al. The neural basis of impulsive discounting in pathological gamblers. *Brain Imaging Behav*. 2015;9(4):887-898. doi:10.1007/s11682-015-9352-1

41. Claus ED, Kiehl KA, Hutchison KE. Neural and behavioral mechanisms of impulsive choice in alcohol use disorder. *Alcohol Clin Exp Res*. 2011;35(7):1209-1219.

42. Christakou A, Brammer M, Rubia K. Maturation of limbic corticostriatal activation and connectivity associated with developmental changes in temporal discounting. *NeuroImage*. 2011;54(2):1344-1354.

43. Waegeman A, Declerck CH, Boone C, Van Hecke W, Parizel PM. Individual differences in self-control in a time discounting task: An fMRI study. *J Neurosci Psychol Econ*. 2014;7(2):65-79. doi:10.1037/npe0000018

44. Power JD, Cohen AL, Nelson SM, et al. Functional network organization of the human brain. *Neuron*. 2011;72(4):665-678. doi:10.1016/j.neuron.2011.09.006

45. Acikalin MY, Gorgolewski KJ, Poldrack RA. A Coordinate-Based Meta-Analysis of Overlaps in Regional Specialization and Functional Connectivity across Subjective Value and Default Mode Networks. *Front Neurosci*. 2017;11. doi:10.3389/fnins.2017.00001

46. Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT. Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: a random-effects approach based on empirical estimates of spatial uncertainty. *Hum Brain Mapp*. 2009;30(9):2907-2926. doi:10.1002/hbm.20718

47. Hoffman WF, Schwartz DL, Huckans MS, et al. Cortical activation during delay discounting in abstinent methamphetamine dependent individuals. *Psychopharmacology (Berl)*. 2008;201(2):183-193.

48. Lim AC, Cservenka A, Ray LA. Effects of Alcohol Dependence Severity on Neural Correlates of Delay Discounting. *Alcohol Alcohol Oxf Oxfs*. 2017;52(4):506-515.

49. Meade CS, Lowen SB, MacLean RR, Key MD, Lukas SE. fMRI brain activation during a delay discounting task in HIV-positive adults with and without cocaine dependence. *Psychiatry Res Neuroimaging*. 2011;192(3):167-175. doi:10.1016/j.pscychresns.2010.12.011

50. Stanger C, Elton A, Ryan SR, James GA, Budney AJ, Kilts CD. Neuroeconomics and adolescent substance abuse: individual differences in neural networks and delay discounting. *J Am Acad Child Adolesc Psychiatry*. 2013;52(7):747-755.e6.

51. Wang Y, Hu Y, Xu J, et al. Dysfunctional prefrontal function is associated with impulsivity in people with internet gaming disorder during a delay discounting task. *Front Psychiatry*. 2017;8:10. doi:10.3389/fpsyt.2017.00287

52. Mavrogiorgou P, Enzi B, Steinmann S, Mulert C, Juckel G. Relationship Between Neuroanatomical and Serotonergic Hypotheses of Obsessive-Compulsive Disorder: A Combined Functional Magnetic Resonance Imaging-Evoked Potential Study. *J Clin Psychiatry*. 2018;79(6). Accessed January 1, 10AD. https://pubmed.ncbi.nlm.nih.gov/30326190/

53. Avsar KB, Weller RE, Cox JE, Reid MA, White DM, Lahti AC. An fMRI investigation of delay discounting in patients with schizophrenia. *Brain Behav*. 2013;3(4):384-401. doi:10.1002/brb3.135

54. Vanyukov PM, Szanto K, Hallquist MN, et al. Paralimbic and lateral prefrontal encoding of reward value during intertemporal choice in attempted suicide. *Psychol Med*. 2016;46(2):381-391. doi:10.1017/S0033291715001890

55. Kishinevsky FI, Cox JE, Murdaugh DL, Stoeckel LE, Cook EW III, Weller RE. fMRI reactivity on a delay discounting task predicts weight gain in obese women. *Appetite*. 2012;58(2):582-592. doi:10.1016/j.appet.2011.11.029

56. Morys F, Bode S, Horstmann A. Dorsolateral and medial prefrontal cortex mediate the influence of incidental priming on economic decision making in obesity. *Sci Rep*. 2018;8(1):17595.

57. Deshpande HU, Mellis AM, Lisinski JM, et al. Reinforcer pathology: Common neural substrates for delay discounting and snack purchasing in prediabetics. *Brain Cogn*. 2019;132:80-88.

58. Martin LE, Pollack L, McCune A, Schulte E, Savage CR, Lundgren JD. Comparison of obese adults with poor versus good sleep quality during a functional neuroimaging delay discounting task: A pilot study. *Psychiatry Res Neuroimaging*. 2015;234(1):90-95. doi:10.1016/j.pscychresns.2015.08.011

59. MacKillop J, Amlung MT, Wier LM, et al. The neuroeconomics of nicotine dependence: a preliminary functional magnetic resonance imaging study of delay discounting of monetary and cigarette rewards in smokers. *Psychiatry Res*. 2012;202(1):20-29.

60. Weygandt M, Spranger J, Leupelt V, et al. Interactions between neural decision-making circuits predict long-term dietary treatment success in obesity. *NeuroImage*. 2019;184:520-534. doi:10.1016/j.neuroimage.2018.09.058

61. Weygandt M, Mai K, Dommes E, et al. Impulse control in the dorsolateral prefrontal cortex counteracts post-diet weight regain in obesity. *NeuroImage*. 2015;109:318-327. doi:10.1016/j.neuroimage.2014.12.073

62. Levy DJ, Glimcher PW. The root of all value: a neural common currency for choice. *Curr Opin Neurobiol*. 2012;22(6):1027-1038. doi:10.1016/j.conb.2012.06.001

63. Schmaal L, Goudriaan AE, Joos L, et al. Neural substrates of impulsive decision making modulated by modafinil in alcohol-dependent patients. *Psychol Med*. 2014;44(13):2787-2798. doi:10.1017/S0033291714000312

64. Marco-Pallarés J, Mohammadi B, Samii A, Münte TF. Brain activations reflect individual discount rates in intertemporal choice. *Brain Res*. 2010;1320:123-129.

65. Kobiella A, Ripke S, Kroemer NB, et al. Acute and chronic nicotine effects on behaviour and brain activation during intertemporal decision making. *Addict Biol*. 2014;19(5):918-930.

66. Mavrogiorgou P, Enzi B, Klimm A, et al. Serotonergic modulation of orbitofrontal activity and its relevance for decision making and impulsivity. *Hum Brain Mapp*. 2017;38(3):1507-1517. doi:10.1002/hbm.23468

67. Hamilton KR, Smith JF, Gonçalves SF, et al. Striatal bases of temporal discounting in early adolescents. *Neuropsychologia*. 2020;144:107492.

68. Weber BJ, Huettel SA. The neural substrates of probabilistic and intertemporal decision making. *Brain Res*. 2008;1234:104-115. doi:10.1016/j.brainres.2008.07.105

69. van den Bos W, Rodriguez CA, Schweitzer JB, McClure SM. Adolescent impatience decreases with increased frontostriatal connectivity. *Proc Natl Acad Sci U S A*. 2015;112(29):E3765-74.

70. Wittmann M, Leland DS, Paulus MP. Time and decision making: differential contribution of the posterior insular cortex and the striatum during a delay discounting task. *Exp Brain Res*. 2007;179(4):643-653.

71. Sripada CS, Gonzalez R, Phan KL, Liberzon I. The neural correlates of intertemporal decision-making: contributions of subjective value, stimulus type, and trait impulsivity. *Hum Brain Mapp*. 2011;32(10):1637-1648.

72. Miedl SF, Büchel C, Peters J. Cue-induced craving increases impulsivity via changes in striatal value signals in problem gamblers. *J Neurosci*. 2014;34(13):4750-4755. doi:10.1523/JNEUROSCI.5020-13.2014

73. Salimi-Khorshidi G, Smith SM, Keltner JR, Wager TD, Nichols TE. Meta-analysis of neuroimaging data: A comparison of image-based and coordinate-based pooling of studies. *NeuroImage*. 2009;45(3):810-823. doi:10.1016/j.neuroimage.2008.12.039

74. Albrecht K, Volz KG, Sutter M, Laibson DI, von Cramon DY. What is for me is not for you: Brain correlates of intertemporal choice for self and other. *Soc Cogn Affect Neurosci*. 2011;6(2):218-225. doi:10.1093/scan/nsq046

75. Albrecht K, Volz KG, Sutter M, von Cramon DY. What do I want and when do I want it: brain correlates of decisions made for self and other. *PloS One*. 2013;8(8):e73531.

76. Ballard K, Knutson B. Dissociable neural representations of future reward magnitude and delay during temporal discounting. *NeuroImage*. 2009;45(1):143-150. doi:10.1016/j.neuroimage.2008.11.004

77. Banich MT, De La Vega A, Andrews-Hanna JR, Mackiewicz Seghete K, Du Y, Claus ED. Developmental trends and individual differences in brain systems involved in intertemporal choice during adolescence. *Psychol Addict Behav*. 2013;27(2):416-430. doi:10.1037/a0031991

78. Bickel WK, Pitcock JA, Yi R, Angtuaco EJC. Congruence of BOLD response across intertemporal choice conditions: Fictive and real money gains and losses. *J Neurosci*. 2009;29(27):8839-8846. doi:10.1523/JNEUROSCI.5319-08.2009

79. Blain B, Schmit C, Aubry A, Hausswirth C, Le Meur Y, Pessiglione M. Neuro-computational Impact of Physical Training Overload on Economic Decision-Making. *Curr Biol CB*. 2019;29(19):3289-3297.e4.

80. Chantiluke K, Christakou A, Murphy CM, et al. Disorder-specific functional abnormalities during temporal discounting in youth with Attention Deficit hyperactivity Disorder (ADHD), Autism and comorbid ADHD and Autism. *Psychiatry Res Neuroimaging*. 2014;223(2):113-120. doi:10.1016/j.pscychresns.2014.04.006

81. Clithero JA, Carter RM, Huettel SA. Local pattern classification differentiates processes of economic valuation. *NeuroImage*. 2009;45(4):1329-1338.

82. Cox KM, Kable JW. BOLD Subjective Value Signals Exhibit Robust Range Adaptation. *J Neurosci*. 2014;34(49):16533-16543. doi:10.1523/JNEUROSCI.3927-14.2014

83. Eppinger B, Nystrom LE, Cohen JD. Reduced sensitivity to immediate reward during decision-making in older than younger adults. *PloS One*. 2012;7(5):e36953.

84. Eppinger B, Heekeren HR, Li SC. Age Differences in the Neural Mechanisms of Intertemporal Choice Under Subjective Decision Conflict. *Cereb Cortex N Y N 1991*. 2018;28(11):3764-3774.

85. Fröhner JH, Teckentrup V, Smolka MN, Kroemer NB. Addressing the reliability fallacy in fMRI: Similar group effects may arise from unreliable individual effects. *NeuroImage*. 2019;195:174-189. doi:10.1016/j.neuroimage.2019.03.053

86. Gardiner CK, Karoly HC, Thayer RE, Gillman AS, Sabbineni A, Bryan AD. Neural activation during delay discounting is associated with 6-month change in risky sexual behavior in adolescents. *Ann Behav Med Publ Soc Behav Med*. 2018;52(5):356-366.

87. Hare TA, Hakimi S, Rangel A. Activity in dlPFC and its effective connectivity to vmPFC are associated with temporal discounting. *Front Neurosci*. 2014;8:50.

88. Ikink I, Engelmann JB, van den Bos W, Roelofs K, Figner B. Time ambiguity during intertemporal decision-making is aversive, impacting choice and neural value coding. *NeuroImage*. 2019;185:236-244. doi:10.1016/j.neuroimage.2018.10.008

89. Koffarnus MN, Deshpande HU, Lisinski JM, Eklund A, Bickel WK, LaConte SM. An adaptive, individualized fMRI delay discounting procedure to increase flexibility and optimize scanner time. *NeuroImage*. 2017;161:56-66. doi:10.1016/j.neuroimage.2017.08.024

90. Laube C, Lorenz R, van den Bos W. Pubertal testosterone correlates with adolescent impatience and dorsal striatal activity. *Dev Cogn Neurosci*. 2020;42:100749.

91. Liu L, Feng T, Wang J, Li H. The neural dissociation of subjective valuation from choice processes in intertemporal choice. *Behav Brain Res*. 2012;231(1):40-47. doi:10.1016/j.bbr.2012.02.045

92. Liu L, Feng T. The neural predictors of choice preference in intertemporal choice. *Brain Res*. 2012;1436:92-100.

93. Luo S, Ainslie G, Pollini D, Giragosian L, Monterosso JR. Moderators of the association between brain activation and farsighted choice. *NeuroImage*. 2012;59(2):1469-1477. doi:10.1016/j.neuroimage.2011.08.004

94. Massar SAA, Libedinsky C, Weiyan C, Huettel SA, Chee MWL. Separate and overlapping brain areas encode subjective value during delay and effort discounting. *NeuroImage*. 2015;120:104-113. doi:10.1016/j.neuroimage.2015.06.080

95. Meade CS, Cordero DM, Hobkirk AL, Metra BM, Chen NK, Huettel SA. Compensatory activation in fronto-parietal cortices among HIV-infected persons during a monetary decision-making task. *Hum Brain Mapp*. 2016;37(7):2455-2467.

96. Murawski C, Harris PG, Bode S, Domínguez D JF, Egan GF. Led into temptation? Rewarding brand logos bias the neural encoding of incidental economic decisions. *PloS One*. 2012;7(3):e34155.

97. Neukam PT, Deza-Araujo YI, Marxen M, et al. No evidence for the involvement of serotonin or the 5-HTTLPR genotype in intertemporal choice in a larger community sample. *J Psychopharmacol Oxf Engl*. 2019;33(11):1377-1387.

98. O'Connell G, Hsu CT, Christakou A, Chakrabarti B. Thinking about others and the future: Neural correlates of perspective taking relate to preferences for delayed rewards. *Cogn Affect Behav Neurosci*. 2018;18(1):35-42.

99. Peters J, Büchel C. Overlapping and distinct neural systems code for subjective value during intertemporal and risky decision making. *J Neurosci Off J Soc Neurosci*. 2009;29(50):15727-15734.

100. Peters J, Büchel C. Episodic future thinking reduces reward delay discounting through an enhancement of prefrontal-mediotemporal interactions. *Neuron*. 2010;66(1):138-148. doi:10.1016/j.neuron.2010.03.026

101. Pine A, Seymour B, Roiser JP, et al. Encoding of marginal utility across time in the human brain. *J Neurosci Off J Soc Neurosci*. 2009;29(30):9575-9581.

102. Pine A, Shiner T, Seymour B, Dolan RJ. Dopamine, time, and impulsivity in humans. *J Neurosci*. 2010;30(26):8888-8896. doi:10.1523/JNEUROSCI.6028-09.2010

103. Piva M, Velnoskey K, Jia R, Nair A, Levy I, Chang SW. The dorsomedial prefrontal cortex computes task-invariant relative subjective value for self and other. *eLife*. 2019;8. Accessed January 1, 6AD. https://pubmed.ncbi.nlm.nih.gov/31192786/

104. Ripke S, Hübner T, Mennigen E, et al. Reward processing and intertemporal decision making in adults and adolescents: The role of impulsivity and decision consistency. *Brain Res*. 2012;1478:36-47. doi:10.1016/j.brainres.2012.08.034

105. Samanez-Larkin G, Mata R, Radu P, Ballard I, Carstensen L, McClure S. Age Differences in Striatal Delay Sensitivity during Intertemporal Choice in Healthy Adults. *Front Neurosci*. 2011;5:126. doi:10.3389/fnins.2011.00126

106. Sasse LK, Peters J, Büchel C, Brassen S. Effects of prospective thinking on intertemporal choice: The role of familiarity. *Hum Brain Mapp*. 2015;36(10):4210-4221.

107. Sasse LK, Peters J, Brassen S. Cognitive control modulates effects of episodic simulation on delay discounting in aging. *Front Aging Neurosci*. 2017;9:11.

108. Schneider S, Peters J, Peth JM, Büchel C. Parental inconsistency, impulsive choice and neural value representations in healthy adolescents. *Transl Psychiatry*. 2014;4(4):e382.

109. Steinbeis N, Haushofer J, Fehr E, Singer T. Development of behavioral control and associated vmPFC–DLPFC connectivity explains children's increased resistance to temptation in intertemporal choice. *Cereb Cortex*. 2016;26(1):32-42. doi:10.1093/cercor/bhu167

110. van den Bos W, Rodriguez CA, Schweitzer JB, McClure SM. Connectivity strength of dissociable striatal tracts predict individual differences in temporal discounting. *J Neurosci Off J Soc Neurosci*. 2014;34(31):10298-10310.

111. Vetter NC, Steding J, Jurk S, Ripke S, Mennigen E, Smolka MN. Reliability in adolescent fMRI within two years - a comparison of three tasks. *Sci Rep*. 2017;7(1):2287. doi:10.1038/s41598-017-02334-7

112. Wang Q, Luo S, Monterosso J, et al. Distributed value representation in the medial prefrontal cortex during intertemporal choices. *J Neurosci*. 2014;34(22):7522-7530. doi:10.1523/JNEUROSCI.0351-14.2014

113. Xu L, Liang ZY, Wang K, Li S, Jiang T. Neural mechanism of intertemporal choice: From discounting future gains to future losses. *Brain Res*. 2009;1261:65-74.

114. Zha R, Bu J, Wei Z, et al. Transforming brain signals related to value evaluation and self-control into behavioral choices. *Hum Brain Mapp*. 2019;40(4):1049-1061.

115. Zhuang JY, Wang JX, Lei Q, Zhang W, Fan M. Neural Basis of Increased Cognitive Control of Impulsivity During the Mid-Luteal Phase Relative to the Late Follicular Phase of the Menstrual Cycle. *Front Hum Neurosci*. 2020;14:568399.

PRISMA 2009 Flow Diagram

Records identified through
database searching
(n = 590)

Additional records identified
through other sources
(n = 3)

Records after duplicates removed
(n = 393)

Records screened
(n = 393)

Records excluded
Animal studies (n = 5)
Not a primary source (n = 40)

Full-text articles assessed
for eligibility
(n = 348)

Full-text articles excluded
(n = 274)

No fMRI during delay
discounting (n=188)
MVPA or connectivity (n=13)
Non-monetary reward or delay
less than 1 day (n=22)
No whole-brain contrast (n=6)
No coordinates available (n=45)

Studies included in
quantitative synthesis
(meta-analysis)
(n = 74)

Figure 2. Whole-brain meta-analysis of common analytic strategies for BOLD signals during delay discounting task. There were no suprathreshold clusters for SSR > LLR and Now.

A Task

L

 R

x = -2   z = 44   z = 10   x = -50   z = 0

B Now

x = -8   z = -4

C LLR > SSR

z = 54

D Subjective value

E Magnitude (red) and delay (blue)

F Hard (red) vs. Easy (blue)

10%   64%

Notes: LLR = larger-later reward, SSR = smaller-sooner reward.

Figure 3. Spatial overlap analysis of clusters by individual contrasts.

A Valuation system

B SV and Easy

C Task and Hard

D Task and Delay

Notes: SV = subjective value, Mag = magnitude, Del = delay.

Table 1. Meta-analytic sample (N=72 studies; 136 contrasts).

Study	n	Diagnosis	Contrast	Number of foci								
				Task	LLR	SSR	SV	Mag	Del	Hard	Easy	Now
Albert et al., 2012[37]	40	NS	Choice evaluation > baseline Positive effects of differential magnitude	4				6				
Albrecht et al., 2011[74]	28	NS	Today trials > delay trials									18
Albrecht et al., 2013[75]	30	HC	SS > LL				3					
Avsar et al., 2013[53]	14	HC	Task > SMC trials Easy > hard	18							9	
	14	Schizophrenia	Task > SMC trials Easy > hard	8							9	
Ballard et al., 2009[76]	16	HC	Magnitude Negative effect of delay					3	3			
Banich et al., 2013[77]	29	HC	Task > fixation Now > later	22		17						
Bickel et al., 2009[78]	30	HC	Discounting > control (real money)	21								
Blain et al., 2019[79]	37	NS	Hard > easy							18		
Chantiluke et al., 2014[80]	18	HC	Delayed > immediate choices		2							
Christakou et al., 2011[42]	40	HC	Delay > immediate Immediate > delay		13	4						
Claus et al., 2011[41]	160	Heavy drinkers	Later > now Now > later		17	18						
Clithero et al., 2009[81]	11	HC	Delay task	3								
Cox et al., 2014[82]	20	HC	SV (narrow and wide)				4					
Deshpande et al., 2019[57]	54	Prediabetics (high BMI)	Hard > easy Easy > hard Immediate available vs. unavailable							6	11	18
Eppinger et al., 2012[83]	30	NS	All choice > baseline Immediate options > delayed options	5								2
Eppinger et al., 2018[84]	50	NS	SV Hard > easy Easy > hard				3			2	4	
Fröhner et al., 2019[85]	126	NS	SV				3					

Study	n	Diagnosis	Contrast	Task	LLR	SSR	SV	Mag	Del	Hard	Easy	Now
Gardiner et al., 2018[86]	177	NS	Hard > easy							4		
Hamilton et al., 2020[67]	27	HC	All choices vs. baseline	8								
Hare et al., 2014[87]	25	HC	SV Del > imm		24		11					
Hoffman et al., 2008[47]	36	Methamphetamine dependence[1]	DD task > magnitude estimation task	11								
Ikink et al., 2019[88]	24	NS	Conjunction of ambiguous and exact DD	7								
Kable et al., 2007[21]	10	NS	SV using HDR Objective value Inverse delay Delayed > immediate Negative effects of SVD		1		14	2	4	8		
Kable et al., 2010[20]	25	NS	NOW - 60 DAY									9
Kishinevsky et al., 2012[55]	19	Obese women	Hard > easy							2		
Koffarnus et al., 2017[89]	18	NS (excluded substance use disorder)	Hard > easy Easy > hard							1	5	
Laube et al., 2020[90]	70	HC	Delayed > immediate Immediate > delayed Now-Later options		8	8						8
Lempert et al., 2017[13]	35	NS	SV (memory and control choice)				24					
Lim et al., 2017[48]	17	Alcohol use disorder	SS > LL			3						
Liu et al., 2012a[91]	19	HC	Free choice > forced choice SV	11			17					
Liu et al., 2012b[92]	18	NS	Task > baseline during valuation	7								
Luo et al., 2012[93]	21	HC	LL > SS		7							

NEURAL SYSTEMS IN DELAY DISCOUNTING

Study	n	Diagnosis	Contrast	Task	LLR	SSR	SV	Mag	Del	Hard	Easy	Now
							Number of foci					
Marco-Pallarés et al., 2010[64]	17	HC	Indifference > non-indifference									
			Non-indifference > indifference							1	11	
Martin et al., 2015[58]	19	Poor quality sleepers[1]	Immediate > baseline	12								
Massar et al., 2015[94]	23	HC	SV				21					
			Negative effects of delay level						12			
Mavrogiorgou et al., 2017[66]	20	HC	Immediate > delay				8					
Mavrogiorgou et al., 2018[52]	38	OCD[1]	Immediate > delay				7					
McClure et al., 2004[18]	14	NS	Delta system									
			Beta system	8								5
Meade et al., 2011[49]	39	Cocaine dependence[1]	Task > baseline	1								
			Hard > easy							9		
Meade et al., 2016[95]	35	HC	Task > baseline									
			Hard > easy	8						8		
Miedl et al., 2012[25]	32	Pathological gamblers[1]	SV				25					
Miedl et al., 2015[40]	15	HC	LDR > SIR			2						
	15	Pathological gamblers	LDR > SIR			8						
	30	Pathological gamblers[1]	Indifferent > sure									
			Sure > indifferent							20	13	
Monterosso et al., 2007[31]	29	Methamphetamine users[1]	Hard > easy	6						2		
Morys et al., 2018[56]	36	Obese[1]	Delay discounting > perceptual decisions									
			SS > LL	7		4						
Murawski et al., 2012[96]	13	NS	Task > baseline									
			SV									
			Easy > hard	13				6				5
Neukam et al., 2019[97]	90	HC	Parametric amount									
			Negative effects of delay					7	32			

Study	n	Diagnosis	Contrast	Task	LLR	SSR	SV	Mag	Del	Hard	Easy	Now
Norman et al., 2017[39]	20	HC	Delay > immediate (both directions)		2	1						
	26	ADHD	Delay > immediate (both directions)		2	7						
	20	OCD	Delay > immediate (both directions)		2	3						
O'Connell et al., 2018[98]	26	NS	DEL > IMM Positive effects of net SV		3						14	
Peters et al., 2009	22	NS	SV Reward magnitude Inverse delay				25	15	13			
Peters et al., 2010[100]	30	HC	SV (control and episodic condition)				15					
Pine et al., 2009[101]	24	HC	Discounted utility (V) Undiscounted utility (U) Temporal discount factor (D) Hard > easy				7	16	22	7		
Pine et al., 2010[102]	14	HC	Discounted utility (V) Undiscounted utility (U) Temporal discount factor (D)				9	25	28			
Piva et al., 2019[103]	20	NS	Hard > easy Positive effect of relative SV							13	2	
Ripke et al., 2012[104]	222	HC	Task SV	18			19					
Samanez-Larkin et al., 2011[105]	25	NS	SV				19					
Sasse et al., 2015[106]	23	HC	SV (episodic condition)				12					
Sasse et al., 2017[107]	22	HC	SV				8					
Schneider et al., 2014[108]	48	HC	SV				4					
Sripada et al., 2011[71]	20	HC	LL > SS Immediacy			15						6
Stanger et al., 2013[50]	30	Substance abuse	LL > control task SS vs. LL	14		2						
Steinbeis et al., 2016[109]	20	HC	SV during valuation						4			
van den Bos et al., 2014[110]	22	NS	Choice > baseline LL-SS	7	5							

Study	n	Diagnosis	Contrast	Number of foci								
				Task	LLR	SSR	SV	Mag	Del	Hard	Easy	Now
van den Bos et al., 2015[69]	50	NS	LL-SS				4					
Vanyukov et al., 2016[54]	48	Major depression[1]	Task > fixation SIR-DDR Negative effect of longer vs. shorter delay Difficult > easy Negative effect of difficult > easy	11				1	1	3	1	
Vetter et al., 2017[111]	174	HC	Decision phase > baseline	1								
Waegeman et al., 2014[43]	41	HC	Delayed > immediate				41					
Wang et al., 2014[112]	28	HC	Amount of delayed reward Positive effects of SVD					7			11	
Wang et al., 2017[51]	21	HC	Delay - Immediate				3					
	18	Internet gambling disorder	Delay - Immediate				1	2				
Weber et al., 2008[68]	23	NS	Delayed > immediate				1					
Wiehler et al., 2017[26]	48	Pathological gamblers[1]	SV (conjunction of HC and gamblers)				3					
Wittmann et al., 2007[70]	13	HC	<1 versus >1 year Delayed vs. immediate				7		1			
Xu et al., 2009[113]	20	HC	Temporal discounting (gains) > baseline Available immediately (G-TD)	19								12
Zha et al., 2019[114]	46	HC	SV Difficulty						10	8		
Zhuang et al., 2020[115]	24	HC	Delayed > immediate Immediate > delayed				3	8				

Notes: HC = healthy controls, NS = not specified, Task = Task activations, LLR = larger later reward > smaller sooner reward, SSR = smaller sooner reward > larger later reward, SV = positive correlation with subjective value, Mag = positive correlation with objective (undiscounted) magnitude of reward, Del = negative correlation with delay or positive correlation with inverse delay, Hard = hard > easy, Easy = easy > hard, Now = trials with immediate option > trials with both options delayed. [1] includes HC (pooled across the entire study sample).

Table 2. Meta-analytic test statistic (% of studies that reported a peak within a 10mm radius).

#	Volume (mm^3)	MNI coordinates			Value	Brain area
		X	Y	Z		
Task						
1	1222	-2	16	48	48.0	L. Superior frontal gyrus
2	671	34	-62	44	32.0	R. Intraparietal sulcus
3	486	-28	-60	46	32.0	L. Intraparietal sulcus
4	440	44	36	10	32.0	R. Frontal pole
5	476	-50	10	28	28.0	L. Inferior frontal gyrus
6	399	38	24	0	28.0	R. Insula
7	375	26	-94	-6	28.0	R. Occipital pole
Now						
1	232	-6	48	-12	50.0	L. vmPFC
2	253	-2	12	-8	50.0	L. Striatum
SSR > LLR						
No suprathreshold clusters						
LLR > SSR						
1	715	-34	-18	54	38.1	L. Precentral gyrus
Hard > Easy						
1	756	2	26	42	50.0	R. Superior frontal gyrus
2	342	50	-36	38	25.0	R. Supramarginal gyrus
Easy > Hard						
No suprathreshold clusters						
SV						
1	1452	-10	4	-6	63.6	L. Striatum
2	1333	-2	48	-4	45.5	L. vmPFC
3	400	-6	-40	34	36.4	L. Posterior cingulate
4	377	-58	-36	-6	27.3	L. Middle temporal gyrus
Magnitude						
1	490	14	4	-6	55.6	R. Striatum
Delay						
1	461	46	28	4	55.6	R. Inferior frontal gyrus
2	537	2	36	10	44.4	R. Medial prefrontal cortex
3	356	50	14	24	44.4	R. Middle frontal gyrus
4	346	-40	32	0	44.4	L. Inferior frontal gyrus
5	269	-12	4	2	44.4	L. Striatum

Notes: Task = Task activations, Now = trials with immediate option > trials with both options delayed, SSR = smaller-sooner reward > larger-later reward, LLR = larger-later reward > smaller-sooner reward, Hard = hard > easy, Easy = easy > hard, SV = positive correlation with subjective value, Magnitude = positive correlation with objective (undiscounted) magnitude of reward, Delay = negative correlation with delay or positive correlation with inverse delay, L = left, R = right.