Inulin prebiotic reinforces host cancer immunosurveillance via ɣδ T cell activation

Emilie Boucher[1], Caroline Plazy[2], Mathias L. Richard[3,4], Antonia Suau[1,5], Irène Mangin[5], Muriel Cornet[2], Delphine Aldebert[1], Bertrand Toussaint[2], Dalil Hannani[1*]

[1] Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, TIMC, 38000 Grenoble, France

[2] Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, CHU Grenoble Alpes, TIMC, 38000 Grenoble, France

[3] Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78352 Jouy-en-Josas, France.

[4] Paris Center for Microbiome Medicine, Fédération Hospitalo-Universitaire, 75012 Paris, France.

[5] USC Cnam-ANSES Metabiot, Conservatoire National des Arts et Métiers, 75003 Paris, France

[*] <u>Corresponding author</u>:

Dr. Dalil HANNANI

Faculté de Médecine de Grenoble – Laboratoire TIMC Equipe TrEE

Bâtiment Jean Roget 38700 La Tronche, France

Dalil.Hannani@univ-grenoble-alpes.fr

Word Count: 3233 (Abstract, References, Material and Method excluded)

Number of Figures: 6

Abstract:

Gut microbiota is now recognized as a key parameter impacting the host's anti-cancer immunosurveillance and ability to respond to immunotherapy. Therefore, its optimal modulation for preventive and therapeutic purposes is very appealing. Diet is one of the most potent modulators of the microbiota, and thus nutritional intervention could be exploited to improve host anticancer immunity. Here we show that inulin-enriched diet, a prebiotic known to promote bacteria described as immunostimulatory, triggers an enhanced Th1-polarized CD4$^+$ and CD8$^+$ αβ T cell-mediated anti-tumor response and attenuates tumor growth in three preclinical tumor-bearing mouse models. We highlighted that inulin-mediated anti-tumor effect relies on the activation of both intestinal and tumor-infiltrating ɣδ T cells that are indispensable for αβ T cell activation and subsequent tumor growth control, in a microbiota-dependent manner. Overall, our data identified these cells as a critical immune subset, mandatory for inulin-mediated anti-tumor immunity *in vivo*, further supporting and rationalizing the use of such prebiotic approaches as well as the development of immunotherapies targeting ɣδ T cells in cancer prevention and immunotherapy.

Introduction:

According to the World Health Organization (WHO), cancer is the second leading cause of death globally, and increasing cancer incidence is a major public health issue worldwide (1). Despite the development of innovative treatments such as immunotherapy, including immune checkpoint blockers (ICB) that has recently revolutionized patient care, quality of life and survival, huge efforts are still needed to prevent and treat cancers. It is worth noting that between 30% and 50% of cancer deaths could be prevented by modifying or avoiding major risk factors and implementing existing evidence-based prevention strategies, including eating a healthy diet, primarily fruits and vegetables (2). A healthy diet helps prevent the development of cancer in a number of ways that may be synergistic. It limits host cell oxidation and metabolism, as well as genetic instability and chronic inflammation. It acts also on gut microbiota composition and functions, and subsequently on host immune functions (3).

Indeed, host immune tonus is a key parameter in cancer prevention through immunosurveillance, which relies on the ability of different immune cell subsets to patrol, recognize and eliminate transformed cells (4). Both innate and adaptive immunity participate in this process and prevent cancer development by recognizing stress-induced ligands, tumor cells metabolic dysregulation, or peptides derived from tumor antigens. Effector CD8$^+$ T cells are one of the most important subsets involved in cancer surveillance and are key targets in immunotherapy. Furthermore, γδ T cells are unconventional T cells representing a unique population displaying multiple functions, including potent anti-tumor activity (5,6). Indeed, γδ T cells are key players in cancer immunosurveillance, as they can recognize transformed cells expressing stress-ligands via innate receptors (e.g. NKG2D) or metabolic products derived from dysregulated tumors (e.g. mevalonate pathway). When activated, γδ T cells produce pro-inflammatory cytokines, such as interferon γ (IFNγ), which in turn promote anti-cancer CD8$^+$ T cell immunity. They can also directly kill cancer cells via perforin/granzyme-mediated cytotoxicity. Therefore, γδ T cell infiltration within the tumor bed is a favorable prognostic marker in several human cancers (7,8), and they represent a promising target in cancer immunotherapy (9,10).

A growing body of evidences has also highlighted that gut microbiota may have a positive impact on cancer prevention or treatment, especially via improved host immunosurveillance of cancer (11,12) as well as patients' ability to respond to chemotherapy (13) or immunotherapy (11,14–16). These seminal studies have identified certain families, genera, or species of bacteria associated with the gut microbiota that appear to correlate with a strong host immune status and subsequent favorable response to cancer therapy. For instance, *Bifidobacterium, Ruminoccocaceae, Akkermansia muciniphila, Enterococcus irae* and *Allistipes* have all been positively associated with ICB responses. As a preclinical proof of concept, it has been shown that either Fecal Microbiota Transplantation (FMT) from responding patients or a direct supplementation with immunostimulatory bacteria can restore the ability to respond to ICB in mice (14,15). Taken together, these studies strongly suggest that modulation of gut microbiota may represent a major strategy to prevent and/or treat cancers, by optimizing host immunity and subsequent antitumor responses.

It is noteworthy that 70% of host immune cells are located within the gut, close to microbiota-derived signals (17). Among Intra Epithelial Lymphocytes (IELs) in the gut, the γδ T cells represent the largest immune subset. Interestingly, their early development is largely dependent of gut microbiota after birth (18). Unlike conventional αβ TcR[+] CD4[+] or CD8[+] T cells that recognize antigen-derived peptides, these cells recognize metabolic products derived from bacteria or tumor cells (19,20).

Diet is probably the most powerful microbiota modulator, both on its composition and metabolic functions (21). Thus, modulation of the gut microbiota by nutritional intervention and in particular via prebiotics could be of great interest to optimize host immune responses including cancer immunosurveillance. Prebiotics are defined as non-digestible food ingredients that benefit the host by selectively stimulating the growth and activity of one species or a limited number of bacterial species in the colon (22). Among prebiotics, inulin, a Fructo-Oligo Saccharide derived from chicory roots, is well known to promote some colonic bacteria including *Bifidobacterium* genus which has been associated with optimal host immunity in the context of cancer immunosurveillance and immunotherapy (11,23). Recently, the use of inulin to stimulate immunosurveillance has been confirmed by Li *et al.* who showed that the inulin diet attenuated tumor growth in a CD8[+] T cell-dependent manner in mice (24). To date, how a nutritional intervention might affect systemic immunity, including the anti-tumor CD8[+] T cell response remains largely unknown.

Because inulin-enriched diet is known to alter the metabolic activity of the gut microbiota (25), we therefore hypothesized that intestinal and systemic γδ T cells might subsequently be altered. In the present study we therefore examined the role of γδ T cells in inulin-mediated anti-tumor effect and highlighted their pivotal role in CD8[+] T cell–mediated immunosurveillance in several transplantable mouse tumor models.

Results:

Inulin enriched diet leads to potent tumor growth control and enhanced γδ TILs activity

Inulin has been described as a promoter of immunostimulatory bacteria associated with strong anti-tumor immunity (11,26). Therefore, we evaluated whether 15 days of preconditioning in WT mice by adding inulin in drinking water could promote host anti-tumor immunity after subcutaneous injection of a syngeneic B16-OVA melanoma tumor (**Fig1A**). Consistent with a previous study in another tumor model (24), inulin resulted in a strong attenuation of the aggressive B16-OVA melanoma tumor growth (**Fig 1B left panel**). Of note, the inulin-mediated anti-tumor effect was also confirmed in other tumor models (i.e. MCA205 fibrosarcoma and MC38 colorectal cancer cell lines, **Fig 1B middle and right panel**). To better understand the mechanisms behind these observations, we analyzed the Tumor Infiltrating Lymphocytes (TILs) when the control diet group reached maximum ethic tumor size (day 13 post B16-OVA tumor inoculation, **Fig 1A**). Flow cytometry analyses revealed that inulin triggered a greater infiltration of immune cell (CD45$^+$ cells) within the tumor bed (**Fig 1C**) as well as a greater tumor infiltration of dendritic cells (**Fig S1**). Consistent with the anti-tumor activity observed (**Fig 1B**), we found that inulin-enriched diet triggered a potent Th1 anti-tumor response as illustrated by greater amount of IFNγ-producing CD4$^+$ and CD8$^+$ TILs (**Fig 1 D and E**). In agreement with this, immunization against OVA protein adjuvanted with PolyI:C (a Th1 polarizing adjuvant) led to a higher frequency of OVA-specific CD8$^+$ T cells in lymph nodes of inulin-supplemented mice (**Fig S2**). This suggests that inulin potentiated the Th1 polarizing activity of PolyI:C.

In addition, we assessed the frequency and activity of γδ TILs, and found that inulin-enriched diet also promoted γδ TIL infiltration (**Fig 1C**) and IFNγ-production in residual tumors (**Fig 1F**). Notably, among both MCA205 and MC38 TILs, only γδ T cells infiltration was significantly increased under the inulin diet (**Fig S3**). Taken together, these data indicate that inulin triggered a greater tumor infiltration by immune cells, potent Th1-polarized anti-tumor immunity, and in particular, high IFNγ production by conventional T cells as well as γδ TILs, which may ultimately lead to tumor growth control.

Inulin-mediated anti-tumor immunity relies on γδ T cell

γδ T cells are unconventional T cells recognizing metabolic-related molecules such as phosphoantigens, and represent an interesting target/tool in cancer immunotherapy because they exhibit potent anti-tumor activity (6). Since their tumor infiltration and IFNγ production are enhanced by inulin-enriched diet, we therefore evaluated whether these cells could play a central role in inulin-mediated anti-tumor immunity. To assess their contribution, we used repeated injections of γδ TcR-blocking antibody starting the day before the tumor implantation (**Fig 2A**). Of note, while inulin-treated mice well-controlled tumor growth at day 13, after implantation, administration of γδ TcR-blocking antibody completely abrogated the inulin-mediated anti-tumor effect (**Fig 2B**). Interestingly, analysis of TILs revealed that under systemic γδ T cell blockade, the inulin-enriched diet failed to trigger greater IFNγ-production by CD4$^+$ and CD8$^+$ TILs (**Fig 2 C and D**). Furthermore, under γδ TcR blockade and despite the inulin diet, the frequency of IFNγ-producing γδ TILs was similar to that of the control group,

suggesting that γδ TcR signaling is required in inulin-mediated immunoactivation (**Fig2E**). Taken together, these data indicate that γδ T cells, and most likely their TcR engagement, are mandatory for the inulin-mediated anti-tumor effect. In addition, we observed a significant correlation between the level of tumor infiltration of IFNγ-producing γδ T cells and tumor growth, for all three tumor models (**Fig 2F**), confirming their pivotal role in inulin-mediated anti-tumor effect.

Inulin-enriched diet leads to gut-associated γδ T cell activation and gut inflammation

It is now well known that gut microbiota affects local (gut) and systemic immune tone (27). Since γδ T cells represent the most abundant immune cell population within the gut epithelium (28), we evaluated the impact of the inulin diet on IFNγ-production by gut-associated immune cells as well as on the level of gut epithelium inflammation. For this purpose, we collected intestines from mice after a 15-day inulin diet (**Fig 3A**). Fractions of Intra Epithelial Lymphocytes (IELs) and Lamina Propria Cells (LPCs) were analyzed using flow cytometry. Under inulin, while the frequency of IFNγ-producing CD4$^+$ and CD8$^+$ T cells was comparable to that of the control group in IELs and LPCs (**Fig 3B-C and Fig S4A, respectively**), γδ T cell IELs showed a higher proportion of IFNγ-producing cells (**Fig 3D**).

To further characterize inulin-mediated intestinal inflammation, we sorted the epithelial (CD45-) fraction from LPCs (CD45$^+$) to perform RT-qPCR on genes involved in inflammation, tissue repair, and tight junctions (**Fig 3 E-G, Fig S5 and Table S2**). Of the 19 analyzed genes, three inflammation-related genes were significantly upregulated under the inulin regime in epithelial cells, namely Tumor Necrosis Factor (TNF)-α, Cyclo-Oxygenase (Cox)-2 and Matrix Metalloprotease (MMP)-9 (**Fig 3F**). In the CD45$^+$ compartment (LPCs), inulin triggered significant overexpression of Macrophage Inflammatory Protein (MIP)-2, Interleukin (IL)-22 and the transcriptional factor Forkhead Box P (FoxP)-3 (**Fig 3G**). Similarly, we observed a trend toward higher IL-10 production, although not significant, by LPC-associated T cells (**Fig S4B**). Altogether, these results indicate that inulin diet triggered both targeted γδ T cell activation in IELs and inflammation in gut epithelial cells, in parallel with tissue repair signaling and tolerance induction in the *lamina propria*.

Inulin-enriched diet alters the gut microbiota and promotes Bifidobacterium *growth*

To establish a link between the observed intestinal immune activation (**Fig 3**) and the inulin-enriched diet, we then characterized the composition of the fecal bacterial microbiota after inulin administration, using 16S rDNA sequencing. Metagenomics analysis indicated that 5 phyla composed the microbiota of these mice: *Firmicutes* (58% of the sequences), *Bacteroidota* (37% of the sequences), *Desulfobacterota* (1.3% of the sequences), *Proteobacteria* (1.2% of the sequences) and *Actinobacteriota* (2.4% of the sequences). A 15-day Inulin diet profoundly altered the gut microbiota composition as revealed by the relative abundance of these bacterial phyla observed in inulin-treated mice compared to control mice (**Fig 4A**), with a visible increase of bacteria of the *Actinobacteriota* phylum, and more precisely to the genus *Bifidobacterium*. This diet altered the alpha diversity, the total number of species was not significantly different (387±53 OTUs for the control group and 378±43 for the inulin group) but the distribution of sequences within the OTUs was significantly different (Shannon

index, p value = 0.00582). A Bray-Curtis β-diversity analysis illustrated by PCoA representation confirmed a significant clustering of the overall bacterial microbiota with a p value < 0.001 (**Fig 4B**). To identify whether particular bacteria were responsible for this dysbiosis and potentially involved in anti-tumor immunity, we performed differential abundance correlation using a linear discriminant analysis (LDA) effect size algorithm (LEfSe see Material and Methods). The LDA scores plotted in a phylogenic cladogram showed significant differences in several taxonomic ranks (**Fig 4C**). One of the strongest signals, LDA score>4, was the increase in *Bifidobacterium animalis* subsp. *animalis,* a member of the *Actinobacteriota* phylum in inulin treated mice (4.67%), compared with control mice (0.03%) (**Fig4D, Table S1**). Importantly, consistent with our results, the *Bifidobacterium* genus has been shown to exhibit immunostimulatory properties in the contexts of cancer immunosurveillance and immunotherapy (11,23). Therefore, *Bifidobacterium* may be involved in the stimulation of inulin-mediated anti-tumor immunity. The second significant signal concerned the family Clostridiaceae, and in particular the genus *Clostridium sensu stricto 1*. Our sequences were close to *Clostridium saudiense*, *Clostridium disporicum* and *Clostridium celatum* (these species have nearly identical 16S rDNA sequences). This OTU statistically decreased in the mice microbiota due to treatment from 6% in the control group to 0,09 % in the inulin group (data not shown).

Inulin-mediated increased anti-tumor immunity relies on gut microbiota

Inulin, as a polysaccharide molecule, has been shown to have adjuvant properties *in vivo* (29,30). Such direct immune activation could itself be responsible for the antitumor immunity observed under the inulin regime. To address this issue, we administrated broad-spectrum antibiotics cocktail to the mice 2 days before the inulin diet to significantly reduce the number of bacteria in their gut, as previously described (31) (**Fig 5A**). After tumor implantation, we observed that concomitant antibiotic treatment totally abolished the ability of inulin to attenuate tumor growth (**Fig 5B**). Furthermore, inulin failed to trigger a higher frequency of IFNγ-producing γδ TILs and subsequent CD4[+] and CD8[+] TILs (**Fig 5 C, D and E**). Altogether, these findings indicate that inulin-mediated anti-tumor activity cannot be attributed to a direct adjuvant effect; rather, it relies on host immune activation mediated by the gut microbiota.

Because γδ T cells are obligatory for inulin-mediated anti-tumor immunity (**Fig 2**), and that γδ T IELs are the only T cell subset activated by inulin diet in the gut (**Fig 3**), we hypothesized that γδ TILs could have an intestinal origin. To address this question, we analyzed the expression of the gut-homing chemokine receptor CCR9 on blood-circulating γδ T and γδ TILs. Whereas the number of CCR9-expressing γδ T decreased in the blood (**Fig 5F**), their frequency tended to increase in the tumor bed, despite a possible loss of expression in this environment (**Fig 5G**). Of note, under a concomitant inulin diet and antibiotic treatment, γδ TILs expressed similar level of CCR9 compared with the control group (**Fig 5H**). Taken together, these data suggest that inulin-induced activated γδ TILs may be of intestinal origin.

Inulin regimen is as efficient as αPD-1 Immunotherapy

A large body of evidences suggests that certain immunostimulatory bacteria – including *Bifidobacterium* – may facilitate the efficacy of Immune Checkpoint Inhibitors, including those

targeting the anti-PD1/PD-L1 axis (11,14,15,32). Because the inulin diet alters the gut microbiota (**Fig5**) and favors *Bifidobacterium* (**Fig 4**) we evaluated whether this diet could be synergistic with anti-PD1 treatment (**Fig 6A**). Tumor growth monitoring indicated that inulin alone allowed the tumor growth control to the same extent as anti-PD-1 alone or in combination (**Fig 6B**). The individual mouse curves clearly showed that inulin alone triggered the strongest anti-tumor effect compared with anti-PD1 alone or even in combination (**Fig 6C**). TILs analysis revealed that inulin led to a greater IFNɣ production by ɣδ T cells (**Fig 6D**) and CD4[+] cells (**Fig 6E**), compared with anti-PD1 alone. The combination did not lead to a potentiation of this effect, despite similar PD-1 expression among TILs (**Fig S6**). Anti-PD1 immunotherapy is known to reinvigorate CD8[+] TILs and increase their ability to produce IFNɣ (33). In agreement with this, anti-PD1 treatment led to a higher proportion of IFNɣ-producing CD8[+] TILs within the tumor bed (**Fig 6F**). Interestingly, inulin alone also triggered such activation of CD8[+] TILs, at least to a similar extent.

Discussion:

The gut microbiota is emerging as a powerful host immune modulator that could be exploited in both cancer prevention and treatment. Consequently, there is an urgent need to better understand the cellular and molecular mechanisms that govern this modulation in order to rapidly translate these findings into innovative preventive or therapeutic strategies.

Prebiotics contribute to the modification of the composition and functions of microbiota, and subsequently, to the modification of host immunity. The importance of a prebiotic-enriched diet was recently highlighted in humans by Spencer *et al.*, as they reported that a fiber-enriched diet positively influences the response to immunotherapy in melanoma patients (34).

Inulin is a widely consumed prebiotic, either in the diet or as a dietary supplement. Recently, inulin has been reported to attenuate transplantable melanoma tumor growth (24) in a CD8[+] T cell dependent manner. In the current study, we reported that inulin-enriched diet triggers not only potent αβ T cell anti-tumor immunity, but also the accumulation and activation of intratumoral ɣδ T cell in a microbiota-dependent manner. We emphasized that these cells are indispensable for the increase of IFNɣ-producing CD4[+] and CD8[+] TILs and subsequent tumor growth control. We also reported that inulin-enriched diet remarkably alters the composition of the microbiota and, as expected, significantly promotes the selective growth of *Bifidobacterium* species, known to be immunostimulatory. Consistent with this, such a diet promotes the activation of ɣδ T IELs in the gut and we found that the frequency of CCR9[+] tumor-infiltrating cells tends to increase, suggesting a possible intestinal origin of ɣδ TILs. Interestingly, CCR9-expressing T cells have been shown to contribute to anti-tumor immunity, and their recruitment into the tumor-bed (by CCL25 intratumoral delivery, (35)) improves immunotherapy efficacy.

Regarding a possible synergy with immunotherapy, we found that a prophylactic inulin diet was as effective as anti-PD-1 immunotherapy in our setting, and that PD-1 expression was equivalent between groups. Because PD-1 expression identifies exhausted T cells only when combined with other inhibitory receptors such as Tim-3, LAG-3 or TIGIT (36), it is possible that

the PD-1$^+$ TILs under inulin were not in an exhausted state, in contrast to TILs in the control group. Furthermore, in our experiment, we only harvested the tumors 13 days after inoculation to characterize the functionality of the TILs, so it is possible that the combination of inulin and anti-PD-1 treatment would prevent later tumor escape, compared to anti-PD-1 alone in B16 melanoma model. To fully address this question, further phenotyping of TILs and prolonged monitoring of tumor growth would be required.

To date, how intestinal and/or systemic ɣδ T cells are activated by inulin-enriched diet remains an open question. Importantly, we observed that ɣδ TcR blockade abrogates inulin-mediated enhanced immunosurveillance. Beyond demonstrating that ɣδ T cells are obligate mediators, these data highlighted that their activation was TcR-dependent and thus that microbiota-derived ɣδ T cell activation signals could be metabolite-mediated. Altogether these data allow us to propose a scenario in which intestinal ɣδ T cells could be activated, in a TcR-dependent manner, most likely by metabolites derived from inulin-mediated remodeling of the microbiota and then recirculate to the tumor to promote CD8$^+$ T cell anti-tumor activity.

In this study, we report in three transplantable tumor models that inulin triggered anti-tumor immune responses. As a result, the growth of melanoma, colorectal and fibrosarcoma tumors were attenuated and accompanied by an increased in ɣδ T cell infiltration. To fully investigate the impact of nutritional intervention on cancer immunosurveillance mediated by ɣδ T cells, the use of more physiological models such as genetically engineered mouse models, or chemically induced cutaneous tumor models (4) may be useful as they allow for the establishment of proper immunoediting, unlike transplantable tumors.

Despite these limitations, our data provide a proof of concept showing that intestinal and systemic ɣδ T cell functions can be modulated by nutritional intervention. In particular, we have identified these cells as critical immune subset, mandatory for inulin-mediated antitumor immunity and subsequent control of the tumor growth *in vivo*. These results further support and rationalize the use of such prebiotic approaches as well as the development of immunotherapies targeting ɣδ T cells in cancer prevention and immunotherapy. Notably, determining which bacteria-derived metabolite(s) is responsible for this activation would be of great interest developing future postbiotic immunotherapeutic approaches, and rapidly translating these findings for cancer prevention and treatment in humans.

References:

1. World Health Organization - Cancer [Internet]. Available from: https://www.who.int/health-topics/cancer#tab=tab_1

2. Fadnes LT, Økland J-M, Haaland ØA, Johansson KA. Estimating impact of food choices on life expectancy: A modeling study. Fontana L, editor. PLOS Med [Internet]. 2022;19:e1003889. Available from: https://dx.plos.org/10.1371/journal.pmed.1003889

3. Kapsetaki SE, Marquez Alcaraz G, Maley CC, Whisner CM, Aktipis A. Diet, Microbes, and Cancer Across the Tree of Life: a Systematic Review. Curr Nutr Rep [Internet]. 2022;11:508–25. Available from: https://link.springer.com/10.1007/s13668-022-00420-5

4. Vesely MD, Kershaw MH, Schreiber RD, Smyth MJ. Natural Innate and Adaptive Immunity to Cancer. Annu Rev Immunol [Internet]. 2011 [cited 2019 Nov 2];29:235–71. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21219185

5. Liu Z, Eltoum I-EA, Guo B, Beck BH, Cloud GA, Lopez RD. Protective Immunosurveillance and Therapeutic Antitumor Activity of γδ T Cells Demonstrated in a Mouse Model of Prostate Cancer. J Immunol [Internet]. 2008;180:6044–53. Available from: http://www.jimmunol.org/lookup/doi/10.4049/jimmunol.180.9.6044

6. Hayday AC. γδ T Cell Update: Adaptate Orchestrators of Immune Surveillance. J Immunol [Internet]. 2019;203:311–20. Available from: http://www.jimmunol.org/lookup/doi/10.4049/jimmunol.1800934

7. Girard P, Charles J, Cluzel C, Degeorges E, Manches O, Plumas J, et al. The features of circulating and tumor-infiltrating γδ T cells in melanoma patients display critical perturbations with prognostic impact on clinical outcome. Oncoimmunology [Internet]. 2019;8. Available from: https://www.tandfonline.com/doi/full/10.1080/2162402X.2019.1601483

8. Gentles AJ, Newman AM, Liu CL, Bratman S V, Feng W, Kim D, et al. The prognostic landscape of genes and infiltrating immune cells across human cancers. Nat Med [Internet]. 2015;21:938–45. Available from: http://www.nature.com/articles/nm.3909

9. Saura-Esteller J, de Jong M, King LA, Ensing E, Winograd B, de Gruijl TD, et al. Gamma Delta T-Cell Based Cancer Immunotherapy: Past-Present-Future. Front Immunol [Internet]. 2022;13. Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2022.915837/full

10. Yazdanifar M, Barbarito G, Bertaina A, Airoldi I. γδ T Cells: The Ideal Tool for Cancer Immunotherapy. Cells [Internet]. 2020;9:1305. Available from: https://www.mdpi.com/2073-4409/9/5/1305

11. Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, Earley ZM, et al. Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. Science (80-). 2015;350:1084–9.

12. Zitvogel L, Ayyoub M, Routy B, Kroemer G. Microbiome and Anticancer Immunosurveillance. Cell [Internet]. 2016;165:276–87. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0092867416302562

13. Viaud S, Saccheri F, Mignot G, Yamazaki T, Daillère R, Hannani D, et al. The intestinal microbiota modulates the anticancer immune effects of cyclophosphamide. Science (80-). 2013;342.

14. Routy B, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillère R, et al. Gut microbiome influences efficacy of PD-1–based immunotherapy against epithelial tumors [Internet]. Science (80-.). 2017. Available from: http://www.sciencemag.org/lookup/doi/10.1126/science.aan3706

15. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets T V., et al. Gut microbiome modulates response to anti–PD-1 immunotherapy in melanoma patients. Science (80-) [Internet]. American Association for the Advancement of Science; 2017 [cited 2017 Nov 3];eaan4236. Available from: http://www.sciencemag.org/lookup/doi/10.1126/science.aan4236

16. Matson V, Fessler J, Bao R, Chongsuwat T, Zha Y, Alegre M-L, et al. The commensal microbiome is

associated with anti-PD-1 efficacy in metastatic melanoma patients. Science [Internet]. 2018 [cited 2018 Jan 8];359:104–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29302014

17. Ganusov V V., De Boer RJ. Do most lymphocytes in humans really reside in the gut? Trends Immunol [Internet]. 2007;28:514–8. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1471490607002311

18. Constantinides MG, Belkaid Y. Early-life imprinting of unconventional T cells and tissue homeostasis. Science (80-) [Internet]. 2021;374. Available from: https://www.science.org/doi/10.1126/science.abf0095

19. Vantourout P, Hayday A. Six-of-the-best: unique contributions of γδ T cells to immunology. Nat Rev Immunol [Internet]. 2013;13:88–100. Available from: https://www.nature.com/articles/nri3384

20. Willcox BE, Willcox CR. γδ TCR ligands: the quest to solve a 500-million-year-old mystery. Nat Immunol [Internet]. 2019;20:121–8. Available from: http://www.nature.com/articles/s41590-018-0304-y

21. Holmes E, Li J V., Marchesi JR, Nicholson JK. Gut Microbiota Composition and Activity in Relation to Host Metabolic Phenotype and Disease Risk. Cell Metab [Internet]. 2012;16:559–64. Available from: https://linkinghub.elsevier.com/retrieve/pii/S155041311200407X

22. Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ, et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. Nat Rev Gastroenterol Hepatol [Internet]. 2017;14:491–502. Available from: http://www.nature.com/articles/nrgastro.2017.75

23. Matson V, Fessler J, Bao R, Chongsuwat T, Zha Y, Alegre M-L, et al. The commensal microbiome is associated with anti–PD-1 efficacy in metastatic melanoma patients. Science (80-) [Internet]. 2018;359:104–8. Available from: https://www.science.org/doi/10.1126/science.aao3290

24. Li Y, Elmén L, Segota I, Xian Y, Tinoco R, Feng Y, et al. Prebiotic-Induced Anti-tumor Immunity Attenuates Tumor Growth. Cell Rep. 2020;30:1753-1766.e6.

25. Uerlings J, Schroyen M, Willems E, Tanghe S, Bruggeman G, Bindelle J, et al. Differential effects of inulin or its fermentation metabolites on gut barrier and immune function of porcine intestinal epithelial cells. J Funct Foods [Internet]. 2020;67:103855. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1756464620300797

26. Meyer D, Stasse-Wolthuis M. The bifidogenic effect of inulin and oligofructose and its consequences for gut health. Eur J Clin Nutr [Internet]. 2009;63:1277–89. Available from: https://www.nature.com/articles/ejcn200964

27. Belkaid Y, Harrison OJ. Homeostatic Immunity and the Microbiota. Immunity [Internet]. 2017 [cited 2018 Oct 25];46:562–76. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28423337

28. Nielsen MM, Witherden DA, Havran WL. γδ T cells in homeostasis and host defence of epithelial barrier tissues. Nat Rev Immunol [Internet]. 2017;17:733–45. Available from: http://www.nature.com/articles/nri.2017.101

29. Skwarczynski M. Inulin: A New Adjuvant With Unknown Mode of Action. EBioMedicine [Internet]. 2017;15:8–9. Available from: https://linkinghub.elsevier.com/retrieve/pii/S2352396416305230

30. Rosales-Mendoza S, Salazar-González JA, Decker EL, Reski R. Implications of plant glycans in the development of innovative vaccines. Expert Rev Vaccines [Internet]. 2016;15:915–25. Available from: https://www.tandfonline.com/doi/full/10.1586/14760584.2016.1155987

31. Viaud S, Saccheri F, Mignot G, Yamazaki T, Daillère R, Hannani D, et al. The Intestinal Microbiota Modulates the Anticancer Immune Effects of Cyclophosphamide. Science (80-) [Internet]. 2013;342:971–6. Available from: https://www.science.org/doi/10.1126/science.1240537

32. Matson V, Fessler J, Bao R, Chongsuwat T, Zha Y, Alegre M-L, et al. The commensal microbiome is associated with anti-PD-1 efficacy in metastatic melanoma patients. Science [Internet]. American

Association for the Advancement of Science; 2018 [cited 2018 Jan 30];359:104–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29302014

33. Ribas A, Shin DS, Zaretsky J, Frederiksen J, Cornish A, Avramis E, et al. PD-1 Blockade Expands Intratumoral Memory T Cells. Cancer Immunol Res [Internet]. 2016;4:194–203. Available from: https://aacrjournals.org/cancerimmunolres/article/4/3/194/468497/PD-1-Blockade-Expands-Intratumoral-Memory-T

34. Spencer CN, McQuade JL, Gopalakrishnan V, McCulloch JA, Vetizou M, Cogdill AP, et al. Dietary fiber and probiotics influence the gut microbiome and melanoma immunotherapy response. Science (80-) [Internet]. 2021;374:1632–40. Available from: https://www.science.org/doi/10.1126/science.aaz7015

35. Chen H, Cong X, Wu C, Wu X, Wang J, Mao K, et al. Intratumoral delivery of CCL25 enhances immunotherapy against triple-negative breast cancer by recruiting CCR9 + T cells. Sci Adv [Internet]. 2020;6. Available from: https://www.science.org/doi/10.1126/sciadv.aax4690

36. Simon S, Labarriere N. PD-1 expression on tumor-specific T cells: Friend or foe for immunotherapy? Oncoimmunology [Internet]. 2018;7:e1364828. Available from: https://www.tandfonline.com/doi/full/10.1080/2162402X.2017.1364828

37. Godon JJ, Zumstein E, Dabert P, Habouzit F, Moletta R. Molecular microbial diversity of an anaerobic digestor as determined by small-subunit rDNA sequence analysis. Appl Environ Microbiol [Internet]. 1997;63:2802–13. Available from: https://journals.asm.org/doi/10.1128/aem.63.7.2802-2813.1997

38. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows analysis of high-throughput community sequencing data. Nat Methods [Internet]. 2010;7:335–6. Available from: http://www.nature.com/articles/nmeth.f.303

39. Escudié F, Auer L, Bernard M, Mariadassou M, Cauquil L, Vidal K, et al. FROGS: Find, Rapidly, OTUs with Galaxy Solution. Berger B, editor. Bioinformatics [Internet]. 2018;34:1287–94. Available from: https://academic.oup.com/bioinformatics/article/34/8/1287/4708232

40. Bernard M, Rué O, Mariadassou M, Pascal G. FROGS: a powerful tool to analyse the diversity of fungi with special management of internal transcribed spacers. Brief Bioinform [Internet]. 2021;22. Available from: https://academic.oup.com/bib/article/doi/10.1093/bib/bbab318/6354026

41. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. Watson M, editor. PLoS One [Internet]. 2013;8:e61217. Available from: https://dx.plos.org/10.1371/journal.pone.0061217

Acknowledgements:

We warmly thank Sylvie Berthier (Cytometry Platform, CHUGA), Dr. Hervé Lerat, Kevin Escot and Laurie Arnaud (PHTA Animal facility), Clément Caffaratti and Hélène Coradin (TIMC Lab), and Théo Ziegelmeyer (Inserm U1209) for their help on some technical aspects.

This work (D.H.) was supported by GEFLUC Dauphiné-Savoie, Ligue contre le Cancer Comité Isère, Ligue contre le Cancer Comité Savoie, Université Grenoble Alpes IDEX Initiatives de Recherche Stratégiques. E.B. is supported by a grant salary from the French Ministry of Higher Education, Research and Innovation. The funders had no role in study design, data collection or analysis.

This work has benefited from the facilities and expertise of @BRIDGe for 16S sequencing (Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350 Jouy-en-Josas, France)

Conflicts of Interests:

Authors declare no conflict of interest

Author's contribution:

Conceptualization: DH, EB, BT

Performed experiment: DH, EB, CP

Analyzed data: DH, EB, CP, MLR, IM, AS

Wrote the paper: DH

Contribute to paper edition/writing: EB, CP, BT, DA, MLR, IM, AS

Grants acquisition: DH

Material and Methods:

Animals
Female C57Bl/6 mice (5 to 6 weeks aged) were provided by Janvier SA Laboratory (Le Genest-Saint-Isle, France) and housed at "Plateforme de Haute Technologie Animale (PHTA)" UGA core facility (Grenoble, France), EU0197, Agreement C38-51610006, under specific pathogen-free conditions, temperature-controlled environment with a 12-h light/dark cycle and ad libitum access to water and diet. Animal housing and procedures were conducted in accordance with the recommendations from the Direction des Services Vétérinaires, Ministry of Agriculture of France, according to European Communities Council Directive 2010/63/EU and according to recommendations for health monitoring from the Federation of European Laboratory Animal Science Associations. Protocols involving animals were reviewed by the local ethic committee "Comité d'Ethique pour l'Expérimentation Animale no.#12, Cometh-Grenoble" and approved by the Ministry of Research (APAFIS#12905-2018010411002729-v5). Mice were distributed in groups according to their diet. For inulin treatment, mice received standard diet and drinking water supplemented with 7.2% inulin starting 15 days before tumor inoculation. Drinking bottles supplemented with inulin were renewed 3 times a week.

Cell lines
Ovalbumin-expressing B16 melanoma (B16-OVA), MCA 205 fibrosarcoma cell lines (MCA 205 OVA), MC38 colorectal cancer cell line (MC38 OVA) were kindly provided by Pr. L. Zitvogel, Gustave Roussy (B16-OVA, MCA205-OVA) , and C. Fournier, Inserm U1209 (MC38-OVA). B16-OVA and MC38-OVA cell lines were cultured in 10% Fetal Bovine Serum (FBS) (Gibco) complete (supplemented with 1% non-essential amino acids, 1 mM sodium pyruvate, 50 U/mL penicillin and 50 µg/mL streptomycin (all from Gibco)) Dulbecco's modified Eagle's medium (DMEM) (Gibco), and MCA 205 OVA cell line was cultured in 10% FBS complete Roswell Park Memorial Institute (RPMI) (Gibco). For plasmid selection, G418 (500µg/ml, Sigma) and Hygromycin B (50µg/ml, Sigma) antibiotics were added to B16-OVA/MC38-OVA and MCA 205-OVA cultures, respectively. All cell lines were tested mycoplasma-free.

Tumor implantation
Mice received subcutaneous implantation of either: 2×10^5 B16-OVA cells, 2×10^5 MCA 205-OVA cells or 5×10^5 MC38-OVA cells in 100 µL PBS, in the right flank. Once palpable, tumors were measured using a caliper, and tumor volume determined with the following formula: $V_{tumor} = 0.5 \times (\text{width} \times \text{lenght}^2)$.

In vivo injection of anti-γδ TCR and anti-PD-1 antibodies
Mice received 100 µg of monoclonal antibody anti-γδ TCR (clone UC7-13D5, Bio X Cell) or anti-PD-1 (clone RMP1-14, Bio X Cell) intraperitoneally, in PBS. Anti- γδ TCR antibodies were injected from the day before tumor implantation, then every 2 days until the end of the experiment. Anti-PD-1 immunotherapy was administrated on days 4, 7 and 10 post-tumor implantation.

In vivo bacterial microbiota depletion
Mouse gut microbiota was depleted using a cocktail of antibiotics diluted in drinking water: Vancomycin 0.25 g/L (Mylan), Colimycin 12 x 10^6 U/L (Medac), Ampicillin 1 g/L (Sigma) and Streptomycin 5 g/L (Sigma), as previously described (13).

Tumor digestion
Tumors were collected in complete RPMI medium, lacerated with scalpels and digested with Liberase™ 2.5 mg/mL (Roche). Finally, tumors were crushed through a 70 μm cell strainer, washed and resuspended in 10% FBS complete RPMI.

Colonic IELs and LPCs isolation and CD45$^{+/-}$ cell sorting
Colons were collected in Hank's Balanced Salt Solution without Ca^{2+} and Mg^{2+} (HBSS$^-$) (Gibco), then digested twice with 5 mM HEPES, 5 mM EDTA, 1 mM DTT and 5 % FBS, for IELs isolation. Remaining colon pieces were then lacerated with scalpels and digested with collagenase VIII 0.5 mg/mL, DNAse I 10 U/mL (Sigma) and 5 % FBS in complete RPMI for LPCs isolation. Finally, digested colon pieces were crushed through a 70 μm cell strainer and cells were resuspended in 10 % FBS complete RPMI. CD45$^+$ and CD45$^-$ cells from the LPCs cell fraction were sorted using MACS (Miltenyi Biotec) with CD45 microbeads according to the manufacturer's protocol.

Lymph node digestion
Lymph nodes were collected in complete RPMI, then crush through a 70 μm cell strainer. Cells were later suspended in 10% FBS complete RPMI.

Spleen digestion
Spleens were collected in complete RPMI, then crushed through a 70 μm cell strainer and washed with 10% FBS RPMI. The pellets were then suspended in 500 μL of Red Blood Cell Lysis Buffer 1X (Sigma) and washed with 10% FBS complete RPMI.

Blood sample preparation
Blood was collected via retro-orbital sampling in K2E tubes (BD Medical). After centrifugation blood pellets were re-suspended in 1 mL Red Blood Cell Lysis Buffer 1X (Sigma) and washed with 10% FBS complete RPMI.

Flow cytometry
To allow intracellular labelling of cytokines, cell suspensions were stimulated 4 hours at 37°C with 50 ng/mL phorbol 12-myristate 13-acetate (PMA) (Sigma), 1 μg/mL ionomycin (Sigma) and Golgi Stop™ (BD Biociences). For extracellular labelling, cells were incubated with 200 ng of each antibody. Antibodies targeting extracellular proteins were: CD45 (30-F11), PD-1 (RMP1-30), (BD Biosciences); CD3 (17A2), CD4 (GK1.5), CD44 (IM7), CCR9 (9B1), CD11c (N418), CD8a (53-6.7), PD-L2 (24F.10C12), PD-L1 (10F.9G2), MHC-class II (M5/114.5.2), CD80 (16-10A1), CD40 (3/23) (Biolegend); CD11b (M1/70), γδ-TcR (eBioGL3 (GL-3, GL3); OVA-Dextramer(H-2 Kb) (Immudex).

To allow intracellular labelling, cells were first permeabilized using FoxP3 staining buffer kit (eBioscience), before intracellular labelling. Used antibodies targeting intracellular cytokines were: IFNγ (XMG1.2) (Biolegend), IL-10 (JES5-16E3), IL-17A (TC11-18H10) (BD Biosciences). After intracellular labelling, cells were fixed and stored using FACS Lysing Solution 1X (BD

Biosciences). All data were collected using BD Biosciences FACSCanto II or Lyric, and analyzed on FlowJo software.

Bacterial DNA extraction and 16S sequencing from mouse feces

Fecal samples were collected and stored at -80°C. DNA was extracted from feces following optimized JJ Godon protocole (37). After nucleic acid precipitation with isopropanol, DNA suspensions were incubated overnight at 4 °C and centrifuged at 20,000 × g for 30 min. The supernatants were transferred to a new tube containing 2 µL of RNase (RNase A, 10 mg/ml; EN0531; Fermentas, Villebon sur Yvette, France) and incubated at 37 °C for 30 min. Nucleic acids were precipitated by the addition of 1 ml of absolute ethanol and 50 µl of 3 M sodium acetate and centrifuged at 20,000 × g for 10 min. The DNA pellets were washed with 70% ethanol 3 times, dried and resuspended in 100 µl of Tris-EDTA (TE) buffer (10 mM Tris-HCl, 1 mM EDTA, adjusted pH 8). The DNA suspensions were stored at -20 °C for 16S rDNA sequences analysis.

16S sequencing data processing

Bacterial diversity was determined for each sample by targeting a portion of the ribosomal genes. PCR was performed to prepare amplicons using V3-V4 oligonucleotides (PCR1F_460: 5' CTTTCCCTACACGACGCTCTTCCGATCTACGGRAGGCAGCAG 3', PCR1R_460: 5' GGAGTTCAGACGTGTGCTCTTCCGATCTTACCAGGGTATCTAATCCT 3'), with 30 amplification cycles and an annealing temperature of 65°C. Amplicon quality was verified using gel electrophoresis, and PCR products were sent to the @BRIDGe platform for the sequencing protocol on an Illumina MiSeq (Illumina, San Diego, CA, USA).

The 16S sequences were demultiplexed and quality filtered using the QIIME version 2.1.0 software package (38). . Sequences were analyzed and normalized with the pipeline FROGS (Find Rapidly OTU with Galaxy Solution) (39,40). PCR primers were removed and sequences with sequencing error in the primers were excluded. Reads were clustered into Operational Taxonomic Units (OTUs) using the Swarm clustering method. Chimerae were removed and 580 OTUs were assigned at different taxonomic levels (from phylum to species) using RDP classifier and NCBI Blast+ on Silva_138.1_16S database. When needed, phylogeny was checked using Phylogeny browser (https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi).

35538 reads were randomly selected for each sample to normalize data. The raw reads were deposited into the SRA database (ID: PRJNA888063). The R packages "Biom" and "Phyloseq" were used for data analysis and plotting (41). Rarefaction analysis was performed and used to compare the relative abundance of OTUs across samples. Alpha diversity was measured using Observed, Chao1, Shannon, Simpson and Inverse Simpson indexes. Beta diversity was measured using the Bray-Curtis distance matrix and was used to build principal coordinates analysis (PCoA) plots. The linear discriminant analysis (LDA) effect size (LEfSe) algorithm was used to identify taxa that were specific to diet and/or treatment (39,40).

Real-time quantitative Reverse Transcription (qRT PCR)

Total RNA was extracted from IELs and LPCs cells with RNEASY-QIAGEN kit according to the manufacturer's protocol. Purity and concentration of extracted RNA were determined by reading at 260 and 280 nm in a NanoDrop spectrophotometer (Thermo Fisher). Extracted RNA was then retro-transcribed using kit Superscript II (Invitrogen). qRT-PCR analysis was performed using a SYBR Green Master Mix (Applied Biosystem) using StepOne[TM] RealTime

PCR system (Thermo Fischer). PCR conditions were: 2 minutes at 95°C, then 40 cycles of 15 seconds at 95°C and 1 minute at 60°C. The sequence-specific primers used are shown in **Table S2** (GAPDH and β2M primers are from Eurofins Scientific; other primers are from Merck). Expression levels were normalized to GAPDH and β2M endogenous gene levels and each value was then compared to a reference sample named calibrator. Finally, the relative Quantification (RQ) value was calculated.

Table S2: Sequence-specific primers used in qRT-PCR. FW = Forward sequence; RV = Reverse sequence.

Primer	Sequence (5' → 3')	Primer	Sequence (5' → 3')
GAPDH FW	GGTGAAGGTCGGTGTGAACG	MMP-9 FW	TGGGGGCAACTCGGC
GAPDH RV	CTCGCTCCTGGAAGATGGTG	MMP-9 RV	GGAATGATCTAAGCCCAG
β2M FW	GTATACTCACGCCACCCACC	iNOS FW	GTTGAAGACTGAGACTCTGG
β2M RV	TCCCGTTCTTCAGCATTTGG	iNOS RV	ACTAGGCTACTCCGTGGA
IL-1β FW	TGATGAGAATGACCTCTTCT	Cox-2 FW	GGGTTGCTGGGGGAAGAAATG
IL-1β RV	CTTCTTCAAAGATGAAGGAAA	Cox-2 RV	GGTGGCTGTTTTGGTAGGCTG
IL-6 FW	TAGTCCTTCCTACCCCAATTCC	TFF-3 FW	CCTGGTTGCTGGGTCCTCTG
IL-6 RV	TTGGTCCTTAGCCACTCCTTCC	TFF-3 RV	GCCACGGTTGTTACACTGCTC
IL-10 FW	TCCTTAATGCAGGACTTTAAGGG	Occludin FW	ACGGACCCTGACCACTATGA
IL-10 RV	GGTCTTGGAGCTTATTAAAAT	Occludin RV	TCAGCAGCAGCCATGTACTC
IL-12 FW	CCTGGGTGAGCCGACAGAAGC	ZO-1 FW	GGGGCCTACACTGATCAAGA
IL-12 RV	CCACTCCTGGAACCTAAGCAC	ZO-1 RV	TGGAGATGAGGCTTCTGCTT
IL-17 FW	GCTCCAGAAGGCCCTCAGACTACC	FoxP3 FW	CCTATGGCTCCTTCCTTGGC
IL-17 RV	CTTCCCTCCGCATTGACACAGC	FoxP3 RV	CCTTGGGTGCAGTCTTCCAG
TNF-α FW	AACTAGTGGTGCCAGCCGAT	RegIIIγ FW	TGGAGGTGGATGGGAATGGA
TNF-α RV	CTTCACAGAGCAATGACTCC	RegIIIγ RV	GCCACAGAAAGCACGGTCTA
IFNγ FW	GAACTGGCAAAAGGATGGTGA	IL-22 FW	GTGCTCAACTTCACCCTGGA
IFNγ RV	TGTGGGTTGTTGACCTCAAAC	IL-22 RV	GGCTGGAACCTGTCTGACTG
MIP-2 FW	TCAATGCCTGAAGACCCTGC	IDO1 FW	TGGGACATTCCTTCAGTGGC
MIP-2 RV	CGTCACACTCAAGCTCTGGA	IDO1 RV	TCTCGAAGCTGCCCGTTCT

A

Tumor inoculation

D-15   D0   D13

+/- Start Inulin   Tumor harvest

B

B16 OVA | **MC38 OVA** | **MCA 205 OVA**

Tumor volume (mm³) vs Days post tumor inoculation

- Control
- Inulin

B16 OVA: ****
MC38 OVA: **
MCA 205 OVA: *

C Tumor-Infiltrated Leukocytes
Positive cells among all cells(%)
CD45⁺
- Control
- Inulin
*

Tumor-Infiltrated T Lymphocytes
Positive cells among CD3⁺ (%)
CD4⁺, CD8⁺, γδ
*

D CD4⁺ T cells
Control: 12 | Inulin: 23
IFNγ vs CD3
IFNγ positive cells (%)
- Control
- Inulin
*

E CD8⁺ T cells
Control: 13 | Inulin: 28
IFNγ vs CD3
IFNγ positive cells (%)
*

F γδ T cells
Control: 8 | Inulin: 21
IFNγ vs CD3
IFNγ positive cells (%)
*

Figure 1: Inulin-enriched diet inhibits tumor growth and reinforces tumor infiltrated immunity.
(A) Experimental schedule. C57BL/6 mice were fed with a control or an inulin-enriched diet (7.2% in drinking water) (n=6 mice per group) starting 15 days before subcutaneous (s.c.) inoculation of $2x10^5$ B16 OVA melanoma cells, or $5x10^5$ MC38 OVA colorectal cancer cells, or $2x10^5$ MCA 205 OVA fibrosarcoma cells (n=6 mice per group). (B) Tumor growth curves of mice treated as described in (A) and implanted with indicated tumors. (C) Frequency of B16 OVA tumor-infiltrated total CD45⁺ and effector CD4⁺, CD8⁺ or γδ TcR⁺ T Lymphocytes from mice treated as described in (A) when control tumors reached the maximal ethic size. (D-F) Frequency of B16 OVA tumor-infiltrated IFNγ-producing cells gated on CD45⁺ CD3⁺ (D) CD4⁺, (E) CD8⁺ or (F) γδ TcR⁺ from mice treated as in (A). Graphs show the mean ± SEM. Statistically significant results are indicated by: *p < 0.05, **p < 0.005, ****p < 0.0001 by two-way ANOVA with Geisser Greenhouse's correction (B) or by Mann-Whitney tests (C-F).

Figure 2: Inulin-triggered anti-tumor effect depends on γδ T lymphocytes.
(A) Experimental schedule. C57BL/6 mice were fed with a control diet or inulin-enriched diet (7.2% in drinking water) (n=12 mice per group) starting 15 days before s.c. inoculation of $2x10^5$ B16 OVA melanoma cells. Anti-γδ TcR antibodies were injected intraperitoneally (i.p.) to 6 mice per group every two days starting the day before tumor inoculation. (B) Tumor growth curves of mice treated as described in (A). (C-E) Frequency of B16 OVA tumor-infiltrated IFNγ-producing cells gated on $CD45^+$ $CD3^+$ (C) $CD4^+$, (D) $CD8^+$ or (E) γδ TcR^+ from mice treated as described in (A) when control tumors reached the maximal ethic size. (F) Correlation between IFNγ-producing γδ T cell infiltration in tumors and the tumor growth in mice fed on a control or inulin-enriched diet (n=6 mice per group), bearing the indicated tumor. (A-E) Graphs show the mean ± SEM, and (F) IFNγ-producing γδ T cell infiltration in the tumor depending to the tumor growth. ns = not-significant, ****p < 0.0001 by two-way ANOVA with Geisser greenhouse's correction (B), by Mann-Whitney tests (C-E), or Spearman test (F).

A

D-15
+/- Start Inulin

D0
Colon harvest

B CD4⁺ T cells

Control   Inulin

IFNγ / CD3

3   3

ns

● Control
● Inulin

IFNγ positive cells (%)

C CD8⁺ T cells

Control   Inulin

IFNγ / CD3

2   2

ns

IFNγ positive cells (%)

D γδ T cells

Control   Inulin

IFNγ / CD3

5   12

*

IFNγ positive cells (%)

E

Colon

→ IELs
Analysis by
Flow Cytometry

Digestion

CD45 sorting

CD45⁺ cells   CD45⁻ cells
Lamina Propria cells   Epithelial cells

Gene expression analysis

F CD45⁻ colonic cells

TNFα
MMP9
COX2

Control mice   Inulin mice

G CD45⁺ colonic cells

FoxP3
IL-22
MIP2

Control mice   Inulin mice

Figure 3: Inulin consumption promotes γδ T IELs and mucosa cell activation in the colon

(A) Experimental schedule. C57BL/6 mice were fed with a control diet or inulin-enriched diet (7.2% in drinking water) (n=12 mice per group) for 15 days before the analysis of their gut immunity. (B-D) Frequency of IFNγ-producing IntraEpithelial Lymphocytes (IELs) gated on CD45⁺ CD3⁺ (B) CD4⁺, (C) CD8⁺, or (D) γδ TcR⁺ from mice treated as described in (A). (E) Experimental plan of colon treatment after harvest. After isolation of IELs, remaining colon pieces were digested before sorting of CD45⁻ and CD45⁺ cells for further analysis. (F-G) qRT-PCR analysis of inflammation, tissue repair and tight junction -related genes in (F) CD45⁻ and (G) CD45⁺ colon cells sorted as described in (E). (B-D) Graphs show the mean ± SEM. ns = not-significant, *p < 0.05 by Mann-Whitney tests (B-D). (F-G) Graphs show the expression levels of significantly increased genes (p < 0.05 by Mann Whitney test).

Figure 4: Inulin modulates gut microbiota and promotes *Bifidobacterium* growth.
Feces from C57BL/6 mice fed with a control (n=10 mice) or an inulin-enriched diet (7.2% in drinking water) (n=11 mice) for 15 days were collected and analysed by 16S rDNA sequencing. (A) Relative abundance of bacterial phyla (B) Bray-Curtis analysis of β-diversity, (C) cladogram representation of Linear discriminant analysis Effect Size (LEfSe) analysis, and (D) relative abundance of Bifidobacterium genus. The graph shows the mean ± SEM. ****p < 0.00001 by by Mann-Whitney test.

Figure 5: Inulin-triggered anti-tumor effect relies on intact bacterial gut microbiota.
(A) Experimental schedule. C57BL/6 mice were fed with a control diet or inulin-enriched diet (7.2% in drinking water) (n=6 mice per group) starting 15 days before s.c. inoculation of $2x10^5$ B16 OVA melanoma cells. Microbial gut microbiota was depleted from 2 days before the beginning of the diet by adding vancomycin (0.25 g/L), colimycin ($12x10^6$ U/L), ampicillin (1 g/L) and streptomycin (5 g/L) to drinking water. (B) Tumor growth curves of mice treated as described in (A). (C-E) Frequency of B16 OVA tumor-infiltrated IFNγ-producing cells gated on CD45+ CD3+ (C) CD4+, (D) CD8+, or (E) γδ TcR+ from mice treated as described in (A). (F-G) Frequency of CCR9+ cells among CD45+ CD3+ γδ TcR+ in (F) the blood and in (G) the tumor of mice fed with a control diet or inulin-enriched diet (7.2% in drinking water) (n=6 mice per group) starting 15 days before s.c. inoculation of B16 OVA melanoma cells. (H) Frequency of CCR9+ cells among CD45+ CD3+ γδ TcR+ in tumor of mice treated as described in (A). Graphs show the mean ± SEM. ns = not-significant, *$p < 0.05$ by two-way ANOVA with Geisser greenhouse's correction (B) or by Mann-Whitney tests (C-H).

Figure 6: Inulin treatment is as efficient as anti-PD-1 immunotherapy.
(A) Experimental schedule. C57BL/6 mice were fed with a control diet or inulin-enriched diet (7.2% in drinking water) (n=12 mice per group) starting 15 days before s.c. inoculation of 2×10^5 B16 OVA melanoma cells. Anti-PD-1 immunotherapy was injected i.p. to 6 mice per group on days 4, 7 and 10 post-tumor inoculation. (B) Mean and (C) individual tumor growth curves of mice treated as described in (A). (D-F) Frequency of B16 OVA tumor-infiltrated IFNγ-producing cells gated on CD45+ CD3+(C) CD4+, (D) CD8+, or (E) γδ TcR+ from mice treated as described in (A). Graphs show the mean ± SEM. ns = not-significant, *p < 0.05, **p < 0.005, ***p < 0.001 by two-way ANOVA with Geisser greenhouse's correction (B) or by Mann-Whitney tests (D-F).