Phylogenomic analysis of the understudied *Neisseriaceae* species reveals a poly- and paraphyletic

Kingella genus

Daniel P. Morreale[1,2], Joseph W. St Geme III[1,2,*], and Paul J. Planet [1,2,3,*,#]

Affiliations:

[1]Perelman School of Medicine, University of Pennsylvania, Philadelphia PA.

[2]Division of Infectious Diseases, Children's Hospital of Philadelphia, Philadelphia PA.

[3]Comparative Genomics, American Museum of Natural History, New York, New York, United States of America

*Contributed Equally

#Corresponding Author: Paul J. Planet (planetp@chop.edu)

Running title: Polyphyly in the genus *Kingella*

Keywords: *Kingella, Simonsiella, Alysiella, Neisseriaceae,* Polyphyly

Abbreviations: ANI: Average Nucleotide identity; dDDH: digital DNA-DNA hybridization, OGRI: overall genome relatedness index, COGs: clusters of orthologous gene groups; MuLDi: Multicellular, longitudinally-dividing; DUS: DNA-uptake sequences, MALDI-TOF MS: Matrix-assisted laser desorption-ionization time of flight mass spectrometry, CA: common ancestor

Abstract

Taxonomic classification and phylogenetic analysis of the *Neisseriaceae* family has focused on the

pathogens *Neisseria meningitidis* and *Neisseria gonorrhoeae*. Less is known about the

relationships of commensal *Neisseria* species and other *Neisseriaceae* genera, raising the

possibility that the phylogeny of this family may not agree with taxonomy. In this study we used

available nucleotide sequences and a phylogenetic approach to assess the *Kingella* genus and its

relatives. We found that this genus is both paraphyletic and polyphyletic. *Kingella potus* is more

closely related to *Neisseria bacilliformis* than other *Kingella* species. The *Alysiella* and *Simonsiella*

genera form a distinct clade within the *Kingella* genus that is closely related to the pathogens *K.*

kingae and *K. negevensis*. We find a phylogenetic relationship between *Conchiformibius,*

Alysiella, Simonsiella, and *Kingella,* which we name the CASK clade. Finally, we define the gene

sets that differentiate each genus of the CASK clade from one another and from the rest of the

Neisseriaceae family.

Importance

Understanding the evolutionary relationships between the species in the *Neisseriaceae* has been a

persistent challenge in bacterial systematics due to high recombination rates in these species. Previous

studies of this family have focused on *N. meningitidis* and *N. gonorrhoeae*. However, previously

understudied *Neisseriaceae* species are gaining new attention, with *K. kingae* now recognized as a

common human pathogen and with *Alysiella* and *Simonsiella* being unique in the bacterial world as

multicellular organisms. A better understanding of the genomic evolution of the *Neisseriaceae* can lead

to identification of the specific genes and traits that characterize the remarkable diversity of this family.

Introduction

Defining taxonomic relationships is a cornerstone of microbial systematics, but in species

that reproduce asexually and trade genetic information through active horizontal gene transfer

across the species boundary, defining the relationships between species can be challenging. As

in many taxonomic disciplines, the relationships amongst and within species have traditionally

been established using bacterial cell morphology and other phenotypic properties. With the

advent of molecular techniques such as DNA-DNA hybridization, 16S rRNA gene sequencing,

multilocus sequence typing (MLST), and matrix-assisted laser desorption/ionization time of flight

(MALDI-TOF), this system gave way to the so called polyphasic classification strategy that seeks

to consider both phenotypic and molecular data (1, 2). As whole genome sequences became

readily available, a newer version of this system, taxonomogenomics, seeks to incorporate this

rich genomic data into taxonomical descriptions (3–6). This approach also has the benefit of being

explicitly phylogenetic, providing a non-arbitrary approach to classification (5, 7, 8). While there

is still controversy about how best to integrate phenotypic and phylogenomic data in bacterial

taxonomy, there is a general consensus that taxonomy should be reflective of the evolutionary

relationships and that monophyly is a desirable characteristic of taxonomic groups.

The *Neisseriaceae* family is a group of Gram-negative, β-proteobacteria that currently

includes 20 genera: *Neisseria, Wielerella, Vitreoscilla, Uruburuella, Stenoxybacter, Snodgrassela,*

Simonsiella, Rivicola, Prolinoborus, Paralysiella, Morococcus, Kingella, Eikenella, Crenobacter,

Craterilacuibacter, Conchiformibius, Bergeriella, Aquella, Amantichitinum, and *Alysiella* (9–11).

The majority of prior work devoted to understanding the taxonomic relationships between these

genera has focused primarily on the pathogenic *Neisseria* species, *N. meningitidis* and *N.*

gonorrhoeae (2, 8, 12–17). Early work using 16S rRNA failed to properly resolve species

boundaries, resulting in several polyphyletic genera (18). In addition, pervasive horizontal gene

transfer (HGT) and recombination in the *Neisseriaceae* further increases taxonomic ambiguity,

with clonal lineages punctuated by large recombination events between distant relatives (14, 19).

HGT results in the formation of ill-defined species with ambiguous species boundaries (20). To

help resolve this ambiguity, recent studies of the commensal *Neisseria* species employed core

genome MLST (cgMLST) and proposed genus specific cut-offs for defining species using genome

relatedness indices (8, 19, 20). However, these studies did not comprehensively address other

polyphyletic genera within the *Neisseriaceae*.

The *Kingella* genus includes five taxa: *K. potus, K. oralis, K. denitrificans, K. negevensis,*

and *K. kingae* (21–27). Recently, two novel species have been proposed, namely *K. bonacorsii*

and K. pumchi (28, 29). All of the *Kingella* species are typically associated with the oral or

oropharyngeal microbiome, a common niche for other *Neisseriaceae*. Isolates of *K. kingae*, *K.*

negevensis, and *K. potus* have all been associated with invasive disease in immunocompetent

individuals (23, 25, 30). In contrast, *K. oralis* and *K. denitrificans* are present in dental plaque and

in gingival and periodontal samples and only rarely cause invasive disease, typically in

immunocompromised individuals (26, 31, 32). *Kingella* species are oxidase positive, catalase

negative, rods and typically form short chains (33).

A review of published phylogenetic relationships among the commensal *Neisseria* species

reveals a high level of variability in the proposed taxonomic relationships, primarily owing to the

genes selected and the method used to perform the analysis (15, 34–36). In many of these studies

the *Kingella* genus is polyphyletic, and the genera that interrupt a monophyletic clade with

Kingella species differ in different published analyses (for examples see 24, 33, 34, 37, 38). Most

isolates used for defining taxonomic nomenclature in this genus were collected prior to pervasive

whole genome sequencing and were assigned to *Kingella* as a result of phenotypic and/or

minimal sequence-based analyses. Additionally, as the frequency of culture-dependent and

culture-independent microbiome studies have become more prevalent, the number and

diversity of formally accepted species in the *Neisseriaceae* have increased (13). Here, we seek to

reassess the relatedness of the five species in the *Kingella* genus using whole genomes and a

phylogenomic approach.

Results

***Kingella potus* is distinct from the other species in the *Kingella* genus**

79   To examine the *Kingella* species, 16S rRNA gene sequences for isolates from the

Neisseriaceae were collected for phylogenetic reconstruction (Fig. 1). By 16S rRNA relatedness

alone, it is clear that a major driver of the polyphyletic structure of *Kingella* is *K. potus.* This

species was recovered from an infected wound caused by an animal bite and is more closely

related to *Neisseria bacilliformis* than it is to any currently described *Kingella* species (25).

Phenotypically, *K. potus* and *N. bacilliformis* are characterized for a limited number of

characteristics, though both species are non-hemolytic, indole-negative coccobacilli that produce

acid from glucose, maltose, and fructose (Table S3).

87   To better understand the relationship between *K. potus* and *N. bacilliformis*, we

calculated average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) between

the type strains of these species. The ANI between *K. potus* and *N. bacilliformis* was calculated at

86.75%, below the established species cut-off of 95% (39). However, the ANI between *K. potus*

and *N. bacilliformis* is appreciably greater than the ANI calculated between *K. potus* and other

Kingella species, which ranges between <75% (*K. kingae*) and 78.8% (*K. denitrificans*) (Table 1,

Fig. 2). The calculated dDDH between these species falls below species thresholds at 30% (95%

confidence interval 27.6-32.5%) (Table 2).

The *Kingella* genus is paraphyletic

To further define the relationships among *Kingella* and closely related species, whole

genome sequences for select species in the *Neisseriaceae* were downloaded from NCBI. As 16S

rRNA gene sequences have been historically used to define species in this group, all following

analyses were limited to species most closely related to *Kingella* by 16S rRNA gene phylogeny

(Fig. 1). This group includes *Neisseria elongata, Neisseria sheyganii, N. bacilliformis, Eikenella*

corrodens, Alysiella filiformis, Alysiella crassa, Simonsiella muelleri, Conchiformibius steedae, K.

oralis, K. bonacorsii, K. potus, K. denitrificans, K. kingae, and *K. negevensis. N. elongata*, *N.*

sheyganii, and *E. corrodens* were included as outgroups for these analyses. In total, 134 whole

genome sequences were downloaded from NCBI and annotated with Prokka.

To minimize the impact of HGT and variability in the pangenome, Roary was used to

identify the core genome (40). Core genes were clustered at the 75%, 80%, 85%, 90%, and 95%

sequence similarity level and were limited to only those clusters of orthologous groups (COGs)

found in all genomes. Core gene alignments were generated and used to reconstruct maximum-

likelihood phylogenies (Fig. 3). Both the Robinson-Foulds distance and normalized Nye Similarity

metrics were used to quantify tree similarity. Each phylogeny was highly congruent with the

others based on these metrics, and each included a polyphyletic clade for the genus *Kingella* (Fig.

3B and C). We therefore proceeded with the lowest similarity cutoff of 75% for further analyses

(Fig. 3A). In each of these analyses, <u>C</u>onchiformibius spp., <u>A</u>lysiella spp., <u>S</u>imonsiella spp, and

<u>K</u>ingella spp. (with the exception of *K. potus*) form a closely related group that we will refer to as

the CASK clade. Phenotypic data on the species in the CASK clade are extremely limited, limiting

the utility of phenotypic properties in understanding relationships within the group (Table S3).

Based on a whole genome approach, *Alysiella* and *Simonsiella* form a clade within the

Kingella genus (Fig. 3A), dividing *K. oralis* and *K. denitrificans* from *K. negevensis* and *K. kingae.*

ANI, dDDH, and G+C% were calculated for type strains of *Kingella*, *Alysiella*, and *Simonsiella* and

are shown in Tables 1 and 3. All of these overall genome relatedness indices (OGRI) support a

close relationship between the species of these three genera*.*

Additionally, we calculated pairwise ANI scores for each of the genomes listed above. The

ANI scores confirm the majority of the proposed species classifications except for *K. bonacorsii*,

an unclassified *Kingella* isolate that was recently proposed to be a new species (28)*.* This isolate

was recovered from a gingival sample, and the authors report a 16S rRNA similarity of 98.7%, ANI

of 95.83%, and dDDH of 63.6% relative to *K. oralis* UB-38, as well as non-identification by MALDI-

TOF mass-spectrometry (28). We replicated each of the genomic calculations reported as well as

additional dDDH calculations using alternative formulae and found that choice of formula is

essential, as only *d4* yielded a dDDH below the accepted values for species cutoffs (Tables 1 and

4, Fig. 2) (39, 41)*.*

Kingella species are unified by gene presence/absence in the pangenome

Finally, another recent evolutionary study of these taxa has used less stringent similarity

and gene presence cutoffs to define the core genome for species in the *Neisseriaceae* for

subsequent phylogenetic analyses (34). We replicated this analysis by reanalyzing the data using

a sequence similarity cut off of 50% for genes present in 80% of analyzed genomes. By lowering

the threshold for a gene to be considered part of the core genome, we increase the number of

genes considered from 109 at the 75% similarity level up to 620 genes. Notably, as we increased

the number of genes considered part of the core genome by decreasing the stringency of our

Roary analysis, the presence of a polyphyletic *Kingella* clade remains constant (Fig. 4A),

demonstrating the robust relationships between these genera that are obtained when we

consider the core genome of species in the *Neisseriaceae*.

To better understand how the pangenome impacts the relationships described in the

above analyses, we reconstructed the phylogeny considering only gene presence and absence in

the pangenome. This analysis was performed with a binary COG presence/absence matrix as

defined by Roary with a 75% similarity cut-off with phylogenetic inference in IQ-Tree. For the first

time in any of our analyses, all named *Kingella* species except for *K. potus* are monophyletic,

excluding the *Simonsiella* and *Alysiella* genomes. To identify genes that drive the unification of

Kingella sensu stricto (*Kingella s.s.*), IQ-Tree was used to perform ancestral reconstructions of the

species in this tree. These ancestral reconstructions facilitate the identification of the genes that

drive the unification of *Kingella* into a monophyletic clade. Discounting statistically ambiguous

sites, monophyly is driven by the loss of eight genes in *Kingella s.s.* (Table 4), all of which fall in

the accessory genome. There are no gene gains that support this clade, and therefore this

monophyly is not well supported.

These data further suggest that horizontal gene transfer (HGT) can confound our ability to

define species relationships and that limits on HGT may be important to consider when defining

species in the *Neisseriaceae.* HGT is driven by DNA uptake sequences (DUS) in the *Neisseriaceae*

(42–44), which are largely thought to be species specific. The DUS in the CASK clade have been

previously characterized (42). Type genomes for each species were used to define DUS

preference by word search. *Alysiella filliformis*, *Alysiella crassa*, *Simonsiella muelleri*, and *Kingella*

oralis genomes are enriched for the AG-SimDUS (AGGCTGCCTGAA) and the AG-KingDUS

(AGGCAGCCTGAA). *Kingella kingae, Kingella negevensis,* and *Kingella denitrificans* genomes are

enriched for the AA-King3DUS (AAGCAGCCTGCA). *Conchiformibius steedae* and *Conchiformibius*

kuhniae do not appear to use any of these previously defined DUS.

The CASK species are unique based on cellular morphology and lifestyle.

To identify the genes that differentiate the species in the CASK clade and to define clusters of

orthologous groups (COGs) unique to groups of interest, Scoary was used to analyze the pangenome

analysis generated by Roary at the 75% sequence similarity level. The presence of genes is defined as

diagnostic for the in-group if they are 100% specific, are 100% sensitive, and have a *p* > 0.001. These

cutoffs reduce the impact of shared mobile genetic elements (MGEs) such as prophages or genomic

islands.

First, the CASK clade was compared against the rest of the species included in the phylogenetic

analyses (Fig. 5A, table S5). This clade is enriched for 1360 COGs that are absent in the rest of the analyzed

species, providing strong evidence supporting the evolutionary relatedness of this group. The average

genome size for the species in this group is approximately 2,000 coding sequences (CDS). To understand

putative roles of these COGs in the CASK species, COG groups and Gene Ontology (GO) functional

annotations were defined with eggnog mapper (Fig. 5B). Approximately half of the COGs were assigned

to the "function unknown" group (n = 656). Of the remaining COGs, the majority play putative roles in cell

wall, membrane, and envelope biogenesis (n = 276); DNA replication, recombination, and repair (n = 224);

or amino acid transport and metabolism (n = 205).

Next, *Kingella sensu stricto (Kingella s.s.)*, comprised of *K. oralis, K. denitrificans, K. negevensis,*

and *K. kingae*, was compared to the rest of the CASK clade. Only 67 COGs meet the criteria to be diagnostic

for *Kingella s.s* (Table S6)*. When *Alysiella* and *Simonsiella* or the pathogenic *Kingella* spp. were compared

to the rest of CASK clade, we identified only 41 or 79 diagnostic COGs, respectively (Fig. 5A). Again, we

assigned putative functional roles for the COGs in each of these groups (Fig. 5C). The COGs diagnostic for

Kingella s.s. and the pathogenic *Kingella* spp (*K. negevensis* and *K. kingae*) are primarily involved in

carbohydrate transport and metabolism or nucleotide transport and metabolism. By contrast, *Alysiella*

and *Simonsiella* diagnostic COGs are involved in DNA replication, recombination, and repair or cell wall,

membrane, and envelope biogenesis (Fig. 5C).

Finally, we compared the genomes of the type-strain for each species in the CASK clade using

BLASTn against *K. kingae* ATCC 23330 (Fig. 5D). We found high levels of nucleotide sequence similarity

across the *K. kingae* genome relative to the related species. There are few notable regions of dissimilarity,

which appear to be MGEs in the *K. kingae* genome.

Discussion

Classification of the species in the *Neisseriaceae* has been difficult using traditional metrics of

relatedness, both phenotypically and phylogenetically. As a result, the family has been restructured

several times since its inception to include or exclude genera (15, 24, 36, 45–47). Species in this family

resist robust phylogenetic classification due to pervasive horizontal gene transfer and recombination,

including in the 16S rRNA sequence (18–20, 42). Due to the large diversity of species and variability,

phenotypic classification using strict phenotypic classification is either difficult or impossible without

additional sampling. For example, presence of catalase is conserved among *Neisseria* isolates, and yet

isolates of *N. elongata* have been recovered that lack the catalase enzyme(33). Furthermore, as more

species are proposed using culture-independent methods or based on a single isolate, it will be essential

to establish phylogenomic species definitions for the *Neisseriaceae* (1, 18, 20)*.*

In this study, we employed a phylogenetic approach to define the evolutionary relationships

within a major subgroup of the *Neisseriaceae*, defined by the genera *Eikenella, Conchiformibius, Alysiella,*

Simonsiella, Kingella, and select *Neisseria* species. Examination by 16S rRNA sequence unveiled a very

recent common ancestor between *K. potus* and *N. bacilliformis*, which was further supported by several

other genome relatedness indices (Fig. 1, Tables 1,3). *K. potus* is a catalase-negative species collected from

an infected wound that resulted from a bite by a captive South American Kinkajou (*Potus flavus*) (25). *N.*

bacilliformis is an opportunistic pathogen that has been recovered from human infections, varies in the

production of catalase (48), and is phenotypically identical to *N. elongata* subsp. *elongata.* These species

were reported nearly simultaneously, preventing their direct comparison. Our reanalysis suggests that *K.*

potus is much more closely related to *N. bacilliformis* than to any other *Kingella* species, though it is likely

a distinct species from *N. bacilliformis*. Inclusion in the genus *Kingella* was likely due to a negative catalase

test, underscoring the importance of a genomic approach to classification of these isolates.

The phylogenomic analyses also uncovered close evolutionary relationships between

<u>C</u>onchiformibius sp., <u>A</u>lysiella sp.*,* <u>S</u>imonsiella sp., and <u>K</u>ingella sp.*,* which we have named the

"CASK" clade (Fig. 3). The CASK clade is a major branch of the *Neisseriaceae* that includes both

invasive pathogens (*K. negevensis* and *K. kingae*) and oral-associated commensals. *Alysiella* and

Simonsiella are genera associated with the oral microbiome of non-human mammals (15, 47, 49).

Conchiformibius spp., *Alysiella* spp., and *Simonsiella* spp. are remarkable for their multicellularity,

which is the result of incomplete septation by the forming peptidoglycan during bacterial division

(34).

The reproducibility of the core genome phylogenetic structure very strongly supports the

close relationship between the species of the CASK clade and the exclusion of *K. potus* (Fig. 3-4).

We hypothesized that prior misclassification could be due to high levels of horizontal gene

transfer of accessory genes between *Kingella* (*sensu stricto*) genomes. To test this hypothesis, we

generated a presence/absence matrix generated by Roary at 75% sequence similarity level, a

technique that emphasizes accessory genes over core genes, and we reconstructed a phylogeny

using the RAxML binary model with gamma model of rate heterogeneity and bootstrapped 100

times (Fig. 3B). This was the only analysis in which we observed a monophyletic *Kingella* genus,

though *Kingella potus* was still excluded. We favor the analyses that are based on nucleotide

sequence of the core genome because they contain much more genomic information (~76,500

variable sites compared to ~29,000 variable genes) and likely lessen confounding signal from

recombined areas. A separate whole gene-based analysis also identified more diagnostic genes

uniting *Alysiella*, *Simonsiella,* and the pathogenic *Kingella* species (n=79) than genes uniting the

Kingella genus *sensu strictu* (n=69). Additionally, as *Simonsiella*, *K. oralis*, and *K. kingae* utilize

specific DNA-uptake sequences (DUS), suggesting they may be able to share DNA more easily

among themselves as compared to the rest of the *Neisseriaceae*. Given that members of the

Neisseriaceae are highly recombinogenic and undergo significant HGT of accessory genes, we

believe that the core genome is a higher fidelity evolutionary record for this family (20).

The CASK clade has a large set of diagnostic genes that differentiate these species from

the remainder of the *Neisseriaceae*. Interestingly, this group is enriched for processes that impact

cell wall and envelope biogenesis (Fig. 5B), likely driven by the unusual cellular morphologies of

Conchiformibius spp., *Alysiella* spp., and *Simonsiella* spp. These genera are distinctive for being

multicellular, longitudinally-dividing (MuLDi) species that comprise the only known animal

multicellular bacterial symbionts (34). Nyongesa and co-workers hypothesize that the MuLDi

phenotype evolved twice in the *Neisseriaceae,* first in *Conchiformibius* species and then in the

common ancestor of *Alysiella* and *Simonsiella*, a hypothesis that is strongly supported by our

phylogenetic analyses. This hypothesis is further supported by consideration of the diagnostic

genes that differentiate *Alysiella* and *Simonsiella* from the rest of this group (Fig. 5C). We again

find an enrichment for COGs implicated in cellular morphology, as we would expect if these

genera evolved the MuLDi phenotype independently of *Conchiformibius* spp. By contrast,

Kingella s.s. is enriched for COGs associated with nutrient transport and metabolism (Fig. 5C).

Our study suggests that the current taxonomic classifications of species in the *Kingella* genus

may not reflect evolutionary history. Based on our results, *K. potus* is more related to *N.*

bacilliformis than to other *Kingella* species by 16S rRNA, OGRI, and whole genome analyses and

should probably be assigned to the genus *Neisseria* or another genus entirely. Additionally,

phylogenomic analysis of the core genomes of *Alysiella* and *Simonsiella* indicates their close

relationship to the *Kingella* genus and supports their reassignment to *Kingella*, resulting in a

monophyletic genus.

Methods

Phylogenetic Analysis

840 of the available 16S rRNA gene sequences for isolates from the *Neisseriaceae* and in

the RefNR database were downloaded from the SILVA r138.1 database on September 08, 2022

(50, 51). 16S sequence accession numbers are listed in table S1. 16S rRNA sequences were aligned

with MAFFT, and a phylogeny was reconstructed using RAxML v. 8.4.2 with the general time

reversible model of nucleotide substitution and gamma model of rate heterogeneity, and

bootstrapped 100 times (52, 53).

Whole genome sequences for *Neisseria elongata, Neisseria sheyganii, N. bacilliformis,*

Eikenella corrodens, Alysiella filiformis, Alysiella crassa, Simonsiella muelleri, Conchiformibius

steedae, K. oralis, K. bonacorsii, K. potus, K. denitrificans, K. kingae, and *K. negevensis* were

downloaded from the NCBI Genome database on September 09, 2022. Sequence accession

numbers are listed in table S2*.* In total, 134 genomes were reassembled with Prokka v1.14.6,

using default settings for bacterial sequences (54). Phenotypic characteristics for each of these

species were collected by literature review and are listed in table S3. Assemblies were analyzed

with Roary v3.13.0. to identify the core genome (40). Core genes were clustered at the 75%, 80%,

85%, 90%, and 95% sequence similarity level and were limited to only those clusters of

orthologous groups (COGs) found in all genomes (-cd 100), unless otherwise noted. Additionally,

other recent evolutionary study of these taxa has used less stringent similarity and gene presence

cutoff to define the core genome for species in the *Neisseriaceae* for subsequent phylogenetic

analyses (8, 34). We replicated this analysis by reanalyzing the data using a sequence similarity

cut off of 50% for genes present in 99% or 80% of analyzed genomes. Nucleotide sequences of

core genes were compared in a codon-aware alignment by PRANK within Roary, and core gene

alignments were used to reconstruct phylogenies using RAxML as above. Phylogenetic trees were

annotated using Figtree v.1.4.4 and ITOL (55). Bootstrap values >70 are shown.

To calculate phylogenetic tree similarity, final trees were analyzed in R using the Ape v5.6-

2 (56), ggtree v3.1.2. (57), and TreeDist v.2.5.0 (58, 59) packages . The Robinson-Fould distance

and normalized Nye Simillarity Score were determined for each pair of trees (58–61).

Ancestral reconstructions were performed using the Roary gene presence/absence (PA)

matrix generated at the 75% sequence similarity cut-off. The matrix was converted into a FASTA

formatted file of binary data using R. This file was used for phylogenetic reconstruction in IQ-Tree

v 1.6.12 with automatic model calling (62, 63). Branch support was determined via ultrafast

bootstrapping (-bb 1000) (64), and ancestral states (-asr) were calculated with default cut-offs.

Full inferred ancestral state matrices are available in Supplementary File 2. To determine the

COGs that unite *Kingella s.s.*, the inferred PA matrix of common ancestor of *Alysiella* spp.,

Simonsiella spp., and *Kingella s.s.* (Node 7), was compared to the inferred PA matrix of common

ancestor of *Kingella* spp. Statistically ambiguous sites present in either PA matrix were removed

from this analysis.

Gene Set Enrichment

Scoary v 1.6.16 was used to further analyze the gene presence absence matrix generated

by Roary at the 75% sequence similarity cutoff (65). The -r flag was used to limit the set of strains

considered in this analysis. COGs were eliminated from this analysis if the Bonferroni-corrected

p-value >0.0001. Diagnostic genes are only those genes which 1) meet this statistical threshold,

2) are enriched in the clade of interest, and 3) have specificity and sensitivity of >80%.

To identify pathways that are enriched in each phylogenetic group, the representative

COG sequence was collected from the Roary output files and compiled into a gene set. Gene sets

were analyzed with the COG database using eggNOG-mapper v 2.1.9 (66, 67), with a DIAMOND

alignment (68) and default settings for bacterial species (Table S4-9). Results were plotted using

Graphpad Prism.

Genome Relatedness Indices

Average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) between the

type strains of these species using FastANI v.1.33 and the Type Strain Genome Server (TYGS),

respectively (69–71).

References

1. Ramasamy D, Mishra AK, Lagier JC, Padhmanabhan R, Rossi M, Sentausa E, Raoult D, Fournier PE.

2014. A polyphasic strategy incorporating genomic data for the taxonomic description of novel

bacterial species. Int J Syst Evol Microbiol 64:384–391.

2. Vandamme P, Pot B, Gillis M, Vos P de, Kersters K, Swings J. 1996. Polyphasic taxonomy, a

consensus approach to bacterial systematics. Microbiol Rev 60:407.

3. Fournier PE, Lagier JC, Dubourg G, Raoult D. 2015. From culturomics to taxonomogenomics: A

need to change the taxonomy of prokaryotes in clinical microbiology. Anaerobe 36:73–78.

4. Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species

definition. Proc Natl Acad Sci U S A 106:19126.

5. Chung M, Munro JB, Tettelin H, Hotopp JCD. 2018. Using Core Genome Alignments To Assign

Bacterial Species. mSystems 3.

6. Caputo A, Fournier PE, Raoult D. 2019. Genome and pan-genome analysis to classify emerging

bacteria. Biol Direct 14.

7. Moldovan MA, Gelfand MS. 2018. Pangenomic definition of prokaryotic species and the

phylogenetic structure of *Prochlorococcus spp.* Front Microbiol 9:428.

8. Khoder M, Osman M, Kassem II, Rafei R, Shahin A, Fournier PE, Rolain J-M, Hamze M. 2022.

Whole Genome Analyses Accurately Identify *Neisseria* spp. and Limit Taxonomic Ambiguity. Int J

Mol Sci 23:13456.

9. Parte AC, Carbasse JS, Meier-Kolthoff JP, Reimer LC, Göker M. 2020. List of prokaryotic names

with standing in nomenclature (LPSN) moves to the DSMZ. Int J Syst Evol Microbiol 70:5607–

5612.

10. Parker CT, Tindall BJ, Garrity GM. 2019. International code of nomenclature of Prokaryotes. Int J

Syst Evol Microbiol 69:S1.

11.　Oren A, Garrity G. 2022. Notification of changes in taxonomic opinion previously published

outside the IJSEM. List of changes in taxonomic opinion no. 36. Int J Syst Evol Microbiol

72:005413.

12.　Bennett JS, Jolley KA, Earle SG, Corton C, Bentley SD, Parkhill J, Maiden MCJ. 2012. A genomic

approach to bacterial taxonomy: An examination and proposed reclassification of species within

the genus *Neisseria*. Microbiol (United Kingdom) 158:1570–1580.

13.　Maiden MCJ, Harrison OB. 2016. Population and functional genomics of *Neisseria* revealed with

gene-by-gene approaches. J Clin Microbiol 54:1949–1955.

14.　Vidovic S, Horsman GB, Liao M, Dillon JAR. 2011. Influence of Conserved and Hypervariable

Genetic Markers on Genotyping Circulating Strains of *Neisseria gonorrhoeae.* PLoS One 6:e28259.

15.　Adeolu M, Gupta RS. 2013. Phylogenomics and molecular signatures for the order *Neisseriales*:

proposal for division of the order Neisseriales into the emended family *Neisseriaceae* and

Chromobacteriaceae fam. nov. Antonie Van Leeuwenhoek 104:1–24.

16.　Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA

hybridization values and their relationship to whole-genome sequence similarities. Int J Syst Evol

Microbiol 57:81–91.

17.　Diallo K, MacLennan J, Harrison OB, Msefula C, Sow SO, Daugla DM, Johnson E, Trotter C,

MacLennan CA, Parkhill J, Borrow R, Greenwood BM, Maiden MCJ. 2019. Genomic

characterization of novel *Neisseria* species. Sci Rep 9.

18.　Rossi-Tamisier M, Benamar S, Raoult D, Fournier PE. 2015. Cautionary tale of using 16s rRNA

gene sequence similarity values in identification of human-associated bacterial species. Int J Syst

Evol Microbiol 65:1929–1934.

19.　Tibayrenc M, Ayala FJ. 2015. How clonal are *Neisseria* species? The epidemic clonality model

revisited. Proc Natl Acad Sci U S A 112:8909.

20.   Hanage WP, Fraser C, Spratt BG. 2005. Fuzzy species among recombinogenic bacteria. BMC Biol

3.

21.   Dewhirst FE, Paster BJ, Bright PL. 1989. *Chromobacterium*, *Eikenella*, *Kingella*, *Neisseria*,

Simensiella, and *Vitreoscilla* species comprise a major branch of the beta group Proteobacteria by

16S ribosomal ribonucleic acid sequence comparison: Transfer of *Eikenella* and *Simonsiella* to the

family *Neisseriaceae* (emend.). Int J Syst Bacteriol 39:258–266.

22.   Henricksen SD, Bovre K. 1976. Transfer of *Moraxella kingae* Henriksen and Bovre to the genus

Kingella gen. nov. in the family *Neisseriaceae*. Int J Syst Bacteriol 26:447–450.

23.   El Houmami N, Bakour S, Bzdrenga J, Rathored J, Seligmann H, Robert C, Armstrong N, Schrenzel

J, Raoult D, Yagupsky P, Fournier PE. 2017. Isolation and characterization of *Kingella negevensis*

sp. nov., a novel *Kingella* species detected in a healthy paediatric population. Int J Syst Evol

Microbiol 67:2370–2376.

24.   Dewhirst FE, Chen CKC, Paster BJ, Zambon JJ. 1993. Phylogeny of species in the family

Neisseriaceae isolated from human dental plaque and description of K*ingella oralis sp. nov*

[corrected]. Int J Syst Bacteriol 43:490–499.

25.   Lawson PA, Malnick H, Collins MD, Shah JJ, Chattaway MA, Bendall R, Hartley JW. 2005.

Description of *Kingella potus sp. nov.*, an Organism Isolated from a Wound Caused by an Animal

Bite. J Clin Microbiol 43:3526.

26.   Snell JJS, Lapage SP. 1976. Transfer of some saccharolytic *Moraxella* species to *Kingella Henriksen*

and Bovre 1976, with descriptions of *Kingella indologenes sp. nov.* and *Kingella denitrificans sp.*

nov. Int J Syst Bacteriol 26:451–458.

27.   Hollis DG, Weaver RE, Riley PS. 1983. Emended description of *Kingella denitrificans* (Snell and

Lapage 1976): correction of the maltose reaction. J Clin Microbiol 18:1174.

28.   Antezack A, Boxberger M, Rolland C, Monnet-Corti V, La Scola B. 2021. Isolation and

392   Characterization of *Kingella bonacorsii sp. nov.,* A Novel *Kingella* Species Detected in a Stable

393   Periodontitis Subject. Pathogens 10:1–13.

394 29. Xiao M, Liu R, Du J, Liu R, Zhai L, Wang H, Yao S, Xu YC. 2022. *Kingella pumchi sp. nov.*, an

395   organism isolated from human vertebral puncture tissue. Antonie Van Leeuwenhoek

396   https://doi.org/10.1007/S10482-022-01786-W.

397 30. Yagupsky P. 2015. *Kingella kingae:* Carriage, transmission, and disease. Clin Microbiol Rev.

398   American Society for Microbiology https://doi.org/10.1128/CMR.00028-14.

399 31. Chen C, Chen C. 1996. Distribution of a newly described species, *Kingella oralis*, in the human oral

400   cavity. Oral Microbiol Immunol 11:425–427.

401 32. Kopyt N, Kumar A, Agrawal V. 2015. A Case of Suppurative Peritonitis by a Commensal Oral

402   Organism, *Kingella denitrificans*, in an Adult Peritoneal Dialysis Patient. Perit Dial Int 35:105.

403 33. Tønjum T. 2005. Order IV. *Neisseriales ord. nov.*, p. 575–922. *In* Garrity, GM, Bell, JA, Lilburn, T

404   (eds.), Bergey's Manual of Systematic Bacteriology, 2nd ed. Springer, Boston.

405 34. Nyongesa S, Weber PM, Bernet È, Pulido F, Nieves C, Nieckarz M, Delaby M, Viehboeck T, Krause

406   N, Rivera-Millot A, Nakamura A, Vischer NOE, vanNieuwenhze M, Brun Y V., Cava F, Bulgheresi S,

407   Veyrier FJ. 2022. Evolution of longitudinal division in multicellular bacteria of the *Neisseriaceae*

408   family. Nat Commun 2022 131 13:1–18.

409 35. Garrity GM, Bell JA, Lilburn T. 2005. Class II. Betaproteobacteria class. nov. Bergey's Manual® Syst

410   Bacteriol 575–922.

411 36. Enright MC, Carter PE, MacLean IA, McKenzie H. 1994. Phylogenetic relationships between some

412   members of the genera *Neisseria*, *Acinetobacter*, *Moraxella*, and *Kingella* based on partial 16S

413   ribosomal DNA sequence analysis. Int J Syst Bacteriol 44:387–391.

414 37. Adeolu M, Gupta RS. 2013. Phylogenomics and molecular signatures for the order *Neisseriales*:

415   proposal for division of the order *Neisseriales* into the emended family *Neisseriaceae* and

Chromobacteriaceae fam. nov. Antonie Van Leeuwenhoek 104:1–24.

38.   Lawson PA, Malnick H, Collins MD, Shah JJ, Chattaway MA, Bendall R, Hartley JW. 2005.

Description of *Kingella potus sp. nov.*, an Organism Isolated from a Wound Caused by an Animal

Bite. J Clin Microbiol 43:3526.

39.   Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. 2013. Genome sequence-based species

delimitation with confidence intervals and improved distance functions. BMC Bioinformatics

14:1–14.

40.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D, Keane JA,

Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics

31:3691–3693.

41.   Auch AF, von Jan M, Klenk HP, Göker M. 2010. Digital DNA-DNA hybridization for microbial

species delineation by means of genome-to-genome sequence comparison. Stand Genomic Sci

2:117.

42.   Frye SA, Nilsen M, Tønjum T, Ambur OH. 2013. Dialects of the DNA Uptake Sequence in

Neisseriaceae. PLOS Genet 9:e1003458.

43.   Weir S, Lee LIW, Marrs C. 1996. Identification of four complete type 4 pilin genes in a single

Kingella denitrificans genome. Infect Immun 64:4993–4999.

44.   Treangen TJ, Ambur O, Tonjum T, Rocha EPC. 2008. The impact of the *Neisserial* DNA uptake

sequences on genome evolution and stability. Genome Biol 9.

45.   Xie CH, Yokota A. 2005. Phylogenetic analysis of *Alysiella* and related genera of *Neisseriaceae*:

proposal of *Alysiella crassa comb. nov., Conchiformibium steedae gen. nov., comb. nov.,*

Conchiformibium kuhniae sp. nov. and *Bergeriella denitrificans gen. nov., comb. nov.* J Gen Appl

Microbiol 51:1–10.

46.   Kuhnert P, Brodard I, Bock S, Hemphill A, Akarsu H, Engelhardt A, Kutzer P. 2022. *Wielerella bovis*

gen. nov., sp. nov. a member of the family *Neisseriaceae* associated with bovine endocarditis. Int

J Syst Evol Microbiol 72.

47. Xie CH, Yokota A. 2005. Phylogenetic analysis of *Alysiella* and related genera of *Neisseriaceae*:

proposal of A*lysiella crassa comb. nov., Conchiformibium steedae gen. nov., comb. nov.,*

Conchiformibium kuhniae sp. nov. and *Bergeriella denitrificans gen. nov., comb. nov.* J Gen Appl

Microbiol 51:1–10.

48. Han XY, Hong T, Falsen E. 2006. *Neisseria bacilliformis sp. nov.* Isolated from Human Infections. J

Clin Microbiol 44:474.

49. Hedlund BP, Tønjum T. 2015. *Simonsiella* . Bergey's Man Syst Archaea Bact 1–12.

50. Glöckner FO, Yilmaz P, Quast C, Gerken J, Beccati A, Ciuprina A, Bruns G, Yarza P, Peplies J,

Westram R, Ludwig W. 2017. 25 years of serving the community with ribosomal RNA gene

reference databases and tools. J Biotechnol 261:169–176.

51. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The

SILVA ribosomal RNA gene database project: improved data processing and web-based tools.

Nucleic Acids Res 41:D590–D596.

52. Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics 30:1312.

53. Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol Biol Evol 30:772.

54. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

55. Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. Nucleic Acids Res 49:W293–W296.

56. Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. Bioinformatics 35:526–528.

57.   Yu G. 2020. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc Bioinforma

465   69:e96.

466   58.   Smith MR. 2022. Robust Analysis of Phylogenetic Tree Space. Syst Biol 71:1255–1270.

467   59.   Smith MR. 2020. Information theoretic generalized Robinson-Foulds metrics for comparing

468   phylogenetic trees. Bioinformatics 36:5007–5013.

469   60.   Robinson DF, Foulds LR. 1981. Comparison of phylogenetic trees. Math Biosci 53:131–147.

470   61.   Nye TMW, Liò P, Gilks WR. 2006. A novel algorithm and web-based tool for comparing two

471   alternative phylogenetic trees. Bioinformatics 22:117–119.

472   62.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, Lanfear R,

473   Teeling E. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

474   Genomic Era. Mol Biol Evol 37:1530–1534.

475   63.   Wang HC, Minh BQ, Susko E, Roger AJ. 2018. Modeling Site Heterogeneity with Posterior Mean

476   Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. Syst Biol 67:216–235.

477   64.   Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the

478   Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

479   65.   Brynildsrud O, Bohlin J, Scheffer L, Eldholm V. 2016. Rapid scoring of genes in microbial pan-

480   genome-wide association studies with Scoary. Genome Biol 17:1–9.

481   66.   Cantalapiedra CP, Herṇandez-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-mapper

482   v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic

483   Scale. Mol Biol Evol 38:5825–5829.

484   67.   Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR,

485   Letunic I, Rattei T, Jensen LJ, Von Mering C, Bork P. 2019. eggNOG 5.0: a hierarchical, functionally

486   and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses.

487   Nucleic Acids Res 47:D309–D314.

68. Buchfink B, Reuter K, Drost HG. 2021. Sensitive protein alignments at tree-of-life scale using

DIAMOND. Nat Methods 2021 184 18:366–368.

69. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun 2018 91 9:1–

8.

70. Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Göker M. 2022. TYGS and LPSN: a database

tandem for fast and reliable genome-based classification and nomenclature of prokaryotes.

Nucleic Acids Res 50:D801–D807.

71. Meier-Kolthoff JP, Göker M. 2019. TYGS is an automated high-throughput platform for state-of-

the-art genome-based taxonomy. Nat Commun 2019 101 10:1–10.

72. Kuhn DA, Gregory DA, Buchanan GE, Nyby MD, Daly KR. 1978. Isolation, characterization, and

numerical taxonomy of *Simonsiella* strains from the oral cavities of cats, dogs, sheep, and

humans. Arch Microbiol 118:235–241.

73. Yagupsky P, Peled N, Katz O. 2002. Epidemiological Features of Invasive *Kingella kingae* Infections

and Respiratory Carriage of the Organism. J Clin Microbiol 40:4180.

74. Hedlund BP, Kuhn DA. 2006. The Genera *Simonsiella* and *Alysiella*. The Prokaryotes 828–839.

75. Hedlund BP, Staley JT. 2002. Phylogeny of the genus *Simonsiella* and other members of the

Neisseriaceae. Int J Syst Evol Microbiol 52:1377–1382.

76. Andersen BM, Weyant RS, Steigerwalt AG, Moss CW, Hollis DG, Weaver RE, Ashford D, Brenner

DJ. 1995. Characterization of *Neisseria elongata subsp. glycolytica* isolates obtained from human

wound specimens and blood cultures. J Clin Microbiol 33:76.

77. Kuhn DA, Gregory DA, Buchanan GE, Nyby MD, Daly KR. 1978. Isolation, characterization, and

numerical taxonomy of *Simonsiella* strains from the oral cavities of cats, dogs, sheep, and

humans. Arch Microbiol 118:235–241.

78.   Bernard KA, Burdz T, Wiebe D, Bernier AM. 2020. Description of *Eikenella halliae sp. nov.* and

513   *Eikenella longinqua sp. nov.,* derived from human clinical materials, emendation of *Eikenella*

514   *exigua* Stormo et al. 2019 and emendation of the genus *Eikenella* to include species which are

515   strict anaerobes. Int J Syst Evol Microbiol 70:3167–3178.

Tables.

Supplemental Tables are attached.

Table 1. Average ANI scores calculated among taxa of interest. ANI scores were calculated using FastANI and are shown only if the ANI >75%. Empty cells denote data excluded from the table.

	N. bacilliformis	K. denitrificans	K. oralis	K. negevensis	K. kingae	N. elongata	A. crassa	A. filiformis	S. muelleri
K. potus	86.75	78.88	78.74	<75	<75	80.7	-	-	-
K. bonacorsii	-	79.84	96.41	78.06	77.9	78.76	77.98	77.87	77.14
A. crassa	-	77.81	77.96	79.37	79.74	<75	100	85.43	79.79
A. filiformis	-	77.81	77.96	79.37	79.74	<75	85.43	100	78.75
S. muelleri	-	<75	76.9	79.16	79.96	<75	79.79	78.75	100

523 **Table 2. dDDH values of *K. potus* calculated against closely related taxa.** dDDH was calculated
524 using TYGS using each of the possible algorithms, as well as the difference in G+C content
525 between *K. potus* and each of the subject type strains.

Query strain	Subject strain	dDDH (d0, in %)	C.I. (d0, in %)	dDDH (d4, in %)	C.I. (d4, in %)	dDDH (d6, in %)	C.I. (d6, in %)	G+C content difference (in %)
K. potus NCTC13336	*N. bacilliformis* ATCC BAA-1200	45.6	[42.2 - 49.0]	30	[27.6 - 32.5]	41	[38.0 - 44.0]	1.86
K. potus NCTC13336	*N. elongata* subsp. *glycolytica* ATCC 29315	20.3	[17.1 - 23.9]	22.5	[20.2 - 25.0]	19.7	[17.0 - 22.8]	3.71
K. potus NCTC13336	*K. negevensis* Sch538	12.9	[10.2 - 16.2]	23.5	[21.2 - 25.9]	13.3	[10.9 - 16.0]	12.24

528 **Table 3. OGRI of *Kingella*, *Alysiella*, and *Simonsiella*.** dDDH was calculated using TYGS using each
529 of the possible algorithms, as well as the difference in G+C content between type strains of
530 *Alysiella* or *Simonsiella* species and each of the closest related subject type strains.

Query strain	Subject strain	dDDH (d0, in %)	C.I. (d0, in %)	dDDH (d4, in %)	C.I. (d4, in %)	dDDH (d6, in %)	C.I. (d6, in %)	G+C content difference (in %)
A. crassa NCTC10283	*A. filiformis* DSM 16848	21.7	[18.5 - 25.3]	34.4	[31.9 - 36.9]	21.9	[19.1 - 25.0]	1.29
A. crassa NCTC10283	*W. bovis* CCUG 44465T	18.4	[15.3 - 22.0]	33.4	[31.0 - 35.9]	18.8	[16.1 - 21.8]	2.67
A. crassa NCTC10283	*S. muelleri* ATCC 29453	16.9	[13.8 - 20.4]	23.2	[20.9 - 25.6]	16.9	[14.3 - 19.8]	3.83
A. crassa NCTC10283	*K. kingae* ATCC 23330	16.3	[13.3 - 19.8]	23.8	[21.5 - 26.3]	16.4	[13.8 - 19.3]	1.45
A. crassa NCTC10283	*K. negevensis* Sch538	15.6	[12.7 - 19.0]	23.3	[21.0 - 25.8]	15.7	[13.2 - 18.6]	0.17
A. crassa NCTC10283	*K. bonacorsii* Marseille-Q4569	14.2	[11.4 - 17.6]	22.7	[20.5 - 25.2]	14.5	[12.0 - 17.3]	8.71
A. crassa NCTC10283	*K. oralis* ATCC 51147	14.2	[11.4 - 17.6]	22	[19.7 - 24.4]	14.5	[12.1 - 17.3]	9.01
A. filiformis DSM 16848	*A. crassa* NCTC10283	21.8	[18.6 - 25.5]	34.1	[31.7 - 36.6]	22	[19.2 - 25.1]	1.25
A. filiformis DSM 16848	*W. bovis* CCUG 44465T	17.2	[14.2 - 20.8]	30.5	[28.1 - 33.0]	17.5	[14.9 - 20.5]	3.95
A. filiformis DSM 16848	*K. kingae* ATCC 23330	16.5	[13.5 - 20.0]	24	[21.7 - 26.5]	16.5	[14.0 - 19.5]	0.17
A. filiformis DSM 16848	*K. negevensis* Sch538	15.1	[12.2 - 18.6]	24.1	[21.8 - 26.6]	15.3	[12.8 - 18.2]	1.11
A. filiformis DSM 16848	*S. muelleri* ATCC 29453	15.1	[12.2 - 18.6]	23.7	[21.4 - 26.1]	15.3	[12.8 - 18.2]	5.12
A. filiformis DSM 16848	*C. steedae* DSM 2580	15.1	[12.2 - 18.5]	22.6	[20.3 - 25.0]	15.3	[12.8 - 18.2]	4.17
A. filiformis DSM 16848	*K. bonacorsii* Marseille-Q4569	14.6	[11.7 - 18.0]	22.8	[20.5 - 25.2]	14.8	[12.3 - 17.6]	7.43
A. filiformis DSM 16848	*K. oralis* ATCC 51147	14.6	[11.8 - 18.0]	22.2	[19.9 - 24.7]	14.8	[12.4 - 17.7]	7.72

Table 4. COGs responsible for the unification of *Kingella s.s.* in the accessory genome.

	COG	State	p_0[1]	p_1[2]		State	p_0[1]	p_1[2]	COG name[3]	Predicted Function[3]
Last Common Ancestor Alysiella, Simonsiella, and Kingella s.s.	742	1	0.15414	0.84586	*Last Common Ancestor of Kingella s.s.*	0	0.98197	0.01803	group_2532	Protein IscX
	766	1	0.02993	0.97007		0	0.97924	0.02076	group_2994	hypothetical protein
	771	1	0.02993	0.97007		0	0.97924	0.02076	*purU*	hypothetical protein
	923	1	0.24673	0.75327		0	0.97821	0.02179	group_3650	Sulfur carrier protein TusA
	3159	1	0.06887	0.93113		0	0.96883	0.03117	group_6852	Isocitrate lyase
	3277	1	0.35969	0.64031		0	0.96819	0.03181	*thiG*	Thiazole synthase
	3284	1	0.35969	0.64031		0	0.96819	0.03181	group_5996	Imidazole glycerol phosphate synthase subunit HisH
	3586	1	0.13975	0.86025		0	0.95545	0.04455	group_5655	Isocitrate dehydrogenase [NADP]

1 – Maximum likelihood posterior probability that this COG was absent at this site in the inferred ancestral reconstruction. Genes were called as absent if p_0 >= 0.95.

2 - Maximum likelihood posterior probability that this COG was present at this site in the inferred ancestral reconstruction. Genes were called as present if p_0 >= 0.95.

3 – Predicted names and functions of COGs of interest were recovered from the associated Roary matrix. Sequential COGs are not necessarily syntenic in any given genome.

Figures and Legends.

Figure 1. High quality 16S rRNA sequences were downloaded from the SILVA database for the family Neisseriaceae (taxid: 408). Sequences were aligned with MAFFT, and a maximum likelihood phylogeny was constructed using RAxML. Species that belong a species in the genus *Kingella* are colored red. By 16S rRNA relatedness, *Kingella* forms a polyphyletic clade with *K.*

549   *potus* falling outside of the rest of the genus. *K. potus* forms a clade with the oral-associated

550   *Neisseria bacilliformis*. Bootstrap values greater than 70% are shown.

Figure 2. ANI scores comparing sequenced isolates from species in the *Neisseriaceae* most closely

related to the genus *Kingella*. ANI scores were calculated between publicly available genomes

downloaded from NCBI using FastANI. Several regions of interest are indicated on the heatmap,

including the species delimiters, and regions of interest comparing *K. bonacorsii* and *K. oralis* (iv)

and *K. potus* and *N. bacilliformis* (v). The resulting ANI matrix was rendered as a heatmap using

R.

Figure 3. A. The core genome of the strains from the 13 species most closely related to *Kingella kingae* was determined using Roary with a minimum homology cutoff of 75%. The core genomes, which encompass approximately two million sites, were aligned with MAFFT and used to reconstruct a maximum likelihood phylogeny. As above, *K. potus* is only distantly related to the rest of the species in the genus *Kingella.* Additionally, the genera *Alysiella* and *Simonsiella* fall within the *Kingella* clade, suggesting the core genome of the species in each of these genera is very closely related to that of *Kingella.* Bootstraps greater than 70% are shown. **B-C.** The Robinson-Foulds and Normalized Nye similarity score congruency metrics were used to quantify changes in tree architecture between phylogenies reconstructed using core genome alignments. The Nye similarity score was normalized against the number of possible rearrangements given the number of leaves in the tree.

Figure 4. A. The core genome of the strains from the 13 species most closely related to *Kingella*

kingae was determined using Roary with a minimum homology cutoff of 50% with COGs present

in at least 80% of the species. The core genomes, which encompass approximately two million

sites, were aligned with MAFFT and used to reconstruct a maximum likelihood phylogeny. As

above, *K. potus* is only distantly related to the rest of the species in the genus *Kingella.*

Additionally, the genera *Alysiella* and *Simonsiella* fall within the *Kingella* clade, suggesting the

core genome of the species in each of these genera is very closely related to that of *Kingella.*

Bootstraps greater than 70% are shown. The phylogeny is rooted by the *Neisseria elongata* clade.

B. A binary gene presence/absence matrix was generated using Roary at the 75% similarity cut-

off, and includes both the core and accessory genes of all species. A phylogenetic tree based on

this matrix was constructed using RAxML via the GAMMA model of rate heterogeneity. This

analysis, based solely on gene content unites *Kingella sensu stricto* into a monophyletic group.

Bootstraps greater than 70% are shown. The phylogeny is rooted by the *Neisseria elongata* clade.

Figure 5. A. Scoary was used to generate lists of genes differentially encoded genes comparing

(i) CASK species against the *Neisseriaceae*, (ii) *Kingella sensu stricto* (iii) *Alysiella* and *Simonsiella*

or (iii) pathogenic *Kingella* against the CASK clade. Finally, *Alysiella* and *Simonsiella* COGs were

compared against the pathogenic *Kingella.* Diagnostic COGs are defined as COGs with 100%

sensitivity and specificity to the in-group, and filtered to *p < 0.001*. Please see Supplemental

tables 5-9 list for full COGs sets. **B-C.** Diagnostic gene sets for each group were annotated using

eggNOG mapper, and classified by Gene Ontology functional groups using the NCBI COG

database. B. The CASK clade is enriched for unique COGs with putative roles in cell wall,

membrane and envelope biogenesis, as well as DNA replication, recombination and repair. COGs

with no functional annotations were omitted from this graphic (n = 656 COGs). **C.** Similarly, the

majority of the unique COGs present in *Alysiella* and *Simonsiella* may function as part of DNA

replication, recombination and repair pathways and cell wall, membrane and envelope

biogenesis. *Kingella sensu stricto* and the pathogenic *Kingella* species encode more unique COGs

implicated in carbohydrate, amino acid, or nucleotide transport and metabolism. **D.** Type strains

of each species in the CASK clade was compared to the *K. kingae* ATCC 23330 genome (inner

circle) using BLASTn. Bars in each ring represent regions of sequence similarity between

genomes. Each of the CASK species shows significant sequence similarity to nearly all of the *K.*

kingae reference, further demonstrating how closely related each of these species are. Regions

of dissimilarly map to regions that encode putative MGEs, particularly prophages.

608