Phylogenomic analyses of understudied *Neisseriaceae* species support the reclassification of the polyphyletic genera *Kingella, Simonsiella, and Alysiella*.

Daniel P. Morreale[1,2], Joseph W. St Geme III[1,2†], and Paul J Planet [1,2,3*†]

Affiliations:

[1] Perelman School of Medicine, University of Pennsylvania, Philadelphia PA.

[2] Division of Infectious Diseases, Children's Hospital of Pennsylvania, Philadelphia PA.

[3] Comparative Genomics, American Museum of Natural History, New York, New York, United States of America

*Corresponding Author: Paul J. Planet (pjplanet@chop.edu)

† Contributed equally

Keywords: *Kingella, Simonsiella, Alysiella, Neisseriaceae,* Polyphyly,

Abbreviations: ANI: Average Nucleotide identity; dDDH: digital DNA-DNA hybridization, OGRI: overall genome relatedness index, COGs: clusters of orthologous gene groups; MuLDi: Multicellular, longitudinally-dividing; DUS: DNA-uptake sequences, MALDI-TOF MS: Matrix-assisted laser desorption-ionization time of flight mass spectrometry

Abstract.

Taxonomic classification of the *Neisseriaceae* family has focused primarily on the pathogens

Neisseria meningitidis and *Neisseria gonorrhoeae*. As a consequence, the commensal *Neisseria*

species and other *Neisseriaceae* genera such as *Kingella* have been only loosely classified,

resulting in several polyphyletic genera. In this study, using available 16S rRNA sequences and a

phylogenetic approach, we found that the genus *Kingella* is polyphyletic, due to misclassification

of *Kingella potus*. Calculation of multiple genome relatedness indices revealed that *K. potus* is

more similar to *Neisseria bacilliformis* than it is to other *Kingella* species. To better understand

the evolution of this clade, we examined the core genome of *Kingella* and closely related genera

and discovered that the related genera *Simonsiella* and *Alysiella* form a distinct clade within the

genus *Kingella* that is closely related to the pathogens *K. kingae* and *K. negevensis*. Based on

these analyses, we propose incorporation of *Simonsiella* and *Alysiella* species into the genus

Kingella.

Main text.

The *Neisseriaceae* family is a group of Gram-negative, β-proteobacteria that currently

includes 20 genera: *Neisseria*, *Wielerella*, *Vitreoscilla*, *Uruburuella*, *Stenoxybacter*, *Snodgrassela*,

Simonsiella, *Rivicola*, *Prolinoborus*, *Paralysiella*, *Morococcus*, *Kingella*, *Eikenella*, *Crenobacter*,

Craterilacuibacter, *Conchiformibius*, *Bergeriella*, *Aquella*, *Amantichitinum*, and *Alysiella* [1–3].

The majority of prior work devoted to understanding the taxonomic relationships between these

genera has focused primarily on the pathogenic *Neisseria* species, *N. meningitidis* and *N.*

gonorrhoeae. To date, the commensal *Neisseria* species and related genera have been only

loosely classified based on preliminary studies. The genus *Kingella* includes five taxa: *K. potus, K.*

oralis, K. denitrificans, K. negevensis, and *K. kingae*. Recently, a new species was proposed, *K.*

bonacorsii [4–10]. Each of the *Kingella* species is typically associated with the oral or

oropharyngeal microbiome, a common niche of other *Neisseriaceae*. Isolates of *K. kingae*, *K.*

negevensis, and *K. potus* have all been associated with bacteremia, osteoarticular infections,

and/or endocarditis in immunocompetent individuals [6, 8, 11]. In contrast, *K. oralis* and *K.*

denitrificans are present in dental plaque, gingival and periodontal samples, and only rarely cause

invasive infections, typically in immunocompromised individuals [9, 12, 13].

A review of published phylogenetic relationships among the commensal *Neisseria* species

reveals a high level of variability in the proposed taxonomic relationships, primarily owing to the

genes selected and method used to perform the analysis [14–17]. The genus *Kingella* is

polyphyletic, and the genera that interrupt a monophyletic *Kingella* clade differ between

published analyses. Most isolates used for defining taxonomic nomenclature in this genus were

collected prior to pervasive whole genome sequencing and were assigned to *Kingella* as a result

of phenotypic and/or minimal sequence-based analyses. Additionally, as the frequency of

culture-dependent and independent microbiome studies have become more prevalent, the

number and diversity of formally accepted species in the *Neisseriaceae* has increased [18]. With

this background, we sought to reassess the relatedness of the five species in *Kingella*.

To examine the *Kingella* species, we employed a phylogenetic approach. 840 of the

available 16S rRNA gene sequences for isolates from the *Neisseriaceae* and in the RefNR database

were downloaded from the SILVA r138.1 database on September 08, 2022 [19, 20]. 16S rRNA

sequences were aligned with MAFFT, and a phylogeny was reconstructed using RAxML v. 8.4.2

with the general time reversible model of nucleotide substitution and gamma model of rate

heterogeneity, and bootstrapped 100 times [21, 22]. The resulting tree was annotated using

Figtree v.1.4.4 and ITOL and is shown in Fig. 1 [23]. By 16S rRNA relatedness alone, it is clear that

a major driver of the polyphyletic structure of *Kingella* is *K. potus.* This species was recovered

from an infected wound caused by an animal bite and is more closely related to *Neisseria*

bacilliformis than it is to any currently described *Kingella* species [8, 24]. *N. bacilliformis* was

identified shortly after *K. potus* was defined, is a part of the oral microbiome of non-human

mammals, and is associated with bacteremia in humans after animal bites [24].

To better understand the relationship between the *K. potus* and *N. bacilliformis*, we

calculated average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) between

the type strains of these species using FastANI v.1.33 and the Type Strain Genome Server (TYGS),

respectively [25–27]. The ANI between *K. potus* and *N. bacilliformis* was calculated at 86.75%,

below the established species cut-off of 95% [28]. However, the ANI between *K. potus and N.*

bacilliformis is appreciably greater than the ANI calculated between *K. potus* and other *Kingella*

species (Table 1). Similarly, while the calculated dDDH is below species thresholds (Table 2) [29],

it is clear that *K. potus* is far more similar to *N. bacilliformis* than it is to other species within the

genus *Kingella.*

To further define the relationships between *Neisseriaceae*, whole genome sequences for

select species in this family were downloaded from NCBI (on September 09, 2022). These

analyses were limited to only those species most closely related to *Kingella, by* 16S rRNA

similarity and include *Neisseria elongata, Neisseria sheyganii, N. bacilliformis, Eikenella*

corrodens, Alysiella filiformis, Alysiella crassa, Simonsiella muelleri, Conchiformibius steedae, K.

oralis, K. bonacorsii, K. potus, K. denitrificans, K. kingae, and *K. negevensis* [7, 30]*.* In total, 134

genomes were reassembled with Prokka v1.14.6, using default settings for bacterial sequences

[31]. Subsequent assemblies were analyzed with Roary v3.13.0. to identify the core genome [32].

Core genes were clustered at the 75%, 80%, 85%, 90%, and 95% sequence similarity level and

were limited to only those clusters of orthologous groups (COGs) found in all genomes.

Nucleotide sequences of core genes were compared in a codon-aware alignment by PRANK

within Roary, and core gene alignments were used to reconstruct phylogenies using RAxML as

above. Each phylogeny was overwhelmingly congruent with the others, and we therefore

proceeded with the lowest similarity cutoff for detailed analysis (Fig. 2A). These phylogenies

further support the close relationship between *K. potus* and *N. bacilliformis*.

The phylogenomic analyses also supported a close evolutionary relationship between

Kingella, *Alysiella*, and *Simonsiella*. *Alysiella* and *Simonsiella* are commensal genera associated

with the oral microbiome of mammals [14, 33, 34]. These two genera, along with the genus

Conchiformibius, are distinctive for being multicellular, longitudinally-dividing (MuLDi) species

that comprise the only known animal multicellular bacterial symbionts [15]. Based on a whole

genome approach, *Alysiella* and *Simonsiella* diverged from the rest of the *Kingella* species more

recently than *K. denitrificans* or *K. oralis* diverged. ANI, dDDH, and G+C% were calculated for type

strains of *Kingella*, *Alysiella*, and *Simonsiella* and are shown in Tables 1 and 3. While there is no

universally accepted ANI cutoff for demarcating a bacterial genus, ANI scores greater than 73%

have been proposed [35]. All of these overall genome relatedness indices (OGRI) support the

close relationship between these three genera, and the ANI falls above the proposed 73% genus

cutoff score. Moreover, Nyongesa and co-workers [15] hypothesize that the MuLDi phenotype

evolved twice in the *Neisseriaceae,* first in *Conchiformibius* species and then in the common

ancestor of *Simonsiella and* Alysiella, a hypothesis that is strongly supported by our phylogenetic

analyses.

Additionally, we calculated pairwise ANI scores for each of the genomes listed above. The

ANI scores confirm the majority of the proposed species classifications with one exception.

Recently, an unclassified *Kingella* isolate was proposed to be a new species and was named *K.*

bonacorsii [4]. This isolate was recovered from a gingival sample, and the authors report a 16S

rRNA similarity of 98.7%, ANI of 95.83%, and dDDH of 63.6% relative to *K. oralis* UB-38, as well as

non-identification by MALDI-TOF mass-spectrometry [4]. We replicated each of the genomic

calculations reported, as well as additional dDDH calculations using alternative formulae (Tables

1 and 4) [28, 36]. The 16S rRNA similarity and ANI score are both above the threshold for

classification of *K. bonacorsii* and *K. oralis* as the same species, consistent with the very similar

G+C% and their common niche. Importantly, we note that the algorithm used to calculate dDDH

is critical in this instance. Analysis with either *d0* or *d6* yields dDDH above the threshold of 70%.

Only if dDDH is calculated with *d4,* which is considered most appropriate for highly fragmented

or gapped genomes, the dDDH falls below the accepted threshold of 70% (Table 4). Very few

isolates of *K. oralis* are publicly available for additional comparisons, and without additional

isolates and sequencing data, it is difficult to appreciate the diversity of *K. oralis* or account for

bias in strain selection. Based on the available data, we conclude that *K. bonacorsii* is likely not a

unique species in the genus *Kingella* and instead may be a subspecies of *K. oralis.*

Finally, another recent evolutionary study of these taxa has used less stringent similarity

and gene presence cutoff to define the core genome for species in the *Neisseriaceae* for

subsequent phylogenetic analyses [15]. We replicated this analysis by reanalyzing the data using

a sequence similarity cut off of 50% for genes present in 99% or 80% of analyzed genomes. By

lowering the threshold for a gene to be considered part of the core genome, we increase the

number of genes considered from 109 at the 75% similarity level up to 620 genes. Notably, as we

increase the number of genes considered part of the core genome, the phylogenetic architecture

remains constant (Fig. 3A).

The reproducibility of the core genome phylogenetic structure very strongly supports the

inclusion of *Kingella*, *Alysiella*, and *Simonsiella* in the same genus, and the exclusion of *K. potus*.

We hypothesized that prior misclassification could be due to high levels of horizontal gene

transfer of accessory genes between *Kingella* (*sensu stricto*) genomes. To test this, we generated

a presence/absence matrix generated by Roary at 75% sequence similarity level, a technique that

emphasizes accessory genes over core genes, and we reconstructed a phylogeny using the RAxML

binary model with gamma model of rate heterogeneity, and bootstrapped 100 times (Fig. 3B).

This was the only analysis in which we observed a monophyletic *Kingella* genus, based on current

taxonomy, though *Kingella potus* was still excluded. *Simonsiella*, *K. oralis*, and *K. kingae* utilize

specific DNA-uptake sequences (DUS) to limit horizontal gene transfer (HGT) events, with

Simonsiella, *K. oralis* seemingly sharing the same DUS [37]. Given that members of the

Neisseriaceae are highly recombinogenic and undergo significant HGT of accessory genes, we

assert that the core genome is a higher fidelity evolutionary record for this family [38].

Based on these results, we propose several changes to the genera *Kingella, Alysiella, and*

Simonsiella. As *K. potus* is more related to *N. bacilliformis* than other *Kingella* species by 16S

rRNA, OGRI, and whole genome analyses, we propose revising the taxonomic nomenclature of

this isolate to reflect this close relationship. Additionally, as phylogenomic analysis of the core

genomes of *Alysiella* and *Simonsiella* supports their close relationship to the genus *Kingella,* we

propose incorporating these genera into the genus *Kingella*, as *K. crassa*, *K. filiformis*, and *K.*

muelleri. These changes would adjust the accepted taxonomic nomenclature to better reflect the

phylogenomic relationships between these isolates and result in a monophyletic *Kingella* genus.

Finally, further characterization will be required for confirm if *K. bonacorsii* constitutes its own

species independent of *K. oralis* or if it would be better classified as a subspecies (*K. oralis* subsp*.*

bonacorsii).

Acknowledgements.

We thank the members of the Planet and St. Geme Labs for their useful feedback and discussion.

Funding.

This work was supported by the National Institute of Allergy and Infectious Diseases under the

training grant in Microbial Pathogenesis and Genomics (5T32141493) to DPM and under award

1R01AI121015 to JWS.

Conflicts of Interest.

The authors report no conflicts of interest.

References.

1. **Parte AC, Carbasse JS, Meier-Kolthoff JP, Reimer LC, Göker M**. List of prokaryotic names

with standing in nomenclature (LPSN) moves to the DSMZ. *Int J Syst Evol Microbiol*

2020;70:5607–5612.

2. **Parker CT, Tindall BJ, Garrity GM**. International code of nomenclature of Prokaryotes. *Int*

J Syst Evol Microbiol 2019;69:S1.

3. **Oren A, Garrity G**. Notification of changes in taxonomic opinion previously published

outside the IJSEM. List of changes in taxonomic opinion no. 36. *Int J Syst Evol Microbiol*

2022;72:005413.

4. **Antezack A, Boxberger M, Rolland C, Monnet-Corti V, La Scola B**. Isolation and

Characterization of Kingella bonacorsii sp. nov., A Novel Kingella Species Detected in a

Stable Periodontitis Subject. *Pathogens* 2021;10:1–13.

5. **Henricksen SD, Bovre K**. Transfer of *Moraxella kingae* Henriksen and Bovre to the genus

Kingella gen. nov. in the family *Neisseriacea*e. *Int J Syst Bacteriol* 1976;26:447–450.

6. **El Houmami N, Bakour S, Bzdrenga J, Rathored J, Seligmann H, *et al.*** Isolation and

characterization of *Kingella negevensis* sp. nov., a novel *Kingella* species detected in a

healthy paediatric population. *Int J Syst Evol Microbiol* 2017;67:2370–2376.

7. **Dewhirst FE, Chen CKC, Paster BJ, Zambon JJ**. Phylogeny of species in the family

Neisseriaceae isolated from human dental plaque and description of *Kingella oralis* sp.

nov [corrected]. *Int J Syst Bacteriol* 1993;43:490–499.

8. **Lawson PA, Malnick H, Collins MD, Shah JJ, Chattaway MA, *et al.*** Description of Kingella

potus sp. nov., an Organism Isolated from a Wound Caused by an Animal Bite. *J Clin*

Microbiol 2005;43:3526.

9.   **Snell JJS, Lapage SP**. Transfer of some saccharolytic *Moraxella* species to *Kingella*

Henriksen and Bovre 1976, with descriptions of *Kingella indologenes* sp. nov. and

Kingella denitrificans sp. nov. *Int J Syst Bacteriol* 1976;26:451–458.

10.   **Hollis DG, Weaver RE, Riley PS**. Emended description of *Kingella denitrificans* (Snell and

Lapage 1976): correction of the maltose reaction. *J Clin Microbiol* 1983;18:1174.

11.   **Yagupsky P**. *Kingella kingae*: Carriage, transmission, and disease. *Clinical Microbiology*

Reviews 2015;28:54–79.

12.   **Chen C, Chen C**. Distribution of a newly described species, *Kingella oralis*, in the human

oral cavity. *Oral Microbiol Immunol* 1996;11:425–427.

13.   **Kopyt N, Kumar A, Agrawal V**. A Case of Suppurative Peritonitis by a Commensal Oral

Organism, *Kingella denitrificans*, in an Adult Peritoneal Dialysis Patient. *Perit Dial Int*

2015;35:105.

14.   **Adeolu M, Gupta RS**. Phylogenomics and molecular signatures for the order Neisseriales:

proposal for division of the order Neisseriales into the emended family Neisseriaceae and

Chromobacteriaceae fam. nov. *Antonie Van Leeuwenhoek* 2013;104:1–24.

15.   **Nyongesa S, Weber PM, Bernet È, Pulido F, Nieves C, *et al.*** Evolution of longitudinal

division in multicellular bacteria of the Neisseriaceae family. *Nat Commun 2022 131*

2022;13:1–18.

16.   **Garrity GM, Bell JA, Lilburn T**. Class II. Betaproteobacteria class. nov. *Bergey's Manual®*

Syst Bacteriol 2005;575–922.

17.   **Enright MC, Carter PE, MacLean IA, McKenzie H**. Phylogenetic relationships between

some members of the genera *Neisseria*, *Acinetobacter, Moraxella*, and *Kingella* based on

partial 16S ribosomal DNA sequence analysis. *Int J Syst Bacteriol* 1994;44:387–391.

18.   **Maiden MCJ, Harrison OB**. Population and functional genomics of *Neisseria* revealed

with gene-by-gene approaches. *J Clin Microbiol* 2016;54:1949–1955.

19.   **Glöckner FO, Yilmaz P, Quast C, Gerken J, Beccati A, *et al.*** 25 years of serving the

community with ribosomal RNA gene reference databases and tools. *J Biotechnol*

2017;261:169–176.

20.   **Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, *et al.*** The SILVA ribosomal RNA gene

database project: improved data processing and web-based tools. *Nucleic Acids Res*

2013;41:D590–D596.

21.   **Stamatakis A**. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* 2014;30:1312.

22.   **Katoh K, Standley DM**. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol Biol Evol* 2013;30:772.

23.   **Letunic I, Bork P**. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Res* 2021;49:W293–W296.

24.   **Han XY, Hong T, Falsen E**. *Neisseria bacilliformis* sp. nov. Isolated from Human Infections.

J Clin Microbiol 2006;44:474.

25.   **Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S**. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun 2018*

91 2018;9:1–8.

26.   **Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Göker M**. TYGS and LPSN: a database

tandem for fast and reliable genome-based classification and nomenclature of

prokaryotes. *Nucleic Acids Res* 2022;50:D801–D807.

27.   **Meier-Kolthoff JP, Göker M**. TYGS is an automated high-throughput platform for state-

of-the-art genome-based taxonomy. *Nat Commun 2019 101* 2019;10:1–10.

28.   **Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M**. Genome sequence-based species

delimitation with confidence intervals and improved distance functions. *BMC*

Bioinformatics 2013;14:1–14.

29.   **Wayne LG, Brenner DJ, Colwell RR, Grimont PAD, Kandler O, *et al.*** Report of the Ad Hoc

Committee on Reconciliation of Approaches to Bacterial Systematics. *Int J Syst Evol*

Microbiol 1987;37:463–464.

30.   **Dewhirst FE, Paster BJ, Bright PL**. *Chromobacterium, Eikenella, Kingella, Neisseria,*

Simensiella, and *Vitreoscilla* species comprise a major branch of the beta group

Proteobacteria by 16S ribosomal ribonucleic acid sequence comparison: Transfer of

Eikenella and *Simonsiella* to the family Neisseriaceae (emend.). *Int J Syst Bacteriol*

1989;39:258–266.

31.   **Seemann T**. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–

2069.

32.   **Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, *et al.*** Roary: rapid large-scale

prokaryote pan genome analysis. *Bioinformatics* 2015;31:3691–3693.

33.   **Hedlund BP, Tønjum T**. *Simonsiella*. *Bergey's Man Syst Archaea Bact* 2015;1–12.

34.   **Xie CH, Yokota A**. Phylogenetic analysis of *Alysiella* and related genera of Neisseriaceae:

proposal of *Alysiella crassa* comb. nov., *Conchiformibium steedae* gen. nov., comb. nov.,

Conchiformibium kuhniae sp. nov. and *Bergeriella denitrificans* gen. nov., comb. nov. *J*

Gen Appl Microbiol 2005;51:1–10.

35.   **Barco RA, Garrity GM, Scott JJ, Amend JP, Nealson KH, *et al.*** A genus definition for

bacteria and archaea based on a standard genome relatedness index. *MBio*;11. Epub

ahead of print 2020. DOI: 10.1128/MBIO.02475-19/SUPPL_FILE/MBIO.02475-19-

S0001.DOCX.

36.   **Auch AF, von Jan M, Klenk HP, Göker M**. Digital DNA-DNA hybridization for microbial

species delineation by means of genome-to-genome sequence comparison. *Stand*

Genomic Sci 2010;2:117.

37.   **Frye SA, Nilsen M, Tønjum T, Ambur OH**. Dialects of the DNA Uptake Sequence in

Neisseriaceae. *PLOS Genet* 2013;9:e1003458.

38.   **Hanage WP, Fraser C, Spratt BG**. Fuzzy species among recombinogenic bacteria. *BMC*

Biol 2005;3:1–7.

Figures.

Figure 1. High quality 16S rRNA sequences were downloaded from the SILVA database for the

family Neisseriaceae (taxid: 408). Sequences were aligned with MAFFT, and a maximum

likelihood phylogeny was constructed using RAxML. Species that belong to the genus *Kingella* are

colored red. By 16S rRNA relatedness, *Kingella* forms a polyphyletic clade with *K. potus* falling

outside of the rest of the genus. *K. potus* forms a clade with the oral-associated *Neisseria*

bacilliformis. Bootstrap values greater than 70% are shown.

Figure 2. A. The core genome of the strains from the 13 species most closely related to *Kingella*

kingae was determined using Roary with a minimum homology cutoff of 75%. The core genomes,

which include the sequences of 109 genes total, which were aligned with PRANK in Roary and

RAxML was used to reconstruct a maximum likelihood phylogeny. *K. potus* is only distantly

related to the rest of the species in the genus *Kingella.* The genera *Alysiella* and *Simonsiella* fall

within the *Kingella* clade, suggesting that the core genome of the species in each of these genera

is very closely related to that of *Kingella.* Additionally, *K. bonacorsii* is very closely related to *K.*

oralis, and these two species form a well-supported clade. The tree is rooted by *N. elongata,* and

bootstraps greater than 70% are shown. **B.** ANI score between each isolate is shown on the tree.

272   The ANI score when comparing *K. oralis* and *K. bonacorsii* suggests that differentiating between

273   these isolates as two species may not reflect speciation events.

Figure 3. A. The core genome of the strains from the 13 species most closely related to *Kingella*

kingae was determined using Roary with a minimum protein similarity cutoff of 50%, effectively

reducing the stringency for genes to be considered a part of the core genome. The core

genomes, which include the sequences of 620 CDS total, were aligned with MAFFT, and RAxML

was used to reconstruct a maximum likelihood phylogeny with 100 bootstraps. Bootstrap

values >70% are shown, and the tree is rooted by *N. elongata*. Unlike previous analyses with

similar datasets, *Alysiella* and *Simonsiella* are still more closely related to *K. kingae* than to

either *K. oralis* or *K. denitrificans*. **B.** The pangenome of the species closely related to *Kingella*

was determined using Roary, at the 75% similar level and ignoring genes that occur in all

isolates. RAxML was used to reconstruct a phylogeny of gene presence/absence. Bootstrap

values >70% are shown, and the tree is rooted by *N. elongata*. This phylogeny is different than

the 75% sequence similarity tree shown in panel A, suggesting that the pangenome gene

content may heavily influence the phylogenetic relation relationship between these species,

clouding the evolutionary history demonstrated by core genes.

Table 1. Average ANI scores calculated among taxa of interest. ANI scores were calculated using
FastANI and are shown only if the ANI >75%. Empty cells denote data excluded from the table.

	N. bacilliformis	K. denitrificans	K. oralis	K. negevensis	K. kingae	N. elongata	A. crassa	A. filiformis	S. muelleri
K. potus	86.75	78.88	78.74	<75	<75	80.7	-	-	-
K. bonacorsii	-	79.84	96.41	78.06	77.9	78.76	77.98	77.87	77.14
A. crassa	-	77.81	77.96	79.37	79.74	<75	100	85.43	79.79
A. filiformis	-	77.81	77.96	79.37	79.74	<75	85.43	100	78.75
S. muelleri	-	<75	76.9	79.16	79.96	<75	79.79	78.75	100

295 **Table 2. dDDH values of *K. potus* calculated against closely related taxa.** dDDH was calculated
296 using TYGS using each of the possible algorithms, as well as the difference in G+C content
297 between *K. potus* and each of the subject type strains.

Query strain	Subject strain	dDDH (d0, in %)	C.I. (d0, in %)	dDDH (d4, in %)	C.I. (d4, in %)	dDDH (d6, in %)	C.I. (d6, in %)	G+C content difference (in %)
K. potus NCTC13336	*N. bacilliformis* ATCC BAA-1200	45.6	[42.2 - 49.0]	30	[27.6 - 32.5]	41	[38.0 - 44.0]	1.86
K. potus NCTC13336	*N. elongata* subsp. *glycolytica* ATCC 29315	20.3	[17.1 - 23.9]	22.5	[20.2 - 25.0]	19.7	[17.0 - 22.8]	3.71
K. potus NCTC13336	*K. negevensis* Sch538	12.9	[10.2 - 16.2]	23.5	[21.2 - 25.9]	13.3	[10.9 - 16.0]	12.24

Table 3. OGRI of *Kingella*, *Alysiella*, and *Simonsiella*. dDDH was calculated using TYGS using each of the possible algorithms, as well as the difference in G+C content between type strains of *Alysiella* or *Simonsiella* species and each of the closest related subject type strains.

Query strain	Subject strain	dDDH (d0, in %)	C.I. (d0, in %)	dDDH (d4, in %)	C.I. (d4, in %)	dDDH (d6, in %)	C.I. (d6, in %)	G+C content difference (in %)
A. crassa NCTC10283	*A. filiformis* DSM 16848	21.7	[18.5 - 25.3]	34.4	[31.9 - 36.9]	21.9	[19.1 - 25.0]	1.29
A. crassa NCTC10283	*W. bovis* CCUG 44465T	18.4	[15.3 - 22.0]	33.4	[31.0 - 35.9]	18.8	[16.1 - 21.8]	2.67
A. crassa NCTC10283	*S. muelleri* ATCC 29453	16.9	[13.8 - 20.4]	23.2	[20.9 - 25.6]	16.9	[14.3 - 19.8]	3.83
A. crassa NCTC10283	*K. kingae* ATCC 23330	16.3	[13.3 - 19.8]	23.8	[21.5 - 26.3]	16.4	[13.8 - 19.3]	1.45
A. crassa NCTC10283	*K. negevensis* Sch538	15.6	[12.7 - 19.0]	23.3	[21.0 - 25.8]	15.7	[13.2 - 18.6]	0.17
A. crassa NCTC10283	*K. bonacorsii* Marseille-Q4569	14.2	[11.4 - 17.6]	22.7	[20.5 - 25.2]	14.5	[12.0 - 17.3]	8.71
A. crassa NCTC10283	*K. oralis* ATCC 51147	14.2	[11.4 - 17.6]	22	[19.7 - 24.4]	14.5	[12.1 - 17.3]	9.01
A. filiformis DSM 16848	*A. crassa* NCTC10283	21.8	[18.6 - 25.5]	34.1	[31.7 - 36.6]	22	[19.2 - 25.1]	1.25
A. filiformis DSM 16848	*W. bovis* CCUG 44465T	17.2	[14.2 - 20.8]	30.5	[28.1 - 33.0]	17.5	[14.9 - 20.5]	3.95
A. filiformis DSM 16848	*K. kingae* ATCC 23330	16.5	[13.5 - 20.0]	24	[21.7 - 26.5]	16.5	[14.0 - 19.5]	0.17
A. filiformis DSM 16848	*K. negevensis* Sch538	15.1	[12.2 - 18.6]	24.1	[21.8 - 26.6]	15.3	[12.8 - 18.2]	1.11
A. filiformis DSM 16848	*S. muelleri* ATCC 29453	15.1	[12.2 - 18.6]	23.7	[21.4 - 26.1]	15.3	[12.8 - 18.2]	5.12
A. filiformis DSM 16848	*C. steedae* DSM 2580	15.1	[12.2 - 18.5]	22.6	[20.3 - 25.0]	15.3	[12.8 - 18.2]	4.17
A. filiformis DSM 16848	*K. bonacorsii* Marseille-Q4569	14.6	[11.7 - 18.0]	22.8	[20.5 - 25.2]	14.8	[12.3 - 17.6]	7.43
A. filiformis DSM 16848	*K. oralis* ATCC 51147	14.6	[11.8 - 18.0]	22.2	[19.9 - 24.7]	14.8	[12.4 - 17.7]	7.72

305 **Table 4. dDDH calculations for the proposed species *K. bonacorsii*.** dDDH was calculated using
306 TYGS using each of the possible algorithms, as well as the difference in G+C content between *K.*
307 *bonacorsii* and each of the subject type strains.

Query strain	Subject strain	dDDH (d0, in %)	C.I. (d0, in %)	dDDH (d4, in %)	C.I. (d4, in %)	dDDH (d6, in %)	C.I. (d6, in %)	G+C content difference (in %)
K. bonacorsii Marseille-Q4569	*K. oralis* ATCC 51147	84.3	[80.5 - 87.4]	63.6	[60.7 - 66.4]	83.1	[79.8 - 86.0]	0.29
K. bonacorsii Marseille-Q4569	*K. denitrificans* ATCC 33394	15.9	[13.0 - 19.4]	23.2	[20.9 - 25.7]	16	[13.5 - 18.9]	0.06
K. bonacorsii Marseille-Q4569	*N. bacilliformis* ATCC BAA-1200	15.1	[12.2 - 18.5]	25.3	[23.0 - 27.8]	15.3	[12.8 - 18.2]	5.55
K. bonacorsii Marseille-Q4569	*K. potus* NCTC13336	14.4	[11.6 - 17.8]	23.2	[20.9 - 25.7]	14.7	[12.2 - 17.5]	3.84
K. bonacorsii Marseille-Q4569	*K. negevensis* Sch538	14.4	[11.6 - 17.8]	22	[19.7 - 24.4]	14.7	[12.2 - 17.5]	8.54

308