Role of spike in the pathogenic and antigenic behavior of SARS-CoV-2 BA.1 Omicron

Da-Yuan Chen[1,2], Devin Kenney[2,3], Chue Vin Chin[1,2], Alexander H. Tavares[1,2], Nazimuddin Khan[1,2], Hasahn L. Conway[1,2], GuanQun Liu[4], Manish C. Choudhary[5,6], Hans P. Gertje[2], Aoife K. O'Connell[2], Darrell N. Kotton[7,8], Alexandra Herrmann[9], Armin Ensser[9], John H. Connor[2,3], Markus Bosmann[10,11,12], Jonathan Z. Li[5,6], Michaela U. Gack[4], Susan C. Baker[13], Robert N. Kirchdoerfer[14], Yachana Kataria[11], Nicholas A. Crossland[2,11], Florian Douam[2,3], Mohsan Saeed[1,2#]

1. Department of Biochemistry, Boston University School of Medicine, Boston, MA, USA

2. National Emerging Infectious Diseases Laboratories, Boston University, Boston, MA, USA

3. Department of Microbiology, Boston University School of Medicine, Boston, MA, USA

4. Cleveland Clinic Florida Research and Innovation Center, Port St. Lucie, FL, USA

5. Brigham and Women's Hospital, Boston, MA, USA

6. Harvard Medical School, Cambridge, MA, USA

7. Center for Regenerative Medicine of Boston University and Boston Medical Center, Boston, MA, USA

8. The Pulmonary Center and Department of Medicine, Boston University School of Medicine, Boston, MA, USA

9. Institute of Clinical and Molecular Virology, University Hospital Erlangen, Friedrich-

Alexander Universität Erlangen-Nürnberg, Erlangen, Germany

10. Pulmonary Center, Boston University School of Medicine, MA, USA

11. Department of Pathology and Laboratory Medicine, Boston University School of

Medicine, MA, USA

12. Center for Thrombosis and Hemostasis, University Medical Center of the Johannes

Gutenberg-University, Mainz, Germany

13. Department of Microbiology and Immunology, and Infectious Disease and Immunology

Research Institute, Stritch School of Medicine, Loyola University, Chicago, Maywood, IL,

USA

14. Department of Biochemistry, College of Agricultural and Life Sciences, University of

Wisconsin, Madison, WI, USA

[#]Correspondence: msaeed1@bu.edu

The recently identified, globally predominant SARS-CoV-2 Omicron variant (BA.1) is highly transmissible, even in fully vaccinated individuals, and causes attenuated disease compared with other major viral variants recognized to date[1-7]. The Omicron spike (S) protein, with an unusually large number of mutations, is considered the major driver of these phenotypes[3,8]. We generated chimeric recombinant SARS-CoV-2 encoding the S gene of Omicron in the backbone of an ancestral SARS-CoV-2 isolate and compared this virus with the naturally circulating Omicron variant. The Omicron S-bearing virus robustly escapes vaccine-induced humoral immunity, mainly due to mutations in the receptor-binding motif (RBM), yet unlike naturally occurring Omicron, efficiently replicates in cell lines and primary-like distal lung cells. In K18-hACE2 mice, while Omicron causes mild, non-fatal infection, the Omicron S-carrying virus inflicts severe disease with a mortality rate of 80%. This indicates that while the vaccine escape of Omicron is defined by mutations in S, major determinants of viral pathogenicity reside outside of S.

As of March 2022, the successive waves of the coronavirus disease 2019 (COVID-19) pandemic have been driven by five major SARS-CoV-2 variants, called variants of concern (VOC); Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2 and AY lineages), and Omicron (BA lineages)[9]. Omicron is the most recently recognized VOC that was first documented in South Africa, Botswana, and in a traveler from South Africa in Hong Kong in November 2021 (GISAID ID: EPI_ISL_7605742)[10,11]. It quickly swept through the world, displacing the previously dominant Delta variant within weeks and accounting for the majority of new SARS-CoV-2 infections by January 2022[12-16]. Omicron has at least three lineages, BA.1, BA.2, and BA.3, with the former being the most predominant lineage worldwide[13,17-19]. BA.1 (hereinafter referred to as Omicron)

63 exhibits a remarkable escape from infection- and vaccine-induced humoral immunity[4,5,20,21].

64 Further, it is less pathogenic than other VOCs in humans and *in vivo* models of infection[1-3,22-26].

65 Omicron differs from the prototype SARS-CoV-2 isolate, Wuhan-Hu-1, by 59 amino acids; 37 of

66 these changes are in the S protein, raising the possibility that S is at the heart of Omicron's

67 pathogenic and antigenic behavior.

Spike mutations only partially affect the replication of Omicron in cell culture

The Omicron S protein carries 30 amino acid substitutions, 6 deletions, and one three-

amino acid-long insertion compared to Wuhan-Hu-1 (**Fig. 1a,b**). Twenty-five of these changes are

unique to Omicron relative to other VOCs, although some of them have been reported in waste

water and minor SARS-CoV-2 variants[27-29]. To test the role of the S protein in Omicron phenotype,

we generated a chimeric recombinant virus containing the S gene of Omicron (USA-lh01/2021) in

the backbone of an ancestral SARS-CoV-2 isolate (GISAID EPI_ISL_2732373)[30] (**Fig. 1c**). To

produce this chimeric Omi-S virus, we employed a modified form of cyclic polymerase extension

reaction (CPER) (**Extended Data Fig. 1**) that yielded highly concentrated virus stocks, containing

0.5-5 x 10^6 plaque-forming units (PFU) per ml, from transfected cells within two days of

transfection (**Fig. 1d,e**), obviating the need for additional viral amplification[31,32].

We first compared the infection efficiency of Omi-S with an ancestral virus and Omicron in

cell culture (**Fig. 2a**). For this, we infected ACE2/TMPRSS2/Caco-2[33] and Vero E6 cells with Omi-

S, a recombinant D614G-bearing ancestral virus (GISAID EPI_ISL_2732373)[30], and a clinical

Omicron isolate (USA-lh01/2021) at a multiplicity of infection (MOI) of 0.01 and monitored viral

propagation by flow cytometry and the plaque assay. The ancestral virus [hereinafter referred to

as wild-type (WT)] and Omi-S spread fast in ACE2/TMPRSS2/Caco-2 cells, yielding 89% and

80% infected cells, respectively, at 24 hours post-infection (hpi) (**Fig. 2b**). In contrast, Omicron

replicated slower, leading to 48% infected cells at 24 hpi. A similar pattern was seen in Vero E6

cells, where 60% and 41% of cells were positive for WT and Omi-S, respectively, at 48 hpi, in

contrast to 10% positive cells for Omicron (**Fig. 2c**). The plaque assay showed that although both

Omi-S and Omicron produced lower levels of infectious virus particles compared with WT, the

viral titer of Omi-S was significantly higher than that of Omicron. In ACE2/TMPRSS2/Caco-2 cells,

Omi-S produced 5.1-fold ($p = 0.0006$) and 5.5-fold ($p = 0.0312$) more infectious particles than

Omicron at 12 hpi and 24 hpi, respectively (**Fig. 2d**). Similarly, in Vero E6 cells, the infectious

virus titers of Omi-S were 17-fold ($p = 0.0080$) and 11-fold ($p = 0.0078$) higher than that of Omicron

at 24 hpi and 48 hpi, respectively (**Fig. 2e**). The difference between viruses became less obvious

at later time points due to higher cytotoxicity caused by Omi-S compared with Omicron (**Fig. 2f**).

The higher infection efficiency of Omi-S relative to Omicron was also reflected in the plaque size;

while WT produced the largest plaques (~ 4.1 mm), the size of Omi-S plaques (~2.2 mm) was 2-

fold ($p < 0.0001$) larger than that of Omicron plaques (~1.1 mm) (**Fig. 2g**). These results indicate

that while mutations in the S protein influence the infection efficiency of Omicron, they do not fully

explain the infection behavior of Omicron in cell culture.

We next expanded our studies to lung epithelial cells, which are a major viral replication

site in patients with severe COVID-19. Accordingly, we employed human induced pluripotent stem

cell-derived lung alveolar type 2 epithelial (iAT2) cells. AT2 cells represent an essential cell

population in the distal lung and constitute one of the primary targets of SARS-CoV-2 infection[34-

36]. We infected iAT2 cells, grown as an air-liquid interface (ALI) culture, at an MOI of 2.5 and

monitored the secretion of viral progeny on the apical interface of cells at 48 hpi and 96 hpi. In

congruence with the results obtained from cell lines, WT SARS-CoV-2 produced the highest levels

of infectious virus particles (**Fig. 2h**). Among the Omi-S and Omicron, the former yielded ~5-fold

(p = 0.0008) higher infectious viral titer at 48 hpi. The viral titers for WT and Omi-S decreased at

96 hpi compared with 48 hpi due to the cytopathic effect (CPE) of infection. However, no CPE

was seen for Omicron, leading to sustained production of infectious virions. Overall, these results

corroborate the conclusion that mutations in S do not fully account for the attenuated replication

capacity of Omicron in cultured human cells.

Spike has an appreciable but minimal role in Omicron pathogenicity in K18-hACE2 mice

To examine if Omi-S exhibits higher *in vivo* fitness compared with Omicron, we investigated

the infection outcome of Omi-S relative to WT SARS-CoV-2 and Omicron in K18-hACE2 mice. In

agreement with the published literature[3,37-39], intranasal inoculation of mice (aged 12-20 weeks)

with Omicron (10^4 PFU per animal) caused no significant weight loss, whereas inoculation with

WT virus triggered a rapid decrease in body weight with all animals losing over 20% of their initial

body weight by 8 days post-infection (dpi) (**Fig. 3a**). Importantly, 80% of animals infected with

Omi-S also lost over 20% of their body weight by 9 dpi (**Fig. 3a and Extended Data Fig. 2a**). The

evaluation of clinical scores (a cumulative measure of weight loss, abnormal respiration, aberrant

appearance, reduced responsiveness, and altered behavior) also revealed a similar pattern; while

Omicron-infected mice displayed little to no signs of clinical illness, the health of those infected

with WT and Omi-S rapidly deteriorated, with the former inflicting a more severe disease (p =

0.0102) (**Fig. 3b and Extended Data Fig. 2b**). Since SARS-CoV-2 causes fatal infection in K18-

hACE2 mice[3,40,41], we leveraged this situation to compare the animal survival after viral infection.

In agreement with the results of body-weight loss and clinical score, WT and Omi-S caused

mortality rates of 100% (6/6) and 80% (8/10), respectively. In contrast, all animals infected with

Omicron survived (**Fig. 3c**). These findings indicate that the S protein is not the primary

determinant of Omicron's pathogenicity in K18-hACE2 mice.

Next, we compared the virus propagation of Omi-S with Omicron and WT SARS-CoV-2 in

the lungs of K18-hACE2 mice. The mice (12-20 weeks old) were intranasally challenged with 10^4

PFU (7 mice per virus), and their lungs were collected at 2 and 4 dpi for virological and histological

analysis. Consistent with *in vitro* findings, the infectious virus titer in the lungs of WT-infected mice

was higher than that detected in mice infected with other two viruses (**Fig. 3d**). Notably however,

Omi-S-infected mice produced 30-fold (p = 0.0286) more infectious virus particles compared with

Omicron-infected mice at 2 dpi. The titer decreased at 4 dpi for WT- and Omi-S-infected mice, yet

it showed an increasing trend for Omicron-infected animals, pointing to the possibility of mild but

persistent infection by Omicron in K18-hACE2 mice.

To evaluate the viral pathogenicity in the lungs, we performed histopathological analysis of

the lung tissue of infected K18-hACE2 mice. As previously reported[3,42], an extensive near-diffused

immunoreactivity of the SARS-CoV-2 S protein was detected in lung alveoli of mice infected with

WT virus (**Fig. 3e**). In contrast, Omi-S and Omicron infection produced localized foci of alveolar

staining with fewer foci for Omicron compared with Omi-S. The most striking phenotype was seen

in bronchiolar epithelium. While Omi-S virus caused a severe bronchiolar infection with around

15-20% of bronchioles being positive for the S protein in all mice examined at 2 dpi, less than 1%

bronchioles were S-positive in Omicron-infected mice (**Fig. 3f**). Further, bronchiolar infection was

associated with epithelial necrosis in Omi-S-infected mice, as determined through serial

hematoxylin and eosin (H&E) section analysis, whereas no histological evidence of airway injury

was observed in Omicron-infected mice (**Extended Data Fig. 3**). This suggests that the replication

of Omicron in mice lungs, particularly in bronchioles, is substantially attenuated compared with

Omi-S, supporting our conclusion that mutations in the S protein are only partially responsible for

the attenuated pathogenicity of Omicron.

Mutations in the spike RBM are major drivers of Omicron's escape from neutralization

Next, we examined if Omi-S captures the immune escape phenotype of Omicron. A large

body of literature has demonstrated extensive escape of Omicron from vaccine-induced humoral

immunity [4,10,43]. We compared the *in vitro* neutralization activity of sera obtained from vaccinated

individuals against the SARS-CoV-2 Washington isolate (USA-WA1/2020), Omi-S, and Omicron.

Sera collected within two months of the second dose of mRNA-1273 (Moderna mRNA vaccine; n

= 12) or BNT162b2 (Pfizer-BioNTech mRNA vaccine; n = 12) vaccine were included (**Extended**

Data Table 1). We performed a multicycle neutralization assay using a setting in which the virus

and neutralizing sera were present at all times, mimicking the situation in a seropositive individual.

All sera poorly neutralized Omicron, with 11.1-fold (range: 4.4- to 81.2-fold; $p < 0.0001$) lower half-

maximal neutralizing dilution (ND_{50}) for Omicron compared with WA1 (**Fig. 4a,b**). In fact, around

80% of samples failed to completely neutralize Omicron at the highest tested concentration

(**Extended Data Fig. 4**). Notably, Omi-S exhibited identical ND_{50} values to Omicron (11.5-fold

lower than that of WA1; $p < 0.0001$) (**Fig. 4a,b**), suggesting that the Omicron S protein, when

incorporated into a WT virus, behaves the same way as in Omicron.

The SARS-CoV-2 S protein comprises two domains: the S1 domain, which interacts with

the ACE2 receptor, and the S2 domain, which is responsible for membrane fusion[44]. Within the

S1 domain lie an N-terminal domain (NTD) and a receptor-binding domain (RBD), which harbors

the receptor-binding motif (RBM) that makes direct contact with the ACE2 receptor[45]. The NTD of

Omicron S carries 11 amino acid changes, including 6 deletions and one three-amino acid-long

insertion, whereas the RBD harbors 15 mutations, 10 of which are concentrated in the RBM (**Fig.**

1a,b). Both NTD and RBD host neutralizing epitopes[46-50], but the RBD is immunodominant and

represents the primary target of the neutralizing activity present in SARS-CoV-2 immune sera[50,51].

To determine if the neutralization resistance phenotype of Omicron is caused by mutations in a

particular S domain, we generated two groups of chimeric viruses. The first group comprised the

WA1 virus carrying the NTD, RBD, or RBM of Omicron (**Fig. 4c**), and the second group consisted

of Omi-S virus bearing the NTD, RBD, or RBM of WA1 (**Fig. 4d**). The neutralization assay showed

that mutations in the RBM were the major cause of Omicron's resistance to vaccine-induced

humoral immunity: replacing the RBM of WA1 with that of Omicron decreased ND_{50} by 5.4-fold (*p*

< 0.0001), and conversely, substituting the RBM of Omi-S with that of WA1 increased ND_{50} by

5.6-fold (*p* = 0.0003) (**Fig. 4c,d**). The fact that none of the RBM-swap viruses achieved the

difference of ~11-fold seen between WA1 and Omi-S suggests that mutations in other parts of S

also contribute to vaccine resistance.

To investigate if specific mutations in Omicron RBM drive vaccine escape, we

generated two additional panels of recombinant viruses, one with WA1 spike carrying Omicron

RBM mutations, either singly or in combination (**Fig. 4e**), and the other with Omicron spike lacking

the same set of mutations (**Fig. 4f**). Two WA1 mutants, mutant 3 (carrying E484A substitution)

and mutant 4 (bearing a cluster of five substitutions Q493R, G496S, Q498R, N501Y, Y505H)

exhibited a moderate but statistically significant decrease of 1.4-fold (*p* = 0.0002) and 1.8-fold (*p*

= 0.0003) in ND_{50} values, respectively, compared with WA1 (**Fig. 4e**). The opposite was observed

when these mutations were removed from Omicron S; the Omicron mutant 3 (lacking E484A

substitution) and mutant 4 (lacking Q493R, G496S, Q498R, N501Y, Y505H) had a 1.9-fold (p =

0.0082) and 3.1-fold (p = 0.0025) higher ND_{50} values compared with Omicron (**Fig. 4f**). Since

none of the mutants captured the overall phenotype of Omicron, we assume that the vaccine

escape is a cumulative effect of mutations distributed along the length of the S protein. It is

possible that mutations alter the conformation of Omicron S in such a manner that most of the

immunodominant neutralizing epitopes are disrupted and become unavailable for neutralization.

DISCUSSION

This study provides important insights into Omicron pathogenicity. We show that spike, the

single most mutated protein in Omicron, has an incomplete role in Omicron attenuation. In *in vitro*

infection assays, the Omicron spike-bearing ancestral SARS-CoV-2 (Omi-S) exhibits much higher

replication efficiency compared with Omicron. Similarly, in K18-hACE2 mice, Omi-S contrasts with

non-fatal Omicron and causes a severe disease leading to around 80% mortality. This suggests

that mutations outside of spike are major determinants of the attenuated pathogenicity of Omicron

in K18-hACE2 mice. Further studies are needed to identify those mutations and decipher their

mechanisms of action.

One potential limitation of our study is the use of K18-hACE2 mice for pathogenesis studies

instead of the primate models that have more similarities with humans[52,53]. It should however be

noted that the K18-hACE2 mouse model is a well-established model for investigating the lethal

phenotype of SARS-CoV-2[3,42,54-56]. While these mice develop lung pathology following SARS-

CoV-2 infection, mortality has been associated with central nervous system involvement due to

viral neuroinvasion[42,57]. The fact that infection with Omi-S, but not with Omicron, elicits neurologic

signs, such as hunched posture and lack of responsiveness, in K18-hACE2 mice suggests that

the neuroinvasion property is preserved in Omi-S, and the determinants of this property lie outside

of the spike protein.

We found that while the ancestral virus mainly replicates in lung alveoli and causes only

rare infection of bronchioles in K18-hACE2 mice, Omi-S with isogenic ancestral virus backbone

exhibits higher propensity to replicate in bronchiolar epithelium. This is consistent with a hamster

study demonstrating higher predilection of Omicron for bronchioles[1]. In vitro studies have also

showed that while Omicron replicates poorly in lower lung cells[58], it causes a robust infection in

bronchiolar and nasal epithelial cells[58-60]. Our findings indicate that the higher preference of

Omicron for bronchioles is dictated by mutations in the spike protein. We speculate that both Omi-

S and Omicron enter the bronchiolar epithelium of K18-hACE2 mice, yet only Omi-S replicates to

high enough levels to manifest in overt bronchiolar injury. The preference of Omicron spike for

bronchiolar epithelium is likely mediated by its improved efficiency to utilize Cathepsin B/L[58-62],

which form an active viral entry pathway in bronchioles and other airway cells[59,63]. In contrast,

SARS-CoV-2 entry into alveolar epithelial cells is mainly driven by TMPRSS2[36,64], which Omicron

spike is deficient in utilizing[60,65], leading to poor infection of these cells[3,37,58,60]. These findings

explain the higher transmission and lower lung pathology caused by Omicron.

Our study shows that mutations in the RBM of Omicron spike are the major determinants

of the viral escape from neutralizing antibodies, although mutations in other regions of spike also

contribute. Within the RBM, we identify two hotspots of mutations, which impart on Omicron spike

the ability to resist neutralization: one bearing the E484A substitution and the other harboring a

cluster of five substitutions, Q493R, G496S, Q498R, N501Y and Y505H. The E484A substitution

has been shown to escape neutralization by convalescent sera[66]. Further, structural modeling suggests that some therapeutic monoclonal antibodies establish highly stable salt bridges with the E484 residue, entirely losing their binding when this residue is changed to A or upon Q493K and Y505H changes[67]. Similarly, mapping of RBM residues that directly interact with 49 known neutralizing antibodies revealed N440, G446, S477, and T478 as low-frequently interactors, N501, Y505, and Q498 as medium-frequency interactors, and E484 and Q493 as high-frequency interactors[68], which is in line with our neutralization assay results. Interestingly, while antibody-binding potential of Omicron spike is impaired[69], its receptor-binding capacity is intact. In fact, the Omicron RBD has higher affinity for ACE2 relative to the Wuhan-Hu-1 and Delta RBDs[60]. This indicates that mutations in the Omicron spike have evolved in such a manner that they hinder antibody binding but preserve the receptor engagement. This opens up the possibility of targeting the conserved and structurally constrained regions of spike involved in ACE2 recognition for the design of broad-spectrum vaccines to control the current COVID-19 pandemic.

MATERIALS AND METHODS

Cells, antibodies, and plasmids

The cell lines were incubated at 37ºC and 5% CO_2 in a humidified incubator. Human embryonic kidney HEK293T cells (ATCC; CRL-3216), human lung adenocarcinoma A549 cells (ATCC; CCL-185), human colorectal adenocarcinoma Caco-2 cells (ATCC; HTB-37), and African green monkey kidney Vero E6 cells were maintained in DMEM (Gibco; #11995-065) containing 10% FBS and 1X non-essential amino acids. Lentiviral delivery system was used to generate cells

stably expressing human ACE2 and TMPRSS2. Mycoplasma negative status of all cell lines was

confirmed.

Anti-SARS-CoV nucleocapsid (N) protein antibody (Rockland; #200-401-A50) was used

for detection of the SARS-CoV-2 N protein by IF. Expression plasmid encoding the spike protein

of the SARS-CoV-2 Wuhan isolate, pCSII-SARS-CoV-2 F8, was a kind gift from Yoshiharu

Matsuura[32]. We replaced the Wuhan spike in this plasmid with a chemically synthesized version

of Omicron spike and called the resulting plasmid pCSII-SARS-CoV-2 F8_Omicron. The lentiviral

vectors, pLOC_hACE2_PuroR and pLOC_hTMPRSS2_BlastR, containing human ACE2 and

TMPRSS2, respectively, have been previously described[33].

Omicron stock preparation and titration

All procedures were performed in a biosafety level 3 (BSL3) facility at the National

Emerging Infectious Diseases Laboratories of the Boston University using biosafety protocols

approved by the institutional biosafety committee (IBC). The SARS-CoV-2 BA.1 Omicron virus

stock was generated in ACE2/TMPRSS2/Caco-2 cells. Briefly, 5×10^5 cells, grown overnight in

DMEM/10%FBS/1X NEAA in one well of a 6-well plate, were inoculated with the collection

medium in which the nasal swab from a SARS-CoV-2 patient was immersed. The swab material

was obtained from the Department of Public Health, Massachusetts, and it contained the

sequence-verified Omicron virus (NCBI accession number: OL719310). Twenty-four hours after

infecting cells, the culture medium was replaced with 2 ml of DMEM/2%FBS/1X NEAA and the

cells were incubated for another 72h, at which point the CPE became visible. The culture medium

was harvested, passed through a 0.45 μ filter, and kept at -80ºC as a P0 virus stock. To generate

a P1 stock, we infected 1×10^7 ACE2/TMPRSS2/Caco-2 cells, seeded the day before in a T175

flask, with the P0 virus at an MOI of 0.01. The next day, the culture medium was changed to 25

ml of 2% FBS-containing medium. Three days later, when the cells exhibited excessive CPE, the

culture medium was harvested, passed through a 0.45 μ filter, and stored at -80ºC as a P1 stock.

To titrate the virus stock, we seeded ACE2/TMPRSS2/Caco-2 cells into a 12-well plate at

a density of 2 x 10^5 cells per well. The next day, the cells were incubated with serial 10-fold

dilutions of the virus stock (250 μl volume per well) for 1h at 37ºC, overlayed with 1 ml per well of

medium containing 1:1 mixture of 2X DMEM/4% FBS and 1.2% Avicel (DuPont; RC-581), and

incubated at 37ºC for another three days. To visualize the plaques, the cell monolayer was fixed

with 4% paraformaldehyde and stained with 0.1% crystal violet, with both fixation and staining

performed at room temperature for 30 minutes each. The number of plaques were counted and

the virus titer was calculated.

Recombinant SARS-CoV-2 generation by CPER

SARS-CoV-2 recombinant viruses were generated by using a modified form of the recently

published CPER protocol[32,70]. Full-length SARS-CoV-2 cDNA cloned onto a bacterial artificial

chromosome (BAC)[30] was used as a template to amplify the viral genome into eight overlapping

fragments (F1, F2, F3, F4, F5, F6, F7, and F9). The pCSII-SARS-CoV-2 F8 and pCSII-SARS-

CoV-2 F8_Omicron plasmids, which were used to generate spike mutants, served as templates

for amplification of fragment 8 (F8). A UTR linker containing a hepatitis delta virus ribozyme

(HDVr), the bovine growth hormone polyadenylation signal sequence (BGH-polyA), and a

cytomegalovirus (CMV) promoter was cloned onto a pUC19 vector and used as a template to

amplify the linker sequence. The 5' termini of all ten DNA fragments (F1-F9 and the linker) were

phosphorylated by using T4 PNK (NEB; #M0201), and the equimolar amounts (0.05 pmol each)

of the resulting fragments were subjected to a CPER reaction in a 50 μl volume using 2 μl of

PrimeStar GXL DNA polymerase (Takara Bio; #R050A). The following cycling conditions were

used for CPER: an initial denaturation at 98°C for 2 min; 35 cycles of denaturation at 98°C for 10

s, annealing at 55°C for 15 s, and extension at 68°C for 15 min; and a final extension at 68°C for

15 min. The nicks in the circular product were sealed by using DNA ligase.

To transfect cells with the CPER product, we seeded ACE2/TMPRSS2/Caco-2 cells into a

6-well plate at a density of 5 x10^5 cells per well. The transfection mix was prepared by mixing 26

μl of the original 52 μl CPER reaction volume with 250 μl of Opti-MEM (Thermo Fisher Scientific;

#31985070) and 6 μl of TransIT-X2 Dynamic Delivery System (Mirus Bio; #MIR 6000). Following

incubation at room temperature for 25 min, the transfection mix was added to the cells. The next

day, the culture medium was replaced with fresh DMEM containing 2% FBS. The CPE became

visible in 3-4 days, at which point the culture medium was collected and stored as a P0 virus

stock. The P0 stock was used for experiments described in this manuscript. The spike region of

all CPER-generated viruses was sequenced by either Sanger sequencing or next generation

sequencing to confirm the presence of desired and the absence of adventitious changes.

SARS-CoV-2 neutralization assay

For neutralization assays, initial 1:10 dilutions of plasma, obtained from individuals

who received two shots of either Moderna or Pfizer mRNA-based SARS-CoV-2 vaccine, were

five-fold serial diluted in Opti-MEM over seven or eight dilutions. These plasma dilutions were

then mixed at a 1:1 ratio with 1.25 x 10^4 infectious units of SARS-CoV-2 and incubated for 1h at

37°C. Thereafter, 100 μl of this mixture was directly applied to ACE2/A549 cells seeded the

previous day in poly-L-lysine-coated 96-well plates at a density of 2.5 x 10^4 cells per well in 100

μl volume. Thus, the final starting dilution of plasma was 1:20 and the final MOI was 0.5. The cells

were incubated at 37°C for 24h, after which they were fixed and stained with an anti-nucleocapsid

antibody. When PBS instead of plasma was used as a negative control, these infection conditions

resulted in around 40-50% infected cells at 24 hpi.

Generation and infection of iAT2 cells

The detailed protocol for generation of human iPSC-derived alveolar epithelial type II cells

(iAT2s) has been published in our recent papers[36,71]. The air-liquid interface (ALI) cultures were

established by preparing single cell suspensions of iAT2 3D sphere cultures grown in Matrigel.

Briefly, Matrigel droplets containing iAT2 spheres were dissolved in 2 mg/ml dispase (Sigma) and

the spheres were dissociated in 0.05% trypsin (GIBCO) to generate a single-cell suspension. 6.5

mm Transwell inserts (Corning) were coated with dilute Matrigel (Corning) in accordance with the

manufacturer's protocol. Single-cell iAT2s were plated on Transwells at a density of 520,000

cells/cm2 in 100 μl of CK+DCI medium containing 10 μM of Rho-associated kinase inhibitor ("Y";

Sigma Y-27632). 600 μl of this medium was added to the basolateral compartment. 24h after

plating, the basolateral medium was changed with fresh CK+DCI+Y medium. 48h after plating,

the apical medium was aspirated to initiate ALI culture. 72h after plating, basolateral medium was

replaced with CK+DCI medium to remove the rho-associated kinase inhibitor. Basolateral medium

was changed every two days thereafter. The detailed composition of CK+DCI medium is provided

in our previous publications[36,71].

iAT2 cells in ALI cultures were infected with purified SARS-CoV-2 stock at an MOI of 2.5

based on the titration done on ACE2/TMPRSS2/Caco-2 cells. For infection, 100 μl of inoculum

prepared in 1X PBS (or mock-infected with PBS-only) was added to the apical chamber of each

Transwell and incubated for 2h at 37°C followed by the removal of the inoculum and washing of

the apical side three times with 1X PBS (100 μl/wash). The cells were incubated for two or four

days, after which the newly released virus particles on the apical side were collected by adding

100 μl of 1X PBS twice to the apical chamber and incubating at 37°C for 15 min. The number of

infectious virus particles in the apical washes were measured by the plaque assay on

ACE2/TMPRSS2/Caco-2 cells. For flow cytometry, iAT2 cells were detached by adding 0.2 ml

Accutase (Sigma; #A6964) apically and incubated at room temperature for 15 min. The detached

cells were pelleted by low-speed centrifugation, fixed in 10% formalin, and stained with anti-

SARS-CoV-2 N antibody.

Mice maintenance and approvals

Mice was maintained in a facility accredited by the Association for the Assessment and

Accreditation of Laboratory Animal Care (AAALAC). Animal studies were performed following the

recommendations in the Guide for the Care and Use of Laboratory Animals of the National

Institutes of Health. The protocols were approved by the Boston University Institutional Animal

Care and Use Committee (IACUC). Heterozygous K18-hACE2 C57BL/6J mice (Strain 2B6.Cg-

Tg(K18-ACE2)2Prlmn/J) were purchased from the Jackson Laboratory (Jax, Bar Harbor, ME).

Animals were housed in ventilated cages (Tecniplast, Buguggiate, Italy) and maintained on a

12:12 light cycle at 30-70% humidity, ad-libitum water, and standard chow diets (LabDiet, St.

Louis, MO).

Mice infection

Twelve to twenty weeks old male and female K18-hACE2 mice were inoculated intranasally

with 10^4 PFU of SARS-CoV-2 in 50 μl of sterile 1X PBS. The inoculations were performed under

1-3% isoflurane anesthesia. Twenty-six mice (6 for WT, 10 for Omi-S, and 10 for Omicron) were

enrolled in a 14-day survival study, and another 42 mice (14 for each of the WT, Omi-S, and

Omicron viruses) were used for virological and histological analysis of infected lungs. During the

survival study, the animals were monitored for body weight, respiration, general appearance,

responsiveness, and neurologic signs. An IACUC-approved clinical scoring system was used to

monitor disease progression and define humane endpoints. The score of 1 was given for each of

the following situations: body weight, 10-19% loss; respiration, rapid and shallow with increased

effort; appearance, ruffled fur and/or hunched posture; responsiveness, low to moderate

unresponsiveness; and neurologic signs, tremors. The sum of these individual scores constituted

the final clinical score. Animals were considered moribund and humanly euthanized in case of

weight loss greater than or equal to 20%, or if they received a clinical score of 4 or greater for two

consecutive days. Body weight and clinical score were recorded once per day for the duration of

the study. For the purpose of survival curves, animals euthanized on a given day were counted

dead the day after. Animals found dead in cage were counted dead on the same day. For

euthanization, an overdose of ketamine was administered followed by a secondary method of

euthanization.

For quantification of SARS-CoV-2 infectious particles in lungs by the plaque assay, lung

tissues were collected in 600 μl of RNAlater stabilization solution (ThermoFisher Scientific;

#AM7021) and stored at -80°C until analysis. 20-40 mg of tissue was placed in a tube containing

600 μl of OptiMEM and a 5 mm stainless steel bead (Qiagen; #69989) and homogenized in the

Qiagen TissueLyser II by two dissociation cycles (1,800 oscillations/minute for 2 minutes) with a

one-minute interval between cycles. The homogenate was centrifuged at 15,000 xg for 10 minutes

at room temperature and the supernatant was transferred to a new tube. Ten-fold serial dilutions

of this supernatant were used for the plaque assay on ACE2/TMPRSS2/Caco-2 cells, as

described above.

For IHC and histologic analysis, the insufflated whole lung tissues were inactivated in 10%

neutral buffered formalin at a 20:1 fixative to tissue ratio for a minimum of 72h before removal

from BSL3 in accordance with an approved IBC protocol. Tissues were subsequently processed,

embedded in paraffin and five-micron sections stained with hematoxylin and eosin (H&E) following

standard histological procedures. IHC was performed using a Ventana BenchMark Discovery

Ultra autostainer (Roche Diagnostics, USA). An anti-SARS-CoV-2 S antibody (Cell Signaling

technologies: clone E5S3V) that showed equivalent immunoreactivity against WT and Omicron

spike was used to identify virus-infected cells. Negative and positive controls for IHC included

blocks of uninfected and SARS-CoV-2-infected Vero E6 cells, respectively.

Flow cytometry

For flow cytometry, fixed cells were permeabilized in 1x permeabilization buffer

(ThermoFisher Scientific; #00-5523-00) and stained with SARS-CoV-2 nucleocapsid antibody

(Rockland; #200-401-A50, 1:1,000), followed by donkey anti-rabbit IgG-AF647 secondary

antibody (ThermoFisher Scientific; #A-31573). Gating was based on uninfected stained control

cells. The extent of staining was quantified using a BD LSR II flow cytometer (BD Biosciences,

CA), and the data were analyzed with FlowJo v10.6.2 (FlowJo, Tree Star Inc).

Immunofluorescence

Immunofluorescence was performed as described in our previous publication[33]. Briefly,

virus-infected cells were fixed in 4% paraformaldehyde and permeabilized in a buffer containing

0.1% Triton X-100 prepared in PBS. Following blocking in a buffer containing 0.1% Triton X-100,

10% goat serum, and 1% BSA, the cells were incubated overnight at 4°C with anti-SARS-CoV

Nucleocapsid antibody (1:2,000 dilution). The cells were then stained with Alexa Fluor 568-

conjugated goat anti-rabbit secondary antibody (1:1000 dilution) (Invitrogen; #A11008) in the dark

at room temperature for 1h and counterstained with DAPI. Images were captured using the

ImageXpress Micro Confocal (IXM-C) High-Content Imaging system (Molecular Devices) with a

4x S Fluor objective lens at a resolution of 1.7 micron/pixel in the DAPI (excitation: 400 nm/40 nm,

emission: 447 nm/60 nm) and TexasRed (excitation: 570nm/80nm, emission: 624nm/40nm)

channels. Both channels were used to establish their respective laser autofocus offsets. The

images were analyzed using MetaXpress High Content Image Acquisition and Analysis software

(Molecular Devices). First, the images were segmented using the CellScoring module. The

objects between 7 and 20 microns in diameter and greater than 1800 gray level units in intensity

were identified and classified as nuclei. Positive cells were taken as nuclei having TexasRed

signal of 1500 gray level units or above within 10 to 20 microns of each nucleus. The remaining

objects were set to negative cells. From these objects, the following readouts were measured and

used for downstream analysis: Total number of positive and negative cells, total area of positive

cells, and integrated intensity in the TexasRed channel for positive cells. To calculate the 50%

neutralizing dilution (ND_{50}), we performed a non-linear regression curve fit analysis using Prism 9

software (GraphPad).

ACKNOWLEDGEMENT

We thank Dr. Yoshiharu Matsuura from Osaka University, Japan, for providing the pCSII-

SARS-CoV-2 F8 plasmid; the Department of Public Health, Massachusetts, for providing the

clinical specimen containing Omicron virus; and the ICCB-Longwood Screening Facility of

Harvard Medical School for assistance with IF image acquisition and analysis. This work was

supported by Boston University startup funds (to MS and FD), National Institutes of Health, NIAID

grants R01 AI159945 (to SB and MS) and R37 AI087846 (to MUG), NIH SIG grants S10-

OD026983 and SS10-OD030269 (to NAC), Peter Paul Career Development Award (to FD), and

BMBF SenseCoV2 01KI20172A (AE) and DFG Fokus COVID-19, EN 423/7-1 (AE). We thank the

Clinical & Translational Science Institute (CTSI; 1UL1TR001430) and Evans Center for

Interdisciplinary Biomedical Research at Boston University School of Medicine for their support of

the Affinity Research Collaborative on 'Respiratory Viruses: A Focus on COVID-19'.

AUTHOR CONTRIBUTIONS

M.S. conceptualized the study. DYC, AHT, DK, CVC, NK, HLC, FD, and MS performed

experiments. GL and MUG established and provided the modified CPER system. NAC performed

histopathologic and IHC analysis of mouse lungs. SCB and MB provided scientific input and

helped secure funds. AH and AE provided BAC harboring the SARS-CoV-2 genome. JHC

provided the Omicron isolate. YK provided plasma samples. MS wrote the manuscript, which was

read, edited, and approved by all authors.

REFERENCES

Suzuki, R. *et al.* Attenuated fusogenicity and pathogenicity of SARS-CoV-2 Omicron variant. *Nature*, doi:10.1038/s41586-022-04462-1 (2022).

Halfmann, P. J. *et al.* SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters. *Nature*, doi:10.1038/s41586-022-04441-6 (2022).

Shuai, H. *et al.* Attenuated replication and pathogenicity of SARS-CoV-2 B.1.1.529 Omicron. *Nature*, doi:10.1038/s41586-022-04442-5 (2022).

Dejnirattisai, W. *et al.* SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape from neutralizing antibody responses. *Cell* **185**, 467-484 e415, doi:10.1016/j.cell.2021.12.046 (2022).

VanBlargan, L. A. *et al.* An infectious SARS-CoV-2 B.1.1.529 Omicron virus escapes neutralization by therapeutic monoclonal antibodies. *Nat Med*, doi:10.1038/s41591-021-01678-y (2022).

Planas, D. *et al.* Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. *Nature*, doi:10.1038/s41586-021-04389-z (2021).

Liu, L. *et al.* Striking antibody evasion manifested by the Omicron variant of SARS-CoV-2. *Nature*, doi:10.1038/s41586-021-04388-0 (2021).

Hoffmann, M. *et al.* The Omicron variant is highly resistant against antibody-mediated neutralization: Implications for control of the COVID-19 pandemic. *Cell* **185**, 447-456 e411, doi:10.1016/j.cell.2021.12.032 (2022).

WHO. *Tracking SARS-CoV-2 variants*, <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/> (2022).

Cele, S. *et al.* Omicron extensively but incompletely escapes Pfizer BNT162b2 neutralization. *Nature* **602**, 654-656, doi:10.1038/s41586-021-04387-1 (2022).

Gu, H. *et al.* Probable Transmission of SARS-CoV-2 Omicron Variant in Quarantine Hotel, Hong Kong, China, November 2021. *Emerg Infect Dis* **28**, 460-462, doi:10.3201/eid2802.212422 (2022).

Iuliano, A. D. *et al.* Trends in Disease Severity and Health Care Utilization During the Early Omicron Variant Period Compared with Previous SARS-CoV-2 High Transmission Periods - United States, December 2020-January 2022. *MMWR Morb Mortal Wkly Rep* **71**, 146-152, doi:10.15585/mmwr.mm7104e4 (2022).

CDC. *COVID Data Tracker*, <https://covid.cdc.gov/covid-data-tracker/#variant-proportions> (2022).

CoVariants. *CoVariants*, <https://covariants.org> (2022).

Taylor, L. Covid-19: Omicron drives weekly record high in global infections. *BMJ* **376**, o66, doi:10.1136/bmj.o66 (2022).

Desingu, P. A. & Nagarajan, K. SARS-CoV-2 Omicron variant is spreading in different parts of the world in three different trends. *J Med Virol*, doi:10.1002/jmv.27646 (2022).

WHO. Statement on Omicron sublineage BA.2. (2022).

Desingu, P. A. & Nagarajan, K. Omicron variant losing its critical mutations in the receptor-binding domain. *J Med Virol*, doi:10.1002/jmv.27667 (2022).

Desingu, P. A., Nagarajan, K. & Dhama, K. Emergence of Omicron third lineage BA.3 and its importance. *J Med Virol*, doi:10.1002/jmv.27601 (2022).

Takashita, E. *et al.* Efficacy of Antibodies and Antiviral Drugs against Covid-19 Omicron Variant. *N Engl J Med*, doi:10.1056/NEJMc2119407 (2022).

21   Schmidt, F. *et al.* Plasma Neutralization of the SARS-CoV-2 Omicron Variant. *N Engl J Med* **386**,
599-601, doi:10.1056/NEJMc2119641 (2022).
22   ImperialCollegeLondon. *Hospitalisation risk for Omicron cases in England*,
<https://www.imperial.ac.uk/mrc-global-infectious-disease-analysis/covid-19/report-50-severity-
omicron/> (2022).
23   Madhi, S. A. *et al.* Population Immunity and Covid-19 Severity with Omicron Variant in South
Africa. *N Engl J Med*, doi:10.1056/NEJMoa2119658 (2022).
24   Sigal, A. Milder disease with Omicron: is it the virus or the pre-existing immunity? *Nat Rev*
Immunol **22**, 69-71, doi:10.1038/s41577-022-00678-4 (2022).
25   Wolter, N. *et al.* Early assessment of the clinical severity of the SARS-CoV-2 omicron variant in
South Africa: a data linkage study. *Lancet* **399**, 437-446, doi:10.1016/S0140-6736(22)00017-4
(2022).
26   Dyer, O. Covid-19: Omicron is causing more infections but fewer hospital admissions than delta,
South African data show. *BMJ* **375**, n3104, doi:10.1136/bmj.n3104 (2021).
27   Haseltine, W. A. in *Understanding Omicron: Changes In The Spike Protein And Beyond And What*
They Portend   (2021).
28   Smyth, D. S. *et al.* Tracking cryptic SARS-CoV-2 lineages detected in NYC wastewater. *Nat*
Commun **13**, 635, doi:10.1038/s41467-022-28246-3 (2022).
29   Kirby, A. E. *et al.* Notes from the Field: Early Evidence of the SARS-CoV-2 B.1.1.529 (Omicron)
Variant in Community Wastewater - United States, November-December 2021. *MMWR Morb*
Mortal Wkly Rep **71**, 103-105, doi:10.15585/mmwr.mm7103a5 (2022).
30   Herrmann, A. *et al.* Cloning of a Passage-Free SARS-CoV-2 Genome and Mutagenesis Using Red
Recombination. *Int J Mol Sci* **22**, doi:10.3390/ijms221910188 (2021).
31   Amarilla, A. A. *et al.* A versatile reverse genetics platform for SARS-CoV-2 and other positive-
strand RNA viruses. *Nat Commun* **12**, 3431, doi:10.1038/s41467-021-23779-5 (2021).
32   Torii, S. *et al.* Establishment of a reverse genetics system for SARS-CoV-2 using circular
polymerase extension reaction. *Cell Rep* **35**, 109014, doi:10.1016/j.celrep.2021.109014 (2021).
33   Chen, D. Y. *et al.* SARS-CoV-2 Disrupts Proximal Elements in the JAK-STAT Pathway. *J Virol*
95, e0086221, doi:10.1128/JVI.00862-21 (2021).
34   Mulay, A. *et al.* SARS-CoV-2 infection of primary human lung epithelium for COVID-19 modeling
and drug discovery. *Cell Rep* **35**, 109055, doi:10.1016/j.celrep.2021.109055 (2021).
35   Carcaterra, M. & Caruso, C. Alveolar epithelial cell type II as main target of SARS-CoV-2 virus
and COVID-19 development via NF-Kb pathway deregulation: A physio-pathological theory. *Med*
Hypotheses **146**, 110412, doi:10.1016/j.mehy.2020.110412 (2021).
36   Huang, J. *et al.* SARS-CoV-2 Infection of Pluripotent Stem Cell-Derived Human Lung Alveolar
Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response. *Cell Stem Cell* **27**, 962-
973 e967, doi:10.1016/j.stem.2020.09.013 (2020).
37   Halfmann, P. J. *et al.* SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters.
Nature, doi:10.1038/s41586-022-04441-6 (2022).
38   Winkler, E. S. *et al.* SARS-CoV-2 infection of human ACE2-transgenic mice causes severe lung
inflammation and impaired function. *Nat Immunol* **21**, 1327-1335, doi:10.1038/s41590-020-0778-
2 (2020).
39   Chen, R. E. *et al.* In vivo monoclonal antibody efficacy against SARS-CoV-2 variant strains. *Nature*
596, 103-108, doi:10.1038/s41586-021-03720-y (2021).
40   Cecon, E. *et al.* Therapeutic potential of melatonin and melatonergic drugs on K18-hACE2 mice
infected with SARS-CoV-2. *J Pineal Res* **72**, e12772, doi:10.1111/jpi.12772 (2022).

41   Lutz, C., Maher, L., Lee, C. & Kang, W. COVID-19 preclinical models: human angiotensin-converting enzyme 2 transgenic mice. *Hum Genomics* **14**, 20, doi:10.1186/s40246-020-00272-6 (2020).

42   Mariano Carossino, D. K., Aoife K. O'Connell, Paige Montanaro, Anna E. Tseng, Hans P. Gertje, Kyle A. Grosz, Maria Ericsson, Bertrand R. Huber, Susanna A. Kurnick, Saravanan Subramaniam, Thomas A. Kirkland, Joel R. Walker, Kevin P. Francis, Alexander D. Klose, Neal Paragas, Markus Bosmann, Mohsan Saeed, Udeni B. R. Balasuriya, Florian Douam, Nicholas A. Crossland Fatal Neurodissemination and SARS-CoV-2 Tropism in K18-hACE2 Mice Is Only Partially Dependent on hACE2 Expression. *Viruses* **14** (2022).

43   Ai, J. *et al.* Omicron variant showed lower neutralizing sensitivity than other SARS-CoV-2 variants to immune sera elicited by vaccines after boost. *Emerg Microbes Infect* **11**, 337-343, doi:10.1080/22221751.2021.2022440 (2022).

44   Huang, Y., Yang, C., Xu, X. F., Xu, W. & Liu, S. W. Structural and functional properties of SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19. *Acta Pharmacol Sin* **41**, 1141-1149, doi:10.1038/s41401-020-0485-4 (2020).

45   Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **581**, 215-220, doi:10.1038/s41586-020-2180-5 (2020).

46   Cerutti, G. *et al.* Potent SARS-CoV-2 neutralizing antibodies directed against spike N-terminal domain target a single supersite. *Cell Host Microbe* **29**, 819-833 e817, doi:10.1016/j.chom.2021.03.005 (2021).

47   Chi, X. *et al.* A neutralizing human antibody binds to the N-terminal domain of the Spike protein of SARS-CoV-2. *Science* **369**, 650-655, doi:10.1126/science.abc6952 (2020).

48   Voss, W. N. *et al.* Prevalent, protective, and convergent IgG recognition of SARS-CoV-2 non-RBD spike epitopes. *Science* **372**, 1108-1112, doi:10.1126/science.abg5268 (2021).

49   Premkumar, L. *et al.* The receptor binding domain of the viral spike protein is an immunodominant and highly specific target of antibodies in SARS-CoV-2 patients. *Sci Immunol* **5**, doi:10.1126/sciimmunol.abc8413 (2020).

50   Ju, B. *et al.* Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* **584**, 115-119, doi:10.1038/s41586-020-2380-z (2020).

51   Piccoli, L. *et al.* Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell* **183**, 1024-1042 e1021, doi:10.1016/j.cell.2020.09.037 (2020).

52   Vogel, A. B. *et al.* BNT162b vaccines protect rhesus macaques from SARS-CoV-2. *Nature* **592**, 283-289, doi:10.1038/s41586-021-03275-y (2021).

53   Chang, M. C., Hild, S. & Grieder, F. Nonhuman primate models for SARS-CoV-2 research: Consider alternatives to macaques. *Lab Anim (NY)* **50**, 113-114, doi:10.1038/s41684-021-00755-6 (2021).

54   Oladunni, F. S. *et al.* Lethality of SARS-CoV-2 infection in K18 human angiotensin-converting enzyme 2 transgenic mice. *Nat Commun* **11**, 6122, doi:10.1038/s41467-020-19891-7 (2020).

55   Yinda, C. K. *et al.* K18-hACE2 mice develop respiratory disease resembling severe COVID-19. *PLoS Pathog* **17**, e1009195, doi:10.1371/journal.ppat.1009195 (2021).

56   Gan, E. S. *et al.* A mouse model of lethal respiratory dysfunction for SARS-CoV-2 infection. *Antiviral Res* **193**, 105138, doi:10.1016/j.antiviral.2021.105138 (2021).

57   Kumari, P. *et al.* Neuroinvasion and Encephalitis Following Intranasal Inoculation of SARS-CoV-2 in K18-hACE2 Mice. *Viruses* **13**, doi:10.3390/v13010132 (2021).

58   Hui, K. P. Y. *et al.* SARS-CoV-2 Omicron variant replication in human bronchus and lung ex vivo.
Nature, doi:10.1038/s41586-022-04479-6 (2022).
59   Thomas P. Peacock, J. C. B., Jie Zhou, Nazia Thakur, Joseph Newman, Ruthiran Kugathasan,
Ksenia Sukhova, Myrsini Kaforou, Dalan Bailey, Wendy S. Barclay. The SARS-CoV-2 variant,
Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently uses the
endosomal route of entry. *bioRxiv*, doi:https://doi.org/10.1101/2021.12.31.474653 (2022).
60   Meng, B. *et al.* Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts tropism and
fusogenicity. *Nature*, doi:10.1038/s41586-022-04474-x (2022).
61   Du, X. *et al.* Omicron adopts a different strategy from Delta and other variants to adapt to host.
Signal Transduct Target Ther **7**, 45, doi:10.1038/s41392-022-00903-5 (2022).
62   Pranesh Padmanabhan, N. M. D. Evidence of increased Cathepsin B/L and decreased TMPRSS2
usage   for   cell   entry   by   the   SARS-CoV-2   Omicron   variant.   *bioRxiv*,
doi:https://doi.org/10.1101/2022.01.13.476267 (2022).
63   Kawase, M., Shirato, K., van der Hoek, L., Taguchi, F. & Matsuyama, S. Simultaneous treatment
of human bronchial epithelial cells with serine and cysteine protease inhibitors prevents severe acute
respiratory syndrome coronavirus entry. *J Virol* **86**, 6537-6545, doi:10.1128/JVI.00094-12 (2012).
64   Grau-Exposito, J. *et al.* Evaluation of SARS-CoV-2 entry, inflammation and new therapeutics in
human lung tissue cells. *PLoS Pathog* **18**, e1010171, doi:10.1371/journal.ppat.1010171 (2022).
65   Zhao, H. *et al.* SARS-CoV-2 Omicron variant shows less efficient replication and fusion activity
when compared with Delta variant in TMPRSS2-expressed cells. *Emerg Microbes Infect* **11**, 277-
283, doi:10.1080/22221751.2021.2023329 (2022).
66   Liu, Z. *et al.* Identification of SARS-CoV-2 spike mutations that attenuate monoclonal and serum
antibody neutralization. *Cell Host Microbe* **29**, 477-488 e474, doi:10.1016/j.chom.2021.01.014
(2021).
67   Shah, M. & Woo, H. G. Omicron: A Heavily Mutated SARS-CoV-2 Variant Exhibits Stronger
Binding to ACE2 and Potently Escapes Approved COVID-19 Therapeutic Antibodies. *Front
Immunol* **12**, 830527, doi:10.3389/fimmu.2021.830527 (2021).
68   Ye, G., Liu, B. & Li, F. Cryo-EM structure of a SARS-CoV-2 omicron spike protein ectodomain.
Nat Commun **13**, 1214, doi:10.1038/s41467-022-28882-9 (2022).
69   Carreno, J. M. *et al.* Activity of convalescent and vaccine serum against SARS-CoV-2 Omicron.
Nature **602**, 682-688, doi:10.1038/s41586-022-04399-5 (2022).
70   Tamura, T. *et al.* Generation and characterization of genetically and antigenically diverse infectious
clones   of   dengue   virus   serotypes   1-4.   *Emerg   Microbes   Infect*   **11**,   227-239,
doi:10.1080/22221751.2021.2021808 (2022).
71   Jacob, A. *et al.* Derivation of self-renewing lung alveolar epithelial type II cells from human
pluripotent stem cells. *Nat Protoc* **14**, 3303-3332, doi:10.1038/s41596-019-0220-0 (2019).
625

FIGURE LEGENDS

Fig. 1: Generating Recombinant SARS-Co-2 by CPER. a, Schematic overview of mutations in

Omicron spike (in comparison to the SARS-CoV-2 Wuhan-Hu-1 isolate; NCBI accession number:

NC_045512). Numbering is based on Wuhan-Hu-1 sequence. Mutations not reported in previous

variants of concern are shown in red. NTD, N-terminal domain; RBD, receptor-binding domain;

RBM, receptor-binding motif. **b,** Location of Omicron mutations on the trimeric spike protein.

Domains are colored according to a. **c,** Schematic of recombinant SARS-CoV-2 generated by

CPER. S, spike; N, nucleocapsid. **d,** ACE2/TMPRSS2/Caco-2 cells transfected with the SARS-

CoV-2 CPER product were stained with an anti-nucleocapsid antibody on indicated days post-

transfection. DAPI was used to stain the cell nuclei. NC, negative control generated by omitting

Fragment 9 from the CPER reaction. **e,** Virus titer in the culture medium of the transfected cells

at indicated days post-transfection, as measured by the plaque assay. The data are plotted as

mean ± SEM of two experimental repeats.

Fig. 2: Effect of spike on *in vitro* growth kinetics of Omicron. a, Schematic of viruses used in

this figure. S, spike; N, nucleocapsid. **b-e,** ACE2/TMPRSS2/Caco-2 and Vero E6 cells were

infected at an MOI of 0.01, and the percentage of nucleocapsid (N)-positive cells (n = 6) (**b,c**) and

levels of infectious virus production (n = 3) (**d,e**) were determined by flow cytometry and the

plaque assay, respectively. **f,** The cell viability of SARS-CoV-2-infected ACE2/TMPRSS2/Caco-2

cells (MOI of 0.1) was quantified by the CellTiter-Glo assay at indicated time points. The *P* values

reflect a statistically significant difference between Omi-S and Omicron. **g,** Plaque sizes. Left,

representative images of plaques on ACE2/TMPRSS2/Caco-2 cells. Right, diameter of plaques

is plotted as mean ± SD of 20 plaques per virus. **h,** Human induced pluripotent stem cell-derived

alveolar type 2 epithelial cells were infected at an MOI of 2.5 for 48h or 96h. The apical side of

cells was washed with 1X PBS and the levels of infectious virus particle were measured by the

plaque assay. n = 4. Data are mean ± SD from the indicated number of biological replicates.

Experiments were repeated twice, with each experimental repeat containing 2 (**h**) or 3 (**b-g**)

replicates. p values were calculated by a two-tailed, unpaired t-test with Welch's correction. *p

<0.05, **p <0.01, ***p <0.001, and ****p < 0.0001; ns, not significant.

Fig. 3: Role of spike in Omicron pathogenicity. a-c, Male and female K18-hACE2 mice (aged
12-20 weeks) were intranasally inoculated with 1 x 10^4 PFU of WT (n = 6), Omi-S (n = 10), or
Omicron (n = 10). Two independently generated virus stocks were used in this experiment. The
body weight (**a**), clinical score (**b**), and survival (**c**) were monitored daily for 14 days. Animals
losing 20% of their initial body weight were euthanized. **d,e,** K18-hACE2 mice were intranasally
inoculated with 1 x 10^4 PFU of WT (n = 7), Omi-S (n = 7), and Omicron (n = 7). Lung samples of
the infected mice were collected at 2 or 4 dpi to determine the viral titer (n = 4) (**d**) or for
immunohistochemistry (IHC) detection of the S protein (n = 3) (**e**). In e, representative images of
IHC staining for the detection of the SARS-CoV-2 S protein (brown color) in alveoli (arrows) and
bronchioles (arrowheads) in the lungs of the infected mice at 2 dpi are shown. (Scale bar = 100
μm). **f,** The percentage of S-positive bronchioles in the lungs of infected mice. Each dot represents
an infected animal. Data are presented as mean ± SD from the indicated number of biological
replicates. Statistical significance was determined using two-tailed, unpaired t-test with Welch's
correction (**a,b,d,f**) and log-rank (Mantel-Cox) test (**c**). *p <0.05, **p <0.01, ***p <0.001, and ****p
< 0.0001; ns, not significant.

Fig. 4: Role of spike in immune resistance of Omicron. a, ND_{50} values for WA1, Omi-S, and
Omicron in sera from individuals who received two shots of Moderna (donor 1-12) or Pfizer (donor
13-24) vaccine (further details of sera are provided in Extended Data Table 1; individual curves
are shown in Extended Data Fig. 4). **b,** Trajectories of ND_{50} values against WA1, Omi-S, and
Omicron (the data from a is plotted). Fold-change in ND_{50} values is indicated. **c,d,e,f,** Schematic

of the chimeric (**top panels; c,d**) and mutant (**top panels; e,f**) viruses. The amino acid numbering

for WA1 mutants in e is based on the WA1 spike sequence, whereas the numbering for Omicron

mutants in f is based on the Omicron spike sequence. Six of the 24 sera (three from Moderna and

three from Pfizer) were tested. Each serum sample is represented by a dot of specific color. The

data are plotted as fold-change of the parental virus. Statistical significance was determined using

a two-tailed, unpaired t test with Welch's correction. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$; ns, not significant.

EXTENDED DATA FIGURES

Extended Data Fig. 1: Schematic representation of CPER to generate recombinant SARS-CoV-2. The SARS-CoV-2 genome was amplified into nine overlapping fragments. These fragments and a linker (containing a hepatitis delta virus ribozyme, a poly-A signal, and a CMV promoter) were treated with PNK to phosphorylate 5' ends. The 5'-end phosphorylated fragments were then stitched together by CPER, and the nicks in the resulting circular DNA molecule were closed by treatment with DNA ligase. The CPER product was transfected into cells to rescue virus particles.

Extended Data Fig. 2: Clinical signs of Omi-S-infected mice. K18-hACE2 mice ($n = 10$) inoculated intranasally with 1×10^4 PFU of Omi-S and described in Fig. 3a-c were monitored for body weight (**a**) and clinical score (**b**). Animals losing 20% of their body weight (8 out of 10) were euthanized. The surviving animals did not show any signs of distress.

Extended Data Fig. 3: Lung pathology induced by Omi-S. The lungs of the male and female K18-hACE2 mice intranasally inoculated with 1×10^4 PFU of WT, Omi-S, and Omicron were collected at 2 dpi for histological analysis. **a,** Representative images of hematoxylin and eosin

(H&E) staining for the detection of bronchiolar damage in the lungs of the infected mice. The

bronchiolar epithelial necrosis is shown with arrows. Note that the necrosis was no longer evident

at 4 dpi in any cohort. **b,** Immunohistochemistry (IHC) staining for the detection of SARS-CoV-2

S protein in the same area where bronchiolar necrosis was seen. The only bronchiole found to be

positive for Omicron is shown. No evidence of necrosis was seen for this bronchiole. (Scale bar

= 100 μm).

Extended Data Fig. 4: Individual neutralization data. Individual neutralization curves for the

data presented in Fig. 4a,b are shown. The data represent the mean ± SD of three technical

replicates. The curves were calculated based on a non-linear regression curve fit analysis in

Prism. The dotted lines represent the limit of detection.

Extended Data Table 1: Overview of serum samples used for the analysis of antibody

neutralization of WA1, Omi-S, and Omicron. *Days after the second vaccine shot. **The spike

antibody titer was measured by Abbott's SARS-CoV-2 immunoassays.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Extended Data Fig. 1

Extended Data Fig. 2

Survived ■
Dead ●

a

b

Extended Data Fig. 3

Extended Data Fig. 4

Extended Data Table 1

Serum no.	Sex	Race	Age	Days post-vaccination*	Vaccine (Manufacturer)	Spike Ab titer (AU/ml)**
1	Male	White	59	18	mRNA-1273 (Moderna)	39823.0
2	Male	Black	26	37	mRNA-1273 (Moderna)	26978.7
3	Male	Asian	55	34	mRNA-1273 (Moderna)	24880.7
4	Male	White	39	32	mRNA-1273 (Moderna)	23816.7
5	Male	Asian	45	38	mRNA-1273 (Moderna)	21659.5
6	Male	White	30	32	mRNA-1273 (Moderna)	18986.5
7	Female	Asian	47	35	mRNA-1273 (Moderna)	100000.0
8	Female	White	62	47	mRNA-1273 (Moderna)	69680.0
9	Female	White	39	14	mRNA-1273 (Moderna)	54996.6
10	Female	White	38	32	mRNA-1273 (Moderna)	46494.7
11	Female	White	34	30	mRNA-1273 (Moderna)	43784.0
12	Female	White	57	42	mRNA-1273 (Moderna)	42140.5
13	Male	Mixed	28	51	BNT162b2 (Pfizer-BioNTech)	17623.8
14	Male	White	30	54	BNT162b2 (Pfizer-BioNTech)	16154.5
15	Male	White	29	54	BNT162b2 (Pfizer-BioNTech)	14261.5
16	Male	Asian	48	48	BNT162b2 (Pfizer-BioNTech)	10593.6
17	Male	White	46	60	BNT162b2 (Pfizer-BioNTech)	9752.3
18	Male	White	31	53	BNT162b2 (Pfizer-BioNTech)	8715.2
19	Female	White	55	52	BNT162b2 (Pfizer-BioNTech)	100000.0
20	Female	White	43	47	BNT162b2 (Pfizer-BioNTech)	44385.4
21	Female	White	56	48	BNT162b2 (Pfizer-BioNTech)	39998.5
22	Female	Mixed	44	49	BNT162b2 (Pfizer-BioNTech)	31141.9
23	Female	White	56	50	BNT162b2 (Pfizer-BioNTech)	25969.6
24	Female	White	55	51	BNT162b2 (Pfizer-BioNTech)	23539.1