# Obscurin Rho GEF domains are phosphorylated by MST-family kinases but do not exhibit nucleotide exchange factor activity towards Rho GTPases *in vitro*

Daniel Koch[1,2,#], Ay Lin Kho[1], Atsushi Fukuzawa[1], Alexander Alexandrovich[1], Andrew Beavil[1], Mark Pfuhl[1], Martin Rees[1,#], Mathias Gautel[1,#]

[1]Randall Centre for Cell and Molecular Biophysics, King's College London, United Kingdom. [2]Current address: Max Planck Institute For Neurobiology of Behavior - caesar, Bonn, Germany. [#] correspondence to: DK (dkoch.research@protonmail.com), MR (martin.rees@kcl.ac.uk) and MG (mathias.gautel@kcl.ac.uk)

## Abstract:

Obscurin is a giant muscle protein (>800 kDa) featuring multiple signalling domains, including an SH3-DH-PH domain triplet from the Trio-subfamily of guanosine nucleotide exchange factors (GEFs). While previous research suggests that these domains can activate the small GTPases RhoA and RhoQ in cells, *in vitro* characterization of these interactions using biophysical techniques has been hampered by the intrinsic instability of obscurin GEF domains. To study substrate specificity, mechanism and regulation of obscurin GEF function by individual domains, we successfully optimized recombinant production of obscurin GEF domains and found that MST-family kinases phosphorylate the obscurin DH domain at Thr5798. Despite extensive testing of multiple GEF domain fragments, we did not detect any nucleotide exchange activity *in vitro* against 9 representative small GTPases. Bioinformatic analyses show that obscurin differs from other Trio-subfamily GEFs in several important aspects. While further research is necessary to evaluate obscurin GEF activity *in vivo*, our results indicate that obscurin has atypical GEF domains that, if catalytically active at all, are subject to complex regulation.

**Keywords**: sarcomere, small GTPases, guanosine nucleotide exchange factors, enzyme kinetics, pseudo-enzymes

## 1. Introduction

Obscurin is a giant muscle protein of yet poorly understood functions [1]. Its N-terminus links the proteins titin and myomesin at the sarcomeric M-band and the C-terminus of obscurin isoform A binds to small ankyrins at the sarcoplasmic reticulum membrane and at subsarcolemmal protein complexes, thereby creating multiple physical links that contribute to M-band stability and membrane integrity [2–8]. In addition to these structural roles, obscurin is likely involved in multiple signalling contexts as it features two kinase domains, an IQ-motif, two calmodulin binding regions and an SH3-DH-PH RhoGEF domain triplet [1,9]. Small GTPases of the Rho-family are a class of important signalling proteins that cycle between a biologically inactive GDP-bound form and a functional GTP-bound form and regulate diverse cellular functions including gene expression, cytoskeletal remodelling and contractility [10]. Similar to phosphorylation being catalyzed by a kinase, transition to the active GTP state is promoted by guanosine nucleotide exchange factors (GEFs) [11]. Previous studies showed that the obscurin DH-PH domains may act as a GEF on GTPases RhoA and RhoQ/TC10 but not Rac1 or Cdc42 and that these interactions are important for myofibril organization and may alter gene expression in response to mechanical signals [12,13]. However, these insights were obtained using co-IP techniques involving nucleotide-state modifying Rho mutants and effector-domain pulldown assays from cell homogenates. While these approaches are suitable for a first characterization, they offer little information on substrate specificity and kinetics and can be prone to unintended consequences resulting from GTPase-mutants, overexpression of the GEF and unidentified cellular factors [14,15]. Characterising the regulation and mechanism of GEF activity itself is subject to similar experimental constraints. Moreover, obscurin has

only been tested against four GTPases although the Rho-family features more than 20
members, many of which could be potential substrates of the obscurin RhoGEF domains [10,16].

In contrast, biochemical, biophysical and structural *in vitro* methods allow for quantitative and
precisely controlled characterization of kinetics, mechanism and regulation of GTPase:GEF
interactions. Here, we report the production of soluble, stable recombinant obscurin GEF
domains necessary for such studies. We identified kinases and phosphatases that can
regulate the phosphorylation status of the obscurin RhoGEF region at a physiological site and
tested over 10 different obscurin RhoGEF fragments against RhoA, RhoB, RhoC, Rac1, Rac2,
Rac3, RhoG, Cdc42 and RhoQ/TC10 to determine kinetics, substrate specificity and
regulation of the obscurin DH domain by adjacent domains and phosphorylation. Surprisingly,
we found that none of the obscurin RhoGEF fragments exhibited any GEF activity.
Bioinformatic analyses suggest that obscurin is an atypical member of the trio-subfamily of
RhoGEFs to which it belongs and differs from other subfamily members at several important
and conserved residues implicated in interactions with substrate GTPases, potentially
explaining the lack of activity *in vitro*. We discuss implications for our understanding of
obscurin RhoGEF function and outline avenues for future research.

## 2. Results

*Construct design and purification of human obscurin RhoGEF domains*

Initial attempts to purify recombinant, His$_6$-tagged obscurin RhoGEF domains (SH3-DH-PH,
DH-PH, DH and PH) following expression in *E. coli* were not successful in our hands. Domain
boundaries of these constructs were based on exon boundaries. We were able, however, to
successfully express and purify obscurin SH3 and SH3-DH at high purity and at a yield of >5-
10 mg/L culture (**Figure S1**). Since these protein fragments were stable at high concentrations
without precipitation, we reasoned that the insolubility of the catalytic DH domain may be a
result of suboptimal domain boundaries. We thus performed a bioinformatic analysis of the
amino acid sequence of obscurin RhoGEF domains using multiple sequence and structure
prediction algorithms and found that strict adherence to exon boundaries may disrupt the first
N-terminal alpha-helix of the obscurin DH domain and excludes several highly conserved
residues (**Figure 1A**). Using this information, we designed and screened 39 additional
constructs of the DH, PH and DH-PH domains and found that DH domain constructs that fully
include the predicted N-terminal extension resulted in soluble protein fragments at the correct
molecular weight in small-scale purification assays (**Figure S2-3**). DH-PH or PH domain
constructs, however, did not result in soluble protein, despite different domain boundaries,
indicating that the PH domain is intrinsically unstable and difficult to purify. Larger scale
purification of two selected DH domain fragments resulted in milligrams of highly pure, stable
and folded protein (**Figure 1B-C and S4**).

*Human obscurin RhoGEF domains do not exhibit catalytic activity towards Rho GTPases in*
*vitro*

The availability of highly pure and folded protein domains prompted us to characterize the
catalytic activity of the obscurin RhoGEF domains towards Rho GTPases using enzyme
kinetic methods. Using GTPases prepared with the fluorescent nucleotide 2'/3'-O-(N-Methyl-
anthraniloyl)-GDP as a substrate, we tested whether obscurin SH3-DH and DH fragments can
facilitate nucleotide exchange on any of the typical Rho GTPases that cycle between
GDP/GTP states and are thus regulated by GEFs (with the exception of RhoJ, which was not

**Figure 1**. **A**, obscurin RhoGEF region. Exon boundaries and bioinformatic analysis and structure prediction based on amino acid sequence of human obscurin B isoform (NCBI ref. seq. NM_001098623). **B,** gel-filtration elution profile of DH domain fragment comprising residues 5681-5889. **C**, SDS-PAGE analysis of samples at different steps of the purification process of the DH$_{5681-5889}$ fragment. **D**, 1D-NMR analysis of purified DH$_{5681-5889}$ fragment. NMR spectrum exhibits wide peak dispersal and peaks below 0 ppm (black arrow), indicating that the protein is folded.

soluble in our hands). Surprisingly, we found that the nucleotide exchange rate after addition of obscurin RhoGEF fragments was indistinguishable from the intrinsic nucleotide exchange rate – in stark contrast to the positive controls (**Figure 2** and **S5**). In other words, neither of the obscurin RhoGEF fragments exhibited nucleotide exchange activity towards any of the tested Rho GTPases, including RhoA and RhoQ/TC10.

To confirm these results, we performed different pull-down experiments using a previously described protocol to stabilize the nucleotide-free high-affinity GEF-GTPase complex to specifically enrich GEFs [17]. Using the putative substrate RhoA as a bait, we found that regardless of whether RhoA is bound to nucleotides, the bare DH domain does not bind to RhoA, suggesting RhoA is neither a substrate, nor binding partner of obscurin's DH domain *in vitro* (**Figure S6**). Our findings thus disagree with previous reports of obscurin RhoGEF domains activating RhoA and RhoQ/TC10 [12,13].

To understand the reason for the discrepancy posed by our results, we further analyzed the sequence and predicted structure of obscurin RhoGEF domains, allowing us to better interpret our results in the context of the currently available mechanistic paradigms for Rho GEFs. The general mechanisms of DH-domain GEFs are well characterized and reviewed in the literature [11,18–20]. Due to the amino acid sequence of its RhoGEF domains, obscurin belongs to the Trio-subfamily of RhoGEFs, which includes the GEFs dbl and dbs, for which detailed structural and kinetic data are available [21]. Interestingly, the DH domains of dbl, dbs and trio have a much lower nucleotide exchange factor activity in the absence of their corresponding PH domain [22].

119
**Figure 2**. Guanosine nucleotide exchange factor activity of obscurin RhoGEF fragments towards RhoA, Cdc42,
Rac1 and RhoQ/TC10. Left panels show the intrinsic nucleotide activity as a negative control, EDTA and GEFs
Vav2 and Larg as positive controls for each GTPase. Right panels show the nucleotide exchange activity after
addition of obscurin. Black arrows indicate addition of buffer/GEF/EDTA. Data represent mean of n = 2-3
experiments.

Subsequent work demonstrated that residues at the DH-PH domain interface (including
residues from the PH domain) make contact with the switch II region of the substrate GTPase
and contribute to the catalytic activity of the GEF (**Figure 3A, B**) [19,20]. These residues are also
present in most other trio-subfamily GEFs and are conserved in the obscurin sequence in
humans, mice, chicken and zebrafish, but not in drosophila or nematodes (**Figure S7**) [19].

This suggests the catalytic contribution of PH-domain residues is evolutionary conserved both
in the trio-subfamily and across species. We thus hypothesized that these residues may need
to be present for obscurin to exert GEF activity towards Rho GTPases. Since we were unable
to purify any proteins containing the human obscurin PH domain, we cloned and attempted to
purify several chimeric obscurin DH-PH fragments containing the human DH domain followed
by the PH domain from human dbs and trio and the obscurin PH domain from chicken or
zebrafish, all of which feature the aforementioned residues involved in supporting GEF activity.
Of these chimeric proteins, only human/zebrafish chimeras (henceforth called "mermaid"
obscurin) could be successfully purified (**Figure S8**). Many zebrafish RhoGTPases (in
particular the putative substrate RhoA) have >90% sequence identity with human
RhoGTPases [23]. Given the high sequence identity of the substrates across species, we
reasoned that zebrafish obscurin GEF fragments should likely exhibit activity towards human
RhoGTPases.

We therefore cloned and successfully purified additional mermaid as well as several zebrafish
obscurin DH and DH-PH fragments (**Figure S9**). Next, we tested mermaid and zebrafish
RhoGEF fragments for nucleotide exchange activity towards RhoA, Cdc42, Rac1 and
RhoQ/TC10. Again, we found no increase in the nucleotide exchange rate after addition of
these GEF domains (**Figure 3D**). We also tested the nucleotide exchange activity of zebrafish
obscurin DH-PH$_{5884-6217}$ towards RhoB, RhoC, Rac2, Rac3 and RhoG but did not detect any
increase in the nucleotide exchange rate (**Figure S10**).

Our data show, therefore, that the inclusion of the PH domain was not sufficient to obtain
catalytically active obscurin RhoGEF *in vitro*, at least for the tested substrates.

*Obscurin RhoGEF domains can be phosphorylated by MST kinases and CaMKs*

The complete absence of guanosine nucleotide exchange activity of DH-PH fragments is
peculiar. Flanking SH3 and PH domains as well as linker regions are known to sometimes
inhibit the activity of the DH domain [24-28]. Typically, release of such autoinhibitory regions is
mediated by phosphorylation (reviewed e.g. in Hodge and Ridley, 2016; Rossman et al.,
2005). In the case of obscurin, simple autoinhibition of the obscurin DH domain by the SH3 or
PH domain as the sole explanation for the lack of obscurin GEF activity is unlikely given the
absence of activity of the isolated DH domain alone and the conserved positive catalytic
contribution of the PH domain in trio-subfamily RhoGEFs. Interestingly, however, the murine
obscurin RhoGEF region, too, has been reported to become phosphorylated at the residues
corresponding to human Ser5669 (in the SH3-DH interdomain linker) and Thr5798 (within the
DH domain) upon muscle exercise [31]. Ser5669 is conserved among human, mice and
zebrafish, Thr5798 only among human and mice (**Figure S11**). We thus speculated that
phosphorylation might still be important to activate obscurin RhoGEF function, possibly via
conformational changes in one of its domains.

**Figure 3**. **A-C**, Structural determinants of nucleotide exchange activity in Trio-subfamily GEFs. **A**, the aligned structures of Dbs bound to RhoA (PDB-ID: 1LB1) and Cdc42 (PDB-ID: 1KZ7) show a contact between the DH-PH domain interface and the switch II region (red) of the respective substrate GTPase. **B**, a closer view of this contact shows a polar interaction between a switch II arginine and aspartate (red) and three conserved residues of the GEF (grey), two of which belong to the PH domain (Tyr889 and Gln834). **C**, Alignment of the predicted structure of the human obscurin DH-PH to the Dbs RhoA/Cdc42 complex shows that the corresponding obscurin residues are in a similar position as in the Dbs/substrate GTPase complexes. **D**, Guanosine nucleotide exchange factor activity of chimeric and zebrafish obscurin RhoGEF fragments towards RhoA, Cdc42, Rac1 and RhoQ/TC10. Black arrows indicate addition of buffer/GEF/EDTA. Data represent mean of n = 2-3 experiments. See Figure 2 for controls (experiments have been performed simultaneously in a multi-well plate reader).

To identify kinases that can phosphorylate the RhoGEF region, we used the largest human RhoGEF fragment available to us (SH3-DH) as a substrate in a commercial screen comprising a library of 245 Ser/Thr-protein kinases. Of these, 42 kinases exhibited a significant activity towards obscurin SH3-DH. Interestingly, the list of significant hits contained many CaMK-family kinases (CAMK1D, CAMK2B, CAMK2D and CAMK4) and MST-family kinases (MST1, MST2 and MST4), which also exhibited high total phosphorylation levels of obscurin SH3-DH (**Figure S12A**). A follow-up assay using the three kinases with the highest activity ratios (TBK1, CaMK4 and MST2) confirmed the activity (**Figure S12B**). While the relevance of CaMKs to cardiovascular biology is well documented, MST2 has been implicated e.g. in cardiac hypertrophic signalling in response to pressure overload [32].

To independently validate the phosphorylation of obscurin and narrow down the phosphorylation sites, we analysed phosphorylation of different GEF fragments (SH3, SH3-DH and DH) after addition of PKA, CaMK1d, CaMKII or MST2 using a phospho-protein specific staining method. In contrast to the screen, we found that only MST2 can robustly phosphorylate obscurin SH3-DH (**Figure 4A**). Interestingly, CaMKII did not phosphorylate the SH3-DH fragment but only the isolated SH3 domain, suggesting that the phosphorylation site is sterically inaccessible in the SH3-DH fragment, potentially due to an intramolecular interaction between SH3 and DH domain. None of the four tested kinases could phosphorylate the DH domain.

We further tested whether phosphatases PP1 and PP2A, both of which are crucial regulators of myofilament and SR proteins, can dephosphorylate obscurin SH3-DH following phosphorylation by MST2. We found that both PP1 and PP2A can dephosphorylate the SH3-DH fragment (**Figure S12C**).

Next, we tested whether MST2 can phosphorylate the zebrafish DH-PH fragment and a human N- and C-terminally slightly extended DH domain featuring most of the interdomain linkers including the Ser5669 site. We found that MST2 can phosphorylate the extended human DH domain, but not the zebrafish DH-PH fragment (**Figure 4B**). Finally, we directly mapped the MST2 phosphorylation site using mass-spectrometry and confirmed that it is likely Thr5798 (**Figure S13**).

Having identified a kinase that phosphorylates the DH domain at a phosphorylation site reported in physiological context, we tested whether this modification leads to the activation of obscurin GEF activity by repeating the GEF activity assays with obscurin that had been phosphorylated *in vitro* by MST2. We found that phosphorylation of either SH3-DH or the N-terminally extended DH domain did not lead to discernible GEF activity towards RhoA, Cdc42, Rac1 or RhoQ/TC10 (**Figure 4C**).

Given the lack of activity *in vitro*, we wondered whether the obscurin DH-PH tandem could change the localization of endogenous RhoA and RhoQ, which might indicate GEF activity in cells. We thus overexpressed the GFP-tagged human obscurin DH-PH domains in neonatal rat and adult murine cardiomyocytes using adenoviral vectors and studied the localization of RhoA and RhoQ using immunofluorescence microscopy (**Figure S14**). In uninfected neonatal rat cardiomyocytes, RhoQ/TC10 exhibited a striated, RhoA a mostly diffuse localisation pattern (although very rarely also a striated pattern). RhoQ/TC10 and RhoA localisation in the obscurin DH-PH overexpressing cells did not appear any different to the cells expressing GFP alone, both in the presence and absence of the adrenergic agonist phenylephrine. Thus, we found no evidence for activity of obscurin RhoGEF domains in living cells. However, we cannot rule out the possibility that obscurin might activate RhoA or RhoQ/TC10 without changing their localisation, or that under different culture conditions or by using alternative cell types a difference would be revealed, warranting further work to test these possibilities.

**Figure 4**. **A,** phosphorylation of different human obscurin RhoGEF domain combinations by different kinases. **B**, MST2-phosphorylation of an N- and C-terminally extended human DH domain and a zebrafish DH-PH fragment compared to human SH3-DH. **C**, Guanosine nucleotide exchange factor activity of obscurin RhoGEF (10 µM) fragments phosphorylated at Thr5798 towards RhoA, Cdc42, Rac1 and RhoQ/TC10. Black arrows indicate addition of buffer/GEF/EDTA. Data represent mean of n = 2-4 experiments.

## 3. Discussion

Although trio-subfamily GEFs are known for their reliance on the PH-domain for effective nucleotide exchange (Rossman and Campbell, 2000; Snyder et al., 2002), the complete absence of any catalytic activity of the recombinant obscurin GEF domains towards any of the 10 tested GTPases is surprising, given the previously reported activity of obscurin towards RhoA and RhoQ/TC10 using co-IP/pulldown assays [12,13]. The putative activating effect of obscurin on RhoA seems to be conserved for the invertebrate homologues unc89 and Rho-1 [33]. The inclusion of the PH domain or Thr5798 phosphorylation in vertebrate obscurin did not lead to activation of its RhoGEF function either. However, when used at sufficiently high concentrations such as in our study, even inefficient DH domains typically show some degree of discernible activity towards their substrate GTPases [22].

There are multiple reasons which could account for this discrepancy. Firstly, while our experiments tested many combinations of GEF fragments and RhoGTPases, they were, due to practical limitations, not exhaustive. Thus, we might have just missed the right GEF/substrate pair which would lead to discernible GEF activity (e.g. pThr5798-SH3-DH and RhoG). However, in a systematic kinetic study of 21 DH-domain RhoGEFs, all of the tested GEFs exhibited at least some discernible residual GEF activity towards RhoA, Rac1, or Cdc42, even when these were not the main physiological substrates [18]. Since our minimal tested set

of potential substrate GTPases included RhoA, Rac1, Cdc42 and RhoQ/TC10, we consider
this possibility unlikely.

Another possibility is that the use of zebrafish obscurin or human/zebrafish chimeras might
not work due to critical differences in the sequence and mechanism between species. Human
and zebrafish RhoA exhibit a very high (>90%) sequence identity [23] and zebrafish obscurin
PH residues known to be important in the GTPase/PH-domain interaction, too, are conserved
as shown above. The conservation of molecular function is a feature often observed in the
evolution of Ras-family GTPases. Yeast Ypt1, for instance, can be substituted by its mouse
homologue without loss of function [34]. For RhoGTPases in particular, evolutionary
diversification of function rather occurs via diversification of their 'regulatome' [21]. For these
reasons, we think species differences are not the most likely explanation for the lack of
observed activity either.

A more likely explanation, in our view, is that the activation of obscurin requires other,
potentially even multiple factors to obtain an active GEF. For instance, we do not yet know
what effect phosphorylation of Ser5669 has on obscurin function or what the responsible
kinase is for this site. Perhaps activation of obscurin RhoGEF function even requires both DH
and PH domains plus phosphorylation at either or both Ser5669 and Thr5798 or
phosphorylation-dependent recruitment of additional cellular cofactors. We conducted
molecular dynamics simulations using a structural model of the SH3-DH-PH domains that
suggested that phosphorylation at Ser5669, Thr5798, or at both positions does not lead to
large conformational changes, inter-domain motions or significantly altered dynamics for
individual residues (**Figure S15**). Thus, phosphorylation at these sites might change the
availability of bindings sites for interaction partners rather than altering intramolecular
conformation. Furthermore, it is worth pointing out that we used cardiomyocytes for cell
biological experiments, whereas Ford-Speelman and colleagues used *tibialis anterior* muscles
[13], suggesting that such additional factors required for GEF activity might even be tissue
specific. It is interesting to note in this context that the complete knockout of obscurin in mice
leads to a fairly mild muscle phenotype with disrupted organisation of the sarcoplasmic
reticulum and altered cellular calcium-handling[5,35], while human loss-of-function mutations in
OBSCN, abrogating the C-terminal GEF domain, display a compatible phenotype with
susceptibility to severe rhabdomyolysis due to defective calcium-handling[36]. Neither
phenotype offers an obvious mechanistic link to defective small GTPase signalling, other than
potentially in membrane remodelling during SR-formation.

Another possibility to consider is that the obscurin RhoGEF domains might have lost its
catalytic function during evolution. Enzymes which lost their catalytic function during evolution
include proteins from various classes including GTPases, kinases and phosphatases [37,38]. The
N-terminal kinase of the *C. elegans* obscurin homologue unc89, for example, is predicted to
be inactive due to the alteration of key residues involved in catalysis and the absence of the
N-terminal kinase lobe [39]. Such enzymes are often called pseudo-enzymes and obscurin might
thus be the first instance of a "pseudo-GEF". In this case, the observed effect of obscurin on
RhoGTPase activity[12,13] could be indirect. Obscurin might, for example via a non-catalytic
interaction, recruit GTPases to specific membrane compartments where they could be
subsequently activated by another GEF.

A more detailed bioinformatic analysis of the trio-subfamily DH-PH domain sequences
suggests that obscurin does not cluster with other trio-subfamily members and shows the least
similarity to other subfamily members (**Figure S16A**). Furthermore, obscurin exhibits several
differences at multiple highly conserved and functionally important residues for the interaction
with GTPases [18] (**Figure S16B**). These analyses, together with our experimental data,

demonstrate that obscurin is a GEF that, if catalytically active, appears to be regulated in a more complex fashion than other RhoGEFs. Clearly, these results warrant further investigation into the activity of the obscurin DH-PH domains in living cells. A promising approach would be to use FRET-biosensors, which are available for both RhoA and RhoQ/TC10 (as well as many other RhoGTPases) to monitor the activity of RhoA or RhoQ/TC10 in response to obscurin in real-time [40,41].

Despite the lack of observed catalytic activity, the availability of highly pure recombinant protein fragments paves the way for the identification and structural or biophysical characterization of further interaction partners. Moreover, the identification of multiple kinases such as CaMKs and MST-kinases as potential regulators of obscurin RhoGEF function opens new avenues to study the role of these domains in cells.

**4. Material and Methods:**

Details on material and methods, including a list primers used for cloning (**Table S1**), can be found in the online supplementary information.

**Author Contributions:**

DK, MR and MG designed the experiments. DK, ALK and AF performed experiments and analysed the data. AA developed the small-scale purification assay. AB and MR performed the molecular dynamics simulations. MP performed the NMR experiments. MP, MR and MG supervised the study. DK wrote the paper with help from MR and ALK. MG conceived the study and acquired funding. All authors proofread the manuscript.

**Acknowledgements:**

We thank Dr Thomas Kampourakis for access to the plate reader. DK was funded by a PhD studentship from the British Heart Foundation (grant [FS/17/65/33481]). This work was supported by funds from the Wellcome Trust (Collaborative Award in Sciences 201543/Z/16/Z to MG), the European Research Council under the European Union's Horizon 2020 Programme (ERC-2019-SyG, grant no. 856118 to MR and MG) and the Medical Research Council (MR/R003106/1 to MG and ALK). Earlier work leading to the present paper was supported by the Volkswagen Foundation Research grant I/81 797 and I/78 989 to MG. MG holds the BHF Chair of Molecular Cardiology. We acknowledge support by CREATE[42]. The authors declare that they have no conflict of interest.

**References**

1.      Young, P., Ehler, E., and Gautel, M. (2001). Obscurin, a giant sarcomeric Rho guanine nucleotide exchange factor protein involved in sarcomere assembly. J Cell Biol *154*, 123–136. 10.1083/jcb.200102110.

2.      Bagnato, P., Barone, V., Giacomello, E., Rossi, D., and Sorrentino, V. (2003). Binding of an ankyrin-1 isoform to obscurin suggests a molecular link between the sarcoplasmic reticulum and myofibrils in striated muscles. The Journal of Cell Biology *160*, 245–253. 10.1083/jcb.200208109.

3.      Blondelle, J., Marrocco, V., Clark, M., Desmond, P., Myers, S., Nguyen, J., Wright, M., Bremner, S., Pierantozzi, E., Ward, S., et al. (2019). Murine obscurin and Obsl1 have functionally redundant roles in sarcolemmal integrity, sarcoplasmic reticulum organization, and muscle metabolism. Communications Biology *2*, 178. 10.1038/s42003-019-0405-7.

4.      Fukuzawa, A., Lange, S., Holt, M., Vihola, A., Carmignac, V., Ferreiro, A., Udd, B., and Gautel, M. (2008). Interactions with titin and myomesin target obscurin and obscurin-like 1 to the M-band – implications for hereditary myopathies. J. Cell Sci. *121*, 1841. 10.1242/jcs.028019.

5.      Lange, S., Ouyang, K., Meyer, G., Cui, L., Cheng, H., Lieber, R.L., and Chen, J. (2009). Obscurin determines the architecture of the longitudinal sarcoplasmic reticulum. J. Cell Sci. *122*, 2640. 10.1242/jcs.046193.

6.      Pernigo, S., Fukuzawa, A., Pandini, A., Holt, M., Kleinjung, J., Gautel, M., and Steiner, R.A. (2015). The Crystal Structure of the Human Titin:Obscurin Complex Reveals a Conserved yet Specific Muscle M-Band Zipper Module. Journal of Molecular Biology *427*, 718–736. 10.1016/j.jmb.2014.11.019.

7.      Randazzo, D., Giacomello, E., Lorenzini, S., Rossi, D., Pierantozzi, E., Blaauw, B., Reggiani, C., Lange, S., Peter,
A.K., Chen, J., et al. (2013). Obscurin is required for ankyrinB-dependent dystrophin localization and sarcolemma integrity. J
Cell Biol *200*, 523–536. 10.1083/jcb.201205118.

8.      Randazzo, D., Blaauw, B., Paolini, C., Pierantozzi, E., Spinozzi, S., Lange, S., Chen, J., Protasi, F., Reggiani, C., and
Sorrentino, V. (2017). Exercise-induced alterations and loss of sarcomeric M-line organization in the diaphragm muscle of
obscurin knockout mice. American Journal of Physiology - Cell Physiology *312*, C16–C28. 10.1152/ajpcell.00098.2016.

9.      Fleming, J.R., Rani, A., Kraft, J., Zenker, S., Börgeson, E., and Lange, S. (2021). Exploring Obscurin and SPEG
Kinase Biology. Journal of Clinical Medicine *10*, 984. 10.3390/jcm10050984.

10.      Ridley, A.J. (2001). Rho family proteins: coordinating cell responses. Trends in Cell Biology *11*, 471–477.
10.1016/S0962-8924(01)02153-5.

11.      Cherfils, J., and Zeghouf, M. (2013). Regulation of small GTPases by GEFs, GAPs, and GDIs. Physiol Rev *93*, 269–
309. 10.1152/physrev.00003.2012.

12.      Coisy-Quivy, M., Touzet, O., Bourret, A., Hipskind, R.A., Mercier, J., Fort, P., and Philips, A. (2009). TC10 controls
human myofibril organization and is activated by the sarcomeric RhoGEF obscurin. J. Cell Sci. *122*, 947–956.
10.1242/jcs.040121.

13.      Ford-Speelman, D.L., Roche, J.A., Bowman, A.L., and Bloch, R.J. (2009). The rho-guanine nucleotide exchange
factor domain of obscurin activates rhoA signaling in skeletal muscle. Mol Biol Cell *20*, 3905–3917. 10.1091/mbc.E08-10-1029.

14.      Langemeyer, L., Bastos, R.N., Cai, Y., Itzen, A., Reinisch, K.M., and Barr, F.A. (2014). Diversity and plasticity in Rab
GTPase nucleotide release mechanism has consequences for Rab activation and inactivation. eLife *3*, e01623.

15.      Stanley, R.J., and Thomas, G.M.H. (2016). Activation of G Proteins by Guanine Nucleotide Exchange Factors Relies
on GTPase Activity. PLOS ONE *11*, e0151861. 10.1371/journal.pone.0151861.

16.      Ridley, A.J. (2016). Open questions: what about the 'other' Rho GTPases? BMC Biology *14*, 64. 10.1186/s12915-
016-0289-7.

17.      Koch, D., Rai, A., Ali, I., Bleimling, N., Friese, T., Brockmeyer, A., Janning, P., Goud, B., Itzen, A., Müller, M.P., et al.
(2016). A pull-down procedure for the identification of unknown GEFs for small GTPases. Small GTPases *7*, 93–106.
10.1080/21541248.2016.1156803.

18.      Jaiswal, M., Dvorsky, R., and Ahmadian, M.R. (2013). Deciphering the Molecular and Functional Basis of Dbl Family
Proteins: A NOVEL SYSTEMATIC APPROACH TOWARD CLASSIFICATION OF SELECTIVE ACTIVATION OF THE Rho
FAMILY PROTEINS. Journal of Biological Chemistry *288*, 4486–4500. 10.1074/jbc.M112.429746.

19.      Rossman, K.L., Worthylake, D.K., Snyder, J.T., Siderovski, D.P., Campbell, S.L., and Sondek, J. (2002). A
crystallographic view of interactions between Dbs and Cdc42: PH domain-assisted guanine nucleotide exchange. The EMBO
Journal *21*, 1315–1326. 10.1093/emboj/21.6.1315.

20.      Snyder, J.T., Worthylake, D.K., Rossman, K.L., Betts, L., Pruitt, W.M., Siderovski, D.P., Der, C.J., and Sondek, J.
(2002). Structural basis for the selective activation of Rho GTPases by Dbl exchange factors. *9*, 468.

21.      Fort, P., and Blangy, A. (2017). The Evolutionary Landscape of Dbl-Like RhoGEF Families: Adapting Eukaryotic
Cells to Environmental Signals. Genome Biol Evol *9*, 1471–1486. 10.1093/gbe/evx100.

22.      Rossman, K.L., and Campbell, S.L. (2000). Bacterial expressed DH and DH/PH domains. In Methods in Enzymology
Regulators and Effectors of Small GTPases - Part D: Rho Family., W. E. Balch, C. J. Der, and A. Hall, eds. (Academic Press),
pp. 25–38. 10.1016/S0076-6879(00)25428-1.

23.      Salas-Vidal, E., Meijer, A.H., Cheng, X., and Spaink, H.P. (2005). Genomic annotation and expression analysis of the
zebrafish Rho small GTPase family during development and bacterial infection. Genomics *86*, 25–37.
10.1016/j.ygeno.2005.03.010.

24.      Aghazadeh, B., Lowry, W.E., Huang, X.-Y., and Rosen, M.K. (2000). Structural Basis for Relief of Autoinhibition of
the Dbl Homology Domain of Proto-Oncogene Vav by Tyrosine Phosphorylation. Cell *102*, 625–633. 10.1016/S0092-
8674(00)00085-4.

25.      Ahmad, K.F., and Lim, W.A. (2010). The Minimal Autoinhibited Unit of the Guanine Nucleotide Exchange Factor
Intersectin. PLOS ONE *5*, e11291. 10.1371/journal.pone.0011291.

26.      Barreira, M., Fabbiano, S., Couceiro, J.R., Torreira, E., Martínez-Torrecuadrada, J.L., Montoya, G., Llorca, O., and
Bustelo, X.R. (2014). The C-Terminal SH3 Domain Contributes to the Intramolecular Inhibition of Vav Family Proteins. Sci.
Signal. *7*, ra35–ra35. 10.1126/scisignal.2004993.

27.      Kintscher, C., Wuertenberger, S., Eylenstein, R., Uhlendorf, T., and Groemping, Y. (2010). Autoinhibition of GEF
activity in Intersectin 1 is mediated by the short SH3-DH domain linker. Protein Sci. *19*, 2164–2174. 10.1002/pro.500.

28.      Mitin, N., Betts, L., Yohe, M.E., Der, C.J., Sondek, J., and Rossman, K.L. (2007). Release of autoinhibition of ASEF
by APC leads to CDC42 activation and tumor suppression. Nature Structural & Molecular Biology *14*, 814. 10.1038/nsmb1290.

29.      Hodge, R.G., and Ridley, A.J. (2016). Regulating Rho GTPases and their regulators. Nature Reviews Molecular Cell
Biology *17*, 496–510. 10.1038/nrm.2016.67.

30.      Rossman, K.L., Der, C.J., and Sondek, J. (2005). GEF means go: turning on RHO GTPases with guanine nucleotide-
exchange factors. *6*, 167.

31.      Potts, G.K., McNally, R.M., Blanco, R., You, J.-S., Hebert, A.S., Westphall, M.S., Coon, J.J., and Hornberger, T.A.
(2017). A map of the phosphoproteomic alterations that occur after a bout of maximal-intensity contractions. J Physiol *595*,
5209–5226. 10.1113/JP273904.

32.      Zi, M., Maqsood, A., Prehar, S., Mohamed, T.M.A., Abou-Leisa, R., Robertson, A., Cartwright, E.J., Ray, S.G., Oh,
S., Lim, D.-S., et al. (2014). The Mammalian Ste20-like Kinase 2 (Mst2) Modulates Stress-induced Cardiac Hypertrophy. J. Biol.
Chem. *289*, 24275–24288. 10.1074/jbc.M114.562405.

33.      Qadota, H., Blangy, A., Xiong, G., and Benian, G.M. (2008). The DH-PH Region of the Giant Protein UNC-89
Activates RHO-1 GTPase in Caenorhabditis elegans Body Wall Muscle. Journal of Molecular Biology *383*, 747–752.
10.1016/j.jmb.2008.08.083.

34.      Haubruck, H., Prange, R., Vorgias, C., and Gallwitz, D. (1989). The ras-related mouse ypt1 protein can functionally
replace the YPT1 gene product in yeast. EMBO J *8*, 1427–1432. 10.1002/j.1460-2075.1989.tb03524.x.

35.      Pierantozzi, E., Szentesi, P., Paolini, C., Dienes, B., Fodor, J., Oláh, T., Colombini, B., Rassier, D.E., Rubino, E.M.,
Lange, S., et al. (2022). Impaired Intracellular Ca2+ Dynamics, M-Band and Sarcomere Fragility in Skeletal Muscles of
Obscurin KO Mice. International Journal of Molecular Sciences *23*, 1319. 10.3390/ijms23031319.

36.      Cabrera-Serrano, M., Caccavelli, L., Savarese, M., Vihola, A., Jokela, M., Johari, M., Capiod, T., Madrange, M.,
Bugiardini, E., Brady, S., et al. (2021). Bi-allelic loss-of-function OBSCN variants predispose individuals to severe recurrent
rhabdomyolysis. Brain, awab484. 10.1093/brain/awab484.

37.      Reiterer, V., Eyers, P.A., and Farhan, H. (2014). Day of the dead: pseudokinases and pseudophosphatases in
physiology and disease. Trends in Cell Biology *24*, 489–505. 10.1016/j.tcb.2014.03.008.

38.      Stiegler, A.L., and Boggon, T.J. (2020). The pseudoGTPase group of pseudoenzymes. The FEBS Journal *287*,
4232–4245. https://doi.org/10.1111/febs.15554.

39.      Small, T.M., Gernert, K.M., Flaherty, D.B., Mercer, K.B., Borodovsky, M., and Benian, G.M. (2004). Three New
Isoforms of Caenorhabditis elegans UNC-89 Containing MLCK-like Protein Kinase Domains. Journal of Molecular Biology *342*,
91–108. 10.1016/j.jmb.2004.07.006.

40.      Kawase, K., Nakamura, T., Takaya, A., Aoki, K., Namikawa, K., Kiyama, H., Inagaki, S., Takemoto, H., Saltiel, A.R.,
and Matsuda, M. (2006). GTP Hydrolysis by the Rho Family GTPase TC10 Promotes Exocytic Vesicle Fusion. Developmental
Cell *11*, 411–421. 10.1016/j.devcel.2006.07.008.

41.      Yoshizaki, H., Ohba, Y., Kurokawa, K., Itoh, R.E., Nakamura, T., Mochizuki, N., Nagashima, K., and Matsuda, M.
(2003). Activity of Rho-family GTPases during cell division as visualized with FRET-based probes. Journal of Cell Biology *162*,
223–232. 10.1083/jcb.200212049.

42.      King's College London. (2022). King's Computational Research, Engineering and Technology Environment
(CREATE). Retrieved March 2, 2022, from https://doi.org/10.18742/rnvf-m076