REDUCED FOLATE CARRIER 1 IS PRESENT IN RETINAL MICROVESSELS AND CRUCIAL FOR THE INNER BLOOD RETINAL BARRIER INTEGRITY

Gokce Gurler[1], Nevin Belder[2], Mustafa Caglar Beker[3], Melike Sever-Bahcekapili[1], Gokhan Uruk[1], Ertugrul Kilic[3,5], Muge Yemisci[1,4,5]

1. The Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara, Turkey

2. Biotechnology Institute, Ankara University, Ankara, Turkey

3. Istanbul Medipol University, Physiology, Istanbul, Turkey

4. Faculty of Medicine, Department of Neurology, Hacettepe University, Ankara, Turkey

5. Neuroscience and Neurotechnology Center of Excellence (NÖROM) Ankara, Turkey

Corresponding Author:

Muge Yemisci (MD, PhD), Professor of Neurology

Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara, Turkey

Faculty of Medicine, Department of Neurology, Hacettepe University, Ankara, Turkey

Neuroscience and Neurotechnology Center of Excellence (NÖROM) Ankara, Turkey

myemisciozkan@gmail.com

ABSTRACT

Background: Reduced folate carrier 1 (RFC1; SLC19a1) is the main responsible transporter for the B9 family of vitamins named folates, which are essential for normal tissue growth and development. While folate deficiency resulted in retinal vasculopathy, the expression and the role of RFC1 in blood-retinal barrier (BRB) are not well known.

Methods: We used whole mount retinas and trypsin digested microvessel samples of adult mice. To knockdown RFC1, we delivered RFC1-targeted short interfering RNA (RFC1-siRNA) intravitreally; while, to upregulate RFC1 we delivered lentiviral vector overexpressing RFC1. Retinal ischemia was induced 1-hour by applying $FeCl_3$ to central retinal artery. We used RT-qPCR and Western blotting to determine RFC1. Endothelium (CD31), pericytes (PDGFR-beta, CD13, NG2), tight-junctions (Occludin, Claudin-5 and ZO-1), main basal membrane protein (Collagen-4), endogenous IgG and RFC1 were determined immunohistochemically.

Results: Our analyses on whole mount retinas and trypsin digested microvessel samples of adult mice revealed the presence of RFC1 in the inner BRB and colocalization with endothelial cells and pericytes. Knocking down RFC1 expression via siRNA delivery resulted in the disintegration of tight junction proteins and collagen-4 in twenty-four hours, which was accompanied by significant endogenous IgG extravasation. This indicated the impairment of BRB integrity after an abrupt RFC1 decrease. Furthermore, lentiviral vector-mediated RFC1 overexpression resulted in increased tight junction proteins and collagen-4, confirming the structural role of RFC1 in the inner BRB. Acute retinal ischemia decreased collagen-4 and occludin levels and led to an increase in RFC1. Besides, the pre-ischemic overexpression of RFC1 partially rescued collagen-4 and occludin levels which would be decreased after ischemia.

Conclusion: In conclusion, our study clarifies the presence of RFC1 protein in the inner BRB, which has recently been defined as hypoxia–immune-related gene in other tissues and offers a novel perspective of retinal RFC1. Hence, other than being a folate carrier, RFC1 is an acute regulator of the inner BRB in healthy and ischemic retinas.

Keywords: RFC1, blood-retina barrier, retinal ischemia, siRNA, lentiviral vector, trypsin digestion, pericyte

BACKGROUND

The Reduced Folate Carrier 1 (RFC1), also called solute carrier family 19 member 1 (SLC19A1/SLC19a1) is responsible for transporting folates via Blood Brain Barrier (BBB) to the brain parenchyma (1, 2). Folates belong to the B9 family of vitamins and are essential for proper functioning of the Central Nervous System (CNS) as they are crucial for many key processes such as histone and DNA methylation, DNA replication and repair, RNA synthesis, methionine production, homocysteine remethylation, aminoacid metabolism, neurotransmitter, and phospholipid synthesis (3-6). Several clinical studies in adults determined that lower serum folate levels could cause neurological dysfunction (7, 8). Moreover, folate deficiency was shown to result in microvascular complications in the retina (9, 10). However, the findings of these studies are chronic consequences that occur with long-term changes and/or deficiency of folates.

Retina is the extension of the CNS, displaying similarities in terms of anatomy, function, and response to insults (11). The retina has the blood-retina barrier (BRB) that precludes the free transfer of substances from the capillary blood, and is composed of two parts (12). The outer BRB is constituted of the retinal pigmented epithelium (RPE), which are specialized epithelial cells sealed with tight junctions, residing between the neural retina and a fenestrated capillary network called choriocapillaris. It partly functions to determine the transfer of nutrients from the choriocapillaris to the retina (13). The inner BRB is established in the inner retinal microvessels and comprises specialized endothelial cells secured by tight junctions, enclosed by basal lamina, pericytes, and perivascular astrocytes; hence it is essential for the maintenance of homeostasis through its selective properties such as harboring specialized nutrient carriers (12). To date the expression of RFC1 and its transporter function in the retina was primarily evaluated only in the RPE, thus the outer BRB (14-17).

The recent cerebral microvessel transcriptome data showed prominent RFC1 presence in pericytes, the cells crucial with various roles in the regulation of microvascular blood flow and barrier integrity. The

retinal microvessels have similar features to the brain (18-20) and are unique for having the highest

pericyte ratio in the body. Therefore, our first aim was to investigate the presence of RFC1 protein in

the inner BRB, and especially in the pericytes and endothelial cells.

RFC1 has been recently shown in the cell lines to be an importer of immunotransmitter cGAMP and

other cyclic dinucleotides with robust antitumoral immune response (21) and considered to be the

hypoxia–immune-related gene in multiple myeloma patients (22). The expression and activity of RFC1

was shown to differ during insults like hyperglycemia, nitric oxide exposure, hyperhomocysteinemia

and oxidative stress in the RPE (14, 15, 17). However, no *in vivo* studies were carried out in the inner

BRB. Therefore, our second aim was to investigate the role of RFC1 protein in the inner BRB in healthy

retinas and in acute retinal ischemia which is a world-wide problem for blindness.

In this study, we elucidated the presence of RFC1 protein in the endothelial cells and pericytes of the

inner BRB. We employed a custom-designed RFC1-targeted Accell short interfering RNA (siRNA) to

silence the RFC1 gene, and a lentiviral vector to overexpress RFC1 in the retinas *in vivo*. Abrupt and

brief changes in RFC1 caused a disruption in the barrier properties of the retinal microvessels. Thus, we

identified the immediate role of RFC1 protein in maintaining the inner BRB in health and in acute retinal

ischemia.

METHODS

Animals

Experiments were performed according to the approval provided by Hacettepe University Animal

Experimentations Local Ethics Board (No: 2019/13-06 and 2021/03-19). Adult female and male Swiss

albino mice (25–35 g) housed under 12 hours of light and 12 hours of darkness at standard room

temperature (22±2°C) and 50-60% humidity was used for *in vivo* experiments. Mice were allowed water

and food ad libitum. The recommendations of ARRIVE guidelines and institutional instructions were

considered in housing, care, and while performing experiments, the experimental groups were decided

in a randomized and blinded manner. For all the surgical procedures intraocular injections, the animals

were anesthetized with ketamine (80 mg/kg, intraperitoneal injection, Ketalar®, Pfizer, Cambridge,

United Kingdom) and xylazine (8 mg/kg, intraperitoneal injection, Alfazyne® 2%, Alfasan

International, Woerden, the Netherlands). A rectal probe connected to the homeothermic blanket with a

control unit (Harvard Apparatus, U.S.A.) was utilized to control the body temperature (37.0 ± 0.5°C).

Pulse rate and oxygen saturation were monitored from the right lower limb via an oximeter (The

LifeSense® VET Pulse Oximeter, Nonin Medical Inc., USA). Vital signs (rectal temperature, blood

pressure, oxygen saturation, and pulse rate) were kept within physiological limits and recorded. After

operations, mice were kept on a homoeothermic blanket until they recovered from anesthesia fully. Next,

the mice were kept individually in their cages until sacrificed.

Retinal Whole Mount Preparation

The eyeballs harvested were immersed in 4% paraformaldehyde (PFA) for 1 hr, and under a

stereomicroscope, we made a puncture at the corneal border with an insulin needle while eyeballs were

resting in one drop of PBS at room temperature. A circular cut along the limbus was made with the help

of a micro-dissector. Following the extraction of the lens and vitreous body, the intact retina was

detached from the sclera via a 45° thick curved tip tweezer by going around the edges. Then, the

flattened retina was placed into a centrifuge tube consisting of 200 μL PBS for immunohistochemistry.

Retinal Coronal Section Preparations

121   Obtained eyeballs were immersed in 4% paraformaldehyde (PFA) for 24 hr followed by incubation in

122   30% sucrose for 2 days. Next, 20 μm sections were obtained by cryostat.

Retinal Trypsin Digest Preparation

We used modified retinal trypsin digestion protocol to isolate retinal microvessels from the surrounding

tissue (23) . Retinas were separated in cold Phosphate Buffered Saline (PBS), then immersed in 250 μL

filtered ddH$_2$0, and, at room temperature, kept on a shaker until they disintegrated. Then kept in 0.3%

trypsin (PAN Biotech 1:250) 0.1 M Tris buffer (pH 7.8) in the incubator adjusted to 37°C. The separation

of microvessels were seen clearly, tissue was washed with dH$_2$O, and the procedure was repeated until

no debris remained. Isolated retinal microvessels were transferred to poly-L-lysine coated slides without

disrupting their integrity, and air-dried. This method allowed isolation of the retinal vessels containing

endothelial cells, basal lamina, pericytes, the attached perivascular astrocyte processes, without the

presence of other cell types like photoreceptor cells.

Immunohistochemical Studies

For immunohistochemical studies whole mount retinas, coronal sections and isolated retinal

microvessels were used. PFA fixed whole mount retinas were kept in in 0.5% PBS-Triton-X, freezed 15

min at -80°C and thawed for 15 min at room temperature following overnight incubation in 2% TritonX-

100 (Merck Millipore, 1086031000) PBS for permeabilization. They were immersed in 10% normal

goat serum for blocking steps for 2 hr at room temperature or overnight at +4°C. Next, whole mount

retinas were incubated at 4°C with primary antibodies against RFC1 (SLC19a1, MyBioSource

MBS9134642 or Sigma-Aldrich AV44167) for three consecutive overnights. We used two commercial

polyclonal antibodies developed against different non-overlapping epitopes of RFC1. RFC1/SLC19a1

antibody from MyBioSource (MBS9134642) was produced against a recombinant fusion protein

containing a sequence related to amino acids 452-591 of human SLC19A1 (NP_919231.1), while

RFC1/SLC19a1antibody from Sigma-Aldrich (AV44167) was produced against Synthetic peptide

directed towards the N terminal region of human SLC19A1 (NP_919231). Then retinas were

subsequently incubated with 'Fluorescein' or 'Texas Red' labeled Lectin (Vector Laboratories,

Burlingame, CA). Retinal microvessels obtained via retinal trypsin digestion were permeabilized with

0.3% TBS-Triton-X for 30 min, blocked in 10% normal serum of the host of the associated secondary

antibody for 1 hr at room temperature. Vessel webs were incubated at +4°C overnight with primary

antibodies against RFC1; for mural cells against PDGFR-β (R&D Systems, AF1042), NG2 (Sigma

AB5320A4), CD13 (Acris Antibodies, AM26636AF-N), for endothelial CD31 (BD Bioscience,

550274); for tight junctions ZO-1 (Sigma, MAB1520), Claudin-5 (Invitrogen™, 35-2500), and for

Collagen-4 (Abcam, ab6586) (n=3-6/marker). After several washes, the vessel webs were incubated for

1 hr at room temperature with appropriate Cy2-, Cy3- or Cy5-conjugated anti-rabbit IgG (Jackson

ImmunoResearch 111-165-144 or 711-175-152) or Cy3-conjugated anti-rat IgG (Thermo Scientific™,

A10522, 1:200). To assess the functional inner BRB integrity, PFA fixed whole mount retinas were

incubated with Cy3-conjugated anti-mouse IgG (Jackson ImmunoResearch, 115-165-003) at +4°C

overnight. Later, retinas were incubated with 'Fluorescein' or 'Texas Red' labeled Lectin (Vector

Laboratories, Burlingame, CA) at +4°C overnight to visualize the vessels. Washed with TBS for 3 times,

5 min each; retinal tissues or retinal microvessels were mounted with a PBS/Glycerol mounting medium

consisting of the nuclear staining Hoechst 33258 (Invitrogen™, 1:1000 dilution).

Western Blotting

Whole retinas were lysed in Radioimmunoprecipitation assay (RIPA) buffer, sonicated, and

homogenized on ice, then centrifuged at 10,000xg at +4°C for 20 min. The concentration of the lysates

was assessed using BCA protein assay kit (Thermo Scientific™, 23225). After loading proteins (35

μg/well) to a homemade SDS-PAGE gel, 120 Voltage were applied for 1.5 hr. Proteins were then

transferred to PVDF membranes by applying 120 mAmp/membrane for 2 hr at room temperature.

Membranes were incubated with 5% BSA in 0.1% TBS-Tween for 1 hr at room temperature for blocking

step and incubated with 1:1000 anti-RFC1 antibody produced in rabbit (Sigma-Aldrich; AV44167) at

+4°C overnight. After washing with 0.1% TBS-Tween, we incubated the membranes with 1:5000 goat

anti-rabbit HRP conjugated Invitrogen™, 31460) solution for 2 hr at room temperature. For detection,

the membranes were treated with a high sensitivity enhanced chemiluminescent (ECL) substrate

(Thermo Scientific™, 34096). Imaging was done by Kodak 4000M Image Station. β-Actin and β-tubulin

III (Chemicon, MAB1637) was employed as an internal standard for cell culture homogenates and tissue

homogenates respectively. Densitometric measurements were done by ImageJ 1.52 version (NIH,

Bethesda, Maryland), and expressed as ratios to the internal standard.

Imaging and Analysis

The images of the stained whole mount retinas and retinal microvessels were obtained with a Leica TCS

SP8 confocal laser scanning microscope (Leica, Wetzlar, Germany) with a diode laser 405, 638 and

OPSL 488, 552 nm, with a X, Y, and Z-movement controller, and a high-resolution PMT (Zeiss,

Oberkochen, Germany) and HyD (Leica) detectors. Images were either acquired in a single focal plane

while keeping settings constant between experimental or control groups for analysis or Z-stack mode

with 0.80 μm wide steps along the Z axis. For analysis of immunostained retinal microvessels,

1024x1024 pixels and 276.79x276.79 μm sized, middle of the Z axis level images was used. Acquired

images were exported as 8-bit grayscale .tiff formats from Leica Application Suite X (LAS X; version

3.5.5.19976) and opened in ImageJ 1.52 version. First, the lectin frame of each image was filtered by

Gaussian blur (sigma=4) to reduce the random noise, and Huang Thresholding was applied to mask the

lectin-positive microvessel areas of the images. After de-speckling (Process tab>Noise>Despeckle) and

eroding (Process tab>Binary>Erode), the binary image was divided by 255 using Math function (Process

tab>Math>Divide) to convert microvessel pixels to "1" and background pixels to "0". Later, either

"RAW" lectin image or the overlapping grayscale image belonging to tight junctions or collagen-4 was

divided by the binary of the lectin image. This created 32-bit grayscale images in which microvessels

had singular intensity and the background had infinity. Finally, the mean grey value (RFU, range

between 0-255) was obtained which corresponded to the mean intensity of the pixels occupying the

microvessel area. Number of pericytes were quantified from Lectin and Hoechst labelled microvessel

preparations by researchers blinded to the experimental or control groups. As pericytes are well known

to have protruding nuclei, they were easy to distinguish morphologically and spatially from endothelial

cells having flat nuclei. The total microvessel length was measured with AngioTool (v 0.6a (64 bits),

October 2014) software, and pericyte number per millimeter (mm) of microvessel length was

calculated (24). IgG extravasation was calculated from the 150x150 μm sized ROIs of 8-bit grayscale

.tiff format images of wholemount retinas labeled with Cy3-conjugated anti-mouse IgG and lectin. The

IgG and lectin frames were applied Huang Thresholding to detect the IgG and lectin-positive pixel

counts. After thresholding, the binary images were divided by 255 to convert IgG positive and lectin

positive pixels to "1" and background pixels to "0". Next, the IgG positive pixel counts are divided into

lectin positive pixel counts.

Gene Silencing *in vivo* by Short Interfering RNA (siRNA)

To silence RFC1 gene, we used two custom-designed mouse RFC1 (Slc19a1) Accell siRNAs targeted

to two different regions of RFC1 mRNA and two Accell scrambled (control) siRNAs (Horizon

Discovery, Waterbeach, United Kingdom). Accell siRNA has several advantages as it does not need any

transfection reagents which may have unwanted effects when applied *in vivo*. Furthermore, they could

be internalized by any type of mammalian cells (25). Accell siRNAs have been previously proven to be uniformly distributed across rat retinal tissue and successfully inhibit various genes following intravitreal delivery (26). Product codes of siRNAs with sequences are given in Table 1. RFC1-siRNA-1 and RFC1-siRNA-2 or scrambled (control) siRNA-1 and 2 were pooled in equal volumes for administration to improve the potency and specificity.

We delivered 100 µM pooled RFC1-siRNAs or Control-siRNAs intravitreally to the eyes (2.3 µL/ eye) of animals via 33 G Neuros Syringe (1701 RN, Hamilton). We did not utilize contralateral eyes as controls to avoid the possible transfer of siRNAs to the opposite side (27, 28). After 24 hr, we either proceeded to retinal ischemia or sacrificed mice, and determined the levels of mRNA or protein.

	Product codes	Sequences of sense strand
RFC1-siRNA-1	A-044248-13	C.C.A.G.C.C.U.A.C.U.U.C.A.U.G.C.U.U.U.U.U
RFC1-siRNA-2	A-044248-15	C.C.A.G.G.A.A.A.C.U.A.G.A.U.C.G.C.A.U.U.U
Scrambled siRNA-1	D-001910-01	U.G.G.U.U.U.A.C.A.U.G.U.C.G.A.C.U.A.A.U.U
Scrambled siRNA-2	D-001910-03	U.G.G.U.U.U.A.C.A.U.G.U.U.U.U.C.C.U.A.U.U

Table 1. The product codes and the sequences of sense strands of RFC1-siRNAs and scrambled (control) siRNAs.

Gene Overexpression *in vivo* by Lentiviral Vector (LV)

Lentiviral vectors (LV) that express either green fluorescent protein (GFP) (Control-LV) or GFP with RFC1 (RFC1-LV) driven by EF-1 alpha promoter were used to overexpress RFC1 gene in the retina.

The lentiviral packaging system used for RFC1-LV or control-LV was second-generation and was an effective and safe technique in production. Total RNA from cultured mouse neuro 2a cells was extracted using AllPrep DNA/RNA/Protein Mini Kit (80004, Qiagen, Hilden, Germany) following the manufacturer's instructions. Transcriptor First Strand cDNA Synthesis Kit (04896866001, Roche, Basel, Switzerland) was used to obtain complementary DNA (cDNA). The coding region of Mus musculus SLC19a1 variant 1/RFC1 (NCBI Reference Sequence: NM_031196.3) was amplified with specific primers (forward 5'-agtcagaattcatggtgcccactggccag-3'and reverse 5'-AGTCAGGATCCTCAAGCCTTGGCTTCGACTCTT -3') with fast digest restriction enzymes EcoRl (FD0274, Thermo Fisher Scientific, Massachusetts, USA) and BamHI (FD0054, Thermo Fisher

Scientific). Restriction enzymes BamHI and EcoRI were used to digest the PCR product of SLC19a1

and expression plasmids (pLenti-EF1α-GFP-2A-Puro; Applied Biological Materials, Richmond,

Canada). Subsequently, ligation was realized with T4 DNA ligase (EL0014, Thermo Fisher Scientific).

Sequencing pMD2.G and psPAX plasmids, kindly provided by Dr. Didier Trono (Ecole Polytechnique

Federale, Lausanne, Switzerland), were used to confirm the insert. They were employed as

complementary vectors of the packaging of the lentiviral system (12259; 12260, Addgene, United

Kingdom). HEK293T cell line (6×10^6 cells) was chosen to be seeded on 10 cm plates (CLS3294,

Corning, New York, USA). The following day, the vector transfection Lipofectamine 3000 (L3000015,

Thermo Fisher Scientific) was applied according to the manufacturer's instructions. Shortly, 7 μg

lentiviral vector, 3.5 μg pMD2.G, and 7 μg psPAX were used to prepare DNA-lipid complexes. The

DNA-lipid complex was added slowly after 10 min of incubation at room temperature. Six hours after

transfection, the medium was changed with fresh DMEM (P04–01158, Pan Biotech, Bavaria Germany)

and was incubated at 37°C in a humid atmosphere consisting of 5% CO_2. Twenty-four and fifty-two

hours following the transfection, the medium was harvested, centrifuged (10 min at 2000 rpm), and

filtered with a low binding filter with a 0.45 μm pore size. After ultra-centrifuging (120,000 g for 2 hr),

viral particles were dissolved in Dulbecco's Phosphate Buffered Saline (DPBS) without calcium and

magnesium (P04–3650, Pan Biotech). The plasmid with no DNA inserted was packaged with the same

procedures to use as a control. The viral titer (1×10^8 lentivirus particles in 1 μl 0.1 M PBS) was

measured by a previously published protocol (29).

LV (1×10^8 lentivirus particles in 1 μl 0.1 M PBS) were intravitreally delivered to the retina (2 μL/eye)

via 33 G Neuros Syringe (1701 RN, Hamilton). Ten days after the delivery, we either proceeded to

retinal ischemia or sacrificed mice to obtain eyeballs.

Quantitative Reverse Transcriptase-Polymerase Chain Reaction (qRT-PCR)

Eyeballs were freshly dissected in ice-cold sterile PBS under a surgical microscope to get the retina

intact. The RNA from whole retinas was extracted via TRIzol Reagent (Invitrogen) according to the

manufacturer's guidelines. The retinal RNA was then reverse transcribed to cDNA with a high-capacity

reverse transcription cDNA kit (Applied Biosystems). Primer pairs specific to mouse RFC1/SLC19a1

(Mm00446220_m1) and Tubb4a (Mm00726185_s1, beta 4A class IVA as a housekeeping gene) were

constructed and verified by Life Technologies to use with TaqMan qPCR chemistry (TaqMan™ Gene

Expression Assay, Applied Biosystems™, 4331182). Amplicon context sequences for SLC19a1 and

beta 4A class IVA are shown in Table 2. qRT-PCR analysis was performed to calculate the relative fold

change in mRNA expression level of the SLC19a1 after siRNA administration. The critical threshold

cycle (CT) of SLC19a1 gene was first normalized to the beta-tubulin 4A with the comparative CT

method. The ΔCT which is the difference in CT values between the SLC19a1 and beta-tubulin 4A genes

were then normalized to the corresponding ΔCT of the control siRNA group, expressed as $\Delta\Delta$CT and

calculated as fold change ($2^{-\Delta\Delta CT}$).

Symbols	The full names of genes	Accession numbers of transcripts	Amplicon context sequences
RFC1/SLC19a1	Mus musculus solute carrier family 19 (folate transporter), member 1 [Mus musculus (house mouse)]	NM_031196.3:1240-1390	CGGGACATCTGGGTGTGCT ACGTGACCTTTGTGCTTTT CCGTGGGGCCTACCAGTTC CTTGTGCCCATTGCCACTT TTCAGATTGCGTCTTCCCT GTCTAAAGAGCTCTGTGCA TTGGTCTTTGGGATCAACA CTTTCCTAGCTACTGCGC
Tubb4a	Mus musculus tubulin, beta 4A class IVA (Tubb4a)	NM_009451.3:1852-2030	CCACCTTCTTAGATCTTGA AAATCCTTTCCTTTATGCC CTGTCCCTTCCCCAGCACT CCTGAACCATTCTCCTCCA CCTCTGACATGCCCAGATA TTTCTCTAGATCTTAACAT TTTCCCCAGCCTCCAAGTC CTCTCACCTTGCCTTACCT CTGACCTCACAGCTACCCA CTCCTGAC

Table 2. Gene symbols, the full names of the genes, accession number of transcripts and amplicon

context sequences of SLC19a1 and beta 4A class IVA.

Retinal Ischemia Model

The retinal ischemia model used here was previously developed in our laboratory (30). In brief,

anesthetized mice were put in a prone position under a stereoscope (SMZ1000, Nikon Instruments Inc.,

Amsterdam, The Netherlands) and their heads were immobilized with a nosepiece. Central retinal

arteries located in the central optic nerve were exposed via cautious retroorbital dissection. A small strip

(0.3×1 mm) of 30% $FeCl_3$–soaked filter paper was placed on the optic nerve for 3 min to trigger clot

formation and occlude the central retinal artery. Based on our previous observations, we used 1 hr of

ischemia, as it was sufficient to induce ischemic changes in the microcirculation such as mural cell

contractions (18, 19). One hr later, eyeballs were collected under anesthesia, and animals were sacrificed

by cervical dislocation.

Statistical Analyses

All in vivo experiments were repeated in at least n=3 mice/group. The number of mice per group for *in*

vivo experiments are mentioned in the figure legends. All results were conveyed as mean ± standard

error of the mean (S.E.M.). Data were analyzed using IBM SPSS 23. Non-normally distributed data

were compared using the Mann-Whitney U test (for two groups). The expression ratio in a treatment

group was expressed as a percentage of the ratio in the control group. p≤0.05 was considered significant.

RESULTS

RFC1 protein is abundantly expressed in the endothelial cells and pericytes of the inner BRB

The scarcity of antibodies for RFC1 immunohistochemistry (21) until recently was a challenge for

studies; hence, most former studies used RFC1 antiserum or homemade antibodies to mark RFC1

protein in the tissues (21, 31, 32). The availability of commercial RFC1 antibodies that were confirmed

by immunohistochemistry and by Western Blotting (1, 2) facilitated the research in the field.

We first immunohistochemically examined RFC1 in the coronal retina sections, and as expected,

determined that RPE was immunopositive (Fig. 1A). This was also confirmed in whole mount retina

preparations (Fig. 1B) (33). RFC1 immunopositivity was also found in the ganglion cell layer (GCL),

inner and outer plexiform layers (INL and OPL) that coincided with the superficial and deep

microvascular plexus. Hence, the RFC1 immunosignal was identified in the retinal layers comprising

microvessels (Fig. 1A, C). We used two different antibodies, high concentrations, and long incubation

durations like three overnights at +4°C with the primary antibody for the 200 μm thick whole mount

retinal preparations to visualize retinal microvessels. Likewise, isolating retinal microvessels with the

trypsin digestion method provided us with an easier and more consistent preparation for

immunostainings.

We observed that RFC1 was distributed continuously and diffusely throughout the retinal microvessels

(Fig. 1D). RFC1 was immunohistochemically co-labeled with the endothelial marker, cluster of

differentiation 31 (CD31), whose immunoreactivity exclusively outlined the endothelial cells. As a

convincing finding that there was no crosstalk in immunoreactivity, RFC1 labeling was determined to

colocalize with both the endothelial cells, and the 'bump on a log' shaped cells (Fig. 1E, arrows), which

designates pericytes.

We then labeled the microvessels with well-defined pericyte markers including the neural/glial antigen

2 (NG2), aminopeptidase-N (CD13), and platelet-derived growth factor receptor beta (PDGFR-β) all of

which showed colocalization with RFC1 (Fig. 1F-H). These indicated that RFC1 protein was present

both in the endothelial cells and pericytes.

Retinal RFC1 is downregulated *in vivo* by RFC1 targeted Accell siRNA, and leads to disruption

of the inner BRB

We delivered a combination containing the same amount of specifically designed two RFC1 targeted

Accell siRNAs (RFC1-siRNA) or two scrambled siRNAs intravitreally and sacrificed the mice 24 hr

later. To confirm the knockdown of RFC1, we obtained fresh whole retinas and determined a

pronounced decrease in RFC1 protein via Western Blotting (p= 0.028; Fig. 2A, B). Additionally, RFC1-

siRNA significantly decreased RFC1 mRNA by 24.75% compared to scrambled siRNA (p= 0.004, Fig.

2C). In line with these, RFC1 immunosignal intensity was diminished in the microvessels of RFC1-

siRNA injected retinas when mean grey value of RFC1 immunosignal (i.e., fluorescence intensity) was

measured (72.1%, p<0.0001; Fig. 2C, D). We also noticed that RFC1-siRNA treated retinas were fragile,

and prone to damage during the procedures.

The immunostaining of the whole mount retinas also confirmed RFC1-siRNA knockdown with

diminished RFC1 immunosignal, and further showed that Lectin staining was weaker and interrupted in

RFC1-siRNA treated preparations, indicating that microvessels might be structurally damaged (Fig.

2G). In addition, these retinal microvessels demonstrated lower NG2 signal compared to scrambled

siRNA treated ones, which suggested that pericytes might also be damaged by RFC1-siRNA (Fig. 2E).

However, to address if the reduction in RFC1 immunoreactivity in siRNA treated microvessels might

be associated with the loss of pericytes, we assessed pericyte density (soma per millimeter capillary).

We found no significant decline in pericyte density in RFC1-siRNA administered retinas suggesting

that RFC1 decrease may not affect pericyte maintenance, as it did not reduce pericytes in number at

least at this time point (Fig. 2F). However, we could not exclude disruption of pericyte functions via

RFC1-siRNA since there is marked decline in the expression of pericyte marker NG2.

As we determined that retinal microvessels express RFC1 protein abundantly, and we could efficiently

knock down RFC1 *in vivo*, we elucidated whether RFC1 suppression may lead to any changes in the

inner BRB.

Strikingly, knocking down retinal RFC1 diminished immunoreactivity of the tight junction-associated

transmembrane proteins occludin, claudin-5, and cytoplasmic adaptor protein zonula occludens-1 (ZO-

1), as well as the main basement membrane protein collagen-4 in trypsin digested microvessels (Fig.

2H-K). In RFC1-siRNA treated microvessels, immunoreactivity of collagen-4 that surrounds the

abluminal membrane of endothelial cells and covers pericytes residing in the basement membrane

decreased by 68% (p=0.0043), occludin decreased by 57.03% (p=0.0043), and claudin-5 decreased by

76.26% (*p*= 0.0006). ZO-1, the intracellular adaptor protein establishing a link between occludin and

intracellular actin cytoskeleton decreased by 53.22% (p= 0.0012).

Retinal RFC1 is upregulated *in vivo* by LV overexpressing RFC1, and this upregulation changes

the inner BRB properties

Next, we aimed to increase the expression of RFC1 in mouse retinas via LV carrying RFC1 as described

in detail in the methods section. We preferred the LV gene delivery method due to its efficiency to

induce stable gene expression, its tropism to the inner retina and the cells of retinal microvessels when

delivered intravitreally (34, 35).

The efficient RFC1 protein overexpression was shown by Western Blotting in cell lysates (Fig. 3A).

Subsequently, we proceeded to validate the *in vivo* LV-mediated transduction. Mice had unilateral

intravitreal LV injections. No animals developed any local infections. The eyeballs of mice (n=3) that

received intravitreal Control-LV bearing GFP were harvested 10 days after. The retinal and

perimicrovascular cells showed GFP expression disclosing that the cells were infected with LV (Fig.

3B).

The overexpression of RFC1 by LV in the retina was also observed immunohistochemically. RFC1

immunoreactivity revealed a marked intensity increase in both the retinal whole mount and retinal

microvessel preparations in RFC1-LV compared to Control-LV (Fig. 3C, D) with 167.5% increase in

RFC1 immunosignal (p= 0.0159; Fig. 3D).

As the suppression of RFC1 levels led to an extensive structural damage in retinal microvessels, we

further investigated whether RFC1 overexpression had any effects. Immunohistochemistry of tight

junction proteins (occludin, claudin-5), intracellular adaptor protein ZO-1, and main basement

membrane protein collagen-4 in retinal microvessels (from n=3 retina/per marker, Fig. 3E-H) were done

ten days after administration. RFC1-LV administered retinal microvessels displayed increased

immunosignal in occludin (p= 0.0059), claudin-5 (p= 0.0040), and collagen-4 (p= 0.0159) compared to

Control-LV ones (Fig. 3E-G). However, the immunosignal of ZO-1 did not differ between the groups

(Fig. 3H).

Retinal ischemia alters RFC1 protein and decreases the inner BRB proteins which can be

ameliorated by RFC1 overexpression

After observing the potential role of RFC1 protein in maintaining the inner BRB, we investigated

whether RFC1 protein has a role in retinal ischemia. We made 1 hr permanent retinal ischemia, as it is

considered sufficient to observe microvessel-related changes such as capillary constrictions or

protrusion of the contracted pericyte soma from the microvessel wall (Fig. 4C, white asterisk), and yet

an early time point to allow us to minimize the effects of inflammation (18-20).

Following ischemia, the expression of RFC1 protein was enhanced significantly compared to controls

detected by Western blotting (p= 0.025; Fig. 4A, B) and RFC1 immunopositivity (p <0.0001; Fig. 4C).

The pronounced RFC1 signal was detected in the protruding pericytes (Fig. 4C, white arrows) probable

of an increased signal in affected cells.

We studied whether retinal ischemia caused any alterations in the inner BRB proteins and observed that

1 hr retinal ischemia decreased collagen-4 and occludin immunosignal compared to controls ($p =0.0140$,

$p =0.0317$ respectively; Fig. 4D, E).

Finally, we studied whether RFC1-siRNA knockdown induced disruption of the inner BRB proteins

resulted in barrier dysfunction. We stained vessels with lectin and determined endogenous IgG

extravasation by fluorescently labeled anti-mouse IgG antibody (Fig. 5A, red arrows). RFC1-siRNA

treated mice displayed endogenous IgG leakage (Fig. 5A, red arrows), while scrambled-siRNA treated

retinas did not show any. This indicated that RFC1 is essentially required for the structural and

functional integrity of the inner BRB in health.

On the other hand, we observed no IgG extravasation after 1 hr retinal ischemia, so it significantly did

not compromise the integrity of inner BRB (Fig. 5B). This observation was consistent with the literature

since 1 hr retinal ischemia causes early changes such as capillary constrictions (white arrows, Fig. 5B),

but it does not suffice for the inner BRB disruption to the extent of allowing endogenous IgG (150 kDa)

leakage through paracellular barrier even if disintegration of tight junctions commenced (36-38). We

then studied whether knocking-down RFC1 before ischemia may provoke disruption of the inner BRB.

When RFC1-siRNA was administered 24 hr before retinal ischemia, a significant amount of endogenous

IgG extravasation was determined (Fig. 5D, red arrows). As scrambled siRNA administration 24 hr

before retinal ischemia led to no apparent IgG extravasation, this indicated that decreased RFC1 levels

accelerated the inner BRB disruption in acute retinal ischemia.

DISCUSSION

In this report, for the first time, we demonstrated the presence of RFC1 protein in the endothelial cells

and pericytes of the inner BRB. We also showed that RFC1 - once considered only as a transporter

protein - is an essential component for maintaining the inner BRB and has fundamental roles in

pathological conditions affecting the inner BRB, such as retinal ischemia.

RFC1 expression at the protein level in endothelial cells and pericytes of retina has never been studied

before, despite its consistent detection in transcripts of microvascular mural cells (pericytes and vascular

smooth muscle cells) of the brain (39). Recently, a study characterized the localization of folate

transporters, hence RFC1 expression at CNS barriers in detail focusing especially on BBB cells

including the endothelial cells; however, this study explored neither the retina nor pericytes (40). So,

our study bears another focus, and offers a recognition for the roles of RFC1 in the CNS and deepens

our understanding of other critical functions as well as novel treatment strategies for brain disorders.

Isolating retinal microvessels with the trypsin digestion method provided us an easier and more

consistent preparation for immunostainings. Our results highlight that, retinal endothelial cells, like their

counterparts in the CNS, express RFC1 protein. As an additional novel finding we detected the

colocalization of RFC1 with the well-defined pericyte markers. Most studies evaluating the role of RFC1

in microvessels concentrated on only the endothelial cells, but not the neighboring pericytes (1, 2, 31,

41). In this regard, RFC1 protein was never characterized in CNS pericytes as well. Our findings actually

match the results of a recent meta-analysis combining five different mouse brain pericyte transcriptomic

studies where the RFC1 gene was one of the only three consistently detected genes in brain pericytes

out of the 1180 enriched genes (39).

Our study provides some initial insights regarding the functional importance of RFC1 in retinal

microvessels. The genetic knockout of RFC1 is lethal (42) and no conditional knockout animals had

been used in the studies before, so we decided to utilize siRNA technology to manipulate RFC1 levels

specifically in the retina of adult mice. Two previous *in vitro* studies using RFC1-siRNA which were

performed in primary cultures of human differentiated adipocytes (43), and in rat choroidal epithelial

Z310 cells (44), and both did not focus on the microvessels or the retina. Hence, our method to

knockdown RFC1 *in vivo* by RFC1 targeted custom designed Accell RFC1-siRNA could be considered

as one of the initial studies. RFC1-siRNA administration to the retina *in vivo* via intravitreal delivery

offers some advantages: 1) the retina is the easily reachable, well-known extension of the brain 2) the

vitreous cavity is an enclosed space that minimizes the adverse effects of systemic spread and maximizes

the effect of applied concentration 3) delivery is fast and feasible (45). The application of RFC1-siRNA

deranged the structure of retinal microvessels, specifically the deep plexus comprising blood barrier

properties, as shown by lectin stainings (Fig. 2E). Interestingly, knocking down RFC1 led to a significant

decrease in the barrier proteins occludin, claudin-5, intracellular adaptor protein ZO-1, and the main

basement membrane protein collagen-4. This induced the inner BRB disintegration and led to functional

failure of the barrier properties. Although it may be plausible to interpret this striking result with caution,

there is a study showing that the tight junction proteins like occludin, claudin-1, and ZO-1 were

substantially decreased in the BBB of capillaries isolated from proton coupled folate transporter (PCFT)

null mice in comparison to controls (46). However, this mouse with deleted PCFT gene - another

functionally distinct folate transporter- is accepted as a model for hereditary folate malabsorption that

leads to systemic folate deficiency. Hence, it is not possible to fully exclude the contribution of chronic

folate deficiency to these effects. However, in our study, we observed inner BRB changes after acute

modifications of the transporter in naive mice, which circumvents the confounding developmental

effects of folate deficiency. In line with these observations, our experiments with the retinal

overexpression of RFC1 via LV have shown an upregulation of occludin, claudin-5 and collagen-4,

supporting the role of RFC1 in inner BRB protein expressions. As an interesting observation, RFC1 was

recently determined as a cGAMP importer in the cyclic GMP–AMP synthase (cGAS)–stimulator of

interferon genes (STING) pathway that has a role in inflammation, infection, cellular stress, tissue

damage, and tumor angiogenesis (21, 47, 48). In mouse tumor models, intratumoral cGAMP treatment

led to a 40% decrease in the density of the vessels, a 1.7-fold rise in pericyte coverage, and a 1.5-fold

rise in collagen-4 coverage (47), which may indicate that RFC1 may indirectly involved in remodeling

the tumor vasculature in consistent with our observations in inner retinal microvasculature.

Based on reports that show pericytes are eminent for BBB and inner BRB maintenance and contribute

to tight junction formation and preservation (49-52), we also excluded pericyte deficiency in RFC1-

siRNA treated retinas. Although we found no significant decrease in pericyte density by counting

pericyte soma per mm capillary in RFC1-siRNA treated mice, we observed immunosignal decrease in

pericyte marker NG2, which involves in many functions of pericytes including proliferation, motility

and importantly, endothelial cell junction assembly by activating integrin signaling (53). For example,

the treatment of human microvascular pericytes with NG2-siRNA hindered collagen-4 coverage, ZO-1

expression, and endothelial junction maintenance in endothelial cells. Likewise, tumors from mice

whose pericytes do not express NG2 showed similar changes in their endothelial junctions in the same study (28). Although the underlying mechanisms require clarification, we suggest that NG2 might be a mediator of the effects observed in tight junction proteins and collagen-4 via RFC1-siRNA. Alternatively, the disintegration of NG2 in RFC1-siRNA treated retinal microvessels may indicate more comprehensive dysfunction throughout the inner BRB, to which NG2 disruption is only a contributing factor.

Aside from its role in the maintenance of inner BRB under physiologic conditions, we further investigated the effect of RFC1 in acute retinal ischemia. Retinal ischemia which is a major underlying condition of blindness worldwide is a severe condition with a poor prognosis generally caused by acute occlusion of retinal arteries (54). We induced retinal ischemia by an established method in our laboratory (19, 30), and showed an increase in RFC1 protein after 1 hr permanent ischemia. A previous gene profiling study showed RFC1 mRNA upregulation following 1 hr ischemia / 24 hr recanalization induced by high intraocular pressure retinal ischemia model in rats (55). In contrast, the same study observed no significant change in RFC1 mRNA in 1 hr permanent retinal ischemia. Another study showed that 45 min retinal ischemia followed by 48 hr recanalization upregulated RFC1 mRNA in rats (56). The possible explanation for these results may be that the retinal ischemia models were different, and the authors only determined mRNA levels, not the protein levels of the samples. The estimation of protein levels from mRNA levels might be unreliable, as post-translational changes could affect the protein levels in early time-points (57). There is only one study performed by kidney ischemia-reperfusion model that determined RFC1 protein levels decreased *in vivo* focusing on the proximal tubules where folate reabsorption occurs, emphasizing its importance in adjusting serum folate levels (58). Contrarily, the increase of metabolic needs due to acute phase of ischemia might necessitate the upregulation of RFC1 after 1 hr of retinal ischemia for the preservation of the barrier properties. In line with that, further augmentation of RFC1 expression by LV-RFC1 intervention before ischemia salvaged the decreased occludin and collagen-4 levels, otherwise which would be attenuated by ischemia.

Also, knocking down RFC1 prior to ischemia impaired barrier functions which were demonstrated by endogenous IgG leakage, supporting our findings that RFC1 is essential for the protection of inner BRB

during retinal ischemia. This result might be interpreted in the light of the human meta-analysis showing

that the SNPs of RFC1 gene differed between ischemic stroke and control groups, and some genotypes

were found to be associated with small vessel occlusions and silent brain infarctions (59). Since the

changes in RFC1 function or structure with these polymorphisms had not been defined, it is possible

that our finding of changes in inner BRB via the genetic modifications of RFC1 could carry the potential

of being the experimental evidence of the importance of RFC1 in ischemia and small vessel disease.

Thus, intriguing questions regarding the therapeutic potential of intervening RFC1 before, during or

after ischemia rises.

Our study can be considered as an initial step to investigate the role of 'once an unnoticed transporter

protein' in the inner BRB. We took a step-by-step approach, defined the presence of the RFC1 protein

in the retinal microvessels and established *in vivo* genetic tools to clarify the potential new roles of RFC1

under physiologic conditions, ischemia. However, some limitations of our study merit consideration.

The genetic manipulations we used were not targeted to cells. In addition, we did not study the

underlying pathways of how RFC1 regulates barrier proteins or BRB integrity, and the interplay of folate

and other folate transport systems (FRα and PCFT) or the concert of the efflux systems (Pgp/ABCB1,

MRPs/ABCC, BCRP/ABCG2) (60), which warrants further investigations.

Despite these limitations, our data suggest several implications. We propose a conceptual model of

workflow for the other proteins like RFC1 whose presence and roles have not been elucidated so far.

Nowadays, the roles of these proteins have been investigated through advanced mathematical models,

pathway analyses and the -omics approaches, but the old phrase "seeing is believing" should not be

underestimated and *in vivo* studies may provide surprising insights into the role of a protein. The

upregulation of RFC1 may be attempted for diseases other than ischemia where the impairment of BRB

or BBB is involved in the pathophysiology. Although merely an initial step, our results are encouraging

as the presence of RFC1 in inner BRB may be exploited for targeted drug delivery such as RFC1-

targeted nanodrugs, folate conjugated nanodrugs or even clinically widely used RFC1 substrate,

methotrexate, can be exploited not only in ischemia, but in diseases like cancer.

CONCLUSION

In the present study, we investigated the presence of RFC1 in the retinal endothelial cells and pericytes, and its role in the regulation of inner BRB integrity. For the first time, we introduced and discussed the tools for the manipulation of RFC1 levels or function *in vivo* which included RFC1-targeted Accell siRNA to knockdown RFC1, and LV to overexpress RFC1. By using these tools, we tried to elucidate the role of RFC1 in the abrupt regulation of inner BRB during physiological conditions and acute retinal ischemia and highlighted its potential role for preserving BRB integrity. This study aimed to precede the further investigation of RFC1 in other tissues and diseases by suggesting a conceptual and methodological framework. Future research may extend this work by integrating pathways which RFC1 is associated with to address the concern of how RFC1 regulates inner BRB.

Abbreviations:

BBB: blood-brain barrier; BRB: blood-retina barrier; CD13: Aminopeptidase N; cGAMP: Cyclic guanosine monophosphate–adenosine; cGAS: Cyclic GMP-AMP synthase; GCL: ganglion cell layer; IgG: Immunoglobulin G; INL: inner nuclear layer; IPL: inner plexiform layer; LSCI: Laser speckle contrast imaging; LV: Lentiviral vectors; mRNA: messenger RNA; MTX: Methotrexate; NG2: Neural/glial antigen 2; ONL: outer nuclear layer; OPL: outer plexiform layer; PBS: phosphate buffered saline; PDGFR-β: platelet-derived growth factor receptor beta; PFA: paraformaldehyde; qRT-PCR: Quantitative Reverse Transcriptase-Polymerase Chain Reaction; RBF: Relative blood flow; RFC1: Reduced Folate Carrier 1; RIPA: Radioimmunoprecipitation assay; RPE: retinal pigmented epithelium; siRNA: short interfering RNA; SLC19a1:solute carrier family 19 (folate transporter), member 1; SNPs: single nucleotide polymorphisms; STING: stimulator of interferon genes; UV: ultraviolet; ZO-1: Zonula occludens 1

Data availability

All data generated or analyzed during this study are included in this published article. Raw data and analyses can be provided upon request. There are no restrictions on data availability.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by The Scientific and Technological Research Institution of Turkey (TÜBİTAK; Grant No: 120N690) and Hacettepe University Scientific Research Coordination Unit (Project No: TDK-2020-18590).

Authors' contributions

GG contributed to the study design, performed surgeries and experiments, obtained tissues, performed immunohistochemistry, collected images and data, analyzed the data, prepared figures, wrote the draft of the manuscript and contributed to editing. NB contributed to the design and execution of Western Blottings and collected membrane images. MCB designed and synthesized the lentiviral vector and performed cell-line Western Blottings. MSB contributed to the design and execution of qRT-PCR experiments and performed related analysis. GU contributed to the in vivo application of retinal ischemia model, retinal trypsin digestion and intravitreal injections, and writing of the paper. EK provided lentiviral vector as a courtesy and contributed to editing. MY acquired funds and administered the project, coordinated, and supervised the work, wrote the manuscript, and contributed to figures and edited the paper. All authors read and approved the final manuscript and provided minor modifications.

Acknowledgments

We thank Dr. Canan Cakir Aktas (The Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara, Turkey) for assisting the financial management of the project, Dr. Buket Nebiye Demir (The Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara, Turkey) for technical advice on in vivo procedures, Dr. Cetin Demir (Faculty of Medicine, Department of Pediatrics, Division of Pediatric Oncology, Drug Resistance Laboratory, Hacettepe University, Ankara, Turkey) for his assistance in the execution of qRT-PCR experiments and our laboratory technician

Mesut Firat (The Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara,

Turkey) for providing technical assistance throughout project. We are grateful to our undergraduate

researchers Derin Nalcakan and Neslihan Nisa Gecici (Hacettepe University Medical School), who were

officially added as interns to the project with "TÜBİTAK-Intern Researcher Scholarship Programme",

for their contribution to immunohistochemistry and data analysis.

Supplementary information is available for this paper.

Correspondence and requests for materials should be addressed to MY or GG.

REFERENCES

1.   Alam C, Hoque MT, Finnell RH, Goldman ID, Bendayan R. Regulation of Reduced Folate Carrier
(RFC) by Vitamin D Receptor at the Blood-Brain Barrier. Mol Pharmaceut. 2017;14(11):3848-58.
2.   Alam C, Aufreiter S, Georgiou CJ, Hoque MT, Finnell RH, O'Connor DL, et al. Upregulation of
reduced folate carrier by vitamin D enhances brain folate uptake in mice lacking folate receptor alpha.
Proc Natl Acad Sci U S A. 2019;116(35):17531-40.
3.   Figueiredo JC, Levine AJ, Grau MV, Midttun O, Ueland PM, Ahnen DJ, et al. Vitamins B2, B6,
and B12 and risk of new colorectal adenomas in a randomized trial of aspirin use and folic acid
supplementation. Cancer Epidemiol Biomarkers Prev. 2008;17(8):2136-45.
4.   Balashova OA, Visina O, Borodinsky LN. Folate action in nervous system development and
disease. Dev Neurobiol. 2018;78(4):391-402.
5.   Desai A, Sequeira JM, Quadros EV. The metabolic basis for developmental disorders due to
defective folate transport. Biochimie. 2016;126:31-42.
6.   Mentch SJ, Locasale JW. One-carbon metabolism and epigenetics: understanding the
specificity. Ann N Y Acad Sci. 2016;1363:91-8.
7.   Lehmann M, Regland B, Blennow K, Gottfries CG. Vitamin B12-B6-folate treatment improves
blood-brain barrier function in patients with hyperhomocysteinaemia and mild cognitive impairment.
Dement Geriatr Cogn Disord. 2003;16(3):145-50.
8.   Snowdon DA, Tully CL, Smith CD, Riley KP, Markesbery WR. Serum folate and the severity of
atrophy of the neocortex in Alzheimer disease: findings from the Nun study. Am J Clin Nutr.
2000;71(4):993-8.
9.   Sofi F, Marcucci R, Bolli P, Giambene B, Sodi A, Fedi S, et al. Low vitamin B6 and folic acid levels
are associated with retinal vein occlusion independently of homocysteine levels. Atherosclerosis.
2008;198(1):223-7.
10.   Cahill MT, Stinnett SS, Fekrat S. Meta-analysis of plasma homocysteine, serum folate, serum
vitamin B(12), and thermolabile MTHFR genotype as risk factors for retinal vascular occlusive disease.
Am J Ophthalmol. 2003;136(6):1136-50.
11.   Kaur C, Foulds WS, Ling EA. Blood-retinal barrier in hypoxic ischaemic conditions: basic
concepts, clinical features and management. Prog Retin Eye Res. 2008;27(6):622-47.
12.   London A, Benhar I, Schwartz M. The retina as a window to the brain-from eye research to CNS
disorders. Nat Rev Neurol. 2013;9(1):44-53.
13.   Cunha-Vaz J, Bernardes R, Lobo C. Blood-retinal barrier. Eur J Ophthalmol. 2011;21 Suppl 6:S3-
9.
14.   Naggar H, Ola MS, Moore P, Huang W, Bridges CC, Ganapathy V, et al. Downregulation of
reduced-folate transporter by glucose in cultured RPE cells and in RPE of diabetic mice. Invest
Ophthalmol Vis Sci. 2002;43(2):556-63.

15.   Naggar H, Fei YJ, Ganapathy V, Smith SB. Regulation of reduced-folate transporter-1 (RFT-1) by
homocysteine and identity of transport systems for homocysteine uptake in retinal pigment epithelial
(RPE) cells. Exp Eye Res. 2003;77(6):687-97.
16.   Bridges CC, El-Sherbeny A, Ola MS, Ganapathy V, Smith SB. Transcellular transfer of folate
across the retinal pigment epithelium. Curr Eye Res. 2002;24(2):129-38.
17.   Smith SB, Huang W, Chancy C, Ganapathy V. Regulation of the reduced-folate transporter by
nitric oxide in cultured human retinal pigment epithelial cells. Biochem Biophys Res Commun.
1999;257(2):279-83.
18.   Yemisci M, Gursoy-Ozdemir Y, Vural A, Can A, Topalkara K, Dalkara T. Pericyte contraction
induced by oxidative-nitrative stress impairs capillary reflow despite successful opening of an occluded
cerebral artery. Nature Medicine. 2009;15(9):1031-U82.
19.   Alarcon-Martinez L, Yilmaz-Ozcan S, Yemisci M, Schallek J, Kilic K, Villafranca-Baughman D, et
al. Retinal ischemia induces alpha-SMA-mediated capillary pericyte contraction coincident with
perivascular glycogen depletion. Acta Neuropathol Commun. 2019;7(1):134.
20.   Alarcon-Martinez L, Yilmaz-Ozcan S, Yemisci M, Schallek J, Kilic K, Can A, et al. Capillary
pericytes express alpha-smooth muscle actin, which requires prevention of filamentous-actin
depolymerization for detection. Elife. 2018;7.
21.   Ritchie C, Cordova AF, Hess GT, Bassik MC, Li L. SLC19A1 Is an Importer of the
Immunotransmitter cGAMP. Mol Cell. 2019;75(2):372-81 e5.
22.   Li W, Yuan P, Liu W, Xiao L, Xu C, Mo Q, et al. Hypoxia-Immune-Related Gene SLC19A1 Serves
as a Potential Biomarker for Prognosis in Multiple Myeloma. Front Immunol. 2022;13:843369.
23.   Chou JC, Rollins SD, Fawzi AA. Trypsin digest protocol to analyze the retinal vasculature of a
mouse model. J Vis Exp. 2013(76):e50489.
24.   Zudaire E, Gambardella L, Kurcz C, Vermeren S. A computational tool for quantitative analysis
of vascular networks. PLoS One. 2011;6(11):e27385.
25.   Discovery H. Accell™ siRNA References 8100 Cambridge Research Park, Waterbeach,
Cambridge, CB25 9TL, United Kingdom2021 [Available from: https://horizondiscovery.com/-
/media/Files/Horizon/resources/Reading-lists/accell-sirna-ref-reading.pdf.
26.   Taniguchi T, Endo KI, Tanioka H, Sasaoka M, Tashiro K, Kinoshita S, et al. Novel use of a
chemically modified siRNA for robust and sustainable in vivo gene silencing in the retina. Sci Rep.
2020;10(1):22343.
27.   Calkins DJ, Yu-Wai-Man P, Newman NJ, Taiel M, Singh P, Chalmey C, et al. Biodistribution of
intravitreal (lenadogene) nolparvovec gene therapy in nonhuman primates. Mol Ther Methods Clin
Dev. 2021;23:307-18.
28.   You WK, Yotsumoto F, Sakimura K, Adams RH, Stallcup WB. NG2 proteoglycan promotes tumor
vascularization via integrin-dependent effects on pericyte function. Angiogenesis. 2014;17(1):61-76.
29.   Beker M, Caglayan AB, Beker MC, Altunay S, Karacay R, Dalay A, et al. Lentivirally administered
glial cell line-derived neurotrophic factor promotes post-ischemic neurological recovery, brain
remodeling and contralesional pyramidal tract plasticity by regulating axonal growth inhibitors and
guidance proteins. Exp Neurol. 2020;331:113364.
30.   Karatas H, Erdener SE, Gursoy-Ozdemir Y, Gurer G, Soylemezoglu F, Dunn AK, et al. Thrombotic
distal middle cerebral artery occlusion produced by topical FeCl(3) application: a novel model suitable
for intravital microscopy and thrombolysis studies. J Cereb Blood Flow Metab. 2011;31(6):1452-60.
31.   Hinken M, Halwachs S, Kneuer C, Honscha W. Subcellular localization and distribution of the
reduced folate carrier in normal rat tissues. Eur J Histochem. 2011;55(1):e3.
32.   Trippett TM, Garcia S, Manova K, Mody R, Cohen-Gould L, Flintoff W, et al. Localization of a
human reduced folate carrier protein in the mitochondrial as well as the cell membrane of leukemia
cells. Cancer Res. 2001;61(5):1941-7.
33.   Chancy CD, Kekuda R, Huang W, Prasad PD, Kuhnel JM, Sirotnak FM, et al. Expression and
differential polarization of the reduced-folate transporter-1 and the folate receptor alpha in
mammalian retinal pigment epithelium. J Biol Chem. 2000;275(27):20676-84.

34.   Wei Y, Gong J, Xu Z, Thimmulappa RK, Mitchell KL, Welsbie DS, et al. Nrf2 in ischemic neurons
promotes retinal vascular regeneration through regulation of semaphorin 6A. Proc Natl Acad Sci U S A.
2015;112(50):E6927-36.
35.   Auricchio A, Kobinger G, Anand V, Hildinger M, O'Connor E, Maguire AM, et al. Exchange of
surface proteins impacts on viral vector cellular specificity and transduction characteristics: the retina
as a model. Hum Mol Genet. 2001;10(26):3075-81.
36.   Wang X, Liu Y, Sun Y, Liu W, Jin X. Blood brain barrier breakdown was found in non-infarcted
area after 2-h MCAO. J Neurol Sci. 2016;363:63-8.
37.   Chen B, Friedman B, Cheng Q, Tsai P, Schim E, Kleinfeld D, et al. Severe blood-brain barrier
disruption and surrounding tissue injury. Stroke. 2009;40(12):e666-74.
38.   Jin X, Liu J, Yang Y, Liu KJ, Yang Y, Liu W. Spatiotemporal evolution of blood brain barrier
damage and tissue infarction within the first 3h after ischemia onset. Neurobiol Dis. 2012;48(3):309-
16.
39.   He L, Vanlandewijck M, Raschperger E, Andaloussi Mae M, Jung B, Lebouvier T, et al. Analysis
of the brain mural cell transcriptome. Sci Rep. 2016;6:35108.
40.   Sangha V, Hoque MT, Henderson JT, Bendayan R. Novel localization of folate transport systems
in the murine central nervous system. Fluids Barriers CNS. 2022;19(1):92.
41.   Geier EG, Chen EC, Webb A, Papp AC, Yee SW, Sadee W, et al. Profiling solute carrier
transporters in the human blood-brain barrier. Clin Pharmacol Ther. 2013;94(6):636-9.
42.   Zhao R, Russell RG, Wang Y, Liu L, Gao F, Kneitz B, et al. Rescue of embryonic lethality in
reduced folate carrier-deficient mice by maternal folic acid supplementation reveals early neonatal
failure of hematopoietic organs. J Biol Chem. 2001;276(13):10224-8.
43.   Petrus P, Bialesova L, Checa A, Kerr A, Naz S, Backdahl J, et al. Adipocyte Expression of SLC19A1
Links DNA Hypermethylation to Adipose Tissue Inflammation and Insulin Resistance. J Clin Endocrinol
Metab. 2018;103(2):710-21.
44.   Grapp M, Wrede A, Schweizer M, Huwel S, Galla HJ, Snaidero N, et al. Choroid plexus
transcytosis and exosome shuttling deliver folate into brain parenchyma. Nat Commun. 2013;4:2123.
45.   Meyer CH, Krohne TU, Charbel Issa P, Liu Z, Holz FG. Routes for Drug Delivery to the Eye and
Retina: Intravitreal Injections. Dev Ophthalmol. 2016;55:63-70.
46.   Wang X, Cabrera RM, Li Y, Miller DS, Finnell RH. Functional regulation of P-glycoprotein at the
blood-brain barrier in proton-coupled folate transporter (PCFT) mutant mice. FASEB J.
2013;27(3):1167-75.
47.   Yang H, Lee WS, Kong SJ, Kim CG, Kim JH, Chang SK, et al. STING activation reprograms tumor
vasculatures and synergizes with VEGFR2 blockade. J Clin Invest. 2019;129(10):4350-64.
48.   Luteijn RD, Zaver SA, Gowen BG, Wyman SK, Garelis NE, Onia L, et al. SLC19A1 transports
immunoreactive cyclic dinucleotides. Nature. 2019;573(7774):434-8.
49.   Al Ahmad A, Gassmann M, Ogunshola OO. Maintaining blood-brain barrier integrity: pericytes
perform better than astrocytes during prolonged oxygen deprivation. J Cell Physiol. 2009;218(3):612-
22.
50.   Yang T, Guo L, Fang Y, Liang M, Zheng Y, Pan M, et al. Pericytes of Indirect Contact Coculture
Decrease Integrity of Inner Blood-Retina Barrier Model In Vitro by Upgrading MMP-2/9 Activity. Dis
Markers. 2021;2021:7124835.
51.   Daneman R, Zhou L, Kebede AA, Barres BA. Pericytes are required for blood-brain barrier
integrity during embryogenesis. Nature. 2010;468(7323):562-6.
52.   Bell RD, Winkler EA, Sagare AP, Singh I, LaRue B, Deane R, et al. Pericytes control key
neurovascular functions and neuronal phenotype in the adult brain and during brain aging. Neuron.
2010;68(3):409-27.
53.   Stallcup WB. The NG2 Proteoglycan in Pericyte Biology. Adv Exp Med Biol. 2018;1109:5-19.
54.   Dattilo M, Newman NJ, Biousse V. Acute retinal arterial ischemia. Ann Eye Sci. 2018;3.

55.   Andreeva K, Zhang M, Fan W, Li X, Chen Y, Rebolledo-Mendez JD, et al. Time-dependent Gene
Profiling Indicates the Presence of Different Phases for Ischemia/Reperfusion Injury in Retina.
Ophthalmol Eye Dis. 2014;6:43-54.
56.   Abcouwer SF, Lin CM, Wolpert EB, Shanmugam S, Schaefer EW, Freeman WM, et al. Effects of
ischemic preconditioning and bevacizumab on apoptosis and vascular permeability following retinal
ischemia-reperfusion injury. Invest Ophthalmol Vis Sci. 2010;51(11):5920-33.
57.   Fortelny N, Overall CM, Pavlidis P, Freue GVC. Can we predict protein from mRNA levels?
Nature. 2017;547(7664):E19-E20.
58.   Yang C, Wijerathne CUB, Tu GW, Woo CWH, Siow YL, Madduma Hewage S, et al. Ischemia-
Reperfusion Injury Reduces Kidney Folate Transporter Expression and Plasma Folate Levels. Front
Immunol. 2021;12:678914.
59.   Cho Y, Kim JO, Lee JH, Park HM, Jeon YJ, Oh SH, et al. Association of reduced folate carrier-1
(RFC-1) polymorphisms with ischemic stroke and silent brain infarction. PLoS One.
2015;10(2):e0115295.
60.   Assaraf YG. The role of multidrug resistance efflux transporters in antifolate resistance and
folate homeostasis. Drug Resist Updat. 2006;9(4-5):227-46.

Figure Legends

Fig. 1. RFC1 protein is abundantly expressed in endothelial cells and pericytes of the retinal

microvessels. A) Ex-vivo labeling of 20 μm coronal cryo-section from PFA fixated eyeballs of naive

Swiss Albino mice with anti-RFC1 antibody (red). Abundant RFC1 immunopositivity is observed along

ganglion cell layer (GCL), inner plexiform layer (IPL), outer plexiform layer (OPL) where retinal

microvessels forms vascular horizontal vascular beds; as well as the retinal pigment epithelium (RPE).

However, this preparation limited the observation of microvessels, hence inner BRB, as it commonly

included microvessels circularly rather than longitudinally. Scale Bar=25 μm B) RPE which is known

to express RFC1 previously is well stained with anti-RFC1 antibody (red) as our positive control. C)

The microvessels constituting inner BRB form the deep vascular plexus of the retina were

immunohistochemically labelled with anti-RFC1 antibody in PFA fixated whole-mount retinas D)

Retinal microvessels which were obtained via retinal trypsin digestion method that allowed to get only

microvessels (< 9 μm diameter) were immunofluorescently labeled with anti-RFC1 antibody (n=6

retina; red). Nuclei were labeled with Hoechst 33258 (blue) in all the rows. E) the endothelial marker

CD31 (green) (n=3). White arrow show "bump-on-a log" shaped pericyte body was positively stained

with anti-RFC1 antibody, but negative for endothelial marker CD31. Hence, RFC1 staining was specific.

F-H) RFC1 (red) also colocalized with accepted pericyte markers NG2, CD13, PDGFR-β (green) shown

respectively (n=3/marker). Nuclei were labeled with Hoechst 33258 (blue). Scale bar: 10 μm.

Fig. 2. The validation of Accell siRNA mediated *in vivo* RFC1 knockdown, which led to a reduction

in critical proteins of the inner BRB. A) Western blotting image of retinas (n=4 mice/group) depicting

the robust decrease in RFC1 protein of RFC1-siRNA delivered ones compared to scrambled-siRNA. β-

Tubulin III was loading control. 35 μg protein was loaded for each retina. B) The graph illustrates the

quantification of the relative band densities of given Western blotting image which expresses RFC1

band density in proportion to β-Tubulin III, which shows decrease in relative protein levels in RFC1-

siRNA treated retinas (*p= 0.028). C) The graph shows that RFC1-siRNA delivery reduced retinal

RFC1 mRNA levels by 24.75% when compared to scrambled-siRNA (*p=0.004). D) The graph

illustrates the percentage of mean grey value of RFC1 frame in lectin positive microvessel area in RFC1-

siRNA treated group normalized to Scrambled-siRNA treated group as described in Methods section.

RFC1-siRNA delivery significantly reduced the percent of the mean grey values of RFC1 by 72.10%

(n=4). Representative confocal images of Scrambled-siRNA or RFC1-siRNA delivered retinal

microvessels stained by anti-RFC1 antibody (red). E) RFC1-siRNA treated retinal microvessels showed

less NG2 immunosignal compared to Scrambled-siRNA treated ones, indicating that pericytes might

also be damaged by RFC1-siRNA. F) However, pericyte body counts per microvessel length (mm) were

not different between the groups. n = 3 images were analyzed per animal. G) Whole mount retinas were

labeled by anti-RFC1 antibody (red) and Fluorescein Lectin (green). RFC1 immunosignal was decreased

and interrupted along deep retinal microvessels. Also, lectin signal was weakened in RFC1-siRNA

treated retinas representing the structural decomposition of microvessels. H-K) RFC1-siRNA delivery

significantly reduced the percent of the mean grey values of Collagen-4 68% (n=4), Occludin 57.03%

(n=3), Claudin-5 76.26% (n=4), and ZO-1 53.22% (n=4). The reductions that are statistically significant

represented as *p ≤ 0.05. Immunosignal decrease was also observed collagen-4 (magenta), Occludin

(cyan), Lectin (Yellow) as well as G, H) Claudin-5 (magenta), and ZO-1 (cyan) via RFC1-siRNA

treatment compared to Scrambled-siRNA treatment. Nuclei are labelled with Hoechst 33258 (blue),

Data are mean ± S.E.M Mann-Whitney U; Scale bar: 10 μm.

Fig. 3. The validation of lentiviral vector-mediated overexpression of RFC1 protein *in vivo*, which led to an increase in the proteins of BRB.

A) The Western Blotting image shows corresponding overexpression of RFC1 protein in RFC1-LV transduced neuroblastoma cells (N2a) compared to Control-LV transduced cells. For cell culture lysates, Beta-Actin was the loading control. B) The graph illustrates the quantification of the relative band densities of given Western blotting image which expresses RFC1 band density in proportion to β-Actin, which shows increase in relative protein levels in RFC1-LV treated retinas (*p= 0.016). B) The confocal images from whole mount retinas confirmed the intravitreal delivery of Control-LV into the mouse eye showing lentiviral vectors infected various retinal cells, as reporter gene GFP (green) signal indicated. Also, as magnified image showed, perivascular cells and pericytes expressed GFP protein in LV-RFC1 or Control-LV injected retinas, microvessel trace was shown by dashed line. C) The representative images of whole mount retinas which had been treated by LV-RFC1 showed increased RFC1 immunosignal (red) compared to Control-LV injected control retinas (n=3). D) The graph illustrates the percentage of mean grey value of RFC1 frame in lectin positive microvessel area in RFC1-LV treated group normalized to Control-LV treated group as described in Methods section. RFC1-LV delivery significantly increased the percent of the mean grey values of RFC1 (n=3). E-H) RFC1-LV delivery significantly increased the percent of the mean grey values of Collagen-4 (n=4), Occludin (n=3), Claudin-5 (n=3), but not ZO-1 compared to Control-LV delivered groups. (*p ≤ 0.05). Collagen-4 (magenta), Occludin (cyan) Lectin (Yellow) immunosignal increased as well as Claudin-5 (magenta) except ZO-1 (cyan) via RFC1-LV treatment compared to Control-LV treatment. Nuclei were labelled with Hoechst 33258 (blue) Data are mean ± S.E.M. Mann-Whitney U; Scale bar: 10 μm.

Fig. 4. RFC1 protein increased after retinal ischemia, and the overexpression of RFC1 before ischemia by LV salvaged decreased collagen-4 and occludin levels.

A) The representative Western blotting image from Control (n=5) and ischemic (n=3) retinas. Housekeeping gene β-Tubulin III was used as loading control. B) The graph shows the relative density

measurements of bands which were calculated by proportioning to loading control. One-hour permanent retinal ischemia increased relative RFC1 protein levels compared to control. C) Retinal ischemia significantly augmented the percentage of mean grey value of RFC1 by 114.5% (n=3). Representative confocal images of control or ischemic retinal microvessels stained by anti-RFC1 antibody (red). Significant immunosignal increase in ischemic microvessels was observed compared to controls. Also, ischemic microvessels demonstrated expected characteristics such as constrictions and protruding pericyte bodies (asterisk). Of note, ischemic pericyte bodies showed denser RFC1 immunosignal (white arrows). D, E) In contrast, retinal ischemia decreased collagen-4 by 42.07% (n=3), Occludin by 50.94% (n=3). F, G) In addition, microvessels treated with LV-RFC1, 10 days before ischemia showed 176% increase in Collagen-4 (n=3) and 60.9% in Occludin (n=3) immunosignal compared to Control-LV delivered ones indicating RFC1 overexpression before ischemia might retrieve loss of collagen-4 and occludin in ischemia. (*p $\leq$ 0.05). Nuclei were labelled with Hoechst 33258 (blue) in all images. Data are mean ± S.E.M. Mann-Whitney U; Scale bar: 10 μm.

Fig. 5. siRNA mediated RFC1 knockdown led to inner BRB breakdown and endogenous IgG extravasation. The wholemount retinas were labeled by Lectin (cyan) and incubated with Cy3 labeled goat anti-mouse antibody (yellow) to visualize the extravasated endogenous mouse IgG. A) RFC1-siRNA mediated knockdown led to significant IgG extravasation (red arrows) in microvessels while B) Control-siRNA treated retinas showed no extravascular IgG signal (p=0.001; n=3). C) 1 hr ischemia led to microvessel constrictions (white arrows) but no IgG extravasation. D) However, knocking down RFC1 before ischemia provoked inner BRB disruption leading to endogenous IgG extravasation (red arrows) (p=0.05; n=3). Nuclei were labelled with Hoechst 33258 (blue) in all images. Scale bar: 10 μm

Fig. 1

Reduced Folate Carrier 1 expression in retinal cells

Retinal Layers and RPE

NG2 (+) pericytes

Retinal microvesels

CD13 (+) pericytes

Endothelial cells

PDGFR-β (+) pericytes

Fig. 2

Effects of siRNA mediated RFC1 knockdown

RFC1 Expression after siRNA mediated knockdown

A

Scrambled-siRNA | RFC1-siRNA
n=1 n=2 n=3 n=4 | n=1 n=2 n=3 n=4

RFC1 — 43 kDA
β-Tubulin III — 55 kDA

B — Relative Band Density (RFC1/β-Tubulin III), * , Scrambled siRNA vs RFC1 siRNA

C — RFC1 mRNA Expression (Relative to β-tubulin 4A), * , Scrambled siRNA vs RFC1 siRNA

RFC1 Immunoreactivity after siRNA mediated knockdown

D — % RFC1 mean grey value (R.F.U), * , Scrambled siRNA vs RFC1 siRNA
RFC Hoechst — Scrambled siRNA
RFC1 Hoechst — RFC1 siRNA

NG2 Immunoreactivity after siRNA mediated knockdown

E — NG2 Hoecsth Scrambled siRNA ; NG2 Hoecsth RFC1 siRNA

F — Pericyte Number/Vessel Length (mm), Scrambled siRNA vs RFC1 siRNA

RFC1 and Lectin Immunoreactivity in wholemount retinas after siRNA mediated knockdown

G — RFC1 / Lectin / Merged (Scrambled siRNA) ; RFC1 / Lectin / Merged (RFC1 siRNA)

Collagen-4, Occludin, Claudin-5, and ZO-1 immunoreactivity in RFC1 knockdown

H — % Collagen-4 mean grey value (R.F.U), * , Scrambled siRNA vs RFC1 siRNA ; Col-4 Scrambled siRNA / Col-4 RFC1 siRNA

I — % Occludin mean grey value (R.F.U), * , Scrambled siRNA vs RFC1 siRNA ; Occludin Scrambled siRNA / Occludin RFC1 siRNA

J — % Claudin-5 mean grey value (R.F.U), * , Scrambled siRNA vs RFC1 siRNA ; Claudin-5 Scrambled siRNA / Claudin-5 RFC1 siRNA

K — % ZO-1 mean grey value (R.F.U), * , Scrambled siRNA vs RFC1 siRNA ; ZO-1 Scrambled siRNA / ZO-1 RFC1 siRNA

Fig. 3

Effects of LV mediated overexpression of RFC1 protein

RFC1 Expression after LV delivery

A

Control LV, Control LV, Control LV, RFC1 LV, RFC1 LV, RFC1 LV

RFC1 43kDA

β-Actin 42 kDA

Relative Band Density (RFC1/β-Actin)

Control-LV   RFC1-LV

B GFP Expression after LV delivery

GFP Hoechst   Control-LV

GFP Hoechst   Control-LV

RFC1 Immunoreactivity in wholemount retinas after LV delivery

C

RFC1   Control-LV

RFC1   RFC1-LV

D RFC1 Immunoreactivity after LV delivery

% RFC1 mean grey value (R.F.U)

Control-LV   RFC1-LV

RFC1 Hoechst   Control-LV

RFC1 Hoechst   RFC1-LV

Collagen-4, Occludin, Claudin-5, and ZO-1 Immunoreactivity in RFC1 Overexpression

E

%Collagen-4 mean grey value (R.F.U)

Control-LV   RFC1-LV

Col-4 Control-LV   Col-4 RFC1-LV

F

%Occludin mean grey value (R.F.U)

Control-LV   RFC1-LV

Occludin Control-LV   Occludin RFC1-LV

G

%Claudin-5 mean grey value (R.F.U)

Control-LV   RFC1-LV

Claudin-5 Control-LV   Claudin-5 RFC1-LV

H

%ZO-1 mean grey value (R.F.U)

ns

Control-LV   RFC1-LV

ZO-1 Control-LV   ZO-1 RFC1-LV

Fig. 4

The role of RFC1 protein in retinal ischemia

RFC1 Expression after retinal ischemia

RFC1 Immunoreactivity after retinal ischemia

Collagen-4 and Occludin Immunoreactivity after retinal ischemia

Collagen-4 and Occludin Immunoreactivity in ischemic retinas pre-treated with LV

Fig. 5

siRNA mediated RFC1 knockdown leads to disruption of inner BRB integrity