IMPA1 dependent regulation of plasma membrane phosphatidylinositol 4,5-bisphosphate turnover and calcium signalling by lithium

Sankhanil Saha, Harini Krishnan and Padinjat Raghu*

National Centre for Biological Sciences-TIFR

GKVK Campus, Bellary Road, Bangalore 560065, India

*corresponding author

E-mail: praghu@ncbs.res.in

Running Title- Lithium regulates PIP$_2$ and calcium signaling in human neurons

Abstract

Lithium (Li) is a widely used as a mood stabilizer in the clinical management of Bipolar Affective Disorder (BPAD). However, the molecular targets of Li in neural cells that underpin its therapeutic effect remain unresolved. Inositol monophosphatase (IMPA1), is an enzyme involved in the resynthesis of phosphatidylinositol 4,5-bisphosphate (PIP_2) following receptor-activated phospholipase C (PLC) signalling. *In vitro*, Li inhibits IMPA1, but the relevance of this inhibition within neural cells remains unknown. Here we report that in human cells, treatment with therapeutically relevant concentrations of Li reduces receptor activated calcium release from intracellular stores and also delays the resynthesis of PIP_2 following receptor activated PLC signalling. Both these effects of Li are abrogated in cells where IMPA1 has been deleted. We also observed that in human forebrain cortical neurons, treatment with Li results in reduced neuronal excitability as well as reduced calcium signals following receptor activated PLC signalling. Following Li treatment of human forebrain cortical neurons, transcriptome analyses reveal downregulation of multiple components of the glutamate receptor signalling system. Glutamate is a key excitatory neurotransmitter in the human brain and thus our findings provide an insight into the mechanisms underlying the dampening of neuronal excitability following Li treatment. Collectively, our findings suggest that Li inhibits receptor activated PLC signalling leading to an altered transcriptional response and reduced neuronal excitability.

Introduction

Bipolar affective disorder (BPAD) is a human psychiatric illness characterized by disruptive mood swings; the defining characteristics being the alteration between mania and depression [1] associated with an elevated rate of suicide among patients [2]. BPAD has been regarded by the World Health Organization (WHO) as one of the leading causes of disability worldwide [3]. Mood stabilizers like lithium (Li), Carbamazepine and Valproate remain the first-line mode of medication for bipolar disorder. Among these drugs, Li stands out as the go-to drug for treating acute manic phases and also for its preventive effect against relapses of manic episodes [4, 5]. Although Li remains the mainstay in BPAD treatment, individual patient responses towards Li arevariable- it remains ineffective for a large proportion (approximately 30%) of BPAD patients, the so-called 'Li non-responders', whereas another 30% of patients are only partially responsive to Li [6]. The reason for this variable response is unknown and remains a topic of intense interest.

Li responsiveness is determined clinically by functional improvement in brain function as determined using the Alda score following treatment [7]. Despite substantial research over several decades, the molecular and biochemical mechanism by which Li exerts its effects on human brain cells that are relevant to functional improvements remains unclear [8]. A large number of molecular targets for Li have been described [9]. These include Inositol monophosphatase (IMPase) [10], Glycogen synthasekinase-3β (GSK-3β) [11], Sodium myo-inositol co-transporter (SMIT1) [12] and Bisphosphate- 3'-nucleotidase (BPNT1) [13]. The two most well studied targets of Li are Inositol monophosphatase (IMPase) and Glycogen synthase kinase-3β (GSK-3β).

A long-standing hypothesis for the therapeutic effect of Li is its impact on inositol lipid turnover during receptor activated G-protein coupled signalling through inhibition of IMPase [10]. During GPCR signalling, many receptors activate Phospholipase C (PLC) leading to the hydrolysis of the signalling lipid Phosphatidylinositol 4,5-bisphosphate (PIP_2) generating inositol 1,4,5-trisphosphate (IP_3) and diacylglycerol (DAG) [11] ultimately leading to changes in intracellular calcium $[Ca^{2+}]_I$. Alterations in $[Ca^{2+}]_I$ have previously been reported in cell lines derived from BPAD patients [12]. PIP_2 is a low abundance membrane lipid and its resynthesis is required to sustain PLC signalling during high rates of PLC activity as is seen in the brain [13] during the activation of many GPCR required for neurotransmission such as the metabotropic glutamate receptor [14]. The resynthesis of PIP_2 occurs via a set of biochemical reactions referred to as the PIP_2 cycle [15]. The sequential dephosphorylation of IP_3 produced during PLC activity to generate inositol, is required for PIP_2 resynthesis (Fig

1A). IMPase is the enzyme required for the final step in the conversion of inositol 1-
phosphate to inositol. Biochemical analyses showed a decrease in myo-inositol
concentration in cerebral cortex of lithium-treated rats accompanied by an increase in the
concentration of inositol monophosphate suggesting that Li might inhibit IMPase [16, 17].
Subsequent studies showed that Li binds to IMPase and inhibits the enzyme in a non-
competitive manner [18]. These observations led Berridge to propose the "Inositol depletion
hypothesis" [19] which posits that during PLC signalling in the brain, the inhibition of IMPase
by Li restricts the supply of inositol for PIP_2 resynthesis leading to down regulation of
neurotransmitter receptor signalling, thus reducing excitability in the brain and the control of
mania.
119
The mammalian genome includes two genes encoding IMPase activity, IMPA1 and IMPA2.
However, IMPA1 is more abundant in the human brain [20, 21] and is inhibited by Li at
therapeutically relevant concentrations [22]. *In vivo* studies have also suggested a role of Li
in regulating IMPase function. Briefly, (i) in rodent models, Li treatment reduced brain inositol
levels and elevated Inositol monophosphate (IP_1), the substrate of IMPase in the cerebral
cortex of rat [23] (ii) treatment of human BPAD patients with Li led to a reduction in PIP_2
levels in platelet membranes [24–26]. However, there are also findings that do not support
this hypotheses (i) IMPA1 knockout mice showed only a modest reduction in brain inositol
levels [27] and the impact of IMPA1 knockout on receptor activated PIP_2 turnover remains
unknown, (ii) genome-wide association studies have not revealed polymorphisms in IMPA1
that are linked to BPAD or Li sensitivity [28][6].
131
In this study, we have tested the hypothesis that treatment of human cells with Li diminishes
calcium signalling. We find that this change is underpinned by reductions in G-protein
coupled PIP_2 turnover. These changes in calcium signalling were also observed in human
iPSC derived forebrain cortical neurons and accompanied by reduction in neuronal
excitability. Transcriptomic analysis revealed that Li treatment exerts its effect on neuronal
excitability through multiple mechanisms including but not restricted to GPCR activated
calcium signalling. Our findings open new approaches to predict Li responsiveness in BPAD
patients and develop novel strategies for the clinical management of this disorder.
140
141
142

Materials and Methods

DNA constructs: For PIP_2 measurement, PH-PLCδ::GFP probe was used which was a gift from Tamas Balla (Addgene plasmid # 51407). The cDNA for human muscarinic acetylcholine receptor m1AchR was obtained from cDNA Resource Center (MAR0100000) and was cloned via Gibson assembly into a lentiviral pHR transfer backbone (having Puromycin resistance cassette) for lentiviral generation. The Nir2 shRNA was obtained from GE Dharmacon (RHS4430-200186441: Clone id V2LHS_62836).

Mammalian cell culture, treatment and transfection: HEK293T cells were maintained under standard conditions at ($37^{o}C$, with 5% CO_2) in DMEM high Glucose media (Dulbecco's Modified Eagle Medium; Life Technologies), supplemented with 10% Foetal bovine serum. For Li treatment, the cells were grown and maintained in this media, supplemented with 1 mM LiCl (Sigma) for 5 days.The cells were tested for mycoplasma, prior to the experiments. MycoAlert™ (Lonza) was used per the manufacturer's protocol to test the spent media for mycoplasma contamination. Transfection with the plasmid expressing PH-PLCδ::GFP probe was done 24 hours before the imaging experiments using Polyethylenimine solution (PEI; Polysciences Inc.; 1 mg/ml) when the cells were around 70% confluent.

Lentiviral transduction to generate HEK293T cell line stably expressing muscarinic acetylcholine receptor m1AchR: Recombinant Lentiviral particles were packaged in HEK293T cells grown in DMEM high Glucose media, supplemented with 10% FBS. The HEK293T cells cultured in the 6 well plates were transfected with a solution consisting of 1.2 µg of pHR-IRES plasmid encoding muscarinic acetylcholine receptor *CHRM1* cDNA, along with 1.2 µg of each of the helper plasmids (psPAX and pMD2.G) in 500 µl of Opti-MEM (Invitrogen) and 110 µl Polyethylenimine (PEI, 1 mg/ ml). Medium was changed after 12 hours and the virus was harvested at 72 h after transfection. Then HEK293T cells were transduced with these recombinant viral particles and selected via Puromycin treatment (1.5 µg/ ml) for 1 week- to obtain the HEK293T cell line, expressing m1AchR. Western Blot confirming the expression of m1AchR is shown in supplementary figures (Supp. Fig 3C,D).

Generation of Neural Stem Cells and differentiation and maintenance of neuronal culture: Neural Stem Cells (NSCs) were generated as previously described [61], with slight modifications. Human iPSCs were differentiated to embryoid bodies in E6 medium (Gibco). After that, primary neural rosettes that were formed, were manually picked, passaged and eventually plated as NSC monolayer in Neural Expansion Medium (NEM; DMEM/F12 with 2mM Glutamax, supplemented with N2, B27 without Vitamin A, 20 ng/ml bFGF, 100 µM non-

essential amino acids, 2 µg/ml Heparin Sulphate, 100 U/ml penicillin-streptomycin) (all components procured from Life Technologies). The NSCs were subjected to CD133+ selection as previously described via fluorescence-activated cell sorting (FACS) [62] to eliminate cells of other lineages. The sorted NSCs were further characterized by immunofluorescence for the specific markers. Neural Stem Cells (NSCs) were grown on Matrigel coated tissue dishes in NEM at 37°C and 5% CO_2.

For differentiation, NSCs were seeded onto Matrigel- Growth Factor Reduced (Corning) coated cover-slip dishes or 35 mm dishes and expanded in NEM till the cells reached 70% confluency. NEM was then withdrawn and replaced with Neural Differentiation Medium (NDM; DMEM/F12 with 2mM Glutamax, supplemented with N2, B27 containing Vitamin A, 100 µM non-essential amino acids, 100 U/ml penicillin-streptomycin, 10 ng/ml BDNF, 10 ng/ml GDNF, 10 ng/ml IGF-1, 50 µg/ml Ascorbic acid (Sigma) and 2 µM DAPT (Sigma)). DAPT was supplemented for the first 14 days. Media was changed every 2 days during the differentiation process. All the experiments were done 45 DIV post differentiation (The day of addition of NDM was considered to be DIV 0). For chronic Li treatment, the neuronal cultures were maintained in NDM supplemented with 1 mM LiCl for the last 10 days (DIV35 to DIV45).

Live cell imaging for the PIP$_2$ probe: To analyse the dynamics of PIP_2 at the plasma membrane by following the change in translocation of the PH-PLCδ::GFP probe, live cell imaging was performed as stated in *Varnai et al* [31], with minor modifications. 20000 cells were seeded in Poly-L Ornithine (PLO) coated cover-slip dishes and grown for 96 hours with or without Li. 24 hours prior to imaging, the cells were transfected with PH-PLCδ::GFP plasmid using Lipofectamine 3000 as per manufacturer's protocols. Carbamoylcholine chloride mediated changes in the membrane localization of PH-PLCδ::GFP probe was examined by a time-lapse method using an Olympus FV 3000 confocal microscope. The cells were kept in pre-warmed Kreb's Ringer Buffer (120 mM NaCl, 4.7mM KCl, 1.2 mM $CaCl_2$, 0.7 mM $MgSO_4$, 10 mM Glucose, 10 mM HEPES; pH-7.4) and at 37°C while imaging. Changes in the PIP_2 membrane localization were quantified by taking a fluorescence intensity ratio of the plasma membrane (F_{PM}) and the cytosol (F_{Cyt}) across the time-lapse and normalizing it to the first value.

Calcium Imaging for Carbamoylcholine stimulation: For studying the calcium physiology, HEK293T-M1 cells were seeded in Poly-L Ornithine (PLO) coated cover-slip dishes while iPSC derived cortical neurons (DIV45) were grown on Matrigel coated cover-slip dishes.

Before imaging, the cells were washed twice with Hank's Buffered Salt Solution (HBSS) (10
mM HEPES, 118 mM NaCl, 4.96 mM KCl, 1.18 mM MgSO$_4$, 1.18 mM KH$_2$PO$_4$, 10 mM
Glucose, 2 mM CaCl$_2$; pH-7.4) and then loaded with the ratiometric Ca^{2+} indicator Fura-2 AM
(4 µM) (Acetoxymethylester, Invitrogen) along with 0.002% Pluronic F-127 for 30 minutes.
The excess dye was removed by washing the cells thrice with HBSS and cells were
incubated for an additional 20 minutes to equilibrate the intracellular dye concentration and
allow de-esterification. Imaging was performed for 10 min using a 40X objective (N.A. 0.75)
of wide-field fluorescence microscope Olympus IX-83 for HEK293T cells, while a 20X
objective (N.A. 0.50) was used for neuronal cultures. Free- and Ca^{2+}-bound Fura-2
fluorescence intensities (excitation 340/380 nm, emission 510 nm) were recorded at every 5
seconds intervals using an EM-CCD camera (Evolve 512 Delta, Photometrics). Basal
cytosolic Ca^{2+} was recorded in HBSS buffer for the initial 18 frames. Carbamoylcholine
chloride (20 µM and 50 µM respectively for HEK and neuronal cultures, as determined in
preliminary concentration–response experiments) (Sigma Aldrich-C4382) was added to the
cells for activating PLC and thereby, inducing Ca^{2+} release from the endoplasmic reticulum
store. After recording for 42 frames, 20 µM Ionomycin (Calbiochem) was added and
recorded for 24 frames for recording R$_{max}$. 24 more frames were recorded, post addition of 4
mM EGTA (containing 0.01% Triton-X-100) for chelation of Ca^{2+} ions and determination of
R$_{min}$.

Calcium Imaging for SOCE (Store Operated Calcium Entry): For the SOCE experiment,
the cells were kept in 'zero Ca^{2+} HBSS' (10 mM HEPES, 118 mM NaCl, 4.96 mM KCl, 1.18
mM MgSO$_4$, 1.18 mM KH$_2$PO$_4$, 10 mM Glucose; pH-7.4) just before the imaging. After
recording the initial 12 frames for basal cytosolic Ca^{2+}, an optimal concentration of
Thapsigargin (10 µM) (Invitrogen) was added for the store depletion and 60 frames were
recorded. Then, 2 mM CaCl$_2$ was then added to the extracellular buffer to induce SOCE- 60
frames were recorded post induction of SOCE. Subsequently, 20 µM Ionomycin
(Calbiochem) was added and recorded for 24 frames; then 4 mM EGTA (containing 0.01%
Triton-X-100) was added and 24 frames were recorded. For studying PLC activated calcium
mobilization from stores and SOCE, Carbamoylcholine chloride (20 µM and 50 µM
respectively for HEK and neuronal cultures) was added instead of Thapsigargin.

Analysis of Calcium recordings: The CellSens Olympus software was used for the
analysis. A dynamic region of interest (ROI) was drawn for each cell which exhibited rise in
[Ca^{2+}]$_I$ (as seen by increase in fluorescence post-stimulation) to track the fluorescence
changes over time. The emission intensities corresponding to 340 nm and 380 nm

excitations were used to calculate the 340/380 ratio for each cell across all time points. The

$[Ca^{2+}]_I$ was estimated using the Grynkiewicz equation [63] as follows:

$[Ca^{2+}]_I$ (nM) = K_d X SF X (R - R_{min})/ (R_{max}- R)

where R_{min} and R_{max} refer to the minimum and maximum 340/380 ratio respectively. 224 nM

was taken as K_d for Fura-2 AM in human cells and SF (scaling factor) was calculated for

each cell by dividing the fluorescence emission intensity at Ca^{2+} free form with the

fluorescence emission intensity at Ca^{2+} bound form of the dye after excitation at 380 nm. The

final Ca^{2+} traces were plotted using GraphPad Prism 9.0.

Generation of the IMPA1 deletion in cell lines: For generation of HEK293T-IMPA1$^{-/-}$ cell

line, the initial region of the coding sequence (CDS) (exon1) of the IMPA1 gene was targeted

by a pair of guide RNAs (sgRNAs) and deleted via CRISPR-Cas9 technology, then these

cells were transduced to stably express Muscarinic receptor M1. The sgRNAs targeting the

start of the CDS of IMPA1 gene, were designed via the crispr.mit.edu. The sequences of the

sgRNAs are as follows:

5' GGAGAACCACCTTGTTGGCCAAGCTGGAATGTAGTGGCGTTTTAGAGCTAGAAATAGCAAGTT 3'

5' GGAGAACCACCTTGTTGGGTAATATGGTACAGACACACGTTTTAGAGCTAGAAATAGCAAGTT 3'

These sgRNAs were cloned individually into humanized pgRNA plasmids (pgRNA-

humanized was a gift from Stanley Qi (Addgene plasmid # 44248). The cells were co-

transfected with the three plasmids (both the pgRNA humanized plasmids encoding the

sgRNAs and pLentiCas9 blast encoding for the Cas9 protein endonuclease from

Streptococcus pyogenes) and selected with Puromycin (2 µg/ ml) for 48 hours. Then the

genomic DNA was isolated and a Surveyor PCR was performed to check the occurrence of

the deletion. Once confirmed that the sgRNAs were able to target the exon 1 of IMPA1 gene,

the transfected cells were seeded in 96 wells as single cells and gradually expanded. After

selection via Surveyor PCR using Surveyor primers (primers flanking the region ought to be

deleted), a HEK293T cell line was selected which had showed only the shorter amplicon

(amplicon after deletion) in the Surveyor PCR. This HEK293T cell line was then expanded

and the expression of IMPA1 was checked by Western blotting using the anti-IMPA1

antibody. No corresponding bands for IMPA1 were observed, indicating that IMPA1 was

knocked out in this cell line. This HEK293T-IMPA1-/- cell line did not show any

morphological difference with respect to HEK293T cells and did not require inositol

supplementation in the media for maintenance.

Western Blotting: For estimating the expression of the IMPA1, cells were suspended in

appropriate volume of lysis buffer and kept on ice for lysis; 10 µl of it was used for Bradford

protein assay (BioRad) for quantification. Post quantification, equal amount of samples were

heated at 95°C with Laemli loading buffer for 5 min, then loaded and separated on 10%

SDS-PAGE gel. For estimating the expression of the muscarinic acetylcholine receptor

m1AchR, cells were pelleted and directly lysed by adding Laemmli's loading buffer and

repeated syringing with insulin syringes (31 gauge). Then the samples were heated at 95°C

for 5 min and loaded onto a pre-cast Polyacrylamide 4-12% gradient gel (Bolt, Invitrogen).

The proteins were then transferred onto a nitrocellulose membrane and then blocked with

10% Blotto (Santa Cruz Biotechnology) diluted in Tris Buffer Saline containing 0.1% Tween-

20 (0.1% TBST) for 40 minutes. Subsequently the membrane was incubated with the rabbit

anti-IMPA1 antibody (1:10000 diluted in 5% BSA in 0.1% TBST) (Abcam ab184165) or goat

anti-CHRM1 antibody (1 µg/ ml, diluted in 5% BSA in 0.1% TBST) (Abcam ab77098)

overnight at 4°C. For loading control, the blot was incubated with mouse anti-β-Tubulin

antibody (1:4000 diluted in 5% BSA in 0.1% TBST) (DSHB Hybridoma Product E7). The

blots were then washed thrice with 0.1% TBST and then incubated with the appropriate

HRP-conjugated secondary antibodies (1:10000 dilutions; Jackson Laboratories, Inc.) for 1.5

hrs. Post incubation with the secondary antibodies, the blot was washed thrice with 0.1%

TBST and developed using Clarity Western ECL substrate (Biorad) on a GEImageQuant

LAS 4000 system.The ImageJ software was used for the densitometric analysis of the blots.

Firstly, the background intensities were subtracted from the images using an average of

mean background intensities of a few ROIs adjacent to the band of interest. Then ROIs were

drawn around the bands of interest and the integrated density of each ROI was extracted

from the image. Finally, the ratio of the integrated density of the IMPase to that of the loading

control (β-Tubulin) was calculated.

Calcium imaging for measuring transients: For studying the effect of Li on the excitability

of cortical neurons, spontaneous calcium events (calcium transients) were measured in

D149 cortical neuronal cultures at 45 DIV for both untreated and Li treated neurons,

according to previously published lab protocol [44]. The neuronal cultures were washed

twice for 10 minutes with HBSS and then loaded with 4 µM of Fluo-4 AM (Molecular probes,

F14201) in HBSS having 0.002% Pluronic F-127 for 30 min at room temperature. Imaging

was performed for a time-span of 10 min using a 20X objective (N.A. 0.50) with a time

interval of 1 second at 488 nm illumination. The baseline measurement was recorded for a

period of 4 min to visualize the calcium transients. Then 10 µM Tetrodotoxin (TTX, Hellobio)

was added to abolish calcium transients and recorded for another 4 min. A dynamic region

of interest (ROI) was manually drawn around each neuronal soma; individual Ca^{2+} traces

were then obtained using the CellSens Dimensions software. The fluorescence intensity

value from each neuronal soma was normalized to the first baseline signal and the individual

traces were plotted using GraphPad Prism 9.0. The frequency of calcium transients was measured manually from first-derivative filter traces.

Post-hoc immunofluorescence for neuronal characterization: Post-calcium imaging, the neuronal cultures were washed thoroughly a few times with HBSS buffer and then fixed with 4% PFA dissolved in PBS for 20 mins. Post-fixation, the cultures were washed twice with PBS for 5 min each and then permeabilized with 0.1% Triton-X (dissolved in PBS). The cultures were then blocked with 5% BSA diluted in PBS, for 1 hr at room temperature. Subsequently the neurons were incubated with the Primary antibodies diluted in 5% BSA in PBS overnight at 4°C. Next day, the cultures were subjected to three washes with PBS (10 minutes each) to remove non-specifically bound primary antibody. Respective secondary antibodies at a dilution of 1:300 in 5% BSA in PBS were used for incubation for 2 hours. DAPI was used as a nuclear marker at a dilution of 1:1000.

Primary antibody	Source	Company	Catalogue number	Dilution
SOX1	Rabbit	Abcam	ab87775	1:400
Nestin	Mouse	Thermo Fisher	MA1-110	1:100
SOX2	Mouse	Thermo Fisher	MA1-014	1:500
Pax6	Rabbit	Thermo Fisher	MA1-109	1:100
GFAP	Rabbit	Abcam	ab7260	1:1000
MAP2	Chicken	Novus Biologicals	NB300-213	1:1000
CTIP2	Rat	Abcam	ab18465	1:800
Synapsin	Rabbit	Abcam	ab64581	1:200

Secondary antibody	Company	Catalogue number	Dilution
Alexa 488- Goat α-Rabbit IgG	Life Technology	A11034	1:300
Alexa 568- Goat α-Chicken IgG	Life Technology	A11041	1:300
Alexa 488-Goat α-Rat IgG	Life Technology	A11006	1:300
Alexa 568- Goat α-Mouse IgG	Life Technology	A11004	1:300

RNA isolation, cDNA synthesis and Real-Time Quantitative PCR: The total RNA was extracted from HEK293T cells or mature neuronal cultures, in multiple biological replicates

using TRIzol (Ambion, Life Technologies) following manufacturer's protocol and thereby, quantified via Nanodrop 1000 spectrophotometer (Thermo Fisher Scientific). For cDNA synthesis, 1 μg of the RNA from each replicate was treated with DNase I (amplification grade, Thermo Fisher Scientific) and then incubated with Superscript II reverse transcriptase (Invitrogen) along with random hexamers and dNTPs. A no reverse transcription control sample was also included for each sample. Real-Time Quantitative PCR analysis was performed on an Applied Biosystems 7500 fast qPCR system using diluted cDNA samples and primers against genes of interest and GAPDH as internal controls. The C_t values obtained for different genes were normalized to those of GAPDH from the same sample. The relative expression levels were calculated using ΔC_t method, whereas the fold change was calculated using $\Delta\Delta C_t$ method. The primers used for qRT-PCR are listed in the Table 2.

LCMS based PIP and PIP$_2$ measurements: To validate whether the PH-PLCδ-GFP probe based imaging of PIP$_2$ in the HEK293T cells reflected the changes in PIP$_2$ level, mass spectrometry measurements of PIP$_2$ from the cells was carried out following existing lab protocols[64]. Reverse phase liquid chromatography was coupled with high sensitivity mass spectrometry (LCMS) and a Multiple Reaction Monitoring (MRM) method was employed to detect PIP$_2$ levels from the whole cells.

Lipid extraction—Equal number of cells from a single well of a 12 well plate (counted by a haemocytometer while seeding) were pelleted and then gently resuspended into 170 μl of 1X PBS in a 2 ml low-bind polypropylene centrifuge tube. To this, 750 μl of ice-cold quench mixture (MeOH/CHCl$_3$/1M HCl in the volumetric ratio of 484/242/23.55), followed by 15 μl of a pre-mixed ISD (Internal Standard) mixture containing 25 ng of 37:4 PIP [**PI4P 17:0/20:4 (LM-1901)**], 25 ng of 37:4 PIP$_2$ [**PI 4,5P$_2$ 17:0/20:4 (LM-1904)**] and 50 ng of 31:1 Phosphatidylethanolamine [**PE 17:0/14:1 (LM-1104)**] was added (All the lipid standards were procured from Avanti Lipids). For Carbamoylcholine chloride mediated PLC activation, cells were suspended in 85 μl of 1XPBS and then 85 μl of 20 μM Carbamoylcholine chloride (made in PBS) (final concentration – 10 μM) was added for 1 min at 37°C in shaking conditions. For the regeneration of PIP$_2$ post PLC mediated hydrolysis, 1 ml of ice-cold PBS was added to dilute the agonist Carbamoylcholine chloride and stop the stimulation. Cells were pelleted and washed and then resuspended in pre-warmed buffer (37°C) and kept for 15 mins for recovery (time given for regeneration of PIP$_2$). Post 15 mins of incubation, cells were pelleted and then resuspended in 170 μl of PBS, to which 750 μl of quench mixture and 15 μl of a pre-mixed ISD were added as stated previously. The mixture was vortexed and 725 μl of CHCl$_3$ and 170 μl of 2.4 M HCl were added. After vortexing again for 2 min at 1500 rpm, the phases were separated by centrifugation for 3 min at 1500 g. The lower

organic phase was then collected and added to fresh tubes containing 708 μl of Lower Phase Wash Solution (LPWS; MeOH/1M HCl/CHCl$_3$ in the volumetric ratio of 235/245/15). The tubes were then vortexed and spun at 1500 rpm again for 3 min to separate the phases. The resultant lower organic phase was collected into a fresh tube and subjected to the Derivatization reaction.

Derivatization of Lipids— 50 μl of 2M TMS-Diazomethane was added to the collected organic phase (TMS-Diazomethane is toxic and is to be used with utmost precautions and following safety guidelines). The reaction was allowed for 10 min at 600 rpm at room temperature, post which 10 μl of Glacial acetic acid was added to quench the reaction. 500 μl of Post Derivatization Wash Solution (PDWS- CHCl$_3$/MeOH/H$_2$O in the volumetric ratio of 24/12/9; the upper phase was used) was then added to the sample, vortexed and spun down at 1500 g for 3 min. The upper aqueous phase was discarded and the wash step was repeated. To the final organic phase, 45 μl MeOH and 5 μl H$_2$O was added, mixed and spun down. The samples were then dried in a SpeedVac at 400 rpm for 150 mins under vacuum and 90 μl of MeOH was added to reconstitute the sample to a final volume of about 100μl.

Chromatography and Mass spectrometry—The chromatographic separation was performed on an Acquity UPLC BEH300 C4 column (100 X 1.0 mm; 1.7 μm particle size - Waters Corporation, USA) using a Waters Aquity UPLC system connected to an ABSCIEX 6500 QTRAP mass spectrometer for ion detection. All the samples were injected in duplicates and the flow rate was set to 100 μl /min. Solvent gradients were set, starting from 55% Solvent A (Water + 0.1% Formic Acid)- 45% Solvent B (Acetonitrile + 0.1% Formic acid) from 0-5 min; then 45% B to 100% B from 5-10 min; 100% B was held from 10-15 min; between 15-16 min, 100% B was lowered to 45% B and held there till 20th min to re-equilibrate the column. On the mass spectrometer, Neutral Loss Scans was employed during pilot standardization experiments on biological samples to identify the parent ions that would lose neutral fragments corresponding to 490 a.m.u and 382 a.m.u- these were indicative of PIP$_2$ and PIP species respectively and likewise 155 a.m.u for PE species [64]. Thereafter, we quantified PIP, PIP$_2$ and PE species in biological samples using the selective Multiple Reaction Monitoring (MRM) method in the positive ion mode. Area under the peaks was calculated via the Sciex MultiQuant software. For each run, area under the peak for each species of PIP$_2$, PIP and PE was normalized to PIP$_2$, PIP and PE internal standard peak respectively. Thereafter, the sum of normalized areas for all the species of PIP$_2$ or PIP was taken and divided by the sum of normalized areas for all the species of PE in each of the biological samples to account for differences in total phospholipid extracted across samples. The MRM

415 mass pairs used for PIP, PIP$_2$ and PE species identification and quantification are listed
416 below:

Q1 Mass (Da)	Q3 Mass (Da)	Dwell time (msec)	ID	DP (volts)	EP (volts)	CE (volts)	CXP (volts)
1041.5	551.5	35	32_0 PIP2	60	11	37	15
1039.5	549.5	35	32_1 PIP2	60	11	37	15
1069.5	579.5	35	34_0 PIP2	60	11	37	15
1067.5	577.5	35	34_1 PIP2	60	11	37	15
1065.5	575.5	35	34_2 PIP2	60	11	37	15
1103.5	613.5	35	37_4 PIP2	60	11	37	15
995.5	613.5	35	37_4 PIP	140	12	29	12
1097.5	607.5	35	36_0 PIP2	60	11	37	15
1095.5	605.5	35	36_1 PIP2	60	11	37	15
1093.5	603.5	35	36_2 PIP2	60	11	37	15
1091.5	601.5	35	36_3 PIP2	60	11	37	15
1089.5	599.5	35	36_4 PIP2	60	11	37	15
1125.5	635.5	35	38_0 PIP2	60	11	37	15
1123.5	633.5	35	38_1 PIP2	60	11	37	15
1121.5	631.5	35	38_2 PIP2	60	11	37	15
1119.5	629.5	35	38_3 PIP2	60	11	37	15
1117.5	627.5	35	38_4 PIP2	60	11	37	15
1115.5	625.5	35	38_5 PIP2	60	11	37	15
933.5	551.5	35	32_0 PIP	140	12	29	12
931.5	549.5	35	32_1 PIP	140	12	29	12
961.5	579.5	35	34_0 PIP	140	12	29	12
959.5	577.5	35	34_1 PIP	140	12	29	12
957.5	575.5	35	34_2 PIP	140	12	29	12
989.5	607.5	35	36_0 PIP	140	12	29	12
987.5	605.5	35	36_1 PIP	140	12	29	12
985.5	603.5	35	36_2 PIP	140	12	29	12
983.5	601.5	35	36_3 PIP	140	12	29	12
981.5	599.5	35	36_4 PIP	140	12	29	12
1017.5	635.5	35	38_0 PIP	140	12	29	12
1015.5	633.5	35	38_1 PIP	140	12	29	12

1013.5	631.5	35	38_2 PIP	140	12	29	12
1011.5	629.5	35	38_3 PIP	140	12	29	12
1009.5	627.5	35	38_4 PIP	140	12	29	12
1007.5	625.5	35	38_5 PIP	140	12	29	12
706.5	551.5	35	PE 32_0	140	12	29	12
704.5	549.5	35	PE 32_1	140	12	29	12
734.5	579.5	35	PE 34_0	140	12	29	12
732.5	577.5	35	PE 34_1	140	12	29	12
730.5	575.5	35	PE 34_2	140	12	29	12
762.5	607.5	35	PE 36_0	140	12	29	12
760.5	605.5	35	PE 36_1	140	12	29	12
758.5	603.5	35	PE 36_2	140	12	29	12
756.5	601.5	35	PE 36_3	140	12	29	12
754.5	599.5	35	PE 36_4	140	12	29	12
790.5	635.5	35	PE 38_0	140	12	29	12
788.5	633.5	35	PE 38_1	140	12	29	12
786.5	631.5	35	PE 38_2	140	12	29	12
784.5	629.5	35	PE 38_3	140	12	29	12
782.5	627.5	35	PE 38_4	140	12	29	12
780.5	625.5	35	PE 38_5	140	12	29	12
690.5	535.5	65	PE(31:1)	140	12	29	12

The other mass spectrometer parameters were:

ESI voltage: +5100-5200 V; Source Temperature: 300 C, Curtain Gas (CUR): 35-37, Ion

Source Gas 1 (GS1):15-20, Ion Source Gas 2 (GS2): 15-20.

RNA sequencing and transcriptomic analysis: RNA extracted from mature neurons (45DIV with and without Li treatment) and NSCs (wild-type, Li treated wild-type, IMPA knock-out and Li treated IMPA knock-out) as mentioned above was submitted for paired-end sequencing. Quality check and sequencing was carried out at the NCBS Sanger sequencing facility using Illumina Hiseq 2500 system. Single stranded RNA sequencing libraries were prepared for sequencing using NEBNext® Ultra™ II Directional RNA Library Prep with Sample Purification Beads (Catalog no-E7765L). Paired-end reads, (125 base pair for each neuronal sample and 100 base pair for each NSC samples) was obtained after sequencing. Post adapter trimming, FastQC was used to check the quality of reads. The reads were then aligned to human genome (GRCh38) using hisat2-build and hisat2-align module. The module for single strand sequencing was used in the Hisat-2 protocol. The output generated

436 in the bam format, was further used to obtain the counts of each of the individual gene using
437 HT-Seq pipeline. The differential expression of genes was then calculated using DeSeq2.
438 GO annotation analysis was done using the DAVID tool.
439
Sampling and statistical analysis: Each experiment was performed unblinded on different
biological groups with multiple biological replicates. No statistical analysis was done a priori
to determine the sample sizes. For the calcium imaging experiments, quantification was from
a minimum of 100 cells for all genotypes from multiple dishes of each type. For the PIP_2
probe experiments, approximately 50 cells were taken for the quantitative analysis.
Statistical significance was computed using Student's unpaired t-test to check for differences
in means between samples of different genotypes. Two-way ANOVA with post hoc Tukey's
pairwise comparison was used for the grouped analysis. All statistical analyses were
performed on Graph Pad Prism (version 9) and schematic representations were created with
biorender.com.
450

451 Results

452 An *in vivo* model system to study the impact of Li on PLC induced PIP_2 turnover

In order to study the turnover of PIP_2 following receptor activated Phospholipase C (PLC)
turnover, we created a cell line which heterologously expressed two components. To
stimulate high levels of PLC activity, we expressed the human Muscarinic acetylcholine
receptor (m1AchR) encoded by the gene *CHRM1*; previous studies have shown that
activation of heterologously expressed m1AchR results in high rates of PIP_2 hydrolysis
leading to IP_3 production [29, 30]. To monitor the levels of PIP_2 at the plasma membrane we
expressed the PH-domain of PLC-δ fused to green fluorescent protein (PH-PLCδ::GFP).
This probe binds to PIP_2 and in imaging experiments; its relative distribution between the
plasma membrane and the cytosol has been used by us and others to monitor plasma
membrane PIP_2 levels [31, 32]. A cell line expressing both m1AchR and PH-PLCδ::GFP can
therefore be used to stimulate PLC and monitor the turnover of PIP_2 at the plasma
membrane. We generated stable HEK293T cell lines expressing m1AchR (control-M1) and
PH-PLCδ::GFP was transiently expressed in those cells (Fig 1B); when stimulated with the
m1AchR agonist Carbamoyl choline (carbachol: Cch), the levels of PH-PLCδ::GFP at the
plasma membrane drop following which they gradually recover back towards pre-stimulation
levels (Fig 1C). The changes in plasma membrane PIP_2 was quantified as the ratio of
fluorescence intensity of the PH-PLCδ::GFP probe at the plasma membrane to its
fluorescence intensity in the underlying cytosol in the same part of the cell (Fig 1C). As a

positive control, for our assay to monitor PIP_2 regeneration following PLC activation, we
used shRNA to deplete Nir2; Nir2 (PITPNM1/RdgBαI) encodes for a phosphatidylinositol
transfer protein required for the resynthesis of PIP_2 following PLC stimulation [33–35]. Under
our experimental conditions, we found that depletion of Nir2 in our HEK293T reporter line
resulted in a decreased rate in the recovery of plasma membrane PIP_2 levels following
stimulation with carbachol (Fig 1D,E).

To measure the effect of Li treatment on PIP_2 resynthesis, we incubated our reporter line
with 1 mM Li; this concentration of LiCl was used in our experiments since the therapeutic
range for Li medication in BD patients was 0.75 to1.5 mEq/L [36]. Following Li treatment, we
stimulated the cells with carbachol and compared the dynamics of plasma membrane PIP_2
levels with that in non-Li treated cells. We found that in Li treated cells the rate with which
PIP_2 levels at the plasma membrane recovered to pre-stimulation levels was significantly
slower than in untreated controls (Fig 1 F,G). These findings strongly suggest that treatment
with Li results in a reduction in the rate of PIP_2 resynthesis following PLC stimulation.

In an alternative approach, we quantified total cellular PIP_2 mass using liquid
chromatography coupled with tandem mass spectrometry (LC-MS/MS) at three distinct time
points- corresponding to the basal state, one minute post-stimulation of PLC via carbachol
addition (hydrolysis of PIP_2) state and post-stimulation (recovery of PIP_2) state. From these
experiments, the time for recovery of PIP_2 (to pre-stimulation levels) was standardized at 15
minutes post stimulation with carbachol (Fig 1H). Using these conditions, we compared the
total PIP_2 mass between untreated cells and those treated with Li. We found that the mass
of PIP_2 in Li treated cells during the recovery phase was lower in Li treated cells compared
to controls (Fig 1I); likewise, the level of phosphatidylinositol phosphate (PIP) in Li treated
cells was also lower during the recovery phase (Fig 1J). Thus, Li treatment lowers the
recovery of PIP and PIP_2 levels following PLC activation.

Chronic Li treatment reduces PLC dependent intracellular Ca^{2+} mobilization
A key outcome of agonist triggered PIP_2 hydrolysis by PLC activation is intracellular Ca^{2+}
signalling [11]. Since we noted a reduction in the turnover of PIP_2 following Li treatment, we
tested the impact of Li treatment on intracellular Ca^{2+} levels $[Ca^{2+}]_I$. Using the control-M1 cell
line, we monitored $[Ca^{2+}]_I$ under both resting conditions as well as stimulation with carbachol,
an agonist of m1AchR. Cells were incubated in 1 mM Li prior to the experiment: basal $[Ca^{2+}]_I$
were no different between treated and untreated cells (Fig 2A). Stimulation of untreated cells
with carbachol evoked a rise in $[Ca^{2+}]_I$ which then decayed back to resting levels (Fig 2B). In

cells treated with Li, the carbachol induced rise in $[Ca^{2+}]_I$ was reduced compared to that seen

in untreated cells (Fig 2C). Receptor activated PLC mediated Ca^{2+} signalling is a biphasic

process with two components, and an initial release of Ca^{2+} from intracellular stores followed

by store-operated influx of Ca^{2+} into the cell from the extracellular medium [37]. We

performed experiments to separately monitor each of these components. During the initial

phase of the assay, carbachol stimulation was performed in the presence of zero

extracellular Ca^{2+}, thus selectively monitoring release of Ca^{2+} from intracellular stores (Fig

2D,E). Following this, the extracellular solution was supplemented with Ca^{2+} and the influx of

Ca^{2+} into the cytoplasm was monitored (Fig 2D,F). This analysis revealed that in Li treated

cells both intracellular Ca^{2+} release (Fig 2E) as well as the subsequent store operated Ca^{2+}

influx into cells was reduced (Fig 2F) compared to Li untreated controls. This reduction in

receptor activated release of Ca^{2+} from intracellular stores was not due to a reduction in the

size of intracellular Ca^{2+} stores as emptying of stores by the application of thapsigargin

resulted in equivalent rises in $[Ca^{2+}]_I$ in Li treated cells and untreated controls (Fig 2G,H).

According to the "inositol depletion hypothesis" [19], Li can restrict the available cytoplasmic

inositol pool available for phosphatidylinositol (PI) (and thence PIP_2) resynthesis by inhibiting

IMPase- thereby, lowering down the synthesis of the pool of PIP_2 that is required for PLC

mediated generation of DAG and IP_3. This cytoplasmic inositol pool can also be replenished

by transport of inositol from the extracellular medium via plasma membrane inositol

transporters (summarized in [38]). This implies that if a deficit in the available pool of

cytosolic inositol is responsible for the attenuated calcium signalling in Li treated cells, this

might be rescued by supplementation of the extracellular medium with inositol. We grew

HEK293T cells in an inositol rich DMEM media (inositol concentration is ca. 28 mg/litre;

which is similar to the inositol concentration in the cerebrospinal fluid [39–41]). These cells

were subjected to Li treatment and agonist induced Ca^{2+} influx was quantified. As controls in

this experiment, we used Li treated control-M1cells grown in DMEM not supplemented with

inositol. As in previous experiments, we found that when cells were grown in DMEM, Li

treatment resulted in a reduction of agonist induced Ca^{2+} influx (Fig 2I,J). However, when Li

treatment was done on cells grown in inositol supplemented DMEM, the Li induced reduction

in Ca^{2+} influx was rescued compared to that seen in cells grown under equivalent Li

treatment but in DMEM not supplemented with inositol (Fig 2I,J), compared to Li non-treated

cells.

IMPA1 is required for the effect of Li on PIP$_2$ resynthesis and agonist mediated calcium signalling

IMPase, the enzyme that dephosphorylates inositol monophosphate to generate myo-inositol has been proposed as a direct target of the action of Li, inhibiting its activity [42, 43]. There are two genes encoding IMPase in the human genome *IMPA1* and *IMPA2* of which *IMPA1* is the predominant isoform expressed in HEK293T cells. To test the requirement of *IMPA1*, we generated a HEK293T cell line in which *IMPA1* was deleted by CRISPR/Cas9 genome editing (Supp. Fig 3A); m1AchR was expressed in these cells by lentiviral transduction; these cells will be denoted as IMPA1$^{-/-}$-M1. Western blot analysis of the IMPA1$^{-/-}$ line showed the complete absence of the IMPA1 protein (Fig 3A), IMPA2 levels were not altered (Supp. Fig 3B) and the levels of m1AchR protein was comparable between control-M1 cells and IMPA1$^{-/-}$-M1 cells (Supp. Fig 3C,D). We tested the effect of Li treatment on IMPA1$^{-/-}$-M1 compared to untreated IMPA1$^{-/-}$-M1 cells. This analysis revealed that the rate of regeneration of PIP$_2$ at the plasma membrane was unchanged in IMPA1$^{-/-}$-M1 cells subjected to Li treatment, compared to untreated IMPA1$^{-/-}$-M1 cells (Fig 3B). In addition, the rate of PIP$_2$ regeneration was much slower in IMPA1$^{-/-}$-M1 cells compared to control-M1 cells (Fig 3C). These observations suggest that ability of Li to slow PIP$_2$ resynthesis following PLC activation requires an intact IMPA1.

We also analysed IMPA1$^{-/-}$-M1 cells for receptor activated Ca^{2+} influx, following Li treatment; this was not altered compared to untreated IMPA1$^{-/-}$-M1 cells (Fig 3D,E); in the same experiment, treatment of control-M1 cells with Li resulted in a big reduction in agonist activated Ca^{2+} influx. We also analyzed the two phases of receptor activated Ca^{2+} influx and found that while release of [Ca^{2+}]$_I$ from stores was not affected by Li treatment in IMPA$^{-/-}$-M1 cells (Fig 3F,G), the reduction of SOCE previously noted on Li treatment in control-M1 cells was still present (Fig 3F,H). When we expressed IMPA1 cDNA in the IMPA1$^{-/-}$-M1 cells, the receptor activated Ca^{2+} influx became similar to that of control-M1 cells (Fig 3J,K). These observations suggest that intact IMPA1 is required for the ability of Li treatment to reduce intracellular Ca^{2+} release following PLC activation.

Li reduces excitability in human cortical neurons

To test the relevance of our findings to human cortical neurons whose activity presumably underlies the therapeutic effect of Li in bipolar disorder patients, we generated human forebrain cortical neurons from induced pluripotent stem cells (iPSC) *in vitro* [44]. We differentiated forebrain cortical neurons from neural stem cells (NSC) derived from an Indian control iPSC line, D149 (Fig 4A) [45]. The generated NSCs expressedpreviously described

markers characteristic of NSC such as Nestin, SOX1, SOX2 and PAX6 (Fig 4B) and when differentiated, expressed the neuronal marker MAP2;a proportion of cells in the culture expressed the glial marker GFAP (Fig 4C). When these NSCs were differentiated into cortical neurons, they exhibited spontaneous $[Ca^{2+}]_I$ elevations, referred to as transients, that increase as a function of age *in vitro* [44, 46]. We measured $[Ca^{2+}]_I$ transients from such neuronal cultures at 45 *days in vitro* (DIV 45) and quantified their frequency. We observed that the frequency of $[Ca^{2+}]_I$ transients was decreased following Li treatment (1 mM) for 10 days (Fig 4D,E). We also measured changes in $[Ca^{2+}]_I$ following stimulation with carbachol in DIV45 cortical neurons treated with Li. In our cultures carbachol stimulation results in a rise in $[Ca^{2+}]_I$ that peaks before gradually decaying towards the baseline; in Li treated cultures the peak of this response was reduced (Fig 4F,G). We also performed experiments to measure Ca^{2+} mobilization from internal stores; these revealed that carbachol activated release of Ca^{2+} in Li treated neurons was reduced compared to those that were untreated (Fig 4H-J). By contrast there was only a modest difference in thapsigargin mediated rise in $[Ca^{2+}]_I$ between Li treated and untreated cortical neurons (Supp. Fig 4A-C). These observations suggest that both neuronal excitability as well as PLC mediated Ca^{2+} signalling is attenuated in cortical neurons following Li treatment.

Li treatment induces transcriptional changes in pathways involved in neuronal excitability

To understand the molecular mechanisms underlying the reduced excitability following Li treatment in human iPSC derived cortical neurons, we performed a transcriptomic analysis comparing untreated 45 DIV neurons and Li treated 45 DIV neurons (Fig 5A). Differentially expressed genes were obtained using DeSeq2; a cut-off of log_2FC change greater than 0.2 was used along with a p-value and FDR significance at less than 0.05 were considered. Using these criteria, 335 upregulated and 361 downregulated genes were obtained (Table 1 S1,S2). Gene Ontology (GO) analysis of the KEGG pathway showed substantial enrichment for genes annotated as being involved in calcium signalling or in glutamatergic synapse (Table 1 S3). In addition, GO KEGG analysis of the downregulated gene set also showed enrichment for several other categories of neurotransmission and neuromodulation related pathways (cholinergic, GABAergic, dopaminergic, glutamatergic and serotonergic) (Table 1 S2) (Fig 5B).

Since GO analysis revealed an enrichment in Ca signalling genes, we analysed molecular mechanisms through which Li might downregulate neuronal Ca signalling and hence excitability. There are multiple molecular mechanisms by which neuronal calcium signalling

is modulated [47] including voltage gated Ca channels and receptor activated Ca signalling, a mechanism used by several neurotransmitters and neuromodulators. Using the transcriptomic data, we tested the likely contribution of each of these mechanisms to the effects of Li in our model system.

Physiological and pharmacological evidence suggests that the $[Ca^{2+}]_I$ transients seen during neuronal development arise from the activity of voltage gated calcium channels (VGCC) [48] and this also seems to underlie the $[Ca^{2+}]_I$ transients seen in our experiments when human iPSC are differentiated into cortical neurons [44][this study]. To understand the mechanism by which Li could reduce the frequency of $[Ca^{2+}]_I$ transients we specifically studied the expression of genes encoding VGCC. Our transcriptomic analysis suggested that expression of genes encoding several subunits of VGCC might be downregulated (Table 1 S3). We experimentally tested the expression of genes for *CACNA1A, CACNA1B, CACNA1C, CAC1A1D* and *CACN1A1E* by qRT-PCR analysis; this showed that the expression of CACNA1A,B,C,D were all downregulated in Li treated DIV45 neurons compared to controls (Fig 5C) whereas CACNA1E was unchanged. In the same analysis, there was no downregulation in transcripts for *SCN1A (Nav1.1)* or *SCN9A (Nav7.1)* that encode the brain enriched, a-subunit of voltage gated sodium channels, involved in plasma membrane depolarization that trigger VGCC dependent $[Ca^{2+}]_I$ transients (Fig 5C).

Glutamate is a key excitatory neurotransmitter in the human central nervous system, exerting its effects through both metabotropic and ionotropic mechanisms [14] and changes in $[Ca^{2+}]_I$ isa key mechanism in glutamate signalling in the human brain. Our qRT-PCR analysis of transcript levels of genes encoding key components in the glutamate signalling system in Li treated neurons revealed that transcript levels for metabotropic glutamate receptors (*GRM1, GRM3, GRM5, GRM7*) and the ionotropic glutamate receptors (*GRIA1, GRIA2, GRIN1, GRIN2A, GRIN2B*) were downregulated (Fig 5D).

Many neurotransmitters and modulators transduce signal via G-protein coupled, PLC mediated store-operated Ca^{2+} influx [49]. Key molecular components of this pathway include the IP_3 receptor, ER Ca^{2+} sensor STIM and the Ca influx channel ORAI; we found that transcript levels of *PLCb4, ITPR2, STIM1* and *ORAI2* were all downregulated in Li treated neurons compared to controls (Fig 5E).

Mechanisms of Li induced transcriptional changes

To understand the mechanisms underlying Li induced transcriptome changes we analysed the likely transcriptional control mechanisms. For this we used the htFtarget database that

has information on all human transcription factors (TFs) and the genes they control, tissue

specific TF-target information, data on TFs for non-coding RNA and information on co-

regulation for a target gene and its TFs [50]. TFs controlling genes differentially expressed in

Li treated neurons (DIV 45) were identified from the htFtarget database. Out of 650 up

regulated genes in Li treated neurons, 587 were identified as associated with one or more

TFs in this database. Likewise, out of 525 downregulated genes, 333 genes were linked to a

TF. 485 and 323 unique TFs were identified that control the upregulated and downregulated

genes respectively. We then further looked for transcription factors that control differentially

expressed genes that are involved in calcium signalling (Table1 S3) (Fig 5F). Of the

upregulated genes, only RYR receptor was associated with three different transcription

factors (CTCF, SPI1 and USF1) that could control its expression. On the other hand, several

genes involved in calcium signalling that were downregulated were controlled by

transcription factors. Expression of *CACNA1C, CACNA1D, CAMK2B, CAMK4, CHRM3,*

GRIN1, ITPR2, RYR3, SCN9A and *SLC2A13* genes is regulated by transcription factors. Of

these *ITPR2* and *CAMK2B* are regulated by 10 different TFs, while *CACNA1D, CHRM3,*

RYR3, SCN9A and SLC2A13 are regulated by single transcription factor (Fig 5G). This

suggests that some of the genes that are differentially expressed in Li treated neurons are

under transcriptional control which can be a direct or indirect effect of Li treatment.

Discussion

Li is a monovalent ion with a remarkable therapeutic effect in the management of BPAD.

However only a proportion of human patients with BPAD respond to treatment and the

reasons underlying this variable response to Li remains unknown. In part this is due to a lack

of clear understanding on the mechanisms of action of Li in the brain. Although many

molecular targets of Li have been described in the literature, their role in mediating the action

of Li *in vivo* has not been directly tested.

One of the earliest cellular effects described for Li is its ability to slow Phosphoinositide

turnover in neural cells [51] and it has been proposed that the consequent slowing of

neurotransmitter activated neuronal excitability may underlie the effectiveness of the

compound[19] in managing BPAD. Since Li inhibits the enzyme IMPA1, one long-standing

proposal is that the application of Li to cells disrupts the receptor activated PIP_2 cycle and its

downstream effect on $[Ca^{2+}]_I$ signallingleading to reduced neuronal excitability. However,

there has not been a direct test of the model that the ability of Li to modulate inositol turnover

and $[Ca^{2+}]_I$ signalling *in vivo* requires the *IMPA1* gene product.

In this study, we found that in the application of Li to human cells resulted in a reduction in

$[Ca^{2+}]_l$ signalling following PLC activation. This reduction in $[Ca^{2+}]_l$ was underpinned by a

reduction in the release of Ca^{2+} from intracellular stores although the size of stores itself was

not altered by Li treatment, implying that Li affects the release mechanism. During PLC

activation, the principal mechanism by which Ca^{2+} is released is via the IP_3 receptor;

however, we found that the transcript levels for this gene were not altered by Li treatment

(Supp. Fig 2C). Thus, it seems likely that Li treatment affects IP_3 receptor activation, perhaps

by altering IP_3 production during receptor mediated PLC stimulation, a process that requires

adequate levels of the substrate for PLC, PIP_2. During this study we found that Li treatment

slows that rate of synthesis of PIP_2, the substrate from which PLC produces IP_3 and this

observation likely provides an explanation for the reduced intracellular Ca^{2+} release

phenotype seen following PLC stimulation.

We generated a cell line in which IMPA1 was deleted and found that in the absence of

IMPA1, Li was unable to exert its effects on both $[Ca^{2+}]_l$ signalling and PIP_2 resynthesis

during receptor activated PLC signalling. These findings imply that an intact IMPA1 is

required for the action of Li on these processes. Consistent with this idea, we noted that

receptor activated $[Ca^{2+}]_l$ signalling (Fig 3D,E) and the rate of PIP_2 resynthesis (Fig 3C)

following PLC activation was reduced in IMPA1$^{-/-}$ cells without Li treatment and in brain

cells[42]leading to the therapeutic effects of Li.

IMPA1 is an enzyme that dephosphorylates inositol 1-phosphate to generate inositol. Hence

biological effects resulting from Li inhibition could arise either from an accumulation of the

substrate or a deficiency of product. Indeed, Li treatment in rodent models has shown both

an elevation of inositol 1-phosphate levels as well as a modest reduction in inositol levels

[52][53]. In our analysis, we found that the lowered $[Ca^{2+}]_l$ influx following receptor activation

in human cells could be rescued by supplementation of the culture medium with inositol. This

observation implies that the signalling defect arising from Li inhibition of IMPA1 is a

consequence of inositol depletion rather than an accumulation of the substrate inositol 1-

phosphate. During the PLC activated PIP_2 cycle, inositol generated by the action of IMPA1 is

condensed with cytidine diphosphate diacylglycerol to form phosphatidylinositol[54] which is

then sequentially phosphorylated to generate PIP_2. Consistent with a role for reduced

inositol levels in the $[Ca^{2+}]_l$ signalling phenotype resulting from Li treatment, we found that

the resynthesis of PIP_2 was also slowed by treatment of wild type cells with Li. Together, our

data provide compelling evidence that Li treatment results in inhibition of IMPA1, a depletion

of the inositol pool leading to a reduced rate of PIP_2 resynthesis during PLC signalling.

When Li is used to treat BPAD in human patients, presumably it acts by reducing the
excitability of neurons in the cerebral cortex. In this study, we tested the effect of Li treatment
on human forebrain cortical neuronal cultures differentiated from iPSC. Consistent with
previous reports [55], we found that treatment of human forebrain cortical neurons with
therapeutically relevant concentrations of Li reduced the frequency of $[Ca^{2+}]_i$ transients that
are underpinned by voltage gated Ca^{2+} channel (VGCC) activity [44]. Since acute application
of Li does not affect $[Ca^{2+}]_i$ transients (Supp. Fig 4D,E), it seems unlikely that Li affects
neuronal excitability by directly inhibiting the VGCC. However, VGCC mediated $[Ca^{2+}]_i$
transients are activated by neuronal action potentials which themselves are triggered by
activation of post-synaptic G-protein coupled receptors that bind neurotransmitters. Several
of these receptors (mGluR1, mGluR5, mAchR and 5HT-2A) when bound to their respective
neurotransmitter ligand, use PLC activation as part of their signalling mechanism. Thus, Li
could influence neuronal excitability by modulating signalling through such PLC linked
receptors. In support of this hypothesis, this study we found that in human forebrain cortical
neurons, Li application diminished the elevations of $[Ca^{2+}]_i$ triggered by application of the
mAchR ligand carbachol (Fig 4 F-I). Thus, a key mechanism by which Li modulates neuronal
excitability may be through downregulation of neurotransmitter signalling through PLC linked
GPCRs.
742
Given the extended timeframe over which Li exerts its therapeutic effect it has been
proposed that in addition to directly impacting cellular biochemistry, transcriptional
mechanisms may play a role in its biological effects. SOCE that occurs downstream of
receptor activated PLC signalling has been shown to regulate transcription in neurons
[reviewed in [56] and a proposed target of Li, GSK-3β exerts its signalling effects via the
transcriptional regulator β-catenin [57]. In a transcriptomic analysis of human forebrain
cortical neurons, we found that Li treatment on a clinically relevant time and concentration
range results in a large transcriptional response. Bioinformatics analysis of Li treated
neurons did not reveal features of β-catenin dependent transcriptome changes and qRT-
PCR analysis of selected β-catenin regulated genes did not reveal any alteration in their
transcript levels in Li treated cells relative to controls (Supp. Fig 5A,B). Further, a
comparison of differentially expressed genes from genetic ablation of GSK-3β or the
chemical inhibition of this molecule[58] to the dataset of Li induced differential gene
expression (this study) revealed only a minimal overlap in the identity of genes altered in
both datasets (Supp. Fig 5F). Consistent with this, treatment of HEK293 cells with GSK-3β
inhibitor did not result in any alteration in receptor mediated $[Ca^{2+}]_i$ (Supp. Fig 5D,E).
Therefore, it seems that the transcriptome changes induced by Li treatment occur through a
non-GSK-3β/ β-catenin mediated mechanism. Likewise, a comparison of Li induced

differential gene expression (this study) to that elicited by genetic inhibition of SOCE [59, 60] also revealed only a modest overlap of genes that were differentially regulated (Supp Fig 4F,G). Therefore, it is likely that Li exerts transcriptional changes in neurons independent of its effects on SOCE.

Our analysis of the Li induced transcriptome changes in human forebrain cortical neurons however provide an insight into the mechanisms by which Li might mediate its effects both in cultured forebrain cortical neurons and also its clinical effects. We found that the levels of three of the four subunits of VGCC were downregulated in Li treated cells and this could explain in part the ability of Li to reduce $[Ca^{2+}]_i$ transients in forebrain cortical neurons in culture. A key feature of mania in BPAD patients is heightened neural activity manifested behaviorally, EEG and fMRI studies and presumably Li exerts its therapeutic effect through downregulating this neural activity. Glutamate is the principle excitatory neurotransmitter in the brain [14] and consistent with this we found that transcripts for multiple glutamate receptor subtypes were downregulated in forebrain cortical neurons treated with Li. These included multiple subunits of the metabotropic, AMPA and NMDA subtypes of the glutamate receptor family. Downregulation of these will likely contribute to reducing neural activity following Li treatment manifest clinically as behavioral improvement. Overall, this study demonstrates that the effects of Li on neuronal Ca^{2+} signalling are mediated through an IMPA dependent step and involves a large transcriptional response that downregulates molecular processes that are required for neuronal excitability. The link between Li and transcriptional responses are unclear although studies in yeast have shown that treatment with Li is associated with changes in the transcriptome that also includes changes in inositol biosynthesis.

Acknowledgements: This work was supported by the Department of Atomic Energy, Government of India, under Project Identification No. RTI 4006, the Department of Biotechnology, Government of India through the Accelerator Program for Discovery in Brain Disorders (BT/PR17316/MED/31/326/2015), the Pratiksha Trust and a Wellcome-DBT India Alliance Senior Fellowship to PR (IA/S/14/2/501540). We thank the NCBS Imaging, Mass spectrometry, Genomics, High performance computing, Stem cell and Biosafety facilities for support.

Conflict of Interest:

The authors declare no competing interests**.**

Bibliography

1. Toker L, Bersudsky Y, Plaschkes I, Chalifa-Caspi V, Berry GT, Buccafusca R, et al. Inositol-related gene knockouts mimic lithium's effect on mitochondrial function. Neuropsychopharmacology. 2014;39:319–328.

2. Osby U, Brandt L, Correia N, Ekbom a, Sparén P. Excess mortality in bipolar and unipolar disorder in Sweden. Arch Gen Psychiatry. 2001;58:844–850.

3. Lopez AD, Murray CC. JL. The global burden of disease , 1990 – 2020. Nat Med. 1998;4:1241–1243.

4. Geddes JR, Burgess S, Hawton K, Jamison K, Goodwin GM. Long-term lithium therapy for bipolar disorder: systematic review and meta-analysis of randomized controlled trials. Am J Psychiatry. 2004;161:217–222.

5. Cipriani A, Hawton K, Stockton S, Geddes JR. Lithium in the prevention of suicide in mood disorders: updated systematic review and meta-analysis. Bmj. 2013;3646:1–13.

6. Hou L, Heilbronner U, Degenhardt F, Adli M, Akiyama K, Akula N, et al. Genetic variants associated with response to lithium treatment in bipolar disorder: A genome-wide association study. The Lancet. 2016;387:1085–1093.

7. Manchia M, Adli M, Akula N, Ardau R, Aubry J-M, Backlund L, et al. Assessment of Response to Lithium Maintenance Treatment in Bipolar Disorder: A Consortium on Lithium Genetics (ConLiGen) Report. PLoS One. 2013;8:e65636.

8. Haggarty SJ, Karmacharya R, Perlis RH. Advances toward precision medicine for bipolar disorder: mechanisms & molecules. Mol Psychiatry. 2021;26:168–185.

9. Roux M, Dosseto A. From direct to indirect lithium targets: a comprehensive review of omics data. Metallomics. 2017;9:1326–1351.

10. Harwood a J. Lithium and bipolar mood disorder: the inositol-depletion hypothesis revisited. Mol Psychiatry. 2005;10:117–126.

11. Berridge MJ. Inositol trisphosphate and calcium signalling mechanisms. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 2009;1793:933–940.

12. Wasserman MJ, Corson TW, Sibony D, Cooke RG, Parikh S V., Pennefather PS, et al. Chronic lithium treatment attenuates intracellular calcium mobilization. Neuropsychopharmacology. 2004;29:759–769.

13. Yang YR, Kang D-S, Lee C, Seok H, Follo MY, Cocco L, et al. Primary phospholipase C and brain disorders. Adv Biol Regul. 2016;61:80–85.

14. Reiner A, Levitz J. Glutamatergic Signaling in the Central Nervous System: Ionotropic and Metabotropic Receptors in Concert. Neuron. 2018;98:1080–1098.

15. Cockcroft S, Raghu P. Topological organisation of the phosphatidylinositol 4,5-bisphosphate-phospholipase C resynthesis cycle: PITPs bridge the ER-PM gap. Biochemical Journal. 2016;473:4289–4310.

16. Allison, J.H. SMA. Reduced inositol in rat brain on lithium injection. Nat New Biol. 1971. 1971.

17. Allison JH, Blisner ME, Holland WH, Hipps PP, Sherman WR. Increased brain myo-
inositol 1-phosphate in lithium-treated rats. BiochemBiophys Res Commun.
1976;71:664–670.

18. Allison JH, Blisner ME, Holland WH, Hipps PP, Sherman WR. Increased brain myo-
inositol 1-phosphate in lithium-treated rats. BiochemBiophys Res Commun.
1976;71:664–670.

19. Berridge MJ, Downes CP, Hanley MR. Neural and developmental actions of lithium: a
unifying hypothesis. Cell. 1989;59.

20. Colantuoni C, Lipska BK, Ye T, Hyde TM, Tao R, Leek JT, et al. Temporal dynamics
and genetic control of transcription in the human prefrontal cortex. Nature.
2011;478:519–523.

21. Schubert CR, O'Donnell P, Quan J, Wendland JR, Xi HS, Winslow AR, et al.
BrainSeq: Neurogenomics to Drive Novel Target Discovery for Neuropsychiatric
Disorders. Neuron. 2015;88:1078–1083.

22. Ohnishi T, Ohba H, Seo KC, Im J, Sato Y, Iwayama Y, et al. Spatial expression
patterns and biochemical properties distinguish a second myo-inositol
monophosphatase IMPA2 from IMPA1. Journal of Biological Chemistry.
2007;282:637–646.

23. Sherman WR, Leavitt AL, Honchar MP, Hallcher LM, Phillips BE. Evidence that
Lithium Alters Phosphoinositide Metabolism: Chronic Administration Elevates
Primarily d-myo-Inositol-1-Phosphate in Cerebral Cortex of the Rat. J Neurochem.
1981;36:1947–1951.

24. Soares JC, Dippold CS, Mallinger AG. Platelet membrane phosphatidylinositol-4,5-
bisphosphate alterations in bipolar disorder--evidence from a single case study.
Psychiatry Res. 1997;69:197–202.

25. Soares JC, Mallinger AG, Dippold CS, Frank E, Kupfer DJ. Platelet membrane
phospholipids in euthymic bipolar disorder patients: are they affected by lithium
treatment? Biol Psychiatry. 1999;45:453–457.

26. Soares JC, Mallinger AG, Dippold CS, Forster Wells K, Frank E, Kupfer DJ. Effects of
lithium on platelet membrane phosphoinositides in bipolar disorder patients: a pilot
study. Psychopharmacology (Berl). 2000;149:12–16.

27. Cryns K, Shamir A, Van Acker N, Levi I, Daneels G, Goris I, et al. IMPA1 is essential
for embryonic development and lithium-like pilocarpine sensitivity.
Neuropsychopharmacology. 2008;33:674–684.

28. Sjoholt G, Gulbrandsen AK, Lovlie R, Berle JO, Molven A, Steen VM. A human myo-
inositol monophosphatase gene (IMPA2) localized in a putative susceptibility region
for bipolar disorder on chromosome 18p11.2: genomic structure and polymorphism
screening in manic- depressive patients. Mol Psychiatry. 2000;5:172–180.

29. Parker EM, Kameyama K, Higashijima T, Ross EM. Reconstitutively active G protein-
coupled receptors purified from baculovirus-infected insect cells. J Biol Chem.
1991;266:519–527.

30. Raghu P, Habib S, Hasnain SE, Kasan G. Development of a functional assay for
Ca^{2+}release activity of IP$_3$R and expression of an IP$_3$R gene
fragment in the baculovirus-insect cell system. Gene. 1997;190.

31. Várnai P, Balla T. Visualization of phosphoinositides that bind pleckstrin homology
domains: calcium- and agonist-induced dynamic changes and relationship to myo-
[3H]inositol-labeled phosphoinositide pools. J Cell Biol. 1998;143:501–510.

32. Chakrabarti P, Kolay S, Yadav S, Kumari K, Nair A, Trivedi D, et al. A dPIP5K
dependent pool of phosphatidylinositol 4,5 bisphosphate (PIP2) is required for G-
protein coupled signal transduction in Drosophila photoreceptors. PLoS Genet.
2015;11:e1004948.

33. Chang C-L, Liou J. Phosphatidylinositol 4,5-Bisphosphate Homeostasis Regulated by
Nir2 and Nir3 Proteins at Endoplasmic Reticulum-Plasma Membrane Junctions. J Biol
Chem. 2015;290:14289–14301.

34. Kim S, Kedan A, Marom M, Gavert N, Keinan O, Selitrennik M, et al. The
phosphatidylinositol-transfer protein Nir2 binds phosphatidic acid and positively
regulates phosphoinositide signalling. EMBO Rep. 2013;14:891–899.

35. Kim YJ, Guzman-Hernandez M-L, Wisniewski E, Balla T. Phosphatidylinositol-
Phosphatidic Acid Exchange by Nir2 at ER-PM Contact Sites Maintains
Phosphoinositide Signaling Competence. Dev Cell. 2015;33:549–561.

36. Francis J, Hamzeh RK, Lantin-Hermoso MR. Lithium toxicity-induced wide-complex
tachycardia in a pediatric patient. Journal of Pediatrics. 2004;145:235–240.

37. Prakriya M, Lewis RS. Store-Operated Calcium Channels. Physiol Rev.
2015;95:1383–1436.

38. Balla T. Phosphoinositides: Tiny Lipids With Giant Impact on Cell Regulation. Physiol
Rev. 2013;93:1019–1137.

39. Nixon DA. Cerebrospinal fluid inositol and its rise in post-mortem specimens. J
Physiol. 1955;129:272–280.

40. Swahn C -G. Gas chromatographic-mass spectrometric determination of myo-inositol
in human cerebrospinal fluid. J Neurochem. 1985;45:331–334.

41. Shetty HU, Holloway HW, Schapiro MB. Cerebrospinal fluid and plasma distribution of
myo-inositol and other polyols in Alzheimer disease. Clin Chem. 1996;42:298–302.

42. Hallcher LM, Sherman WR. The effects of lithium ion and other agents on the activity
of myo- inositol-1-phosphatase from bovine brain. J Biol Chem. 1980;255:10896–
10901.

43. V S, P V, QT D, RA A, V S, PD P, et al. 7Li nuclear-magnetic-resonance study of
lithium binding to myo-inositolmonophosphatase. Eur J Biochem. 1996;240:288–291.

44. Sharma Y, Saha S, Joseph A, Krishnan H, Raghu P. In vitro human stem cell derived
cultures to monitor calcium signaling in neuronal development and function [version 1;
peer review: 3 approved]. Wellcome Open Res. 2020;5:16:doi:
10.12688/wellcomeopenres.15626.1. eCollection.

45. Iyer S, Bhatia P, Rao M, Mukherjee O. Developing two reference control samples for
the Indian population. Stem Cell Res. 2018;30:38–42.

46.   Akhtar BM, Bhatia P, Acharya S, Sharma S, Sharma Y, Bhuvanendran Nair Suseela
921   Devi A, et al. A human stem cell resource to decipher the biochemical and cellular
922   basis of neurodevelopmental defects in Lowe syndrome. Biol Open. 2022;11.

923   47.   Berridge MJ. Neuronal calcium signaling. 1998;21:13–26.

924   48.   Rosenberg SS, Spitzer NC. Calcium Signaling in Neuronal Development. Cold Spring
925   HarbPerspect Biol. 2011;3:a004259–a004259.

926   49.   Moccia F, Zuccolo E, Soda T, Tanzi F, Guerra G, Mapelli L, et al. Stim and Orai
927   proteins in neuronal Ca(2+) signaling and excitability. Front Cell Neurosci. 2015;9.

928   50.   Zhang Q, Liu W, Zhang HM, Xie GY, Miao YR, Xia M, et al. hTFtarget: A
929   Comprehensive Database for Regulations of Human Transcription Factors and Their
930   Targets. Genomics Proteomics Bioinformatics. 2020;18:120–128.

931   51.   Berridge MJ, Downes CP, Hanley MR. Lithium amplifies agonist-dependent
932   phosphatidylinositol responses in brain and salivary glands. Biochem J.
933   1982;206:587–595.

934   52.   Sherman WR, Leavitt AL, Honchar MP, Hallcher LM, Phillips BE. Evidence that
935   Lithium Alters Phosphoinositide Metabolism: Chronic Administration Elevates
936   Primarily d-myo-Inositol-1-Phosphate in Cerebral Cortex of the Rat. J Neurochem.
937   1981;36:1947–1951.

938   53.   Sade Y, Toker L, Kara NZ, Einat H, Rapoport S, Moechars D, et al. IP3 accumulation
939   and/or inositol depletion: two downstream lithium's effects that may mediate its
940   behavioral and cellular changes. Transl Psychiatry. 2016;6:e968.

941   54.   Lykidis A, Jackson PD, Rock CO, Jackowski S. The Role of CDP-Diacylglycerol
942   Synthetase and Phosphatidylinositol Synthase Activity Levels in the Regulation of
943   Cellular Phosphatidylinositol Content. 1997;272:33402–33409.

944   55.   Mertens J, Wang Q-W, Kim Y, Yu DX, Pham S, Yang B, et al. Differential responses
945   to lithium in hyperexcitable neurons from patients with bipolar disorder. Nature.
946   2015;527:95–99.

947   56.   Mitra R, Hasan G. Store-operated Ca 2+ entry regulates neuronal gene expression
948   and function. CurrOpinNeurobiol. 2022;73.

949   57.   He R, Du S, Lei T, Xie X, Wang Y. Glycogen synthase kinase 3β in tumorigenesis and
950   oncotherapy (Review). Oncol Rep. 2020;44:2373–2385.

951   58.   Chen X, Wang R, Liu X, Wu Y, Zhou T, Yang Y, et al. A Chemical-Genetic Approach
952   Reveals the Distinct Roles of GSK3α and GSK3β in Regulating Embryonic Stem Cell
953   Fate. Dev Cell. 2017;43:563-576.e4.

954   59.   Gopurappilly R, Deb BK, Chakraborty P, Hasan G. Stable STIM1 Knockdown in Self-
955   Renewing Human Neural Precursors Promotes Premature Neural Differentiation.
956   Front Mol Neurosci. 2018;11.

957   60.   Dhanya SK, Hasan G. Purkinje Neurons with Loss of STIM1 Exhibit Age-Dependent
958   Changes in Gene Expression and Synaptic Components. J Neurosci. 2021;41:3777–
959   3798.

960   61.   Mukherjee O, Acharya S, Rao M. Making NSC and Neurons from Patient-Derived
961   Tissue Samples. Methods Mol Biol, vol. 1919, 2019. p. 9–24.

962   62.   Peh GSL, Lang RJ, Pera MF, Hawes SM. CD133 expression by neural progenitors
963   derived from human embryonic stem cells and its use for their prospective isolation.
964   Stem Cells Dev. 2009;18:269–282.

965   63.   Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca2+ indicators with greatly
966   improved fluorescence properties. J Biol Chem. 1985;260:3440–3450.

967   64.   Sharma S, Mathre S, Ramya V, Shinde D, Raghu P. Regulates Plasma-Membrane
968   PIP 3 Turnover and Regulates Plasma-Membrane PIP 3 Turnover and Insulin
969   Signaling. CellReports. 2019;27:1979-1990.e7.

Figure legends:

Figure 1- An *in vivo* model system to study the impact of Li on PLC induced PIP$_2$ turnover:

(A) The Phosphatidylinositol bisphosphate (PIP$_2$) cycle. (B) Schematic representation of an *in vivo* system elucidating the role of Nir2 and m1AchR in the PIP$_2$ cycle and the PH-PLCδ::GFP probe binding to the PIP$_2$ at the plasma membrane. (C) The PH-PLCδ::GFP probe translocates to the cytoplasm and back to the plasma membrane, corresponding to the hydrolysis and regeneration of PIP$_2$ respectively (control-M1; HEK293T cells expressing m1AchR). (D, E) Nir2 depletion leads to a delayed regeneration of PIP$_2$ at the plasma membrane. Changes in membrane localization of the PH-PLCδ::GFP probe after Cch stimulation (denoted by the ↑), are quantified via the fluorescence intensity ratio of the plasma membrane and the cytosol (PM/ Cyt) at various times. This ratio is normalized to the first time-point and the mean ± SD is plotted (from three experiments each performed in replicates) for control cells (orange line, n =33), and for cells treated with Nir2 shRNA (blue line, n =28). (F, G) Rate of regeneration of PIP$_2$ at the plasma membrane is shown for control-M1 cells (HEK293T cells expressing m1AchR) subjected to lithium treatment, compared to untreated cells. Mean ± SD is plotted from four experiments, each performed in replicates (untreated cells- orange line, n = 58; cells treated with Li- green line, n =53). [Scale Bar- 20 μM]. (H) Total PIP$_2$ levels recovered after various time intervals post stimulation with Cch.

(I,J) Total PIP$_2$ and PIP levels using LCMS from whole cell lipid extract of untreated and lithium treated cells (n=6 for both).

[Statistical test: (H) one-way ANOVA with post hoc Tukey's multiple pairwise comparisons. *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001 and (I,J) Multiple unpaired t test. *p value < 0.05; **p value <0.01; ***p value < 0.001].

Figure 2- Chronic Lithium treatment reduces PLC dependent intracellular Ca^{2+} mobilization:

(A) Basal [Ca^{2+}]$_I$, calculated in nM using Grynkiewicz equation, is unchanged in control-M1 cells due to lithium treatment (untreated cells- represented by orange, n = 133; cells treated with Li -represented by green line, n =156). (B-C) PLC dependent calcium mobilization is decreased in lithium treated cells. Quantification is done from multiple experiments, each with replicates (untreated cells (orange), n =133; cells treated with Li (green), n =156). (D-F) In zero Ca^{2+} buffer, agonist dependent calcium mobilization is reduced and subsequent Store Operated Calcium Entry (SOCE) is also decreased in lithium treated cells. Quantification: untreated cells (orange), n =145; cells treated with Li (green), n =146. (G-H) Intracellular store measured by Tg, is unaltered by lithium treatment. Quantification: untreated cells (orange), n =114; cells treated with Li (green), n =118. (I-J) Inositol supplementation reversed the lithium mediated decrease in intracellular Ca^{2+} release. Quantification: untreated cells (orange), n =98; cells treated with Li (green), n =114, cells treated with Li and high inositol (blue), n =104.

[Statistical tests: (C,E,F) Student's unpaired t test. *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001 and (J) one-way ANOVA with post hoc Tukey's multiple pairwise comparisons. *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001].

Figure 3- IMPA1 is required for the effect of Li on PIP$_2$ resynthesis and agonist mediated Ca^{2+} signalling:

(A) Western blot confirmation of IMPA1 knock-out cells. (B) Lithium does not alter the rate of regeneration of PIP$_2$ at the plasma membrane in the IMPA1$^{-/-}$-M1 cells (HEK293T-IMPA1$^{-/-}$ cells expressing m1AchR), as monitored by the changes in the translocation of the PH-PLCδ::GFP probe. Mean ± SD is plotted from four experiments, each performed in replicates (IMPA1$^{-/-}$ cells without Li treatment- blue line, n =50; IMPA1$^{-/-}$ cells treated with Li- purple line, n =47). (C) PIP$_2$ turnover post PLC stimulation in control-M1 (HEK293T cells expressing m1AchR) cells compared to IMPA1$^{-/-}$-M1 cells [control-M1 cells- orange line, n =50, IMPA1$^{-/-}$-M1 cells- blue line, n =51]. (D,E) Cch mediated calcium release in IMPA1$^{-/-}$-M1 cells are not altered by lithium. Quantification: control-M1 cells (orange), n =126; control-M1 cells treated with Li (green), n =127, IMPA1$^{-/-}$-M1 cells without Li treatment (blue), n =132; IMPA1$^{-/-}$-M1 cells treated with Li (purple), n =112. (F-H) In zero Ca^{2+} buffer, agonist dependent calcium mobilization is unaltered, but subsequent Store Operated Calcium Entry (SOCE) is decreased in lithium treated IMPA1$^{-/-}$-M1 cells. Quantification: IMPA1$^{-/-}$-M1 cells without Li

treatment (blue), n =179; IMPA1$^{-/-}$-M1 cells treated with Li (purple), n =193. (I) Western blot showing IMPA1 expression via lentiviral transduction of cDNA in IMPA1 knock-out cells. (J,K) IMPA1 expression via lentiviral transduction of cDNA in IMPA1$^{-/-}$-M1 cells cells reversed the decreased intracellular Ca^{2+} release. Quantification: control-M1 cells (orange), n =155; IMPA1$^{-/-}$-M1 cells (light blue), n =143, IMPA1$^{-/-}$-M1 cells + IMPA1 cDNA (dark blue), n =162.

[Statistical tests: (E,K) one-way ANOVA with post hoc Tukey's multiple pairwise comparisons. *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001 and (H) Student's unpaired t test. *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001].

Figure 4- Lithium reduces excitability in human cortical neurons:

(A) Overview of expansion and differentiation process of the D149 lineage Neural Stem Cells (NSCs) and the cortical neurons. (B) Characterization of NSCs via immunocytochemistry of known markers like Nestin, Sox2, Sox1 and Pax6. (C) The mature cortical neurons are characterized via immunocytochemistry of markers like MAP2, CTIP2 and GFAP (Astrocytes marker) [Scale Bar – 25 µM]. (D) [Ca^{2+}]$_i$ traces from individual cells (soma; Y-axis shows ΔF and X-axis is time in seconds. The baseline was recorded for 270 secs followed by addition of 10 µM Tetrodotoxin (TTX) (as indicated by the arrows) to block Ca^{2+} transients. Transients are higher in frequency in the neuronal cultures without lithium treatment, compared to the treated cultures. The number of spikes/ 270 secs are counted from individual soma and plotted, each dot representing events from a single soma. (E) Data is plotted from multiple experiments, each done with biological replicates (untreated neurons (orange), n =114; neurons treated with Li (green), n =113). (F-G) Cch dependent calcium mobilization was decreased in lithium treated neuronal cultures (DIV 45). Quantification: untreated neurons (orange), n =97; neurons treated with Li (green), n =106. (H-J) Decrease in the Cch dependent calcium mobilization as well as in Store Operated Calcium Entry (SOCE) cells in lithium treated neuronal cultures (DIV 45). Quantification: untreated neurons (orange), n =154; neurons treated with Li (green), n =156.

[Statistical tests: (E,G,I,J) Student's unpaired t test . *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001].

Figure 5- Lithium treatment induces transcriptional changes in pathways involved in neuronal transmission:

(A) Volcano plot showing expression of genes altered due to lithium treatment in human cortical neuronal cultures 45 DIV (p value <0.05, p adj value <0.01). (B) The GO KEGG pathway enrichment analysis for downregulated genes. (C-E) Quantitative analysis showing several genes related to voltage calcium channels (C), Glutamatergic pathway (D) and calcium homeostasis (E), are downregulated in cortical neurons (DIV 45) due to lithium treatment. (F) Transcription factors that regulate genes involved in calcium signalling were identified using hTFtarget database. The graph represents number of calcium signalling genes controlled by each transcription factor in the lithium treated neurons. (G) The graph represents number of transcription factors that control each gene involved in calcium signalling.

[Statistical tests: (C,D,E) Multiple unpaired t test . *p value < 0.05; **p value <0.01; ***p value < 0.001].

Table legends:

Table 1:

Table 1 S1: List of upregulated genes in Li treated neurons DIV 45.

The table shows Gene ID (ENSEMBL ID), Gene names, base Mean, $\log_2$ fold change, p value, p adjusted value and gene description (as derived from Deseq2 analysis)

Table 1 S2: List of downregulated genes in Li treated neurons DIV 45.

The table shows Gene ID (ENSEMBL ID), Gene names, base Mean, $\log_2$ fold change, p value, p adjusted value and gene description (as derived from Deseq2 analysis)

Table 1 S3: List of genes involved in calcium signalling.

The table shows Gene ID (ENSEMBL ID), Gene names, base Mean, $\log_2$ fold change, p value, p adjusted value and gene description (as derived from Deseq2 analysis)

Table 1 S4: GO pathway analysis for the upregulated genes in Li treated neurons DIV 45.

The table shows GO-Category, GO-Term, Count, Percentage of genes, p value, Gene IDs (ENSEMBL IDs), Fold Enrichment and its significance (as derived from DAVID GO analysis).

Table 1 S5: GO pathway analysis for the downregulated genes in Li treated neurons DIV 45.

The table shows GO-Category, GO-Term, Count, Percentage of genes, p value, Gene IDs (ENSEMBL IDs), Fold Enrichment and its significance (as derived from DAVID GO analysis).

Table 2:

1107   List of primers used for qRT-PCR.

1108   The table shows the gene names and the primer sequence in 5' to 3' orientation.

1109

Supplementary Figure legends:

1111

Supplementary Figure 1- An in vivo model system to study the impact of Li on PLC induced PIP$_2$ turnover:

(A) Nir2 downregulation due to shRNA. (B) Rate of regeneration of PIP$_2$ at the plasma membrane is shown for control-M1 cells transfected with Nir2 shRNA, compared to control-M1 cells transfected with scrambled shRNA. Mean ± SD is plotted from two experiments, each performed in replicates control cells (orange line, n =18), and for cells treated with Nir2 shRNA (blue line, n =21). (C,D) Relative abundance of the PIP$_2$ and PIP species shown, as detected by LC-MS/MS (MRM method in the positive mode).

[Statistical test: (A) Student's unpaired t test. *p value < 0.05; **p value <0.01; ***p value < 0.001.]

1122

Supplementary Figure 2- Chronic Lithium treatment reduces PLC dependent intracellular Ca^{2+} mobilization:

(A-C) Expression studies were done to check whether the decrease in SOCE in the lithium treated HEK293T cells was due to alteration of the genes involved in SOCE. No significant change occurred in terms of the expression of ITPR, STIM and ORAI in HEK293T cells due to lithium treatment.

1129

1130

Supplementary Figure 3- IMPA1 is required for the effect of Li on PIP$_2$ resynthesis and agonist mediated Ca^{2+} signalling:

(A) The starting region of the CDS of *IMPA1* gene is targeted by guide RNAs (gRNAs) and deleted by spCas9; (B) IMPA2 expression is not altered in these knock-out cells. *IMPA1* is the predominant gene in HEK293T cells that encodes for IMPase. (C, D) Western blot showing that both the reporter lines (control-M1 and IMPA1$^{-/-}$-M1 cells) exhibit similar expression of the m1AchR and thereby should achieve similar PC stimulation due to same concentration of Cch.

1139

1140

Supplementary Figure 4- Lithium reduces excitability in human cortical neurons:

(A-C) Intracellular store measured by Tg, is unaltered by lithium treatment in human cortical neurons. Quantification: untreated neurons (orange), n =145; neuronal cultures treated with Li (green), n =145. (D) $[Ca^{2+}]_I$ traces from individual cells (DIV 30 neurons) pre- and post-acute lithium treatment (1 mM Li for 5 mins) (soma; Y-axis shows ΔF and X-axis is time in seconds). Initially, the baseline was recorded for 300 secs- then 1 mM Li was added to the bath, incubated for 5 minutes and then recorded again for 300 secs. (E) The number of spikes/ 270 secs are counted from individual soma and plotted, each dot representing events from a single soma. Quantification- neurons before Li treatment (orange), n =41; neurons treated with Li for 5 mins (green), n =38. (F) The number of differentially expressed genes that are common between lithium treated neurons and in mouse Purkinje neurons where STIM1 is knocked out. (G) The number of differentially expressed genes that are common between lithium treated neurons and in neuronal progenitor cells where STIM1 is knocked down.

[Statistical tests: (B,C) Student's unpaired t test . *p value < 0.05; **p value <0.01; ***p value < 0.001; ****p value < 0.0001].

Supplementary Figure 5- Lithium treatment induces transcriptional changes in pathways involved in neuronal transmission:

(A,B) Transcriptomic analyses were done to check whether lithium treatment in mature neuronal affected genes involved in the Wnt signalling/ GSK-3β pathway. No significant alteration occurred in terms of the expression of the relevant genes due to lithium treatment. (C) Western blot showing CHIR99021 (an inhibitor of GSK-3β) treatment led to the high expression of β-Catenin, one of the targets of GSK-3β destruction complex. (D,E) Inhibition of GSK-3β by CHIR99021 did not alter agonist mediated Ca^{2+} release. Quantification: untreated cells (orange), n =116; cells treated with CHIR99021 (blue), n =118. (F) The number of differentially expressed genes that are common between lithium treated neurons and in mouse ES cells (Embryonic Stem cells) where GSK-3β is inhibited.

Saha Figure 1

Saha Figure 2

Saha Figure 3

DIV45 - Li

DIV45 + Li

Saha Figure 4

Saha Figure 5

Saha Supp Figure 1

A

B

C

Saha Supp Figure 2

A

B

C

D

Saha Supp Figure 3

A

Gene ID	Gene name	base Mean	log2 FoldChange	pvalue	padj
ENSG00000110092	cyclinD1	1317.99	0.28	0.00	0.00
ENSG00000078018	MAP2	48889.94	-0.14	0.00	0.01
ENSG00000177606	c-Jun	4851.97	0.15	0.00	0.02
ENSG00000141510	P53	571.73	0.24	0.00	0.05
ENSG00000080815	Presenilin1	1901.77	0.15	0.01	0.09
ENSG00000103197	TSC2	2096.37	-0.15	0.01	0.10
ENSG00000070018	LRP6	4202.20	0.14	0.01	0.11
ENSG00000198561	δ-catenin	8187.44	0.09	0.02	0.13
ENSG00000134982	APC	7104.11	-0.10	0.04	0.20
ENSG00000082701	GSK3B	0.00	0.08	0.04	0.22
ENSG00000149294	NCAM	32954.36	-0.06	0.08	0.30
ENSG00000178403	Neurogenin2	286.20	0.20	0.15	0.44
ENSG00000162992	NeuroD	722.22	0.11	0.17	0.46
ENSG00000142192	APP	37645.74	0.04	0.19	0.49
ENSG00000197971	Myelin basic protein	386.66	-0.12	0.23	0.55
ENSG00000118513	c-Myb	132.05	-0.16	0.33	0.64
ENSG00000168036	β-catenin	11020.97	-0.03	0.37	0.68
ENSG00000171862	PTEN	4465.42	-0.04	0.37	0.68
ENSG00000118260	CREB	2453.46	0.05	0.40	0.71
ENSG00000161202	Dvl	3265.50	0.03	0.42	0.72
ENSG00000186868	Tau	27952.23	-0.03	0.45	0.74
ENSG00000103126	Axin	958.57	0.05	0.50	0.77
ENSG00000170365	SMAD1	738.14	0.05	0.51	0.78
ENSG00000111361	eIF2B	1238.25	0.03	0.68	0.88
ENSG00000092964	CRMP2	31852.85	-0.01	0.78	0.92
ENSG00000074054	CLASP1	6068.76	-0.01	0.83	0.94
ENSG00000131196	NFAT	68.44	-0.04	0.87	0.96
ENSG00000131711	MAP1B	98292.60	0.01	0.87	0.96
ENSG00000162946	DISC1	117.71	-0.03	0.87	0.96
ENSG00000164885	CDK5	445.92	-0.01	0.88	0.96
ENSG00000130294	Kinesin	29314.79	0.00	0.94	0.98
ENSG00000136997	c-Myc	437.65	0.00	0.98	1.00

B

C

D

E

F

Saha Supp Figure 5