Robustness vs productivity during evolutionary community assembly: short-term synergies and long-term trade-offs

Vasco J. Lepori[1], Nicolas Loeuille[2], and Rudolf P. Rohr[1]

vasco.lepori@unifr.ch, nicolas.loeuille@sorbonne-universite.fr, and rudolf.rohr@unifr.ch

[1]Department of Biology – Ecology and Evolution, University of Fribourg

Chemin du Musée 10, CH-1700 Fribourg, Switzerland.

[2]Sorbonne Université, UPEC, CNRS, IRD, INRA

Institute of Ecology and Environmental Sciences, IEES

F-75005 Paris, France.

Abstract

The realization that evolutionary feedbacks need to be considered to fully grasp ecological dynamics has sparked recent interest in the effect of evolution on community properties like coexistence and productivity. However, modern coexistence theory being limited to pairwise interactions, little is known about coevolution and diversification in rich communities. Here we study such properties in a general trait-based model of competition on a niche axis. We leverage the structural approach to show that the effects of coevolution on coexistence are two-fold. On the short-term our results show synergies emerging between productivity and coexistence, while on the long-term diversification and niche-packing destabilize communities, thus inducing a long-term trade-off between productivity and coexistence. In light of classical and recent work, our findings help advance understanding of evolutionary effects in high-dimensional systems. We illustrate how our theoretical predictions echo in observed empirical patterns. Finally, we discuss their implications and provide testable hypotheses.

KEYWORDS: ECO-EVOLUTIONARY DYNAMICS, EVOLUTIONARY STABLE COMMUNITIES, DIVERSIFICATION, COEXISTENCE THEORY, STRUCTURAL STABILITY, NICHE DIFFERENCE, FITNESS DIFFERENCE, PRODUCTIVITY, PARETO OPTIMALITY

1   Introduction

While ecology and evolutionary biology have long developed as separate disciplines (Lewontin, 2003), recent years have seen an effort to better grasp the feedbacks that link variations in the biotic environment (inter- and intraspecific interactions) and evolutionary trajectories (McPeek, 2017). The realization that evolutionary changes can happen over relatively short time-scales (Lankau and Strauss, 2007; Fussmann et al., 2007; Yoshida et al., 2003; Hairston Jr et al., 2005) fostered a renewed interest in the interface between ecology and evolution, an area of research now named eco-evolutionary dynamics (Hendry, 2009). As species are involved in inter- and intraspecific interactions, these exert selective pressures, just like abiotic factors, that help determine the fitness of different phenotypes. In turn, evolutionary changes in ecological traits affect population dynamics and species interactions. This feedback loop can give rise to complex evolutionary dynamics beyond the simple optimization of growth rates, such as disruptive selection and evolutionary branching (Geritz et al., 1998).

Understanding the effects of evolution on the maintenance of species diversity remains an important, but difficult question. At the single species level, evolutionary adaptation may save species from extinction under specific conditions, a phenomenon named evolutionary rescue (Bell and Gonzalez, 2009). But accounting for density- or frequency-dependent selection can open up scenarios where a species evolve towards self-extinction (Matsuda and Abrams, 1994), a case dubbed evolutionary suicide in the adaptive dynamics literature (Ferriere and Legendre, 2013). More complex communities lead to complex eco-evo feedbacks, for which we have no reason to expect that evolution, which is driven by individual phenotypes fitness, will lead to optimization of emergent properties at the community scale (Metz et al., 2008). In a two species scenarios, there can be both evolution towards stronger niche differentiation (Lankau, 2009) or one species can push the other to extinction (Dercole et al., 2006). Recently, Pastore et al. (2021) studied the effect of co-evolution of niche positions in a model of two competing species in the framework of

Chesson's coexistence theory (Chesson, 2000; Barabás et al., 2018a). Their work shows that evolution tends to have a negative effect on coexistence by increasing competitive imbalance, an outcome matching the experimental results of Hart et al. (2019).

In species-rich communities, Brännström et al. (2012) highlighted the dual nature of competition in evolutionary models, which can lead to the increase of diversity at different levels through generation of polymorphism and speciation, but also to competitive exclusion and evolutionary murder (a situation in which evolution of a given species leads to the extinction of another species in the ecological network). Generally, classical ecological theory predicts that stability in arbitrary large systems may be difficult, with May's seminal paper showing that in randomly interacting communities dynamical stability (local stability analysis) decreased with richness (May, 1972), but adding non-random interactions such as adaptive foraging or eco-evo dynamics can counter this effect (Kondoh, 2003). Loeuille (2010) showed that evolution stabilizes moderately rich communities but destabilizes more species-rich systems.

However, studying local stability - the return to equilibrium after a state perturbation - is only half the picture (Rohr et al., 2014), the other half being feasibility - the existence of an equilibrium state where all species have positive abundances. The structural framework developed by Saavedra et al. (2017) allows us to quantify the tolerance to perturbations in the parameter space (such as changes in growth rates due to environmental factors). The study of feasibility of competing species on a niche axis is tightly linked to what classical theory calls species packing, a concept which dates back to Hutchinson's work on species niches (Hutchinson, 1957, 1979) and was mathematically formalized by MacArthur (1969). In their paper, they studied the limit in similarity between species, i.e., how close species can be on the niche axis while still coexisting. Case (1981) extended this approach by including coevolutionary forces, and studied limiting similarity under the constraint imposed by evolution.

If species interactions are important drivers of evolutionary change we expect that communities arising from co-evolution, for which Edwards et al. (2018) suggested the term

Evolutionary Stable Communities (ESCs), should comprise a highly non-random subset of all possible combinations of species. This has been advocated already by Rummel and Roughgarden (1983), which analyzed island communities modeled through coevolution processes versus random colonization. More recently, Aubree et al. (2020) showed that coevolved communities were generally more productive, more stable, more resistant to invasion than collections which were randomly assembled from the species pool.

Here, we aim to study how evolution and the emergence of polymorphism interplay with coexistence and productivity constraints. Using the structural approach to coexistence theory (Saavedra et al., 2017), we expand on previous approaches based on species pairs and dynamical stability, to focus on multispecies co-evolutionary communities. We start with a monomorphic population determined by its position on a niche axis representing resources. Niche position is the trait under selection, and we let it evolve until it reaches an evolutionary singular strategy, which may be invadable (a branching point) or evolutionary stable (ESS, Smith and Price (1973)). We let the system coevolve and undergo a series of branching events until it reaches a polymorphic evolutionary endpoint (Evolutionary Stable Community in the sense of Edwards et al. (2018)). Along co-evolutionary trajectories, we follow the structural indicators of niche and fitness difference. For a given niche difference, small fitness differences increase robustness through equalizing effects; similarly, for a given fitness difference, niche differences increase robustness due to stabilizing effects. However, as is widely recognized, these two metrics do not exist independently (Barabás et al., 2018b; Song et al., 2019). These two indicators are however insufficient to provide a complete picture of extinction risks. Therefore, we introduce a new structural metric of community robustness, η, and track its change along the evolutionary trajectory. Finally, we consider changes in community productivity, by studying the evolution of the total biomass and of the total relative yield – two common currencies of the biodiversity-ecosystem functioning literature (Loreau and Hector, 2001).

We expect evolution to cause character displacement along the resource axis (Macarthur and Levins, 1967; Dieckmann and Doebeli, 1999; Grant and Grant, 2006), which should

first result in an increase of niche differentiation. At the same time, divergence of niche positions ultimately results in a decrease in the growth rate of phenotypes situated further from the resource optimum, and in a larger imbalance of growth rates in the community and greater fitness differences. We therefore expect that effects of evolution on robustness and productivity may vary depending on the timescale considered, from short term gains, to longer term costs.

2 Materials and methods

2.1 Ecological dynamics

We study a niche-based model of competition based on generalized Lotka-Volterra dynamics (Case, 2000), following the α-r parametrization (Mallet, 2012). Each phenotype is defined by its position μ_i on a niche axis representing resources. Position on the niche axis affects both the intrinsic growth rates $r(\mu_i)$ and the strength $\alpha(\mu_i, \mu_j)$ of density-dependent competition interactions between types. This kind of model has been commonly used to model competition on a niche axis (MacArthur, 1972), and has been shown to readily lead to the evolutionary emergence of polymorphism (Dieckmann and Doebeli, 1999). We use this behavior to explore the conditions of coexistence throughout the diversification process. Population dynamics of a type i then follow:

$$\frac{dN_i}{dt} = N_i \cdot \left(r(\mu_i) - \sum_{j=1}^{S} \alpha(\mu_i, \mu_j) \cdot N_j \right) \qquad i = 1, \ldots, S. \tag{1}$$

We assume a Gaussian function for $r(\mu_i)$, with fecundity decreasing with distance from a resource optimum μ_R, while including a small density-independent intrinsic mortality m to prevent phenotypes with unreasonably large or small trait values from persisting at virtually nil abundances:

$$r(\mu_i) = f_{\max} \, e^{-\frac{(\mu_i - \mu_R)^2}{2\sigma_R^2}} - m. \tag{2}$$

The parameter $f_{\max}$ represents the maximum fecundity rate at the resource optimum, while σ_R depicts the width of resources availability on the niche axis.

In line with previous works (Macarthur and Levins, 1967), we suppose that competition strength is defined by the similarity among types. It then follows a Gaussian function centered in μ_i, so that $\alpha(\mu_i, \mu_j) = \alpha(\mu_j, \mu_i)$, and interaction strength reaches its maximum $\alpha_{\max}$ for $\mu_i = \mu_j$:

$$\alpha(\mu_i, \mu_j) = \alpha_{\max} \; e^{-\frac{(\mu_i - \mu_j)^2}{2\sigma_\alpha^2}}. \tag{3}$$

The parameter σ_α controls the width of the niche. This model is conceptually a size S extension of the model used by Dieckmann and Doebeli (1999); Doebeli and Dieckmann (2000); Taper and Case (1992). The form of the competition function presents the agreeable property of being dissipative *sensu* Volterra (Logofet, 1994; Volterra, 1931). This implies that ecological dynamics possess one and only one globally stable equilibrium point. Moreover, if there exists a feasible equilibrium, i.e., $N_i^* > 0$ for all i, all ecological dynamics will converge to it regardless of initial abundances. Note that a feasible equilibrium must fulfill $r(\mu_i) = \sum_{j=1}^{S} \alpha(\mu_i, \mu_j) \cdot N_j^*$ for each i.

2.2   Evolutionary dynamics

We study evolution within the adaptive dynamics framework (Metz et al., 1995; Brännström et al., 2013), thereby accounting for both frequency- and density-dependent selection. The adaptive dynamics framework assumes clonal reproduction where mutations are infinitesimally small and rare, and a separation of timescales where ecology is assumed much faster than evolution. This implies that advantageous mutations always go to fixation (invasion implies replacement). The evolution of a quantitative trait is determined by the invasion fitness function, defined as the *per capita* growth rate of a rare mutant of traits μ_i' in a resident population of S phenotypes with traits μ_i, $i = 1, \ldots, S$ at their ecological equilibrium $N_i^* > 0$, $i = 1, \ldots, S$ (Metz et al., 1992):

$$\omega_i(\mu_i' | \mu_1, \ldots, \mu_S) = r(\mu_i') - \sum_{j=1}^{S} \alpha(\mu_i', \mu_j) \cdot N_j^*. \tag{4}$$

Note, that each phenotype i has its invasion fitness function ω_i.

The evolution over time of niche positions μ_i, $i = 1, \ldots, S$ is determined by the canonical equation of adaptive dynamics (Dieckmann and Law, 1996):

$$\frac{d\mu_i}{dt} = c \cdot N_i^* \cdot \left. \frac{\partial \omega_i(\mu_i' | \mu_1, \ldots, \mu_S)}{\partial \mu_i'} \right|_{\mu_i' = \mu_i} \qquad i = 1, \ldots, S. \qquad (5)$$

The partial derivatives

$$\left. \frac{\partial \omega_i(\mu_i' | \mu_1, \ldots, \mu_S)}{\partial \mu_i'} \right|_{\mu_i' = \mu_i} \qquad i = 1, \ldots, S \qquad (6)$$

correspond to selection gradients, and they determine the direction of co-evolution of the S niche positions. The coefficient c incorporates the mutation rate and the mutational variance, which we assume to be equal for all phenotypes. Without loss of generality, we can set $c = 1$ by rescaling the time axis. We integrate the canonical equation (Equ. 5) numerically until an evolutionary singular strategy is reached, i.e, a set of traits μ_i^*, $i = 1, \ldots, S$, at which all selection gradients vanish

$$\left. \frac{\partial \omega_i(\mu_i' | \mu_1, \ldots, \mu_S)}{\partial \mu_i'} \right|_{\mu_i' = \mu_i = \mu_i^*} = 0 \qquad i = 1, \ldots, S. \qquad (7)$$

We start with a single monomorphic population and once we reach an evolutionary singular strategy, we evaluate the evolutionary stability condition (invasibility of the strategy) for each phenotype (Brännström et al., 2013; Metz et al., 1995). Invasibility happens when the second derivative of the invasion fitness function (Equ. 4) with the mutant trait is positive at the singular strategy (Dieckmann and Law, 1996; Geritz et al., 1998). Evolutionary branching ensues, and we introduce a new phenotype at a small distance of the singularity, thereby augmenting the dimensionality of the system. We continue the simulation until all strategies are evolutionary stable, thereby making an Evolutionary Stable Community (ESC) *sensu* Edwards et al. (2018). The algorithm is detailed in Supporting Information S1.

2.3 Coexistence metrics

Along co-evolutionary trajectories where $S > 2$, we assess coexistence metrics. Within the structural approach, coexistence is quantified by structural niche differences Ω and structural fitness differences θ. These two metrics are extensions of the stabilizing niche difference and the fitness ratio, defined within the modern coexistence theory (Chesson, 2000), to species-rich systems. The niche difference Ω quantifies the possibility of coexistence and is mathematically defined as the solid angle of the domain of intrinsic growth rates leading to coexistence given the interaction strength $\alpha(\mu_i, \mu_j)$, defining the feasibility domain (Figure 1). In turn, the fitness difference θ quantifies to what extent a phenotype dominates the system, up to the point that it may lead others to extinction. The fitness difference θ depends on both the intrinsic growth rates vector $\mathbf{r}(\mu_i)$ and the competition strength $\alpha(\mu_i, \mu_j)$. To compare values of niche difference across communities of different sizes S, we consider the standardized niche difference $\hat{\Omega} = \sqrt[s]{\Omega}$ (Song et al., 2018).

We also measure η, the smallest angle between the vector of intrinsic growth rates $\mathbf{r}$ and the border of the feasibility domain. This angle, which we call "robustness" gives an indication of the fragility of the community with respect to perturbations in $\mathbf{r}$, i.e., how close the community is to losing of one or more of its components. Figure 1 provides a representation of Ω, θ, and η in a system of $S = 2$. Metric computations are detailed in the Supporting Information S2.

2.4 Productivity metrics

Beside coexistence metrics, we focus on productivity indicators using two standard proxies: total community biomass (or abundance) $N_{\text{tot}} = \sum_i N_i^*$ is (Tilman et al., 1997) and total relative yield $Y_{\text{tot}} = (\sum_i N_i^*/K_i)$ Vandermeer (1989); Loreau and Hector (2001), where $K_i = r_i/\alpha_{ii}$ corresponds to the carrying capacity of phenotype i (monoculture abundance at equilibrium). The relative yield is a measure that has the advantage of quantifying over- ($Y_{\text{tot}} > 1$) or underyielding ($Y_{\text{tot}} < 1$), i.e., whether a mixture produces

Figure 1: **Structural coexistence metrics**. Geometric representation of the structural coexistence metrics for $S = 2$. The green cone, called domain of feasibility, determines the set of intrinsic growth rates leading to a positive equilibrium for each species, i.e., to coexistence. The niche difference Ω is given by the amplitude of this cone, while the fitness difference θ measures the deviation of the vector of intrinsic growth rates ($\mathbf{r}$) from the centroid of the cone. Finally, η – our new measure of robustness – is given by the smallest angle between the vector of intrinsic growth rates and the border of the feasibility domain. Note that we illustrated the structural metrics for $S = 2$, but they are effectively defined and computed for rich system ≥ 2, (see Supporting Information S2 for a three-dimensional version of this figure)

more or less than monocultures. Overyielding can be due to complementarity in resource use along an axis, which can be easily captured under our model, and is expected to vary along coevolutionary time.

2.5   Randomizations

We can follow the coexistence metrics and productivity along co-evolutionary trajectories and branching points. However, this does not tell us whether those properties are maximized by evolution. To assess this, we generate between 4'000 and 128'000 (depending on S) communities for each singular strategy (branching point or ESS) of each simulation, by sampling uniformly sets of niche positions μ_i, conditioned on the resulting community being feasible ($N_i^* > 0$ for all $i = 1, ..., S$). We then compare the metrics (niche and fitness difference, total biomass, total relative yield) of the sampled communities with the eco-evolutionary trajectories. We use two different rules to define the range of niche positions from which we sample. For the first rule, we sample communities within the range of niche positions observed at the ESC (restricted range), which tells us if the evolutionary community is unique within this bracket of niche positions. For the second rule, the range of sampled μ_i encapsulates all niche positions leading to positive intrinsic growth rate $r(\mu_i) > 0$ (full range). Such bounds are given by $\mu_R \pm \sqrt{2\sigma_R^2(\log f_{\max} - \log m)}$ and allow us to compare strategies with respect to the whole trait space.

2.6   Choice of parameter values

To reduce the number of free parameters, we first transform the dynamical system into a nondimensional form. That is, we freely chose the time unit, the abundance unit, and the scale of the niche axis. Regarding the abundance and time unit, we can set without loss of generality $\alpha_{\max} = 1$, $f_{\max} = 1$. The latter defines the ecological timescale, but adaptive dynamics assumes that ecological equilibrium is reached before the next mutation occurs. The combination of both determines the scale of species abundance and can arbitrarily be rescaled. The additional mortality rate m remains, and thus cannot be freely chosen. We show that our results are robust to changes in m.

Regarding the niche axis, we can rescale, without loss of generality, its origin to $\mu_R = 0$. Then we can also choose its scale by setting arbitrarily the resource width $\sigma_R = 1$ and explore the effect of the niche width σ_α, or alternatively, we can also set arbitrarily the niche width $\sigma_\alpha = 1$ and explore the effect of the resource width σ_R. In the main text, we set $m = 0.01$ and either 1) set $\sigma_\alpha = 1$ and vary σ_R, or 2) set $\sigma_R = 1$ and vary σ_α.

3   Results

3.1   Co-evolutionary trajectories and branching events

Figure (2.A) shows a typical co-evolutionary trajectory, which undergoes several diversification events until it reaches a stable and convergent community (ESC). After each branching, we observe a divergence in the niche positions, which leads to a decrease in the level of interspecific competition. The evolutionary endpoint for the monomorphic situation (i.e., before the first branching) can be determined analytically. It converges to an evolutionary singular strategy that is always located at the resource optimum $\mu^* = \mu_R$. Invasibility depends on the width of the competition function (σ_α), the width of the resources (σ_R), the maximum fecundity rate ($f_{\max}$), and the mortality rate (m). We demonstrate that branching occurs if and only if $\sigma_R^2 \cdot (f_{\max} - m) > \sigma_\alpha^2 \cdot f_{\max}$ (Supporting Information S3). Branching will therefore occur if the resource range is wide enough (as defined by σ_R) and/or limiting similarity strong enough (small σ_α). Note also that mortality has an evolutionary stabilizing effect. This formula is alike the one derived by Dieckmann and Doebeli (1999); Doebeli and Dieckmann (2000) but with an additional mortality term.

Figure 2B shows the evolution of the niche difference $\hat{\Omega}$. Between each branching point, the niche difference increases. This can be expected from panel A showing that niche positions diverge, which implies an increase in the niche difference and relaxed competition among phenotypes. But at the branching points, the addition of a new phenotype very close to an existing one causes the average niche difference to abruptly decrease.

Figure 2: **Evolutionary trajectory**. Panel **A** shows the evolutionary trajectory of the niche positions μ_i. The vertical dotted lines indicate evolutionary branching. Panel **B** shows the evolution of the niche difference ($\hat{\Omega}$). Panel **C** shows the evolution of fitness difference (θ) and the robustness (η). For this figure, the resource width has been set to $\sigma_R = 1.7$, the niche width to $\sigma_\alpha = 1$, and the mortality term to $m = 0.1$. Video 1 shows these changes over time in the coexistence space (Fig. 1), for $S = 3$.

Hence, for a period, coexistence is partly driven by neutral (fitness-equivalent) mechanisms, rather than by strong niche separation. More specifically, branching points are the only contiguous trait regions where indefinitely close phenotypes can coexist, creating a niche-neutrality continuum *sensu* Song et al. (2019). Conversely, fitness differences θ increase all along evolutionary trajectories, which undermines community robustness (Fig. 2C). Fitness differences θ are initially zero when only two types coexist, as their trait positions are symmetric around the resource center. They later tend to increase for $S > 2$ as phenotypes start differing in their intrinsic growth rates, reflecting imbalances between strong and weak competitors. Note that because evolution here increases both niche differences (stabilizing effect) and fitness differences (unequalizing effect) between branching points, its overall effect on robustness is not obvious. We therefore use our new indicator of robustness – the angle η – to assess the evolution of the distance to the border to the feasibility domain. Figure 2C shows that the robustness η tends to increase between the branching points, so that the overall effect of a simultaneous increase in both fitness and niche difference ultimately results in more robust communities. As logically expected, η decreases at each branching point. Moreover, as S increases and the community becomes saturated, the value of η becomes very small. Once maximum phenotype packing is achieved (at ESC), small perturbations in $\mathbf{r}$ suffice to lead to non-feasibility, and the community is structurally more fragile than an undersaturated one (fewer phenotypes than at the ESC).

Supporting Information S4 and S5 show that the results are consistent when varying the morality rate σ_R and the niche width σ_α.

3.2   *Effect of resource width on diversity at the ESC*

As expected, a larger resource space allows the emergence of a greater number of phenotypes (Figure 3A). Moreover, for a given number of phenotypes S, the standardized niche difference at the evolutionary singular strategy increases with resource width, as there is then more room for niche differentiation. Vice-versa, Panel B illustrates the effect of

varying the strength of similarity dependence in competition by changing niche widths σ_α of phenotypes (for a fixed resource width). For smaller niches, more phenotypes can be packed at the ESC, while for a given number of phenotypes, narrow niches allow a larger domain of feasibility $\hat{\Omega}$.

Figure 3: **Number of phenotypes at the ESC and at evolutionary branching points.** Panel **A** shows, for a fixed niche width $\sigma_\alpha = 1$, how varying the resource with σ_R modulates the number of consecutive branching points and, therefore, the number of phenotypes at the ESC. Cases marked with "ESC" represent evolutionary endpoints, while all the others are invasible singular strategies (branching points). The color represents the level of standardized niche difference at the evolutionary singular strategy. Making the resource space larger, for a fixed niche width, results in greater diversity at the ESC. Moreover, for a given number of phenotypes at the evolutionary singular strategy, the structural niche difference increases with resource width. Conversely, panel **B** shows the effect of changing the niche width σ_α for a fixed resource width $\sigma_R = 1$: the smaller the niche width, the more phenotypes can fit on a given resource axis.

3.3   Projection into coexistence space

Figure 4 illustrates how coexistence metrics $\hat{\Omega}$ and θ change along the evolutionary trajectory of figure 2 when $S \geq 2$. Remember that large niche differences $\hat{\Omega}$ and small fitness differences θ enhance persistence. In the background, we show the coexistence metrics of randomized communities that were produced according to the two rules presented in the Methods.

Considering all randomized communities, it appears that communities that have small fitness differences most often also have small niche differences. This implies a global trade-off at the community level between the two coexistence metrics. Points that optimize one of the two properties relative to the other lie on a Pareto front. Figure 4 shows that evolution leads to this Pareto-optimality, and more specifically to the point on this front that also maximizes the standardized niche difference within the restricted range (first randomization rule). More extreme niche positions would also allow larger niche differences, but this decrease in competition would come at the cost of decreasing intrinsic growth rates for the phenotypes further from the resource optimum, thereby increasing fitness differences θ.

Regarding community robustness (η), figure 5 indicates that evolution optimizes it within the range of niche positions explored by evolution (first randomization rule), but greater robustness could (rarely) be reached for the second randomization rules. However, among the niche positions explored by evolution and for a given number of phenotypes, evolution converges to more robust communities, by optimizing the niche difference Ω and the robustness η.

Figure 4: **Variations of coexistence metrics along evolutionary trajectories.**
The figure is split into six panels according to the number of phenotypes S. Each panel
therefore represent a slice (in between two branching points) of the trajectory presented
in figure 2. The colored line represents the projection of the evolutionary trajectory
of figure 2 into the space of coexistence metrics for a given S, variations of color from
purple to orange indicating time variations. The X-axis represents the standardized
niche difference ($\hat{\Omega}$), while the Y-axis stands for the fitness difference (θ). Gray points
represent randomized feasible communities, with niche position value sampled from the
restricted range explored by evolution (1st rule; dark gray) or the full range leading to
positive intrinsic growth rates (2nd rule; light gray). Pareto fronts among randomized
communities are shown using dashed lines.

Figure 5: **Evolution of community robustness.** The red line shows the evolution of η. Gray points represent randomized feasible communities, with niche position value sampled from the restricted range explored by evolution (1st rule; dark gray) or the full range leading to positive intrinsic growth rates (2nd rule; light gray). At the singular strategy, the evolved community reaches a value of η greater or equal to than a proportion $P1$ and $P2$ of randomized communities according to the first and second randomization rules, respectively.

3.4   Evolution of productivity

In general, co-evolutionary dynamics increase productivity. While strict optimization can be proven in monomorphic situation (Supporting Information S3), productivity increase also happens throughout diversification, as illustrated by figure 6. For the total abundance (panel A), the co-evolutionary trajectory, though not strictly optimal, is among the most productive communities. This holds for the two randomization rules. On the other hand, the total relatively yield (panel B) shows a contrasting pattern between the two randomization rules. When the communities are sampled within the restricted range of niche positions explored by evolution (1st rule, panel B), the total relative yield reached by evolution is larger than almost all randomized communities. If communities are how-

ever sampled within the full range, the total relative yield reached by evolution is inferior to most randomized situations. This can be explained as follows: the total relative yield is the sum of abundance for each phenotype relative to its carrying capacity. It correlates with the strength of interspecific competition (relative to intraspecific competition). For S phenotypes the maximum total relative yield is equal to S, and can be achieved only in absence of interspecific competition. We have seen on figure 4, that within the restricted range of randomization, co-evolution tends to maximize niche differentiation, and thus to decrease interspecific competition. Given a fuller range of randomization, however, communities with larger niche differentiation can be sampled.

Figure 6: **Productivity along evolutionary trajectories.** Evolution of productivity. Here the numbers $P1$ and $P2$ in the second row of text indicate the percentage of randomized communities that are less productive than the ESS. For S=1, biomass production is optimized at the ESS. For S=6, the ESS is in the 80th percentile of the most productive communities. Point clouds represent randomization in the narrow (1st rule; dark gray) or the full range (2nd rule; light gray).

4   Discussion

In light of global changes, understanding the way evolution shapes and affects community robustness is paramount. Communities are currently reshuffled by the arrival of non-native species and by range shifts (Alexander et al., 2016). How these novel, evolutionary non-equilibrium communities differ from ESCs is unclear. We also need to assess under which conditions and how quickly evolution may prevent species from going extinct following environmental change, considering not only population-level processes (as in evolutionary rescue) but also larger spatial or organizational scales (Urban, 2016). Previous studies showed that mechanisms such as adaptive foraging, which can be evolutionary in nature, can stabilize complex systems (Kondoh, 2003; Valdovinos et al., 2013). Recently a handful of studies, both theoretical and experimental, have explored this question for pairs of coevolving species within the framework of modern coexistence theory Pastore et al. (2021); Hart et al. (2019). Here we go further by investigating the question for species-rich systems ($S > 2$) where diversity arises through subsequent branching events. We show that evolution of coexistence properties follows two distinct trends on two different timescales, while productivity systematically increases along evolutionary trajectories.

The short-term trend, in-between branching events, allows for increased efficiency in resource partitioning and decreased competition. On this timescale, evolution promotes niche differentiation ($\hat{\Omega}$), but this effect is counterbalanced by a simultaneous increase in fitness difference (θ) (Fig. 2). Hence, coexistence is enhanced by favoring niche partitioning, rather than by neutral mechanisms. Since we observe both an increase in the stabilizing mechanisms (increase in niche differences) and a reduction of the equalizing mechanism (in fitness differences), as in (Pastore et al., 2021), those two metrics are not enough to conclude whether evolution tend to more or less robust communities. Our new measure of robustness η shows that between branching communities, evolution selects less fragile communities with respect to environmental perturbations that impact growth

rates **r** (Saavedra et al., 2017). The fact that coexistence metrics are usually not independent is already appreciated (Barabás et al., 2018a; Song et al., 2019), but we here show how they are readily coupled throughout evolutionary dynamics. This link is made obvious in our trait-based model (Fig. 4), where niche position affects both growth rates and competition strength, in turn constraining Ω and θ. This interdependence makes it impossible to minimize simultaneously niche overlap and fitness differences: randomized communities with high Ω values tend to have high θ (Fig. 4). Hence, evolution leads to communities that are not strict optima of Ω nor minima of θ but rather lie on a Pareto front, an optimal compromise that is reached between niche increasing differentiation and restricting fitness imbalances. In a rare direct empirical test of this short term effect, Hart et al. (2019) find an increase in fitness difference after evolution, but not one in niche difference, which they explain may have been prevented by competition for a single discrete, non-substitutable resource. Still, evidence of (evolutionary) niche differentiation measured as character displacement abounds in other natural and experimental settings (Grant and Grant, 2006; Dayan and Simberloff, 2005).

Our results suggest that costs in terms of community robustness may happen on a longer timescale, where evolution leads to diversification along with evolutionary niche packing. Branching points, by virtue of addition of one more phenotype, whose trait value and fitness are initially very close to an existing one, have an equalizing effect on the fitness difference θ and a destabilizing effect on the niche difference $\hat{\Omega}$ and robustness η. As the community size increases, robustness and niche differences peak at increasingly lower levels. This is a consequence of trying to pack a greater amount of phenotypes S in the same resource width σ_R (Fig. 3). Classical theory shows that the number of phenotypes that can be packed on a resource axis is a function of resource width σ_R relative to the niche width of species (Macarthur and Levins, 1967). But when evolution is allowed in the community, the richness at the ESC is lower than under strict maximum (non-evolutionary) packing. In fact, ESCs are by definition uninvadable, because the fitness landscape for any possible invading trait value μ_m is zero at the resident trait values

$\mu_1, ..., \mu_S$ and negative everywhere else; however, richer, feasible non-evolutionary configurations do exist for a given environment. This result has been shown to be consistent across a range of models, for both continuous and discrete resources (e.g., Case (1981); Rummel and Roughgarden (1983); Shoresh et al. (2008)). It is worth pointing out that the answer to whether evolution helps or hinders coexistence is context-dependent: when the community is undersaturated (fewer members than at the ESC), evolution drives an increase in robustness η. Conversely, by starting with feasible but supersaturated communities (more diversity than allowed at the ESC), η will repeatedly drop to zero and we observe a sequence of extinction events until the ESC is reached. For example, Shoresh et al. (2008) found that evolution destabilizes communities, which can be explained by the fact that they started with supersaturated communities. This also helps explain why (Loeuille, 2010) found evolution to be usually stabilizing for small communities (likely undersaturated), while its effect is reversed for rich communities (likely oversaturated). Empirical evidence for the longer term effects of evolution on community properties is difficult to acquire, but could be found in phylogenetic patterns of niche conservatism (Mayfield and Levine, 2010). For example, Yguel et al. (2016) report patterns consistent with increased productivity due to niche diversification and filling.

Ultimately, the evolutionary process results in ESCs that are highly non-random. Coevolved communities also show more evenly-spaced trait distributions than non-evolutionary communities in the model of Barabás and D'Andrea (2016) (but see Bennett et al. (2013)). Our results show that they are also more structurally robust. This has important implications in the context of global change: if coevolved communities are more robust to environmental stresses (changes in temperature, water, nutrients levels) than non-coevolved communities, this could have important consequences in terms of managing global changes that may need to be considered when guiding conservation efforts. Indeed, studies on the effects of global change often focus on single-species responses, but many have argued that community-level responses should be given more attention (Walther, 2010; Alexander et al., 2016; Gilman et al., 2010). Because community composition and interactions are

modified by climate change and invasive species, leading to new assemblages that likely depart from coevolved structures (David et al., 2017; Blowes et al., 2019), our results suggest that the robustness of these new communities may be relatively poor. Further experimental testing of community-level responses to environmental stress in co-evolved versus randomized assemblages should prove an exciting and critically needed avenue of research, and the structural approach provides a useful theoretical framework to tackle the question. Experimental testing could for instance be undertaken using systems that rapidly diversify (e.g., the *Pseudomonas* system in Rainey and Travisano (1998)), where the diversifying community could be stressed and compared to isolated assemblages of phenotypes.

Regarding productivity, rich randomized communities tend to be more productive on average than poor ones (Fig. 6). By requiring randomized communities to be feasible, we however introduce a selection bias: in this sense, the positive slope between richness and productivity is indeed a byproduct of coexistence (Pillai and Gouhier, 2019), since the conditions that promote coexistence (namely niche differences), are also those that promote complementarity and greater productivity. Thus, we observe the emergence of trade-offs between community properties: higher total biomass values are possible among randomized communities, but these tend to have larger fitness imbalance and/or lower community robustness. This finding echoes results of Rohr et al. (2016) showing that very productive communities have low evenness, while those with lower deviation from the feasibility domain center (θ) have intermediate levels of biomass production. In agreement with Aubree et al. (2020), our results also show that evolutionary communities are often more productive on average than randomized ones, especially when species richness is low. Total abundance increases monotonically along evolutionary trajectories, first because of selection for higher intrinsic growth rates in the monomorphic case (selection effect), then due to a better use of the resource space by the diversifying phenotypes (complementarity effect) (Fig. 2). This is consistent with the expectation that increased diversification within and across species should lead to occupation of vacant niche spaces, leading to

increased niche complementarity and utilization, and ultimately increased abundances at the consumer level (a hypothesis supported by Yguel et al. (2016)). For instance, some of the largest biodiversity ecosystem-functioning experiments have consistently reported an increase in the net effect of biodiversity on biomass production and of niche complementarity across a decade (Fargione et al. (2007); Marquard et al. (2009); Cardinale et al. (2007)). Similarly, Stefan et al. (2022) showed that plant-plant interactions shifted towards increased complementarity and yield over just a few generations of coexistence, while van Moorsel et al. (2018) report higher productivity in polycultures with an 8-year co-evolution history compared to identical-composition, but evolutionary naive plant communities. This evidence suggests that the complementarity effect often observed in biodiversity-ecosystem functioning experiments is expected from an evolutionary point of view, and likely reinforcing along evolutionary trajectories.

It is, however, possible to find more productive combinations among the random communities. Indeed, strict evolutionary optimization of total biomass at equilibrium only arises for the monomorphic case in this class of model and is lost in higher-dimensional systems (Rohr and Loeuille, 2020). Mathematically, this is because for $S = 1$, the selection gradient coincides with the gradient of $r(\mu)$ (Supporting Information S3). In polymorphic systems, maximizing biomass productivity would require further niche differentiation, at the expense of increasing fitness imbalance, which is not achieved under the Pareto-optimality engendered by evolution. This runs contrary to a widespread belief that evolution begets optimality: while growth rate optimization might be the norm in simple systems with weak or no interactions, this assumption does not generally hold true if we introduce realistic eco-evolutionary feedback loops (see also Lion and Metz (2018) for a treatment in epidemiological models and Metz and Geritz (2016) for a discussion on optimization principles). Our results show that evolution largely departs from optimality when considering relative yields. Indeed, ESC communities have markedly lower values than most of the randomized communities. This could have meaningful implications for biodiversity-ecosystem functioning experiments (Loreau and Hector, 2001; Tilman et al.,

2014). Such experiments often assemble communities at random from a species pool, then use relative yields to evaluate effects on functioning. Our results suggest that overyielding in natural, coevolved communities may be substantially lower. This should be further tested in experimental systems, for instance by studying the changes in overyielding across time in biodiversity ecosystem-functioning experiments.

Several questions remain open for investigation. We considered only horizontal communities competing on a single resource axis, where packing more than two species with the same amount of niche overlap between all of them is impossible Macarthur and Levins (1967). But if phenotypes coevolve in a multidimensional trait space (two resource axes), neutral configurations with multiple species are possible. In addition, we considered only evolving niche positions in our model, but niche width could also evolve, leading to possible diversification between generalists and specialists species. Nevertheless, the theoretical predictions of our study need to be further experimentally tested. Although there is empirical support in biodiversity ecosystem-functioning studies regarding the increase in niche complementarity and biomass production over time, it remains unclear whether those predictions would hold in co-evolved communities emerging from diversification. Experimentally, comparison of properties of ESCs to non-evolutionary communities is complicated by the fact that ESCs are conceptually useful, but whether they are frequent in nature, and how to go about identifying them, is unclear (Edwards et al., 2018). Despite these hurdles, experimental tests of our theoretical results could provide timely evidence that would contribute to our understanding of evolution of communities properties and their interplay in rich systems. Such experiments could be undertaken in microbial systems, where rapidly evolving communities could be compared to control treatments of non-coevolved assemblages (Rainey and Travisano, 1998; Altermatt et al., 2015).

Acknowledgements

This work has been funded by the Swiss National Science Foundation grant 31003A_182386.

Data and code

Numerical simulations of eco-evolutionary trajectories were performed in Julia 1.5.1 (Bezanson, Edelman, Karpinski, and Shah, 2017), while computation of metrics of co-existence and plotting of results were done in R 4.0.3 (R Core Team, 2020). Code to reproduce the analyses will be made available on github.

References

Alexander, J. M., Diez, J. M., Hart, S. P., and Levine, J. M. 2016. When Climate Reshuffles Competitors: A Call for Experimental Macroecology. Trends in Ecology and Evolution 2147:1–11. ISSN 01695347.

Altermatt, F., Fronhofer, E. A., Garnier, A., Giometto, A., Hammes, F., Klecka, J., Legrand, D., Mächler, E., Massie, T. M., Pennekamp, F., Plebani, M., Pontarp, M., Schtickzelle, N., Thuillier, V., and Petchey, O. L. 2015. Big answers from small worlds: A user's guide for protist microcosms as a model system in ecology and evolution. Methods in Ecology and Evolution 6:218–231. ISSN 2041210X.

Aubree, F., David, P., Jarne, P., Loreau, M., Mouquet, N., and Calcagno, V. 2020. How community adaptation affects biodiversity–ecosystem functioning relationships. Ecology Letters 23:1263–1275. ISSN 14610248.

Barabás, G. and D'Andrea, R. 2016. The effect of intraspecific variation and heritability on community pattern and robustness. Ecology Letters 19:977–986. ISSN 14610248.

Barabás, G., D'Andrea, R., and Stump, S. M. 2018a. Chesson's coexistence theory. Ecological Monographs 0:1–27. ISSN 00129615.

Barabás, G., D'Andrea, R., and Stump, S. M. 2018b. Chesson's coexistence theory. Ecological Monographs 88:277–303. ISSN 15577015.

Bell, G. and Gonzalez, A. 2009. Evolutionary rescue can prevent extinction following environmental change. Ecology Letters 12:942–948. ISSN 1461-0248.

Bennett, J. A., Lamb, E. G., Hall, J. C., Cardinal-McTeague, W. M., and Cahill Jr., J. F. 2013. Increased competition does not lead to increased phylogenetic overdispersion in a native grassland. Ecology Letters 16:1168–1176. ISSN 1461-0248.

Bezanson, J., Edelman, A., Karpinski, S., and Shah, V. B. 2017. Julia: A fresh approach to numerical computing. SIAM review 59:65–98. URL https://doi.org/10.1137/141000671.

Blowes, S. A., Supp, S. R., Antão, L. H., Bates, A., Bruelheide, H., Chase, J. M., Moyes, F., Magurran, A., McGill, B., Myers-Smith, I. H., Winter, M., Bjorkman, A. D., Bowler, D. E., Byrnes, J. E. K., Gonzalez, A., Hines, J., Isbell, F., Jones, H. P., Navarro, L. M., Thompson, P. L., Vellend, M., Waldock, C., and Dornelas, M. 2019. The geography of biodiversity change in marine and terrestrial assemblages. Science 366:339–345. ISSN 0036-8075, 1095-9203.

Brännström, Å., Johansson, J., Loeuille, N., Kristensen, N., Troost, T. A., Lambers, R. H. R., and Dieckmann, U. 2012. Modelling the ecology and evolution of communities: A review of past achievements, current efforts, and future promises. Evolutionary Ecology Research 14:601–625. ISSN 1522-0613.

Brännström, Å., Johansson, J., and von Festenberg, N. 2013. The Hitchhiker's Guide to Adaptive Dynamics. Games 4:304–328.

Cardinale, B. J., Wright, J. P., Cadotte, M. W., Carroll, I. T., Hector, A., Srivastava, D. S., Loreau, M., and Weis, J. J. 2007. Impacts of plant diversity on biomass production increase through time because of species complementarity. Proceedings of the National Academy of Sciences 104:18123–18128.

Case, T. J. 1981. Niche packing and coevolution in competition communities. Proceedings of the National Academy of Sciences of the United States of America 78:5021–5025. ISSN 00278424.

Case, T. J., 2000. An illustrated guide to theoretical ecology, volume 449. Oxford University Press New York.

Chesson, P. 2000. Mechanisms of Maintenance of Species Diversity. Annual Review of Ecology and Systematics 31:343–366. ISSN 0066-4162.

David, P., Thébault, E., Anneville, O., Duyck, P. F., Chapuis, E., and Loeuille, N., 2017. Chapter One - Impacts of Invasive Species on Food Webs: A Review of Empirical Data. Pages 1–60 *in* D. A. Bohan, A. J. Dumbrell, and F. Massol, eds. Advances in Ecological Research, volume 56 of *Networks of Invasion: A Synthesis of Concepts.* Academic Press.

Dayan, T. and Simberloff, D. 2005. Ecological and community-wide character displacement: The next generation. Ecology Letters 8:875–894. ISSN 1461-0248.

Dercole, F., Ferrière, R., Gragnani, A., and Rinaldi, S. 2006. Coevolution of slow–fast populations: Evolutionary sliding, evolutionary pseudo-equilibria and complex Red Queen dynamics. Proceedings of the Royal Society B: Biological Sciences 273:983–990.

Dieckmann, U. and Doebeli, M. 1999. On the origin of species by sympatric speciation. Nature 400:354–357. ISSN 00280836.

Dieckmann, U. and Law, R. 1996. The dynamical theory of coevolution: A derivation from stochastic ecological processes. Journal of Mathematical Biology 34:579–612. ISSN 14321416.

Doebeli, M. and Dieckmann, U. 2000. Evolutionary branching and sympatric speciation caused by different types of ecological interactions. American Naturalist 156. ISSN 00030147.

Edwards, K. F., Kremer, C. T., Miller, E. T., Osmond, M. M., Litchman, E., and Klausmeier, C. A. 2018. Evolutionarily stable communities: A framework for understanding the role of trait evolution in the maintenance of diversity. Ecology Letters 21:1853–1868. ISSN 14610248.

Fargione, J., Tilman, D., Dybzinski, R., Lambers, J. H. R., Clark, C., Harpole, W. S., Knops, J. M., Reich, P. B., and Loreau, M. 2007. From selection to complementarity: Shifts in the causes of biodiversity–productivity relationships in a long-term biodiversity experiment. Proceedings of the Royal Society B: Biological Sciences 274:871–876.

Ferriere, R. and Legendre, S. 2013. Eco-evolutionary feedbacks, adaptive dynamics and evolutionary rescue theory. Philosophical Transactions of the Royal Society B: Biological Sciences 368:20120081.

Fussmann, G. F., Loreau, M., and Abrams, P. A. 2007. Eco-evolutionary dynamics of communities and ecosystems. Functional Ecology 21:465–477. ISSN 02698463.

Geritz, S. A., Kisdi, É., Meszéna, G., and Metz, J. A. 1998. Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. Evolutionary Ecology ISSN 02697653.

Gilman, S. E., Urban, M. C., Tewksbury, J., Gilchrist, G. W., and Holt, R. D. 2010. A framework for community interactions under climate change. Trends in Ecology and Evolution 25:325–331. ISSN 01695347.

Grant, P. R. and Grant, B. R. 2006. Evolution of character displacement in Darwin's finches. Science 313:224–226. ISSN 00368075.

Hairston Jr, N. G., Ellner, S. P., Geber, M. A., Yoshida, T., and Fox, J. A. 2005. Rapid evolution and the convergence of ecological and evolutionary time. Ecology Letters 8:1114–1127. ISSN 1461-0248.

Hart, S. P., Turcotte, M. M., and Levine, J. M. 2019. Effects of rapid evolution on species coexistence. Proceedings of the National Academy of Sciences of the United States of America 116:2112–2117. ISSN 10916490.

Hendry, A. P. A. P., 2009. Eco-evolutionary dynamics. Princeton University Press, Princeton. ISBN 9781400883080.

Hutchinson, G. E. 1957. Concluding remarks. Cold Spring Harbor Symposia on Quantitative Biology 22:415–427. URL `http://symposium.cshlp.org/content/22/415.short`.

Hutchinson, G. E., 1979. An introduction to population ecology. Yale University Press, New Haven, Conn., 1st ed. 1978, 3rd print. 1978 ; 1st ed., 4th print. 1979 edition. ISBN 0300021550.

Kondoh, M. 2003. Foraging Adaptation and the Relationship Between Food-Web Complexity and Stability. Science 299:1388–1391.

Lankau, R. A. 2009. Genetic Variation Promotes Long-Term Coexistence of Brassica nigra and Its Competitors. The American Naturalist 174:E40–E53. ISSN 0003-0147.

Lankau, R. A. and Strauss, S. Y. 2007. Mutual feedbacks maintain both genetic and species diversity in a plant community. Science 317:1561–1563. ISSN 00368075.

Lewontin, R., 2003. Building a science of population biology. Pages 7–20 *in* R. S. Singh and M. K. Uyenoyama, eds. The Evolution of Population Biology. Cambridge University Press, Cambridge.

Lion, S. and Metz, J. A. 2018. Beyond R0 Maximisation: On Pathogen Evolution and Environmental Dimensions. Trends in Ecology and Evolution 33:458–473. ISSN 01695347.

Loeuille, N. 2010. Influence of evolution on the stability of ecological communities. Ecology Letters 13:1536–1545. ISSN 14610248.

Logofet, D. O., 1994. Matrices and Graphs: Stability Problems in Mathematical Ecology, CRC Press, New York (1993)., volume 56. CRC Press. URL `http://link.springer.com/10.1016/S0092-8240(05)80313-1`.

Loreau, M. and Hector, a. 2001. Partitioning selection and complementarity in biodiversity experiments. Nature 412:72–6. ISSN 0028-0836.

MacArthur, R. 1969. Species packing, and what competition minimizes. Proceedings of the National Academy of Sciences 64:1369–1371. ISSN 0027-8424.

Macarthur, R. and Levins, R. 1967. The Limiting Similarity , Convergence , and Divergence of Coexisting Species Author (s): Robert Macarthur and Richard Levins Source : The American Naturalist , Vol . 101 , No . 921 (Sep . - Oct ., 1967), pp . 377-385 Published by : The University of C. American Naturalist 101:377–385.

MacArthur, R. H., 1972. Geographical ecology. Patterns in the distribution of species. Harper and Row.

Mallet, J. 2012. The struggle for existence: How the notion of carrying capacity, K, obscures the links between demography, Darwinian evolution, and speciation. Evolutionary Ecology Research 14:627–665. ISSN 15220613.

Marquard, E., Weigelt, A., Temperton, V. M., Roscher, C., Schumacher, J., Buchmann, N., Fischer, M., Weisser, W. W., and Schmid, B. 2009. Plant species richness and functional composition drive overyielding in a six-year grassland experiment. Ecology 90:3290–3302. ISSN 1939-9170.

Matsuda, H. and Abrams, P. A. 1994. Runaway Evolution to Self-Extinction Under Asymmetrical Competition. Evolution 48:1764–1772. ISSN 0014-3820.

May, R. M. 1972. Will a large complex system be stable? Nature 238:413–414. ISSN 00280836.

Mayfield, M. M. and Levine, J. M. 2010. Opposing effects of competitive exclusion on the phylogenetic structure of communities. Ecology Letters 13:1085–1093. ISSN 1461023X.

McPeek, M. A., 2017. Evolutionary Community Ecology, Volume 58. Monographs in Population Biology ; 77. Princeton University Press,, Princeton, NJ. ISBN 1-4008-8821-2.

Metz, J., Geritz, S., Meszena, G., Jacobs, F., and van Heerwaarden, J., 1995. Adaptive dynamics: A geometrical study of the consequences of nearly faithful reproduction. Iiasa working paper, IIASA, Laxenburg, Austria. URL `http://pure.iiasa.ac.at/id/eprint/4497/`.

Metz, J., Mylius, S., and Diekmann, O. 2008. When does evolution optimize? Evol. Evolutionary Ecology Research 10.

Metz, J. A. and Geritz, S. A. 2016. Frequency dependence 3.0: An attempt at codifying the evolutionary ecology perspective. Journal of Mathematical Biology 72:1011–1037. ISSN 14321416.

Metz, J. A. J., Nisbet, R. M., and Geritz, S. A. H. 1992. How should we define 'fitness' for general ecological scenarios? Trends in Ecology & Evolution 7:198–202. ISSN 0169-5347.

Pastore, A. I., Barabás, G., Bimler, M. D., Mayfield, M. M., and Miller, T. E. 2021. The evolution of niche overlap and competitive differences. Nature Ecology and Evolution Pages 21–23. ISSN 2397334X.

Pillai, P. and Gouhier, T. C. 2019. Not even wrong: The spurious measurement of biodiversity's effects on ecosystem functioning. Ecology 100:1–12. ISSN 00129658.

R Core Team, 2020. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. URL `https://www.R-project.org/`.

Rainey, P. B. and Travisano, M. 1998. Adaptive radiation in a heterogeneous environment. Nature 394:69–72. ISSN 1476-4687.

Rohr, R. P. and Loeuille, N. 2020. Effects of evolution on niche displacement and emergent population properties , a discussion on optimality. BioRxiv Pages 1–46.

Rohr, R. P., Saavedra, S., and Bascompte, J. 2014. On the structural stability of mutualistic systems. Science 345. ISSN 10959203.

Rohr, R. P., Saavedra, S., Peralta, G., Frost, C. M., Bersier, L. F., Bascompte, J., and Tylianakis, J. M. 2016. Persist or produce: A community trade-off tuned by species evenness. American Naturalist 188:411–422. ISSN 00030147.

Rummel, J. D. and Roughgarden, J. 1983. Some Differences between Invasion-Structured and Coevolution-Structured Competitive Communities: A Preliminary Theoretical Analysis. Oikos 41:477. ISSN 00301299.

Saavedra, S., Rohr, R. P., Bascompte, J., Godoy, O., Kraft, N. J., and Levine, J. M. 2017. A structural approach for understanding multispecies coexistence. Ecological Monographs 87:470–486. ISSN 15577015.

Shoresh, N., Hegreness, M., and Kishony, R. 2008. Evolution exacerbates the paradox of the plankton. Proceedings of the National Academy of Sciences of the United States of America 105:12365–12369. ISSN 00278424.

Smith, J. M. and Price, G. R. 1973. The Logic of Animal Conflict. Nature 246:15–18. ISSN 1476-4687.

Song, C., Barabás, G., and Saavedra, S. 2019. On the Consequences of the Interdependence of Stabilizing and Equalizing Mechanisms. The American Naturalist 194:000–000. ISSN 0003-0147.

Song, C., Rohr, R. P., and Saavedra, S. 2018. A guideline to study the feasibility domain of multi-trophic and changing ecological communities. Journal of Theoretical Biology 450:30–36. ISSN 10958541.

Stefan, L., Engbersen, N., and Schöb, C. 2022. Rapid transgenerational adaptation in response to intercropping reduces competition. eLife 11:e77577. ISSN 2050-084X.

Taper, M. L. and Case, T. J. 1992. Models of character displacement and the theoretical robustness of taxon cycles. Evolution 46:317–333. ISSN 00143820.

Tilman, D., Isbell, F., and Cowles, J. M. 2014. Biodiversity and ecosystem functioning. Annual Review of Ecology, Evolution, and Systematics 45:471–493. ISSN 15452069.

Tilman, D., Knops, J., Wedin, D., Reich, P., Ritchie, M., and Siemann, E. 1997. The Influence of Functional Diversity and Composition on Ecosystem Processes. Science 277:1300–1302.

Urban, M. 2016. Improving the forecast for biodiversity under climate change. Science (New York, N.Y.) 353:1293–1310. ISSN 1095-9203.

Valdovinos, F. S., Moisset de Espanés, P., Flores, J. D., and Ramos-Jiliberto, R. 2013. Adaptive foraging allows the maintenance of biodiversity of pollination networks. Oikos 122:907–917. ISSN 1600-0706.

van Moorsel, S. J., Hahl, T., Wagg, C., De Deyn, G. B., Flynn, D. F. B., Zuppinger-Dingley, D., and Schmid, B. 2018. Community evolution increases plant productivity at low diversity. Ecology Letters 21:128–137. ISSN 1461-0248.

Vandermeer, J. H., 1989. The Ecology of Intercropping. Cambridge University Press.

Volterra, V., 1931. Leçons sur la théorie mathématique de la lutte pour la vie. Les grands classiques Gauthier-Villars. J. Gabay, Sceaux. ISBN 2876470667.

Walther, G.-r. 2010. Community and ecosystem responses to recent climate change. Philosophical Transactions of the Royal Society B: Biological Sciences 365:2019–24. ISSN 1471-2970.

Yguel, B., Jactel, H., Pearse, I. S., Moen, D., Winter, M., Hortal, J., Helmus, M. R., Kühn, I., Pavoine, S., Purschke, O., Weiher, E., Violle, C., Ozinga, W., Brändle, M., Bartish, I., and Prinzing, A. 2016. The evolutionary legacy of diversification predicts ecosystem function. American Naturalist 188:398–410. ISSN 00030147.

Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F., and Hairston, N. G. 2003. Rapid evolution drives ecological dynamics in a predator-prey system. Nature 424:303–306. ISSN 00280836.