1   Alternating selection for dispersal and multicellularity favors regulated life cycles
2
3   Julien Barrere[1], Piyush Nanda[1], Andrew W. Murray[1,2]
4
5

6   Affiliations:
7
8   [1] Department of Molecular and Cellular Biology, Harvard University, 52 Oxford Street,
9   Cambridge, MA 02138, USA
10 [2] Lead Contact
11
12 Correspondence: awm@mcb.harvard.edu (A.W.M.)

14 Keywords: life cycle, evolution of multicellularity, *Saccharomyces cerevisiae* wild isolates,
15 clustering, patchy resources, public goods.

17 **Summary:**
18
The evolution of complex multicellularity opened paths to increased morphological diversity and
organizational novelty. This transition involved three processes: cells remained attached to one
another to form groups, cells within these groups differentiated to perform different tasks, and the
groups evolved new reproductive strategies[1–5]. Recent experiments identified selective pressures
and mutations that can drive the emergence of simple multicellularity and cell differentiation[6–11]
but the evolution of life cycles, in particular, how simple multicellular forms reproduce has been
understudied. The selective pressure and mechanisms that produced a regular alternation
between single cells and multicellular collectives are still unclear[12]. To probe the factors regulating
simple multicellular life cycles, we examined a collection of wild isolates of the budding yeast, *S.
cerevisiae*[12]. We found that all these strains can exist as multicellular clusters, a phenotype that
is controlled by the mating type locus and strongly influenced by the nutritional environment.
Inspired by this variation, we engineered inducible dispersal in a multicellular laboratory strain
and demonstrated that a regulated life cycle has an advantage over constitutively single-celled or
constitutively multicellular life cycles when the environment alternates between favoring
intercellular cooperation (a low sucrose concentration) and dispersal (a patchy environment
generated by emulsion). Our results suggest that simple multicellularity in wild isolates could be
under selection and is regulated by their genetic composition and the environments they
encounter and that alternating patterns of resource availability may have played a role in the
evolution of life cycles.

Results and Discussion:

During the evolution of multicellularity, single cells start forming clusters of multiple cells. For this single cell to cluster transition to be complete, the group must reproduce. The simplest strategy is random breakage of the cluster into two or more smaller clusters, creating a minimal life cycle. While early life cycles might have been rudimentary, more complex life cycles have evolved resulting in a plethora of reproductive strategies with different group formation mechanisms, group features (size, shape, etc.), and propagation modes (propagule size and the signal, mechanism, and timing of offspring release)[13]. Here we explore life cycle variation within a species and the factors influencing these life cycles. We then ask which conditions could select for the evolution of regulated life cycles where multicellular growth is followed by a phase of dispersal into single cells.

The conditions that favor the evolution of multicellularity impose high costs on reproducing through single cells[6,7,14,15], suggesting that additional selective pressures would be required to evolve regulated life cycles with single-celled intermediates. Patchily distributed resources have been proposed as a factor that could select for dispersal and thus favor the production of single-cell propagules[16]: at the same total number of cells, single cells can colonize more resource patches than multicellular clusters. Theoretical work[17–21] supports the idea that alternation between selection for multicellularity and selection for dispersal could result in the evolution of regulated multicellular life cycles but few experimental studies have tested these theoretical predictions[22,23] or examined factors influencing multicellularity within natural isolates of a single species.

Here, we characterized wild isolates of the budding yeast, *S. cerevisiae*, and showed that they form clonal multicellular clusters as part of their life cycles. These experiments also revealed that both the mating type locus and the identity of nutrients regulate the presence and size of multicellular clusters. Inspired by the environmental variation, we used a laboratory strain, W303, to engineer and compete three life cycles: constitutively single-celled, constitutively multicellular, and a regulated alternation between single cells and multicellular clusters. The single-celled life cycle was most fit in a patchy environment, the multicellular life cycle was fittest when the extracellular hydrolysis of sucrose selected for intercellular cooperation, and the regulated life cycles was fittest when conditions fluctuated between these two environments.

The mating type locus regulates *S. cerevisiae's* multicellularity

Laboratory strains of *S. cerevisiae* were selected to be unicellular when they were domesticated, a feature that makes them useful for studying evolutionary forces that can select for the evolution of multicellularity[6,7]. We examined wild isolates to determine the effect of genetics and the environment on simple forms of multicellularity. Here we focus on clonal multicellularity (groups formed by persistence of the shared cell wall that links mother and the daughter cells). To characterize life cycles, we focused on 1) quantifying the size and size distribution of multicellular clusters, and 2) identifying factors that influence cluster size.

S. cerevisiae can grow as clonal, undifferentiated multicellular clusters[7,24,25]. This simple
multicellular phenotype has been observed in engineered and laboratory evolved strains[6,7,25].
Clusters have also been described in the wild isolates RM11-a and YL1C[26,27], prompting us to ask
if this phenotype is also present in other *S. cerevisiae* isolates. We examined 22 phylogenetically
diverse wild strains from the SGRP-2 collection, sampled from various environments (wine, sake,
soil, baking, etc.)[12,28]. As a control, we included *S*288c, the unicellular lab strain whose sequence
is the *S. cerevisiae* reference genome.
96
We began by validating forward scatter, in flow cytometry, as a measure of cross sectional area
and a tool to distinguish single cells from clusters. We verified that the forward scatter of
polystyrene beads is linearly correlated with their measured cross sectional area (Figure S1A)
and compared forward scatter measurements of single cells to strains known to form clusters.
Cultures of these clustering strains are known to include clusters of different sizes as well single
cells[7]. In addition, the area of a cluster is influenced both by the number and the size of cells it
contains. To detect cluster formation, we use a "clustering score" that is the ratio of the mean
forward scatter of the 10% largest objects in the population to the mean forward scatter of the
10% smallest objects (Figure 1A). By focusing on the 10% largest objects, this score identifies if
clusters are formed in a population. Dividing by the smallest 10% of the objects, which are almost
all single cells, corrects for the differences in cell size between environments and strains. This
clustering score is therefore a relative value to compare cluster formation in different conditions
and genetic backgrounds. We validated this score using diploid and haploid single cells as well
as haploid clusters: single cells of different sizes show similar clustering scores, while clusters
have a higher value (Figure 1A). To ensure that we were only detecting clonal multicellularity, we
prevented   aggregative   multicellularity   (often   referred   to   as   flocculation)   by   using
ethylenediaminetetraacetic acid (EDTA) to chelate divalent cations before measuring the
clustering score[29].
115
We measured the clustering score of all 22 wild isolates and S288c both as diploids and haploids
of both mating types (**a** and α) in rich, glucose-containing medium (YPD). Our measurements
showed that in their diploid state, the strains have low clustering scores, comparable to the values
of *S*288c (Figure 1B). The haploids derived from wild isolates, however, showed a much greater
clustering score, suggesting the presence of multicellular clusters (Figure 1B). Microscopy on a
subset of these strains confirmed that they form multicellular clusters (Figure 1C) and reveal that
natural isolates harbor a wide diversity of cluster morphology and size.
123
The contrasting phenotypes of haploids and diploids suggested that the mating type locus, which
is heterozygous in diploids *(MATa/MATα)* and present in a single copy in haploids *(MATa* or
MATα), plays a role in controlling multicellularity. We tested this hypothesis by randomly selecting
5 diploid isolates and creating derivatives homozygous at the mating type locus (*MATa/MATa* or
MATα/MATα diploids). Figure 1D shows that becoming homozygous at the mating type greatly
increased the size and frequency of clusters. Our results reveal that the formation of clonal
multicellular clusters in wild isolates is controlled at least in part by the mating type locus.
131
Environment modulates multicellular life cycle in *S. cerevisiae* wild isolates

Environmental cues often drive transitions between different phases of life cycles[30]. Hence, we
asked if the formation of clusters is influenced by the environment in which they grow.

We grew our strains in rich medium (YEP, yeast extract and peptone) containing five different
carbon sources (glucose, sucrose, galactose, glycerol or ethanol). For each carbon source, we
tested 2 different concentrations (0.2 and 2% (w/v)), and measured the clustering score in each
of these environments.

Figure 2A and 2B show that the clustering score of some strains changes drastically between
different environments. The effect of the carbon source was highest at a concentration of 2%
(Figure 2C and Figure S2). The strongest effect was observed with the diploids in galactose: while
diploids form single cells in glucose, many strains form clusters in galactose. Our results show
that the multicellularity of *S. cerevisiae* is regulated by the environment, and suggest that the
environment can act as a signal for switching between single-celled and multicellular stages of
the yeast life cycle.

Fluctuation between bulk sucrose and glucose emulsion selects for regulated life cycles

Observing that the nutritional environment could influence the degree of multicellularity prompted
us to ask if environmental oscillations could select for alternation between single-celled and
multicellular phases of a life cycle. To address this question, we engineered the standard yeast
laboratory strain, W303, previously used to study multicellularity[7,25]. We analyzed how the
environment influenced competitions between strains with three reproductive modes: 1) a
constitutively single-celled life cycle; 2) a constitutively multicellular life cycle and 3) a regulated
life cycle able to switch between single cells and clusters (Figure 3A).

For the constitutively single-celled life cycle, we used a prototrophic, single-celled W303 strain.
For constitutive multicellularity we deleted *ACE2*, which encodes a transcription factor that
induces genes that separate daughter cells from their mothers, in the same background. Finally,
to create a regulated life cycle, we placed *ACE2* under the control of a promoter that is activated
by β-estradiol[31] allowing the strain to grow as clusters when uninduced and as single cells in the
presence of β-estradiol. To confirm that we could control cluster size in the *ACE2*-inducible strain,
we measured cluster size at increasing β-estradiol concentrations, and the number of generations
needed for clusters to become single cells and vice versa. Figure 3C shows that increasing β-
estradiol decreases cluster size and Figure 3D reveals that the switches between single cells and
clusters happens in a few generations.

We used the three strains to ask how different environments would affect their fitness. It has been
suggested that environmental cycles (tidal, seasonal, trophic, etc.) could have driven the evolution
of regulated life cycles alternating between two phenotypes (i.e. single cell and cluster)[32]. Hence,
we asked if an environment cycle that alternated between favoring single cells and favoring
clusters would favor a regulated life cycle.

Because previous experiments[7,25] showed that yeast clusters grow better than single cells in low
concentrations of sucrose, we used low sucrose as a selection for clusters. We tested the
advantage of clusters by competing them against single cells in different concentration of sucrose.
Because extracellular invertase hydrolyses sucrose to glucose and fructose, we used a mixture
of glucose and fructose as a control, with both media having the same hexose monomer
concentration (ranging from 10 to 40 mM). We also tested the effect of dilution at each transfer
during the competition experiment (ranging from 100-fold to 10,000-fold). These experiments
confirmed that clusters, grown on sucrose, have a fitness advantage over single cells and that
this advantage is stronger as the sucrose concentration decreases (Figure S4A). We also
confirmed that the dilution regime (strength of dilution between transfers of the competition
experiment) greatly influences the fitness. Dilutions lower than 10,000-fold do not result in any
fitness differences between single cells and clusters (Figure S4B).
189
These and previous results[7,25] suggest that increased cluster size should correlate with an
increased fitness on low sucrose concentrations. To directly test this hypothesis we measured
the fitness of the inducibly multicellular strain expressing different levels of Ace2, allowing us to
directly measure fitness as a function of size (Figure 4C). Figure 4A shows a positive relationship
between size and fitness in low sucrose, confirming that size directly correlates with fitness
advantage in low sucrose. In the control, where a mixture of glucose and fructose replaced
sucrose, there was no difference in fitness as a function of size as the clusters are too small to
limit the diffusion of nutrients to the center of the cluster[33].
198
To favor single cells, we created a patchy environment, in which resources were concentrated in
small, separated patches by emulsifying glucose-containing medium in an inert oil. When
populations are seeded in this environment, at a density of less than one cell per glucose-
containing droplet, single cells should outcompete clusters thanks to their dispersal advantage: a
cluster of 10 cells can only colonize 1 patch, while 10 individual cells can colonize 10 patches.
Because the yield in a patch is set by the amount of the limiting resource in the patch, not the
number of cells that invade it, more colonized patches result in a greater final population size.
206
We used an emulsion, whose droplets contained glucose and fructose, as a patchy environment.
The starting cell density was adjusted so that on average a droplet does not contain more than
one cell, which means that clusters will seed a much smaller fraction of the patches than single
cells. The cultures were grown to saturation for 48 h in emulsion before breaking the emulsion,
diluting the culture and starting a new cycle of emulsion growth. We confirmed the stability of the
emulsion by measuring the size of droplets just after emulsification and after 48h (Figure S4D).
In addition, we confirmed that the emulsification protocol itself did not bias the frequency of single
cells and clusters: we broke the emulsion immediately after its creation and measured the
frequency of single cells and clusters (Figure S4C). Figure 4B shows that as the size of the
clusters increases their fitness relative to single cells decreases. As a control, we performed the
same experiments in a homogeneous and well-stirred environment and did not observe any
fitness difference as a function of size.
219

With two environments, one favoring single cells and one favoring clusters, we could finally test
whether fluctuation between them would favor regulated life cycles. We therefore competed our
inducible strain against single cells or clusters in an alternating selection regime. Cells were grown
for one cycle of growth in low sucrose followed by a cycle in the glucose emulsion before being
transferred back in sucrose for the cycle to start again. To simulate a regulated life cycle, we
induced our *ACE2* inducible strain to break the clusters into single cells before they entered the
emulsion and allowed them to grow as clusters for the remainder of the competition experiment
(Figure 4C). Figure 4D shows that this fluctuating environment caused regulated life cycles to
outcompete both single-celled and multicellular life cycles, confirming our hypothesis. Directly
competing the 3 life cycles together, also confirms that the regulated life cycle outcompetes the
two other life cycles (S4E).
231
232 Taken together, these fitness assays show that the outcome of competition between clusters and
233 single cells is determined by the environment: in the homogenous, low sucrose environment,
234 clusters are fitter, but in a patchy resource environment, it is single cells that are fitter. Most
235 provocatively, our results suggest that alternating selection for dispersal and multicellularity favors
236 regulated life cycles.
237
Regulated, multicellular life cycles have evolved, independently, in many branches of the tree of
life, including the fungi[1,4,13]. Many of these cycles pass through single-celled intermediates. We
examined wild isolates of the budding yeast, *Saccharomyces cerevisiae*, and found that they form
clonal multicellular clusters whose size is regulated by the mating type locus and their
environment. These two forms of regulation suggested that some isolates of this species may use
regulated multicellularity to increase their fitness in addition to the well characterized alternation
between haploid and diploid states and the formation of spores to survive starvation and
desiccation and promote dispersal. This hypothesis prompted us to ask if a regulated
multicellularity confers an advantage over being constitutively single-celled or constitutively
multicellular. We engineered a lab strain to exist in three states: constitutively single-celled,
constitutively multicellular and capable of an environmentally regulated life cycle. Competing
these strains with each other reveal that the single-celled life cycle is favored in a patchy resource
environment, multicellular clusters are favored in an environment requiring metabolic cooperation,
and that an alternation between these two environments favors a regulated life cycle.
252
Clonal multicellularity in yeast is due to the failure to degrade the primary septum, the specialized
portion of the cell wall that holds mother and daughter cells together after cytokinesis has
separated their cytoplasms. The increased cluster size in haploids strains suggests that the
expression of haploid specific genes may hinder the production of enzymes that degrade the
primary septum. For example Ste12, a haploid specific transcription factor, is required for the
expression of *AMN1*, a gene known to inhibit cell separation and the binding of the Mata1/Mata2
heterodimer prevents Ste12 from accessing the *AMN1* promoter in diploid cells[27].
260
261 Clonal multicellularity has an intrinsic advantage, resistance to cheating, over the agglomerative
262 multicellularity exhibited by slime molds and some bacteria[1,2]. However, the group formation
263 mechanisms constrain clonal multicellularity's ability to rapidly switch from single cell to

multicellularity. Under the fluctuating regime selection described here, we would predict that aggregative multicellularity carries an advantage as the switch from single cell to multicellularity can happen in less than a generation.

Most natural isolates of *S. cerevisiae* are diploid and these diploids are primarily formed by mating between sister spores produced from the same meiosis. The budding pattern of haploid cells has been proposed to favor the mating that occurs when a germinated spore lacks an appropriate partner and uses mating type switching to produce descendants that have opposite mating types[34,35]. By forming multicellular clusters, these clonally related cells would still be able to mate, even in environments where fluid flow and other physical forces would otherwise separate the cells.

In a single environment, the clonal clusters produced by experimental evolution or found in natural isolates usually break into smaller clusters rather than giving rise to populations of single cells. This mode of reproduction is favored in environments where metabolic cooperation or direct selection for cluster size gives clusters higher fitness than single cells. But if the environment alternates between favoring clusters and single cells, life cycles that regulate propagule size in response to the environmental fluctuation should be favored. We created this fluctuation by alternating between an environment that selected for metabolic cooperation and one that favored dispersal in a patchy environment. Our engineered strains add evidence that experimentally constructed, alternating environments can select for a regulated life cycle[22,36]. Outside the lab, natural oscillations, such as tidal, seasonal, or ecological fluctuations could have selected for the evolution of regulated life cycle. Examining wild isolates in their natural habitats and correlating their phenotype with fluctuations in their environments, over time and space, may reveal that *S. cerevisiae* is capable of more sophisticated and ecologically adaptive life cycles than have been discovered so far.

Acknowledgements:

The authors thank members of the Murray lab, especially Thomas LaBar and Tanush Jagdish for discussions, John Koschwanez for help with developing the project and Michael Desai, Michael Laub, Silvia De Monte, and William Ratcliff for their feedback on the manuscript. J.B, P.N and AWM are supported by the grant NIH/NIGMS R01 GM043987. A.W.M. is also supported by the NSF-Simons Center for the Mathematical and Statistical Analysis of Biology (NSF #1764269 , Simons #594596).

Author contributions:

Conceptualization, J.B., and A.W.M.; Methodology, J.B., and A.W.M.; Investigation, J.B. and P.N.; Writing – Original Draft, J.B.; Writing – Review & Editing, J.B., P.N., and A.W.M.; Funding Acquisition, A.W.M; Supervision, A.W.M.

Declaration of interests:

The authors declare no competing interests.

Main-text Figure Legends:

See below figures.

STAR Methods:

RESOURCE AVAILABILITY:

Lead Contact:

Further information and requests for resources and reagents should be directed to and will be
fulfilled by the Lead Contact, Andrew Murray (awm@mcb.harvard.edu).

Materials availability:

Yeast strains created in this study are available upon request from the Lead Contact.

Data and code availability:

All data and analysis codes in this study are available upon request from the Lead Contact.

EXPERIMENTAL MODEL AND SUBJECT DETAILS:

Yeast strains and media:

All experiments with the engineered strains were conducted in minimal (no amino acids or
nucleotides) synthetic media made by combining refrigerated stocks of 10X Yeast Nitrogen Base
(with ammonium sulfate) from BD Difco™ with the appropriate sugar (sucrose stocks were kept
at -20°C). Once made, the minimal synthetic media was stored at 4°C and protected from light.
β-estradiol stocks were made at a concentration of 10mM in ethanol and stored at -20°C. To
obtain the final working concentration for the experiments, the β-estradiol stock was diluted in
ultrapure water. Experiments involving the wild isolates experiments were performed in rich
medium, YEP (1% Yeast-Extract, 2% Peptone) to which we added dextrose, sucrose, ethanol,
galactose or glycerol at the appropriate concentration. YPD is YEP containing 2% w/v glucose.

Engineered strains: Engineered strains were constructed by first inserting the LexA-ER-AD
system at the *HIS3* locus as described by Ottoz et al.[31]. We then replaced the *ACE2* promoter
with an inducible promoter containing 1 LexA binding site to create the inducible strain. *ACE2* was
deleted to create the positive control: the constitutively multicellular strain. Nothing was changed
for the negative control: the constitutively single-celled strain. Each of these three strains were

marked with a different fluorescent protein, expressed from the *ACT1* promoter, to allow
competition essays.
353
Original wild isolates: The wild isolates used are from the SGRP-2 collection created by the Liti
lab[12]. Before performing the experiments, we confirmed the mating type of all the strains used
both by mating and by PCR[37]. Inconsistencies between reported and observed genotypes of the
haploids strains were corrected by combining mating type switching and sporulation. We created
a new set of diploids by mating the two heterothallic haploids, allowing us to have isogenic
haploids and diploids. We used all the strains from the collection that we could make available as
diploids, and both *MAT***a** and *MAT*α haploid*s*: 273614N, BC187, DBVPG1106, DBVPG1373,
DBVPG6044, 322134S, L_1374, L_1528, NCYC110, SK1, 378604,X UWOPS05-217.3, S288c,
UWOPS83-787.3, UWOPS87-2421, Y12, Y55, YIIc17_E5, YJM975, YJM978, YJM981, YPS128,
and YPS606.
364
365 *Homozygosing the* MAT *locus:* to create diploids homozygous at the mating type locus, we
366 transformed diploids with a plasmid carrying the *HO* gene under an inducible promoter[38]. Colonies
367 were selected on uracil plates and allowed to lose the plasmid on YPD before being tested for
368 mating type both by mating and PCR using methods described here[37].
369
370 Table S1 lists laboratory yeast strains used in this study:
371
372 All strains were derived from yJK089, a derivative of the W303 strain background
373

Strain Name	Purpose	Genotype
yJB064	Cluster forming strain	*can1-100 HIS3Δ::prACT1-yCerulean-tADH1-His3MX6 BUD4-S288C ACE2Δ::bleMX*
yJB077	Reference strain for competition experiment	*can1-100 HIS3Δ::prACT1-ymCherry-tADH1-His3MX6 BUD4-S288C*
yJB128	Single-celled strain with β-estradiol construct.	*can1-100 his3-11,15::PACT1(-1-520)-LexA-ER-haB42-TCYC1 BUD4-S288C HOΔ::prACT1-yCerulean-ADH1tr-KanMX*
yJB130	Inducible strain with β-estradiol construct.	*can1-100 his3-11,15::PACT1(-1-520)-LexA-ER-haB42-TCYC1 BUD4-S288C prACE2Δ:: PTEF-HygMX-TTEF-insul-(lexA-box)1-PminCYC1 HOΔ::prACT1-ymCherry-ADH1tr-KanMX*

yJB136	Single-celled strain with β-estradiol construct.	*can1-100 his3-11,15::PACT1(-1-520)-LexA-ER-haB42-TCYC1 BUD4-S288C HOΔ::prACT1-ymCitrine-ADH1tr-KanMX*
yJB138	Cluster forming strain with β-estradiol construct.	*can1-100 his3-11,15::PACT1(-1-520)-LexA-ER-haB42-TCYC1 BUD4-S288C HOΔ::prACT1-ymCitrine-ADH1tr-KanMX ACE2Δ::HphMX*

METHOD DETAILS

Competition essays:

All competition experiments were performed in YNB complemented with the appropriate carbon source. Experiments were performed in plastic culture tubes. Fluorescently labeled strains were individually pre-grown overnight in media in which the competition experiment would be performed. Strains were then mixed in 1:1 ratios and diluted in PBS before being inoculated into the appropriate medium. Cultures were passaged for three cycles of growth and frequency of each strain was measured at each transfer with a flow cytometer (Fortessa, BD Bioscience, RRID:SCR_013311, US). Because the distribution of number of cells per cluster was stable from one transfer from another we were able to estimate the relative fitness between single cells and clusters by extracting the ratio between fluorescent events (an event here can be a single cell, or a cluster of any size). The frequency of each strain was extracted with the CytoExploreR[39] package. To quantify the relative fitness we performed a linear regression between the number of generation elapsed between each transfer and the log of the ratio between the two genotypes. The relative fitness is the slope of the regressed line. Errors bars reflect the standard error of the mean from at least 3 independent replicates. Note that the reference strain varies for different experiments and is indicated for each plot.

Emulsions:

Emulsions were produced by mixing 150 ul of culture with 350 ul of FC-40 oil containing 2% (w/v) 008-FluoroSurfactant (Ran Biotechnologies) in centrifuge tubes (VWR 76332-074). Note that emulsion stability is influenced by the type of centrifuge tube used. The tubes were then gently tapped before being vortexed at max speed for 30 sec. The tubes were allowed to rest before being opened and sealed with parafilm for the 48h incubation time. To break the emulsion, the tubes were first spun down with a counter top centrifuge, for 30 sec, the lower oil phase was removed before the addition of 100 ul of 1H,1H,2H,1H-Perfluoro-1-octanol (Sigma Aldrich) mixed by pipetting up and down and gently turning the tube. After 10 min of rest, tubes were spun down for 30 sec on a counter top centrifuge, the lower phase was removed before gently mixing the aqueous phase and recovering the cells by pipetting.

409　Cycling experiments:
410
For the cycling experiments (Figure 4C), the strains were first pre-grown overnight in YNB with
100 mM glucose before being mixed in a 1:1 ratio and diluted 1:100 in YNB with 100 mM glucose
to start the first step of the cycle. After 24h of incubation at 30°C on a roller drum, culture were
diluted 1:10,000 in sucrose for the second step and incubated for 4 days. For the third step, the
cells were diluted 1:100 in 100 mM glucose and β-estradiol was added for induction when needed.
After 24h, the culture was diluted 1:10,000 and transferred into emulsion and allowed to grow to
saturation before the emulsion was broken (as described above) and the diluted 1:100 in YNB
with 100 mM glucose to restart the cycle. The frequency of each strain was measure at each
transfer using flow cytometry.
420
421　Size measurements on wild isolates:
422
A colony of each wild isolates was inoculated in 300 ul of YPD in a 2 ml deep 96 well plate and
allowed to grow overnight on a shaker at 1000 rpm. Cultures were then diluted 1:100 in the
appropriate media and incubated for 48h before being transferred again 1:100 in the same media.
Samples were measured in early stationary phase, after 24h or 48 depending on the media. To
prevent flocculation, cultures were diluted in EDTA to a final concentration of 100 mM. Each of
the 3 replicates of these measurement was perform as an independent batch.
429
430　Clustering Score:
431
To allow us to compare the wild isolates size measurements we developed a clustering score
allowing us to quantify the clustering level of each strain and to compare between different strains
and different environment (Figure 1A). Cultures of strains that form clusters are composed of
clusters of different sizes but also of single cells. We therefore divide the forward scatter
measurements by the mean of the 10% smallest forward scatter values of a population (usually
mostly single cells), which corrects for differences in cell size between environments and strains.
We then take the mean of the 10% largest corrected forward scatter values as the clustering
score. This score is used as a relative value to compare the formation of clusters by different
strains in different environments.
441
442　Microscopy:
443
All images of cells and clusters were taken in a 96-well glass-bottomed plate (Greiner bio-one,
www.gbo.com) using a Nikon Ti inverted microscope (www.nikoninstruments.com) with
MetaMorph software (www.metamorph.com). Contrast of images was adjusted, and images were
annotated with scale bars using the Fiji distribution of ImageJ[40]. Contrast was changed for visibility
only and does not impact the results. Images of the emulsion for droplet size quantification were
taken in a hemocytometer (Bulldog Bio).
450
451
452

453 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
β-estradiol	SIGMA ALDRICH INC	Cat#E8875
Tryptone to 2xYT Broth	Difco™	Cat#90004-150
1H,1H,2H,1H-Perfluoro-1-octanol	SIGMA ALDRICH INC	Cat#370533
Surfactant	Ran Biotechnologies	Cat#008-FluoroSurfactant
Fluorinert™ FC-40	SIGMA ALDRICH INC	Cat#F9755
Experimental models: Organisms/strains		
Laboratory Yeast strains	See Table S1	N/A
Wild isolates	[13]	SGRP-2
Recombinant DNA		
Plasmids	[31]	FRP1639 & FRP1642
Software and algorithms		
Fiji	[40]	https://imagej.net/software/fiji/
CytoExploreR	[39]	https://dillonhammill.github.io/CytoExploreR/

455
456 **Supplementary data legends:**

457
458 See bellow figures.

459
460 **References:**

461
462

1. Grosberg, R.K., and Strathmann, R.R. (2007). The Evolution of Multicellularity: A Minor Major
Transition? Annual Review of Ecology, Evolution, and Systematics *38*, 621–654.
10.1146/annurev.ecolsys.36.102403.114735.

2. Brunet, T., and King, N. (2017). The Origin of Animal Multicellularity and Cell Differentiation.
Developmental Cell *43*, 124–140. 10.1016/j.devcel.2017.09.016.

3. Knoll, A.H. (2011). The Multiple Origins of Complex Multicellularity. Annual Review of Earth
and Planetary Sciences *39*, 217–239. 10.1146/annurev.earth.031208.100209.

4. BONNER, J.T. (1993). Life Cycles: Reflections of an Evolutionary Biologist (Princeton
University Press).

5. Kirk, D.L. (2005). A twelve-step program for evolving multicellularity and a division of labor.
Bioessays *27*, 299–310. 10.1002/bies.20197.

6. Ratcliff, W.C., Denison, R.F., Borrello, M., and Travisano, M. (2012). Experimental evolution
of multicellularity. PNAS *109*, 1595–1600. 10.1073/pnas.1115323109.

7. Koschwanez, J.H., Foster, K.R., and Murray, A.W. (2013). Improved use of a public good
selects for the evolution of undifferentiated multicellularity. eLife *2*, e00367.
10.7554/eLife.00367.

8. Wahl, M.E., and Murray, A.W. (2016). Multicellularity makes somatic differentiation
evolutionarily stable. PNAS *113*, 8362–8367. 10.1073/pnas.1608278113.

9. Kirk, M.M., Stark, K., Miller, S.M., Muller, W., Taillon, B.E., Gruber, H., Schmitt, R., and Kirk,
D.L. (1999). regA, a Volvox gene that plays a central role in germ-soma differentiation,
encodes a novel regulatory protein. Development *126*, 639–647. 10.1242/dev.126.4.639.

10. Dudin, O., Wielgoss, S., New, A.M., and Ruiz-Trillo, I. (2022). Regulation of sedimentation
rate shapes the evolution of multicellularity in a close unicellular relative of animals. PLoS
biology *20*, e3001551.

11. Oud, B., Guadalupe-Medina, V., Nijkamp, J.F., Ridder, D. de, Pronk, J.T., Maris, A.J.A. van,
and Daran, J.-M. (2013). Genome duplication and mutations in ACE2 cause multicellular,
fast-sedimenting phenotypes in evolved Saccharomyces cerevisiae. PNAS *110*, E4223–
E4231. 10.1073/pnas.1305949110.

12. Cubillos, F.A., Louis, E.J., and Liti, G. (2009). Generation of a large set of genetically
tractable haploid and diploid Saccharomyces strains. FEMS Yeast Res *9*, 1217–1225.
10.1111/j.1567-1364.2009.00583.x.

13. van Gestel, J., and Tarnita, C.E. (2017). On the origin of biological construction, with a focus
on multicellularity. PNAS *114*, 11018–11026. 10.1073/pnas.1704631114.

14. Boraas, M.E., Seale, D.B., and Boxhorn, J.E. (1998). Phagotrophy by a flagellate selects for
colonial prey: A possible origin of multicellularity. Evolutionary Ecology *12*, 153–164.
10.1023/A:1006527528063.

15. Tong, K., Bozdag, G.O., and Ratcliff, W.C. (2022). Selective drivers of simple
multicellularity. Current Opinion in Microbiology *67*, 102141. 10.1016/j.mib.2022.102141.

16. Black, A.J., Bourrat, P., and Rainey, P.B. (2020). Ecological scaffolding and the evolution of
individuality. Nat Ecol Evol, 1–11. 10.1038/s41559-019-1086-9.

17. Pichugin, Y., Peña, J., Rainey, P.B., and Traulsen, A. (2017). Fragmentation modes and the
evolution of life cycles. PLOS Computational Biology *13*, e1005860.
10.1371/journal.pcbi.1005860.

18. Pichugin, Y., and Traulsen, A. (2020). Evolution of multicellular life cycles under costly
fragmentation. PLOS Computational Biology *16*, e1008406. 10.1371/journal.pcbi.1008406.

19. Pichugin, Y., Park, H.J., and Traulsen, A. (2019). Evolution of simple multicellular life cycles
in dynamic environments. Journal of The Royal Society Interface *16*, 20190054.
10.1098/rsif.2019.0054.

20. Staps, M., van Gestel, J., and Tarnita, C.E. (2019). Emergence of diverse life cycles and life
histories at the origin of multicellularity. Nature Ecology & Evolution, 1. 10.1038/s41559-
019-0940-0.

21. Ispolatov, I., Ackermann, M., and Doebeli, M. (2012). Division of labour and the evolution of
multicellularity. Proceedings of the Royal Society B: Biological Sciences *279*, 1768–1776.
10.1098/rspb.2011.1999.

22. Hammerschmidt, K., Rose, C.J., Kerr, B., and Rainey, P.B. (2014). Life cycles, fitness
decoupling and the evolution of multicellularity. Nature *515*, 75–79. 10.1038/nature13884.

23. Ratcliff, W.C., Herron, M.D., Howell, K., Pentz, J.T., Rosenzweig, F., and Travisano, M.
(2013). Experimental evolution of an alternating uni- and multicellular life cycle in
Chlamydomonas reinhardtii. Nature Communications *4*, 2742. 10.1038/ncomms3742.

24. Ratcliff, W.C., Fankhauser, J.D., Rogers, D.W., Greig, D., and Travisano, M. (2015). Origins
of multicellular evolvability in snowflake yeast. Nature Communications *6*, 6102.
10.1038/ncomms7102.

25. Koschwanez, J.H., Foster, K.R., and Murray, A.W. (2011). Sucrose Utilization in Budding
Yeast as a Model for the Origin of Undifferentiated Multicellularity. PLOS Biology *9*,
e1001122. 10.1371/journal.pbio.1001122.

26. Yvert, G., Brem, R.B., Whittle, J., Akey, J.M., Foss, E., Smith, E.N., Mackelprang, R., and
Kruglyak, L. (2003). Trans -acting regulatory variation in Saccharomyces cerevisiae and the
role of transcription factors. Nature Genetics *35*, 57–64. 10.1038/ng1222.

27. Fang, O., Hu, X., Wang, L., Jiang, N., Yang, J., Li, B., and Luo, Z. (2018). Amn1 governs
post-mitotic cell separation in Saccharomyces cerevisiae. PLoS Genet *14*, e1007691.
10.1371/journal.pgen.1007691.

28. Liti, G., Carter, D.M., Moses, A.M., Warringer, J., Parts, L., James, S.A., Davey, R.P.,
Roberts, I.N., Burt, A., Koufopanou, V., et al. (2009). Population genomics of domestic and
wild yeasts. Nature *458*, 337–341. 10.1038/nature07743.

29. Stratford, M. (1989). Yeast flocculation: Calcium specificity. Yeast *5*, 487–496.
10.1002/yea.320050608.

30. Coelho, S.M., Peters, A.F., Charrier, B., Roze, D., Destombe, C., Valero, M., and Cock, J.M.
(2007). Complex life cycles of multicellular eukaryotes: New approaches based on the use
of model organisms. Gene *406*, 152–170. 10.1016/j.gene.2007.07.025.

31. Ottoz, D.S.M., Rudolf, F., and Stelling, J. (2014). Inducible, tightly regulated and growth
condition-independent transcription factor in Saccharomyces cerevisiae. Nucleic Acids Res
42, e130. 10.1093/nar/gku616.

32. De Monte, S., and Rainey, P.B. (2014). Nascent multicellular life and the emergence of
individuality. J Biosci *39*, 237–248. 10.1007/s12038-014-9420-5.

33. Lavrentovich, M.O., Koschwanez, J.H., and Nelson, D.R. (2013). Nutrient shielding in
clusters of cells. Phys. Rev. E *87*, 062703. 10.1103/PhysRevE.87.062703.

34. Wang, Y., Lo, W.-C., and Chou, C.-S. (2017). A modeling study of budding yeast colony
formation and its relationship to budding pattern and aging. PLOS Computational Biology
13, e1005843. 10.1371/journal.pcbi.1005843.

35. Nasmyth, K.A. (1982). Molecular genetics of yeast mating type. Annu Rev Genet *16*, 439–
500. 10.1146/annurev.ge.16.120182.002255.

36. Beaumont, H.J.E., Gallie, J., Kost, C., Ferguson, G.C., and Rainey, P.B. (2009).
Experimental evolution of bet hedging. Nature *462*, 90–93. 10.1038/nature08504.

37. Huxley, C., Green, E.D., and Dunham, I. (1990). Rapid assessment of S. cerevisiae mating
type by PCR. Trends Genet *6*, 236. 10.1016/0168-9525(90)90190-h.

38. [8] Putting the HO gene to work: Practical uses for mating-type switching - ScienceDirect
https://www-sciencedirect-com.ezp-
prod1.hul.harvard.edu/science/article/pii/007668799194011Z?via%3Dihub.

39. Interactive Analysis of Cytometry Data https://dillonhammill.github.io/CytoExploreR/.

40. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676–682. 10.1038/nmeth.2019.

592   In progress

594   **Figures:**

596
597

598   *Figure 1: Wild S. cerevisiae isolates grow as multicellular clusters in their haploid state.*

599   *See also Figure S1.*

A) Description of the clustering score. FSC stands for flow cytometry "Forward SCatter". Scale bar represents 25 μm. Data for the single-celled diploid and haploid are prototrophic W303 strains. The haploid cluster is the evolved cluster Evo2 from Koschwanez et al.

B) Clustering score measurements of the wild isolates grown in YPD. Haploids MATa and MATα are statistically different from the diploids (p-value < 1e-13), while haploids are not different from one another (p-value = 0.3) (Wilcoxon test). Error bars represent the standard error of the mean for 3 independent replicates.

C) Representative DIC (Differential Interference Contrast) images and forward scatter (FSC) distributions of a subset of the strains. Scale bar on the images represents 25 μm.

*D) Clustering score measurements for a subset of strains comparing haploid, diploids and diploids engineered to be homozygous at the mating type locus. Error bars represent the standard error of the mean for 3 independent replicates and p-values are the results of a Wilcoxon test: ****: p < 1e-4 for comparisons against the a/α diploid.*

Figure 2: Environment modulates multicellularity in wild S. cerevisiae isolates:

See also Figure S2.

A) Clustering score for diploid isolates. Environments are hierarchically grouped by clustering
score similarity. Strains were grown in yeast extract and peptone (YEP) complemented with 5
different carbon source (concentration of 2% (w/v)).

B) Representative DIC images of selected diploid strains. Scale bar represents 50 μm

C) Same as A but for MATa strains. See Figure S2 for MATα heatmap.

D) Same as B but for MATa strains

624
625

Figure 3: Regulated life cycles can be engineered

See also Figure S3.

A) Cartoon representation of the different engineered life cycles, Single-celled life cycle:
reproduction as single cells (most lab strains) . Multicellular life cycle: multicellular yeast clusters
that reproduce by breaking into smaller clusters. Regulated life cycle: a strain engineered to be
able to reproduce either as a single cell, or a multicellular cluster.

B) To create a regulated life cycle, we placed the ACE2 gene under a promoter regulated by β-
estradiol. In the absence of β-estradiol, ACE2 is not transcribed, and the strain reproduces as
clusters. Increasing the β-estradiol concentration produces a single-celled life cycle.

C) Mean forward scatter of the ACE2-inducible strain grown in yeast nitrogen base (YNB) with 10
mM glucose at increasing concentration of β-estradiol. Error bars represent the standard error of
the mean for 3 independent replicates. Scale bar represents 50 μm.

D) Kinetics of cluster formation and dissolution on the removal or addition of β-estradiol. Error
bars represent the standard error of the mean for 3 independent replicates.

641
642
643

Figure 4: Resource availability controls the fitness differences between different life cycles.

See also Figure S4.

A) Competition experiment between the ACE2-inducible strain and a constitutively single-celled reference strain in YNB with 10 mM sucrose (blue) or YNB with 10 mM glucose and 10 mM fructose (red) as a control. Each point shows the fitness of the inducible strain relative to the constitutively single-celled strain at a given ACE2 induction level. FSC-A stands for Forward SCatter Area and correlates with cluster size. Competitions performed for 3 cycles of growth and dilution. Error bars represent the standard error of the mean for 3 independent replicates.

B) Competition experiment between the ACE2-inducible strain and a constitutively single-celled reference strain in YNB with 100 mM glucose in emulsion (green) or in bulk as a control (grey).

Each point shows the fitness of the inducible strain relative to the constitutively single-celled strain
at a given ACE2 induction level. Error bars represent the standard error of the mean for 3
independent replicates. Cells were seeded at density that corresponded to a mean of less than 1
cell per drop of the emulsion. Competitions performed for 3 cycles of growth and dilution.

C) Protocol for the fluctuating selection regime. Cells are first grown for a cycle in YNB with 10
mM sucrose before being transferred to YNB with 100 mM glucose and induced with β-estradiol
for 24 hours. After induction, cells are put in an emulsion of YNB with 100 mM glucose for 48h.
The emulsion is then broken, and cells are allowed to recover in 100 mM glucose YNB before the
cycle starts again. Error bars represent the standard error of the mean for 3 independent
replicates.

D) Competition experiment in the fluctuating selection regime between the ACE2-inducible strain
and strains that are either constitutively single-celled or constitutively form clusters. Error bars
represent the standard error of the mean for 3 independent replicates. Competitions were
performed for 2 full selection cycles.

673
674 *Supplementary Figure 1:*

675 *A) correlation between bead cross-sectional area and mean forward scatter measurements.*
676 *FSC for each bead size was measured in 3 independent measurements.*

677 *B) Clustering score in YPD of all 23 strains. The indicated p-values are the results of Wilcoxon*
678 *test.*

Supplementary Figure 2:

A) *Clustering score for haploid MATα isolates. Environments are hierarchically grouped. Strains were grown in YEP complemented with 5 different carbon source (concentration of 2% (w/v)).*

B) *Same as A but for diploid strains in sugar concentration of 0.2%.*

C) *Same as A but for MAT**a** strains in sugar concentration of 0.2%.*

D) *Same as A but for MATα strains in sugar concentration of 0.2%.*

690
691

Supplementary Figure 3:

A) Mean forward scatter of the engineered, ACE2- inducible strain in glucose + fructose or
sucrose at increasing concentration of β-estradiol. Error bars represent the standard error of the
mean for 3 independent replicates.

B) DIC images of the ACE2-inducible strain when uninduced or induced with 10 nM of β-
estradiol. Cells were grown in YNB with 10 mM sucrose or YNB with 10 mM glucose and 10 mM
fructose. Scale bar represents 50 μm.

Supplementary Figure 4:

A) Competition experiment between an ace2Δ cluster and a single cell in different concentrations
of sucrose or glucose + fructose. These results confirm work by Koschwanez et al. (2013) and
quantify the fitness advantage of clusters in sucrose. Error bars represent the standard error of
the mean for 3 independent replicates.

B) Similar to A but competitions were done in 10 mM sucrose and 10 mM Glucose + 10 mM
Fructose with different dilution factors between transfers (100 stands for a 1:100 dilution). A large
dilution factor is needed to observe fitness differences between single cells and clusters. Error
bars represent the standard error of the mean for 3 independent replicates.

C) Measurements of ratio of single cells to clusters before introducing them into an emulsion and
after immediately breaking the emulsion, with no intervening growth, to confirm that the emulsion
protocol does not create a bias in favor of clusters or single cells.

D) Size distribution of emulsion droplets. The emulsion was imaged and droplet diameters were
measured using Fiji.

715