Main Manuscript for

Neuronal Morphology Enhances Robustness to Perturbations of
Channel Densities

Yunliang Zang, Eve Marder

Volen Center and Department of Biology, Brandeis University, Waltham, MA 02454, USA.

*Eve Marder

Email: marder@brandeis.edu

Author Contributions: Y.Z. designed this project; Y.Z. implemented model simulations, data analysis, and figure visualization; Y.Z. and E.M. wrote the manuscript.

Competing Interest Statement: The authors declare no competing interest.

Keywords: compartmental models; ion channels; individual variability; neuronal resilience; parameter space

This PDF file includes:

      Main Text
      Figures 1 to 6

Abstract

Biological neurons show significant cell-to-cell variability but have the striking ability to maintain their key firing properties in the face of unpredictable perturbations and stochastic noise. Using a population of multi-compartment models consisting of soma, neurites, and axon for the lateral pyloric (LP) neuron in the crab stomatogastric ganglion, we explored how rebound bursting is preserved when the 14 channel conductances in each model are all randomly varied. The soma-axon coupling is critical for the ability of the axon to spike during bursts and consequently determines the set of successful solutions. When the coupling deviates from a biologically realistic range, the neuronal tolerance of conductance variations is significantly lessened. Thus, the gross morphological features of these neurons enhance their robustness to perturbations of channel densities and expands the space of individual variability that can maintain a desired output pattern.

1   **Main Text**
2
Introduction
It is now well-established that multiple sets of maximal conductances of ion channels can produce
very similar neuronal activity patterns in both individual neurons and circuits (1-9). Nonetheless, it
is also clear that models of desired intrinsic properties are relatively rarely found by randomly
sampling within a space of parameters (5, 6). Thus, it is likely that biological neurons use
homeostatic and other developmental mechanisms to successfully find rare sets of values of ion
channel conductances that give rise to specific patterns of intrinsic excitability (10-14).

Although single-compartment, isopotential, conductance-based models have been
enormously useful in understanding the relationships among ion channels that give rise to
electrical excitability, most biological neurons have complex structures, with differential
distributions of channels in the membranes of different neuronal regions. While there are specific
examples of conductance-based models of complex neurons (15-20), these models have often
been finely tuned to achieve their desired activity patterns. Thus, it is less clear in these more
complex models the extent to which complex structures aid or diminish the likelihood of finding
vastly different degenerate sets of conductance values that support characteristic activity
patterns.

In this study, we exploit a previously developed 4-compartment model database of the Lateral
Pyloric (LP) neuron of the crustacean stomatogastric ganglion (STG) to look at the influence of
LP neuron structure on its robustness to perturbations of its channel densities. This is partly
motivated by recent studies that show that the detailed neuropilar projections of the LP neuron
have relatively little impact on how slow synaptic events are processed because of the large
length constants of the major neurites (21-23). In these 4-compartment models (soma, axon, and
two neurite compartments) only the axon compartment has voltage-gated Na^+ channels (5). To
create this model database, Taylor et al (5) generated ~ 600,000 randomly selected sets of
conductance densities and then sorted them to find ~1,300 models with the desired firing
patterns matching the biological properties of the LP neuron. Specifically, when isolated, the
biological LP neuron fires tonically, but shows robust rebound bursts in response to inhibitory
inputs from its presynaptic partners. These rebound bursts are important in ensuring that the LP
neuron fires at an appropriate time during the triphasic pyloric rhythm (24).

The successful ~1,300 models vary considerably in their maximal conductances and are
examples of degenerate or multiple solutions that produce a characteristic behavior. In this paper,
we look at the effect of neuronal structure on the robustness of this population of LP neuron
models in response to perturbations of conductance densities.

Results

Figure 1A shows a diagram of the chemical synaptic inputs to the LP neuron. The top trace in Figure 1B is an intracellular recording from the LP neuron, with its sequential inhibition from the PY (Pyloric), PD (Pyloric Dilator) and AB (Anterior Burster) neurons (note that AB and PD slow waves and spikes are synchronized), followed by the rebound burst in the LP neuron. The timing of the inhibitory inputs is shown by the extracellular recordings from the lvn (lateral ventricular nerve, Figure 1B, bottom). The LP neuron somatic action potentials that ride on the top of the slow wave are highly attenuated by the LP neuron's cable, as they are generated in the distant axon.

Figure 1C shows the composition of ion channels in each of the 4 compartments in the model. The soma has a leak current (I_{leak}), a modulator current (I_{MI}), three potassium currents (I_{Kd}, delayed-rectifier K^+ current; I_A, A-type K^+ current; I_{KCa}, calcium-activated K^+ current), a calcium current (I_{Ca}) and a hyperpolarization-activated inward current (I_h). In addition to these currents, three inhibitory synaptic currents, from the AB, PD, and PY neurons (synAB, synPD and synPY) are also distributed in the neurites. The axon has a I_{leak}, I_{Kd}, I_A, and a voltage-gated Na^+ current (I_{Na}). A characteristic waveform from all LP neuron models is seen in Figure 1D. The top trace is the model behavior in the soma compartment, while the bottom trace is the model behavior in the axon. The axon generates overshooting action potentials, and only a very small slow wave is visible, while the soma compartment replicates the somatic recordings seen in the biological neurons (Fig. 1B).

Neuronal Robustness to Channel Conductance Variations.

We used 1292 of Taylor's LP neuron models (5), hereafter referred to as "parent" models whose parameters show large cell-to-cell variations despite similar output patterns. We first varied all of the 14 channel conductances at the same time by different random values in each "parent" model, based on their range of values for all of the 1292 models. Then, for each "parent" model, multiple runs, with different sets of channel perturbations, we generated a number of "child" models, which displayed a variety of intrinsic patterns of activity (Fig. 2A; methods).

The "child" models showed a variety of behaviors, some of which can be generally classified

as tonic spiking, rebound bursting, aperiodic spiking, depolarization block or non-spiking (Fig.

2A). The simulation results of 100 trials for each "parent" model are summarized in Figure 2B. For

each "parent" model, the ratio of "child" models that are also bursters with at least 2 spikes/burst

was calculated to capture its robustness to channel conductance variations. A higher ratio

suggests the "parent" model is more robust to channel conductance variations. When channel

conductances were simultaneously varied within ±10% of their corresponding parameter ranges,

the rebound bursting pattern was remarkably robust to the conductance variations. For 989 of the

"parent" models over 90% of the "child" models generated rebound bursting and for 649 of the

"parent" models all the "child" models reliably generated rebound bursting. The overall high ratios

of "child" models being bursters indicates that the pattern of rebound bursting is relatively

insensitive to the conductance variations in these "parent" models. When channel conductances

were varied in larger ranges, more and more "parent" models showed decreased ratios of "child"

models being bursters, as reflected by their left-shifted distribution in the histograms (Fig. 2B).

It is always interesting to estimate the size and structure of the 14-dimension manifold in

which these models exist. Previous work on similar conductance-based models argued that

models with similar behavior are in connected regions of parameter space (3, 7). For many

"parent" models in these previous simulations (n = 649), the perturbation-generated "child"

models reliably generate rebound bursting, suggesting that there are reasonably sized areas

around those "parent" models with relatively consistent behaviors. We estimate that the size of

the manifold in each channel conductance direction is at least 20% of their corresponding

conductance ranges.

Therefore, we selected 37 elite "parent" models whose "child" models reliably generate

rebound bursting for 100 trials of perturbations and have 3 – 11 spikes per burst when the

variation range is ±10%. Then, we challenged each of them with 10,000 trials of perturbations.

For 23 models, all their "child" models reliably generated rebound bursting and in the other 14

models the lowest ratio of the "child" models generating rebound bursting was 98.95%. These

results again support the interpretation that this bursting behavior is relatively insensitive to

random channel conductance variations. We plotted the spike waveforms and channel

conductances for 8 of the 23 models in Figure 3. Except for larger values of the PY neuron

synaptic input, the other channel conductance values spread broadly within their corresponding

conductance ranges, suggesting that robust models exist widely in the 14-dimension parameter

space.

If one group of neurons shows an advantage in maintaining rebound bursting relative to

another group within a small variation range, is this advantage preserved with larger variations?

When the variation range increases, the two groups of neurons have similar ratios of maintaining

the bursting pattern regardless of their ratios with smaller variation ranges (Fig. 4A).

We also explored the effect of varying individual channel conductance on the robustness of

rebound bursting. Compared with the effect of varying 14 channel conductances simultaneously,

the models are more resilient to single channel conductance variations. In the soma and neurites,

the model's performance shows modest sensitivity to I_A and I_{MI} conductances. In contrast, varying

each of the conductances in the axon changes the model's firing pattern more efficiently.

The Effect of Soma-axon Coupling on Axonal Spiking during Bursts

In the LP neuron, the soma and neurites generate rebound slow waves, that evoke bursts of Na^+

spikes in the axon. Therefore, the soma-axon coupling is key to regulating these bursts of spikes.

In the whole model database (n = 1292), we varied the axial resistivity, Ra, between the soma

and the axon to change their degree of coupling, while the Ra values in other compartments were

fixed at the control value of 100 Ωcm. When the soma to axon Ra was changed, some models no

longer produced rebound bursts (Fig. 5A). All models generated rebound bursting when the Ra

was in the range of 95 Ωcm – 125 Ωcm (Fig. 5B). For Ra in the range of 40 Ωcm – 300 Ωcm,

1175 (91%) models generated rebound bursting. These results suggest that the bursting pattern

is resilient to Ra changes around the control value of 100 Ωcm. When Ra deviated considerably

from the control value, the number of rebound bursting neurons decreased significantly. The

models that generate rebound bursting with larger Ra deviations are usually bursters when the

Ra deviation is lower. For example, in the range of 2 Ωcm – 100 Ωcm, if a model is a burster with

a smaller Ra, it is always a burster when the Ra has a larger value, with rare exceptions for the

Ra from 15 Ωcm to 20 Ω cm (2 exceptions) and from 35 Ωcm to 40 Ωcm (1 exception).

The simulation results above show that the degree of soma-axon coupling affects the

population size of bursting neurons. We selected 130 models that always generate the pattern of

rebound bursting for Ra within 2 Ωcm – 500 Ωcm. The number of spikes per burst reflects the

ability of the axon to spike during bursts. For most of these neurons, the number of spikes per

burst shows biphasic changes with Ra, as also reflected by the average changes (Fig. 5C).

However, in several neuron models, this number monotonically increases within the tested Ra

range.

When Ra is small, the soma and the axon are almost isopotential, as revealed by the nearly

overlapped somatic and axonal membrane potentials (Fig. 5D, left). When the soma and axon are

strongly coupled, this reduces the ability of the axon to spike during bursts for two reasons. First,

the axonal membrane potential after each spike is closer to the somatic slow-wave peak (bottom

Vm during the bursts), which reduces the recovery of Na^+ channels from inactivation. Second, the

soma exerts a larger capacitance load on the axon during spiking (top Vm during the bursts).

Therefore, when the Ra was 2 Ωcm, the number of spikes per burst was low. Increasing Ra to

make the soma and the axon more isolated should increase the axonal ability to spike during

bursts, considering just the above 2 factors. However, the increased isolation between the soma

and the axon also reduces the axial current from the somatic slow wave that triggers axonal

spiking during bursts. Consequently, changes in the number of spikes per burst depend on the

balance of these 3 factors when Ra changes (Fig. 5D).

The Effect of Soma-axon Coupling on Neuronal Robustness

We explored how soma-axon coupling affects neuronal robustness for all of the 1292 "parent"

models. When the soma to axon Ra is near the control value of 100 Ωcm, the average ratio of

"child" models that are bursters is insensitive to Ra changes irrespective of conductance variation

ranges (Fig. 6A1, Ra = 50, 100 and 125 Ωcm). However, when the soma-axon coupling gets too

strong (Ra = 2 Ωcm) or too weak (Ra = 500 Ωcm), the average ratio of "child" models that are

bursters is significantly reduced. Under these conditions, for many "parent" models that lose

rebound bursting, their "child" models become bursters when the channel conductances are

perturbed to appropriate ranges (Fig. 6A2, A3). When Ra is 2 Ωcm, the average ratio of "child"

models that are bursters shows biphasic changes. Under these conditions, some of the original

"parent" models are not bursting neurons, but in the neighboring parameter space they may

remain bursting (Fig. 6A2, A3). Figure 6 suggests that the "density" of bursting neuron models in

the whole parameter space is nearly unchanged when Ra is near 100 Ωcm but significantly

decreases when Ra deviates further from its control value.

To explore how the degree of soma-axon coupling affects the ability of individual neurons to

maintain rebound bursting when faced with channel conductance variations, we selected 131 elite

models from the 1292 "parent" models. These 131 elite "parent" models reliably generate

rebound bursting when the soma to axon Ra is 2 Ωcm, 100 Ωcm, and 500 Ωcm. Strikingly, for

each of these "parent" models, the ratio of "child" models that are bursters consistently decreases

when Ra is changed from 100 Ωcm to 2 Ωcm and from 100 Ωcm to 500 Ωcm (Fig. 6B, only 80

cells were plotted, but all cells show the same trend), regardless of the conductance variation

ranges. These simulation results suggest that the degree of soma-axon coupling determines the

size of the manifold that corresponds to bursting neurons and therefore the neuronal robustness

to channel conductance variations.

Discussion
166

Most neurons have complex structures over which ion channel distributions must be established

such that the neuron's characteristic electrical properties are maintained, given the inevitable

"noise" that must accompany the ongoing turnover of channels in the membrane. It is known that

some neurons display specific patterns of channel distributions that are thought to aid in the

neuronal computations that these neurons perform (25-27). Therefore, it becomes important to

determine the extent to which the spatial structure of neurons makes them more or less sensitive

to fluctuations in channel densities across their structures.

In this work, we used a population of LP neuron compartmental models developed previously

(5) to explore neuronal robustness to channel conductance variations. All members of this

population met a series of criteria that ensure that they capture many of the essential features of

the biological neurons whose behavior they were intended to capture (5). At the same time, these

models show considerable degeneracy in that the specific values of their conductances are highly

variable (5). While these models separate the soma, axon, and dendritic compartments, and are

therefore more biologically "realistic" than single compartment, point, neurons (28), they are not

intended to capture all of the detailed structures of individual STG neurons.

In models of one or two dimensions, it is easy to find the parameter sets that define the

boundary between different types of neuronal activity patterns (29, 30). However, as the number

of dimensions increases, it becomes more challenging to characterize the parameter space for

neurons with specific properties (7, 16). For the 14-dimension parameter space here, it would be

a daunting task to locate all possible bursting models with a grid search, because it is impossible

to know whether the sampling step size is small enough (3). Instead, we used random parameter

perturbations. This approach does not provide an unambiguous map of the entire manifold.

Nonetheless, if all "child" models are bursters after exerting random and simultaneous

conductance perturbations for sufficiently many trials, it argues the existence of a multi-dimension

manifold that contains the very large majority of models of similar behavior (7). We did this in a

subset of neuron models for 10,000 trials and most of them reliably generated rebound bursting

after perturbations (Fig. 3).

Multiple mechanisms likely exist that can allow neurons to compensate for perturbation-

caused changes. In homeostatic models of intrinsic excitability and regulation of synaptic

strength, both ion channel conductances and synaptic strength can be altered to recover

neuronal and circuit function properties in response to deviations from a set-point (14, 31-36).

Some homeostatic mechanisms require new protein synthesis, and therefore recovery may be

slow and require hours or even days, whereas other processes may be more rapid (33, 35, 37,

38).

Neurons constantly face diverse perturbations such as temperature changes,

neuromodulation, and stochastic changes in protein synthesis, but they rarely fail in healthy

animals under physiological conditions. This argues that most neurons are not sitting too close to

bifurcations, but neurons and circuits can tolerate some changes in their parameters without loss

of function. The effect of morphology on the stability of physiological activity has received less

attention than it probably merits. Nonetheless, there are examples of structural alterations that

can be homeostatically regulated to control activity (39).

In many neurons, including those of the STG, neuronal Na$^+$ spikes are initiated in the axon.

In STG neurons, the slow waves that underlie bursting depend on channels in the soma and

major neurites; Synaptic inputs and outputs occur on the processes, and these various regions of

the neuron are coupled appropriately (5). Biological ranges of soma-axon coupling can increase

the density and the size of manifolds that allow rebound bursting (Figs. 5, 6). Recent data argue

that the detailed morphology of STG neurons are not important for their overall function (21-23).

Thus, instead of requiring the detailed specification of the number and location of each ion

channel, this strategy requires only the specification of a generic structure, that in turn depends

on an appropriate range of values of the electrical coupling between the soma and the axon. The

control value of 100 Ωcm for the Ra in this model falls in the middle range of experimentally

reported values (40) and was originally chosen to make the voltage waveforms of the LP neuron

model roughly match experimental data (5). While individual LP and other STG neurons show

substantial animal-to-animal variability in channel conductances and channel mRNAs (1, 41-46),

it is important to remember that the motor patterns produced by these neurons in the pyloric

circuit are tightly maintained, and the LP neuron's rebound burst is critical for its proper function

(24).

The implications in this work amplify those of the sloppy morphology observed in crab

stomatogastric ganglion neurons (21-23). Although the morphological structures of these neurons

exhibit significant cell-to-cell variations, they are electrotonically compact with gradually tapering

neurites from the soma to the distal tips. This elegant, but not too rigid, solution makes graded

transmission resilient to electrotonic decrement across sparsely distributed, synchronous

presynaptic sites. The separation of synaptic integration and slow waves from axonal spike

initiation zones may be a morphological strategy established to avoid shunting of synaptic

currents (21-23, 47). Here, we use models to demonstrate that this division of labor can make

neurons more robust to perturbations.

The findings in this study may be relevant to other neuronal systems. Spike bursts are widely

observed in the cerebellum (48, 49), the neocortex (50), and the hippocampus (51). For

example, the inferior olive neurons in the cerebellum show significant similarities to the LP

neuron. Inferior olive neurons show subthreshold oscillations in the soma and then trigger axonal

spike bursts to guide cerebellum-related movement and learning (48). Additionally, soma-axon

coupling is also critical for the initiation of axonal spike bursts (complex spikes) in cerebellar

Purkinje cells (15, 52). Because of the functional importance of these signals in cerebellar coding

and learning, the reliability of triggering axonal bursting determines the robustness of

corresponding behavioral functions. Moreover, neurons may utilize their entire structure to

maintain a robust functional output in other spiking patterns (53, 54).

Materials and Methods
244

We used the LP neuron multicompartment models constructed previously (5) and implemented

them in NEURON 7.6 (55). These models were constructed to replicate the rebound bursting

properties observed in the LP neuron of the crab stomatogastric ganglion. We adopted 1292 out

of the original 1304 cell models. The excluded cell models either need a long time to generate

rebound bursting or didn't generate this pattern after reaching a steady state, possibly because of

NEURON version changes.

The current types and compartmental distributions are shown in Fig. 1. The detailed

mathematical description of these currents can be found in the original work (5). All current

models were models as ohmic currents, except I_{Ca}. For the ohmic current, $I = \bar{g} * w * (V - E_{rev})$,

$\bar{g}$ is the maximal conductance, w is the product of activation and inactivation gate state variables,

V is the transmembrane potential and E_{rev} is the reversal potential. The calcium current was

modeled by Goldman-Hodgkin-Katz equations, $I = \overline{P_{Ca}} * d_{Ca}$, where $d_{Ca} = m_{Ca}^3 * h_{Ca} * N_A * q_e *$

$z_{Ca} * ([Ca^{2+}]_i * -\xi/(\exp(-\xi) - 1) - [Ca^{2+}]_o * \xi/(\exp(\xi) - 1)), \xi = z_{Ca} * q_e/(k * T) * V$, $\overline{P_{Ca}}$ is the

maximal permeability of the calcium ions, m_{Ca} is the calcium channel activation gate, h_{Ca} is the

calcium channel inactivation gate, N_A is the Avogadro's constant, q_e is the elementary charge, z_{Ca}

is the valence of a calcium ion, $[Ca^{2+}]_i$ and $[Ca^{2+}]_o$ are intracellular and extracellular calcium

concentrations, k is the Boltzmann's constant and T is the Kelvin temperature. The three types of

inhibitory synaptic inputs were approximated by fixed rhythmic patterns of synaptic conductances.

For each type of synapse, the synaptic conductance was approximated by $g_{syn} = \overline{g_{syn}} * s, \tau_{syn} *$

$\dot{s} = s_\infty * ((V_{pre} - V_{thresh})/v_{scale})$, where $\overline{g\_syn}$ is the maximal synaptic conductance, s is the

fraction of synaptic activation, τ_{syn} is the time constant, V_{pre} is the presynaptic voltage (smoothed

experimental recordings), V_{thresh} is the synaptic threshold, and v_{scale} is the voltage sensitivity of

the synaptic activation. The AB neuron and PY neuron synapses have instantaneous dynamics

and a reversal potential of -70 mV. The PD neuron synapse has slower dynamics (with a fixed

time constant of 50 ms) and a reversal potential of -80 mV, in keeping with the known properties

of these synapses (56).

For conductance perturbation simulations within a variation range, all 14 channel

conductances were simultaneously and independently varied by values randomly sampled in the

range of [-1, +1]*variation_range*($g_{x,max}$ - $g_{x,min}$). *x* represents a type of ionic current; $g_{x,max}$ and

$g_{x,min}$ represent the maximal and the minimal conductances of current *x* in the parameter space,

respectively. A conductance was set to zero if the value became negative after a random

perturbation. A "child" model was classified as a burster if it fires periodically and there were at

least 2 spikes per burst. For the elite "parent" models in Fig. 3, the first criterion is all their "child"

models need to be bursters when the variation range is ±10%. The second criterion is the number

of spikes per burst in the "parent" models and their corresponding "child" models need to be in the

range of 3 – 11 (5). For the results in Figs. 5 and 6, the axial resistivity Ra between the soma and

the axon was varied to simulate different degrees of soma-axon coupling. We constrained the Ra

within the range of 2 Ωcm – 500 Ωcm, to maintain a sufficient number of neurons in Fig. 5C and

Fig. 6B to compare the changes in individual neurons.

Data Availability Model code will be available in ModelDB.

Acknowledgments
286
This work was supported by NIH Grants R35NS097343 and R01MH046742. Y.Z. would like to

thank Adam Taylor for sharing the model code and Sonal Kedia for sharing experimental data.

References

1.   M. S. Goldman, J. Golowasch, E. Marder, L. F. Abbott, Global structure, robustness, and
modulation of neuronal models. *J. Neurosci.* **21**, 5229-5238 (2001).
2.   J. Golowasch, M. S. Goldman, L. F. Abbott, E. Marder, Failure of averaging in the
construction of a conductance-based neuron model. *J. Neurophysiol.* **87**, 1129-1131
(2002).
3.   A. A. Prinz, C. P. Billimoria, E. Marder, Alternative to hand-tuning conductance-based
models: construction and analysis of databases of model neurons. *J. Neurophysiol.* **90**,
3998-4015 (2003).
4.   L. M. Alonso, E. Marder, Visualization of currents in neural models with similar behavior
and different conductance densities. *Elife* **8**, e42722 (2019).
5.   A. L. Taylor, J. M. Goaillard, E. Marder, How multiple conductances determine
electrophysiological properties in a multicompartment model. *J. Neurosci.* **29**, 5573-
5586 (2009).
6.   J. S. Caplan, A. H. Williams, E. Marder, Many parameter sets in a multicompartment
model oscillator are robust to temperature perturbations. *J. Neurosci.* **34**, 4963-4975
(2014).
7.   A. L. Taylor, T. J. Hickey, A. A. Prinz, E. Marder, Structure and visualization of high-
dimensional conductance spaces. *J. Neurophysiol.* **96**, 891-905 (2006).
8.   A. A. Prinz, D. Bucher, E. Marder, Similar network activity from disparate circuit
parameters. *Nat. Neurosci.* **7**, 1345-1352 (2004).
9.   A. Doloc-Mihu, R. L. Calabrese, Identifying crucial parameter correlations maintaining
bursting activity. *PLoS. Comput. Biol.* **10**, e1003678 (2014).
10.   S. Gorur-Shandilya, E. Marder, T. O'Leary, Activity-dependent compensation of cell size
is vulnerable to targeted deletion of ion channels. *Sci. Rep.* **10**, 15989 (2020).
11.   G. LeMasson, E. Marder, L. F. Abbott, Activity-dependent regulation of conductances in
model neurons. *Science* **259**, 1915-1917 (1993).
12.   Z. Liu, J. Golowasch, E. Marder, L. F. Abbott, A model neuron with activity-dependent
conductances regulated by multiple calcium sensors. *J. Neurosci.* **18**, 2309-2320 (1998).

13.   T. O'Leary, A. H. Williams, J. S. Caplan, E. Marder, Correlations in ion channel expression
319   emerge from homeostatic tuning rules. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E2645-2654
320   (2013).

321   14.   T. O'Leary, A. H. Williams, A. Franci, E. Marder, Cell types, network homeostasis, and
322   pathological compensation from a biologically plausible ion channel expression model.
323   *Neuron* **82**, 809-821 (2014).

324   15.   Y. Zang, S. Dieudonne, E. De Schutter, Voltage- and Branch-Specific Climbing Fiber
325   Responses in Purkinje Cells. *Cell. Rep.* **24**, 1536-1549 (2018).

326   16.   P. Achard, E. De Schutter, Complex parameter landscape for a complex neuron model.
327   *PLoS. Comput. Biol.* **2**, e94 (2006).

328   17.   Y. Zang, S. Hong, E. De Schutter, Firing rate-dependent phase responses of Purkinje cells
329   support transient oscillations. *Elife* **9**, e60692 (2020).

330   18.   E. Hay, S. Hill, F. Schurmann, H. Markram, I. Segev, Models of neocortical layer 5b
331   pyramidal cells capturing a wide range of dendritic and perisomatic active properties.
332   *PLoS. Comput. Biol.* **7**, e1002107 (2011).

333   19.   N. Keren, N. Peled, A. Korngreen, Constraining compartmental models using multiple
334   voltage recordings and genetic algorithms. *J. Neurophysiol.* **94**, 3730-3742 (2005).

335   20.   Y. Zang, E. De Schutter, The Cellular Electrophysiological Properties Underlying
336   Multiplexed Coding in Purkinje Cells. *J. Neurosci.* **41**, 1850-1863 (2021).

337   21.   A. G. Otopalik *et al.*, Sloppy morphological tuning in identified neurons of the crustacean
338   stomatogastric ganglion. *Elife* **6**, e22352 (2017).

339   22.   A. G. Otopalik, A. C. Sutton, M. Banghart, E. Marder, When complex neuronal structures
340   may not matter. *Elife* **6**, e23508 (2017).

341   23.   A. G. Otopalik, J. Pipkin, E. Marder, Neuronal morphologies built for reliable physiology
342   in a rhythmic motor circuit. *Elife* **8**, e41728 (2019).

343   24.   D. K. Hartline, D. V. Gassie, Jr., Pattern generation in the lobster (Panulirus)
344   stomatogastric ganglion. I. Pyloric neuron kinetics and synaptic interactions. *Biol.*
345   *Cybern.* **33**, 209-222 (1979).

346   25.   J. Magee, D. Hoffman, C. Colbert, D. Johnston, Electrical and calcium signaling in
347   dendrites of hippocampal pyramidal neurons. *Annu. Rev. Physiol.* **60**, 327-346 (1998).

348   26.   J. C. Magee, Dendritic hyperpolarization-activated currents modify the integrative
349   properties of hippocampal CA1 pyramidal neurons. *J. Neurosci.* **18**, 7613-7624 (1998).

350   27.   J. C. Magee, M. Carruth, Dendritic voltage-gated ion channels regulate the action
351   potential firing mode of hippocampal CA1 pyramidal neurons. *J. Neurophysiol.* **82**, 1895-
352   1901 (1999).

353   28.   F. Buchholtz, J. Golowasch, I. R. Epstein, E. Marder, Mathematical model of an identified
354   stomatogastric ganglion neuron. *J. Neurophysiol.* **67**, 332-340 (1992).

355   29.   G. Drion, T. O'Leary, E. Marder, Ion channel degeneracy enables robust and tunable
356   neuronal firing rates. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E5361-5370 (2015).

357   30.   H. Ori, E. Marder, S. Marom, Cellular function given parametric variation in the Hodgkin
358   and Huxley model of excitability. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E8211-E8218 (2018).

359   31.   J. Cannon, P. Miller, Synaptic and intrinsic homeostasis cooperate to optimize single
360   neuron response properties and tune integrator circuits. *J. Neurophysiol.* **116**, 2004-
361   2022 (2016).

362   32.   J. Cannon, P. Miller, Stable Control of Firing Rate Mean and Variance by Dual
363   Homeostatic Mechanisms. *J. Math. Neurosci.* **7**, 1 (2017).

364   33.   G. W. Davis, Homeostatic control of neural activity: from phenomenology to molecular
365   design. *Annu. Rev. Neurosci.* **29**, 307-323 (2006).
366   34.   G. Turrigiano, Too many cooks? Intrinsic and synaptic homeostatic mechanisms in
367   cortical circuit refinement. *Annu. Rev. Neurosci.* **34**, 89-103 (2011).
368   35.   G. G. Turrigiano, The dialectic of Hebb and homeostasis. *Philos. Trans. R. Soc. Lond. B.*
369   *Biol. Sci.* **372**, 20160258 (2017).
370   36.   D. Kueh, W. H. Barnett, G. S. Cymbalyuk, R. L. Calabrese, Na$^+$/K$^+$ pump interacts with the
371   h-current to control bursting activity in central pattern generator neurons of leeches.
372   *Elife* **5**, e19322 (2016).
373   37.   R. Gowrishankar *et al.*, Region-Specific Regulation of Presynaptic Dopamine
374   Homeostasis by D2 Autoreceptors Shapes the In Vivo Impact of the Neuropsychiatric
375   Disease-Associated DAT Variant Val559. *J. Neurosci.* **38**, 5302-5312 (2018).
376   38.   Y. Kulik, R. Jones, A. J. Moughamian, J. Whippen, G. W. Davis, Dual separable feedback
377   systems govern firing rate homeostasis. *Elife* **8**, e45717 (2019).
378   39.   M. S. Grubb, J. Burrone, Activity-dependent relocation of the axon initial segment fine-
379   tunes neuronal excitability. *Nature* **465**, 1070-1074 (2010).
380   40.   A. Roth, M. Hausser, Compartmental models of rat cerebellar Purkinje cells based on
381   simultaneous somatic and dendritic patch-clamp recordings. *J. Physiol.* **535**, 445-472
382   (2001).
383   41.   J. M. Goaillard, A. L. Taylor, D. J. Schulz, E. Marder, Functional consequences of animal-
384   to-animal variation in circuit parameters. *Nat. Neurosci.* **12**, 1424-1430 (2009).
385   42.   J. Golowasch, L. F. Abbott, E. Marder, Activity-dependent regulation of potassium
386   currents in an identified neuron of the stomatogastric ganglion of the crab Cancer
387   borealis. *J. Neurosci.* **19**, RC33 (1999).
388   43.   A. J. Northcutt *et al.*, Molecular profiling of single neurons of known identity in two
389   ganglia from the crab Cancer borealis. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 26980-26990
390   (2019).
391   44.   A. J. Northcutt *et al.*, Deep sequencing of transcriptomes from the nervous systems of
392   two decapod crustaceans to characterize genes important for neural circuit function and
393   modulation. *BMC. Genomics.* **17**, 868 (2016).
394   45.   D. J. Schulz, J. M. Goaillard, E. Marder, Variable channel expression in identified single
395   and electrically coupled neurons in different animals. *Nat. Neurosci.* **9**, 356-362 (2006).
396   46.   D. J. Schulz, J. M. Goaillard, E. E. Marder, Quantitative expression profiling of identified
397   neurons reveals cell-specific constraints on highly variable levels of gene expression.
398   *Proc. Natl. Acad. Sci. U. S. A.* **104**, 13187-13191 (2007).
399   47.   A. E. Tobin, S. D. Van Hooser, R. L. Calabrese, Creation and reduction of a
400   morphologically detailed model of a leech heart interneuron. *J. Neurophysiol.* **96**, 2107-
401   2120 (2006).
402   48.   A. Mathy *et al.*, Encoding of oscillations by axonal bursts in inferior olive neurons.
403   *Neuron* **62**, 388-399 (2009).
404   49.   Y. Zang, E. De Schutter, Climbing Fibers Provide Graded Error Signals in Cerebellar
405   Learning. *Front. Syst. Neurosci.* **13**, 46 (2019).
406   50.   M. E. Larkum, J. J. Zhu, B. Sakmann, A new cellular mechanism for coupling inputs
407   arriving at different cortical layers. *Nature* **398**, 338-341 (1999).

51. H. Su, G. Alroy, E. D. Kirson, Y. Yaari, Extracellular calcium modulates persistent sodium
current-dependent burst-firing in hippocampal pyramidal neurons. *J. Neurosci.* **21**, 4173-
4182 (2001).
52. J. T. Davie, B. A. Clark, M. Hausser, The origin of the complex spike in cerebellar Purkinje
cells. *J. Neurosci.* **28**, 7599-7609 (2008).
53. E. Hay, F. Schurmann, H. Markram, I. Segev, Preserving axosomatic spiking features
despite diverse dendritic morphology. *J. Neurophysiol.* **109**, 2972-2981 (2013).
54. M. S. Hamada, S. Goethals, S. I. de Vries, R. Brette, M. H. Kole, Covariation of axon initial
segment location and dendritic tree normalizes the somatic action potential. *Proc. Natl.*
Acad. Sci. U. S. A. **113**, 14841-14846 (2016).
55. M. L. Hines, N. T. Carnevale, The NEURON simulation environment. *Neural. Comput.* **9**,
1179-1209 (1997).
56. E. Marder, J. S. Eisen, Transmitter identification of pyloric neurons: electrically coupled
neurons use different transmitters. *J. Neurophysiol.* **51**, 1345-1361 (1984).

Figures and Tables

426
Fig. 1. LP neuron multi-compartment model structure and spike properties. (*A*) Schematic of the
chemical synaptic inhibition that a LP neuron receives. (*B*) Intracellular recordings of a LP neuron
(top) and an extracellular recording at the lvn (bottom, spikes of LP, PY and PD neurons are color
coded, unpublished data from Sonal Kedia). (*C*) Schematic of the LP neuron models, consisting
of an axon, a soma, proximal neurites, and distal neurites with corresponding ion channels and
synapses distributed in each compartment, modified from (5). (*D*) Example somatic (top) and
axonal (bottom) spike waveforms in a model. The green and blue shadings illustrate the timing of
the inhibition from PY neurons and PD neurons (also AB neuron).

Fig. 2. "Noise" perturbation protocol and neuronal resilience to conductance variations. (*A*) 2-D schematic of the channel conductance perturbation protocol (bottom middle, shaded in yellow). *x* and *y* represent different types of ion channels. In each trial, all channel conductances of a "parent" model were simultaneously varied by random values (details in Methods). Each "parent" model (n = 1292) generated new m "child" models after m-trial perturbations. Example firing patterns in "child" models include tonic spiking (child 1), bursting (child 2), aperiodic spiking (child 3), depolarization block (db, child 4), and non-spiking (child 100). (*B*) Histogram of the ratio of "child" models that are bursters for each "parent" model, when models undergo different ranges of random channel conductance variations, ±10%, ±50%, and ±100% from left to right.

445
446
447 Fig. 3. Robust models are widely distributed in parameter space. (*A*) The spike waveforms in the
448 soma (top) and axon (bottom) of 8 example "parent" models (models a – h), whose "child" models
449 reliably produce rebound bursting after 10,000-trial perturbations when the variation range is
450 ±10%. (*B*) The parameter distribution of these 8 example models. Colors code different models
451 (models a – h from panel A).

452
Fig. 4. Factors regulating neuronal robustness. (*A*) The effect of variation range on neuronal

resilience between two groups of neurons. They are grouped by the ratio of "child" models that

are bursters when the variation range is ±10% (orange group, 100%; blue group, <70%). (*B*) The

sensitivity of neuronal resilience to individual channel conductance variations (coded by colors).

Left and right plots show the effect of varying one of the channel conductances in the soma-

neurite and the axon, respectively. The x-axis represents conductance variation range, and the y-

axis represents the average ratio of "child" models that are bursters.

Fig. 5. The effect of soma-axon coupling. (*A*) Example somatic (green) and axonal (black) spike waveforms when Ra was changed from 100 Ωcm (middle) to 2 Ωcm (left) and 500 Ωcm (right). (*B*) The relationship between the soma to axon Ra and the bursting neuron population size (n_burster). n = 1292 when Ra = 100 Ωcm under control condition. The x-axis is a logarithmic scale. (*C*) The relationship between the soma to axon Ra and the number of spikes per burst (spike_burst) in neurons that generate rebound bursting regardless of Ra changes (n = 130). Purple, population average; red and cyan, two example neurons; gray, other neurons. The x-axis is a logarithmic scale. (*D*) The effect of soma to axon Ra on somatic (green) and axonal (black) bursting changes in the example cells (cells a and b) highlighted in C (Ra, 2 Ωcm → 500 Ωcm from left to right). In all simulations, there are no channel conductance variations.

Fig. 6. The effect of soma-axon coupling on neuronal resilience to conductance variations. (*A*) The effect of soma to axon Ra on neuronal resilience to conductance variations for all the "parent" models. A1, The average ratio of "child" models that are bursters with different Ra values (coded by colors). A2 and A3 show individual neuronal ratios of "child" models that are bursters for Ra = 2 Ωcm and 500 Ωcm, respectively. Green traces show typical example plots. Blue and magenta traces show population average ratios, as in A1. Gray traces show other individual neurons. (*B*) The comparison of each neuronal ratio of "child" models that are bursters when Ra is changed from the control value of 100 Ωcm. B1, Ra is changed from 100 Ωcm (black dots) to 2 Ωcm (blue dots); B2, Ra is changed from 100 Ωcm (black dots) to 500 Ωcm (magenta dots). From left to right, conductance variation range increases from ±10% to ±100%. Eighty of the total 131 elite "parent" models are shown.