Exploration of the sensitivity to macrocyclic lactones in the canine heartworm

(*Dirofilaria immitis*) in Australia using phenotypic and genotypic approaches

Rosemonde Isabella Power[a] and Jan Šlapeta[a,b,*]

[a]Sydney School of Veterinary Science, Faculty of Science, University of Sydney, New South

Wales 2006, Australia

[b] The University of Sydney Institute for Infectious Diseases, New South Wales 2006,

Australia

* Corresponding author.

E-mail address: jan.slapeta@sydney.edu.au (J. Šlapeta)

Keywords: disease, circulating microfilariae, moxidectin, Illumina metabarcoding, PCR,

antigen, Knott's test, *Wolbachia*

Abstract

Canine heartworm disease is a potentially deadly cardiopulmonary disease caused by the mosquito-borne filarial nematode *Dirofilaria immitis*. In Australia, the administration of macrocyclic lactone (ML) drugs has successfully reduced the prevalence of *D. immitis* infection. However, the recent re-emergence of *D. immitis* in dogs in Queensland, Australia and the identification of ML-resistant isolates in the USA poses an important question of whether ML-resistance has emerged in this parasite in Australia. The aim of this study was to utilise phenotypic and genotypic approaches to examine the sensitivity to ML drugs in *D. immitis* in Australia. To do this, we surveyed 45 dogs from Queensland and New South Wales across 3 years (2019-2022) for the presence of *D. immitis* infection using an antigen test, quantitative Modified Knott's test, and qPCR targeting both *D. immitis* and the *D. immitis* symbiont *Wolbachia*. A phenotype observed by utilising sequential quantification of microfilariae for 23/45 dogs was coupled with genetic testing of filtered microfilariae for SNPs previously associated with ML-resistance in isolates from the USA. Sixteen (16/45) dogs tested positive for *D. immitis* infection despite reportedly receiving 'rigorous' heartworm prevention for 12 months prior to the study, according to the owners' assessment. The phenotype and genotypic assays in this study did not unequivocally demonstrate the presence of ML-resistant *D. immitis* in Australia. Although the failure of 16 dogs to reduce microfilaremia by >90% after ML treatment was considered a suspect phenotype of ML-resistance, no genotypic evidence was discovered using the genetic SNP analysis. The traditional quantitative Modified Knott's test can be substituted by qPCR targeting *D. immitis* or associated *Wolbachia* endosymbiont DNA for a more rapid measurement of microfilariae levels. More definitive phenotypic evidence of resistance is critically needed before the usefulness of SNPs for the detection of ML-resistance in Australia can be properly assessed.

1. Introduction

Canine heartworm disease is a potentially deadly cardiopulmonary disease caused by the mosquito-borne filarial nematode *Dirofilaria immitis*, which is distributed across many parts of the world (Simón et al., 2012; Genchi and Kramer, 2020; Lau et al., 2021; Noack et al., 2021). Although the prevalence of *D. immitis* in Australia is low, the parasite is endemic to the country and has recently re-emerged in dogs in Queensland (Nguyen et al., 2016; Orr et al., 2020; Panetta et al., 2021). For the past 30 years in Australia and the USA, the main strategy for canine heartworm prevention has been the administration of macrocyclic lactone (ML) anthelmintics (Orr et al., 2020; Diakou and Prichard, 2021; Lau et al., 2021; Noack et al., 2021; Prichard, 2021). These ML preventatives include monthly oral or topical formulations of ivermectin, milbemycin oxime, moxidectin and selamectin, or extended-release injections of moxidectin microspheres (Nguyen et al., 2016; McTier et al., 2019a). When used according to product instructions, these preventatives are approved to be 100% effective against less than 30-day old *D. immitis* (Hampshire, 2005). Over the past 20 years, lack of efficacy (LOE) reports for all heartworm product categories have occurred in the Lower Mississippi River Valley region of the USA (Hampshire, 2005). Most of these LOE reports were associated with compliance issues and extenuating circumstances, which prevented dogs from receiving sufficient product every 30 days (Atkins et al., 2014). A small number of LOE cases were not linked to noncompliance, indicating the potential development of ML-resistance in canine heartworms (Atkins et al., 2014). Laboratory studies have now identified several ML-resistant *D. immitis* isolates circulating in the Lower Mississippi Valley region of the USA (Blagburn et al., 2011; Pulaski et al., 2014; Bourguinat et al., 2015; Blagburn et al., 2016; Maclean et al., 2017; McTier et al., 2017b; Diakou and Prichard, 2021). These studies involved experimentally infecting ML-naïve dogs with infective L3 obtained from field LOE cases (Pulaski et al., 2014). In addition, <100%

efficacy has been experimentally demonstrated for all currently marketed MLs (Bourguinat et

al., 2015; Blagburn et al., 2016; McTier et al., 2019b).

MLs are essentially the only class of drugs currently available for canine heartworm

prevention in Australia, in addition to the daily diethylcarbamazine citrate (Noack et al.,

2021). Hence, the emergence of ML-resistance in *D. immitis* in the USA presents a

significant concern for the veterinary industry by limiting the usefulness of ML products and

jeopardising canine health. Prolonging the efficacy of MLs for canine heartworm prevention

is warranted. With the absence of rapid biological tests that would demonstrate LOE of the

target population (<30-day old *D. immitis*), discovery of heritable DNA molecular markers is

needed for monitoring ML-resistance in adult *D. immitis* populations or their microfilariae

within the dog bloodstream. Previous studies from the USA have compared the genotypes of

different groups of *D. immitis* and identified single nucleotide polymorphisms (SNPs)

potentially linked to ML-resistance (Bourguinat et al., 2011; Bourguinat et al., 2015). A

whole genome approach was used in Bourguinat et al. (2015) to identify 186 SNP loci

associated with the inability of dogs to remove circulating microfilariae after ML

administration. From these 186 SNPs, a subset of 42 SNPs was initially validated as being

highly associated with the ML-resistance phenotype in *D. immitis* (Bourguinat et al., 2015).

The 42 SNPs were further evaluated in 10 additional ML-susceptible and ML-resistant

isolates to generate 2, 3, 5 and 10-SNP predictive models for ML-resistance (Bourguinat et

al., 2017). The 10 SNP approach that most effectively separated ML-susceptible and ML-

resistant isolates was validated in an extensive study from south-east USA with field-

collected *D. immitis* microfilariae, demonstrating high levels of sensitivity and specificity

(Ballesteros et al., 2018). The SNP-based genotype predicted the phenotype which was

defined by the inability of MLs to clear microfilariae using a suppression test and thus

implied the presence of ML-resistant *D. immitis* infection (Ballesteros et al., 2018).

In contrast to the USA, it is currently unknown whether ML-resistance has emerged in *D.*

immitis in Australia. A recent study revealed a large proportion of Australian dogs that were

microfilaria-positive despite 'rigorous' administration of MLs in far-north Queensland

(Nguyen et al., 2016). However, evidence of ML purchase history or ML administration was

not provided by clients (Nguyen et al., 2016). A study that adopted the predictive SNP

models developed in the USA to determine ML-resistance status did not find evidence of

SNPs associated with ML-resistance in adult *D. immitis* from Sydney, Australia (Lau et al.,

2021). As *D. immitis* prevention remains critical and prevalent across tropical Australia,

further inquiry into ML-resistance in Australian *D. immitis* populations is necessary (Orr et

al., 2020; Panetta et al., 2021). New screening tools and surveillance are required that apply

or adapt the microfilarial suppression test originally proposed by Geary et al. (2011) and

subsequently utilised by Ballesteros et al. (2018).

The aim of this study was to utilise phenotypic and genotypic approaches to determine the

presence of ML-resistant *D. immitis* in Australia. To do so, we surveyed dogs across 3 years

(2019-2022) for the presence of *D. immitis* using an antigen test, quantitative Modified

Knott's test, and qPCR targeting both *D. immitis* as well as the *D. immitis* symbiont

Wolbachia. Material was collected prior to ML application and post-ML application at

varying intervals. A Modified Knott's test was coupled with filtration of microfilariae for

genetic testing of SNPs previously associated with ML-resistance in isolates from the USA.

2. Material & Methods

2.1. Blood collection from dogs in Queensland and New South Wales

Blood samples were collected opportunistically from dogs in Queensland and New South

Wales, Australia between September 2019 and January 2022. The dogs selected for this study

were privately-owned animals from participating veterinary practices where the practitioner

had either confirmed or suspected the presence of *D. immitis* infections. Blood samples (≤4

mL) were collected by registered veterinarians and shipped in EDTA tubes to the Sydney

School of Veterinary Science, University of Sydney, Australia for diagnostic *D. immitis*

testing. The following details were obtained for each dog: sex, age, breed, desexing status,

locality, heartworm prevention history for the previous 12 months based on the owner's

assessment, and whether the animal had lived or visited other states or territories. Dog owners

were required to complete a consent form prior to blood collection.

2.2. Sequential Modified Knott's test

To determine the effect of ML administration on *D. immitis* microfilariae counts from

infected dogs, we utilised a Modified Knott's test. This was used to enumerate the number of

D. immitis microfilariae in dog blood before ML treatment and then at a second point

following the ML treatment. The time interval between drug treatment and the follow-up

count varied across the study as necessitated by the need to follow timelines within the

veterinary practices that were providing blood samples for the study, but was generally in the

period 30-40 days after treatment. In some cases, an additional post-treatment microfilariae

count was made at a later timepoint. The type of ML product administered was at the

veterinarian's discretion, according to their best practice in heartworm disease management.

We calculated the percentage (%) reduction of microfilariae between the pre-treatment and

post-treatment samples. In cases in which dogs showed <90% reductions in microfilariae,

additional blood samples were collected where possible to observe whether microfilariae

levels eventually declined.

To perform the Modified Knott's test, 1 mL of whole blood was mixed with 10 mL of 2%

formalin in a 15-mL tube (Knott, 1939; Mylonakis et al., 2004). The mixture was centrifuged

at 2,300 relative centrifugal force (RCF) for 5 min using a benchtop centrifuge (POCD

Scientific, Australia). The pellet was mixed with 1% methylene blue stain and enumerated

microscopically. The slide was examined under a microscope (BX40, Olympus, Australia)

using 10× objective and the number of microfilariae observed in the entire slide was counted.

2.3. Detection of *D. immitis* antigen using DiroCHEK® ELISA

Prior to performing DiroCHEK® ELISA (Zoetis, Australia) testing, the stored plasma

samples were thawed at room temperature and vortexed for 3 sec. Both heated and unheated

(neat) aliquots of each plasma sample were tested. To heat the plasma samples, a 100 μL

aliquot of plasma was diluted with 100 μL of phosphate buffered saline (PBS; pH = 7.4) in a

plain 1.5-mL tube. For samples with insufficient plasma volume, only 50 μL of plasma was

diluted with 50 μL of PBS. The diluted plasma samples were heated at 103°C for 10 min

using a dry heat block (Major Science). Heated samples were immediately centrifuged at

16,100 RCF for 5 min using a benchtop centrifuge (Eppendorf, Australia) set at 4°C. The

supernatant was transferred into a separate 1.5-mL tube for use in DiroCHEK® ELISA

testing. To load the DiroCHEK® ELISA plate, 50 μL of heated and unheated plasma for each

sample were transferred into separate wells. Positive and negative controls provided by the

manufacturer were included, and samples with leftover heated plasma were included as

repeats. The DiroCHEK® ELISA test was then performed according to manufacturer's

instructions. The optical density (OD) of each sample in the DiroCHEK® ELISA plate was

measured at 620 nm in a Halo LED 96 microplate reader (Dynamica). To distinguish between

negative and positive samples, the following equation was applied as a threshold:

$$Threshold = 3 \ x \ SD \ (Mean[OD \ of \ negative \ controls])$$

2.4. Microfilariae extraction

To isolate microfilariae from dog blood, a filtration procedure was adopted and modified

(Bourguinat et al., 2015). Prior to filtration, 1 mL of whole blood was diluted with 35 mL of

Red Blood Cell lysis buffer (8 g/L sodium carbonate, 5 mL/L Triton X-100) in a 50-mL tube.

For one sample (HW52), only 100 µl of blood was diluted. The diluted blood was then

pushed through a syringe containing a polycarbonate membrane filter (pore size 20 µm, filter

diameter 25 mm; Sterlitech[®] Corporation, WA). This allowed the microfilariae to be

collected on the filter membrane whilst other blood components passed through the syringe.

The syringe was then flushed with 50 mL of phosphate buffered saline (PBS). The filter

membrane was removed from the syringe, placed into a 1.5-mL tube, and stored at -20°C

until further use.

2.5. DNA isolation

DNA was isolated from 100 µL of whole blood using the blood protocol from the Monarch[®]

Genomic DNA Purification Kit (New England Biolabs, Australia). In addition, DNA was

isolated from microfilariae on filter membranes using the same kit. DNA isolation from filter

membranes followed the animal tissue sample protocol and used Proteinase K. Each DNA

isolation batch contained an extraction blank. The eluted DNA was stored at -20°C until

further use in the qPCR and next generation sequencing (NGS) assays.

2.6. Quantitative PCR targeting dog, *D. immitis* and *Wolbachia* sp. of *D. immitis*

A qPCR targeting the partial canine glyceraldehyde3-phosphate dehydrogenase (*GAPDH*)

region was performed to verify the presence of dog DNA using primers with a probe, as

previously described (Orr et al., 2020; Panetta et al., 2021) (Supplementary Table S1). A

diagnostic TaqMan probe qPCR targeting the *cox1* gene was performed to detect *D. immitis*

DNA using primers with a probe, as previously described (Panetta et al., 2021)

(Supplementary Table S1). A qPCR targeting the *ftsZ* gene of *D. immitis*-associated

Wolbachia was performed using primers and probe described in Laidoudi et al. (2020)

(Supplementary Table S1).

The dog, *D. immitis* and *Wolbachia* qPCR assays were carried out in 10 μL final volumes of

Luna® Universal Probe qPCR Master Mix (New England Biolabs, Australia) containing 1 μL

of template DNA isolated from whole dog blood and primers and probe at a final

concentration of 400 nM and 100 nM, respectively. To confirm the presence of *D. immitis*,

qPCRs targeting dog and *D. immitis* DNA were repeated using DNA from extracted

microfilariae. For these reactions, the volumes of primer, probe, PCR-grade water, and

template DNA were doubled to reach final volumes of 20 μL. All qPCRs were carried out on

a CFX96 Real-Time PCR system with the corresponding Bio-Rad CFX Maestro Software

(BioRad, Australia). Cycling conditions for all three assays consisted of an initial

denaturation at 95°C for 1 min followed by 40 two-step cycles at 95°C for 15 sec and 60°C

for 30 sec. Each qPCR run included DNA extraction blanks and a no template control (NTC)

of PCR-grade water. The arbitrary qPCR baseline threshold was set to 100 RFU and C_t-

values were reported for all samples. A qPCR targeting the *GAPDH* region of the dog

genome confirmed the presence of dog DNA in all samples. All No Template Control (NTC)

reactions remained negative.

2.7. Statistical analysis of phenotypic data

A paired t-test was performed to compare the OD between neat and heated plasma samples in

the DiroCHEK® ELISA. Additional paired t-tests were performed to observe differences

between pre- and post-ML treatment samples in the qPCRs. To evaluate linear relationships

between phenotypic datasets, Pearson's correlation coefficients were generated. Prior to

performing correlation analyses with the qPCR data, 'N/A' results were replaced with a C_t-

value of '40'. All statistical analyses were performed using GraphPad Prism (version 9.3.1).

2.8. Selection of SNP markers

Studies from the USA identified 10 SNPs that best differentiated ML-susceptible *D. immitis* from ML-resistant isolates (Bourguinat et al., 2017). Seven of these SNPs were selected for analysis in this study as they covered both 5-SNP models generated by Ballesteros et al. (2018) and Bourguinat et al. (2017). The 2 SNP markers (L15709_A, L30575) reported as giving the best prediction of ML-resistance in Ballesteros et al. (2018) were included in these 7 SNPs. Additional information about the 7 SNPs selected for this study including their position in the genome and various names used in other studies are available in Supplementary Table S2.

2.9. Illumina PCR amplicon Next Generation Sequencing targeting 7 SNPs associated with ML-resistance in USA

Illumina overhang adapters were added to all 7 primer sets which targeted regions encompassing the SNPs of interest (L42411, L21554, L45689, L9400, L20587, L15709_A, and L30575) according to Bourguinat et al. (2015). Their reference sequences were mapped to *D. immitis* genome nDi.2.2 scaffolds (Godel et al., 2012). The first stage PCRs for Illumina Next Generation Sequencing (NGS) were performed in-house according to the '16S Metagenomic Sequencing Library Preparation' instructions for the Illumina MiSeq system (Illumina, Australia). Conventional PCR assays were carried out with Q5 Hot Start High-Fidelity 2× Master Mix (New England Biolabs, Australia). For some samples, dilutions of 1:10 or 1:100 were used due to low C_t-values obtained from the *D. immitis*-specific qPCR. Conventional PCRs were performed on a Bio-Rad T100™ Thermal Cycler (Bio-Rad, Australia). The cycling conditions included an initial denaturation step at 98°C for 30 sec followed by 27 – 35 cycles of 98°C for 5 sec, 55°C for 10 sec and 72°C for 20 sec with a final extension at 72°C for 2 min. The number of cycles varied so that each sample reached

the exponential phase of amplification, as previously described (Power et al., 2021). To

increase efficiency of the first stage PCRs, conventional PCR assays were replaced with

qPCR assays with SensiFAST™ SYBR® No-ROX Kit (Bioline, Australia). The cycling

conditions included an initial denaturation at 95°C for 3 min followed by 28 - 40 cycles of

95°C for 5 sec, 55°C for 10 sec and 72°C for 20 sec with a final melt curve. The inclusion of

a melt curve allowed us to confirm the presence of the desired PCR product. DNA of an adult

D. immitis collected from a dog in Sydney (P6/20D), Australia was used as a positive control

(Lau et al., 2021). A no template control (NTC) of PCR-grade water was included for each

SNP assay. Presence of expected PCR products was confirmed by visualisation on agarose

gel for conventional PCR and qPCR products with ambiguous melt profiles. To determine

whether comparable results were obtained using the two PCR approaches, three samples

(HW19, HW20, HW32) were run as both conventional PCR and qPCR reactions for 5 loci

(L42411, 21554, 45689, 9400, 20587), along with additional samples for L42411 (HW33)

and L20587 (HW63). To investigate the impact of DNA concentration on the NGS results,

four samples in the L45689 assay (HW20, HW41, HW45, HW66) and four samples in the

L9400 assay (HW20, HW31, HW66, P6/20D) were repeated using alternative dilutions or

cycle numbers. As one sample (HW52) was analysed using filtered microfilariae from 100

µL of blood, repeats for three loci (L20587, L15709A, L30575) were included using filtered

microfilariae from 1 mL of blood. There was no difference that wound compromised SNP

calls between the approaches irrespective of the dilution, blood volume, cycle number or

PCR approach used.

After the first-stage PCR, products were submitted to the Ramaciotti Centre for Genomics,

University of New South Wales, Australia for indexing (384), library preparation and next

generation sequencing (NGS) using Illumina MiSeq v2 250PE sequencing run.

2.10. SNP analysis of NGS samples

Data as FastQ files were analysed in R 4.0.5 within RStudio 2022.07.1 Build 554 (RStudio

Team, 2022). For each SNP locus, primers were identified and removed from all sequencing

reads using Cutadapt 4.0 (Martin, 2011) followed by DADA2, an open-source R package

(https://github.com/benjjneb/dada2). DADA2 pipeline (1.18.0; Callahan et al. (2016)) was

used to infer the *D. immitis* SNP amplicon sequence variants (ASVs) present in the filtered

microfilariae DNA samples. The quality profiles of the cut sequencing reads were visualised

and inspected using the "plotQualityProfile" tool. The cut forward and reverse reads (FastQ)

were filtered, denoised and trimmed to 180 and 150 nucleotides, respectively, using the

"filterAndTrim" tool. Overall, >92%, >75%, >97%, >71%, >96%, >91%, and >40% of input

reads were retained and 0.03%, 0.01%, 0.58%, 0.01%, 0.01%, 0.00%, and 0.00% of merged

paired reads were removed as chimeras for L42411, L21554, L45689, L9400, L20587,

L15709_A, and L30575, respectively. For data cleaning purposes, the proportion (%) of each

ASV in each sample was calculated. In an effort to remove misassigned reads caused by

sample bleeding or barcode hopping, a threshold of <2.5% was applied (Mitra et al., 2015).

This involved removing the reads of ASVs whose proportions represented <2.5% in a sample

and replacing them with '0'. Alternative thresholds of <1% and <0.5% were tested but not

selected for further downstream analysis. A reference table containing frequencies of

sequence fragments of all 7 SNP loci was created, with options for each possible SNP

polymorphism. This reference table was imported into DADA2 and the "AssignSpecies" tool

was used to assign the remaining ASVs to a reference sequence if they were a perfect match.

ASVs were imported into CLC Main Workbench (version 22.0) (CLC bio, Qiagen, Australia)

and "BLAST at NCBI" was performed. Based on the BLAST results, ASVs that were not

identified as *D. immitis* were excluded and ASV proportions for each sample were re-

calculated. The remaining *D. immitis* ASVs were aligned to their respective SNP reference

sequences and visually inspected using CLC Main Workbench.

2.11. Animal Ethics

Research performed at the University of Sydney was approved by the Animal Research Ethics Committee, which follows the NSW Animal Research Act 1985 and the Australian code for the care and use of animals for scientific purposes 8th edition (2013).

2.12. Data accessibility

Raw FastQ sequences were deposited at SRA NCBI BioProject: PRJNA889241. Additional data for this study is available at LabArchives (https://dx.doi.org/10.25833/ygs4-ha97).

3. Results

3.1. Blood collection pre- and post-ML treatment

A total of 45 dogs (Dog 1 to Dog 45) and their 78 blood samples (HW01 to HW63, HW65 to HW79) were collected and followed up over 2.5 years (2019-2022) as suspect to be infected with *D. immitis* based on the veterinarians' judgement (Supplementary Table S3). At the beginning of the study, infection with *D. immitis* was confirmed in 44/45 dogs using at least one diagnostic test. Dogs testing positive for *D. immitis* were domiciled at eleven locations across eastern Australia, with thirty-eight dogs from Queensland and six dogs from New South Wales (Figure 1). All dog bloods were processed using an exhaustive diagnostic laboratory workflow (Figure 2). Twenty-three (n=23) dogs were sampled at the time of diagnosis (pre-ML treatment) and at least once post-diagnosis; 14 dogs once after treatment (post-ML), eight dogs twice after treatment, and one dog (Dog 40) three times after treatment (Figure 2, 3). In addition, twenty-two dogs were only sampled pre-ML treatment and following samples were not possible to obtain.

3.2. Effects of ML administration on *Dirofilaria immitis* microfilariae counts

A total of 37/44 (84.1%) *D. immitis*-positive dogs were microfilaraemic at the beginning of the study, with a mean initial microfilariae count of 7,689 microfilariae/mL. Sequential Modified Knott's tests were performed to demonstrate the suspect ML-resistance phenotypes of *D. immitis* (Figure 2). The percentage (%) reduction of microfilariae counts after ML administration was obtained for 22 dogs that had sequential blood samples. The mean initial microfilariae count for 21 dogs was 6,617 microfilariae/mL (min. 38 to max. 38,250 microfilariae/mL) plus one had 1 microfilaria/mL (Dog 1) (Figure 3, Table 1). In addition, one dog (Dog 19) had 0 microfilariae/mL, but tested positive on PCR and serology so a post-ML sample was also obtained and processed. Sixteen dogs (73%, 16/22) had ≤ 90% reduction in microfilariae count, including three dogs (Dog 25, Dog 2, Dog 7) which had an increase in microfilariae (Figure 3, Table 1).

Using seven dogs sampled pre-ML which were then administered moxidectin (ProHeart SR-12, s.c.) and sampled again at least twice post-ML, we show a 47% (14%-67%, Figure 3A) reduction of microfilariae by day 27-49. This was reduced to 95% (73%-100%)by 60-136 days. Four of these dogs were subjectively considered to be on 'rigorous' heartworm prevention by their owners (Figure 3A). All remaining dogs but three were administered moxidectin (ProHeart SR-12, s.c.) at the time of diagnosis (Figure 3B,C,D). Three dogs were administered ivermectin (HeartGard Plus, p.o.). Dog 21 on ivermectin (HeartGard Plus) completely reduced their microfilariae by day 28 from a relatively low initial microfilaremia of 175 microfilariae/mL (Figure 3C). The remaining two dogs (Dog 2, 13) on ivermectin (HeartGard Plus) experienced <90% reductions in microfilaremia post-ML, with Dog 2 experiencing an increase in microfilariae by 181% (Figure 3C).

3.3. Presence of heartworm antigen using DiroCHEK® ELISA

Out of all dogs with confirmed *D. immitis* infection at the start of this study, 41/44 (93.2%)

dogs initially tested positive for *D. immitis* antigen using DiroCHEK[®] ELISA with unheated

(neat) plasma samples and 43/44 (97.7%) initially tested positive using DiroCHEK[®] ELISA

with heated plasma samples. Out of all blood samples collected in this study, 88.2% (67/76)

of neat plasma samples and 98.7% (75/76) of heated plasma samples tested positive for *D.*

immitis antigen using DiroCHEK[®] ELISA, respectively (Figure 4A, Table 1). A paired t-test

was performed on the raw DiroCHEK[®] ELISA data, revealing a significant difference in

OD_{620} between neat and heated plasma samples (Figure 4B). Moderate positive correlations

were found between raw microfilariae counts and the OD_{620} of neat and heated dog plasma in

DiroCHEK[®] ELISA (Figure 5A). When performing the modified Knott's test on samples

with microfilariae counts >20,000 mf/mL, we observed the overlap and clumping of

microfilariae, which made counting these samples difficult. To determine whether similar

statistical results were still obtained for samples with microfilariae counts <20,000 mf/mL,

the correlation analyses between microfilariae counts and the DiroCHEK[®] ELISA were

repeated. By doing so, moderate positive correlations were maintained (Figure 5B).

3.4. Presence of *Dirofilaria immitis* and *Wolbachia* sp. of *D. immitis* using qPCR

DNA samples obtained from dog blood were subjected to qPCR targeting the *cox1* gene of *D.*

immitis and the *ftsZ* gene of *D. immitis*-associated *Wolbachia*. *D. immitis* and *Wolbachia*

DNA were initially detected in 39/44 (88.6%) and 34/44 (77.3%) dogs with confirmed *D.*

immitis infection, respectively. At Day 0 (pre-ML), *D. immitis* and *Wolbachia* were detected

in 100% (23/23, C_t mean: 24.63, C_t range: 19.48-37.68) and 82.61% (19/23, C_t mean: 28.21,

C_t range: 25.07-31.75) of dogs, respectively. At ~Day 30 (post-ML), *D. immitis* was detected

in 82.61% (19/23, C_t mean: 24.37, C_t range: 21.80-32.13) of dogs and *Wolbachia* was

detected in 65.22% (15/23, C_t mean: 33.10, C_t range: 28.78-35.48) of dogs. By Day 30+,

77.8% (7/9, C_t mean: 29.00, C_t range: 25.89-36.45) of dogs were positive for *D. immitis* and

33.33% (3/9, C_t mean: 30.27, C_t range: 21.01-35.78) of dogs were positive for *Wolbachia*.

The *D. immitis*-specific qPCR on extracted microfilariae DNA found one additional *D.*

immitis DNA positive dog (Dog 31: 0 microfilariae/mL) that tested negative for *D. immitis*

DNA on a sample from whole blood. In addition, two dogs (Dog 1: 1 microfilaria/mL, Dog

20: 0 microfilariae/mL) were *D. immitis* DNA positive only on whole blood but *D. immitis*

DNA-negative using extracted microfilariae (Supplementary Table S3).

A strong negative correlation was demonstrated between the *D. immitis* qPCR C_t-values of

whole blood DNA and microfilariae counts that were <20,000 mf/mL (Figure 5C). Samples

with microfilariae counts >20,000 mf/mL were excluded from this analysis due to difficulties

with counting clumped and overlapping microfilariae. Similarly, a strong negative correlation

was observed between the *Wolbachia* qPCR C_t-values of whole blood DNA and microfilariae

counts <20,000 mf/mL (Figure 5D). These statistical analyses were repeated to determine

whether similar correlations were still obtained when including samples with microfilariae

counts > 20,000 mf/mL. By doing so, a moderate negative correlation was shown between

the *D. immitis* qPCR C_t-values of whole blood DNA and microfilariae counts (Figure 5E).

For all *Wolbachia* qPCR of whole blood DNA, a moderate negative correlation was observed

with microfilariae counts (Figure 5F). A strong positive correlation was found between the *D.*

immitis and *Wolbachia* qPCR C_t-values of whole blood DNA samples (Figure 5G). Samples

with C_t-values of 40 were considered negative and excluded from the analysis, resulting in a

moderate positive correlation (Figure 5H).

Paired t-tests of the raw *D. immitis* and *Wolbachia* qPCR data revealed statistically

significant increases in C_t-value from pre-ML to first post-ML sampling (Figure 6A,B). To

determine whether similar results were obtained for samples with pre-ML C_t-values <30 in

the *D. immitis* qPCR, the analyses were repeated. By doing so, we confirmed a significant

increase in C_t-value. (*D. immitis* paired t-test, t[19]=2.159, p=0.0439). To determine if we

would obtain similar results for samples with pre-ML C_t-values <35 in the *Wolbachia* qPCR,

the analyses were repeated. This confirmed a significant increase in C_t-value. (*Wolbachia*

qPCR paired t-test, t[18]=9.290, p <0.0001). Similarly, paired t-tests of the raw *D. immitis*

and *Wolbachia* qPCR data showed statistically significant increases in C_t-values from pre-

ML to second post-ML sampling (Figure 6C,D).

3.5. Illumina PCR amplicon NGS identification of ML-susceptible *D. immitis* in

Australian dogs

Illumina NGS amplicon metabarcoding from 64 extracted microfilariae DNA samples from

37 dogs and one DNA sample from an adult *D. immitis* (P6/20D, positive control) was

performed to determine the nucleotide proportions at 7 SNPs of interest (65 DNA samples×7

loci = 455 PCRs) (Figure 7). Successful amplification of the target region was achieved for

96.5% (439/455) of PCR samples. Three DNA samples had loci which failed the first-stage

PCR amplification and were not submitted for Illumina NGS amplicon sequencing (HW34:

1/7 SNPs submitted, 0 microfilariae/mL; HW50: 5/7 SNPs submitted, 1 microfilariae /mL;

HW51: 1/7 SNPs submitted, 0 microfilariae/mL). Only 5/7 SNPs were submitted for P6/20D.

FastQ files were obtained, and SNP calling was successful for 98.2% (431/439) of submitted

PCR samples. Four DNA samples had loci which failed to produce reads for SNP calling

(HW29: 6/7 SNPs successful; HW34: 0/1 SNPs successful; HW50: 0/5 SNPs successful;

HW51: 0/1 SNPs successful). For the purpose of reproducibility, an additional 28 PCR

samples were submitted as 'repeats'. These repeat samples demonstrated identical outcomes

in terms of SNP calling. Overall, a total of 8,742,348 merged non-chimeric paired reads were

obtained to acquire amplicon sequence variants (ASVs) which were matched to the reference

ML 'susceptible' and ML 'resistant' allele for their respective SNP (Figure 7).

3.5.1. *L42411 (Di.2.2.scaf00001:466197)*

A total of 1,368,512 merged non-chimeric paired reads were obtained for L42411 with an

average of 20,426 paired reads per sample. Overall, 62/65 DNA samples displayed 100% of

the susceptible nucleotide at the suspect position. All 62 samples had 100% of their reads

assigned to a single ASV (L42411_ASV1) which was a perfect match to the ML 'susceptible'

reference sequence.

3.5.2. L21554 (nDi.2.2.scaf00046:222254)

A total of 1,167,570 merged, non-chimeric paired reads for 362 ASVs were obtained for

L21554 with an average of 17,690 paired reads per sample. Overall, 62/65 DNA samples

displayed 100% of the susceptible nucleotide at the suspect position. Of these 62 samples, 14

had 100% of their reads assigned to a single ASV (L21554_ASV1) which was a perfect

match to the 'susceptible' reference sequence. The remaining 48 samples contained

L21554_ASV1 along with low levels of other ASVs (L21554_ASV2, L21554_ASV3) which

possessed a SNP at another position in the sequence.

3.5.3. L45689 (nDi.2.2.scaf00185:10639)

A total of 1,664,418 merged, non-chimeric paired reads for 101 ASVs were obtained for

L45689 with an average of 23,777 paired reads per sample. Overall, 61/65 DNA samples had

100% of the susceptible nucleotide at the suspect position. Of these 61 samples, 15 had 100%

of their reads assigned to a single ASV (L45689_ASV1) which was a perfect match to the

'susceptible' reference sequence. The remaining 46 samples contained multiple ASVs

(L45689_ASV1,   L45689_ASV2,   L45689_ASV3,   L45689_ASV4,   L45689_ASV8,

L45689_ASV10, L45689_ASV11) which possessed 1-2 SNPs at other positions in the

sequence. One sample (Dog 25 – HW39) contained low levels (3.8%) of L45689_ASV6

which possessed a nucleotide of unknown ML association at the suspect position (Figure 8).

3.5.4. L9400 (nDi.2.2.scaf00005:662854)

A total of 908,148 merged, non-chimeric paired reads for 543 ASVs were obtained for L9400

with an average of 12,613 paired reads per sample. Overall, 62/65 DNA samples displayed

100% of the susceptible nucleotide at the suspect position. Of these 62 samples, 57 had 100%

of their reads assigned to a single ASV (L9400_ASV1) which was a perfect match to the

'susceptible' reference sequence. The remaining 5 samples contained L9400_ASV1 along

with other ASVs (L9400_ASV2, L9400_ASV30) which possessed a SNP at another position

in the sequence.

3.5.5. *L20587 (nDi.2.2.scaf00185:62174)*

A total of 1,600,533 merged, non-chimeric paired reads for 90 ASVs were obtained for

L20587 with an average of 23,537 paired reads per sample. Overall, 62/65 DNA samples

displayed 100% of the susceptible nucleotide at the suspect position. Of these 62 samples, 57

had 100% of their reads assigned to a single ASV (L20587_ASV1) which was a perfect

match to the 'susceptible' reference sequence. The remaining five samples contained

L20587_ASV1 along with other ASVs (L20587_ASV2, L20587_ASV4, L20587_ASV5,

L20587_ASV6, L20587_ASV8, L20587_ASV11) which possessed a SNP at another position

in the sequence.

3.5.6. *L15709_A (nDi.2.2.scaf00046:76278)*

A total of 1,526,468 merged, non-chimeric paired reads for 78 ASVs were obtained for

L15709_A with an average of 24,620 paired reads per sample. Overall, 61/65 DNA samples

displayed 100% of the susceptible nucleotide at the suspect position. All 61 samples had

100% of their reads assigned to a single ASV. Thirty samples had all reads assigned to

L15709A_ASV1 which was a perfect match to the 'susceptible' reference sequence. The

other 31 samples had all reads assigned to L15709A_ASV2 which possessed a deletion

elsewhere in the sequence.

3.5.7. *L30575 (nDi.2.2.scaf00046:22857)*

A total of 506,699 merged, non-chimeric paired reads for 542 ASVs were obtained for L30575 with an average of 8,173 paired reads per sample. Overall, 60/65 DNA samples displayed 100% of the susceptible nucleotide at the suspect position. All 60 samples had 100% of their reads assigned to a single ASV (L30575_ASV1) which was a perfect match to the 'susceptible' reference sequence.

4. Discussion

The use of MLs over the past 30 years has successfully reduced the prevalence of canine heartworm infection (Orr et al., 2020; Lau et al., 2021). However, the re-emergence of *D. immitis* infection in Queensland, Australia and the confirmation of ML-resistant isolates in the USA poses an important question of whether ML-resistance exists in Australia (Blagburn et al., 2011; Pulaski et al., 2014; Bourguinat et al., 2015; Blagburn et al., 2016; Nguyen et al., 2016; Maclean et al., 2017; McTier et al., 2017b; Orr et al., 2020; Panetta et al., 2021). By considering microfilariae counts as the phenotype and genotypic tests adopted from the USA, this study did not unequivocally demonstrate the presence of ML-resistant *D. immitis* in Australia. Although the sequential Modified Knott's tests showed delayed clearance of circulating microfilariae compared to the > 90% reduction at 28 days described by McTier et al. (2017a), suggestive of suspect ML-resistance, no genotypic evidence was discovered. Our approach was in part modelled to that of Ballesteros et al. (2018), where 27 dogs underwent 'microfilarial suppression testing' *sensu* Geary et al. (2011) and SNP analysis with moxidectin (Elanco, Advantage Multi for dogs, spot on). By doing so, they validated the association between ML microfilaricidal response phenotype and SNP loci genotype (Ballesteros et al., 2018). The current study obtained a comparable sample size to Ballesteros

et al. (2018), and adopted their 5-SNP predictive model which had 94.1% sensitivity and 100% specificity. We utilised Illumina amplicon metabarcoding coupled with DADA2 analysis to detect the susceptible and resistant SNPs, which provided us with reproducible results directly comparable to Ballesteros et al. (2018).

To treat canine heartworm disease, infected dogs are administered doxycycline and ML at the time of diagnosis (Day 0) to target the *Wolbachia* endosymbiont of *D. immitis*, prevent further infection by eliminating *D. immitis* L3/L4 and if possible (depending on the specific ML used) decrease circulating microfilariae (Nelson et al., 2020). To target the adult stages of *D. immitis*, clinicians in Australia typically follow the three-dose protocol recommended by the American Heartworm Society (Nelson et al., 2020). Importantly, for over 10 years, Australian veterinarians had access to a yearly injectable moxidectin (ProHeart SR-12, Zoetis) preventative (Mwacalimba et al., 2021). ProHeart SR-12 does not have a claim for circulating microfilariae in Australia, but it has been shown to reduce circulating microfilariae by >90% 28 days post-administration (McTier et al., 2017a).

The utilised sequential Modified Knott's test to calculate the reduction is not identical to the 'microfilarial suppression testing' *sensu* Geary et al. (2011) with spot-on moxidectin in the USA based on the label claim for microfilariae and 99% clearance within 14 days in an experimental trial (Geary et al., 2011; Bowman et al., 2015). We extended the post-ML interval to ~30 days for two reasons. Firstly, to suit the veterinary practice when owners brought their dog back for a health check-up, and secondly to utilise the published microfilaricidal effect of ProHeart SR-12 (McTier et al., 2017a). While the choice of ML-preventative was at the veterinarian's discretion, most were injected with ProHeart SR-12 in our study. The failure of 16 dogs (ML at diagnosis; moxidectin microspheres: 14 dogs, ivermectin: 2 dogs) to reduce microfilaremia by >90% was considered phenotypic evidence of suspect ML-resistance. This finding was not corroborated by genetic SNP analysis or

phenotypic testing beyond 30 days, as dogs that were re-tested eventually reached a >90%

reduction in microfilaremia by Day 55 to Day 136. Delays in reaching >90% microfilarial

reduction following ML treatment have been previously observed. In McTier et al. (2017a),

single doses of ProHeart 6 (Zoetis, moxidectin microspheres, 170 µg/kg) and oral moxidectin

(250 µg/kg) on Day 0 resulted in ≤90% reductions in mean microfilariae counts for the ML-

resistant isolate ZoeMO-2012 at Day 28 (ProHeart 6: 86.5%, oral moxidectin: 83.6%),

whereas ProHeart 12 (Zoetis, moxidectin microspheres, equivalent to ProHeart SR-12

registered in Australia, 500 µg/kg) led to a 91.2% reduction in microfilariae. At Day 84, all

three MLs reached >90% reductions in microfilariae (ProHeart 6: 96.8%, ProHeart 12:

96.9%, oral moxidectin: 92.6%) (McTier et al., 2017a). In contrast to the experimental trial

by McTier et al. (2017a), the current study sampled heterogeneous privately-owned dogs

which most likely carried a diverse population of *D. immitis*..

The variability of reduction in microfilariae counts post-ML administration in the present

study can be attributed to the varying effectiveness of MLs to clear circulating microfilariae

(Prichard, 2021). In Australia, no product has a label claim for microfilariae, unlike in the

USA where Advantage Multi for dogs (10% imidacloprid + 2.5% moxidectin topical

solution) has such a claim and was therefore used by Ballesteros et al. (2018) in their study

(Bowman et al., 2015). A microfilaricidal effect was noted early on for ivermectin in dogs at

a dose of 50 µg/kg, but due to toxicity in dolichocephalic breeds, this high dose is not on any

label (Blair and Campbell, 1979). The current preventative dose against <30-day old *D.

immitis* with ivermectin is only 6 µg/kg, which is considered insufficient to clear

microfilariae efficiently. Milbemycin oxime at 500 µg/kg is considered microfilaricidal, but

no registered product currently has such a claim (Prichard, 2021). Moxidectin in the spot-on

formulation (Elanco, Advantage Multi for dogs/Advocate for dogs at a minimum dose of

2,500 µg/kg) was demonstrated to be effective against microfilariae, which has been claimed

as such in the Advantage Multi for dogs (Elanco) marketed in the USA (McCall et al., 2014; Bowman et al., 2015).

Sequential quantitative Modified Knott's tests to enumerate the number of circulating *D. immitis* microfilariae pre- and post-ML treatment enables clinicians to determine whether moxidectin treatment is successfully clearing microfilaremia in the infected dog and whether the dog remains a threat to other dogs. However, performing a non-quantitative Knott's test is considered impractical for clinicians as it can be time-consuming, labour-intensive, and prone to human error. Alternative tools that are more rapid and accurate are required to measure changes in post-treatment microfilaremia levels. The blood samples in this study were screened for *D. immitis* using DiroCHEK® and qPCR targeting *D. immitis* and its associated *Wolbachia* endosymbiont. Based on our data, performing DiroCHEK® on neat or heated plasma was not suitable for the assessment of ML-treatment efficacy due to its lower correlation with the Modified Knott's test. The *D. immitis* and *Wolbachia* qPCRs produced higher correlations with the Modified Knott's test and were able to demonstrate the success of ML-treatment for reducing microfilariae levels. We therefore suggest clinicians to use these faster molecular tools as a substitute for the Modified Knott's test when monitoring the success of MLs in canine heartworm cases. The *D. immitis*-specific qPCR detected a larger proportion of infected dogs in this study compared to the *Wolbachia*-specific qPCR and may therefore be a better tool for measuring microfilariae levels. Future studies will benefit from a triplex qPCR targeting *D. immitis*, its associated *Wolbachia*, and dog DNA to reduce time and cost. The same *Wolbachia* qPCR assay was recently used to monitor the *Wolbachia* levels in dogs from Italy following doxycycline and Advocate (moxidectin) treatment (Laidoudi et al., 2020; Louzada-Flores et al., 2022). By doing so, they found that all dogs tested negative for *Wolbachia* after one month. Contrastingly, the current study saw only 8/28 dogs test negative for *Wolbachia* approximately 30 days after ML-treatment, with 4 of these

dogs already being negative on Day 0. When infected dogs were re-sampled on Day 30+, only 6/9 tested negative for *Wolbachia*.

The SNPs analysed in Ballesteros et al. (2018) were able to distinguish between ML-susceptible and ML-resistant *D. immitis* in the USA. However, in the current study and Lau et al. (2021), no evidence of ML-resistance in Australian *D. immitis* was found using these same ML resistance-associated SNPs. In Lau et al. (2021), the genome data of five adult *D. immitis* exhibited marked genetic differences compared to those previous phenotypically characterised *D. immitis* isolates from the USA. Genomics data from ML-susceptible and ML-resistant *D. immitis* are required to associate and specifically predict the ML-resistance in clinical cases in Australia. This will enable the early detection of ML resistance in Australia, allowing clinicians, researchers and the animal pharmaceutical industry to develop effective control measures to limit and manage the spread of resistant parasites.

Successful prevention of *D. immitis* heartworm can only be achieved with rigor in prevention due to inability to successfully remove older than 30-day *D. immitis* with current ML-based preventatives (Prichard, 2021). For veterinarians, however, the inquiry into rigor with ML-prevention will always be difficult with products administered by owners as they can unintentionally skip dosing if dogs regurgitate the dose (Atkins et al., 2014). An unintentionally missed ML-dose may lead to infection – apparent lack of efficacy (LOE) reports - despite owners' assumed rigor with prevention. In our study, despite owners' claims that dogs received 'rigorous' heartworm prevention with either moxidectin, milbemycin oxime or ivermectin for 12 months prior to the veterinarian's diagnosis, 16 dogs tested positive for canine heartworm infection. Similar findings were encountered in Nguyen et al. (2016) where microfilaraemic dogs in Queensland previously received 'rigorous' prevention. In this study, owner compliance could not be confirmed and was solely based on the owner's reports.

5. Conclusion

To confirm the presence of drug resistance in a parasite population, one must demonstrate that resistance is genetically inherited (Kohler, 2001). Since 2005, lack of efficacy complaints for heartworm preventatives led to speculation and controversy on the potential emergence of ML-resistant *D. immitis* in the USA (Hampshire, 2005; Atkins et al., 2014). It was not until 2014 that the first ML-resistant strain (LSU 10) was isolated in the USA and resistance to MLs was unequivocally confirmed (Pulaski et al., 2014). To do this, a laboratory dog was experimentally infected with L3s obtained from a field case, and then treated with ivermectin (Pulaski et al., 2014). Infective larvae from this dog were then successfully passaged into a second laboratory dog, which confirmed the genetic heritability of drug resistance (Pulaski et al., 2014). In Australia, we currently lack these types of specialised facilities where laboratory dogs can be experimentally infected with canine heartworm. Due to this limitation, it remains unknown whether ML-resistance is present in Australia, much like the 2005-2014 time period in the USA. Until we can demonstrate the heritability of ML-resistance in Australian *D. immitis* isolates via experimental infection studies, we can only speculate and test for resistance using the available tools as a proxy.

Acknowledgements

We thank all the veterinary practitioners and owners of the patients, in particular Dr. Tony Phillis and his team at Greencross Deeragun for valiant effort to recruit patients and collect samples for screening. The work was in part supported by the Canine Research Foundation and Dogs Victoria, Australia and by the Australian Companion Animal Health Foundation Research Grant. RP is supported by an Australian Government Research Training Program

Stipend and The Jean Walker Trust Fellowship. We thank Ilze Nel and Richard L'Estrange

(Zoetis Australia Pty Ltd) for discussion around heartworm prevention.

Supplementary Tables

Supplementary Table S1: Primers used throughout the study

Supplementary Table S2: SNPs in this study adopted from previous 5-SNPs models from

the USA

Supplementary Table S3: Summary of Australian dogs infected with *Dirofilaria immitis*

monitored over 2019-2022

References

Atkins, C.E., Murray, M.J., Olavessen, L.J., Burton, K.W., Marshall, J.W., Brooks, C.C., 2014. Heartworm 'lack of effectiveness' claims in the Mississippi delta: computerized analysis of owner compliance-2004-2011. Vet. Parasitol. 206, 106-113. doi: 10.1016/j.vetpar.2014.08.013

Ballesteros, C., Pulaski, C.N., Bourguinat, C., Keller, K., Prichard, R.K., Geary, T.G., 2018. Clinical validation of molecular markers of macrocyclic lactone resistance in *Dirofilaria immitis*. Int. J. Parasitol. Drugs Drug Resist. 8, 596-606. doi: 10.1016/j.ijpddr.2018.06.006

Blagburn, B.L., Arther, R.G., Dillon, A.R., Butler, J.M., Bowles, J.V., von Simson, C., Zolynas, R., 2016. Efficacy of four commercially available heartworm preventive products against the JYD-34 laboratory strain of *Dirofilaria immitis*. Parasit. Vectors 9, 191-191. doi: 10.1186/s13071-016-1476-7

Blagburn, B.L., Dillon, A.R., Arther, R.G., Butler, J.M., Newton, J.C., 2011. Comparative efficacy of four commercially available heartworm preventive products against the MP3 laboratory strain of *Dirofilaria immitis*. Vet. Parasitol. 176, 189-194. doi: 10.1016/j.vetpar.2010.12.049

Blair, L.S., Campbell, W.C., 1979. Efficacy of avermectin B1a against microfilariae of *Dirofilaria immitis*. Am. J. Vet. Res. 40, 1031-1032. doi:

Bourguinat, C., Keller, K., Prichard, R.K., Geary, T.G., 2011. Genetic polymorphism in *Dirofilaria immitis*. Vet. Parasitol. 176, 368-373. doi: 10.1016/j.vetpar.2011.01.023

Bourguinat, C., Keller, K., Xia, J., Lepage, P., McTier, T.L., Woods, D.J., Prichard, R.K., 2017. Genetic profiles of ten *Dirofilaria immitis* isolates susceptible or resistant to macrocyclic lactone heartworm preventives. Parasit. Vectors 10, 504-504. doi: 10.1186/s13071-017-2428-6

Bourguinat, C., Lee, A.C., Lizundia, R., Blagburn, B.L., Liotta, J.L., Kraus, M.S., Keller, K.,

Epe, C., Letourneau, L., Kleinman, C.L., Paterson, T., Gomez, E.C., Montoya-Alonso, J.A.,

Smith, H., Bhan, A., Peregrine, A.S., Carmichael, J., Drake, J., Schenker, R., Kaminsky, R.,

Bowman, D.D., Geary, T.G., Prichard, R.K., 2015. Macrocyclic lactone resistance in

Dirofilaria immitis: Failure of heartworm preventives and investigation of genetic markers

for resistance. Vet. Parasitol. 210, 167-178. doi: 10.1016/j.vetpar.2015.04.002

Bowman, D.D., Charles, S.D., Arther, R.G., Settje, T., 2015. Laboratory evaluation of the

efficacy of 10 % imidacloprid + 2.5 % moxidectin topical solution (Advantage® Multi,

Advocate®) for the treatment of *Dirofilaria immitis* circulating microfilariae in dogs.

Parasitol. Res. 114 Suppl 1, S165-174. doi: 10.1007/s00436-015-4522-z

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P.,

2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods

13, 581-583. doi: 10.1038/nmeth.3869

Diakou, A., Prichard, R.K., 2021. Concern for *Dirofilaria immitis* and macrocyclic lactone

loss of efficacy: current situation in the USA and Europe, and future scenarios. Pathogens 10,

1323. doi: 10.3390/pathogens10101323

Geary, T.G., Bourguinat, C., Prichard, R.K., 2011. Evidence for macrocyclic lactone

anthelmintic resistance in *Dirofilaria immitis*. Top. Companion Anim. Med. 26, 186-192. doi:

10.1053/j.tcam.2011.09.004

Genchi, C., Kramer, L.H., 2020. The prevalence of *Dirofilaria immitis* and *D. repens* in the

Old World. Vet. Parasitol. 280, 108995. doi: 10.1016/j.vetpar.2019.108995

Godel, C., Kumar, S., Koutsovoulos, G., Ludin, P., Nilsson, D., Comandatore, F., Wrobel,

N., Thompson, M., Schmid, C.D., Goto, S., Bringaud, F., Wolstenholme, A., Bandi, C., Epe,

C., Kaminsky, R., Blaxter, M., Mäser, P., 2012. The genome of the heartworm, *Dirofilaria*

immitis, reveals drug and vaccine targets. FASEB J. 26, 4650-4661. doi: 10.1096/fj.12-

205096

Hampshire, V.A., 2005. Evaluation of efficacy of heartworm preventive products at the FDA.

Vet. Parasitol. 133, 191-195. doi: 10.1016/j.vetpar.2005.04.004

Knott, J., 1939. A method for making microfilarial surveys on day blood. Trans. R. Soc.

Trop. Med. Hyg. 33, 191-196. doi: 10.1016/S0035-9203(39)90101-X

Kohler, P., 2001. The biochemical basis of anthelmintic action and resistance. Int. J.

Parasitol. 31, 336-345. doi: 10.1016/S0020-7519(01)00131-X

Laidoudi, Y., Davoust, B., Varloud, M., Niang, E.H.A., Fenollar, F., Mediannikov, O., 2020.

Development of a multiplex qPCR-based approach for the diagnosis of *Dirofilaria immitis*,

D. repens and *Acanthocheilonema reconditum*. Parasit. Vectors 13, 319. doi:

10.1186/s13071-020-04185-0

Lau, D.C.-W., McLeod, S., Collaery, S., Peou, S., Tran, A.T., Liang, M., Šlapeta, J., 2021.

Whole-genome reference of *Dirofilaria immitis* from Australia to determine single nucleotide

polymorphisms associated with macrocyclic lactone resistance in the USA. Curr. Res.

Parasitol. Vector. Borne Dis. 1, 100007. doi: 10.1016/j.crpvbd.2021.100007

Louzada-Flores, V.N., Kramer, L., Brianti, E., Napoli, E., Mendoza-Roldan, J.A., Bezerra-

Santos, M.A., Latrofa, M.S., Otranto, D., 2022. Treatment with doxycycline is associated

with complete clearance of circulating *Wolbachia* DNA in *Dirofilaria immitis*-naturally

infected dogs. Acta Trop. 232, 106513. doi: 10.1016/j.actatropica.2022.106513

Maclean, M.J., Savadelis, M.D., Coates, R., Dzimianski, M.T., Jones, C., Benbow, C.,

Storey, B.E., Kaplan, R.M., Moorhead, A.R., Wolstenholme, A.J., 2017. Does evaluation of

in vitro microfilarial motility reflect the resistance status of *Dirofilaria immitis* isolates to

macrocyclic lactones? Parasit. Vectors 10, 480-480. doi: 10.1186/s13071-017-2436-6

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet J. 17, 10-12. doi: 10.14806/ej.17.1.200

McCall, J.W., Arther, R., Davis, W., Settje, T., 2014. Safety and efficacy of 10%

imidacloprid+2.5% moxidectin for the treatment of *Dirofilaria immitis* circulating

microfilariae in experimentally infected dogs. Vet. Parasitol. 206, 86-92. doi:

10.1016/j.vetpar.2014.09.011

McTier, T.L., Kryda, K., Wachowski, M., Mahabir, S., Ramsey, D., Rugg, D., Mazaleski, M.,

Therrien, C., Adams, E., Wolff, T., Bowman, D.D., 2019a. ProHeart® 12, a moxidectin

extended-release injectable formulation for prevention of heartworm (*Dirofilaria immitis*)

disease in dogs in the USA for 12 months. Parasit. Vectors 12, 369-369. doi:

10.1186/s13071-019-3632-3

McTier, T.L., Pullins, A., Inskeep, G.A., Gagnon, G., Fan, H., Schoell, A., Bidgood, T.,

Login, J., Meeus, P., 2017a. Microfilarial reduction following ProHeart® 6 and ProHeart®

SR-12 treatment in dogs experimentally inoculated with a resistant isolate of *Dirofilaria*

immitis. Parasit. Vectors 10, 485. doi: 10.1186/s13071-017-2430-z

McTier, T.L., Six, R.H., Pullins, A., Chapin, S., Kryda, K., Mahabir, S.P., Woods, D.J.,

Maeder, S.J., 2019b. Preventive efficacy of oral moxidectin at various doses and dosage

regimens against macrocyclic lactone-resistant heartworm (*Dirofilaria immitis*) strains in

dogs. Parasit. Vectors 12, 444-444. doi: 10.1186/s13071-019-3685-3

McTier, T.L., Six, R.H., Pullins, A., Chapin, S., McCall, J.W., Rugg, D., Maeder, S.J.,

Woods, D.J., 2017b. Efficacy of oral moxidectin against susceptible and resistant isolates of

Dirofilaria immitis in dogs. Parasit. Vectors 10, 482. doi: 10.1186/s13071-017-2429-5

Mitra, A., Skrzypczak, M., Ginalski, K., Rowicka, M., 2015. Strategies for achieving high

sequencing accuracy for low diversity samples and avoiding sample bleeding using illumina

platform. PLoS One 10, e0120520. doi: 10.1371/journal.pone.0120520

Mwacalimba, K., Wright, A., Giannakakis, K., L'Estrange, R., Nguyen, T.S., 2021.

Corrigendum: Retrospective Analysis of Heartworm (Dirofilia immitis) Prevention

Medication Compliance and Economic Value in Dogs in Veterinary Practices in Australia.

Front Vet Sci 8, 759537. doi: 10.3389/fvets.2021.759537

Mylonakis, M.E., Papadopoulos, E., Koutinas, A.F., Paitaki, C., Leontides, L., 2004.

Comparative methodology for the detection and differentiation of circulating microfilariae of

Dirofilaria immitis in the dog. J. Helminthol. 78, 137-140. doi: 10.1079/JOH2003210

Nelson, D.C.T., McCall, D.J.W., Jones, D.S., Moorhead, D.A., 2020. Current canine

guidelines for the prevention, diagnosis, and management of heartworm (*Dirofilaria immitis*)

infection in dogs. American Heartworm Society, Wilmington, DE.

Nguyen, C., Koh, W.L., Casteriano, A., Beijerink, N., Godfrey, C., Brown, G., Emery, D.,

Šlapeta, J., 2016. Mosquito-borne heartworm *Dirofilaria immitis* in dogs from Australia.

Parasit. Vectors 9, 535. doi: 10.1186/s13071-016-1821-x

Noack, S., Harrington, J., Carithers, D.S., Kaminsky, R., Selzer, P.M., 2021. Heartworm

disease - Overview, intervention, and industry perspective. Int. J. Parasitol. Drugs Drug

Resist. 16, 65-89. doi: 10.1016/j.ijpddr.2021.03.004

Orr, B., Ma, G., Koh, W.L., Malik, R., Norris, J.M., Westman, M.E., Wigney, D., Brown, G.,

Ward, M.P., Šlapeta, J., 2020. Pig-hunting dogs are an at-risk population for canine

heartworm (*Dirofilaria immitis*) infection in eastern Australia. Parasit. Vectors 13, 69. doi:

10.1186/s13071-020-3943-4

Panetta, J.L., Calvani, N.E.D., Orr, B., Nicoletti, A.G., Ward, M.P., Šlapeta, J., 2021.

Multiple diagnostic tests demonstrate an increased risk of canine heartworm disease in

northern Queensland, Australia. Parasit. Vectors 14, 393-393. doi: 10.1186/s13071-021-

04896-y

Power, R.I., Calvani, N.E.D., Nachum-Biala, Y., Salant, H., Harrus, S., Šlapeta, J., 2021.

Adaptation of *gltA* and *ssrA* assays for diversity profiling by Illumina sequencing to identify

Bartonella henselae, *B. clarridgeiae* and *B. koehlerae*. J. Med. Microbiol. 70. doi:

10.1099/jmm.0.001400

Prichard, R.K., 2021. Macrocyclic lactone resistance in *Dirofilaria immitis*: risks for

prevention of heartworm disease. Int. J. Parasitol. 51, 1121-1132. doi:

10.1016/j.ijpara.2021.08.006

Pulaski, C.N., Malone, J.B., Bourguinat, C., Prichard, R., Geary, T., Ward, D., Klei, T.R.,

Guidry, T., Smith, G.B., Delcambre, B., Bova, J., Pepping, J., Carmichael, J., Schenker, R.,

Pariaut, R., 2014. Establishment of macrocyclic lactone resistant *Dirofilaria immitis* isolates

in experimentally infected laboratory dogs. Parasit. Vectors 7, 494-494. doi: 10.1186/s13071-

014-0494-6

RStudio Team, 2022. RStudio: Integrated development environment for R. RStudio, PBC,

Boston, MA.

Simón, F., Siles-Lucas, M., Morchón, R., González-Miguel, J., Mellado, I., Carretón, E.,

Montoya-Alonso Jose, A., 2012. Human and animal dirofilariasis: the emergence of a

zoonotic mosaic. Clin. Microbiol. Rev. 25, 507-544. doi: 10.1128/CMR.00012-12

Legends to Figures

Fig. 1. Location of dogs infected with *Dirofilaria immitis* included in this study from Eastern

Australia. These dogs (n=44) were sampled between 2019-2022 for presence of *D. immitis*

antigen, *D. immitis* DNA and associated *Wolbachia* DNA. The red circles indicate a location

that is associated with the town name and the number indicates the number of *D. immitis*-

positive dogs using any test. On the right, two close-ups of Townsville, Queensland and

Greater Sydney, New South Wales are shown. Image from Google Earth; data from SIO,

NOAA, U.S Navy, NGA, GEBCO (Image Landsat/Copernicus, CNES/Airbus, Maxar

Technologies).

Fig. 2. Diagnostic methods for the diagnosis of *Dirofilaria immitis* infection and response to

macrocyclic lactone administration in dogs. Blood samples were collected at the time of

diagnosis, followed by the administration of doxycycline antibiotic and macrocyclic lactone,

and then the administration of adulticide injections. Each blood sample was assayed using a

Modified Knott's test to quantify microfilariae, microfilariae were extracted from blood via

filtration for DNA isolation, plasma samples were assayed for the presence of *D. immitis*

antigen and DNA isolated directly from blood was assayed quantitatively for the presence of

D. immitis DNA and endosymbiotic *Wolbachia* DNA. DNA isolated from filtered

microfilariae were tested for the presence of *D. immitis* DNA and endosymbiotic *Wolbachia*

DNA in addition to single nucleotide polymorphisms (SNPs) associated with macrocyclic

lactone resistance from the USA. SNPs were assayed using Illumina amplicon metabarcoding

next generation sequencing (NGS).

Fig. 3. Sequential Modified Knott's tests performed on on 21 dogs positive for *Dirofilaria*

immitis. (A) Dogs (n=7) that exhibited <90% (red square %) microfilarial reduction at the

first post-ML sampling, but consecutive second post-ML sampling demonstrated >90%

microfilarial reduction (green square %). (B) Dog 25 at the first post-ML sampling had an

increased number of microfilariae, but one week later the microfilariae were reduced by

<90% (red squares); however, if comparing the 1st and 2nd post-ML samples, the reduction

was >90%. Dog 35 has had its 1st sample post-ML collected on day 7. (C) Dogs (n=9) that

were sampled only once post-ML with variable reduction of microfilariae. (D) Dogs (n=3)

that were sampled once, yet this sampling was delayed to 70 or more days post-ML. QLD –

Queensland, Australia, NSW – New South Wales, Australia, yellow star – dogs with at least

two consecutive post-ML blood samples tested, ML – macrocyclic lactone.

Fig. 4. *Dirofilaria immitis* antigen testing (DiroCHEK[®] ELISA, Zoetis, Australia) on 76

blood samples collected from 44 dogs in Australia. (A) Neat plasma samples from 41/44 dogs

and heated plasma samples from 43/44 dogs initially tested positive for *D. immitis* antigen.

Out of all blood samples tested, 67/76 neat samples and 75/76 heated samples were positive

for *D. immitis* antigen. A threshold of 3×SD (Mean[optical density of negative controls]) was

applied to each run to distinguish between *D. immitis*-negative and *D. immitis*-positive

samples. Thresholds of 0.0784 (HW1-HW10), 0.0596 (HW11-HW17, HW19), 0.0786

(HW20-HW49, HW51-HW57), and 0.0532 (HW58-HW63, HW65-HW75, HW77-HW79)

were applied. (B) Paired t-test on DiroCHEK[®] ELISA data demonstrated a significant

difference in optical density between neat and heated plasma samples (Paired t-test, $t[74]=$

4.220, $p < 0.0001$). Data analysis and graphs were produced using GraphPad Prism (version

9.3.1).

Fig. 5. Correlations between microfilariae count (mf/mL), DiroCHEK[®] ELISA (Zoetis,

Australia) of neat and heated plasma, and qPCR of whole blood DNA targeting *D. immitis*

and associated *Wolbachia*. (A) Moderate positive correlations between microfilariae counts

and OD_{620} of neat (n=76) and heated (n=76) plasma. (B) Moderate positive correlations

between microfilariae counts and OD_{620} of neat (n=71) and heated (n=71) plasma for samples

with microfilariae counts <20,000 mf/mL. (C) Strong negative correlation between *D.*

immitis qPCR C_t-values of whole blood DNA and microfilariae counts (n=72) for samples

with <20,000 mf/mL. (D) Strong negative correlation between *Wolbachia* qPCR C_t-values of

whole blood DNA and microfilariae counts (n=72) for samples with <20,000 mf/mL. (E)

Moderate negative correlation between *D. immitis* qPCR C_t-values of whole blood DNA and

microfilariae counts (n=77). (F) Moderate negative correlation between *Wolbachia* qPCR of

whole blood DNA and microfilariae counts (n=77). (G) Strong positive correlation between

D. immitis and *Wolbachia* qPCR C_t-values of whole blood DNA (n=78). (H) Moderate

positive correlation between *D. immitis* and *Wolbachia* qPCR C_t-values of whole blood DNA (n=52) for samples with C_t-values <40. Data analysis and graphs were produced using GraphPad Prism (version 9.3.1).

Fig. 6. Paired t-test analyses on *D. immitis* and *Wolbachia* qPCR C_t-values pre- and post-ML treatment. (A) *D. immitis* qPCR demonstrated statistically significant increase in C_t-value between pre-ML and first post-ML sampling (n=23). (B) *Wolbachia* qPCR demonstrated statistically significant increase in C_t-value between pre-ML and first post-ML sampling (n=23). (C) *D. immitis* qPCR demonstrated statistically significant increase in C_t-value between pre-ML and second post-ML sampling (n=9). (D) *Wolbachia* qPCR demonstrated statistically significant increase in C_t-value between pre-ML and second post-ML sampling (n=9). Data analysis and graphs were produced using GraphPad Prism (version 9.3.1).

Fig. 7. Australian dogs infected with *Dirofilaria immitis* and response to ML treatment. Dog identification number is indicated at the top. Upper-most bar chart represents log transformed microfilariae (mf) counts pre-ML and post-ML (black to grey; Modified Knott's test). Yellow stars – dogs with at least two consecutive post-ML blood samples tested (2+ post-ML), blue stars – dogs with one post-ML blood sample tested (1 post-ML). Percent (%) reduction of microfilariae at the first and second post-ML time point is plotted (purple bars, n/a – not applicable due to absence of sequential sample). Percentage (%) reduction at 1^{st} post-ML timepoint is written along the year these samples were collected. Nucleotide proportions at SNP regions (identified according to nDi2.2. genome assembly contigs) are shown for all samples analysed. The blue, red, and orange represent the nucleotides, with the reference nucleotides associated with ML-resistance indicated for each SNP on the right. Dog 25 has microfilariae count and L45689 highlighted by dotted line rectangles.

834　**Fig 8.** Nucleotide percentages at a SNP region (L45689) previously associated with ML-

835　resistance in the USA for Dog 25. Blood samples were collected at 0-, 30- and 36-days post-

836　ML treatment and DNA was isolated from filtered *Dirofilaria immitis* microfilariae. The

837　blue, red, and orange boxes in the key represent the nucleotides and their associated ML-

838　resistance status.

Table 1. Microfilariae quantification, antigen and qPCR diagnostics on *Dirofilaria immitis* infected dogs from Australia

| Dog | Location (Australia) | *D. immitis* prevention | | Microfilariae reduction | | | DiroCHEK® ELISA | | qPCR C$_t$ | |
| | | Compliance (past 12 months) | ML at diagnosis | pre-ML/post-ML | | % mf change | pre-ML/post-ML | | pre-ML/post-ML | |
				ID	mf/ml		Neat	Heat	*D. immitis*	*Wolbachia*
>90% microfilariae (mf) reduction										
1	Cairns, QLD	Sporadic	HeartGard Plus	HW1/HW2	1/0	-100	+/-	+/+	37.7/-	-/-
15	Deeragun, QLD	No	ProHeart SR-12	HW25/HW36	380/0	-100	+/-	+/+	26.0/-	-/-
21	Tugun, QLD	Yes	HeartGard Plus	HW31/HW34	175/0	-100	-/+	+/+	32.2/32.1	-/-
38	Deeragun, QLD	n/a	ProHeart SR-12	HW63/HW75	1,500/0	-100	+/+	+/+	23.4/-	29.5/-
16	Deeragun, QLD	n/a	ProHeart SR-12	HW26/HW33	7,625/600	-92	+/+	+/+	23.1/25.3	28.1/34.7
5	Deeragun, QLD	Sporadic	ProHeart SR-12	HW7/HW13	38,250/3,512	-91	+/+	+/+	20.0/22.4	25.1/32.3
<90% microfilariae (mf) reduction										
33	Deeragun, QLD	Yes	ProHeart SR-12	HW55/HW59	1,570/470	-70	+/+	+/+	24.6/25.2	30.3/-
13	Cairns, QLD	No	HeartGard Plus	HW20/HW22	7,450/2,475	-67	+/+	+/+	20.5/21.8	25.2/32.8
6	Deeragun, QLD	Yes	ProHeart SR-12	HW8/HW15	7,560/2,518	-67	+/+	+/+	21.3/22.3	26.7/32.0
34	Deeragun, QLD	Yes	ProHeart SR-12	HW56/HW58	9,400/3,180	-66.	+/+	+/+	21.2/22.9	27.3/35.5
28	Westbrook, QLD	No	ProHeart SR-12	HW44/HW49	3,180/1,100	-65	+/+	+/+	25.1/24.9	28.8/33.7
27	Westbrook, QLD	No	ProHeart SR-12	HW43/HW48	3,420/1,350	-61	+/+	+/+	25.8/23.9	29.8/33.8
37	Deeragun, QLD	Sporadic	ProHeart SR-12	HW61/HW65	2,300/960	-58	+/+	+/+	22.6/25.4	29.9/-
35	Deeragun, QLD	Yes	ProHeart SR-12	HW57/HW62	11,990/7,760	-35	+/+	+/+	19.5/22.7	27.3/32.6
40	Nana Glen, NSW	No	ProHeart SR-12	HW68/HW71	3,410/2,380	-30	+/-	+/+	22.7/27.0	28.2/-
24	Deeragun, QLD	Yes	ProHeart SR-12	HW38/HW40	420/300	-29	+/+	+/+	26.6/26.0	31.7/33.5
4	Deeragun, QLD	Sporadic	ProHeart SR-12	HW6/HW12	2,856/2,217	-22	+/+	+/+	22.0/22.9	26.6/32.3
3	Deeragun, QLD	Yes	ProHeart SR-12	HW5/HW14	9,540/8,208	-14	+/+	+/+	22.7/24.4	26.1/35.1
43	Deeragun, QLD	n/a	ProHeart SR-12	HW74/HW78	2,590/2,590	0	+/+	+/+	24.7/22.7	30.6/33.1
19	Tugun, QLD	Yes	HeartGard Plus	HW29/HW35	0/0	n/a	+/-	+/+	33.1/-	-/-
Increase in microfilariae (mf)										
7	Townsville, QLD	n/a	ProHeart SR-12	HW9/HW16	38/565	+1,386	+/+	+/+	28.7/24.0	31.8/34.3
2	Thursday Is, QLD	No	HeartGard Plus	HW3/HW4	510/1436	+182	+/+	+/+	21.8/22.9	27.0/31.9
25	Deeragun, QLD	Yes	ProHeart SR-12	HW39/HW41	24,800/6,216 0	+151	+/+	+/+	21.3/24.1	26.1/28.8

Note: n/a – not available, + (DiroCHEK) – positive, - (DiroCHEK) – negative, - (qPCR) – negative (>40), ML – macrocyclic lactone, mf – microfilariae, HeartGard Plus (ivermectin/pyrantel; Boehringer-Ingelheim), ProHeart SR-12 (moxidectin; Zoetis).

Fig. 2

Fig. 3

A

QLD Dog 3* | QLD Dog 6* | QLD Dog 24* | QLD Dog 27* | QLD Dog 28* | QLD Dog 34* | NSW Dog 40*

B

QLD Dog 25* | QLD Dog 35*

C

QLD Dog 2** | QLD Dog 4* | QLD Dog 5* | QLD Dog 7* | QLD Dog 13** | QLD Dog 16* | QLD Dog 21** | QLD Dog 37* | QLD Dog 43*

D

QLD Dog 15* | QLD Dog 33* | QLD Dog 38*

Prevention rigor declared by owner:

R Rigorous prevention   S Sporadic prevention   N No prevention

ML received at diagnosis: * Moxidectin   ** Ivermectin

Number of microfilaria counts post-ML administraton:
⭐ (cyan) 1 post-ML sample
⭐ (yellow) 2+ post-ML samples

Fig. 4

Fig. 5

Fig. 6

A

Paired t-test
t[22]=2.515, p=0.0197

C_t *D. immitis*

pre-ML   post-ML #1   diff.

Mean of differences

B

Paired t-test
t[22]=6.893, p<0.0001

C_t *Wolbachia*

pre-ML   post-ML #1   diff.

Mean of differences

C

C_t *D. immitis*

Pired t-test:
t[8]=5.483, p=0.0006

pre-ML   post-ML #2   diff.

Mean of differences

D

Paired t-test:
t[8]=4.361, p= 0.0024

C_t *Wolbachia*

pre-ML   post-ML #2   diff.

Mean of differences

Fig. 7

Fig. 8