1 **The Golgi complex is a regulatory hub for homologous recombination-mediated DNA**
2 **repair.**
3

4 George Galea[1*], Karolina Kuodyte[1,2], Muzamil M. Khan[1,3], Peter Thul[4], Beate Neumann[5],
5 Emma Lundberg[4,6,7], Rainer Pepperkok[1,3,5*]
6
7
[1]Cell Biology and Biophysics Unit, European Molecular Biology Laboratory (EMBL), 69117
Heidelberg, Germany
[2]Collaboration for joint PhD degree between EMBL and Heidelberg University, Faculty of
Biosciences
[3]Translational Lung Research Center Heidelberg, German Center for Lung Research,
Heidelberg, Germany
[4]Science for Life Laboratory, School of Engineering Sciences in Chemistry, Biotechnology and
Health, KTH - Royal Institute of Technology, Stockholm, 17121, Sweden.
[5]Advanced Light Microscopy Facility, EMBL, Heidelberg, 69117, Heidelberg, Germany
[6]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA
[7]Department of Pathology, Stanford University, Stanford, CA 94305, USA
19
20
21 *corresponding authors
22 e-mail: george.galea@embl.de; rainer.pepperkok@embl.org

Abstract

Cells are constantly exposed to a multitude of DNA damaging agents that can lead to mutation, dysregulation, and possibly cell death. To ensure genomic integrity, DNA Damage Response (DDR) mechanisms are set in motion to repair and mitigate any damage to the DNA structure. Although these pathways are well-studied in the context of nuclear function, relatively little is known of the regulatory function of cytoplasmic organelles. Here we show the first example of DDR regulation at the Golgi complex, coordinating Homologous Recombination (HR)-mediated DNA repair. We found that RAD51C, a regulatory HR protein, localises to the Golgi and nuclear compartments and in response to double-strand DNA breaks, the Golgi protein population of RAD51C redistributes to form nuclear foci. Furthermore, we found that the Golgi localisation of RAD51C is dependent on the Golgin Giantin. Depletion of Giantin induces the redistribution of the RAD51C Golgi pool to form nuclear foci, independent of DNA damage induction, and concurrent with a significant increase in genomic instability and inhibition of HR signalling regulators. This study presents evidence for a novel pathway where the Golgi is a central regulatory hub for HR DDR and potentially other repair pathways, a finding with important therapeutic implications.

Introduction

DNA damage response (DDR) is extensively studied and its many pivotal roles in preservation of genomic integrity such as DNA damage sensing, repair, cell cycle arrest and apoptosis regulation are relatively well-characterised (Chatterjee & Walker, 2017; Ciccia & Elledge, 2010). However, most of this research has exclusively focused on nuclear events, while cytoplasmic organelles' response and regulatory function have only begun to be elucidated. Nevertheless, a few studies have hinted at a link between cytoplasmic organelles, genomic stability and, in turn, cancer, with the Golgi complex emerging as a central theme (Kulkarni-Gosavi et al., 2019; Petrosyan, 2015; Zhang, 2021). The Golgi is an important homeostasis hub, where a multitude of signalling pathways and essential cellular processes intersect (Makhoul et al., 2018; Wilson et al., 2011). Therefore, it is not unexpected that perturbations to the genome would impact the Golgi complex. At a structural level, the Golgi undergoes dramatic morphological changes after the induction of DNA lesions, from ribbon-like perinuclear stack to dispersed fragments (Farber-Katz et al., 2014). A response requiring the phosphorylation of the Golgi oncoprotein Golgi phosphoprotein 3 (GOLPH3) by DDR regulator DNA-dependent protein kinase (DNA-PK). The Golgi dispersion leads to changes in trafficking patterns and is associated with cell survival, although our understanding of this process is still limited (Farber-Katz et al., 2014). Golgi morphology alterations are also a common feature across a wide variety of cancer types and are often reflected in changes in the distribution of Golgi resident proteins (Petrosyan, 2015; Zhang, 2021). Rearrangement of Golgi glycosyltransferases distribution is not uncommon in cancer cells, resulting in defective glycosylation, a process thought to promote cancer development (Bui et al., 2021; Petrosyan, 2015; Zhang, 2021). It remains unclear what the impacts of these Golgi organisational changes are and whether they're part of a response to DNA damage or a cause of carcinogenesis. Further studies are needed to identify the linking pathways that coordinate DDR and Golgi-associated molecular pathways, such as cell cycle progression, apoptosis and membrane trafficking, that ensure cellular homeostasis is maintained during DNA damage.

In an effort to understand the impact of the Golgi complex on DDR, we investigated the
regulation of homologous recombination (HR)-mediated DNA repair through the activity of
RAD51C. The protein is part of the RAD51 recombinase protein family, which is a key active
complex in the recombination-mediated repair of double-strand DNA breaks (Rein et al., 2021;
Sullivan & Bernstein, 2018). RAD51C along with the other RAD51 paralogs assemble into
subcomplexes aiding in the regulation of RAD51 activity, although the precise cellular function
of these paralogs and recruitment mechanisms remain unclear (Sullivan & Bernstein, 2018).
RAD51C is required for DDR signalling through the activation of the checkpoint kinase CHK2
and cell cycle arrest in response to DNA damage and it has been postulated that it can regulate
the recruitment and translocation of the HR complex via its nuclear localization signal (Badie
et al., 2009; French et al., 2003; Gildemeister et al., 2009). The correct functioning of this
protein is crucial for the regulation of HR, with mutations in this gene leading to Fanconi
anaemia-like syndrome (chromosomal-instability disorder associated with a variety of
developmental abnormalities, bone marrow failure and predisposition to leukaemia and other
cancers) and high susceptibility to ovarian and breast cancer (Clague et al., 2011; Loveday et
al., 2012; Somyajit et al., 2010; Vaz et al., 2010).
86
Here, we show the first example of a mechanistic model (**Fig. 1a**) for DDR regulation at the
Golgi complex, activating HR-mediated DNA repair through the interaction of RAD51C and
Giantin. We find that RAD51C, a regulatory HR protein, localises to the Golgi and nuclear
compartments and in response to double-strand DNA breaks, the RAD51C Golgi population
redistributes to form nuclear foci. This response requires importin-beta mediated nuclear
import and the phosphorylation of ATM protein kinase. We further show that the RAD51C
Golgi localisation is dependent on the transmembrane Golgi protein, Giantin (GOLGB1).
Depletion of Giantin induces the redistribution of the RAD51C Golgi population to form nuclear
foci, independent of DNA damage induction. This resulted in a significant increase in genomic
instability concurrently with DNA damage response signalling inhibition. Thus, we identify a
novel regulatory pathway for the timely control and activation of HR DNA repair at the Golgi
complex following DNA damage. We postulate that this novel Golgi-DDR pathway could
extend to other types of DNA repair mechanisms as we validate the localisation of 15 other
Golgi-localising DDR proteins associated with the diverse DNA repair processes.
101
102 **Results**
103
104 **RAD51C Golgi population redistributes to the nucleus in response to DNA damage**
105 **events.**
106
To examine the role of the Golgi complex in homologous recombination (HR)-mediated DNA
repair, we examined the role of the HR regulatory repair protein, RAD51C. We began by
examining the localisation of the protein using immunofluorescence staining in standard model
cell lines, cervical carcinoma HeLa Kyoto (HK) cells, human bone osteosarcoma epithelial
cells, U-2 OS and RAD51C high-expressing, human mammary gland epithelial
adenocarcinoma cell line, MCF7 (**Fig. 1b; Extended Data Fig. 1a**). We observed that
endogenous RAD51C protein was present in a juxta-nuclear compact localisation that
colocalized with the Golgi marker, GM130 and a diffused nuclear distribution with some foci
present (**Fig. 1b**). This localisation pattern was reproduced in all the tested cell lines
(**Extended Data Fig. 1a**). The specificity of the antibody was tested by the depletion of
RAD51C (**Extended Data Fig. 1b-d**). Next, to biochemically validate the subcellular

localisation of RAD51C, we isolated nuclear, membrane and cytoplasmic fractions from HK protein extracts. Fractions were verified using well-established markers (GM130 for Golgi membranes, Lamin B1 for the nuclear compartment, alpha-tubulin for the cytoplasmic fraction) and probed for RAD51C presence in each compartment (**Fig. 1c**). RAD51C was found to be present in all fractions, in agreement with our immunofluorescence data (**Fig. 1b**).

Since RAD51C has been reported to be recruited to DNA repair foci upon the introduction of Double-Strand DNA Breaks (DSBs) (Badie et al., 2009), we induced the formation of these DNA breaks by treating cells with doxorubicin (DOX) and tested whether the Golgi population of RAD51C responds to DNA damage. Upon the addition of the drug (**Fig. 1d-g**), we observed an overall decrease in RAD51C protein level, with the portion of the RAD51C colocalising with the Golgi marker, GM130, reducing at a higher rate than in the nuclear compartment. The RAD51C diffused nuclear pattern changed into very distinct nuclear foci, with this redistribution getting more pronounced with increasing length of treatment (**Fig. 1d, Golgi fraction marked with yellow arrows**). Analysis of these experiments allowed us to measure the changes in RAD51C levels at both the Golgi (**Fig. 1e**) and the nuclear compartment (**Fig. 1f**), and to calculate a RAD51C Golgi-nuclear distribution ratio (**Fig. 1g**). This ratio allowed us to measure robustly and quantitatively the Golgi-nuclear distribution of RAD51C, despite fluctuations of protein levels due to the treatment. After DOX treatment, the ratio value is reduced with increasing length of time when compared to the control, as normalised to 1; giving the RAD51C distribution Golgi:nuclear ratio: 0.86 at 1 h, 0.52 at 2 h, 0.41 at 3 h and 0.32 at 4 h. These results were further validated by subcellular fractionation (**Extended Data Fig. 1e, f**). Similarly, the total protein level of RAD51C is seen to decrease after 3h treatments, with a much larger reduction in the membrane compartment when compared to the nuclear fraction. In both experiments, the RAD51C distribution ratio is seen to decrease significantly after 3 h DOX treatment to 0.41 and 0.69 for the immunofluorescence and biochemical assay, respectively (**Fig. 1g, Extended Data Fig. 1f).**

To further probe this change in the localisation pattern of RAD51C and identify the mechanism of protein redistribution from one compartment to another, we introduced DSBs using DOX while inhibiting the Importin-β-mediated nuclear import, utilising the small peptide inhibitor Importazole (IPZ) (Soderholm et al., 2011). Cells were treated with DMSO (solvent control) or IPZ for 30 min before treatment with DOX (**Fig. 2a, Golgi fraction marked with yellow arrows**). Treatment with IPZ inhibited the previously described DOX-RAD51C localisation pattern change, instead, the majority of the protein population colocalized with the Golgi marker, GM130, and significant inhibition of RAD51C nuclear foci formation was observed (**Fig. 2a-c**). Quantification of the images (**Fig. 2b, c**) showed that the combinatory treatment of DOX and IPZ significantly inhibited the RAD51C protein redistribution with a calculated RAD51C distribution ratio of 0.68, while the treatment with DOX only resulted in a ratio of 0.31. A reduction in the overall RAD51C protein level was observed when cells were treated with IPZ only, but there was no significant change in the RAD51C distribution pattern, where the distribution ratio was calculated to be 0.83.

Having established that the RAD51C Golgi population responds to DSB-inducing treatments, we sought to test whether the phosphorylation of DDR kinases mediates this redistribution. The three main kinases that are active in response to DNA damage are ataxia-telangiectasia mutated (ATM), Ataxia telangiectasia and Rad3-related (ATR) and DNA-dependent protein kinase (DNA-PK) (Ciccia & Elledge, 2010). We treated cells with phosphorylation inhibitors

specific for each kinase and analysed RAD51C redistribution after DOX treatment (**Fig. 2d-f**).
The combination treatment of DOX with the ATM phosphorylation inhibitor, KU55933 (Hickson
et al., 2004) significantly inhibited the redistribution of RAD51C, with most of the protein
population still colocalising with the Golgi marker, GM130 and a calculated RAD51C
distribution ratio of 0.77. Whereas the combined treatment of DOX with ATR and DNA-PK
phosphorylation inhibitor, VE-821 (RAD51C distribution ratio of 0.27) (Fokas et al., 2012) and
NU7441 (RAD51C distribution ratio of 0.26) (Tavecchio et al., 2012) respectively, had no
apparent impact on the redistribution of RAD51C when compared with cells treated with DOX
only (RAD51C distribution ratio of 0.31) (**Fig. 2c, Golgi fraction marked with yellow arrows).**
No apparent changes in the RAD51C localisation pattern were observed with the treatment of
inhibitors only (**Extended Data Fig. 2**).
177
To ensure that these observations are a direct result of DSBs and not the drug itself, we next
investigated the effects of other DSB-causing drugs (Jekimovs et al., 2014): camptothecin
(CPT) (**Extended Data Fig. 3a-c**), etoposide (ETO) (**Extended Data Fig. 3d-f**), and mitomycin
C (MMC) (**Extended Data Fig. 3g-i**). All treatments showed a significant change in RAD51C
distribution, from the Golgi to the nuclear compartment. Furthermore, treatment with CPT
(**Extended Data Fig. 3a**) led to a change in RAD51C distribution in the nuclear compartment,
from a diffused distribution to distinct nuclear foci, while the application of ETO and MMC
(**Extended Data Fig. 3d, g**) resulted in an increase in RAD51C nuclear population but with no
apparent nuclear foci pattern.
187
188 **RAD51C Golgi localisation is dependent on the Golgin Giantin.**
189
To identify potential Golgi candidates that could act as membrane anchors for RAD51C, we
analysed two genome-wide siRNA screens carried out to identify regulators of HR and DDR
(**Extended Data Fig. 4a, b**) (Adamson et al., 2012; Paulsen et al., 2009). Analysis of the
primary data from these screens highlighted the Golgin protein family as potential candidate,
where the knockdown of GMAP210 and Giantin lead to a significant reduction in HR repair
rates (Adamson et al., 2012) (**Extended Data Fig 4a**), with the depletion of Giantin also
leading to a significant inhibition of H2AX phosphorylation, an important DDR signalling
regulator (Paulsen et al., 2009) (**Extended Data Fig 4b**). The Golgin family is composed of
predominantly coiled-coil proteins that are anchored to the Golgi membrane by their carboxy
terminus and are predicted to adopt an extended conformation that projects into the
surrounding cytoplasm. This arrangement is ideal for the capture or tethering of nearby
membranes (Witkos & Lowe, 2015) and also could potentially act as a surface that retains
DDR proteins. Therefore, we tested whether the localization of RAD51C is dependent on these
two Golgins (**Fig. 3a-c**). We transfected HK cells with two siRNA targeting GMAP210 or
Giantin and monitored for any changes in RAD51C localization. Depletion of GMAP210 and
treatment with a control siRNA had no significant effect on RAD51C distribution whereas the
knock-down of the Golgi matrix protein Giantin led to a significant redistribution of RAD51C
(**Fig. 3a**). We observed a significant decrease in the colocalization of RAD51C with the Golgi
marker (**yellow arrows**), GM130 and a change in protein localisation in the nucleus from a
diffused distribution to distinct bright nuclear foci (**white arrows**) (**Fig. 3a**). Quantifications
showed a reduction of the RAD51C Golgi population (**Fig. 3b**) with the distribution ratio (**Fig.
3c**) reduced by more than half and the number of RAD51C foci (**Fig. 3d**) increased by more
than two-fold upon depletion of Giantin with either siRNAs. While there was no significant
difference in either RAD51C Golgi intensity, distribution ratio or foci number (**Fig. 3b-d**)

between the GMAP210-depleted and control cells. In light of these results, we next tested
whether RAD51C physically interacts or forms a complex with Giantin by performing an
immunoprecipitation (IP) assay (**Fig. 3e**). Here, we immunoprecipitated endogenous Giantin,
probed for RAD51C and vice versa (**Fig. 3e; IP: Giantin and IP: RAD51C**). Endogenous
Giantin and RAD51C were found to co-immunoprecipitate from HK protein extracts in both
conditions.
220
To gain mechanistic insight into the nature of these RAD51C foci induced by Giantin depletion,
we carried out colocalization assays to determine whether these structures contain standard
HR markers, phosphorylated H2AX (y-H2AX) (**Fig. 4a**) and phosphorylated ATM (p-ATM)
(**Fig. 4b**) under physiological conditions. Both proteins are well established markers for DSB
repair sites and are important for the recruitment of the HR repair machinery (Vítor et al.,
2020). In cells treated with a control siRNA, we observed that a large proportion of the
RAD51C nuclear foci were decorated with either y-H2AX or p-ATM (**Fig. 4a, b**; **colocalising**
foci are denoted with an arrow), as previously described in literature (Vítor et al., 2020).
RAD51C foci induced by the depletion of Giantin showed significantly lower colocalization with
both markers. Analysis of these images (**Fig. 4c, d**) showed that 48.6% and 39.2% of RAD51C
foci colocalized with y-H2AX and p-ATM respectively under control conditions, whereas
RAD51C foci resulting from the depletion of Giantin were found to colocalise significantly less
with both markers, with only 17.2% and 15.2% colocalising with y-H2AX and p-ATM
respectively.
235
236 **Depletion of Giantin leads to genomic instability and inhibition of DDR signalling**
237
Having tested the requirement of Giantin in the recruitment and localisation of RAD51C, we
next assessed whether the redistribution of the HR protein by the depletion of Giantin would
have any significant impact on the cells' ability to maintain genomic stability and undergo DNA
repair. To gauge the status of the cell genome, we applied the comet assay (Møller, 2018)
which provides a measure of fragmented genomic DNA, an indicator of genomic stability (**Fig.**
5a, b). Cells treated with DNA damaging agents or having impaired DNA repair pathways
would result in distinctive long comet tails, due to significant amounts of fragmented genomic
DNA. In contrast, healthy untreated cells would display shorter or no comet tail. In our
experiment (**Fig. 5a, b**), we observed that samples treated with siRNAs against Giantin and
RAD51C showed significantly longer comet tails when compared to the control siRNA
treatment. The depletion of RAD51C resulted in the strongest phenotype but was closely
followed by the results obtained by the Giantin knockdown. Quantifications of the comets
showed that the knockdown of RAD51C resulted in a 2.5-fold increase in the comet tail length
and fluorescence intensity, here quantified as a standard metric known as olive tail moment
(Olive et al., 2012). Depletion of Giantin using two different siRNAs (siGiantin#1 and
siGiantin#2) resulted in 1.8 and 1.6-fold change in the olive tail moment ratio. Next, we
validated this effect on genomic stability by measuring the incidence of micronuclei (denoted
with red arrows) (**Fig. 5c, d**). We therefore analysed cell populations depleted of Giantin and
RAD51C for their occurrence. Our results highlighted a significantly larger percentage of the
cell population displayed micronuclei formation and aberrant nuclear structure after the
depletion of Giantin and RAD51C when compared to our control samples. We calculated (**Fig.**
5d) that 5% of the population under control conditions displayed micronuclei when compared
with 12.5%, 9% and 10% in cells depleted of RAD51C, and two different siRNAs against
Giantin, respectively.

RAD51C is required for DSB repair by HR and therefore its mislocalization due to Giantin
depletion might lead to inhibition of this pathway and activation of the alternative repair system,
the Non-Homologous End Joining (NHEJ) repair (Brandsma & Gent, 2012). To gain insight
into the mechanism by which the redistribution of RAD51C leads to a decrease in genomic
stability, cells depleted of Giantin were assessed for their ability to phosphorylate and thereby
activate well-characterised regulators of HR and NHEJ repair pathways in response to DNA
damage (**Fig. 5e, f**). In our negative control, HK cells treated with DOXO and CPT, we observe
increased levels of phosphorylated ATM, CHK2, H2AX (**Fig.5e**) and DNA-PK (**Fig. 5f**) in
response to the treatment when compared to our solvent control. Cells depleted of the Golgi
protein Giantin when treated with DOXO and CPT showed significantly lower levels of
phosphorylation of the tested HR DDR proteins except for the NHEJ regulator, DNA-PK, when
compared to our control siRNA (**Fig. 5e, f**). Finally, to assess whether this effect is not a result
of cell cycle perturbations, we assessed the cell cycle profile of cells depleted of Giantin (**Fig.
5g**), using Fluorescence Activated Cell Sorting (FACS) analysis, where no significant effects
were observed.
277
278 **Antibody-based analysis identifies a network of DNA damage response proteins at the
279 Golgi complex.**
280
To identify other double-localising proteins, similar to RAD51C, which we postulate would link
cytoplasmic functions of the secretory pathway to nuclear events, we shortlisted 329 proteins
(339 antibodies) identified by the Human Protein Atlas (HPA) project (Thul et al., 2017) to
localise at both Golgi-like membranes and the cell nucleus (**Supplementary table S1**). To
confirm the localisation data and validate the specificity of the antibody binding, we measured
the changes of antibody staining at the Golgi and nuclear compartment after siRNA-mediated
knockdown of the target proteins. Candidate antibodies that displayed a 25% or more
decrease in fluorescent intensity in both compartments after depletion were considered to be
validated (Stadler et al., 2012) with in total 163 candidates (167 antibodies) validated.
(**Supplementary table S1**).
291
All the validated proteins were interrogated by STRING (Szklarczyk et al., 2021) and
Reactome gene enrichment database (Gillespie et al., 2022). Much to the confirmation of our
proposed hypothesis, two major protein networks were identified, a membrane trafficking and
a DDR cluster (**Fig. 6a**). The latter was found to be composed of 15 proteins that function in
various components of the core DNA repair complexes: HR DNA repair: RAD51C & NBN (**Fig.
6b**); Non-Homologous End Joining (NHEJ): POLQ & USP1; Mismatch DNA Repair (MMR):
MSH6 (**Fig. 6b**) & Base Excision Repair (BER): LIG1 & TOPORS, as well as other integral
regulators of DNA repair cellular response such as chromatin cohesion (ESCO2) (**Fig. 6a, b**),
ubiquitination (USP1), cell cycle (CCAR1) and signalling (PFDN6 & FMN2). To our knowledge,
these findings present the first validation of DNA repair protein clusters to localise to the Golgi
complex.
303
304 **Discussion**
305
The Golgi complex is a key regulatory hub with numerous cellular processes intersecting at
this organelle (Makhoul et al., 2018; Wilson et al., 2011). Therefore, understanding the role of
the Golgi in response to DNA damage is crucial for discerning the system-wide mechanism at
play for maintaining cellular homeostasis. Here we show a novel regulatory pathway that links

the Golgi complex and its cytoplasmic functions to DDR for the activation of HR-mediated DNA
repair. A mechanism driven by the Golgi localisation of RAD51C and its interaction with the
Golgin, Giantin. Previous studies have shown that RAD51C is essential for the recombination-
mediated repair of DSBs and is recruited to the sites of DNA lesions (Badie et al., 2009;
Gildemeister et al., 2009). In line with these findings, we show that the RAD51C Golgi
population redistributes and forms nuclear foci in response to the DNA damage via an
importin-beta mediated transport. Additionally, we show that Giantin is required to maintain
the Golgi localisation of RAD51C and the disruption of this interaction leads to inhibition of HR
DNA damage response, resulting in increased genomic instability. Consistent with previous
literature (Badie et al., 2009), we found that RAD51C Golgi population redistribution requires
ATM activity, although the mechanism of recruitment is still unclear, however, another DDR
kinase, DNA-PKcs (Farber-Katz et al., 2014), has been shown to interact with the Golgi
complex. It is possible that ATM functions in a similar manner for the recruitment of RAD51C
at the Golgi, with Giantin being a possible phosphorylation target as previously predicted (Cara
et al., 2016).
325
Here, we propose a model where the cytoplasmic tail of Giantin anchors RAD51C to the Golgi
complex, hiding its nuclear localisation sequence motif (French et al., 2003), which would
otherwise facilitate its shuttling from the Golgi to the nucleus. When required RAD51C would
be activated by the kinase, ATM and recruited to the DNA damage site, and engage other HR
repair proteins or further propagate DDR signalling. Consistent with our hypothesis, we found
that the mislocalization of RAD51C by Giantin depletion impairs HR signalling response and
results in a significant increase in genomic instability. Previous findings also have shown that
the depletion of Giantin inhibited both HR repair rates (Adamson et al., 2012) and
phosphorylation of the DDR signalling protein, H2AX (Paulsen et al., 2009). However, we
cannot exclude that the downregulation of Giantin has impacts on other nuclear events beyond
the redistribution of RAD51C, although none of our data and previous literature would suggest
this. Interestingly, other members of the Golgin family have also been reported to alter HR
repair rates (Adamson et al., 2012) and phosphorylation of DDR signalling protein, H2AX
(Paulsen et al., 2009). Numerous studies have also implicated the role of the Golgins in cancer
progression (Baschieri et al., 2015; Bhat et al., 2017; Bui et al., 2021; Ghannoum et al., 2022)
and although further studies are required to clarify their role, these observations raise the
intriguing possibility that this protein family can be integral for DDR regulation.
343
Taking in the wide array of novel double-localising proteins identified in our study and their
respective roles in important cellular processes such as membrane trafficking, cell cycle and
DNA damage response, our results propose a high degree of interconnectivity between the
secretory membrane system and nuclear compartment. Although further studies are needed
to investigate Golgi-DDR pathway functions, our data strongly suggests that the Golgi
localisation of RAD51C is crucial for the correct functioning of DSB repair and it is tempting to
speculate that the interaction we find for RAD51C and Giantin could be extrapolated to the
other DDR proteins identified from this study and the Golgin protein family. When taking into
context the established role of the Golgi as a cellular hub for signalling, transport and post-
translational modification (Jackson, 2009; Wilson et al., 2011), it comes to no surprise that this
organelle could have a function in DNA damage response. The Golgi has already been shown
to be a pivotal link between nuclear and cytoplasmic processes such as cholesterol
homeostasis (Brown et al., 2018), trafficking events (Scharaw et al., 2016), and mitosis
(Colanzi & Sütterlin, 2013). Considering the proximity of the two organelles, the Golgi's cellular

influence in numerous pathways and unique compact architecture, the Golgi is ideal to connect
cytoplasmic processes to DDR.
360
Taken together, our study presents strong evidence for a novel Golgi regulatory pathway for
the coordination of HR-mediated repair through the activity of RAD51C and Giantin.
Furthermore, our localisation data also suggests that this regulation is not just restricted to HR
DNA repair but could extend to all other types of DNA repair mechanisms such as NHEJ,
MMR, and BER. This discovery has the potential to open new scientific fields from different
ends and has the potential to significantly change the way we understand DNA repair
regulation.
368
369 **Materials and Methods**
370
Antibodies and chemicals. Several commercially antibodies and chemical were used in the
paper, including RAD51C (ab72063, Abcam; 1:500 for immunostaining) & (ab95069, Abcam,
1:2000 for western blotting; 1:500 for immunoprecipitation), GOLGB1/Giantin (AF8159, R&D
systems; 1:500 for immunostaining and immunoprecipitation, 1:2000 for western blotting) &
(G1/M1, Enzo Life Sciences, 1:500 for immunostaining), GM130 (610822, BD Biosciences,
1:500 for immunostaining, 1:2000 for western blotting), ATM Ser1981 (MA1-2020, Invitrogen;
1:500 for immunostaining; 1:2000 for western blotting), CHK2 pThr68 (PA5-17818, Invitrogen;
1:2000 for western blotting), DNA-PKcs (Ser2056) (PA5-78130, Invitrogen, 1:2000 for western
blotting), gamma-H2AX pSer139 (613402, Biolegend; 1:500 for immunostaining, 1:2000 for
western blotting), alpha-tubulin (MS-581, Thermo Fisher; 1:10,000 for western blotting) Lamin
B1 (ab16048, Abcam; 1:2000 for western blotting) vinculin (ab219649, abcam; 1:2000 for
western blotting), doxorubicin (ab120629, Abcam); Importazole (SML0341, Sigma-Aldrich),
KU55933 (ATMi, SML1109, Sigma-Aldrich); ve-821 (ATRi, HY-14731; MedChemExpress)
NU7441 (DNA-PKi, HY-11006; MedChemExpress), etoposide (ab120227, Abcam), mitomycin
C (M7949, Sigma-Aldrich), and campothecin (ab120115, Abcam).
386
Cell lines, cell culture and siRNA transfection. HeLa Kyoto, U2-O S, MCF7, were cultured
in DMEM (Life Technologies) supplemented with 10% FBS (Invitrogen) and 1% L-glutamine
(Invitrogen). Cells were checked for mycoplasma contamination by PCR. siRNA transfections
were performed with Lipofectamine 2000 (Invitrogen) using SilencerSelect siRNAs (Ambion)
according to the manufacturer's instructions. Transfections were carried out for 72 h and the
final siRNA concentrations used were 15nM for all siRNAs. Giantin siRNA-1: 5951 & siRNA-
2: 5953; GMAP210 siRNA-1: s17811 & siRNA-2: s17812; RAD51C siRNA: s11737.
394
Drug treatments. Cells were treated 24 h after seeding. For camptothecin, etoposide and
mitomycin C experiments, cells were treated for 16 h at a concentration of 0.1 μM, 50μM and
5μM, respectively. After treatment cells were incubated with fresh medium for 2 hours.
Doxorubicin was used at a concentration of 40μM for 3 h unless indicated otherwise. For
kinase and importin-β inhibitor treatments, cells were pretreated for 30 min with the described
inhibitor prior to the addition of doxorubicin. The inhibitors were used at the following
concentrations: importazole (20μM), KU55933 (30μM), NU7441 (10μM) and ve-821 (10μM).
402
Immunofluorescence assay. Cells were fixed with 4% paraformaldehyde in PBS and
permeabilized with 0.1% Triton X-100 for 15 min at room temperature, then cells were blocked
with 5% bovine serum albumin in 0.05% Triton X-100 for 60 min and incubated with primary

antibodies in blocking buffer at room temperature for 3 h. Following 3 washes with PBS, cells
were incubated with fluorescent dye-conjugated secondary antibodies diluted in a blocking
buffer for 1 h at room temperature.
409
Image and data analysis. Confocal microscopy experiments were performed with fixed and
immunostained cell samples on a microscope FV3000, Olympus. Z stacks of images covering
the entire cell thickness were acquired. All image analysis was performed using Cell Profiler
(Carpenter et al., 2006) and Fiji (Schindelin et al., 2012). Briefly, first nuclei were segmented
in the Hoechst channel. When appropriate, the Golgi complex was segmented in the Golgi
marker channel and using the segmented nuclei as seeds, the two structures were associated.
Intensity profiles, morphology features and structuring counting analysis were performed when
required using Cell Profiler (Carpenter et al., 2006).
418
Comet assay. The assay was carried out as previously described (Vodenkova et al., 2020).
Briefly, cells were trypsinized, pelletized and resuspended in ice-cold PBS at a concentration
of 25×10^4 cells per ml. The cells were resuspended in 2% low melting agarose (Sigma) and
spread quickly onto gel bond film (Biozym) covered in 1% agarose (Sigma). Samples were
immersed into a lysis buffer (100mM EDTA, 2.5M NaCl, 10mM Tris-HCl and 1% Triton-X100;
pH 10) overnight at 4°C. Followed by a wash with ice cold water and run in an electrophoresis
chamber (alkaline buffer: 1mM EDTA, 300mM NaOH; pH 13) at 15V, 300mA for 60 min at
4°C. Slides were first washed in a Tris-HCl neutralisation buffer (0.4M; pH 7.5) followed by
water, stained with SYBR Gold (Thermo Fisher Scientific) (1:10000) and finally dried. Comets
were imaged by automated Olympus Scan^R screening microscope and comet tails scored
using OpenComet plugin (Gyori et al., 2014).
430
Western blotting analysis. HK cells were lysed using RIPA buffer (Thermo Scientific) with
complete protease inhibitor cocktail (Roche). SDS-PAGE was performed on pre-cast Tris-
Acetate gels followed by transfer to PVDF transfer membrane (Merck Millipore). Proteins were
detected using primary antibodies as described followed by incubation with secondary
antibodies coupled with HRP (Invitrogen). Detection of protein was performed using Pierce
ECL Plus Western Blotting Substrate reagent (Thermo Scientific) and visualised on Azure 280
chemiluminescent imaging system. Subcellular fractionation was performed using a
subcellular protein fractionation kit for cultured cells (Thermo scientific) according to the
manufacturer's instructions. Fractions were verified using well-established markers (GM130
for Golgi membranes, Lamin B1 for the nuclear compartment, alpha-tubulin for the cytoplasmic
fraction) and probed for RAD51C presence in each compartment. The RAD51C
membrane:nuclear distribution ratio was calculated by first dividing the RAD51C protein level
in each fraction by the protein level of the appropriate control, the resulting membrane (M_{R51C}
/ M_{GM130}) and nuclear (N_{R51C} / $N_{laminB1}$) ratios were further divided to give the ratio = (M_{R51C} /
M_{GM130}) / (N_{R51C} / $N_{laminB1}$). The ratios obtained from the control were normalised to 1 and
compared to the treated group.
447
Immunoprecipitation. HK cells were lysed using a lysis buffer (50mM Hepes, 130mM NaCl,
1mM DTT 1% NP-40) with complete protease inhibitor cocktail (Roche). Cell lysates were
centrifuged at 13,000 rpm for 10 mins at 4°C. For immunoprecipitation, the lysates were
incubated with the primary antibody described and rotated overnight at 4°C. Next, the lysates
were incubated with G-agarose beads (Roche) and rotated for 4 h at 4°C. The samples were

washed with cold lysis buffer and then precipitated proteins were eluted by 2x SDS sample
buffer and analysed by Western blotting.
455
Localisation validation siRNA assay. A custom-designed siRNA library targeting our
proteins of interest (Ambion) was designed and prepared in 96-well glass bottom plates
(Miltenyi Biotec) using a protocol for solid phase reverse transfection as previously described
(Erfle et al., 2008; Stadler et al., 2012). Non-targeting siRNA was used as a negative control.
72 hours after cell seeding, cells were fixed with 4% paraformaldehyde, permeabilized with
0.1% Triton-X100 and immunostained against the HPA antibodies of interest and a Golgi
marker, GM130. Hoechst 33342 was used as a nuclear stain. siRNAs and antibodies details
are available in **Supplementary table S1**. Images were acquired on a fully automated
Molecular Devices IXM with a 10x/0.45 NA P-APO objective. The resulting images were
analysed using Cell Profiler software (Carpenter et al., 2006) for quantitative and automated
measurements of fluorescence from the antibodies as previously described (Stadler et al.,
2012). Briefly, nuclei were segmented in the Hoechst channel, the Golgi complex was
segmented in the Golgi marker channel and using the segmented nuclei as seeds, the two
structures were associated. Intensity profiles of each compartment were acquired. Reduction
of 25% or more of the antibody staining in both Golgi and nuclear compartments was
considered as a validation of the antibody specificity (**Supplementary table S1**).
472
Statistical analysis. All data were obtained from at least three independent experiments if
not otherwise stated. Statistical analyses were performed using two tailed t-test for pairwise
comparison and one-way analysis of variance (ANOVA) for multiple comparisons on
GraphPad Prism 9. Data are expressed as standard error of the mean (s.e.m.). n values
indicate biologically independent samples and experiments. $P < 0.05$ was considered
statistically significant.
479

Acknowledgements. We thank the ALMF and the Pepperkok team for their support. In
addition, we acknowledge Jan Ellenberg, Claudia Lukas, and Simone Köehler for fruitful
discussion for the development of the project and manuscript. G.G. was supported by the
EMBL EIPOD programme, K.K. by the EMBL PhD programme and M.M.K. by the German
Centre for Lung Research..

Figure legend
486
Figure 1. **RAD51C localises to both the Golgi and nuclear compartment and**
redistributes in response to DNA damage events. (a) Proposed mechanistic model for the
regulation of HR-mediated repair through the activation of RAD51C at the Golgi complex. (b)
Representative images of HK cells stained with antibodies against RAD51C (green) and
GM130 (red). DNA was stained with Hoechst 33342 (blue). (c) Western blot showing the
subcellular membrane (M), nuclear (N) and cytoplasmic (C) fractions of RAD51C. (d) HK cells
treated with the DNA damage inducing drug, doxorubicin (DOX), for increasing lengths of time.
Yellow arrows denote the Golgi membrane; Scale bar 10 μm. (e) Quantification of sum
intensity of RAD51C Golgi population, (f) RAD51C nuclear population, and (g) a ratio of
RAD51C Golgi-nuclear distribution after treatment with DOX. Data represent the
mean ± standard error of the mean (s.e.m.). ($n = 3$ biologically independent samples).

Statistical significance was determined using one-way analysis of variance (ANOVA);
*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.001$.
500

Figure 2. Redistribution of RAD51C Golgi fraction is required for the formation of
RAD51C nuclear foci and is dependent on the kinase ATM. HK cells were stained with
antibodies against RAD51C and GM130. (a) HK cells were treated with DMSO or IPZ prior to
a 3-hour treatment with DOX. Yellow arrows denote the Golgi membrane. Results were
quantified as (b) relative sum intensity of RAD51C at the Golgi and (c) ratio of RAD51C Golgi-
nuclear distribution. (d) HK cells were treated with DMSO (control) alone or with ATM inhibitor
(KU55933) or ATR inhibitor (VE-821) or DNA-PK inhibitor (NU7441) prior to a 3-hour treatment
with doxorubicin. Results were quantified as an (e) relative sum intensity of RAD51C at the
Golgi and (f) ratio of RAD51C Golgi-nuclear distribution. Scale bar 10 μm. Data represent the
mean ± standard error of the mean (s.e.m.). ($n = 3$ biologically independent samples).
Statistical significance was determined using one-way analysis of variance (ANOVA);
*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.001$.
513

Figure 3. RAD51C Golgi localization and activation is dependent on the Golgin protein
Giantin. (a) HK cells were transfected with control, or Giantin siRNAs for 72 hours. The cells
were stained with antibodies against RAD51C and Golgi marker GM130. Yellow arrows
indicate the Golgi and white arrows indicate nuclear foci; scale bar 10 μm. Quantification of
the localisation changes of RAD51C were quantified as (b) relative sum intensity of RAD51C
at the Golgi, (c) fluorescent intensity ratio of the RAD51C Golgi-nuclear distribution and (d)
relative number of RAD51C foci per cell. Data represent the mean ± standard error of the mean
(s.e.m.). ($n = 3$ biologically independent samples). Statistical significance was determined
using one-way ANOVA; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.001$. (e)
Immunoprecipitation of Giantin with RAD51C.
524

Figure 4. Colocalization analysis of RAD51C nuclear foci induced by Giantin depletion
with DDR markers. (a-b) Co-localisation experiment of cells treated with control siRNA, or
Giantin siRNA. Cell stain with antibody against RAD51C (green) and HR DDR markers (red):
(a) y-H2AX, and (b) p-ATM. Co-localisation of structure is denoted by an arrow; Scale bar 10
μm. (i) Quantification of percentage RAD51C foci co-localising with y-H2AX and p-ATM. Data
represent the mean ± standard error of the mean (s.e.m.). ($n = 3$ biologically independent
samples). Statistical significance was determined using a two-tailed unpaired Student's t-test;
*$P < 0.05$.
533

Figure 5. Depletion of Giantin led to increased genomic instability and disruption of
DDR signalling. (a) Representative images of comet assay detection genomic DNA
fragmentation in HK cells treated with control, Giantin, or RAD51C targeting siRNA. (b)
Quantification of comet OliveMoment. (c, d) Representative images and quantification of
percentage cells displaying micronuclei after siRNA treatment; scale bar 10 μm. Red arrows
denote micronuclei structures. Data represent the mean ± standard error of the mean (s.e.m.).
($n = 3$ biologically independent samples). Statistical significance was determined using one-
way analysis of variance (ANOVA); *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.001$. (e, f) HK
cells transfected with control or Giantin siRNAs and treated with doxorubicin or camptothecin
after 72 h transfection, the extracts were prepared and immunoblotted as indicated. (g) Cell
cycle profile of HK cell treated with Giantin or control siRNA. Data represent the
mean ± standard error of the mean (s.e.m.). ($n = 3$ biologically independent samples).

Figure 6. **Antibody-based analysis identifies a network of DNA damage response proteins at the Golgi complex.** (a) Protein-protein interaction network showing the DDR proteins identified to localise to both the Golgi complex and nucleus; yellow nodes indicate double-localising proteins, grey nodes are filler untested proteins; experimentally-based string network. (b) Representative images showcasing the antibody-based localisation of 3 DDR proteins, displaying both nuclear and Golgi localisation. Scale bar 10 µm. Yellow arrows indicate the Golgi compartment.

EXTENDED DATA FIGURE 1. (a) Representative images showing U-2 OS and MCF7 cells stained with antibodies against RAD51C (green) and GM130 (red). DNA was stained with Hoechst 33342 (blue). (b) Representative images showing the RAD51C antibody specificity. HK cells are stained with antibodies against RAD51C and GM130; DNA was stained with Hoechst 33342. (c) Quantification of RAD51C sum intensity after RAD51C depletion. Data represent the mean ± standard error of the mean (s.e.m.). ($n = 3$ biologically independent samples). (d) Western blot showing the level of RAD51C protein level, in RAD51C-depleted and control cells. (e) Western blot showing the subcellular localization of RAD51C treated with the DMSO and doxorubicin. (f) Quantification of WB showing a ratio of RAD51C membrane-nuclear distribution. Data represent the mean ± standard error of the mean (s.e.m.). ($n = 2$ biologically independent samples). Statistical significance was determined using a two-tailed unpaired Student's *t*-test; *$P < 0.05$; ****$P < 0.001$.

EXTENDED DATA FIGURE 2. HK cells were stained with antibodies against RAD51C and GM130. HK cells were treated with DMSO (control) alone or with ATM inhibitor (KU55933) or ATR inhibitor (VE-821) or DNA-PK inhibitor (NU7441); scale bar 10 µm.

EXTENDED DATA FIGURE 3. HK cell stained with antibodies against RAD51C and GM130. Cells were treated with (a) Camptothecin (CPT), (d) etoposide (ETO) and (g) mitomycin C (MMC) for 16 h followed by media change for 2 h; scale bar 10 µm. Quantification of RAD51C percentage distribution between the Golgi and nuclear compartment after (b) CPT, (e) ETO and (h) MMC treatments. Quantification of RAD51C percentage distribution between the Golgi and nuclear compartment after (c) CPT, (f) ETO and (i) MMC treatments. treatments. Data represent the mean ± standard error of the mean (s.e.m.). ($n = 3$ biologically independent samples). Statistical significance was determined using a two-tailed unpaired Student's *t*-test; **$P < 0.01$; ****$P < 0.001$.

EXTENDED DATA FIGURE 4. (a) siRNA screen data (Adamson *et al*, 2012) showing the relative homologous recombination (HR) repair rate upon systematic knockdown of the Golgin protein family (grey) and HR complex proteins (black). (b) siRNA screen data (Paulsen et al., 2009) showing the relative percent cell population with phosphorylation of H2AX upon systematic knockdown of the Golgins. The datasets are normalised to the negative control set at 1. (c) HK cells were transfected with control, or Giantin siRNAs for 72 hours. The cells were stained with antibodies against Giantin and nuclei stained with Hoechst 33342; scale bar 10 µm.

References

Adamson, B., Smogorzewska, A., Sigoillot, F. D., King, R. W., & Elledge, S. J. (2012). A

genome-wide homologous recombination screen identifies the RNA-binding protein RBMX as
a component of the DNA-damage response. *Nature Cell Biology*, *14*(3), 318–328.

Badie, S., Liao, C., Thanasoula, M., Barber, P., Hill, M. A., & Tarsounas, M. (2009). RAD51C
facilitates checkpoint signaling by promoting CHK2 phosphorylation. *The Journal of Cell*
Biology, *185*(4), 587–600.

Baschieri, F., Uetz-von Allmen, E., Legler, D. F., & Farhan, H. (2015). Loss of GM130 in breast
cancer cells and its effects on cell migration, invasion and polarity. *Cell Cycle* , *14*(8), 1139–
1147.

Bhat, G., Hothpet, V.-R., Lin, M.-F., & Cheng, P.-W. (2017). Shifted Golgi targeting of
glycosyltransferases and α-mannosidase IA from giantin to GM130-GRASP65 results in
formation of high mannose N -glycans in aggressive prostate cancer cells. In *Biochimica et*
Biophysica Acta (BBA) - General Subjects (Vol. 1861, Issue 11, pp. 2891–2901).
https://doi.org/10.1016/j.bbagen.2017.08.006

Brandsma, I., & Gent, D. C. (2012). Pathway choice in DNA double strand break repair:
observations of a balancing act. *Genome Integrity*, *3*(1), 9.

Brown, M. S., Radhakrishnan, A., & Goldstein, J. L. (2018). Retrospective on Cholesterol
Homeostasis: The Central Role of Scap. *Annual Review of Biochemistry*, *87*, 783–807.

Bui, S., Mejia, I., Díaz, B., & Wang, Y. (2021). Adaptation of the Golgi Apparatus in Cancer
Cell Invasion and Metastasis. *Frontiers in Cell and Developmental Biology*, *9*, 806482.

Cara, L., Baitemirova, M., Follis, J., Larios-Sanz, M., & Ribes-Zamora, A. (2016). The ATM-
and ATR-related SCD domain is over-represented in proteins involved in nervous system
development. In *Scientific Reports* (Vol. 6, Issue 1). https://doi.org/10.1038/srep19050

Carpenter, A. E., Jones, T. R., Lamprecht, M. R., Clarke, C., Kang, I. H., Friman, O., Guertin,
D. A., Chang, J. H., Lindquist, R. A., Moffat, J., Golland, P., & Sabatini, D. M. (2006).
CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome*
Biology, *7*(10), R100.

Chatterjee, N., & Walker, G. C. (2017). Mechanisms of DNA damage, repair, and
mutagenesis. *Environmental and Molecular Mutagenesis*.
https://onlinelibrary.wiley.com/doi/abs/10.1002/em.22087

Ciccia, A., & Elledge, S. J. (2010). The DNA damage response: making it safe to play with
knives. *Molecular Cell*, *40*(2), 179–204.

Clague, J., Wilhoite, G., Adamson, A., Bailis, A., Weitzel, J. N., & Neuhausen, S. L. (2011).
RAD51C germline mutations in breast and ovarian cancer cases from high-risk families. *PloS*
One, *6*(9), e25632.

Colanzi, A., & Sütterlin, C. (2013). Signaling at the Golgi during mitosis. *Methods in Cell*
Biology, *118*, 383–400.

Erfle, H., Neumann, B., Rogers, P., Bulkescher, J., Ellenberg, J., & Pepperkok, R. (2008).
Work flow for multiplexing siRNA assays by solid-phase reverse transfection in multiwell
plates. *Journal of Biomolecular Screening*, *13*(7), 575–580.

Farber-Katz, S. E., Dippold, H. C., Buschman, M. D., Peterman, M. C., Xing, M., Noakes, C.
J., Tat, J., Ng, M. M., Rahajeng, J., Cowan, D. M., Fuchs, G. J., Zhou, H., & Field, S. J. (2014).
DNA damage triggers Golgi dispersal via DNA-PK and GOLPH3. *Cell*, *156*(3), 413–427.

Fokas, E., Prevo, R., Pollard, J. R., Reaper, P. M., Charlton, P. A., Cornelissen, B., Vallis, K.

A., Hammond, E. M., Olcina, M. M., Gillies McKenna, W., Muschel, R. J., & Brunner, T. B.
(2012). Targeting ATR in vivo using the novel inhibitor VE-822 results in selective sensitization
of pancreatic tumors to radiation. In *Cell Death & Disease* (Vol. 3, Issue 12, pp. e441–e441).
https://doi.org/10.1038/cddis.2012.181

French, C. A., Tambini, C. E., & Thacker, J. (2003). Identification of functional domains in the
RAD51L2 (RAD51C) protein and its requirement for gene conversion. *The Journal of*
Biological Chemistry, *278*(46), 45445–45450.

Ghannoum, S., Fantini, D., Zahoor, M., Reiterer, V., Phuyal, S., Netto, W. L., Sørensen, Ø.,
Iyer, A., Sengupta, D., Prasmickaite, L., Mælandsmo, G. M., Köhn-Luque, A., & Farhan, H.
(2022). A combined experimental-computational approach uncovers a role for the Golgi matrix
protein Giantin in breast cancer progression. In *bioRxiv* (p. 2022.04.25.489358).
https://doi.org/10.1101/2022.04.25.489358

Gildemeister, O. S., Sage, J. M., & Knight, K. L. (2009). Cellular Redistribution of Rad51 in
Response to DNA Damage. *The Journal of Biological Chemistry*, *284*(46), 31945–31952.

Gillespie, M., Jassal, B., Stephan, R., Milacic, M., Rothfels, K., Senff-Ribeiro, A., Griss, J.,
Sevilla, C., Matthews, L., Gong, C., Deng, C., Varusai, T., Ragueneau, E., Haider, Y., May,
B., Shamovsky, V., Weiser, J., Brunson, T., Sanati, N., … D'Eustachio, P. (2022). The
reactome pathway knowledgebase 2022. *Nucleic Acids Research*, *50*(D1), D687–D692.

Gyori, B. M., Venkatachalam, G., Thiagarajan, P. S., Hsu, D., & Clement, M.-V. (2014).
OpenComet: an automated tool for comet assay image analysis. *Redox Biology*, *2*, 457–465.

Hickson, I., Zhao, Y., Richardson, C. J., Green, S. J., Martin, N. M. B., Orr, A. I., Reaper, P.
M., Jackson, S. P., Curtin, N. J., & Smith, G. C. M. (2004). Identification and characterization
of a novel and specific inhibitor of the ataxia-telangiectasia mutated kinase ATM. *Cancer*
Research, *64*(24), 9152–9159.

Jackson, C. L. (2009). Mechanisms of transport through the Golgi complex. *Journal of Cell*
Science, *122*(Pt 4), 443–452.

Jekimovs, C., Bolderson, E., Suraweera, A., Adams, M., O'Byrne, K. J., & Richard, D. J.
(2014). Chemotherapeutic compounds targeting the DNA double-strand break repair
pathways: the good, the bad, and the promising. *Frontiers in Oncology*, *4*, 86.

Kulkarni-Gosavi, P., Makhoul, C., & Gleeson, P. A. (2019). Form and function of the Golgi
apparatus: scaffolds, cytoskeleton and signalling. *FEBS Letters*, *593*(17), 2289–2305.

Loveday, C., Turnbull, C., Ruark, E., Xicola, R. M. M., Ramsay, E., Hughes, D., Warren-Perry,
M., Snape, K., Breast Cancer Susceptibility Collaboration (UK), Eccles, D., Evans, D. G.,
Gore, M., Renwick, A., Seal, S., Antoniou, A. C., & Rahman, N. (2012). Germline RAD51C
mutations confer susceptibility to ovarian cancer [Review of *Germline RAD51C mutations*
confer susceptibility to ovarian cancer]. *Nature Genetics*, *44*(5), 475–476; author reply 476.

Makhoul, C., Gosavi, P., & Gleeson, P. A. (2018). The Golgi architecture and cell sensing.
Biochemical Society Transactions, *46*(5), 1063–1072.

Møller, P. (2018). The comet assay: ready for 30 more years. *Mutagenesis*, *33*(1), 1–7.

Olive, P. L., Banáth, J. P., & Durand, R. E. (2012). Heterogeneity in Radiation-Induced DNA
Damage and Repair in Tumor and Normal Cells Measured Using the "Comet" Assay. In
Radiation Research (Vol. 178, Issue 2, pp. AV35–AV42). https://doi.org/10.1667/rrav04.1

Paulsen, R. D., Soni, D. V., Wollman, R., Hahn, A. T., Yee, M.-C., Guan, A., Hesley, J. A.,
Miller, S. C., Cromwell, E. F., Solow-Cordero, D. E., Meyer, T., & Cimprich, K. A. (2009). A

genome-wide siRNA screen reveals diverse cellular processes and pathways that mediate
genome stability. *Molecular Cell*, *35*(2), 228–239.

Petrosyan, A. (2015). Onco-Golgi: Is Fragmentation a Gate to Cancer Progression?
Biochemistry & Molecular Biology Journal, *1*(1). https://doi.org/10.21767/2471-8084.100006

Rein, H. L., Bernstein, K. A., & Baldock, R. A. (2021). RAD51 paralog function in replicative
DNA damage and tolerance. *Current Opinion in Genetics & Development*, *71*, 86–91.

Scharaw, S., Iskar, M., Ori, A., Boncompain, G., Laketa, V., Poser, I., Lundberg, E., Perez, F.,
Beck, M., Bork, P., & Pepperkok, R. (2016). The endosomal transcriptional regulator RNF11
integrates degradation and transport of EGFR. *The Journal of Cell Biology*, *215*(4), 543–558.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri,
K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image
analysis. *Nature Methods*, *9*(7), 676–682.

Soderholm, J. F., Bird, S. L., Kalab, P., Sampathkumar, Y., Hasegawa, K., Uehara-Bingen,
M., Weis, K., & Heald, R. (2011). Importazole, a small molecule inhibitor of the transport
receptor importin-β. *ACS Chemical Biology*, *6*(7), 700–708.

Somyajit, K., Subramanya, S., & Nagaraju, G. (2010). RAD51C: a novel cancer susceptibility
gene is linked to Fanconi anemia and breast cancer. *Carcinogenesis*, *31*(12), 2031–2038.

Stadler, C., Hjelmare, M., Neumann, B., Jonasson, K., Pepperkok, R., Uhlén, M., & Lundberg,
E. (2012). Systematic validation of antibody binding and protein subcellular localization using
siRNA and confocal microscopy. *Journal of Proteomics*, *75*(7), 2236–2251.

Sullivan, M. R., & Bernstein, K. A. (2018). RAD-ical New Insights into RAD51 Regulation.
Genes, *9*(12). https://doi.org/10.3390/genes9120629

Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva, N. T.,
Legeay, M., Fang, T., Bork, P., Jensen, L. J., & von Mering, C. (2021). The STRING database
in 2021: customizable protein-protein networks, and functional characterization of user-
uploaded gene/measurement sets. *Nucleic Acids Research*, *49*(D1), D605–D612.

Tavecchio, M., Munck, J. M., Cano, C., Newell, D. R., & Curtin, N. J. (2012). Further
characterisation of the cellular activity of the DNA-PK inhibitor, NU7441, reveals potential
cross-talk with homologous recombination. *Cancer Chemotherapy and Pharmacology*, *69*(1),
155–164.

Thul, P. J., Åkesson, L., Wiking, M., Mahdessian, D., Geladaki, A., Ait Blal, H., Alm, T.,
Asplund, A., Björk, L., Breckels, L. M., Bäckström, A., Danielsson, F., Fagerberg, L., Fall, J.,
Gatto, L., Gnann, C., Hober, S., Hjelmare, M., Johansson, F., … Lundberg, E. (2017). A
subcellular map of the human proteome. *Science*, *356*(6340).
https://doi.org/10.1126/science.aal3321

Vaz, F., Hanenberg, H., Schuster, B., Barker, K., Wiek, C., Erven, V., Neveling, K., Endt, D.,
Kesterton, I., Autore, F., Fraternali, F., Freund, M., Hartmann, L., Grimwade, D., Roberts, R.
G., Schaal, H., Mohammed, S., Rahman, N., Schindler, D., & Mathew, C. G. (2010). Mutation
of the RAD51C gene in a Fanconi anemia–like disorder. *Nature Genetics*, *42*(5), 406–409.

Vítor, A. C., Huertas, P., Legube, G., & de Almeida, S. F. (2020). Studying DNA Double-Strand
Break Repair: An Ever-Growing Toolbox. *Frontiers in Molecular Biosciences*, *7*, 24.

Vodenkova, S., Azqueta, A., Collins, A., Dusinska, M., Gaivão, I., Møller, P., Opattova, A.,
Vodicka, P., Godschalk, R. W. L., & Langie, S. A. S. (2020). An optimized comet-based in vitro

DNA repair assay to assess base and nucleotide excision repair activity. *Nature Protocols*,
15(12), 3844–3878.

Wilson, C., Venditti, R., Rega, L. R., Colanzi, A., D'Angelo, G., & De Matteis, M. A. (2011).
The Golgi apparatus: an organelle with multiple complex functions. *Biochemical Journal*,
433(1), 1–9.

Witkos, T. M., & Lowe, M. (2015). The Golgin Family of Coiled-Coil Tethering Proteins.
Frontiers in Cell and Developmental Biology, *3*, 86.

Zhang, X. (2021). Alterations of Golgi Structural Proteins and Glycosylation Defects in Cancer.
Frontiers in Cell and Developmental Biology, *9*, 665289.

FIGURE 1

RAD51C — GM130 - nuclei

FIGURE 2

FIGURE 3

RAD51C – GM130 - nuclei

FIGURE 4

FIGURE 5

a

siNEG9   siGiantin#1   siGiantin#2   siRAD51C

b

Relative Olive tail moment

***
*
*

siNEG9
siGiantin#1
siGiantin#2
siRAD51C

c

siNEG9   siGiantin#1   siGiantin#2   siRAD51C

d

% cells with micronuclei

**
*
*

siNEG9
siGiantin#1
siGiantin#2
siRAD51C

e

	control	DOX	CPT
siNeg9	+ - -	+ - -	+ - -
siGiantin#1	- + -	- + -	- + -
siGiantin#2	- - +	- - +	- - +

p-ATM

p-CHK2

TUBA1A

γH2AX

f

p-DNAPK

VLC

g

cell population (%)

☐ siNEG9
☐ siGiantin#1

G1   S   G2/M

FIGURE 6

EXTENDED DATA FIGURE 1

a

	RAD51C	GM130	merge	zoom
U-2 OS				
MCF7				

RAD51C – GM130 - nuclei

b

	RAD51C	GM130	nuclei
siNEG9			
siRAD51C			

c

RAD51C total int. (a.u)

****

siNEG9   siRAD51C

d

siNEG9   siRAD51C

RAD51C

α-tubulin

e

	DMSO		DOX	
	M	N	M	N
RAD51C				
GM130				
Lamin B1				

f

RAD51C distibution ratio

*

DMSO   DOX

EXTENDED DATA FIGURE 2

EXTENDED DATA FIGURE 3

EXTENDED DATA FIGURE 4

a

b

c