The ER tether VAPA is required for proper cell motility and anchors ER-PM contact

sites to focal adhesions

Hugo Siegfried[1], Rémi Le Borgne[1], Catherine Durieu[1], Thaïs De Azevedo Laplace[2], Agathe

Verraes[1], Lucien Daunas[1], Jean-Marc Verbavatz[1*], Mélina L. Heuzé[1*]

[1] Université Paris Cité, CNRS, Institut Jacques Monod, F-75013 Paris, France

[2] Université Paris Cité, INSERM, Centre de Recherche sur l'Inflammation, F-75018 Paris,

France

[*] Co-Corresponding authors

13 ABSTRACT

Cell motility processes highly depend on the membrane distribution of Phosphoinositides

(PInst), giving rise to cytoskeleton reshaping and membrane trafficking events. Membrane

contact sites serve as platforms for lipid exchange and calcium fluxes between two organelles.

Here, we show that VAPA, an ER membrane-resident contact site tether, plays a crucial role

during cell motility. CaCo2 adenocarcinoma epithelial cells depleted for VAPA exhibit several

collective and individual motility defects, disorganized actin cytoskeleton and altered protrusive

activity. During migration, VAPA is required for the maintenance of PI(4,5)P2 levels at the

plasma membrane, but not for PI(4)P homeostasis in the Golgi and endosomal compartments.

Importantly, we show that VAPA regulates the dynamics of focal adhesions (FA) through its

MSP domain, and is essential to stabilize and anchor ventral ER-PM contact sites to FA, thus

mediating microtubule-dependent FA disassembly. To conclude, our results reveal

unprecedented functions for VAPA-mediated membrane contact sites during cell motility and

provides a dynamic picture of ER-PM contact sites connection with FA mediated by VAPA.

INTRODUCTION

The lipid identity of membrane compartments is highly regulated within cells and plays a key role in a diversity of cellular processes. During cell motility, lipids, in particular Phosphoinositides (PInst), are local determinants of molecular events leading to cell polarization, the formation of actin-driven protrusions and turnover of focal adhesions (FA) (Hammond and Burke, 2020; Tsujita and Itoh, 2015). PI(4,5)P2, the most abundant PInst at the plasma membrane, controls FA dynamics by regulating the binding of talin to integrins (Martel et al., 2001; Thapa et al., 2012) and the polarized trafficking of integrins (Nader et al., 2016). PI(4,5)P2 is described as a modulator of actin cytoskeleton organization and dynamics, either directly through the recruitment of several actin-binding proteins (Senju et al., 2017), or indirectly by regulating the activation of RhoA (Lacalle et al., 2007; Xu et al., 2010) and Cdc42 GTPases (Daste et al., 2017). The product of PI(4,5)P2 phosphorylation, PI(3,4,5)P3, also plays a crucial role in cell motility at the plasma membrane (PM). Several i*n vitro* and *in vivo* studies have shown that PI(3,4,5)P3 acts as a "compass lipid" that stabilizes the direction of migration (Funamoto et al., 2002; Hannigan et al., 2002; Lam et al., 2012) through the activation of Rac1 (Kunisaki et al., 2006; Welch et al., 2002; Yoshii et al., 1999).

Lipids are synthesized in the endoplasmic reticulum (ER) and delivered to membranes through vesicular transport, but also through non-vesicular transport at so-named membrane contact sites (MCS) (Jackson et al., 2016). MCS are sites of close apposition between two membranes, often the ER membrane and the membrane of another organelle, at a distance of less than 80nm, where exchanges of lipids, Ca2+ and metabolites take place (Prinz et al., 2020; Scorrano et al., 2019; Wu et al., 2018). While the activity of lipid transfer at MCS was observed decades ago (Vance, 1990), their implication in patho-physiological processes such as cell motility has gained interest only recently (Prinz et al., 2020).

In this work, we studied the role of VAPA, a tethering protein at ER-mediated MCS during cell motility. VAPA is a member of the highly conserved VAP family of proteins, together with VAPB. VAP proteins are integral ER membrane proteins bridging the ER membrane to the

membrane of other compartments by assembling with lipid transfer proteins that bind the target membrane, such as Nir proteins and OSBP-related (ORP) proteins. The assembly of VAP complexes relies on the interaction between the conserved N-terminal MSP (Major Sperm Protein) domain of VAPs and the FFAT (2 phenyl alanines in an acidic tract) motif on their partner. VAPs also contain a central coiled-coil dimerization domain and an ER transmembrane domain in their C-terminus (Murphy and Levine, 2016). VAP proteins are involved in several cellular functions such as lipid transport, membrane trafficking, the unfolded protein response pathway and microtubules organization (Kamemura and Chihara, 2019; Lev et al., 2008). VAP complexes with lipid transfer proteins control the homeostasis of PInst and sterols at various intracellular locations. At ER-PM contact sites, the VAP/Nir2 and VAP/ORP3 complexes regulate the transport of Phosphoinositol (PI) and PI(4)P respectively. Together with membrane-associated kinases and phosphatases, they contribute to modulating locally the pool of PI(4)P, PI(4,5)P2 and PI(3,4,5)P3 at the PM (Chang and Liou, 2015; Gulyás et al., 2020; Kim et al., 2013). In addition, the association of VAPs with ORP proteins modifies the level of PI(4)P at endosomal membranes and at the Golgi, therefore regulating trafficking events in these two compartments (Dong et al., 2016; Kawasaki et al., 2022; Mesmin et al., 2013). Moreover, VAP proteins mediate cholesterol transport at ER-Golgi, ER-endosome and ER-peroxisome MCS (Dong et al., 2019; Hua et al., 2017; Mesmin et al., 2013; Wilhelm et al., 2017). Depending on the cell type and organism, the loss of VAP proteins can lead to either a reduction (Peretti et al., 2008) or an accumulation (Mao et al., 2019) of PI(4)P levels in the Golgi membranes and its redistribution on endosomes (Dong et al., 2016). In yeast, depleting VAP orthologs results in the accumulation of PI(4)P at the plasma membrane (Stefan et al., 2011).

Altogether, these observations prompted us to hypothesize that VAPs, by regulating PInst homeostasis, could contribute to cell motility processes, which heavily rely on regulation of PM PInst. Importantly, two partners of VAPs at ER-PM contact sites, ORP3 and Nir2, have been shown to regulate cell motility processes through different pathways (D'Souza et al., 2020; Keinan et al., 2014; Lehto et al., 2008; Weber-Boyvat et al., 2015). As a first-line approach, we

decided to investigate the effects of VAPA depletion alone on the motility of human adenocarcinoma Caco2 cells. We observed strong motility defects, suggesting that VAPA is required for proper cell motility and that VAPB is not sufficient to compensate for the loss of VAPA. We then aimed to understand the precise function of VAPA in this process. Here, we show that the depletion of VAPA strongly impacts the organization of the actin cytoskeleton, the dynamics of protrusions and the turnover of FA during migration. The role of VAPA stands mainly at the PM where it regulates PI(4,5)P2, the stability of ER-PM contact sites and their local anchoring at FA, thereby regulating FA disassembly.

RESULTS

VAPA is required for proper cell migration and cell spreading

To assess the role of VAPA during cell motility, we generated, using CRISPR/Cas9 gene editing, stable Caco-2 cells either depleted for VAPA (VAPA KO), or expressing non-targeting sequences (Control). VAPA KO cells exhibited a complete and specific depletion of VAPA protein (Fig.1A and 1B) without significantly affecting the levels of VAPB (which on the contrary had a tendency to increase), or its distribution in the ER (Sup Fig.1). We first assessed the capacity of VAPA KO and Control monolayers to migrate collectively upon space release on fibronectin-coated glass. In these conditions, VAPA KO monolayers filled the open space less efficiently than Control monolayers (Fig.1C and 1D, Sup Movie 1). The analysis of leader cells trajectories revealed that VAPA KO leader cells were displacing faster but with a remarkable lack of directionality compared to Control leader cells (Fig.1E and 1F), which could account for the slower colonization capacity VAPA KO monolayers. This collective migration impairment was at least partially cell-intrinsic, as single VAPA KO cells displacing on a 2D surface exhibited identical defects compared to VAPA KO leader cells in monolayers (Fig.1G and 1H). Moreover, upon plating, VAPA KO cells were spreading much faster than Control cells, also indicating a cell-autonomous defect arising in these cells (Fig.1I and 1J). Together, these results identify VAPA as a novel actor of collective and individual cell motility processes.

VAPA regulates focal adhesions and actin cytoskeleton through its MSP domain

In order to characterize the motility defects of VAPA KO cells, we studied the organization of cell-matrix adhesions and actin cytoskeleton in these cells. To facilitate the analysis, we decided to focus on leader cells, standing at the leading edge of migrating monolayers, as these cells move in a given direction with a polarized and reproducible morphology. We observed an alteration of cell-matrix adhesions in VAPA KO cells that formed larger FA compared to Control cells (Fig.2A and 2D). Importantly, the expression of an exogenous wild-type form of VAPA in VAPA KO cells restored the size of FA, but it was not the case with the MSP-mutated form of VAPA (VAPA KDMD), indicating that VAPA regulates FA size through its association with proteins containing a FFAT motif (Fig.2A, 2D and Sup.Fig.2). We then compared the organization of actin cytoskeleton in both Control and VAPA KO cell lines. In Control cells, the F-actin and cortactin stainings revealed the presence of numerous and thick parallel transverse actin arcs at the front, accompanied by cortactin-rich subdomains of branched actin standing at the front edge corresponding to lamellipodial extensions (Fig.2B and 2C). We indeed observed periodic waves of protrusion-retraction arising in these cells every 15 minutes in average (Fig.2C and 2G). By contrast, VAPA KO cells exhibited disorganized and thinner transverse actin arcs, wider cortactin-rich subdomains occupying most of the leading edge and giving rise to long-lasting protrusions (Fig.2B,2C,2E-2G). Similarly to the FA phenotype, we were able to recover the size of cortactin-rich domains by expressing a wild-type form of VAPA but not the MSP mutant VAPA KDMD (Fig.2E). To conclude, we show that VAPA controls different aspects of cell motility, namely the maintenance of FA size and the preservation of the spatial and temporal organization of the actin cytoskeleton. These functions depend on the association of VAPA with FFAT-motif containing proteins such as lipid transfer proteins, suggesting that lipid transfer is probably required.

VAPA controls PI(4,5)P2 levels at the PM, but is not essential for PI(4)P homeostasis in

the Golgi and endosomal compartments

We then intended to decipher the mechanisms by which VAPA could regulate FA size, the

actin cytoskeleton and cell motility. To test the lipid transfer hypothesis, we first characterized

to which extend its deletion affected PInst homeostasis. Previous studies have shown that

depleting both VAPA and VAPB affected the levels of PI(4)P at the Golgi and PM (Mao et al.,

2019; Stefan et al., 2011), and induced the accumulation of PI(4)P in early endosomes (Dong

et al., 2016). Using a GFP-PH-OSBP probe, we detected a major intra-cellular pool of PI(4)P

at similar levels in the Golgi compartment in Control and VAPA KO cells (Fig.3A-C).

Importantly, the depletion of VAPA alone was not sufficient to observe an accumulation of

PI(4)P in early endosomes (Fig.3A-C). This result suggests that VAPB can compensate for the

absence of VAPA at ER-Golgi and ER-endosome contact sites thus maintaining PI(4)P

homeostasis at the Golgi and endosomes in VAPA KO Caco-2 cells. However, at the PM, the

depletion of VAPA induced a decrease of PI(4,5)P2 which is the most abundant PInst in this

membrane compartment (Fig.3D-3G). This decrease was detected at the dorsal side of cells

by immunofluorescence staining (Fig.3D and 3F), and at the protrusive cortactin-rich leading

edge of migrating cells by measuring the PM/cytoplasm ratio of RFP-PH-PLCδ1 probe for

PI(4,5)P2 (Fig.3E and G). Altogether, these results show that in Caco-2 cells, VAPA is required

to maintain a certain level of PI(4,5)P2 specifically at the plasma membrane, but is dispensable

for PI(4)P homeostasis in the Golgi and endosomal compartments.

VAPA stabilizes ventral ER-PM contact sites at the front of migrating cells

Based on our observation pointing to a specific role of VAPA for the PI(4,5)P2 homeostasis at

the PM, we questioned the spatio-temporal distribution of ER-PM contacts sites during cell

migration and the role of VAPA in this organization. To this aim, we took advantage of a

fluorescent probe, GFP-MAPPER, which selectively labels ER-PM contact sites (Chang et al.,

2013). When expressed in Caco-2 cells, GFP-MAPPER distributed as puncta along ER tubules

(Fig.4A) and appeared at ER-PM appositions detected by TIRF microscopy at the cell bottom

(Fig.4B and Sup Movie 2). At the front of leader cells, GFP-MAPPER puncta divided in two

subpools: the dorsal subpool ongoing forward movement with the migrating cell and the ventral

subpool remaining immobile relative to the substrate, suggesting that ventral ER-PM contact

sites might be anchored to cell-matrix adhesions (Fig.4C). Using electron microscopy, we were

indeed able to identify numerous ER-PM contact sites sitting at the ventral leading edge of

migrating cells (Fig.4D). At the ultrastructural level, the depletion of VAPA had no significant

effect on ER-PM contact sites regardless of their location (Sup Fig.3). However, the dynamic

analysis of GFP-MAPPER puncta revealed that ventral ER-PM contact sites, which barely

moved (400nm/min on average) and persisted generally more than 10 minutes in Control cells,

were significantly less stable both spatially and temporally in VAPA KO cells (Fig.4E-G and

Sup Movie 3). To conclude, our results identify a sub-pool of ER-PM contact sites docked to

the substrate at the front of migrating cells, and whose spatio-temporal stability requires VAPA.

Together with the fact that FA size is altered in VAPA KO cells, these results prompted us to

hypothesize that VAPA might accomplish a function at FA.

VAPA promotes microtubule-dependent FA disassembly

To further investigate the role of VAPA in FA turnover, we studied more precisely FA dynamics

in Control and VAPA KO cells during cell migration by TIRF microscopy. While the assembly

rate of FA was similar in both cell lines, their disassembly rate was slower in VAPA KO cells

(Fig.5A, 5B and 5E). In addition, VAPA KO FA exhibited a longer lifetime in average (Fig.5C

and 5E), with a majority lasting at least 45 minutes (Fig.5D and E). To test whether VAPA-

mediated disassembly of FA relied on the microtubule network, we synchronized FA

disassembly by nocodazole washout experiment in migrating cells and measured the evolution

of FA size, as described previously (Ezratty et al., 2005). Our results clearly show that VAPA

KO FA failed to disassemble upon microtubule recovery, contrary to Control FA that lost half

of their size within 15 minutes after nocodazole washout (Fig.5F and 5G). These results

demonstrate that VAPA is required for microtubule-dependent FA disassembly during

migration of Caco2 cells.

VAPA mediates the anchoring of ER-PM contact sites to FA

To understand the function of VAPA at FA, we analysed the repartition and the dynamic of ER-PM contact sites relative to FA by TIRF microscopy. In Control cells, the majority of vinculin-stained FA (80%) established a contact with ventral ER-PM contact sites whereas it was the case for only 30-40% of FA in VAPA KO cells (Fig.6A and 6B). We then monitored the time course of mCherry-Vinculin and GFP-MAPPER signals within the perimeter of single FA (Fig.6C-D and Sup Movie 4). GFP-MAPPER foci started to accumulate in the FA vicinity after its assembly, suggesting that ER-PM contact sites were anchoring to preexisting FA. Importantly, the disassembly of FA started immediately upon GFP-MAPPER recruitment and evolved linearly in parallel to GFP-MAPPER accumulation (Fig.6E and Sup Movie 4). Thus, we show that VAPA is required for the anchoring of ER-PM contact sites to FA, which happens at the time of FA disassembly.

DISCUSSION

In this work, we identify VAPA as a crucial player in cell motility, with implications in different aspects of this process. We show that VAPA is required for both single and collective cell migration, cell spreading, protrusive waves, actin cytoskeleton dynamics and FA turnover. Our study reveals a non-redundant function for VAPA in cell motility among the VAP family of proteins in Caco2 cells. Indeed, the presence of VAPB in VAPA KO cells was not sufficient to compensate the loss of VAPA. The function of VAPA in cell migration seems to arise from its specific role at ER-PM contact sites where it was necessary to maintain high levels of PI(4,5)P2 independently of VAPB. It was not the case at the Golgi or endosomes, where the depletion of VAPA alone had no effect on PI(4)P levels, which is in agreement with previous observations by Dong et al in VAPA/VAPB double knock-out cells re-expressing VAPB (Dong et al., 2016). This suggests that VAPB might compensate for the absence of VAPA at ER-Golgi and ER-endosomes contact sites and that migration defects in VAPA KO cells were not a result of a dysfunction in the secretory pathway. Together with previous studies (Dong et al., 2016; Mao et al., 2019; Peretti et al., 2008; Stefan et al., 2011), our results converge to the idea that VAPA

and VAPB might fulfill both redundant and specific tasks in lipid homeostasis and cellular

functions.

PI(4,5)P2 and its phosphorylation product PI(3,4,5)P3 have been described for decades as

central orchestrators of cell motility through regulation of actin dynamics, FA maturation and

turnover, membrane organization and curvature, and intracellular trafficking (Mandal, 2020;

Tsujita and Itoh, 2015). Thus, the phenotypes that we observed in VAPA KO cells – namely

disorganization of actin, protrusion defects, FA persistence, loss of directional movement and

faster speed– could be solely explained by a global reduction in PI(4,5)P2 levels at the PM.

However, our results clearly show that VAPA not only regulates PI(4,5)P2 homeostasis at the

PM, but also fulfills a function at FA where it anchors ER-PM contact sites and favours their

stability, thereby promoting microtubule-dependent FA disassembly. Importantly, we provide

for the first time a quantitative spatio-temporal connection between ER-PM contact sites and

FA dynamics in polarized and migrating cells. Based on these results, we propose a

hypothetical model in which VAPA, through local regulation of lipid transfer in the vicinity of FA

would induce the internalization of integrins through clathrin-mediated endocytosis which

depends on PI(4,5)P2 levels (Chao et al., 2010; Nader et al., 2016) and on microtubule

polymerization (Ezratty et al., 2009, 2005) (Fig.7). Interestingly, MCS have already been

shown to directly regulate endocytosis in yeast (Encinar del Dedo et al., 2017).

Which proteins could control the anchoring of VAPA-mediated ER-PM contact sites to FA?

Previous studies have highlighted the role of ER proteins in FA turnover and maturation that

could be potential candidates. Kinectin, an integral ER membrane protein interacting with

kinesin, has been shown to control the apposition of ER to FA through a microtubule-

dependent transport (Guadagno et al., 2020; Ng et al., 2016). The phosphatase PTP1B, which

localizes to the cytoplasmic face of the ER and was identified as a partner of VAPA using a

BioID approach (Antonicka et al., 2020), is targeted to newly formed FA and contributes to

their maturation through dephosphorylation of its substrate p130Cas (Dadke and Chernoff,

2002; Hernández et al., 2006; Liu et al., 1998). In addition, we demonstrate that the function

of VAPA at FA depends on its MSP domain of interaction with proteins containing a FFAT

domain. Importantly, two lipid transfer proteins interacting with VAPA at ER-PM contact sites,

Nir2 and ORP3, have been previously described as regulators of cell motility, establishing them

as potential drivers for the specific function of VAPA at FA. Nir2, which interacts with VAPs at

ER-PM and ER-Golgi contact sites and regulates PI transport (Chang and Liou, 2015; Kim et

al., 2013; Peretti et al., 2008), favors epithelial-mesenchymal transition and tumor metastasis

(Keinan et al., 2014). Similarly to VAPA KO cells, the depletion of Nir2 in HeLa cells induces a

decrease of PI(4,5)P2 levels at the PM (Kim et al., 2013), indicating that a part of VAPA effect

on PInst homeostasis could be supported by VAPA/Nir2-mediated lipid transfer. ORP3, which

extracts PI(4)P from the PM at ER-PM contact sites (Gulyás et al., 2020), positively regulates

the R-Ras pathway promoting cell-matrix adhesion (Lehto et al., 2008; Weber-Boyvat et al.,

2015). More recently, ORP3 was found to be recruited to disassembling FA upon calcium influx

and to favour their turnover through lipid exchange. Interestingly, the authors highlight a

crosstalk between ORP3-mediated lipid exchange and STIM1/Orai1-mediated calcium influx

at FA (D'Souza et al., 2020). Therefore, the role of VAPA-mediated contact sites at FA could

not only be the local modulation of lipid transfer, but also the establishment of a common

platform with STIM1, anchored to FA at ER-PM contact sites controlling lipid and calcium

exchanges and driving FA maturation and turnover.

To conclude, our study reveals unprecedented functions of VAPA in cell motility processes and

in local anchoring of ER-PM contact sites to FA. For a long time, we have assumed that PInst

regulation at FA will principally be mediated by the interplay of lipid kinases and phosphatases.

Our work reinforces the idea that lipid transfer would be an additional pathway for lipids to be

controlled at FA and provides evidences for a spatio-temporal connection between MCS and

FA mediated by VAPA. Finally, this work opens new roads to decipher the precise molecular

determinants of VAPA function and to explore the role of VAPA during cancerogenesis.

MATERIALS AND METHODS

Cell culture

CaCo-2 cells, originally acquired from ATCC, were kindly provided by Dr. D. Delacour (Institut Jacques Monod, Paris). CaCo-2 CRISPR control and VAPA knock-out cell lines were maintained in culture in DMEM (1x) GlutMax 5g/L D-glucose + pyruvate medium (Gibco) supplemented with 20% fetal bovine serum (Gibco), 100 units/mL penicillin, 100 µg/mL streptomycin and 2 µg/Ml (Gibco) puromycin in 5% CO_2 at 37°C.

Antibodies, reagents and plasmids

The following primary antibodies were used: mouse anti-EEA1 monoclonal antibody (BD transduction, 610456), rabbit anti-VAPB polyclonal antibody (Atlas antibody HPA 013144), mouse anti-VAPA monoclonal antibody (Sigma 10355-4C12) and rabbit anti-VAPA polyclonal antibody for the immunoblot (Sigma SAB2702059), rabbit anti-TGN46 polyclonal antibody (abcam ab50595), mouse anti-paxilline monoclonal antibody (Merck, 05-417), mouse anti-cortactin monoclonal antibody (Merck, 05-180), mouse anti-tubulin monoclonal antibody (GeneTex GTX628802), mouse anti-PI(4,5)P2 monoclonal IgM antibody (Echelon Z-P045). The following secondary antibodies were used: goat anti-mouse IgM Alexa fluor 555 (Invitrogen, A21426) or Alexa fluor 488 (Invitrogen, A21042); goat anti-mouse IgG Alexa fluor 555 (Invitrogen, A21424) or Alexa fluor 488 (Invitrogen, A31620); goat anti-rabbit IgG Alexa fluor 555 (Invitrogen, A21429) or Alexa fluor 488 (Invitrogen, A11033).

The following reagents were used: fibronectin Hu Plasma Fibronectine EMD Milipore Corp; Methanol-free formaldehyde (Thermofisher, 28908); Fluoromont-G with DAPI (Invitrogen, 00-4959-52); Alexa fluor 647-conjugated Phalloidin (Invitrogen, A22287).

The following plasmids were used; pEGFP-MAPPER was a kind gift from Jen Liou (UT Southwestern Medical Center, USA) (Chang et al., 2013), pmCherry-Vinculin was a gift from Chinten James Lim (Addgene plasmid # 80024 ; http://n2t.net/addgene:80024 ; RRID:Addgene_80024) (Lee et al., 2013); pEGFP-PH-OSBP was a gift from Marci Scidmore

(Addgene plasmid # 49571 ; http://n2t.net/addgene:49571 ; RRID:Addgene_49571)

(Moorhead et al., 2010); pmRFP-PH-PLCδ1 was a kind gift from Francesca Giordano (Institute

for Integrative Biology of the Cell, France) (Giordano et al., 2013); pmRFP-KDEL plasmid was

a kind gift from Nihal Altan-Bonnet (National Heart Lung and Blood Institute, NIH, USA) (Altan-

Bonnet et al., 2006); pmCherry-VAPA and pmCherry-VAPA KD/MD, obtained from pEGFP-

VAPA and pEGFP-VAPA KD/MD, were kind gifts from Fabien Alpy (IGBMC, France) (Alpy et

al., 2013); CRISPR-Cas9-resistant pmCherry-VAPA and pmCherry-VAPA KD/MD plasmids

were obtained by mutagenesis of pmCherry-VAPA and pmCherry-VAPA KD/MD plasmids.

Briefly, pmCherry-VAPA and pmCherry-VAPA KD/MD plasmids were amplified by PCR (30

cycles of 10s at 94°C, 5s at 55°C and 6min at 72°C) using the forward

5'TAAGACCGAATTCCGGTATCATCGATCCAGGGTCAACTGTGACTGTTTCAGT 3' and

reverse 5' CGGAATTCGGTCTTACGCAATATCGTCGAGGTGCTGTAGTCTTCACTTTGA 3'

primers. PCR products were digested by DpnI enzyme for 2h at 37°C (Agilent) and loaded on

a 1% Agarose Gel. PCR amplicons were extracted, purified using the NucleoSpin Gel and

PCR clean up kit (Macherey-Nagel) and ligated using the NEBuilder HiFi DNA assembly kit

(BioLabs).

Transfection

For each transfection, 0.5 million Caco2 cells were nucleofected with 2 µg of DNA in 100uL of

T solution using the B024 program on Amaxa Nuclefector II machine, as recommended by the

manufacturer (Lonza). CaCo-2 transfected cells were then re-suspended in warm culture

medium, replaced 24 hours later.

Production of CRISPR Cas9 cell lines

Cells were transfected as described above with the double nickase CRISPR Cas9 plasmid

(Santa Cruz, sc-424440-NIC) or a CRISPR Control plasmid (Santa Cruz, sc-437281). The

following day, cells were incubated with puromycin at 2 µg/mL. After antibiotic selection, single

GFP positive cells were sorted by flow cytometry. Clonal cells were maintained in culture with

puromycin and the amount of VAPA protein was detected by western blot and

immunofluorescence. VAPA KO cell lines expressing exogenous Cherry-VAPA and Cherry-

VAPAKDMD were obtained by transfection of VAPA KO cells with CRISPR-Cas9-resistant

pmCherry-VAPA and pmCherry-VAPA KD/MD plasmids. Stable cell lines were generated by

selection with Geneticin (Gibco) and sorting of Cherry-positive populations by flow cytometry.

Western Blotting

Confluent cells were lysed in 100 mM Tris pH 7.5,150 mM NaCl, 0.5% NP40, 0.5% triton-X100,

10% glycerol,1X protease inhibitor cocktail (Roche) and 1X phosphatase inhibitor

(Phosphostop, Roche) for 20 minutes at 4°C. After 15 minutes centrifugation at 13000 g,

solubilized proteins were recovered in the supernatant. Protein concentration was measured

using Bradford assay (BioRad). For SDS PAGE, 50ug protein extracts were loaded in 4-12%

Bis-Tris gel (Novex) or poly-acrylamide gels and proteins were transferred overnight at 4°C on

a nitrocellulose membrane using a liquid transfer system (BioRad). Non-specific sites were

blocked with 5% non-fat dry milk in PBS 0.1% Tween 20. Primary antibodies were diluted

(1/1000 to 1/500) in PBS 0.1% Tween 20 and incubated overnight at 4ºC. After three washes

in PBS 0.1% Tween 20, secondary HRP antibodies diluted in PBS 0.1% Tween 20 (1/10000)

were incubated for 1 hour and washed 3 times with PBS 0.1% Tween 20. Immunocomplexes

of interest were detected using Supersignal west femto maximum sensitivity substrate

(ThermoFisher) and visualized with ChemiDoc chemoluminescence detection system

(Biorad). Quantification of Western blots by densitometry was performed using the Gel

analyzer plug in from Image J.

Immunofluorescence

Cells were fixed with pre-warmed 4% formaldehyde in PBS for 15 min at RT and then washed

3 times with PBS. Permeabilization and blocking were performed in 0.05% saponin/0.2% BSA

in PBS for 15 minutes at RT. The primary antibodies diluted (1/100) in Saponin/BSA buffer

were then incubated overnight at 4°C. After 3 washes in saponin/BSA buffer, the samples were

incubated with secondary antibodies (1/2000) and, when indicated, Alexa-coupled phalloidin (1/200) to stain F-Actin in the same buffer for 1 hour at RT. The preparations were washed twice in saponin/BSA buffer, once in PBS, and then mounted with the DAPI Fluoromount-G mounting media.

Immunostaining with anti-PI(4,5)P2 antibody was performed as described by Kim et al. (Kim et al., 2013). Briefly, cells were fixed at 4°C for 1h in 3.7% formaldehyde / 0.1% glutaraldehyde and incubated in PBS containing 0.1M glycine for 15 min. Permeabilization, blocking and staining were performed as described above, at 4°C.

Cell migration assays

Collective migration assays were performed using 2-well silicon inserts (Ibidi). Glass coverslips were coated with fibronectin solution at 20 µg/mL in water, for 1h at room temperature. The cover slip surface was washed with sterile water and air-dried. Ibidi inserts were deposited on the fibronectin coated surface and 40000 to 50000 cells were loaded per well. For fluorescence live imaging, cells were directly plated after transfection in the well. After 3 to 4 hours of incubation, cell division was blocked using mitomycin at 10 µg/mL in CaCo-2 culture medium, for 1 hour. After overnight incubation in a fresh CaCo-2 culture medium, the insert was removed. Experiments were performed 24h to 48h after insert removal.

For individual cell migration, cell division was blocked using mitomycin at 10 µg/mL in CaCo-2 culture medium, for 1 hour. After overnight incubation in a fresh CaCo-2 culture medium, cells were detached and plated on a glass coverslip coated with 20 µg/mL fibronectin. After 6 hours, cells were imaged for 24h.

Collective and individual cell migration assays were performed using a Zeiss Wide-Field Microscope and imaged at 1 frame every 10 minutes. Cell trajectories were analysed using Manual Tracking from Fiji.

Cell spreading assay

Cells were incubated for 1h on fibronectin-coated coverslips and fixed with formaldehyde 4%

solution as described in the immuno labelling section. Coverslips were mounted on slides and

samples were imaged using a Zeiss Apotome fluorescence microscope equipped with a 10x

objective. The spreading area was determined based on Phalloidin-segmented ROI in

Fiji/ImageJ.

Nocodazole experiment

The nocodazole wash-out experiment was adapted from Ezratty et al (Ezratty et al., 2005) .

Briefly, cells were plated in the insert in starvation medium (DMEM GlutaMax medium

containing 1% fetal bovine serum and 1% of penicillin/streptomycin). 24h later, the insert was

removed, cells were left migrating for 24h in the starvation medium and treated with 10 µM

nocodazole in starvation medium for 2h. Then, nocodazole was washed-out and cells were

incubated with fresh starvation medium for 0, 15, 30 and 60 min and then fixed using 4%

formaldehyde. Focal adhesions were labelled using a mouse anti-paxillin antibody as

described above.

Electron microscopy

Migrating cells were fixed in a 2% formaldehyde/1% glutaraldehyde in PBS solution for 1h at

room temperature, then washed in PBS. Cells were embedded in a 10% BSA and 10% gelatin

gel. Post fixation was performed in 1% OsO_4+1% $K_3Fe(CN)_6$ in water at 4°C for 1h. After

extensive washes, cells were incubated in 1% Thiocarbohydrazine in water for 20 min at room

temperature and then washed and incubated with 2% OsO_4 in water for 30 min at room

temperature. After washes, cells were contrasted with 1% uranyl acetate overnight at 4°C. The

following day, the uranyl acetate solution was removed and the samples washed using pure

water. Cells were incubated with a pH 5.5 lead nitrate in aspartic acid solution for 30 min at

room temperature, washed in water and dehydrated in successive ethanol baths. Cells were

embedded in Agar Low Viscosity Resin (R1018 Kit from agar Scientific). 70 nm-thick thin

416 sections were cut using a UC6 ultramicrotome from Leica and deposited on EM grids. Electron

417 microscopy acquisitions were performed using a 120kV Tecnai 12 electron microscope

418 (ThermoFisher) equiped with a OneView 4K camera (Gatan).

Analysis of protrusive activity

421 The analysis of protrusive activity was performed on DIC images acquired every 3 seconds for

422 30 minutes. Kymographs representing the evolution of the leading edge in time along a line

423 were generated and analysed with Fiji. Positive slopes were considered as protrusions and

424 negative slopes as retractions. For each cell, the mean time spent in protrusion and the mean

425 frequency of protrusions.was calculated from 2 to 4 kymographs.

Measurement of focal adhesion assembly and disassembly

Cells were transfected as described above using the mCherry-Vinculine plasmid and migration

assays were performed on fibronectin coated 1.5H Glass bottom Dishes (Ibidi). Cells were

imaged using a TIRF microscope at 1 frame every minute for 1 to 2 hours. Focal adhesion

assembly and disassembly rates were obtained using the focal adhesion analysis server

(https://faas.bme.unc.edu/) (Berginski and Gomez, 2013). Focal adhesion assembly and

disassembly rate tracks were generated. The tracks with a R-squared equal to or greater than

0.8 were used for the analysis.

Quantification of FA in contact with ER-PM contact sites

437 The quantification of FA in contact with ER-PM contact sites was performed on TIRF

438 microscopy acquisitions. 10 to 30 FA were randomly selected per cell. An FA was considered

439 to be in contact with an ER-PM contact sites if there was at least one pixel recovery between

440 GFP-MAPPER signal and mCherry-vinculin signal.

Quantification of PI(4)P and PI(4,5)P2 levels

To characterize the intracellular pool of PI(4)P at the Golgi and on endosomes, colocalization between the GFP-PH-OSBP and TGN46 or EEA1 signals were obtained using Fiji Pearson coefficient plugins. The relative levels of PI(4)P in the Golgi or in endosomes were calculated as the ratio between GFP-PH-OSBP signal in the Golgi or endosomes masks and GFP-PH-OSBP signal in the cytosol, using segmentation tools in Fiji.

Concerning PI(4,5)P2, PM to cytosol signal ratio of RFP-PH-PLCδ1 was measured along lines of 5 to 15 microns at the leading edge of migrating cells, in protrusive area. The ratio was calculated as the maximal peak intensity along the line divided by the mean intensity value of RFP-PH-PLCδ1 in the cytosol. The plots on the graph represent the mean of 3 ratio at 3 different locations per cell.

For the quantification of endogenous PI(4,5)P2, cells were immnuno-stained with anti-PI(4,5)P2 antibodies. Confocal imaging was performed using the same acquisition parameters between control and VAPA KO cells. For one z-section, 3 lines were drawn per cell and the mean of maximal fluorescence intensity at the peak was plotted.

Analysis of ER-PM contact sites on electron microscopy images

Electron microscopy images were analysed using Fiji/ImageJ. For the analysis, only ER-PM appositions of less than 30 nm distance were considered as ER-PM contact sites.

Images acquisition and analysis

For live-microscopy experiments, the samples were placed in a chamber equilibrated at 37°C under 5% CO2 atmosphere.

Fluorescence live microscopy imaging was acquired using a Yokogawa CSU-X1 Spinning Disk confocal mounted on an inverted motorized Axio Observer Z1 (Zeiss) and equipped with a sCMOS PRIME 95 (photometrics) camera and a 63 X oil immersion objective or using an

ELYRA TIRF microscope from Zeiss equipped with EMCCD, iXon 897 (512*512, 16 microns of pixel size) camera and 63 X oil immersion objective.

For fixed samples, images were acquired with a Zeiss Apotome fluorescence microscope equipped with a 63 X oil immersion objective or with a Zeiss LSM 780 confocal microscope equipped with a 63 X oil immersion objective at a resolution of 0.3 µm z-stacks. Image processing and analysis were done on Fiji software.

AKNOWLEDGEMENTS

We thank Delphine Delacour, Fabien Alpy, Simon de Beco and the past and current members of Membrane Dynamics and Intracellular Trafficking team at Institut Jacques Monod for helpful discussions. We acknowledge the ImagoSeine core facility of the IJM, member of the France BioImaging infrastructure (ANR-10-INBS-04) and GIS-IBiSA, with support from La ligue contre le Cancer (R03/75-79), the Region Île-de-France (Sesame), Université Paris Cité (Labex Who Am I), Inserm (Plan Cancer), Région Ile de France (SESAME) and Fondation Bettencourt Schueller, France. This work was supported by grants from Gefluc groupement Les Entreprises Contre le Cancer, Cancéropôle Ile de France and INCa (2021-1-EMERG-51), La Ligue contre le cancer (RS21/75-85), the Agence Nationale de la Recherche (ANR-20-CE13-0021) and the Fondation pour la Recherche Médicale (DEQ. 20150934717), France. H.S was supported by fellowships from La Ligue contre le cancer and the Fondation pour la Recherche Médicale. M.L.H acknowledges René-Marc Mège, Benoît Ladoux and Delphine Delacour for kindly providing antibodies and plasmids during the starting period of this work.

Author contributions

M.L.H. conceived the project. M.L.H and J-M.V supervised the project. H.S, R.LB, C.D, T.DA, A.V. and M.L.H performed experiments. M.L.H and H.S designed experiments. H.S, T.DA, A.V and M.L.H analysed data. M.L.H, H.S and J-M.V wrote the manuscript.

Competing Interests statement

The authors declare no competing interests.

REFERENCES

Alpy F, Rousseau A, Schwab Y, Legueux F, Stoll I, Wendling C, Spiegelhalter C, Kessler P, Mathelin C, Rio MC, Levine TP, Tomasetto C. 2013. STARD3 or STARD3NL and VAP form a novel molecular tether between late endosomes and the ER. *J Cell Sci* **126**:5500–5512. doi:10.1242/JCS.139295

Altan-Bonnet N, Sougrat R, Liu W, Snapp EL, Ward T, Lippincott-Schwartz J. 2006. Golgi inheritance in mammalian cells is mediated through endoplasmic reticulum export activities. *Mol Biol Cell* **17**:990–1005. doi:10.1091/MBC.E05-02-0155/ASSET/IMAGES/LARGE/ZMK0020675010009.JPEG

Antonicka H, Lin ZY, Janer A, Aaltonen MJ, Weraarpachai W, Gingras AC, Shoubridge EA. 2020. A High-Density Human Mitochondrial Proximity Interaction Network. *Cell Metab* **32**:479-497.e9. doi:10.1016/J.CMET.2020.07.017

Berginski ME, Gomez SM. 2013. The Focal Adhesion Analysis Server: a web tool for analyzing focal adhesion dynamics. *F1000Research* **2**:68. doi:10.12688/F1000RESEARCH.2-68.V1

Chang CL, Hsieh TS, Yang TT, Rothberg KG, Azizoglu DB, Volk E, Liao JC, Liou J. 2013. Feedback regulation of receptor-induced ca2+ signaling mediated by e-syt1 and nir2 at endoplasmic reticulum-plasma membrane junctions. *Cell Rep* **5**:813–825. doi:10.1016/j.celrep.2013.09.038

Chang CL, Liou J. 2015. Phosphatidylinositol 4, 5-bisphosphate homeostasis regulated by Nir2 and Nir3 proteins at endoplasmic reticulum-plasma membrane junctions. *J Biol Chem* **290**:14289–14301. doi:10.1074/jbc.M114.621375

Chao W-T, Ashcroft F, Daquinag AC, Vadakkan T, Wei Z, Zhang P, Dickinson ME, Kunz J.

2010. Type I phosphatidylinositol phosphate kinase beta regulates focal adhesion

disassembly by promoting beta1 integrin endocytosis. *Mol Cell Biol* **30**:4463–4479.

doi:10.1128/MCB.01207-09

D'Souza RS, Lim JY, Turgut A, Servage K, Zhang J, Orth K, Sosale NG, Lazzara MJ,

Allegood J, Casanova JE. 2020. Calcium-stimulated disassembly of focal adhesions

mediated by an ORP3/IQSec1 complex. *Elife* **9**. doi:10.7554/eLife.54113

Dadke S, Chernoff J. 2002. Interaction of protein tyrosine phosphatase (PTP) 1B with its

substrates is influenced by two distinct binding domains. *Biochem J* **364**:377–383.

doi:10.1042/BJ20011372

Daste F, Walrant A, Holst MR, Gadsby JR, Mason J, Lee JE, Brook D, Mettlen M, Larsson E,

Lee SF, Lundmark R, Gallop JL. 2017. Control of actin polymerization via the

coincidence of phosphoinositides and high membrane curvature. *J Cell Biol* **216**:3745–

3765. doi:10.1083/jcb.201704061

Dong J, Du X, Wang H, Wang J, Lu C, Chen X, Zhu Z, Luo Z, Yu L, Brown AJ, Yang H, Wu

JW. 2019. Allosteric enhancement of ORP1-mediated cholesterol transport by PI(4,5)P

2 /PI(3,4)P 2. *Nat Commun* **10**. doi:10.1038/s41467-019-08791-0

Dong R, Saheki Y, Swarup S, Lucast L, Harper JW, De Camilli P. 2016. Endosome-ER

Contacts Control Actin Nucleation and Retromer Function through VAP-Dependent

Regulation of PI4P. *Cell* **166**:408–423. doi:10.1016/j.cell.2016.06.037

Encinar del Dedo J, Idrissi FZ, Fernandez-Golbano IM, Garcia P, Rebollo E, Krzyzanowski

MK, Grötsch H, Geli MI. 2017. ORP-Mediated ER Contact with Endocytic Sites

Facilitates Actin Polymerization. *Dev Cell* **43**:588-602.e6.

doi:10.1016/J.DEVCEL.2017.10.031

Ezratty EJ, Bertaux C, Marcantonio EE, Gundersen GG. 2009. Clathrin mediates integrin

endocytosis for focal adhesion disassembly in migrating cells. *J Cell Biol* **187**:733–747.

doi:10.1083/JCB.200904054

Ezratty EJ, Partridge MA, Gundersen GG. 2005. Microtubule-induced focal adhesion

disassembly is mediated by dynamin and focal adhesion kinase. *Nat Cell Biol 2005 76*

7:581–590. doi:10.1038/NCB1262

Funamoto S, Meili R, Lee S, Parry L, Firtel RA. 2002. Spatial and temporal regulation of 3-

phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell* **109**:611–23.

Giordano F, Saheki Y, Idevall-Hagren O, Colombo SF, Pirruccello M, Milosevic I, Gracheva

EO, Bagriantsev SN, Borgese N, De Camilli P. 2013. PI(4,5)P2-Dependent and Ca2+-

Regulated ER-PM Interactions Mediated by the Extended Synaptotagmins. *Cell*

153:1494–1509. doi:10.1016/J.CELL.2013.05.026

Guadagno NA, Margiotta A, Bjørnestad SA, Haugen LH, Kjos I, Xu X, Hu X, Bakke O,

Margadant F, Progida C. 2020. Rab18 regulates focal adhesion dynamics by interacting

with kinectin-1 at the endoplasmic reticulum. *J Cell Biol* **219**. doi:10.1083/jcb.201809020

Gulyás G, Sohn M, Kim YJ, Várnai P, Balla T. 2020. ORP3 phosphorylation regulates

phosphatidylinositol 4-phosphate and Ca 2+ dynamics at PM-ER contact sites . *J Cell*

Sci jcs.237388. doi:10.1242/jcs.237388

Hammond GRV, Burke JE. 2020. Novel roles of phosphoinositides in signaling, lipid

transport, and disease. *Curr Opin Cell Biol* **63**:57–67. doi:10.1016/j.ceb.2019.12.007

Hannigan M, Zhan L, Li Z, Ai Y, Wu D, Huang CK. 2002. Neutrophils lacking

phosphoinositide 3-kinase γ show loss of directionality during N-formyl-Met-Leu-Phe-

induced chemotaxis. *Proc Natl Acad Sci U S A* **99**:3603–3608.

doi:10.1073/pnas.052010699

Hernández M V., Davies Sala MG, Balsamo J, Lilien J, Arregui CO. 2006. ER-bound PTP1B

is targeted to newly forming cell-matrix adhesions. *J Cell Sci* **119**:1233–1243.

doi:10.1242/JCS.02846

Hua R, Cheng D, Coyaud É, Freeman S, Di Pietro E, Wang Y, Vissa A, Yip CM, Fairn GD,

Braverman N, Brumell JH, Trimble WS, Raught B, Kim PK. 2017. VAPs and ACBD5

tether peroxisomes to the ER for peroxisome maintenance and lipid homeostasis. *J Cell*

Biol **216**:367–377. doi:10.1083/JCB.201608128

Jackson CL, Walch L, Verbavatz JM. 2016. Lipids and Their Trafficking: An Integral Part of

Cellular Organization. *Dev Cell* **39**:139–153. doi:10.1016/j.devcel.2016.09.030

Kamemura K, Chihara T. 2019. Multiple functions of the ER-resident VAP and its

extracellular role in neural development and disease. *J Biochem* **165**:391–400.

doi:10.1093/jb/mvz011

Kawasaki A, Sakai A, Nakanishi H, Hasegawa J, Taguchi T, Sasaki J, Arai H, Sasaki T,

Igarashi M, Nakatsu F. 2022. PI4P/PS countertransport by ORP10 at ER–endosome

membrane contact sites regulates endosome fission. *J Cell Biol* **221**.

doi:10.1083/jcb.202103141

Keinan O, Kedan A, Gavert N, Selitrennik M, Kim S, Karn T, Becker S, Lev S. 2014. The

lipid-transfer protein Nir2 enhances epithelial-mesenchymal transition and facilitates

breast cancer metastasis. *J Cell Sci* **127**:4740–4749. doi:10.1242/jcs.155721

Kim S, Kedan A, Marom M, Gavert N, Keinan O, Selitrennik M, Laufman O, Lev S. 2013. The

phosphatidylinositol-transfer protein Nir2 binds phosphatidic acid and positively

regulates phosphoinositide signalling. *EMBO Rep* **14**:891–899.

doi:10.1038/embor.2013.113

Kunisaki Y, Nishikimi A, Tanaka Y, Takii R, Noda M, Inayoshi A, Watanabe K, Sanematsu F,

Sasazuki T, Sasaki T, Fukui Y. 2006. DOCK2 is a Rac activator that regulates motility

and polarity during neutrophil chemotaxis. *J Cell Biol* **174**:647–52.

doi:10.1083/jcb.200602142

Lacalle RA, Peregil RM, Albar JP, Merino E, Martínez-A C, Mérida I, Mañes S. 2007. Type I

phosphatidylinositol 4-phosphate 5-kinase controls neutrophil polarity and directional

movement. *J Cell Biol* **179**:1539–53. doi:10.1083/jcb.200705044

Lam P ying, Yoo SK, Green JM, Huttenlocher A. 2012. The SH2-domain-containing inositol

5-phosphatase (SHIP) limits the motility of neutrophils and their recruitment to wounds

in zebrafish. *J Cell Sci* **125**:4973–4978. doi:10.1242/jcs.106625

Lee HS, Anekal P, Lim CJ, Liu CC, Ginsberg MH. 2013. Two modes of integrin activation

form a binary molecular switch in adhesion maturation. *Mol Biol Cell* **24**:1354–1362.

doi:10.1091/MBC.E12-09-0695

Lehto M, Mäyränpää MI, Pellinen T, Ihalmo P, Lehtonen S, Kovanen PT, Groop PH, Ivaska

J, Olkkonen VM. 2008. The R-Ras interaction partner ORP3 regulates cell adhesion. *J*

Cell Sci **121**:695–705. doi:10.1242/jcs.016964

Lev S, Halevy D Ben, Peretti D, Dahan N. 2008. The VAP protein family: from cellular

functions to motor neuron disease. *Trends Cell Biol*. doi:10.1016/j.tcb.2008.03.006

Liu F, Sells MA, Chernoff J. 1998. Protein tyrosine phosphatase 1B negatively regulates

integrin signaling. *Curr Biol* **8**:173–176. doi:10.1016/S0960-9822(98)70066-1

Mandal K. 2020. Review of PIP2 in cellular signaling, functions and diseases. *Int J Mol Sci*.

doi:10.3390/ijms21218342

Mao D, Lin G, Tepe B, Zuo Z, Tan KL, Senturk M, Zhang S, Arenkiel BR, Sardiello M, Bellen

HJ. 2019. VAMP associated proteins are required for autophagic and lysosomal

degradation by promoting a PtdIns4P-mediated endosomal pathway. *Autophagy*

15:1214–1233. doi:10.1080/15548627.2019.1580103

Martel V, Racaud-Sultan C, Dupe S, Marie C, Paulhe F, Galmiche A, Block MR, Albiges-Rizo

C. 2001. Conformation, Localization, and Integrin Binding of Talin Depend on Its

Interaction with Phosphoinositides. *J Biol Chem* **276**:21217–21227.

doi:10.1074/jbc.M102373200

Mesmin B, Bigay J, Moser Von Filseck J, Lacas-Gervais S, Drin G, Antonny B. 2013. XA

four-step cycle driven by PI(4)P hydrolysis directs sterol/PI(4)P exchange by the ER-

Golgi Tether OSBP. *Cell* **155**:830. doi:10.1016/j.cell.2013.09.056

Moorhead AM, Jung JY, Smirnov A, Kaufer S, Scidmore MA. 2010. Multiple host proteins

that function in phosphatidylinositol-4-phosphate metabolism are recruited to the

chlamydial inclusion. *Infect Immun* **78**:1990–2007. doi:10.1128/IAI.01340-09

Murphy SE, Levine TP. 2016. VAP, a Versatile Access Point for the Endoplasmic Reticulum:

Review and analysis of FFAT-like motifs in the VAPome. *Biochim Biophys Acta - Mol*

Cell Biol Lipids **1861**:952–961. doi:10.1016/J.BBALIP.2016.02.009

Nader GPF, Ezratty EJ, Gundersen GG. 2016. FAK, talin and PIPKI 3 regulate endocytosed

integrin activation to polarize focal adhesion assembly. *Nat Cell Biol* **18**:491–503.

doi:10.1038/ncb3333

Ng IC, Pawijit P, Teo LY, Li H, Lee SY, Yu H. 2016. Kinectin-dependent ER transport

supports the focal complex maturation required for chemotaxis in shallow gradients. *J*

Cell Sci **129**:2660–2672. doi:10.1242/jcs.181768

Peretti D, Dahan N, Shimoni E, Hirschberg K, Lev S. 2008. Coordinated lipid transfer

between the endoplasmic reticulum and the golgi complex requires the VAP proteins

and is essential for Golgi-mediated transport. *Mol Biol Cell* **19**:3871–3884.

doi:10.1091/mbc.E08-05-0498

Prinz WA, Toulmay A, Balla T. 2020. The functional universe of membrane contact sites. *Nat*

Rev Mol Cell Biol **21**:7–24. doi:10.1038/s41580-019-0180-9

Scorrano L, De Matteis MA, Emr S, Giordano F, Hajnóczky G, Kornmann B, Lackner LL,

Levine TP, Pellegrini L, Reinisch K, Rizzuto R, Simmen T, Stenmark H, Ungermann C,

Schuldiner M. 2019. Coming together to define membrane contact sites. *Nat Commun*

10:1287. doi:10.1038/s41467-019-09253-3

Senju Y, Kalimeri M, Koskela E V., Somerharju P, Zhao H, Vattulainen I, Lappalainen P.

2017. Mechanistic principles underlying regulation of the actin cytoskeleton by

phosphoinositides. *Proc Natl Acad Sci U S A* **114**:E8977–E8986.

doi:10.1073/pnas.1705032114

Stefan CJ, Manford AG, Baird D, Yamada-Hanff J, Mao Y, Emr SD. 2011. Osh proteins

regulate phosphoinositide metabolism at ER-plasma membrane contact sites. *Cell*

144:389–401. doi:10.1016/j.cell.2010.12.034

Thapa N, Sun Y, Schramp M, Choi S, Ling K, Anderson RA. 2012. Phosphoinositide

signaling regulates the exocyst complex and polarized integrin trafficking in directionally

migrating cells. *Dev Cell* **22**:116–30. doi:10.1016/j.devcel.2011.10.030

Tsujita K, Itoh T. 2015. Phosphoinositides in the regulation of actin cortex and cell migration.

Biochim Biophys Acta - Mol Cell Biol Lipids **1851**:824–831.

doi:10.1016/j.bbalip.2014.10.011

Vance JE. 1990. Phospholipid synthesis in a membrane fraction associated with

mitochondria. *J Biol Chem* **265**:7248–7256. doi:10.1016/S0021-9258(19)39106-9

Weber-Boyvat M, Kentala H, Lilja J, Vihervaara T, Hanninen R, Zhou Y, Peränen J, Nyman

TA, Ivaska J, Olkkonen VM. 2015. OSBP-related protein 3 (ORP3) coupling with VAMP-

associated protein A regulates R-Ras activity. *Exp Cell Res* **331**:278–291.

doi:10.1016/j.yexcr.2014.10.019

Welch HCE, Coadwell WJ, Ellson CD, Ferguson GJ, Andrews SR, Erdjument-Bromage H,

Tempst P, Hawkins PT, Stephens LR. 2002. P-Rex1, a PtdIns(3,4,5)P3- and

Gbetagamma-regulated guanine-nucleotide exchange factor for Rac. *Cell* **108**:809–21.

Wilhelm LP, Wendling C, Védie B, Kobayashi T, Chenard M, Tomasetto C, Drin G, Alpy F.

2017. STARD 3 mediates endoplasmic reticulum-to-endosome cholesterol transport at

671 membrane contact sites . *EMBO J* **36**:1412–1433. doi:10.15252/embj.201695917

672 Wu H, Carvalho P, Voeltz GK. 2018. Here, there, and everywhere: The importance of ER

673  membrane contact sites. *Science (80-)* **361**. doi:10.1126/science.aan5835

674 Xu W, Wang P, Petri B, Zhang Y, Tang W, Sun L, Kress H, Mann T, Shi Y, Kubes P, Wu D.

675  2010. Integrin-induced PIP5K1C kinase polarization regulates neutrophil polarization,

676  directionality, and in vivo infiltration. *Immunity* **33**:340–50.

677  doi:10.1016/j.immuni.2010.08.015

678 Yoshii S, Tanaka M, Otsuki Y, Wang DY, Guo RJ, Zhu Y, Takeda R, Hanai H, Kaneko E,

679  Sugimura H. 1999. alphaPIX nucleotide exchange factor is activated by interaction with

680  phosphatidylinositol 3-kinase. *Oncogene* **18**:5680–90. doi:10.1038/sj.onc.1202936

 FIGURE LEGENDS

Figure 1. VAPA is required for proper cell migration and cell spreading.

A. Left panel: Representative immunoblots showing the levels of VAPA and Tubulin in Control

and VAPA KO cells. Tubulin expression level was used as loading control. Right panel:

quantification of relative VAPA density in Control and VAPA KO cells normalized to Tubulin

levels (mean±SEM from three independent experiments). **B.** Confocal images of Control and

VAPA KO cells immunostained for VAPA. Scale bar 20 µm. **C.** Phase contrast images of

Control and VAPA KO cells migrating collectively 48h after space release. The trajectories of

a Control and VAPA KO leader cell over 20h are shown in green. Scale bar: 100 µm. **D.**

Analysis of relative colonized surface within the last 20h by Control and VAPA KO cells

(mean±SEM from three independent experiments). Data were analysed using a paired t-test.

E.F. Analysis of Control and VAPA KO leader cell velocity (**E**) and directionality coefficient (**F**)

calculated as the ratio between the net displacement and the trajectory length within the last

20h of collective migration (mean±SEM, Control: n=33 cells; VAPA KO: n=36 cells, from 3

independent experiments). **G.H.** Analysis of cell velocity (**G**) and directionality coefficient (**H**)

of Control and VAPA KO individual cells displacing on fibronectin-coated glass during at least

3h (mean±SEM; Control: n=64 cells; VAPA KO: n=33 cells from 2 independent experiments).

I. Epifluorescence images of Control and VAPA KO cells after 1 hour spreading on fibronectin-

coated glass and stained as indicated. Scale bar: 20 µm. **J.** Analysis of Control and VAPA KO

cells area after 1 hour spreading on fibronectin-coated glass (mean±SEM; Control: n=170

cells; VAPA KO: n=163 cells from 3 independent experiments). All data were analysed using

non parametric Mann-Whitney t-test. (ns: non significant, ***P-values <0.001).

Figure 2. VAPA regulates focal adhesions and actin cytoskeleton through its MSP domain.

A. Confocal images with zoom boxes of paxillin-labeled focal adhesions in leader cells from Control, VAPA KO, VAPA KO stably expressing wild-type mCherry-VAPA, or mCherry-VAPA KD/MD cell lines. Scale bar: 20 µm (5 µm in insets). **B**. Confocal images of actin cytoskeleton network in Control and VAPA KO leader cells stained for cortactin and F-actin. Magenta arrows point to cortactin-enriched domains at the leading edge. Green arrows point to F-Actin transversal arcs. Scale bar: 20 µm. **C.** Differential Interference Contrast (DIC) images (left) of Control and VAPA KO leader cells and kymographs (right) along the yellow lines from a 30 min movie at 1 frame every 3 seconds, showing protrusion and retraction phases of the leading edge. Scale bar: 20 µm (left) and 5 µm (right)**. D.** Analysis of focal adhesion area quantified from images in A, in Control (n=2968 focal adhesions from 20 cells), VAPA KO (n=4329 focal adhesions from 22 cells), VAPA KO expressing mCherry-VAPA, (n= 3818 focal adhesions from 24 cells) and VAPA KO expressing mCherry-VAPA KD/MD (n=2801 focal adhesions from 19 cells) leader cells, from 3 independent experiments **E.** Analysis of leading edge proportion enriched with cortactin staining quantified from images in B, in Control, VAPA KO, VAPA KO expressing wild-type mCherry-VAPA and VAPA KO expressing mCherry-VAPA KD/MD leader cells, (mean±SEM; n=29 cells for each cell line, from 3 independent experiments) .**F.** Analysis of the size of cortactin-enriched domains at the leading edge, quantified from images in B, in Control and VAPA KO leader cells (mean±SEM; Control n= 77 cells; VAPA KO n=72 cells, from 3 independent experiments) **G.** Quantification of protrusion phases frequency per 30min quantified from the kymographs in C, in Control and VAPA KO leader cells (mean±SEM; Control n=16 cells; VAPA KO: n=21 cells from 3 independent experiments). All data were analysed using non parametric Mann-Whitney t-test (ns: non significant, ***P-values <0.001).

Figure 3. VAPA controls PInst homeostasis at the PM, but not in other compartments

A. Confocal images and zoom boxes of PI(4)P distribution in Control and VAPA KO leader cells expressing GFP-PH-OSBP and immunostained for EEA1 or TGN46. Scale Bar: 10 μm. **B.** Analysis of colocalization by Pearson coefficient determination between GFP-PH-OSBP and EEA1 or TGN46 signals, quantified from images in A, in Control and VAPA KO leader cells (mean±SEM; Control: n=9 cells and 12 cells respectively; VAPA KO: n=9 cells and 12 cells respectively, from 3 independent experiments). **C.** Analysis of Golgi(TGN46)/cytosol (left) and Endosomes(EEA1)/Cytosol (right) ratio of GFP-PH-OSBP signal, quantified from images in A, in Control and VAPA KO leader cells (mean ±SEM; Control: n=12 cells; VAPA KO: n=11 cells, from 3 independent experiments) . **D.** Analysis of RFP-PH-PLCδ1 peak intensity, quantified from images in F, in Control and VAPA KO leader cells (mean±SEM; Control: n=58 cells; VAPA KO: n=53 cells, from 3 independent experiments). **E.** Analysis of PM/Cytosol ratio of RFP-PH-PLCδ1, quantified from images in G, at the leading edge of Control and VAPA KO leader cells (mean±SEM; Control: n=27 cells; VAPA KO: n=26 cells, from 3 independent experiments). **F.** XZ view of confocal images of Control and VAPA KO cells in cluster immunostained for PI(4,5)P2. Scale Bar: 5 μm. **G.** Confocal images of PI(4,5)P2 distribution in Control and VAPA KO leader cells expressing RFP-PH-PLCδ1, represented as a color-coded heat map. Scale bar: 10 μm. All data were analysed using non parametric Mann-Whitney t-test (ns: non significant, ***P-values <0.001).

Figure 4. VAPA stabilizes ventral ER-PM contact sites at the front of migrating cells

A. TIRF microscopy images of GFP-MAPPER foci distribution along the ER in Control and VAPA KO leader cells expressing GFP-MAPPER and RFP-KDEL. Scale bar: 5 μm. **B.** Sequential TIRF microscopy images of ER and GFP-MAPPER foci accumulation at the front of a Control leader cell expressing GFP-MAPPER and RFP-KDEL. Scale bar: 1 μm. **C.** Sequential spinning disk images of dorsal (magenta) and ventral (green) GFP-MAPPER foci at the front of a Control leader cell expressing GFP-MAPPER. Top and bottom plane images

were color-coded in magenta and green respectively. Circles highlight single GFP-MAPPER

foci at the dorsal (blue) and ventral (red) sides. Scale bar: 5 µm. **D.** Transmission Electron

Microscopy images of transversal cuts of a Control leader cell, showing the leading edge.

Arrows point to ER-PM contact sites at the bottom of the cell. Scale bar: 1 µm (top) and 100

nm in insets 1 and 2. **E.** Sequential TIRF microscopy images of GFP-MAPPER foci at the front

of Control and VAPA KO leader cells expressing GFP-MAPPER. Individual GFP-MAPPER foci

are pictured in the frames below images. Scale bar: 1 µm. **F.G.** Analysis of the lifetime (**F**) and

the speed (**G**) of ventral GFP-MAPPER foci, quantified from images in E, in Control and VAPA

KO leader cells (mean±SEM; Control: n=8 cells; VAPA KO: n=6 cells; at least 10 ER-PM MCS

were analysed per cell, from 4 independent experiments). All data were analysed using non

parametric Mann-Whitney t-test (***P-values <0.001).

Figure 5. VAPA promotes microtubule-dependent FA disassembly

A-C Analysis of assembly rate (**A**), disassembly rate (**B**) and lifetime (**C**) of focal adhesions

(FA), quantified from time-lapse images in E, in Control and VAPA KO leader cells

(mean±SEM; Panels (A) and (B): Control: n=217 FA from 9 cells; VAPA KO: n=342 FA from 8

cells, from 3 independent experiments. Panel (C): Control: n=133 FA from 9 cells; VAPA KO:

n=126 FA from 8 cells, from 3 independent experiments). **D.** Distribution of focal adhesion (FA)

life times, quantified from time-lapse images in E, in Control and VAPA KO leader cells

(Control: n=133 FA from 9 cells; VAPA KO: n=126 FA from 8 cells, from 3 independent

experiments). **E.** Sequential TIRF microscopy images of focal adhesions in Control and VAPA

KO leader cells expressing mCherry-Vinculin. Scale Bar: 1 µm. **F.** Sequential Confocal images

of focal adhesions after nocodazole treatment and wash-out in migrating Control and VAPA

KO leader cells immunostained for paxillin. Scale bar: 10 µm (2 µm in insets). **G** Analysis of

relative focal adhesions (FA) size, quantified from images in F, in Control and VAPA KO leader

cells after 0 min, 15 min, 30 min and 60 min after nocodazole wash-out (mean±SEM. FA from

22 to 25 cells were analysed, from 3 independent experiments. Control T0min: n=6053 FA;

Control T15 min: n=5146 FA; Control T30 min: n=5543 FA; Control T60 min: n=3913 FA; VAPA

KO T0 min: n=4481 FA, VAPA KO T15 min: n=3878 FA, VAPA KO T30 min: n= 4165 FA ; T60

min : n=4889 FA). All data were analysed using non parametric Mann-Whitney t-test (ns: non

significant, ***P-values <0.001).

Figure 6. VAPA mediates the anchoring of ER-PM contact sites to FA

A. TIRF microscopy images of ER-PM contact sites and focal adhesions in Control and VAPA

KO leader cells expressing GFP-MAPPER and mCherry-Vinculin. Scale bars: 10 µm (1 µm in

insets). **B.** Analysis of the percentage of focal adhesions in contact with ER-PM contact sites,

quantified from images in A, in Control and VAPA KO leader cells (mean±SEM; Control: n=12

cells; VAPA KO: n=10 cells, from 3 independent experiments). **C.** Sequential TIRF microscopy

images of ER-PM contact sites and focal adhesions in a Control leader cell expressing GFP-

MAPPER and mCherry-Vinculin. Scale bar: 1 µm. **D.** Time course of GFP-MAPPER (green)

and mCherry-Vinculin (magenta) signals during the lifetime of the focal adhesion pointed in C,

within a region that comprises the entire FA + 2 pixels. **E.** Time course of mCherry-Vinculin

and GFP-MAPPER signals obtained by TIRF microscopy in Control leader cells, as in C.,

starting 4 minutes before the accumulation of GFP-MAPPER in FA regions (comprising the

entire FA + 2 pixels). The signals were all smoothed, readjusted to the minimal value and

expressed as % of the maximal value (mean±SEM; n=14 FA from 5 cells, from 3 independent

experiments). All data were analysed using non parametric Mann-Whitney t-test (***P-values

<0.001).

Figure 7

Hypothetic model depicting the role of VAPA in FA disassembly, based on our observations

and previously published data (see Discussion part).

Figure 1

Figure 2

Figure 4

Figure 5

A GFP MAPPER | mCherry Vinculin | Merge

Control

VAPA KO

B Percentage of focal adhesion in contact with ER-PM MCS (%)

***

Control VAPA KO

C T=0 T=13min T=20min T=33min T=50min

mCherry-Vinculin

GFP-MAPPER

Merge

D mCherry-Vinculin
GFP MAPPER

Fluorescence intensity (.10³ A.U)

Time (min)

E GFP-MAPPER Anchoring

% of Max Ch-Vinculin

% of Max GFP-MAPPER

Time (min)

FA disassembly mediated by VAPA

Internalization of integrins through
microtubule- and clathrin-dependent endocytosis