**Class IIa HDACs reprogram mitochondrial metabolism to inhibit apoptosis and ferroptosis in response to lipotoxicity**

Sheree D. Martin[1,9], Timothy Connor[1,9], Andrew Sanigorski[1], Kevin A. McEwen[1], Darren C. Henstridge[2,3], Brunda Nijagal[4], David De Souza[4], Dedreia L. Tull[4], Peter J. Meikle[3], Greg M. Kowalski[1,5], Clinton R. Bruce[5], Paul Gregorevic[6], Mark A. Febbraio[7], Fiona M. Collier[8], Ken R. Walder[1], Sean L. McGee[1,10]

[1]Institute for Mental and Physical Heath and Clinical Translation (IMPACT) and Metabolic Research Unit, School of Medicine, Deakin University, Geelong, Victoria 3216, Australia;
[2]College of Health and Medicine, School of Health Sciences, University of Tasmania, Launceston, Australia;
[3]Baker Heart and Diabetes Institute, Melbourne, Victoria 3004, Australia;
[4]Metabolomics Australia, The University of Melbourne, Parkville, Victoria 3010, Australia;
[5]Institute of Physical Activity and Nutrition (IPAN) and School of Exercise and Nutrition Sciences, Deakin University, Geelong, Victoria 3216, Australia;
[6]Centre for Muscle Research, Department of Anatomy and Physiology, The University of Melbourne, Parkville, VIC, Australia.
[7]Monash Institute of Pharmaceutical Sciences, Monash University, Parkville, VIC, Australia.
[8]Barwon Health, Geelong, Victoria 3220, Australia
[9]These authors contributed equally to this work
[10]Lead contact

Lead contact:

Professor Sean L. McGee
Metabolic Remodelling Laboratory
Metabolic Research Unit
School of Medicine
Deakin University, Waurn Ponds, Australia, 3216
Ph: +61 3 5227 2519
Fax: +61 3 5227 2945
Email: sean.mcgee@deakin.edu.au

**ABSTRACT**

Lipotoxicity, the accumulation of lipids in non-adipose tissues, alters the metabolic transcriptome and mitochondrial metabolism in skeletal muscle. The mechanisms involved remain poorly understood. Here we show that lipotoxicity increased histone deacetylase 4 (HDAC4) and histone deacetylase 5 (HDAC5), which reduced the expression of metabolic genes and oxidative metabolism in skeletal muscle. This metabolic reprogramming was linked with reduced expression of p53-dependent genes that mediate apoptosis and ferroptosis, which preserved cell viability in response to lipotoxicity. Mechanistically, impaired mitochondrial metabolism reduced acetylation of p53 at K120, a modification required for transcriptional activation of apoptosis, while redox drivers of ferroptosis were also reduced. Overexpression of loss-of-function HDAC4 and HDAC5 mutants in skeletal muscle of obese *db/db* mice enhanced oxidative capacity, increased apoptosis and ferroptosis and reduced muscle mass. This study identifies HDAC4 and HDAC5 as repressors of the oxidative state of skeletal muscle, and that this metabolic reprogramming, considered deleterious for normal metabolism, is critical to preserve muscle integrity in response to lipotoxicity.

## INTRODUCTION

Obesity is driving an epidemic of chronic disease, including type 2 diabetes, non-alcoholic fatty liver disease, cardiovascular disease and cancer. Linked to many of these diseases are adverse cellular responses arising from the accumulation of lipids in non-adipose tissues, a process that has been termed lipotoxicity[1]. The deleterious effects of lipotoxicity, which include altered cellular signalling, metabolic reprogramming and apoptotic cell death[2], are thought to be mediated by bioactive lipid species such as diglycerides (DG) and ceramides[3]. In contrast, the accumulation of neutral lipids such as triglycerides (TG) is thought to protect against the harmful effects of lipotoxicity[4]. Tissues that are particularly susceptible to lipotoxicity include the liver, kidneys, pancreas, heart and skeletal muscle[2].

Alterations in skeletal muscle metabolism occur in obesity, including insulin resistance and impairments in several aspects of oxidative metabolism. Reduced rates of both glucose and lipid oxidation have been observed in the skeletal muscle of obese patients[5, 6] and data describing alterations in amino acid metabolism are also emerging[7, 8, 9]. While methodological caveats to some of these studies have been raised[10], these impairments in oxidative metabolism are associated with reduced expression of a metabolic and mitochondrial transcriptional program controlled by the peroxisome proliferator-activated receptor-gamma coactivator 1 alpha (PGC-1$\alpha$) transcriptional coactivator[11], encoded by the *PPARGC1A gene*. Furthermore, down-regulation of this PGC-1$\alpha$-dependent transcriptional program is also associated with insulin resistance[12], although studies using several animal models have questioned the relationship between PGC-1$\alpha$ and skeletal muscle insulin action[13, 14].

Experimental induction of acute lipotoxicity in humans through lipid infusion results in many of the same skeletal muscle metabolic alterations that are observed in obesity. These include insulin resistance, reduced substrate oxidation and impairment of mitochondrial function[15], as well as reduced expression of *PPARGC1A* and genes involved in mitochondrial and metabolic processes[16]. Moreover, transcriptional reprogramming of metabolism, like that in obesity, is observed when muscle cells are cultured with elevated concentrations of the saturated fatty acid palmitate[17, 18, 19]. It is not known why repression of the metabolic transcriptome and reduced oxidative metabolism is an adaptation to lipotoxicity and the molecular pathways involved remain unclear. Furthermore, how these metabolic adaptations influence other responses to lipotoxicity, such as cell signalling and cell death responses, are unknown.

Studies from our group and others have found that the class IIa histone deacetylase (HDAC) isoforms 4 and 5 are regulators of PGC-1$\alpha$ expression and oxidative metabolism in skeletal muscle[20, 21]. HDAC4 and 5 act co-operatively as heterodimers to repress the myocyte-enhancer factor 2 (MEF2) family of transcription factors[22, 23]. The ability of the class IIa HDACs to repress MEF2 transcriptional activity is prevented by phosphorylation-dependent nuclear export[24]. HDAC4 and 5 are also sensitive to energetic stress, such as during exercise, through phosphorylation by the AMP-activated protein kinase (AMPK)[25, 26, 27], which in turn activates transcriptional programs that enhance oxidative metabolism and capacity, including expression of PGC-1$\alpha$ and its transcriptional programs[20, 27, 28]. Binding sites for the MEF2 transcription factor are found in the *PPARGC1A* promoter[21] and in the promoters and

enhancer regions of other genes involved in oxidative metabolism[20]. Together, these
findings suggest that the class IIa HDACs are a critical hub regulating skeletal muscle
metabolic transcriptional programs in response to changes in energy balance. However, the
role of the class IIa HDAC transcriptional repressors in regulating metabolic transcriptional
programs and oxidative metabolism in the context of lipotoxicity is unknown. The current
study sought to determine whether HDAC4 and 5 contribute to the suppression of
metabolic genes and oxidative metabolism in response to lipotoxicity and to explain why
these adaptations occur.

**RESULTS**

**HDAC4 and 5 are increased in models of lipotoxicity**

To test the hypothesis that HDAC4 and 5 are increased by lipotoxicity, cultures of C2C12 myotubes were exposed to 0.25 or 0.5mM palmitate, or BSA (vehicle), for 16 hr. Exposure to both 0.25 and 0.5mM palmitate did not negatively impact cell viability, assessed by lactate dehydrogenase (LDH) release (Supplementary Fig. 1A). Lipidomic profiling indicated ceramide and DG species were increased following exposure to 0.5mM palmitate, while increases in some TG species were observed following exposure to both 0.25mM and 0.5mM palmitate (Figure 1A). Total Cer levels were increased by both 0.25mM and 0.5mM palmitate while total DG and TG levels were increased by 0.5mM palmitate (Figure 1B). Consistent with the known effect of palmitate on C2C12 cells, application of 0.5mM palmitate was associated with a non-significant trend for *Ppargc1a* gene expression to be reduced (Fig. 1C) and a significant reduction in cellular respiration (Fig. 1D). The abundance of HDAC5 protein was increased by both 0.25 and 0.5mM palmitate, while HDAC4 protein abundance was increased by 0.5mM palmitate (Figure 1E). There were no differences in *Hdac4* gene expression between groups, while *Hdac5* gene expression was reduced by 0.5mM palmitate (Supplementary Fig. 1B).

To examine these mechanisms *in* vivo, the tibialis anterior skeletal muscle of obese *db/db* and control heterozygous littermate mice was analysed. Specific ceramide, DG and TG species (Fig. 1F) and total ceramides and DG were increased in skeletal muscle of *db/db* mice (Fig. 1G). The increased muscles lipid concentrations in these obese mice were associated with decreased *Ppargc-1α* gene expression (Fig. 1H), reduced respiratory responses in skeletal muscle biopsies (Fig. 1I) and increased HDAC4 and 5 protein abundance (Fig. 1J). Skeletal muscle gene expression levels of *Hdac5*, but not *Hdac4*, were increased in *db/db* mice (Supplementary Fig. 1C). Together, these data suggest that skeletal muscle HDAC4 and 5 are increased in response to lipotoxicity and are associated with reduced *Ppargc1a* expression and impaired oxidative metabolism.

**Overexpression of HDAC4 and 5 in skeletal muscle represses *Pppargc1a* expression and oxidative capacity**

To determine whether increased HDAC4 and 5 reduces *Ppargc1a* gene expression and impairs oxidative capacity in skeletal muscle, a bilateral skeletal muscle HDAC4 and 5 mouse model was developed using recombinant serotype 6 adeno-associated viral (AAV6) vectors expressing HDAC4 and HDAC5 (Fig. 2A). Administration of HDAC4 and 5 AAV6 vectors increased *Hdac4* and *Hdac5* gene expression (Fig. 2B) and HDAC4 and 5 protein (Fig. 2C) in tibialis anterior muscle. Overexpression of HDAC4 and 5 reduced *Ppargc1a* expression (Fig. 2D). Furthermore, HDAC4 and 5 overexpression reduced the expression of a range of MEF2 and PGC-1α target genes involved in oxidative and mitochondrial metabolism (Fig. 2E). These changes in metabolic transcripts were associated with reduced abundance of ATP synthase subunits, but not other subunits of electron transport chain complexes (Fig. 2F). To establish whether reduced ATP synthase components resulted in reduced oxidative capacity, respiration was assessed in skeletal muscle biopsies. There were overall main effects for respiratory responses to be reduced in muscles overexpressing HDAC4 and 5

when driven by both succinate (Fig. 2G) and malate (Fig. 2H) as substrates. The reduction in
oxidative capacity in skeletal muscle overexpressing HDAC4 and 5 was not associated with
the accumulation of any lipid species examined, although all lipid species tended to be
higher in skeletal muscle overexpressing HDAC4 and 5 (Fig. 2I). These data indicate that
increasing HDAC4 and 5 represses *Ppargc1a* gene expression and oxidative metabolism in
skeletal muscle.
168
The class IIa HDACs form heterodimers, which influences their repressive functions and
sensitivity to upstream signalling pathways[22]. To investigate the separate role of HDAC4 and
5 in repression of *Ppargc1a* gene expression, HDAC4 and 5 were individually overexpressed
in skeletal muscle with increasing doses of AAV6 vectors (Supplementary Fig. 2A).
Treatment of muscles with AAV vectors expressing HDAC4 or HDAC5 resulted in dose
dependent increases in the expression of *Hdac4* and *Hdac5* genes (Supplementary Fig. 2B)
and increased HDAC4 and 5 protein abundance (Supplementary Fig. 2C). Overexpression of
HDAC4 at both AAV6 doses reduced *Ppargc1a* gene expression (Supplementary Fig. 2D).
Similarly, overexpression of HDAC5 using both AAV doses was sufficient to reduce *Ppargc1a*
expression (Supplementary Fig. 2E). Together, these data suggest that increases in either
HDAC4 or HDAC5 can repress *Ppargc1a* gene expression.
180
181    **Overexpression of HDAC4 and 5 in skeletal muscle inhibits oxidative glucose utilisation**
182
Impaired skeletal muscle oxidative metabolism and mitochondrial function associated with
lipotoxicity have been linked with reduced insulin action and impaired glucose utilisation[6, 12,
29]. As HDAC4 and 5 overexpression reduced skeletal muscle oxidative capacity, the impact of
HDAC4 and 5 overexpression on glucose metabolism was examined. Bilateral skeletal
muscle HDAC4 and 5 mice were administered vehicle or insulin (1.2U/kg i.p.) with $^3$H-2-
deoxyglucose (2-DG) and $^{14}$C-glucose tracers 30 min prior to tissues collection (Fig. 3A).
Insulin reduced the excursion of both tracers (Supplementary Fig. 3A and B). There was a
main effect for increased glucose clearance by skeletal muscles overexpressing HDAC4 and 5
(Fig. 3B). When expressed relative to control skeletal muscle within each mouse, muscles
with HDAC4 and 5 overexpression increased glucose clearance in vehicle but not insulin-
stimulated conditions (Figure 3C). Overexpression of HDAC4 and 5 increased glucose
incorporation into glycogen (Fig. 3D) and increased glycogen concentration (Supplementary
Fig. 3C). In contrast there was a main effect for insulin to increase glucose incorporation into
lipids (Fig. 3E), which was associated with main effects for insulin to increase Akt
phosphorylation at both of its main regulatory sites at T308 (Fig. 3G) and S473 (Fig. 3H).
However, consistent with the glucose clearance data, there was a main effect for increased
phosphorylation of TBC1D4 at S642 in skeletal muscles overexpressing HDAC4 and 5 (Fig.
3I). These data suggest that overexpression of HDAC4 and 5 increased non-oxidative glucose
utilisation in skeletal muscle.
202
To further examine the effect of HDAC4 and 5 overexpression on oxidative glucose
utilisation, a targeted stable isotope metabolomics approach following an oral bolus of
U$^{13}$C-glucose was used in bilateral skeletal muscle HDAC4 and 5 mice[30] (Fig. 3J).
Overexpression of HDAC4 and 5 increased the abundance of lactate and succinate (Fig. 3K)
and total $^{13}$C labelling of succinate and malate (Fig. 3L). Isotopologue analysis revealed
accumulation of both M+0 and M+2 lactate in muscle overexpressing HDAC4 and 5 (Fig.

3M), suggesting both increased non-oxidative glucose utilisation and increased
malate/pyruvate cycle flux, respectively[31]. Succinate M+0, M+1 and M+2 isotopologues
were increased in muscle overexpressing HDAC4 and 5 (Fig. 3N), consistent with a major
impairment in SDH activity[32], which is bifunctional for both the TCA cycle and the electron
transport chain. These data are also supported by reduced succinate-driven respiration in
muscles overexpressing HDAC4 and 5 (Fig. 2G). Similarly, HDAC4 and 5 overexpression was
associated with increased malate M+0 and M+1 isotopologues (Fig. 3O). Together, these
data indicate that increased HDAC4 and 5 is linked with defective oxidative utilisation of
glucose in skeletal muscle.
218
219  **HDAC4 and 5 enhance cell survival in response to lipotoxicity and supress the apoptosis**
220  **and ferroptosis cell death pathways**
221
Overexpression of HDAC4 and 5 impaired mitochondrial function and reprogrammed
glucose metabolism, increasing non-oxidative glucose utilisation and inhibiting oxidative
glucose utilisation. In doing so, increased HDAC4 and 5 replicate some, but not all, of the
metabolic alterations seen during lipotoxicity. To better understand why this metabolic
reprogramming occurs, a comparative transcriptomics approach was employed. Specifically,
the transcriptome altered in the skeletal muscle of *db/db* mice in which HDAC4 and 5 was
increased (Fig. 1J) was compared to the transcriptome altered in C2C12 myoblasts stably
overexpressing HDAC4 and 5 (Supplementary Fig. 4A). These cultured C2C12 cells were used
to increase the biological diversity in this comparison and were analysed as myoblasts, as
HDAC4 and 5 overexpression prevented their differentiation into myotubes (data not
shown). Nonetheless, HDAC4 and 5 overexpression in C2C12 myoblasts induced phenotypic
characteristics consistent with HDAC4 and 5 overexpression in skeletal muscle, including
reduced *Ppargc1a* gene expression, suppression of a broad metabolic transcriptional
program, including reduced OXPHOS and TCA cycle pathways, and reduced oxidative
capacity (Supplementary Fig. 4B-D). Comparison of the transcriptome altered in the skeletal
muscle of *db/db* mice to that altered in C2C12 myoblasts overexpressing HDAC4 and 5 and
subsequent pathway analysis of coregulated genes revealed the p53 signalling pathway as
the only significantly regulated pathway (Supplementary Fig. 4E).
240
Profiling of p53-dependent genes in C2C12 myoblasts overexpressing HDAC4 and 5 revealed
that several pro-apoptosis genes were reduced, including *Casp9, Cycs, Pmaip1* and *Bbc3*
(Fig. 4A), consistent with suppression of apoptosis. Similarly, genes involved in ferroptosis,
which is a form of iron-mediated cell death involving lipid peroxidation[33], were altered in a
way that indicated suppression of ferroptosis, via increased *Slc7a11* expression and reduced
*Sat1* and *Alox15* expression (Fig. 4A). Together, these data suggest that increased HDAC4
and 5 inhibit apoptotic and ferroptotic cell death pathways. To test this hypothesis, C2C12
myoblasts overexpressing HDAC4 and 5 were treated with agents that induce either
apoptosis or ferroptosis. Overexpression of HDAC4 and 5 in C2C12 myoblasts reduced the
number of early apoptotic cells (Annexin$^+$/PI$^-$) following exposure to the apoptosis-inducing
agent camptothecin (Fig. 4B). Similarly, overexpression of HDAC4 and 5 increased the
percentage of viable cells following exposure to the ferroptosis-inducing agent RSL3 (Fig.
4C). These data raised the possibility that HDAC4 and 5 suppress the apoptosis and
ferroptosis pathways to enhance cell survival in response to lipotoxicity. To test this
hypothesis, cells were exposed to high concentrations of palmitate that were intended to

induce cytotoxicity. Overexpression of HDAC4 and 5 increased the viability of cells exposed
to both 1 and 2mM of palmitate compared with control cells and resulted in cells being
largely resistant to lipotoxicity (Fig. 4D). This protection of cell viability in response to
lipotoxicity was associated with reduced cleavage of caspase 3 and 9, the irreversible events
that trigger apoptosis (Fig. 4E), as well as fewer cells in both early (Fig. 4F) and late (Fig. 4G)
apoptosis in response to increasing concentrations of palmitate. Similarly, overexpression of
HDAC4 and 5 reduced lipid peroxidation, the trigger for ferroptosis induced-cell death, in
response to increasing concentrations of palmitate (Fig. 4H). These data indicate that
increased HDAC4 and 5 inhibit the apoptosis and ferroptosis cell death pathways and
enhance cell survival in response to lipotoxicity.
266
267 **HDAC4 and 5 inhibit apoptosis by reducing p53 K120 acetylation, which is regulated by**
268 **HDAC4 and 5-mediated alterations in metabolism**
269
In several cancer cell types, increased HDAC5 has been linked with reduced acetylation of
p53 at K120, which inhibits p53 DNA binding and the expression of p53-dependent
apoptosis genes, and reduces apoptosis sensitivity[34, 35, 36]. Furthermore, acetylation of p53,
including at K120, is linked to the transcriptional regulation of ferroptosis[37]. To explore
whether reduced acetylation of p53 at K120 by HDAC4 and 5 is involved in suppression of
apoptosis and ferroptosis in myogenic cells, C2C12 myoblasts stably overexpressing HDAC4
and 5 were transfected with wild type p53, or p53 in which K120 was mutated to either
glutamine (K120Q; acetylation mimetic gain-of function) or arginine (K120R; deacetylation
mimetic loss-of-function) (Fig. 5A). The acetylation of p53 at K120 was reduced by the
overexpression of HDAC4 and 5 (Fig. 5A). In cells overexpressing HDAC4 and 5, introduction
of the p53$^{K120Q}$ acetylation mimetic mutant restored the expression of the pro-apoptosis
genes *Casp3, Casp9* and *Cycs* (Fig. 5B). However, the expression of *Slc7a11*, which opposes
ferroptosis, was not influenced by expression of this mutant (Fig. 5B). This suggests that
reduced acetylation of p53 at K120 by increased HDAC4 and 5 is involved in the
transcriptional suppression of apoptosis, but not ferroptosis cell death pathways. Indeed,
expression of p53$^{K120Q}$ also restored apoptosis in HDAC4 and 5 overexpressing cells exposed
to the apoptosis-inducing agent camptothecin (Fig. 5C). Although p53 is also known to
regulate aspects of cellular metabolism[38], gain and loss-of-function K120 mutants had no
effect on either oxidative (Supplementary Fig. 5A) or glycolytic flux (Supplementary Fig. 5B).
These data show that reduced acetylation of p53 at K120 contributes to inhibition of
apoptosis by HDAC4 and 5 in myogenic cells.
291
The class IIa HDACs have a unique amino acid substitution within their HDAC active site that
confers low deacetylase activity against acetyl-lysine[39], indicating that it is unlikely that
HDAC4 or 5 are direct p53 deacetylases. We explored the possibility that K120 p53
acetylation is sensitive to the HDAC4 and 5-mediated reduction in oxidative metabolic flux
that reduces acetyl-CoA availability for acetylation reactions. To test this hypothesis,
substrate tracer incorporation into K120 of p53 was examined. Exposure of control cells to
$^{14}$C-palmitate resulted in isotope incorporation into wild type (WT) p53, with only limited
isotope incorporation into mutant p53$^{K120R}$ (Fig. 5D). There was significantly lower tracer
incorporation into both WT and p53$^{K120R}$ in cells overexpressing HDAC4 and 5 (Fig. 5D).
There was negligible tracer incorporation into p53 when cells were exposed to $^{14}$C-glucose

(Fig. 5D). This suggests that carbon derived from fatty acids but not glucose is used for
apoptosis-inducing post-translational modification of p53 at K120.

The acetyl-CoA required for cytosolic and nuclear acetylation reactions is derived from
citrate that is exported from the TCA cycle by the mitochondrial citrate transporter[40]. Once
exported from the mitochondria, citrate is converted to acetyl-CoA by ATP-citrate lyase,
whose activity is increased by insulin[41]. To determine whether HDAC4 and 5 control p53
K120 acetylation via mitochondrial metabolism, cells were treated with vehicle or insulin in
the presence or absence of the citrate transport inhibitor, 1,2,3-benzenetricarboxylate
(BTC). In the non-mitochondrial fraction of control cells, insulin increased acetylation of p53
at K120, which was reduced by exposure to BTC (Fig. 5E). However, acetylation of p53 at
K120 was completely abrogated in cells overexpressing HDAC4 and 5. These findings show
that K120 acetylation is dependent on mitochondrial metabolism and acetyl-CoA
availability. Consistent with a requirement for acetyl-CoA in p53 K120 acetylation, acetyl-
CoA levels were reduced in cells overexpressing HDAC4 and 5 (Fig. 5F).

To determine whether p53 K120 acetylation determines cell viability in response to
lipotoxicity, cells expressing WT p53 or p53$^{K120R}$ were exposed to 2mM palmitate. However,
the p53$^{K120R}$ mutant did not influence cell viability at any time point after palmitate
exposure (Fig. 5G), suggesting that other mechanisms are also involved in the cell death
response to lipotoxicity. Nonetheless, these data show that HDAC4 and 5 inhibit apoptosis
by reducing the acetylation of p53 at K120, which is sensitive to HDAC4 and 5-mediated
alterations in mitochondrial metabolism. However, this inhibition of p53 K120 acetylation
alone is not sufficient to prevent cell death in response to lipotoxicity.

**HDAC4 and 5 regulate multiple metabolic inputs to the ferroptosis pathway**

As suppression of apoptosis through reduced p53 acetylation alone was not sufficient to
explain the protective effect of HDAC4 and 5 on cell survival in response to lipotoxicity, the
role of ferroptosis-mediated mechanisms of cell death (Fig. 6A) were further explored. The
ferroptosis inhibitor Ferrostatin enhanced C2C12 myoblast viability following exposure to
2mM palmitate (Fig. 6B), suggesting that ferroptosis contributes to lipotoxicity-mediated
cell death. Overexpression of HDAC4 and 5 enhanced cell viability in response to increasing
concentrations of Erastin (Fig. 6C), an agent that induces ferroptosis by inhibiting system $x_c$
and glutamate and cysteine exchange[42], which in turn reduces glutathione (GSH) levels and
results in lipid peroxidation through multiple mechanisms (Fig. 6A). Consistent with HDAC4
and 5 increasing the expression of *Slc7a11* (Fig. 4A), overexpression of HDAC4 and 5
reduced intracellular glutamate concentrations, tended to increase intracellular GSH and
increased the GSH/glutamate ratio (Fig. 6D). This was associated with increased glutathione
peroxidase (GPX) activity in cells overexpressing HDAC4 and 5 (Fig. 6E) and reduced total
$H_2O_2$ production (Supplementary Fig. 6A), including from mitochondria and NADPH oxidases
(Fig. 6F). Together, these data indicate that increased HDAC4 and 5 alters the metabolism-
sensitive redox inputs to ferroptosis consistent with inhibition of this pathway.

Ferroptosis involves peroxidation of polyunsaturated (PU) phosphatidylethanolamine (PE)
and phosphatidylcholine (PC) lipids, particularly those with arachidonic acid (AA) or adrenic
acid (Ada) fatty acid species[43]. Changes in the levels of these lipid species have been linked

to alterations in ferroptosis sensitivity[43]. In lipidomic data from skeletal muscle bilateral
HDAC4 and 5 mice, overexpression of HDAC4 and 5 had no effect on any specific PU-PE or
PU-PC species between groups, although most lipid species tended to be higher in muscle
overexpressing HDAC4 and 5 (Supplementary Fig. 6B). However, it should be noted that
HDAC4 and 5 reduced the gene expression of *Alox15* (Fig. 4A), which catalyses lipid
peroxidation (Fig. 6A), and of *Sat1* (Figure 4A), which is a transcriptional regulator of *Alox15*
and a p53 target gene[44]. The extent of lipid saturation has also been linked to lipotoxicity
induced cell death[45], however there were no differences in lipid saturation when expressed
either as absolute abundance (Supplementary Fig. 6C) or proportion of total lipids
(Supplementary Fig. 6D). This suggests that HDAC4 and 5 have minimal effect on the
abundance of lipid species but reduces expression of enzymes that mediate lipid
peroxidation.  Together, these findings indicate that increased HDAC4 and 5 alter cellular
redox balance and expression of *Alox15*, which contributes to inhibition of ferroptosis.
362
363  **HDAC4 and 5 are required to suppress apoptosis and ferroptosis and maintain muscle**
364  **mass in obese mice**
365
To establish whether HDAC4 and 5 suppress apoptosis and ferroptosis in skeletal muscle in
response to lipoxicity *in vivo*, a bilateral skeletal muscle loss-of-function HDAC4 and 5 model
was developed in obese *db/db* mice using AAV6 vectors (Fig. 7A). This loss-of-function
approach employed active-site HDAC4 and 5 mutants (D832N HDAC4 and D861N HDAC5),
which we have previously described as acting in a dominant negative (DN) manner when
overexpressed[20]. Overexpression of these mutants in skeletal muscle increases the
expression of *Ppargc1a* and other metabolic genes and enhance oxidative metabolism[20].
Intramuscular administration of AAV6 vectors expressing DN HDAC4 and 5 to *db/db* mice
increased HDAC4 and 5 gene (Fig. 7B) and protein (Fig. 7C) expression and increased the
expression of *Ppargc1a* (Fig. 7D). Consistent with these findings, there was a main effect for
respiratory responses to be increased in muscle biopsies expressing DN HDAC4 and 5 (Fig.
7E). Expression of DN HDAC4 and 5 modestly increased caspase 3 activity (Fig. 7F) and lipid
peroxidation (Fig. 7G), functional readouts of apoptosis and ferroptosis, respectively.
Examination of the mechanisms involved in HDAC4 and 5-mediated apoptosis revealed that
p53 acetylation of K120 tended to be higher (Fig. 7H), and the expression of p53-dependent
pro-apoptosis genes, including *Casp3, Casp9* and *Bbc3* was increased (Fig. 7I). The
ferroptosis mechanisms regulated by HDAC4 and 5 were also examined. In contrast to
observations in myogenic cells, the expression of *Slc7a11* in skeletal muscle was low and
close to assay detection limits (Supplementary Fig. 7A). This suggests that system $x_c$ has only
a minor role in the regulation of ferroptosis in skeletal muscle. There were no differences in
GPX activity between groups (Supplementary Fig. 7B). However, expression of DN HDAC4
and 5 increased ROS (Fig. 7J) and increased *Sat1* and *Alox15* expression (Fig. 7K), indicating
that increased capacity for lipid peroxidation. Increased apoptosis and ferroptosis in skeletal
muscles expressing DN HDAC4 and 5 was associated with reduced mass of the
gastrocnemius and the TA muscles (Fig. 7L). In the TA muscle, there was no difference in
fibre diameter between groups (Supplementary Fig. 7C). However, Masson's trichrome
staining showed sporadic areas of localised fibrosis in response to DN HDAC4 and 5
overexpression (Fig. 7M). In skeletal muscles overexpressing DN HDAC4 and 5 there were
also discreet regions of fibres with centralised nuclei (Fig. 7N), which is indicative of

regenerating muscle fibres. These data show that HDAC4 and 5 are required to suppress
apoptosis and ferroptosis and to maintain muscle mass in skeletal muscle of obese mice.

**DISCUSSION**

Lipotoxicity has a major impact on the metabolism, function and survival of various tissues including skeletal muscle. The current study has revealed that increases in HDAC4 and 5 in response to lipotoxicity induce transcriptional repression of oxidative metabolism and of the apoptosis and ferroptosis cell death pathways in both myogenic cells and skeletal muscle (Fig. 8A and B). The impairment in oxidative metabolism is directly linked to reduced p53 acetylation at K120 and inhibition of a pro-apoptosis transcriptional program. Furthermore, metabolic reprogramming to a lower oxidative state also impacted multiple redox inputs to ferroptosis, which inhibited this pathway. These data indicate that metabolic reprogramming in skeletal muscle in response to lipotoxicity is directly linked with cell survival responses.

Previous findings revealed that lipotoxicity is associated with suppression of a pro-apoptosis transcriptional program in skeletal muscle, with consequences for maintenance of muscle mass[46]. Our data show that the class IIa HDACs are essential for this response, as loss of HDAC4 and 5 function in skeletal muscle in a model of lipotoxicity increased apoptosis genes and markers of apoptosis, which was associated with reduced muscle mass. Furthermore, a readout of ferroptosis was also increased. The exact contribution of these cell death pathways to the reduction in muscle mass observed remains to be determined. However, identification that p53 acetylation at K120 is sensitive to substrate supply, metabolic flux and acetyl-CoA availability is an important finding linking metabolism to cell survival responses and has implications for several metabolic diseases, including cancer. As acetyl-CoA operates at the crossroads of catabolic and anabolic metabolism, it seems rational to hypothesise that this metabolite is involved in linking metabolic sensing to cell signalling and adaptive responses. These findings also provide insights into why some substrates are more cytotoxic than others when in excess. It is well established that saturated fatty acids, such as palmitate, are more effective at inducing cell death than unsaturated fatty acids such as oleate[4]. Palmitate is more readily oxidised than oleate[4], thus increasing the acetyl-CoA pool, which would place pressure on K120 p53 acetylation and apoptosis responses. This concept was supported by our tracer data showing much lower overall carbon enrichment into p53 from glucose when compared with that derived from palmitate. Indeed, there are many fates for the carbon derived from glucose in comparison to that derived from fatty acids, which must form mitochondrial acetyl-CoA as a process of β-oxidation.

While there is much debate in the literature about the relationship between mitochondrial capacity and function and insulin action[10], the reduction in oxidative capacity induced by HDAC4 and 5 did not result in overt insulin resistance. Overexpression of HDAC4 and 5 increased basal glucose clearance, with more glucose directed to glycogen synthesis. Oxidative utilisation of glucose was impaired through the TCA cycle, particularly at the level of succinate, which was consistent with decreased succinate-driven respiration. The levels of succinate are thought to reflect the redox state of the mitochondrial coenzyme Q pool, with increased reduced CoQ ($CoQH_2$) preventing the oxidation of succinate by SDH[47]. Impairments in electron transport chain complexes, such as that observed in the current study, are therefore linked to inhibition of succinate oxidation[48]. Furthermore, oxidation of succinate by SDH is tightly associated with reverse electron flow to complex I of the electron transport chain and ROS production[49]. Indeed, succinate oxidation-dependent ROS

production is critical for pro-inflammatory cytokine production by macrophages[50], adipose
tissue thermogenesis[51] and the activity of carotid body oxygen sensing $K^+$ channels[52]. In the
context of the present study, HDAC4 and 5-mediated impairments in succinate oxidation
and ROS production likely contribute to inhibition of the ferroptosis pathway and
preservation of cell viability in response to lipotoxicity. Additionally, impairments in TCA
cycle activity have recently been found to promote cytosolic glutathione synthesis that
increases antioxidant capacity[32], which would also inhibit ferroptosis. Together, these data
indicate that inhibition of oxidative substrate utilisation is an important adaptation to
lipotoxicity.
453
In conclusion, this study provides evidence that the class IIa HDACs link impairments in
skeletal muscle oxidative metabolism in response to lipotoxicity with inhibition of the
apoptosis and ferroptosis cell death pathways to preserve cell viability. This study has
identified new metabolic inputs to transcriptional control of the apoptosis pathway and
show that decreasing the overall oxidative state of skeletal muscle is also important for
inhibition of ferroptosis. These findings provide key new insights into why certain metabolic
adaptations occur in response to excess lipids and advance our understanding of the
aetiology of metabolic diseases driven by lipotoxicity and excess nutrient availability.

**MATERIALS AND METHODS**

**Plasmids and adeno-associated viral vectors**

cDNAs for mouse wild type HDAC4 and 5 were subcloned into pAAV-CMV-M and were packaged into serotype 6 adeno-associated viruses (AAV), purified and titred as previously described [53]. AAV6 vectors expressing DN HDAC4 and 5 have been previously described [20]. cDNAs for human wild type HDAC4 and HDAC5 were subcloned into pIRESneo3 and pIREShyg3 vectors, respectively, to facilitate generation of stable cell lines. pcDNA3 Flag p53 (wild-type, human) was purchased from Addgene (plasmid #10838). Mutant p53 plasmids (K120R and K120Q) were generated using the QuikChange XL Site-Directed Mutagenesis Kit (Agilent Technologies).

**Cell culture**

Mouse C2C12 myoblasts (ATCC) expressing pIRES neo3 HDAC4 (wild-type, human) and pIRES hyg3 HDAC5 (wild-type, human) were produced by transfection of plasmids with Lipofectamine LTX and PLUS reagent (Life Technologies) following manufacturer's instructions. Control cells were generated by transfection with empty plasmids. Stable expression was achieved by multiple rounds of antibiotic selection (500ug/mL Hygromycin B and 500ug/mL Geneticin; Invitrogen). Cells were maintained in growth media consisting of MEM alpha (1g/L glucose; Invitrogen), 10% foetal bovine serum (FBS; In Vitro Technologies) and antibiotic-antimycotic (100units/mL penicillin, 100µg/mL streptomycin and 0.25µg/mL Amphotericin B; Life Technologies), with the addition of selection antibiotics, at 37°C in 10% $CO_2$. For insulin-stimulated experiments, cells were treated for 15min with 0nM or 10nM insulin (Humulin R; Eli Lilly) prior to protein collection. Treatment doses of palmitate (Sigma-Aldrich), camptothecin (Sigma), Erastin (Sigma), 1S,3R-RSL 3 (Sigma), Ferrostatin (Selleckchem) and 1,2,3-benzenetricarboxylate (BTC; Sigma) are reported in Figure legends and were applied for 24h unless otherwise stated. Palmitate was dissolved in ethanol at 70°C and diluted in ultrapure $H_2O$ before being conjugated to fatty acid-free BSA at 1:10 ratio (w/v). An ethanol and BSA solution without palmitate was prepared at the same time for use as a vehicle. Both palmitate and vehicle solutions were aliquoted and stored at -20°C. Each cell culture experiment was performed at least twice.

**Animal models**

All animal experimentation was approved by the Deakin University Animal Ethics Committee (G27-2012 and G21-2015), which is subject to the Australian Code for the Responsible Conduct of Research. Mice were acquired from the Animal Resources Centre (Perth, WA, Australia) and were group housed with a 12hr light/dark cycle, temperature 21± 3°C, humidity: 30-70%. Animals had *ad libitum* access to standard rodent chow and water. Heterozygous *db/+* and homozygous *db/db* mice were fasted for 5 hr prior to humane killing by cervical dislocation for rapid excision of tissues, which were snap frozen in liquid nitrogen for later analysis. C57Bl/6J mice were administered rAAV6 vectors via intramuscular injections into the anterior and posterior compartments of the hind limbs. Each compartment of the left limb received 2 x $10^{10}$ vector genomes (vg) of empty rAAV6, whilst each compartment of the right limb received 1 x $10^{10}$ vg of WT HDAC4 AAV6 and 1 x $10^{10}$ vg of WT HDAC5 AAV6. For expression of individual HDACs in gene dosage experiments, AAV doses were administered as indicated in relevant Figures. Experiments with C57Bl6 mice were performed 2-3 weeks after AAV administration. *db/db* mice were similarly

administered rAAV6 vectors via intramuscular injections into the anterior and posterior
compartments of the hind limbs. Each compartment of the left limb received $5 \times 10^{10}$ vg of
empty rAAV6, whilst each muscle of the right limb received $2.5 \times 10^{10}$ vg of DN HDAC4 AAV6
and $2.5 \times 10^{10}$ vg of DN HDAC5 AAV6. Approximately 8 weeks after rAAV6 administration,
*db/db* mice were fasted for 4 hours before humane killing by cervical dislocation. Tissues
were rapidly excised and snap frozen in liquid nitrogen or fixed in 10% formalin.
515
**Respiration and mitochondrial function analysis**
Cells were treated with vehicle (BSA) or palmitate in 75cm$^2$ flasks for 24 hrs and transferred
to a 24-well XF24 microplate (Seahorse Bioscience) at 50,000 cells/well, with analysis
performed 3 hrs later. Mitochondrial function was assessed as previously described[54] and
respiratory analysis of skeletal muscle biopsies was also performed as previously
described[20].
522
**Lipidomics, metabolomics and acetyl-CoA analysis**
To determine cellular lipid profiles, ~$1 \times 10^6$ cells were collected in 100µl of PBS and frozen at
-80°C, or ~10mg of muscle was homogenised in 100µl of PBS, and lipids were determined as
previously described[55]. Briefly, lipids were extracted using 20 volumes of
chloroform:methanol (2:1) in a single phase extraction process, recovering all lipids in a
single phase suitable for liquid chromatography–mass spectrometry analysis. Lipids were
identified using a HP 1200 liquid chromatography system (Agilent Technologies) coupled to
a PE Sciex API 4000 Q/TRAP mass spectrometer (Applied Biosystems/MDS SCIEX, Mulgrave,
VIC, Australia) with a turbo-ionspray source (350°C) and Analyst 1.5 data system (Applied
Biosystems/MDS SCIEX). Quantification of individual lipid species was performed using
multiple-reaction monitoring (MRM) in positive ion mode. MRM product ions used were
m/z 264 for ceramides, while diacylglycerides and triacylglycerides were monitored by the
neutral loss of individual fatty acid species. Each ion pair was monitored for between 10 and
50 ms (using scheduled MRM mode), with a resolution of 0.7 amu at half-peak height, and
averaged from continuous scans over the elution period. The proportionately higher signals
resulting from diacylglycerol and triacylglycerol consisting of two or more identical fatty
acids were corrected prior to normalisation against internal standards. Lipid concentrations
were calculated by relating the peak area of each species to the peak area of the
corresponding internal standards (100 pmol ceramide 17:0, 200 pmol diacylglycerol
17:0/17:0 and 100 pmol triacylglycerol 17:0/17:0/17:0), which were added prior to
extraction.
544
For metabolite and acetyl-CoA analysis, cells in 6-well plates underwent metabolic arrest by
washing in 1mL ultrapure water before being frozen instantaneously by the addition of
liquid nitrogen. To extract polar metabolites, 600µl of ice-cold 9:1 methanol:chloroform
containing internal standards (0.5nM 13C-sorbitol and 5nM 13C, 15N-valine) was added per
well. Following collection, samples were centrifuged at 16,000 x g for 5min at 4°C and the
supernatant stored at -80°C. Polar metabolites in cellular extracts, including acetyl-CoA,
were assessed using LC and high resolution QTOF mass spectrometry previously described [56]
with the following modifications. Samples (10µL) were injected onto an Agilent 1290 LC
fitted with a SeQuant ZIC®-pHILIC column (2.1 x 150 mm, 5µm; Merck) using 20mM
ammonium carbonate, pH 9.0 (Sigma-Aldrich) (A) and 100% acetonitrile (B) as the mobile
phases. The solvent flow rate of 250 µl/min and a 14 min gradient starting from 90% B to

40% B over 12 minutes, held for 2 min followed by washing at 5% B for 3 minutes and re-
equilibration at 90% B, was used.

**Microarray and gene expression analysis**
For cell culture studies, RNA was extracted from ~1 x $10^6$ cells. For comparative
transcriptomics analysis, RNA was also extracted from ~10mg of muscle from male
heterozygous db/+ and homozygous *db/db* mice. Each treatment group consisted of three
biological replicates and microarray analysis was performed as previously described[20]. For
comparative transcriptomics analyses between cell culture and mouse data, genes found to
be significantly altered in both data sets in the same direction (n = 236) were analysed by
gene set enrichment analysis from all KEGG pathway lists. The microarray datasets
generated conforms to MIAME guidelines and is available at Gene Expression Omnibus
(http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi) series record GSE215185. Gene
expression was measured using real time RT-PCR as previously described [27]. Primer
sequences are available in Table S1.

**Protein fractionation, immunoprecipitation and western blot analysis**
Cells were collected in lysis buffer (50mM Tris pH 7.5, 1mM EDTA, 1mM EGTA, 10% glycerol,
1% Triton X-100, 50mM NaF, 5mM sodium pyrophosphate, 1mM DTT, 1X protease inhibitor
cocktail (Sigma)) and ~20mg skeletal muscle was homogenised in 10v volume of lysis buffer
using a handheld homogeniser. Protein concentration was determined using the BCA assay
(Pierce). Fractionation procedures, immunoprecipitation and western blotting were
performed as previously described[57]. Antibody details are listed in Table S2.

**Stable isotope metabolomics and insulin and isotopic glucose administration**
To assess insulin action, mice were administered 200uCi/kg body weight of 2[1,2-$^3$H]-
deoxyglucose and 100uCi/kg body weight of 1-$^{14}$C glucose, with or without 1.2U/kg
bodyweight of insulin after a 5hr fast. Blood glucose was determined using a handheld
glucometer (Accu-Chek Performa, Roche) and ~5µL blood was obtained from the tail tip
prior to tracer administration and 5, 15 and 30min after tracer administration. Collected
blood was mixed with 75ul of a 2.5% zinc sulfate solution in 1.5ml tubes before being stored
on ice. Mice were humanely killed by cervical dislocation 30 min after tracer administration
and tissues were collected. At the conclusion of the test 25µl of saturated 0.3N barium
hydroxide was added to the blood samples. The samples where then mixed by pipetting
before centrifugation at 1000 x g for 5 min at $4^o$C to pellet any precipitated protein. After
centrifugation 25µl of supernatant was transferred to scintillation vials containing 2ml of
Ultima Gold scintillation liquid (Perkin Elmer) before $^3$H and$^{14}$C counts were measured on a
beta counter (Tri-Carb 2910TR, Perkin Elmer) to determine blood tracer concentration. The
tibialis anterior (TA) skeletal muscle (15-25mg) was used to determine 2-deoxyglucose
clearance as we have previously described[58]. To determine 1-$^{14}$C glucose incorporation into
glycogen, ~10-15mg of TA muscle was digested in 1M KOH at $70^o$C for 20 min and glycogen
was precipitated with saturated $Na_2SO_4$, washed twice with 95% ethanol and resuspended
in acetate buffer (0.84% sodium acetate, 0.46% acetic acid, pH 4.75) containing 0.3mg/mL
amyloglucosidase (Sigma). Glycogen was allowed to digest overnight at $37^o$C before being
assayed for glucose content using the glucose oxidase method[54]. Digested glycogen was also
assessed for $^{14}$C-glucose incorporation by scintillation counting. To determine 1-$^{14}$C glucose
incorporation into lipids, 5-10mg of TA muscle was homogenised in chloroform/methanol
(2:1) and mixed overnight at room temperature. Organic and inorganic phases were

separated by addition of 0.6% NaCl and the lower organic phase was collected and
evaporated under $N_2$ at 45$^o$C. The dried extract was resuspended in absolute ethanol and TG
content was assayed using TG GPO-PAP reagent (Roche) and $^{14}$C-glucose incorporation by
scintillation counting.
607
To assess oxidative glucose utilisation, mice were administered a bolus of 50mg of [U-$^{13}$C]
glucose (Sigma-Aldrich) via oral gavage. At 60 min after [U-$^{13}$C] glucose administration, mice
were humanely killed by cervical dislocation and skeletal muscles were immediately
collected. Targeted metabolomic analysis with [U-$^{13}$C] glucose was performed as we have
previously described[58]. Briefly, approximately 25-30mg of mixed gastrocnemius muscle
tissue was homogenised in 1.2mL of 3:1 methanol:dH$_2$O (-15$^o$C) using a bead cryomill
(Precellys). Extracted metabolites were derivatised before injection into an gas
chromatography system (Agilent 6890N) connected to a VF-5ms capillary column with a 10-
m inert EZ-guard (30m, 0.25mm, 0.25μm) and mass selective detector (Agilent 5975C) with
helium as the carrier gas. Electron ionisation mode was used for the measurement of all
organic and amino acids in their TBDMS derivatized form via analysis of their pseudo-
molecular fragment ions (i.e. M-57). All metabolites were analysed in the selected ion
monitoring mode. Ion abundances, measured as the area under the chromatogram curve,
were determined using the Mass Hunter Workstation (Agilent Technologies, Mulgrave,
Australia). Total metabolite labelling in HDAC4 and 5 over expressing muscle was expressed
relative to labelling in the control muscle within each mouse.
624
**Measurement of apoptosis and cell viability**.
Analysis of apoptotic cells was performed by dual staining with FITC-conjugated Annexin V
(BD Pharmingen) and propidium iodide (PI; Molecular Probes) and was optimised for
adherent cells. Cells in 12-well plates were rinsed with PBS containing 2mM EDTA and
removed from the culture plate using 0.025% trypsin in PBS/2mM EDTA. Trypsin was
inactivated using growth media containing an extra 10% FBS and centrifuged at 500g for
5min. Cells were resuspended and incubated in growth media for ~1h. Staining was
performed in growth media using 5.0μg/mL PI and FITC-conjugated Annexin V according to
the manufacturer's protocol and incubated for 15min in the dark. Annexin binding buffer
(250μl, 10mM HEPES pH 7.4, 140mM NaCl, 2.5mM CaCl$_2$) was added and cells were
immediately analysed by flow cytometry to determine the percentage of viable (Annexin V$^-$
/PI$^-$) and early apoptotic (Annexin V$^+$/PI$^-$) cells. For measurement of apoptosis in tissues,
caspase 3 activity was measured using Caspase-Glo 3/7 assay (Promega,) from
approximately 10mg of EDL tissue from *db/db* mice administered rAAVs.
639
Detection of absolute non-viable cells was performed by staining with 7-aminoactinomycin
D (7-AAD; BD Pharmingen) and flow cytometry, with crystal violet staining and colorimetric
detection, or by LDH release depending on the model used. For 7AAD staining, cells in a 12-
well plate were dislodged with 0.125% trypsin in phosphate buffered saline (PBS) containing
5mM EDTA, centrifuged at 500g for 5min and resuspended in 100μl PBS with 0.2% bovine
serum albumin (BSA). Staining was performed by adding 10μl 7-AAD and incubating in the
dark for 10min. Cells were then rinsed with PBS and fixed in 1% paraformaldehyde
supplemented with 2% FBS and 2μg/ml actinomycin D before being analysed by flow
cytometry using the FACS Calibur® system. The percentage of viable (7-AAD$^-$) and
apoptotic/necrotic (7-AAD$^+$ non-viable) cells was determined using BD CellQuest™ Pro

software (BD Biosciences). For detection of relative non-viable cells, crystal violet staining was performed as we have previously described[59] and cell viability was represented relative to vehicle treated cells. For myotube experiments, cell viability was assessed by LDH release and represented relative to total cellular LDH, as we have previously described[54].

**[14]C-incorporation from palmitate and glucose into p53.**
Cells in a 6-well culture plate were washed twice with PBS and incubated with D-[14C]Glucose (1.14µCi/well; PerkinElmer) or [14C]Palmitic Acid (0.5µCi/well; PerkinElmer) in growth media for 3 hr. Immunoprecipitation of FLAG-tagged p53 was performed with anti-FLAG tag antibody (Cell Signaling). Beads were washed multiple times with ice cold PBS and half of the beads were analysed by scintillation counting. The remaining beads were resuspended in Lane Marker Reducing Sample Buffer (Pierce) and proteins analysed by western blotting. Scintillation counts were normalised to the p53 protein in each immunoprecipitation.

**Reactive oxygen species detection**
In cells, hydrogen peroxide ($H_2O_2$) was detected using Amplex Red assay (Invitrogen). Briefly, cells were seeded into a 96-well plate at a density of 25,000 cells per well. The following day, cells were treated with vehicle (DMSO), 100µM mitoTEMPO or 100µM apocynin for 30 min at 37°C. Cells were then co-incubated with 50µM Amplex Red and 0.1U/ml horseradish peroxidase (HRP) for a further 30 min at 37°C. Production of the fluorescent product resorufin was detected on a Flexstation II[384] fluorometer (Molecular Devices; excitation 544nm, emission 590nm). Mitochondrial ROS was designated as that sensitive to MitoTEMPO, while NADPH oxidase ROS was designated as that sensitive to apocynin. For tissues, the OxiSelect In Vitro ROS/RNS Assay kit (Green Fluorescence) (Cell Biolabs Inc.) was used according to the manufacturer's instructions, using lysates from homogenised extensor digitorum longus (EDL) muscle from *db/db* mice. The assay was performed in a black 96-well plate with DCF standards and was read on a fluorometer (Flexstation II[384] Molecular Devices; excitation 480nm, emission 530nm).

**GPX4 activity and lipid peroxidation**
To measure glutathione dependent peroxidase activity within cultured cells and skeletal muscle, a Glutathione Peroxidase Assay Kit (colorimetric) (Ab Cam Australia) was used. A modified method of the standard protocol was used in both cells and tissue. Cells were plated into 10cm tissue culture plates at $4 \times 10^6$ cells per dish. After 3hrs the growth media was refreshed and cells incubated for 1hr, washed and collected in 250ul of the supplied assay buffer. For tissues, lysate from homogenised gastrocnemius muscle from *db/db* mice was analysed. The assay was performed according to manufacturer's instructions, including generation of the NADPH standards. The reaction was incubated at room temp for 15min. The plate was then read on a plate reader at 340nm (Xmark, Bio-Rad) to ensure all samples had an OD above 1 but below the OD of the highest standard. Next 10ul of cumene hydroperoxide was added to each of the sample and reagent control wells, mixed and the $T_1$ reading taken at 340nm. Following this the plate was incubated in the plate reader at 25°C for 5mins then the $T_2$ reading was taken again at 340nm. Data was then analyzed as described by the manufacturer with all results being normalized to protein concentration.

696    Lipid peroxidation in both cells and muscle lysates was assessed using a lipid peroxidation
697    kit (Sigma). Approximately $1 \times 10^6$ cells were collected in MDA lysis buffer while lysate from
698    homogenised gastrocnemius muscle from *db/db* mice was analysed. The assay was
699    performed according to manufacturer's instructions, including addition of butanol to the
700    reaction.
701
**Histology**
Fixed TA muscles from db/db mice were processed by University of Melbourne Histology.
Tissues were embedded in paraffin before transverse sections were obtained at 5µm
thickness and stained for H&E and Masson's trichrome. Slides were imaged with a slide
scanner (Aperio, Leica Biosystems). Images were analysed using photoshop and image J to
quantify fibre diameter and cell number.
708
**Statistical analysis.**
All data are expressed as mean ± SEM. Individual data points identified as greater than two
standard deviations away from the mean were designated as outliers and removed from the
dataset. Data was assessed for normality using the Shapiro-Wilk test. Differences between
groups were assessed using t-test, Kruskal-Wallis test, one-way ANOVA or two-way ANOVA
as appropriate using GraphPad Prism. Specific differences between groups were identified
using Dunn's or Tukey's multiple comparisons tests. For animal experiments where
opposing hind limbs were administered different rAAVs, such that control and experimental
conditions were contained within the one animal, paired t-tests were used. Differences
were considered statistically significant where $p < 0.05$.

**ACKNOWLEDGEMENTS**

The authors wish to thank all members of the Metabolic Reprogramming Laboratory for helpful insights and technical advice and are grateful to the technical staff of the Deakin University Animal Services team. The authors also wish to thank Hongwei Qian for AAV production. This work was supported by grants from the National Health and Medical Research Council (NHMRC: APP1027227, APP1030474 and APP1127059) to SLM. PJM, PG and MAF are supported by Fellowships from the NHMRC.

**AUTHOR CONTRIBUTIONS**

Conceptualization, SDM, TC and SLM; Methodology, SDM, TC, DDS, GMK, CRB, PG, FMC, KRW and SLM; Investigation, SDM, TC, AS, KAM, DCH, BN, DDS, GMK, FMC and SLM; Resources, DLT, PJM, CRB, PG, MAF and SLM; Writing – original draft, SDM, TC and SLM; Writing – review and editing, all authors; Funding acquisition, SLM.

**DECLARATION OF INTERESTS**

The authors declare no competing interests related to this article.

**SUPPLEMENTARY FILES**
- MDAR checklist
- Supplementary tables file

**DATA AND MATERIALS AVAILABILITY**

All data generated during this study are included in data supplement files. Microarray datasets are available at Gene Expression Omnibus (http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi) series record GSE215185. Any unique or newly created materials will be made available upon request wherever possible.

**REFERENCES**

1.  Lee Y, Hirose H, Ohneda M, Johnson JH, McGarry JD, Unger RH. Beta-cell lipotoxicity in the pathogenesis of non-insulin-dependent diabetes mellitus of obese rats: impairment in adipocyte-beta-cell relationships. *Proceedings of the National Academy of Sciences of the United States of America* **91**, 10878-10882 (1994).

2.  Schaffer JE. Lipotoxicity: when tissues overeat. *Current opinion in lipidology* **14**, 281-287 (2003).

3.  Montgomery MK, De Nardo W, Watt MJ. Impact of Lipotoxicity on Tissue "Cross Talk" and Metabolic Regulation. *Physiology* **34**, 134-149 (2019).

4.  Listenberger LL*, et al.* Triglyceride accumulation protects against fatty acid-induced lipotoxicity. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 3077-3082 (2003).

5.  Kelley DE, Simoneau JA. Impaired free fatty acid utilization by skeletal muscle in non-insulin-dependent diabetes mellitus. *The Journal of clinical investigation* **94**, 2349-2356 (1994).

6.  Simoneau JA, Kelley DE. Altered glycolytic and oxidative capacities of skeletal muscle contribute to insulin resistance in NIDDM. *Journal of applied physiology* **83**, 166-171 (1997).

7.  Baker PR, 2nd*, et al.* Metabolomic analysis reveals altered skeletal muscle amino acid and fatty acid handling in obese humans. *Obesity (Silver Spring)* **23**, 981-988 (2015).

8.  Lerin C*, et al.* Defects in muscle branched-chain amino acid oxidation contribute to impaired lipid metabolism. *Molecular metabolism* **5**, 926-936 (2016).

9.  White PJ*, et al.* Branched-chain amino acid restriction in Zucker-fatty rats improves muscle insulin sensitivity by enhancing efficiency of fatty acid oxidation and acyl-glycine export. *Molecular metabolism* **5**, 538-551 (2016).

10. Muoio DM, Neufer PD. Lipid-induced mitochondrial stress and insulin action in muscle. *Cell metabolism* **15**, 595-605 (2012).

11. Sparks LM*, et al.* A high-fat diet coordinately downregulates genes required for mitochondrial oxidative phosphorylation in skeletal muscle. *Diabetes* **54**, 1926-1933 (2005).

12. Patti ME*, et al.* Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: Potential role of PGC1 and NRF1. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 8466-8471 (2003).

796    13.    Zechner C*, et al.* Total skeletal muscle PGC-1 deficiency uncouples mitochondrial
797           derangements from fiber type determination and insulin sensitivity. *Cell metabolism*
798           **12**, 633-642 (2010).
799

800    14.    Choi CS*, et al.* Paradoxical effects of increased expression of PGC-1alpha on muscle
801           mitochondrial function and insulin-stimulated muscle glucose metabolism.
802           *Proceedings of the National Academy of Sciences of the United States of America*
803           **105**, 19926-19931 (2008).
804

805    15.    Dube JJ*, et al.* Effects of acute lipid overload on skeletal muscle insulin resistance,
806           metabolic flexibility, and mitochondrial performance. *American journal of physiology*
807           *Endocrinology and metabolism* **307**, E1117-1124 (2014).
808

809    16.    Richardson DK*, et al.* Lipid infusion decreases the expression of nuclear encoded
810           mitochondrial genes and increases the expression of extracellular matrix genes in
811           human skeletal muscle. *The Journal of biological chemistry* **280**, 10290-10297 (2005).
812

813    17.    Staiger H, Staiger K, Haas C, Weisser M, Machicao F, Haring HU. Fatty acid-induced
814           differential regulation of the genes encoding peroxisome proliferator-activated
815           receptor-gamma coactivator-1alpha and -1beta in human skeletal muscle cells that
816           have been differentiated in vitro. *Diabetologia* **48**, 2115-2118 (2005).
817

818    18.    Coll T*, et al.* Palmitate-mediated downregulation of peroxisome proliferator-
819           activated receptor-gamma coactivator 1alpha in skeletal muscle cells involves
820           MEK1/2 and nuclear factor-kappaB activation. *Diabetes* **55**, 2779-2787 (2006).
821

822    19.    Crunkhorn S*, et al.* Peroxisome proliferator activator receptor gamma coactivator-1
823           expression is reduced in obesity: potential pathogenic role of saturated fatty acids
824           and p38 mitogen-activated protein kinase activation. *The Journal of biological*
825           *chemistry* **282**, 15439-15450 (2007).
826

827    20.    Gaur V*, et al.* Disruption of the Class IIa HDAC Corepressor Complex Increases Energy
828           Expenditure and Lipid Oxidation. *Cell reports* **16**, 2802-2810 (2016).
829

830    21.    Czubryt MP, McAnally J, Fishman GI, Olson EN. Regulation of peroxisome
831           proliferator-activated receptor gamma coactivator 1 alpha (PGC-1 alpha ) and
832           mitochondrial function by MEF2 and HDAC5. *Proceedings of the National Academy*
833           *of Sciences of the United States of America* **100**, 1711-1716 (2003).
834

835    22.    Backs J, Backs T, Bezprozvannaya S, McKinsey TA, Olson EN. Histone deacetylase 5
836           acquires calcium/calmodulin-dependent kinase II responsiveness by oligomerization
837           with histone deacetylase 4. *Molecular and cellular biology* **28**, 3437-3445 (2008).
838

839    23.    Lemercier C, Verdel A, Galloo B, Curtet S, Brocard MP, Khochbin S. mHDA1/HDAC5
840           histone deacetylase interacts with and represses MEF2A transcriptional activity. *The*
841           *Journal of biological chemistry* **275**, 15594-15599 (2000).
842

843 24. McKinsey TA, Zhang CL, Lu J, Olson EN. Signal-dependent nuclear export of a histone
844   deacetylase regulates muscle differentiation. *Nature* **408**, 106-111 (2000).
845

846 25. McGee SL*, et al.* AMP-activated protein kinase regulates GLUT4 transcription by
847   phosphorylating histone deacetylase 5. *Diabetes* **57**, 860-867 (2008).
848

849 26. Mihaylova MM*, et al.* Class IIa histone deacetylases are hormone-activated
850   regulators of FOXO and mammalian glucose homeostasis. *Cell* **145**, 607-621 (2011).
851

852 27. McGee SL*, et al.* Compensatory regulation of HDAC5 in muscle maintains metabolic
853   adaptive responses and metabolism in response to energetic stress. *FASEB journal :*
854   *official publication of the Federation of American Societies for Experimental Biology*
855   **28**, 3384-3395 (2014).
856

857 28. McGee SL, Fairlie E, Garnham AP, Hargreaves M. Exercise-induced histone
858   modifications in human skeletal muscle. *The Journal of physiology* **587**, 5951-5958
859   (2009).
860

861 29. Petersen KF, Dufour S, Befroy D, Garcia R, Shulman GI. Impaired mitochondrial
862   activity in the insulin-resistant offspring of patients with type 2 diabetes. *The New*
863   *England journal of medicine* **350**, 664-671 (2004).
864

865 30. Kowalski GM*, et al.* Application of dynamic metabolomics to examine in vivo skeletal
866   muscle glucose metabolism in the chronically high-fat fed mouse. *Biochemical and*
867   *biophysical research communications* **462**, 27-32 (2015).
868

869 31. Zhang W, Guo C, Jiang K, Ying M, Hu X. Quantification of lactate from various
870   metabolic pathways and quantification issues of lactate isotopologues and
871   isotopmers. *Sci Rep* **7**, 8489 (2017).
872

873 32. Ryan DG*, et al.* Disruption of the TCA cycle reveals an ATF4-dependent integration of
874   redox and amino acid metabolism. *Elife* **10**,  (2021).
875

876 33. Jiang X, Stockwell BR, Conrad M. Ferroptosis: mechanisms, biology and role in
877   disease. *Nature reviews Molecular cell biology* **22**, 266-282 (2021).
878

879 34. Sykes SM*, et al.* Acetylation of the p53 DNA-binding domain regulates apoptosis
880   induction. *Molecular cell* **24**, 841-851 (2006).
881

882 35. Tang Y, Luo J, Zhang W, Gu W. Tip60-dependent acetylation of p53 modulates the
883   decision between cell-cycle arrest and apoptosis. *Molecular cell* **24**, 827-839 (2006).
884

885 36. Sen N, Kumari R, Singh MI, Das S. HDAC5, a key component in temporal regulation of
886   p53-mediated transactivation in response to genotoxic stress. *Molecular cell* **52**, 406-
887   420 (2013).
888

889    37.    Jiang L*, et al.* Ferroptosis as a p53-mediated activity during tumour suppression.
890           *Nature* **520**, 57-62 (2015).
891

892    38.    Berkers CR, Maddocks OD, Cheung EC, Mor I, Vousden KH. Metabolic regulation by
893           p53 family members. *Cell metabolism* **18**, 617-633 (2013).
894

895    39.    Lahm A*, et al.* Unraveling the hidden catalytic activity of vertebrate class IIa histone
896           deacetylases. *Proceedings of the National Academy of Sciences of the United States*
897           *of America* **104**, 17335-17340 (2007).
898

899    40.    Martinez-Reyes I, Chandel NS. Mitochondrial TCA cycle metabolites control
900           physiology and disease. *Nature communications* **11**, 102 (2020).
901

902    41.    Berwick DC, Hers I, Heesom KJ, Moule SK, Tavare JM. The identification of ATP-
903           citrate lyase as a protein kinase B (Akt) substrate in primary adipocytes. *The Journal*
904           *of biological chemistry* **277**, 33895-33900 (2002).
905

906    42.    Stockwell BR*, et al.* Ferroptosis: A Regulated Cell Death Nexus Linking Metabolism,
907           Redox Biology, and Disease. *Cell* **171**, 273-285 (2017).
908

909    43.    Kagan VE*, et al.* Oxidized arachidonic and adrenic PEs navigate cells to ferroptosis.
910           *Nature chemical biology* **13**, 81-90 (2017).
911

912    44.    Ou Y, Wang SJ, Li D, Chu B, Gu W. Activation of SAT1 engages polyamine metabolism
913           with p53-mediated ferroptotic responses. *Proceedings of the National Academy of*
914           *Sciences of the United States of America* **113**, E6806-E6812 (2016).
915

916    45.    Piccolis M*, et al.* Probing the Global Cellular Responses to Lipotoxicity Caused by
917           Saturated Fatty Acids. *Molecular cell* **74**, 32-44 e38 (2019).
918

919    46.    Turpin SM*, et al.* Examination of 'lipotoxicity' in skeletal muscle of high-fat fed and
920           ob/ob mice. *The Journal of physiology* **587**, 1593-1605 (2009).
921

922    47.    Murphy MP, Chouchani ET. Why succinate? Physiological regulation by a
923           mitochondrial coenzyme Q sentinel. *Nature chemical biology* **18**, 461-469 (2022).
924

925    48.    Burger N*, et al.* A sensitive mass spectrometric assay for mitochondrial CoQ pool
926           redox state in vivo. *Free radical biology & medicine* **147**, 37-47 (2020).
927

928    49.    Chance B, Hollunger G. The interaction of energy and electron transfer reactions in
929           mitochondria. I. General properties and nature of the products of succinate-linked
930           reduction of pyridine nucleotide. *The Journal of biological chemistry* **236**, 1534-1543
931           (1961).
932

933    50.    Mills EL*, et al.* Succinate Dehydrogenase Supports Metabolic Repurposing of
934           Mitochondria to Drive Inflammatory Macrophages. *Cell* **167**, 457-470 e413 (2016).
935

936  51.  Mills EL*, et al.* Accumulation of succinate controls activation of adipose tissue
937       thermogenesis. *Nature* **560**, 102-106 (2018).
938
939  52.  Fernandez-Aguera MC*, et al.* Oxygen Sensing by Arterial Chemoreceptors Depends
940       on Mitochondrial Complex I Signaling. *Cell metabolism* **22**, 825-837 (2015).
941
942  53.  Winbanks CE*, et al.* The bone morphogenetic protein axis is a positive regulator of
943       skeletal muscle mass. *The Journal of cell biology* **203**, 345-357 (2013).
944
945  54.  Martin SD, Morrison S, Konstantopoulos N, McGee SL. Mitochondrial dysfunction has
946       divergent, cell type-dependent effects on insulin action. *Molecular metabolism* **3**,
947       408-418 (2014).
948
949  55.  Genders AJ*, et al.* Reducing hepatic PKD activity lowers circulating VLDL cholesterol. *J
950       Endocrinol* **246**, 265-276 (2020).
951
952  56.  Hortle E*, et al.* Adenosine monophosphate deaminase 3 activation shortens
953       erythrocyte half-life and provides malaria resistance in mice. *Blood* **128**, 1290-1301
954       (2016).
955
956  57.  McGee SL, Hargreaves M. Exercise and myocyte enhancer factor 2 regulation in
957       human skeletal muscle. *Diabetes* **53**, 1208-1214 (2004).
958
959  58.  De Jong KA*, et al.* Loss of protein kinase D activity demonstrates redundancy in
960       cardiac glucose metabolism and preserves cardiac function in obesity. *Molecular
961       metabolism* **42**, 101105 (2020).
962
963  59.  Preston S*, et al.* 1-Methyl-1H-pyrazole-5-carboxamide Derivatives Exhibit
964       Unexpected Acute Mammalian Toxicity. *Journal of medicinal chemistry* **64**, 840-844
965       (2021).

**FIGURE 1**

**Figure 1: HDAC4 and 5 are increased in models of lipotoxicity.** (A) Heat map of ceramide (Cer), diglyceride (DG), phosphatidylcholine (PC) and triglyceride (TG) lipid species; (B) Total Cer, DG and TG lipids (One-way ANOVA; Cer $p<0.0001$ (F(2,14)=73.86, [f]Tukey's multiple comparisons test $p<0.0077$ vs. 0mM, *Tukey's multiple comparisons test $p<0.0001$ vs. 0mM and vs 0.25mM; DG $p<0.0001$ (F(2,14)=38.50 *Tukey's multiple comparisons test $p<0.0001$ vs. 0mM and vs 0.25mM; TG $p<0.0017$ (F(2,14)=10.38 *Tukey's multiple comparisons test $p=0.0333$ vs. 0mM); (C) *Ppargc1a* gene expression; (D) Oxygen consumption rate (OCR; One-way ANOVA $p<0.0255$ (F(2,25)=4.262 *Tukey's multiple comparisons test $p<0.0001$ vs. 0mM and vs 0.25mM), and; (E) HDAC4 and 5 protein in C2C12 myotubes treated with 0mM (BSA vehicle), 0.25mM or 0.5mM palmitate (PA) for 24 hrs (HDAC4, Kruskal-Wallis test, $p=0.049$

($X^2$ = 5.591), *Dunn's multiple comparison test $p$=0.0380; HDAC5, One-way ANOVA, $p$=0.038
(F(2,6)=5.903) $^f$Tukey's multiple comparisons test $p$=0.0318, *Tukey's multiple comparisons
test $p$=0.0251). (F) Heat map of Cer, DG, PC and TG lipid species; (G) Total Cer, DG and TG
lipids (Unpaired t-tests, Cer $p$=0.0094, DG $p$=0.0022, TG $p$=0.0616); (H) *Ppargc1a* gene
expression (Unpaired t-test, $p$=0.0357); (I) Basal, substrate and ADP-driven OCR from muscle
biopsies (Two-way ANOVA, $^\varnothing$genotype $p$=0.0493, $F$(1,33)=4.166), and; (J) HDAC4 and 5
protein in Control and *db/db* mice (Mann-Whitney test, $^\#$HDAC4 $p$=0.0002, Unpaired t-test
$^\#$HDAC5 $p$=0.0308). Data are mean±SEM, n=3-6 biological replicates per group for cell
experiments and 6-8 per group for animal experiments.
994
995
996  **Figure 1 - source data 1:** Source data for Figure 1A
997
998  **Figure 1 - source data 2:** Source data for Figure 1B
999
1000  **Figure 1 - source data 3:** Source data for Figure 1C
1001
1002  **Figure 1 - source data 4:** Source data for Figure 1D
1003
1004  **Figure 1 - source data 5:** Source data and raw blots for Figure 1E
1005
1006  **Figure 1 - source data 6:** Source data for Figure 1F
1007
1008  **Figure 1 - source data 7:** Source data for Figure 1G
1009
1010  **Figure 1 - source data 8:** Source data for Figure 1H
1011
1012  **Figure 1 - source data 9:** Source data for Figure 1I
1013
1014  **Figure 1 - source data 10:** Source data and raw blots for Figure 1J
1015
1016
1017
1018  **Figure 1 Supplement** – Viability and gene expression responses to lipotoxicity.
1019
1020  **Figure 1 supplement – source data 1:** Source data for Figure S1A
1021
1022  **Figure 1 supplement – source data 1:** Source data for Figure S1B
1023
1024  **Figure 1 supplement – source data 1:** Source data for Figure S1C
1025
1026

1029
**Figure 2: Overexpression of HDAC4 and 5 in skeletal muscle suppresses *Pppargc1a***
**expression and oxidative capacity.** (A) The bilateral AAV6 HDAC4 and 5 overexpression
mouse model. (B) *Hdac4* and *Hdac5* gene expression (Paired t-test, #*Hdac4 p*=0.0246,
#*Hdac5 p*=0.0305); (C) HDAC4 and 5 protein; (D) *Pppargc1a* gene expression (Paired t-test,
*p*=0.0006); (E) Metabolic gene expression (Paired t-test, #*Ppard p*=0.0056, #*Cpt1b p*=0.0212,
#*Pdk4 p*=0.0058, #*Cox7a p*=0.0175, #*CytC p*=0.0042, #*Idh1 p*=0.0358, #*Fh p*=0.0030, #*Mcad*
*p*=0.0056, #*Hadh p*=0.0057, #*Echs1 p*=0.0046); (F) Respiratory chain complex subunit protein
(Paired t-test, #Complex V *p*=0.0036); (G) Basal, succinate and ADP-driven oxygen
consumption rate (OCR) from muscle biopsies (Two-way ANOVA, ∅genotype *p*=0.0247,
*F*(1,43)=5.417); (H) Basal, malate and ADP-driven OCR from muscle biopsies (Two-way
ANOVA, ∅genotype *p*=0.0036, *F*(1,37)=9.661), and; (I) Ceramide (Cer), diglyceride (DG),
triglyceride (TG), phosphatidylethanolamine (PE), lysophosphatidylethanolamine (LPE),
phosphatidylcholine (PC), lysophosphatidylcholine (LPC), phosphatidylinositol (PI),
lysophosphatidylinositol (LPI), phosphoglycerol (PG) and phosphatidylserine (PS) lipids in

Control and HDAC4 and 5 overexpressing skeletal muscle. Data are mean±SEM, n=4-6 per
group.
**Figure 2 - source data 1:** Source data for Figure 2B
1047
1048  **Figure 2 - source data 2:** Raw blots for Figure 2C
1049
1050  **Figure 2 - source data 3:** Source data for Figure 2D
1051
1052  **Figure 2 - source data 4:** Source data for Figure 2E
1053
1054  **Figure 2 - source data 5:** Source data and raw blots for Figure 2F
1055
1056  **Figure 2 - source data 6:** Source data for Figure 2G
1057
1058  **Figure 2 - source data 7:** Source data for Figure 2H
1059
1060  **Figure 2 - source data 8:** Source data for Figure 2I
1061
1062
1063
1064  **Figure 2 Supplement** - Individual HDAC isoform and gene dosage effects on *Ppargc1a*
1065  *expression.*
1066
1067  **Figure 2 supplement – source data 1:** Source data for Figure S2B
1068
1069  **Figure 2 supplement – source data 2:** Raw blots for Figure S2C
1070
1071  **Figure 2 supplement – source data 3:** Source data for Figure S2D
1072
1073  **Figure 2 supplement – source data 4:** Source data for Figure S2E
1074

**FIGURE 3**

1076

1077
1078
**Figure 3: Overexpression of HDAC4 and 5 in skeletal muscle inhibits oxidative glucose**
**utilisation.** (A) Experimental overview of insulin and isotopic glucose tracers in the bilateral
AAV6 HDAC4 and 5 overexpression model. (B) Glucose clearance (Two-way ANOVA,
$^{\varnothing}$genotype $p$=0.0328, $F$(1,27)=5.061); (C) Glucose clearance in muscle overexpressing HDAC4

and 5 relative to Control (Paired t-test, [#]Vehicle $p$=0.0355); (D) $^{14}$C-glucose incorporation
into glycogen (Two-way ANOVA, $^{\varnothing}$genotype $p$=0.0372, $F(1,24)$=4.864); (E) $^{14}$C-glucose
incorporation into lipid (Two-way ANOVA, ^Insulin $p$=0.0009, $F(1,21)$=14.94); (F)
Representative images of insulin signalling components; (G) pT308 Akt (Two-way ANOVA,
^Insulin $p$=0.0010, $F(1,27)$=13.75); (H) pS473 Akt (Two-way ANOVA, ^Insulin $p$<0.0001,
$F(1,28)$=33.13), and; (I) pS642 TBC1D4 (Two-way ANOVA, $^{\varnothing}$genotype $p$=0.0011,
$F(1,25)$=13.60) in bilateral AAV6 HDAC4 and 5 mice administered insulin and isotopic
glucose tracers. (J) Experimental overview of stable isotope metabolomics in the bilateral
AAV6 HDAC4 and 5 overexpression model. (K) Total metabolite abundance (Paired t-test,
[#]Lactate $p$=0.0447, [#]Succinate $p$=0.0230); (L) Total $^{13}$C labelling of glycolytic and TCA cycle
metabolites (Paired t-test, [#]Succinate $p$<0.0001, [#]Malate $p$=0.0337); (M) Lactate
isotopologue (Paired t-test, [#]M0 $p$=0.0448, [#]M2 $p$=0.0299); (N) Succinate isotopologues
(Paired t-test, [#]M0 $p$=0.0230, [#]M1 $p$=0.0002, [#]M2 $p$=0.0005), and; (O) Malate isotopologues
(Paired t-test, [#]M1 $p$=0.0178) in bilateral AAV6 HDAC4 and 5 mice administered $^{13}$C- glucose
tracer. (E) Data are mean±SEM, n=5-8 per group.
1098
1099 **Figure 3 - source data 1:** Source data for Figure 3B
1100
1101 **Figure 3 - source data 2:** Source data for Figure 3C
1102
1103 **Figure 3 - source data 3:** Source data for Figure 3D
1104
1105 **Figure 3 - source data 4:** Source data for Figure 3E
1106
1107 **Figure 3 - source data 5:** Raw blots for Figure 3F
1108
1109 **Figure 3 - source data 6:** Source data for Figure 3G
1110
1111 **Figure 3 - source data 7:** Source data for Figure 3H
1112
1113 **Figure 3 - source data 8:** Source data for Figure 3I
1114
1115 **Figure 3 - source data 9:** Source data for Figure 3K
1116
1117 **Figure 3 - source data 10:** Source data and raw blots for Figure 3L
1118
1119 **Figure 3 - source data 11:** Source data for Figure 3M
1120
1121 **Figure 3 - source data 12:** Source data for Figure 3N
1122
1123 **Figure 3 - source data 13:** Source data and raw blots for Figure 3O
1124
1125
1126 **Figure 3 Supplement** – Plasma tracer and muscle glycogen concentrations.
1127
1128 **Figure 3 supplement – source data 1:** Source data for Figure S3A
1129

1130    **Figure 3 supplement – source data 2:** Source data for Figure S3B
1131
1132    **Figure 3 supplement – source data 3:** Source data for Figure S3C
1133
1134
1135
1136

**FIGURE 4**

**Figure 4: HDAC and 5 enhance cell survival in response to lipotoxicity and supress the apoptosis and ferroptosis cell death pathways.** (A) Expression of p53-dependent genes involved in pro-apoptosis, anti-apoptosis, intrinsic apoptosis and ferroptosis (Unpaired t-test, #*Casp9* p=0.0258, #*Cycs* p=0.0049, #*Pmaip1* p=0.0010, #*Bbc3* p=0.0476, #*Bcl2* p=0.0037, #*Bcl2l1* p=0.0005, #*Slc7a11* p<0.0001, #*Sat1* p<0.0001, #*Alox15* p=0.0041); (B) Early apoptotic cells in response to the apoptosis-inducing agent Camptothecin (Two-way ANOVA, genotype p<0.0001 F(1,12)=349.3, treatment p<0.0001 F(2,12)=270.4, interaction p<0.0001 F(2,12)=149.5; #p<0.05 vs Control for same treatment, *p<0.05 vs vehicle by Tukey's multiple comparisons tests); (C) Cell viability in response to the ferroptosis-inducing agent RSL3 (Two-way ANOVA, genotype p<0.0001 F(1,12)=1645, treatment p<0.0001 F(1,12)=36.09, interaction p<0.0090 F(1,12)=9.665; #p<0.05 vs Control for same treatment, *p<0.05 vs 0.25µM RSL by Tukey's multiple comparisons tests); (D) Cell viability in response to palmitate (Two-way ANOVA, genotype p<0.0001 F(1,12)=3374, treatment p<0.0001 F(2,12)=1012, interaction p<0.0001 F(2,12)=684.8; #p<0.05 vs Control for same treatment, *p<0.05 vs vehicle by Tukey's multiple comparisons tests); (E) Full-length (FL) and cleaved (CL) caspase 3 and 9 in response to palmitate; (F) Early apoptotic cells in response to

palmitate (Two-way ANOVA, genotype $p<0.0001$ $F(1,12)=1468$, treatment $p=0.0001$
$F(2,12)=20.35$, interaction $p<0.0050$ $F(2,12)=8.498$; [#]p<0.05 vs Control for same treatment,
*p<0.05 vs vehicle by Tukey's multiple comparisons tests); (G) Late apoptotic cells in
response to palmitate (Two-way ANOVA, genotype $p<0.0001$ $F(1,12)=313.4$, treatment
$p<0.0001$ $F(2,12)=67.31$, interaction $p=0.0001$ $F(2,12)=21.00$; [#]p<0.05 vs Control for same
treatment, *p<0.05 vs vehicle by Tukey's multiple comparisons tests), and (H) Lipid
peroxidation in response to palmitate (Two-way ANOVA, genotype $p<0.0001$ $F(1,12)=313.4$,
treatment $p<0.0103$ $F(2,12)=67.31$; [#]p<0.05 vs Control for same treatment by Tukey's
multiple comparisons tests) in Control and HDAC4 and 5 overexpressing C2C12 myoblasts.
Data are mean±SEM, n=3-4 biological replicates per group.
1167
1168 **Figure 4 - source data 1:** Source data for Figure 4A
1169
1170 **Figure 4 - source data 2:** Source data for Figure 4B
1171
1172 **Figure 4 - source data 3:** Source data for Figure 4C
1173
1174 **Figure 4 - source data 4:** Source data for Figure 4D
1175
1176 **Figure 4 - source data 5:** Raw blots for Figure 4E
1177
1178 **Figure 4 - source data 6:** Source data for Figure 4F
1179
1180 **Figure 4 - source data 7:** Source data for Figure 4G
1181
1182 **Figure 4 - source data 8:** Source data for Figure 4H
1183
1184
1185
1186 **Figure 4 Supplement** - Characterisation of HDAC4&5 overexpression in C2C12 myoblasts.
1187
1188 **Figure 4 supplement – source data 1:** Raw blots for Figure S4A
1189
1190 **Figure 4 supplement – source data 2:** Source data for Figure S4B
1191
1192 **Figure 4 supplement – source data 3:** Source data for Figure S4C
1193
1194 **Figure 4 supplement – source data 4:** Source data for Figure S4D
1195

	**FIGURE 5**

1197
1198

**Figure 5: HDAC4 and 5 suppress apoptosis by reducing p53 K120 acetylation, which is**
**regulated by HDAC4 and 5-mediated alterations in metabolism.** (A) Acetylated p53 at
lysine 120 (aK120 p53); (B) Expression of caspase 3 (*Casp3;* One-way ANOVA $p$=0.0040
$X^2$=0.8408, ^$p$<0.05 vs Control + WT p53 and HDAC4&5 + K120Q p53 by Tukey's multiple
comparisons test), caspase 9 (*Casp9;* One-way ANOVA $p$=0.0025 $X^2$=0.8650, ^$p$<0.05 vs
Control + WT p53 and HDAC4&5 + K120Q p53 by Tukey's multiple comparisons test),
cytochrome c (*Cycs;* One-way ANOVA $p$=0.0033 $X^2$=0.8518, ^$p$<0.05 vs Control + WT p53 by
Tukey's multiple comparisons test) and the cystine/glutamate antiporter xCT (*Slc7a11;* One-
way ANOVA $p$<0.0001 $X^2$=0.9769, ^$p$<0.05 vs Control + WT p53 by Tukey's multiple
comparisons test); (C) Percentage of early apoptotic cells following exposure to
camptothecin (One-way ANOVA $p$=0.0008 $X^2$=0.9065, ^$p$<0.05 vs Control + WT p53 and
HDAC4&5 + K120Q p53 by Tukey's multiple comparisons test); (D) $^{14}$C incorporation into p53
from palmitate (One-way ANOVA, $p$=0.0115 $X^2$=0.5602, ^p<0.05 vs Control + WT p53 by
Tukey's multiple comparisons tests) or glucose; (E) aK120 p53 following expose to vehicle or
insulin for 15 min following 24 hrs in the presence or absence of BTC; (F) Acetyl-CoA levels
(Unpaired t-test, $p$=0.0014), and; (G) Cell viability 6, 8, 12 and 16 hrs after palmitate
exposure (Two-way ANOVA, time $p$=0.0003 $F$(1,6)=52.34; $^*$p<0.05 vs 6hrs and 8hrs for same
genotype by Tukey's multiple comparisons tests) in Control or HDAC4 and 5 overexpressing
C2C12 myoblasts also expressing wild type (WT), K120R (loss-of-function) and K120Q (gain-
of-function) p53. Data are mean±SEM, n=3-6 biological replicates per group.

**Figure 5 - source data 1:** Raw blots for Figure 5A

1220

1221    **Figure 5 - source data 2:** Source data for Figure 5B

1222

1223    **Figure 5 - source data 3:** Source data for Figure 5C

1224

1225    **Figure 5 - source data 4:** Source data for Figure 5D

1226

1227    **Figure 5 - source data 5:** Raw blots for Figure 5E

1228

1229    **Figure 5 - source data 6:** Source data for Figure 5F

1230

1231    **Figure 5 - source data 7:** Source data for Figure 5G

1232

1233

1234

1235    **Figure 5 Supplement** – Bioenergetics in response to wild type and mutant p53
1236    overexpression.

1237

1238    **Figure 5 supplement – source data 1:** Source data for Figure S5A

1239

1240    **Figure 5 supplement – source data 2:** Source data for Figure S5B

1241

    **FIGURE 6**

1243

1244

1245
1246

**Figure 6: HDAC4 and 5 regulate multiple metabolic inputs to the ferroptosis pathway.** (A)
Schematic showing key regulatory points in the ferroptosis pathway. (B) Cell viability
following palmitate exposure and co-incubation with vehicle (DMSO) or increasing
concentrations of Ferrostatin (One-way ANOVA $p$=0.0323 $X^2$=0.5337, *$p$<0.05 vs Vehicle by
Tukey's multiple comparisons test); (C) Cell viability following exposure to increasing
concentrations of Erastin (Two-way ANOVA, genotype $p$<0.0001 $F(1,18)$=293.7, treatment
$p$<0.0001 $F(2,18)$=139.6, interaction $p$<0.0001 $F(2,18)$=50.88; #$p$<0.05 vs Control for same
treatment, *p<0.05 vs Vehicle same genotype by Tukey's multiple comparisons tests); (D)
Glutamate (Glu), glutathione (GSH) and the GSH/Glu ratio (Unpaired t-tests, #Glu $p$=0.0049,
#GSH/Glu $p$=0.0023); (E) Glutathione peroxidase (GPX) activity (Unpaired t-test, $p$<0.0001),
and; (F) $H_2O_2$ derived from mitochondria (mito), NADPH oxidases (NOX) and other sources
(Unpaired t-tests, #Mito $p$=0.0105, #NAPDH Ox. $p$<0.0001, #Other $p$=0.0004) in Control or
HDAC4 and 5 overexpressing C2C12 myoblasts. Data are mean±SEM, n=4-10 biological
replicates per group.

1261

1262    **Figure 6 - source data 1:** Source data for Figure 6B

1263

1264    **Figure 6 - source data 2:** Source data for Figure 6C

1265

1266    **Figure 6 - source data 3:** Source data for Figure 6D

1267

1268    **Figure 6 - source data 4:** Source data for Figure 6E

1269

1270    **Figure 6 - source data 5:** Source data for Figure 6F

1271

1272

1273  **Figure 6 Supplement** – ROS and phospholipid responses to HDAC4&5 overexpression.

1274

1275  **Figure 6 supplement – source data 1:** Source data for Figure S6A

1276

1277  **Figure 6 supplement – source data 2:** Source data for Figure S6B

1278

1279  **Figure 6 supplement – source data 3:** Source data for Figure S6C

1280

1281  **Figure 6 supplement – source data 4:** Source data for Figure S6D

1282

1283
1284 **FIGURE 7**
1285

1286
1287
1288 **Figure 7: HDAC4 and 5 are required to suppress apoptosis and ferroptosis and maintain**
1289 **muscle mass in obese mice.** (A) The bilateral AAV6 dominant negative (DN) HDAC4 and 5
1290 overexpression *db/db* mouse model. (B) *Hdac4* and *Hdac5* gene expression (Paired t-tests,
1291 [#]*Hdac4 p*=0.0004, [#]*Hdac5 p*=0.0002); (C) HDAC4 and 5 protein; (D) *Ppargc1a* gene

expression (Paired t-tests, *p*=0.0191); (E) Basal, substrate and ADP-driven oxygen consumption rate (OCR) from muscle biopsies (Two-way ANOVA, genotype *p*=0.0482 *F*(1,42)=4.141); (F) Caspase 3 activity (Paired t-tests, *p*=0.0292); (G) Lipid peroxidation (Paired t-tests, *p*=0.0013); (H) Acetylated p53 at lysine 120 (aK120 p53); (I) Expression of apoptosis genes (Paired t-tests, [#]*Casp3 p*=0.0072, [#]*Casp9 p*=0.0129, [#]*Bbc3 p*=0.0049); (J) Reactive oxygen species (ROS; Paired t-tests, *p*=0.0399); (K) Expression of ferroptosis genes (Paired t-tests, [#]*Sat1 p*=0.0258, [#]*Casp9 p*=0.0286); (L) Mass of the gastrocnemius (Gastroc.) and tibialis anterior (TA) skeletal muscles (Paired t-tests, [#]*Gastroc. P*<0.0001, [#]*TA p*=0.0004); (M) Masson's trichrome staining, and; (N) H&E staining of sections of the TA muscle in Control and DN HDAC4 and 5 overexpressing skeletal muscle in *db/db* mice. Arrows indicate fibres with centralised nuclei. Data are mean±SEM, n=6-10 per group.

**Figure 7 - source data 1:** Source data for Figure 7B

**Figure 7 - source data 2:** Raw blots for Figure 7C

**Figure 7 - source data 3:** Source data for Figure 7D

**Figure 7 - source data 4:** Source data for Figure 7E

**Figure 7 - source data 5:** Source data for Figure 7F

**Figure 7 - source data 6:** Source data for Figure 7G

**Figure 7 - source data 7:** Source data and raw blots for Figure 7H

**Figure 7 - source data 8:** Source data for Figure 7I

**Figure 7 - source data 9:** Source data for Figure 7J

**Figure 7 - source data 10:** Source data and raw blots for Figure 7K

**Figure 7 - source data 11:** Source data for Figure 7L

**Figure 7 Supplement** – *Slc7a11* expression, GPX activity and fibre diameter in response to DN HDAC4&5 expression in obese *db/db* mice.

**Figure 7 supplement – source data 1:** Source data for Figure S7A

**Figure 7 supplement – source data 2:** Source data for Figure S7B

**Figure 7 supplement – source data 3:** Source data for Figure S7C

1339  **FIGURE 8**

1340
1341

**Figure 8: Proposed model.** (A) Lipotoxicity, through mitochondrial metabolism, increases
acetylation of p53 and a transcriptional program that increases apoptosis. Lipotoxicity also
increases reactive oxygen species (ROS) and lipid peroxidation, leading to ferroptosis.
Activation of both apoptosis and ferroptosis increases myocyte death and loss of muscle
mass. (B) Lipotoxicity increases HDAC4 and HDAC5, resulting in repression of a broad
transcriptional program controlling oxidative metabolism and lipid peroxidation. Impaired
mitochondrial metabolism reduces p53 acetylation at K120, which inhibits apoptosis.
Similarly, p53-dependent genes involved in lipid peroxidation are reduced, which together
with reduced ROS, inhibits lipid peroxidation and ferroptosis. Inhibition of both the
apoptosis and ferroptosis cell death pathways preserves myocyte viability and muscle mass
in response to lipotoxicity.
1353

**FIGURE SUPPLEMENT 1**

**Figure Supplement 1: Viability and gene expression responses to lipotoxicity.** (A) Viability of C2C12 myotubes (One-way ANOVA $p<0.0001$ $X^2=0.6927$, *$p<0.05$ vs Vehicle by Tukey's multiple comparisons test), and; (B) *Hdac4* and *Hdac5* gene expression following exposure to vehicle, 0.25mM or 0.5mM palmitate (PA) for 24hrs (One-way ANOVA $p=0.0195$ $X^2=0.3256$, *$p<0.05$ vs Vehicle by Tukey's multiple comparisons test). (C) *Hdac4* and *Hdac5* gene expression in tibialis anterior skeletal muscle from Control and *db/db* mice (Unpaired t-test, $p=0.0253$). Data are mean±SEM, n=6-8 biological replicates per group for cell experiments and 6-8 per group for animal experiments.

**Figure Supplement 2: Individual HDAC isoform and gene dosage effects on *Ppargc1a* expression.** (A) The bilateral AAV6 HDAC4 and 5 overexpression mouse model. (B) *Hdac4* and *Hdac5* gene expression; (C) HDAC4 and 5 protein; (D) and (E) *Ppargc1a* gene expression in skeletal muscle administered empty AAV6 (Control), HDAC4 AAV6 (Two-way ANOVA, genotype $p<0.0001$ $F(1,28)=52.30$; #$p<0.05$ vs Control for same AAV dose by Tukey's multiple comparisons tests) or HDAC5 AAV6 (Two-way ANOVA, genotype $p<0.0001$ $F(1,28)=112.7$; #$p<0.05$ vs Control for same AAV dose by Tukey's multiple comparisons tests). Data are mean±SEM, n=6-8 per group.

**FIGURE SUPPLEMENT 3**

**Figure Supplement 3: Plasma tracer and muscle glycogen concentrations.** (A) Plasma [3]H at 0, 5, 15 and 30 min (Two-way ANOVA, time $p<0.0001$ $F(2.607,36.5)=68.08$, interaction $p=0.0079$ $F(3,42)=4.511$, [#]$p<0.05$ vs Vehicle for same time point by Sidak's multiple comparisons tests) and plasma [3]H area-under-the-curve (AUC; Unpaired t-test $p=0.0003$) following vehicle or insulin and [3]H-2-deoxyglucose administration. (B) Plasma [14]C at 0, 5, 15 and 30 min (Two-way ANOVA, time $p<0.0001$ $F(3,21)=47.84$, treatment $p=0.0442$ $F(1,7)=5.997$, interaction $p=0.0010$ $F(3,21)=7.944$, [#]$p<0.05$ vs Vehicle for same time point by Sidak's multiple comparisons tests) and plasma [14]C AUC (Unpaired t-test $p<0.0001$) following vehicle or insulin and [14]C-glucose administration. (C) Glycogen (Two-way ANOVA, [Ø]genotype $p=0.0053$ $F(1,28)=9.131$) in Control or HDAC4 and 5 overexpressing skeletal muscle. Data are mean±SEM, n=6-8 per group.

**FIGURE SUPPLEMENT 4**

**Figure Supplement 4: Characterisation of HDAC4&5 overexpression in C2C12 myoblasts.** (A) HDAC4 and 5 protein; (B) *Ppargc1a* gene expression (Unpaired t-test, *p*=0.0475); (C) Bubble plot of metabolic pathway gene expression; (D) Oxygen consumption rate (OCR) linked to basal (Unpaired t-test, *p*<0.0001), ATP turnover (Unpaired t-test, *p*<0.0001), uncoupled (Unpaired t-test, *p*<0.0001) and maximal respiration (Unpaired t-test, *p*<0.0001) in Control and HDAC4 and 5 overexpressing C2C12 myoblasts. (E) Identification of co-differentially expressed genes in skeletal muscle from *db/db* mice and C2C12 myoblasts overexpressing HDAC4 and 5 (n=3/group for skeletal muscle and cell analyses) followed by pathway enrichment analysis revealed the p53 signalling pathway as being significantly enriched in this geneset. Data are mean±SEM, n=6-20 biological replicates per group. # p<0.05 vs Control.

**FIGURE SUPPLEMENT 5**

**Figure Supplement 5: Bioenergetics in response to wild type and mutant p53 overexpression.** (A) Oxygen consumption rate (OCR) linked to basal, ATP turnover, uncoupled and maximal respiration and (B) Basal and maximal extracellular acidification rate (ECAR) in C2C12 myoblasts expressing WT, K120R or K120Q p53. Data are mean±SEM, n=8 biological replicates per group.

**FIGURE SUPPLEMENT 6**

**Figure Supplement 6: ROS and phospholipid responses to HDAC4&5 overexpression.** (A) Total $H_2O_2$ release from Control and HDAC4 and 5 overexpressing C2C12 myoblasts (Unpaired t-test, $p<0.0001$). (B) Phosphatidylcholine (PC), lysophosphatidylcholine (LPC), lysophosphatidyl-ethanolamine (LPE) and phosphatidylethanolamine (PE) lipids; (C) Saturation of PC, PE and phosphatidylinositol (PI) lipids, and; (D) Saturation of PC, PE and PI lipids expressed as a percentage of total lipid within each species in Control or HDAC4 and 5 overexpressing skeletal muscle. Data are mean±SEM, n=6 biological replicates per group for cell experiments and n=7 per group in mouse experiments. # p<0.05 vs Control, ∅ p,0.05 main effect for genotype.

**FIGURE SUPPLEMENT 7**

**Figure Supplement 7:** *Slc7a11* **expression, GPX activity and fibre diameter in response to DN HDAC4&5 expression in obese** *db/db* **mice.** (A) *Slc7a11* expression in C2C12 myoblasts or mouse skeletal muscle. (B) Glutathione peroxidase (GPX) activity, and; (C) Fibre diameter in Control and DN HDAC4 and 5 skeletal muscle of *db/db* mice. Data are mean±SEM, n=7-8 per group.