1   *Brief Report*

2   Reclassification of probiotic *Lactobacillus acidophilus* NCIMB

3   30184 as *Lactobacillus helveticus* and *Lactobacillus casei* NCIMB

4   30185 as *Lacticaseibacillus paracasei*

Rodenes, A[1]., Casinos, B.[1], Alvarez, B.[1], Martinez-Blanch, J.F.[1], Vijayakumar, V.[2*], Superti, G.[3], Day, R.[2],

Chenoll, E[1*].

[1] ADM-BIOPOLIS. ADM Health & Wellness, 46980 Paterna, Valencia, Spain

[2]ADM-PROTEXIN, ADM Health & Wellness, Lopen Head TA13 5JH, UK

[3]ADM do Brasil ADM Health & Wellness, São Paulo, SP, Brazil

*Corresponding authors:

maria.chenoll@adm.com. ADM-BIOPOLIS, 46980 Paterna, Valencia, Spain

vineetha.vijayakumar@adm.com ADM Protexin, Lopen Head TA13 5JH, UK

ABSTRACT

The strains NCIMB 30184 and NCIMB 30185, originally designated as *Lactobacillus*

acidophilus and Lactobacillus casei respectively, have been studied in published pre-

clinical and clinical studies. Their nucleotide sequences of the 16S rRNA were compared

with public databases and were preliminarily found to have the highest similarity with

Lactobacillus helveticus and *Lacticaseibacillus paracasei* species respectively.

Subsequently, total 16S and 23S rRNA genes were compared with their most similar taxa,

and the preliminary identification as *Lactobacillus helveticus* and *Lacticaseibacillus*

paracasei were confirmed.

INTRODUCTION

Initially proposed by Beijerinck in 1901 as Gram-positive, homofermentative,

thermophilic and non-spore- forming rods, genus *Lactobacillus* has experienced multiple

taxonomical reassignments. Initially based on phenotypic traits [1], its taxonomic

assignment criteria has been modified throughout the last decades mostly based on

molecular approaches. Sequencing of 16S rRNA genes has been established in bacterial

taxonomy as a marker gene in phylogenetic studies for classification and nomenclature

purposes [2]. However, *Lactobacillus* is a very complex genus, in which the initial

phenotypic differentiations are highly variable within clades [3,4] with an ever increasing

number of new species and re-designations [5,6]. This heterogeneity was finally studied

as a whole by Zheng and coworkers [7], who divided the genus in 23 different genera. All

of these reassignments and changes have made new taxonomical evaluation of the

strains necessary and have reinforced the taxonomical complexity of this genus in its

identification.

Strain NCIMB 30184, originally designated as lactic acid bacteria *Lactobacillus*

acidophilus, has been studied in published pre-clinical and clinical studies [8,9]. An isolate

from human origin, probiotic strain PXN35 was deposited at National Collection of

Industrial, Food and Marine Bacteria (NCIMB, Aberdeen, United Kingdom) on the 7th of

April 2006 as *L. acidophilus*.

Strain NCIMB 30185 was identified initially as *Lactobacillus casei* and its functionality

was informed in [8,9]. The probiotic strain PXN37 was deposited at NCIMB collection on

the 7th of April 2006 as *L. casei*.

In both NCIMB 30184 and NCIMB 30185 strains, *in vitro* microbiological and molecular

evaluation performed showed that the genetic characteristics presented by the isolates

differed from their original designation. In order to determine the correct identification,

the strains were revived from the original master cell banks and 16s rRNA and 23s rRNA

gene sequencing analysis were performed to allow comparison with closely related

species and determine the correct classification of these probiotic strains.

MATERIALS AND METHODS

Culture and growth conditions The long-term storage stock of the strains from NCIMB

(vial dated 22nd October 2009 for PXN 35 and 04th November 2002 for PXN 37)) were

revived by culturing in MRS (Oxoid) at 30 °C under microaerophilic conditions. Working

Reclassification of probiotic strain NCIMB 30184 as *L. helveticus* and NCIMB 30185 *as L. paracasei*

cell banks were conserved in 20 % (v/v) glycerol at -80 °C. Microbiological evaluation

was applied by standard methodology.

Nucleotide sequence analysis. DNA from pure culture was extracted following the

guanidium thiocyanate method isolation [10], spectrophotometrically quantified

(Nanodrop) and adjusted to the final concentration with ultra-pure water (Sigma-Aldrich,

Madrid, Spain). 16s rRNA gene sequencing was run following Chenoll et al [11]. Almost

full-length 16S rRNA gene fragment was amplified from genomic DNA by using the

following primers: 616 Valt (5'-AGA GTT TGA TYMTGG CTC AG-3') and 630R (5'-CAK AAA

GGA GGT GATCC-3'). PCR amplifications were conducted in a solution containing 1× PCR

buffer, 100 µmol/l of each dNTP, 1 µmol/l of each primer, 1.5 mmol/l $MgCl_2$, 0.5 U of

thermoestable Biotaq™ DNA polymerase (BioLine, Randolph, MA) and 5 µl of DNA

template, in a final volume of 50 µl. Amplification conditions were: 5 min at 94 °C, 35

cycles of 30 sec at 94 °C, 60 sec at 56 °C and 90 sec at 72 °C, and a final extension of 10

min at 72 °C. Almost full-length 23S rRNA gene was in vitro amplified using three

universal primer sets following Elizaquivel and coworkers [12]: 1027V/504R, and Ibrahim

et al. 2001 [13]: 1104F/1930R and 1522F/130R (all amplification primers are listed in

Table 1). PCR amplifications were carried as described above. Amplification conditions

were 5 min at 94 °C, 35 cycles of 30s at 94°C, 45 s at 52 °C and 45 s (with a 5 s increment

in each cycle) at 72 °C for elongation, and a final extension of 10 min at 72 °C. All

reactions were carried out in Mastercycler nexus 2x system thermal cycler (Eppendorf).

Controls devoid of DNA were simultaneously included in the amplification process. The

integrity of the PCR products was checked by single bands detection following

electrophoresis for 1 h at 100 V in 1.4% (wt/vol) agarose gels in Tris-borate-EDTA buffer.

Table 1. Primers sequences to amplify 16S and 23S rRNA.

ID primer	rRNA	Sequence (5'-3')
616 Valt	16S rRNA	5'-AGAGTTTGATYMTGGCTCAG-3'
630R	16S rRNA	5'-CAKAAAGGAGGTGATCC-3'
1522F	23S rRNA	5'-TGCGGTGGATCACCTCCTT-3'
130R	23S rRNA	5'-CCTTGCCCCATTCGG-3'
1104F	23S rRNA	5'-TGCGTAATAGCTCAC-3'
1930R	23S rRNA	5'-CGACAAGGAATTTCGCTAC-3'
1027V	23S rRNA	5'-AAACCGACACAGGTRG-3'
504R	23S rRNA	5'-SWGTTCGRVAWGGGA-3'

Sequencing of the 16S and 23S rRNA genes and sequence data analysis. Almost full

sequences of the 16S and 23S rRNA genes were amplified separately as indicated above.

Purification of amplification products was applied by QIAquick PCR purification kit

(QIAGEN) and subsequent sequencing reactions were performed using the Big Dye

Terminator (version 3.1) cycle sequencing kit (Applied Biosystems) in a premixed format.

The 16S rRNA and 23S rRNA sequences determined in this study have been deposited in

the GenBank data base (PXN35 accession reference ON495975 and ON495977

respectively, and PXN37 accession reference OP623495 and OP623508 respectively).

The 16S and 23S rRNA sequences were aligned separately and initially inspected by eye

and   compared   by   the   use   of   the   online   tool   BLAST

(http://blast.ncbi.nlm.nih.gov/Blast.cgi). The strains were tentatively identified on the

basis of highest scores. For each strain, two phylogenetic trees were constructed based

on 16S and 23 rRNA genes with sequences from type strains (Tables 2-5). Sequences

were downloaded from NCBI database and *Bifidobacterium longum* ATCC 15707 was

used as outgroup. T-Coffee multiple sequence alignment software [14] was used to align

the selected sequences. The jModelTest algorithm [15,16] was used to statistically select

the best nucleotide substitution model. In this case, the selected model was TPM2uf+G.

PhyML software [17] was then used to check and build the 16S rRNA phylogenetic tree.

Finally, the iTOL v5 interactive platform [18] was used for its visualization. Percentage of

identity among sequences was obtained by Clustal Omega v2.1 package with default

parameters [19].

Table 2. PXN35 16s rRNA gene sequences used for the study.

Strain	Sequence ID
L. iners DSM 13335[T]	NR036982
L. hominis DSM 23910[T]	NR125548
L. gasseri ATCC 33323[T]	NR075051
L. delbrueckii subsp *bulgaricus* ATCC 11842[T]	NR075019
L. kefiranofaciens subsp *kefiranofaciens* DSM 5016	AM113781
L. acidophilus TMPC 32111	OM281001
L. acidophilus 10058	OK447646
L. acidophilus JCM 1132[T]	HM162411
L. acidophilus ATCC 4356[T]	AB008203
L. crispatus ATCC 33820[T]	NR041800
L. amylovorus DSM 20531[T]	AY944408
L. kitasatonis JCM 1039[T]	AB107638
L. ultunensis DSM 10647[T]	AY253660
L. helveticus IDCC 3801	GU121903
PXN35	ON495975
L. helveticus D5401	EU183494
L. gallinarum ATCC 33199[T]	NR042111
L. helveticus DSM 20075[T]	NR042439
L. helveticus JCM 1120[T]	LC062899
L. helveticus C09	EU377824
L. pasteurii DSM 23907[T]	NR117058
L. amylolyticus DSM 11664[T]	NR029352

110   Table 3. PXN37 16s rRNA gene sequences used for the study.

Strain	Sequence ID
PXN37	PXN37
Lacticaseibacillus absianus YH-lac23	MT009028.2
Lacticaseibacillus chiayiensis BCRC81062	MF446960.1
Lacticaseibacillus chiayiensis NCYUAS	NZ_MSSM01000072.1
Lacticaseibacillus brantae DSM23927[T]	NZ_AYZQ01000010.1
Lacticaseibacillus casei ATCC393[T]	AF469172.1
Lacticaseibacillus casei BIO5773	NZ_WBOC01000018.1
Lactobacillus crispatus ATCC33820[T]	NZ_AZCW01000112.1
Lacticaseibacillus paracasei LOCK919	NC_021721.1
Lacticaseibacillus paracasei subsp *paracasei* JCM8130[T]	D79212.1

Reclassification of probiotic strain NCIMB 30184 as *L. helveticus* and NCIMB 30185 *as L. paracasei*

Lacticaseibacillus rhamnosus NCTC13764[T]	NZ_LR134331.1
Lacticaseibacillus rhamnosus NRRL B-442	NZ_PKQF01000030.1
Lacticaseibacillus saniviri CJNU3004	NZ_JAIZVU010000062.1
Lacticaseibacillus saniviri DSM24301[T]	AB602569.1

112   Table 4. PXN35 23S rRNA gene sequences used for the study.

Strain	Sequence ID
L. iners DSM 13335[T]	NZ_GG700803.1
L. gasseri 63 AM	NR_076437.1
L. hominis DSM 23910	NZ_AYZP01000029.1
L. delbrueckii subsp *bulgaricus* ATCC 11842[T]	NR_077006.1
L. amylolyticus DSM 11664[T]	Y17360.1
L. pasteurii DSM 23907[T]	NZ_PUFQ01000021.1
L. kefiranofaciens subsp *kefiranofaciens* JCM 6985	NZ_BAMG01000080.1
L. ultunensis Kx293C1[T]	NZ_CP059829.1
L. acidophilus NCFM	NR_076355.1
L. amylovorus GRL 1112	NR_076802.1
L. kitasatonis JCM1039[T]	NZ_BALU01000025.1
L. crispatus ST1	NR_077026.1
L. gallinarum DSM 10532[T]	NZ_AZEL01000029.1
L. helveticus DPC 4571	NR_076471.1
PXN35	PXN35

114   Table 5. PXN37 23S rRNA gene sequences used for the study.

Strain	Sequence ID
PXN37	PXN37
Lacticaseibacillus absianus YH-lac23	NZ_JABEKX010000006.1
Lacticaseibacillus chiayiensis BCRC81062	NZ_NOXN01000109.1
Lacticaseibacillus chiayiensis NCYUAS	NZ_MSSM01000072.1
Lacticaseibacillus brantae DSM23927[T]	NZ_AYZQ01000009.1
Lacticaseibacillus casei ATCC393[T]	NZ_AP012544.1
Lacticaseibacillus casei BIO5773	NZ_WBOC01000014.1
Lactobacillus crispatus ATCC33820[T]	NZ_AZCW01000100.1
Lacticaseibacillus paracasei LOCK919	NC_021721.1
Lacticaseibacillus paracasei subsp *paracasei* JCM8130[T]	NZ_AP012541.1
Lacticaseibacillus rhamnosus NCTC13764[T]	NZ_LR134331.1
Lacticaseibacillus rhamnosus NRRL B-442	NZ_PKQF01000030.1
Lacticaseibacillus saniviri CJNU3004	NZ_JAIZVU010000042.1
Lacticaseibacillus saniviri DSM24301[T]	NZ_JQCE01000031.1

116   **RESULTS AND DISCUSSION**

Reclassification of probiotic strain NCIMB 30184 as *L. helveticus* and NCIMB 30185 *as L. paracasei*

Strain PXN35

Strain PXN35 was originally designated as *L. acidophilus* and deposited at National

Collection of Industrial, Food and Marine Bacteria (NCIMB, Aberdeen, United Kingdom)

as NCIMB 30184.

Due to *in vitro* microbiological and molecular evaluation discrepancies with *L.*

acidophilus designation, almost full 16s rRNA gene sequencing was applied with a

sample from the revived original master cell bank. Initially, *on line* tool BLAST grouped

the strain with *Lactobacillus helveticus* sequences (data not shown), as a preliminary

tentative identification. In order to confirm BLAST identification, a phylogenetic tree was

constructed based on 16S rRNA with its closest phylogenetic neighbors. For the analysis,

we selected the closest members of *Lactobacillus* phylogenetic group, using the most

recent reclassification available and scientifically accepted [7]. The analysis grouped

PXN35 sequence with all the *L. helveticus* sequences studied, including the species type

strain sequence [7]. *Lactobacillus acidophilus* grouped separately. PXN35 and *L.*

helveticus shared 99.43% 16s rRNA gene on average, whereas strain PXN35 only showed

97.83% homology with *L. acidophilus* representatives (Figure 1, Table 6).

Figure 1. Phylogenetic tree constructed using the neighbour-joining method based on 16S rRNA gene sequences. Strain PXN35 sequence is close

to *L. helveticus*. Bar, 10% nucleotide substitution.

Table 6. PXN35 16s rRNA gene % homology.

	L. iners strain DSM 13335	L. hominis DSM 23910	L. gasseri ATCC 33323	L. delbrueckii subsp bulgaricus ATCC 11842	L. kefiranofaciens subsp kefiranofaciens DSM 5016	L. acidophilus strain TMPC 32111	L. acidophilus strain 10058	L. acidophilus ATCC 4356	L. crispatus strain ATCC 33820	L. amylovorus strain DSM 20531	L. kitasatonis	L. ultunensis	L. gallinarum strain ATCC 33199	L. helveticus DSM 20075	L. helveticus strain C09	L. pasteurii DSM 23907	L. amylolyticus DSM 11664	PXN35
L. iners strain DSM 13335	100.00																	
L. hominis DSM 23910	94.50	100.00																
L. gasseri ATCC 33323	95.00	98.38	100.00															
L. delbrueckii subsp *bulgaricus* ATCC 11842	89.29	90.77	90.85	100.00														
L. kefiranofaciens subsp *kefiranofaciens* DSM 5016	89.52	91.00	90.93	92.06	100.00													
L. acidophilus strain TMPC 32111	89.57	91.27	91.13	92.33	96.60	100.00												
L. acidophilus strain 10058	89.87	91.70	91.57	93.05	97.24	99.29	100.00											
L. acidophilus ATCC 4356	89.94	91.78	91.64	93.13	97.31	99.22	99.93	100.00										
L. crispatus strain ATCC 33820	90.16	91.92	91.78	92.98	97.88	97.66	98.30	98.37	100.00									
L. amylovorus strain DSM 20531	89.66	91.71	91.57	92.70	97.45	97.52	98.16	98.23	98.73	100.00								
L. kitasatonis	89.52	91.71	91.57	92.77	97.66	97.87	98.51	98.58	98.80	99.50	100.00							
L. ultunensis	90.30	91.99	91.86	92.63	97.31	97.09	97.73	97.81	98.30	98.23	98.23	100.00						
L. helveticus strain IDCC 3801	89.57	91.27	91.21	91.70	95.96	97.30	97.02	96.95	97.09	96.88	96.88	96.95						
L. gallinarum strain ATCC 33199	90.30	92.20	92.07	93.13	97.31	97.73	98.37	98.44	98.58	98.23	98.37	98.44	100.00					
L. helveticus DSM 20075	90.16	92.06	91.93	92.84	97.03	97.59	98.23	98.30	98.30	97.95	98.09	98.16	99.58	100.00				

L. helveticus strain C09	90.23	92.13	92.00	92.77	97.10	97.66	98.30	98.37	98.37	98.02	98.16	98.23	99.65	99.93	100.00			
L. pasteurii DSM 23907	91.80	93.07	93.30	93.91	96.09	96.17	96.17	96.17	95.86	96.32	96.24	96.17	96.09	95.79	95.86	100.00		
L. amylolyticus DSM 11664	90.15	91.91	91.85	92.48	95.54	95.25	95.89	95.96	96.03	96.60	96.60	95.54	95.82	95.54	95.61	96.84	100.00	
PXN35	90.16	92.06	91.93	92.70	97.10	97.38	98.02	98.09	98.23	98.02	98.02	98.09	99.50	99.36	99.43	95.79	95.68	100.00

After that, we ran a further phylogenetic analysis with the same group of strains by

comparing the almost full-length large ribosomal subunit 23S. Both 16S and 23S

phylogenies are described to be in good agreement and provide an advantage over short

ones like 5S, as local non-random rearrangements do not disturb the overall result [20].

The 23S phylogenetic tree structure closely resembled the 16S, with PXN35 grouped

together with *L. helveticus* representatives and separated from *L. acidophilus*. When

percentage homologies were studied, the analysis showed the highest nucleotide

sequence homology between PXN35 and *L. helveticus* 23S rRNA gene (99.61%), whereas

strain PXN35 showed 97.26% homology with *L. acidophilus* representatives (Figure 2,

Table 7).

Figure 2. Phylogenetic tree constructed using the neighbour-joining method based on 23S rRNA gene sequences. Strain PXN35 sequence is close

to *L. helveticus*. Bar, 10% nucleotide substitution.

Table 7. 23s rRNA gene % homology (only reference sequence is considered).

	L. iners DSM 13335	*L. gasseri* 63 AM	*L. hominis* DSM 23910	*L. delbrueckii* subsp *bulgaricus* strain ATCC 11842	*L. amylolyticus* DSM 11664	*L. pasteurii* DSM 23907	*L. kefiranofaciens* subsp *Kefiranofaciens* JCM 6985	*L. ultunensis* Kx293C1	*L. acidophilus* strain NCFM	*L. amylovorus* GRL 1112	*L. kitasatonis* JCM1039	*L. crispatus* ST1	*L. gallinarum* DSM 10532	*L. helveticus* DPC 4571	PXN35
L. iners DSM 13335	100.00														
L. gasseri 63 AM	93.05	100.00													
L. *hominis* DSM 23910	92.27	95.65	100.00												
L. delbrueckii subsp *bulgaricus* ATCC 11842	89.47	88.97	89.24	100.00											
L. amylolyticus DSM 11664	89.24	90.15	90.07	89.49	100.00										
L. pasteurii DSM 23907	89.89	90.45	91.02	90.75	92.62	100.00									
L. kefiranofaciens subsp *kefiranofaciens* JCM 6985	89.46	90.33	90.47	89.24	92.98	92.12	100.00								
L. ultunensis Kx293C1	89.81	91.32	90.54	88.84	93.01	92.15	96.52	100.00							
L. acidophilus strain NCFM	89.63	90.89	90.37	88.40	92.18	91.71	95.52	97.26	100.00						
L. amylovorus GRL 1112	89.81	91.29	90.47	89.15	92.84	92.64	96.00	96.87	97.43	100.00					
L. kitasatonis JCM1039	89.72	91.33	90.77	88.93	92.84	92.72	95.87	96.69	97.35	99.17	100.00				
L. crispatus ST1	89.72	90.76	90.51	89.15	92.89	92.29	95.78	96.30	96.69	97.00	97.13	100.00			
L. gallinarum DSM 10532	90.07	91.10	90.32	89.31	92.66	92.19	95.99	97.17	97.21	97.82	97.47	97.74	100.00		
L. helveticus DPC 4571	90.07	91.27	90.58	89.44	92.66	92.23	95.86	96.95	97.26	97.65	97.26	97.43	99.22	100.00	
PXN35	90.11	91.22	90.58	89.52	92.48	92.14	95.95	96.91	97.26	97.69	97.26	97.43	99.17	99.61	100.00

Isolated from a human individual and deposited in NCIMB collection as NCIMB 30184,

strain PXN35 is a gram-positive, catalase negative strain. Cells are nonmotile and non-

spore-forming, rod-shaped, isolated or tend to form pairs, with 1.0 μm x 5-15 μm. Strain

is obligately homofermentative and grows well in commercially available media for lactic

acid bacteria (MRS, Mann Rogosa and Sharpe) at 37°C. Colonies of strain PXN35 grown

on MRS agar plates are grey, smooth, slightly domed, and round. The strain PXN35 has

been shown to inhibit growth of *Salmonella Typhimurium*, *Clostridium difficile*,

Staphylococcus aureus, *Escherichia coli* and *Enterococcus faecalis*. Strain PXN35

produces organic acids and significantly reduces pH [21, 22] . Originally designated as *L.*

acidophilus, the present study demonstrates its unequivocal identification as *L.*

helveticus.

Strain PXN37

Originally identified as *L. casei,* strain PXN37 was deposited at NCIMB collection as

NCIMB 30185. Further molecular evaluation with almost full 16s rRNA gene sequencing

with the revived original master cell bank showed some divergences with *L. casei* species

and BLAST tool tentatively grouped the strain with *Lacticaseibacillus paracasei*

sequences (data not shown). The initial BLAST identification was checked by a

phylogenetic tree based on 16S rRNA with is closest phylogenetic neighbors as Zheng

and coworkers established [7]. Strain PXN37 was clearly grouped with all the *L. paracasei*

sequences including the species type strain sequence [7] sharing 100% 16s rRNA gene

on average with *L. paracasei* whereas *L. casei* representatives grouped separately

(Figure 3, Table 8).

Figure 3. Phylogenetic tree constructed using the neighbour-joining method based on 16S rRNA gene sequences. Strain PXN37 sequence is close

to *L. paracasei*. Bar, 10% nucleotide substitution.

Table 8. PXN37 16s rRNA gene % homology.

	L. crispatus ATCC33820[T]	*L. absianus* YH-lac23	*L. rhamnosus* NCTC13764[T]	*L. rhamnosus* NRRL B-442	*L. paracasei* subsp. *paracasei* JCM8130[T]	*L. paracasei* LOCK919	*L. casei* ATCC393T	*L. chiayiensis* NCYUAS	*L. chiayiensis* BCRC81062	*L casei* BIO5773	*L. saniviri* DSM 24301[T]	*L. saniviri* CJNU3004	*L. brantae* DSM23927T	PXN37
Lactobacillus crispatus ATCC33820[T]	100.00													
Lacticaseibacillus absianus YH-lac23	86.38	100.00												
Lacticaseibacillus rhamnosus NCTC13764[T]	86.37	94.03	100.00											
Lacticaseibacillus rhamnosus NRRL B-442	86.37	94.03	100.00	100.00										
Lacticaseibacillus paracasei subsp. *paracasei* JCM8130[T]	86.37	93.83	98.85	98.85	100.00									
Lacticaseibacillus paracasei LOCK919	86.37	93.83	98.85	98.85	100.00	100.00								
Lacticaseibacillus casei ATCC393[T]	86.51	93.83	98.78	98.78	99.19	99.19	100.00							
Lacticaseibacillus chiayiensis NCYUAS	86.43	93.96	98.92	98.92	99.19	99.19	99.73	100.00						
Lacticaseibacillus chiayiensis BCRC81062	86.43	93.89	98.92	98.92	99.19	99.19	99.73	99.93	100.00					
Lacticaseibacillus casei BIO5773	86.5	93.96	98.92	98.92	99.19	99.19	99.8	99.86	99.86	100.00				
Lacticaseibacillus saniviri DSM 24301[T]	87.4	93.96	94.65	94.65	94.51	94.51	94.44	94.58	94.51	94.51	100.00			
Lacticaseibacillus saniviri CJNU3004	87.4	93.96	94.65	94.65	94.51	94.51	94.44	94.58	94.51	94.51	100.00	100.00		
Lacticaseibacillus brantae DSM23927[T]	86.99	94.09	94.65	94.65	94.65	94.65	94.38	94.58	94.51	94.44	97.43	97.43	100.00	
PXN37	86.37	93.83	98.85	98.85	100.00	100.00	99.19	99.19	99.19	99.19	94.51	94.51	94.65	100.00

Reclassification of probiotic strain NCIMB 30184 as *L. helveticus* and NCIMB 30185 *as L. paracasei*

Further phylogenetic analysis with full-length large ribosomal subunit 23S was run as

previously and as in the case of 16s rRNA analysis, PXN37 grouped together with *L.*

paracasei representatives and separated from *L. casei*. When percentage homologies

were studied, the analysis showed the highest nucleotide sequence homology between

PXN37 and *L. paracasei* 23S rRNA gene (100%), and lower (98.59%) with *L. casei*

representatives (Figure 4, Table 9).

201   Figure 4. Phylogenetic tree constructed using the neighbour-joining method based on 23S rRNA gene sequences. Strain PXN37 sequence is close

202   to *L. paracasei*. Bar, 10% nucleotide substitution.

203   Table 9. PXN37 23s rRNA gene % homology.

	L. crispatus ATCC33820T	*L. absianus* YH-lac23	*L. paracasei subsp paracasei* JCM8130[T]	*L. paracasei* LOCK919	*L. rhamnosus* NCTC13764[T]	*L. rhamnosus* NRRL B-442	*L. casei* ATCC393[T]	*L. casei* BIO5773	*L. chiayiensis* BCRC81062	*L. chiayiensis* NCYUAS	*L. brantae* DSM23927[T]	*L. saniviri* CJNU3004	*L. saniviri* DSM24301[T]	PXN37
L. crispatus ATCC33820[T]	100.00													
L. absianus YH-lac23	83.85	100.00												
L. paracasei subsp paracasei JCM8130[T]	83.85	93.48	100.00											
L. paracasei LOCK919	83.85	93.44	99.96	100.00										
L. rhamnosus NCTC13764[T]	83.57	93.44	98.47	98.51	100.00									
L. rhamnosus NRRL B-442	83.65	93.48	98.51	98.55	99.88	100.00								
L. casei ATCC393[T]	83.49	93.20	98.59	98.59	99.15	99.11	100.00							
L. casei BIO5773	83.49	93.24	98.59	98.59	99.15	99.11	99.84	100.00						
L. chiayiensis BCRC81062	83.68	93.15	98.51	98.51	98.87	98.83	99.40	99.48	100.00					
L. chiayiensis NCYUAS	83.57	93.04	98.39	98.39	98.79	98.75	99.36	99.44	99.92	100.00				
L. brantae DSM23927[T]	84.01	93.92	94.28	94.24	94.12	94.12	93.88	94.04	94.07	93.96	100.00			
L. saniviri CJNU3004	84.30	93.60	94.32	94.28	93.92	93.96	93.72	93.88	93.75	93.64	98.07	100.00		
L. saniviri DSM24301[T]	84.30	93.60	94.32	94.28	93.92	93.96	93.72	93.88	93.75	93.64	98.07	100.00	100.00	
PXN37	83.85	93.48	100.00	99.96	98.47	98.51	98.59	98.59	98.51	98.39	94.28	94.32	94.32	100.00

Reclassification of probiotic strain NCIMB 30184 as *L. helveticus* and NCIMB 30185 *as L. paracasei*

The strain PXN37 was isolated from human origin as a gram-positive, catalase negative

strain and deposited in NCIMB collection under reference NCIMB 30185. Cells are

nonmotile and non-spore-forming, rod-shaped. Strain is obligately homofermentative

and grows well in commercially available media for lactic acid bacteria (MRS, Mann

Rogosa and Sharpe) at 37°C. The strain PXN37 has been shown to inhibit growth of

Salmonella Typhimurium, *Clostridium difficile*, *Staphylococcus aureus*, *Escherichia coli*

and *Enterococcus faecalis*. Strain PXN37 produces organic acids and significantly reduces

pH [21,22]. Originally designated as *L. casei*, the present study demonstrates its

unequivocal identification as *Lacticaseibacillus paracasei*.

ACKNOWLEDGEMENTS

228   We thank Janice Rueda and Marta Tortajada for critical revision.

COMPETING INTERESTS

230   A.R., B.C., B.A., J.F.M-B. and E.C. are employed by Biopolis ADM. V.V. and R.D. are

231   employed by Protexin ADM. G.S. is employed by ADM Brazil.

REFERENCES

1.   Orla-Jensen, S. *The lactic acid bacteria*; Fred, A., Son, H.a., Eds.; Copenhagen:, 1919.

2.   Schleifer, K.H.; Stackebrandt, E. Molecular systematics of prokaryotes. *Annu Rev*
Microbiol **1983**, *37*, 143-187, doi:10.1146/annurev.mi.37.100183.001043.

3.   Zheng, J.; Ruan, L.; Sun, M.; Gänzle, M. A Genomic View of Lactobacilli and Pediococci
Demonstrates that Phylogeny Matches Ecology and Physiology. *Appl Environ Microbiol*
2015, *81*, 7233-7243, doi:10.1128/aem.02116-15.

4.   Sun, Z.; Harris, H.M.; McCann, A.; Guo, C.; Argimón, S.; Zhang, W.; Yang, X.; Jeffery, I.B.;
Cooney, J.C.; Kagawa, T.F., et al. Expanding the biotechnology potential of lactobacilli
through comparative genomics of 213 strains and associated genera. *Nat Commun* **2015**,
6, 8322, doi:10.1038/ncomms9322.

5.   Vancanneyt, M.; Naser, S.M.; Engelbeen, K.; De Wachter, M.; Van der Meulen, R.;
Cleenwerck, I.; Hoste, B.; De Vuyst, L.; Swings, J. Reclassification of Lactobacillus brevis
strains LMG 11494 and LMG 11984 as Lactobacillus parabrevis sp. nov. *Int J Syst Evol*
Microbiol **2006**, *56*, 1553-1557, doi:10.1099/ijs.0.64215-0.

6.   Naser, S.M.; Vancanneyt, M.; Snauwaert, C.; Vrancken, G.; Hoste, B.; De Vuyst, L.; Swings,
J. Reclassification of Lactobacillus amylophilus LMG 11400 and NRRL B-4435 as
Lactobacillus amylotrophicus sp. nov. *Int J Syst Evol Microbiol* **2006**, *56*, 2523-2527,
doi:10.1099/ijs.0.64463-0.

7.   Zheng, J.; Wittouck, S.; Salvetti, E.; Franz, C.; Harris, H.M.B.; Mattarelli, P.; O'Toole, P.W.;
Pot, B.; Vandamme, P.; Walter, J., et al. A taxonomic note on the genus Lactobacillus:
Description of 23 novel genera, emended description of the genus Lactobacillus
Beijerinck 1901, and union of Lactobacillaceae and Leuconostocaceae. *Int J Syst Evol*
Microbiol **2020**, *70*, 2782-2858, doi:10.1099/ijsem.0.004107.

8.   Chapman, C.M.; Gibson, G.R.; Todd, S.; Rowland, I. Comparative in vitro inhibition of
urinary tract pathogens by single- and multi-strain probiotics. *Eur J Nutr* **2013**, *52*, 1669-
1677, doi:10.1007/s00394-013-0501-2.

9.   Koradia, P.; Kapadia, S.; Trivedi, Y.; Chanchu, G.; Harper, A. Probiotic and cranberry
supplementation for preventing recurrent uncomplicated urinary tract infections in
premenopausal women: a controlled pilot study. *Expert Rev Anti Infect Ther* **2019**, *17*,
733-740, doi:10.1080/14787210.2019.1664287.

10. Pitcher, D.G.; Saunders, N.A.; Owen, R.J. Rapid extraction of bacterial genomic DNA with
guanidium thiocyanate. *Letters in Applied Microbiology* **1989**, *8*, 151-156,
doi:https://doi.org/10.1111/j.1472-765X.1989.tb00262.x.
11. Chenoll, E.; Macián, M.C.; Aznar, R. Identification of Carnobacterium, Lactobacillus,
Leuconostoc and Pediococcus by rDNA-based techniques. *Syst Appl Microbiol* **2003**, *26*,
546-556, doi:10.1078/072320203770865855.
12. Elizaquível, P.; Chenoll, E.; Aznar, R. A TaqMan-based real-time PCR assay for the specific
detection and quantification of Leuconostoc mesenteroides in meat products. *FEMS*
Microbiol Lett **2008**, *278*, 62-71, doi:10.1111/j.1574-6968.2007.00974.x.
13. Ibrahim, A.; Hofman-Bang, J.; Ahring, B.K. Amplification and direct sequence analysis of
the 23S rRNA gene from thermophilic bacteria. *Biotechniques* **2001**, *30*, 414-416, 418,
420, doi:10.2144/01302rr05.
14. Di Tommaso, P.; Moretti, S.; Xenarios, I.; Orobitg, M.; Montanyola, A.; Chang, J.M.; Taly,
J.F.; Notredame, C. T-Coffee: a web server for the multiple sequence alignment of
protein and RNA sequences using structural information and homology extension.
Nucleic Acids Res **2011**, *39*, W13-17, doi:10.1093/nar/gkr245.
15. Darriba, D.; Taboada, G.L.; Doallo, R.; Posada, D. jModelTest 2: more models, new
heuristics and parallel computing. *Nat Methods* **2012**, *9*, 772, doi:10.1038/nmeth.2109.
16. Guindon, S.; Gascuel, O. A simple, fast, and accurate algorithm to estimate large
phylogenies by maximum likelihood. *Syst Biol* **2003**, *52*, 696-704,
doi:10.1080/10635150390235520.
17. Guindon, S.; Dufayard, J.F.; Lefort, V.; Anisimova, M.; Hordijk, W.; Gascuel, O. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. *Syst Biol* **2010**, *59*, 307-321, doi:10.1093/sysbio/syq010.
18. Letunic, I.; Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic Acids Res* **2021**, *49*, W293-w296,
doi:10.1093/nar/gkab301.
19. Sievers, F.; Higgins, D.G. The Clustal Omega Multiple Alignment Package. *Methods Mol*
Biol **2021**, *2231*, 3-16, doi:10.1007/978-1-0716-1036-7_1.
20. Ludwig, W.; Schleifer, K.H. Bacterial phylogeny based on 16S and 23S rRNA sequence
analysis. *FEMS Microbiology Reviews* **1994**, *15*, 155-173, doi:10.1111/j.1574-
6976.1994.tb00132.x.
21. Tejero-Sariñena, S.; Barlow, J.; Costabile, A.; Gibson, G.R.; Rowland, I. In vitro evaluation
of the antimicrobial activity of a range of probiotics against pathogens: evidence for the
effects of organic acids. *Anaerobe* **2012**, *18*, 530-538,
doi:10.1016/j.anaerobe.2012.08.004.

22. Tejero-Sariñena, S.; Barlow, J.; Costabile, A.; Gibson, G.R.; Rowland, I. Antipathogenic
activity of probiotics against Salmonella Typhimurium and Clostridium difficile in
anaerobic batch culture systems: is it due to synergies in probiotic mixtures or the
specificity of single strains?. *Anaerobe* **2013**, *24*, 60-65, doi:
10.1016/j.anaerobe.2013.09.011.