A marmoset brain cell census reveals influence of developmental origin and functional class on
neuronal identity

Fenna M. Krienen[1,2]*⌂, Kirsten M. Levandowski[2,3,4], Heather Zaniewski[2,3,4], Ricardo C.H. del Rosario[2],
Margaret E. Schroeder[2,3,4], Melissa Goldman[2], Martin Wienisch[2,3,4], Alyssa Lutservitz[1], Victoria F. Beja-
Glasser[2,3,4], Cindy Chen[2,3,4], Qiangge Zhang[2,3,4], Ken Y. Chan[2], Katelyn X. Li[3,4], Jitendra Sharma[3,4], Dana
McCormack[2,3,4], Tay Won Shin[3,4,5], Andrew Harrahill[3,4], Eric Nyase[3,4], Gagandeep Mudhar[6], Abigail
Mauermann[3,4,5], Alec Wysoker[2], James Nemesh[2], Seva Kashin[2], Josselyn Vergara[2], Gabriele Chelini[7],
Jordane Dimidschstein[2], Sabina Berretta[8,9], Benjamin E. Deverman[2], Ed Boyden[3,4,5], Steven A.
McCarroll[1,2#]*, Guoping Feng[2,3#]*

10
11

[1] Department of Genetics, Harvard Medical School; Boston, Massachusetts 02115, USA.
[2] Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard; Cambridge, Massachusetts
02142, USA.
[3] McGovern Institute for Brain Research, Department of Brain and Cognitive Sciences, Massachusetts
Institute of Technology; Cambridge, Massachusetts 02142, USA.
[4] Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology; Cambridge,
Massachusetts 02139, USA.
[5] HHMI, Cambridge, MA, USA.
[6] Princeton Neuroscience Institute, Princeton University. Princeton, NJ 08544, USA.
[7] Center for Mind/Brain Sciences, University of Trento. Piazza della Manifattura n.1, Rovereto (TN).
38068, IT.
[8] Basic Neuroscience Division, McLean Hospital, Belmont MA USA.
[9] Dept. of Psychiatry, Harvard Medical School, Boston, MA USA.
25
[#]These authors contributed equally to this work
*Corresponding authors. Email: fkrienen@princeton.edu, mccarroll@genetics.med.harvard.edu,
fengg@mit.edu
⌂Present affiliation: Princeton Neuroscience Institute. Princeton NJ USA

Abstract:

The mammalian brain is composed of many brain structures, each with its own ontogenetic and
developmental history. Transcriptionally-based cell type taxonomies reveal cell type composition and
similarity relationships within and across brain structures. We sampled over 2.4 million brain cells across
18 locations in the common marmoset, a New World monkey primed for genetic engineering, and used
single-nucleus RNA sequencing to examine global gene expression patterns of cell types within and
across brain structures. Our results indicate that there is generally a high degree of transcriptional
similarity between GABAergic and glutamatergic neurons found in the same brain structure, and there are
generally few shared molecular features between neurons that utilize the same neurotransmitter but reside
in different brain structures. We also show that in many cases the transcriptional identities of cells are
intrinsically retained from their birthplaces, even when they migrate beyond their cephalic compartments.
Thus, the adult transcriptomic identity of most neuronal types appears to be shaped much more by their
developmental identity than by their primary neurotransmitter signaling repertoire. Using quantitative

mapping of single molecule FISH (smFISH) for markers for GABAergic interneurons, we found that the
similar types (e.g. *PVALB*+ interneurons) have distinct biodistributions in the striatum and neocortex.
Interneuron types follow medio-lateral gradients in striatum but form complex distributions across the
neocortex that are not described by simple gradients. Lateral prefrontal areas in marmoset are
distinguished by high relative proportions of *VIP*+ neurons. We further used cell-type-specific enhancer
driven AAV-GFP to visualize the morphology of molecularly-resolved interneuron classes in neocortex
and striatum, including the previously discovered novel primate-specific *TAC3*+ striatal interneurons. Our
comprehensive analyses highlight how lineage and functional class contribute to the transcriptional
identity and biodistribution of primate brain cell types.
52

53 **One-Sentence Summary:** Adult primate neurons are imprinted by their region of origin, more so than by
54 their functional identity.
55

Main Text:
The mammalian brain's complex functional diversity stems from its vast cellular and molecular
repertoire. To provide a more complete understanding of the cell types across major cortical and
subcortical structures in a non-human primate brain, we conducted a census of cell types of the adult
marmoset brain based on their transcriptional profiles. Previous single cell sequencing studies of the
marmoset brain focused on single brain regions (*1*, *2*), or on specific cell classes across regions (*3*, *4*).
However, inclusion of both closely and distantly related brain structures and cell types can yield insights
into the developmental and ontological relationships between them (*5*). Comprehensive transcriptomic
cell type atlases have been produced for the mouse (*6–9*). Complementing these, recent transcriptomic
datasets that sample many regions across humans and nonhuman primates brains (*10–12*) offer powerful
resources for comparative analysis of brain cell type features. We generated single-nucleus RNA
sequencing (snRNA-seq; 10x Genomics 3' v3.1) data from 2.4 million unsorted brain cell nuclei across 8
neocortical and 10 subcortical locations from 10 young adult marmosets (4 M, 6 F), and resolved clusters
from all major neuronal and non-neuronal cell classes. snRNA-seq data were generated as part of the
Brain Initiative Cell Census Network (BICCN, RRID:SCR_015820) and are available on the BICCN
Data Center (RRID:SCR_022815; https://biccn.org/data) as well as the NeMO archive
(RRID:SCR_016152; https://doi.org/10.1101/2022.10.18.512442).

All neuron-containing brain structures in the central nervous system possess both excitatory and
inhibitory neuronal populations, though the proportions and degree of developmental relatedness between
these two populations varies by structure. In the neocortex and other telencephalic structures of
vertebrates, distinct populations of neurons are typically categorized based on their neurotransmitter
status as either inhibitory (GABAergic) or excitatory (glutamatergic). In other brain structures, primary
neurotransmission appears less essential to a cell type's identity. We found that the transcriptional
identities of excitatory and inhibitory neurons within telencephalic brain structures segregate strongly,
consistent with previous studies in other mammalian species (*2*, *6*, *7*, *13*). In contrast, there is much
greater transcriptional similarity between GABAergic and glutamatergic neuronal types in non-
telencephalic compartments. Moreover, few gene expression distinctions present in telencephalic
glutamatergic neurons are shared with glutamatergic neurons in non-telencephalic brain regions. While
primary neurotransmission did not drive transcriptional similarity between neurons, their brain structure

of origin did; the adult transcriptomic identity of most neuronal types appear to be shaped much more by
their developmental identity than by their neurotransmitter signaling repertoire.

In the mouse, overall interneuron/excitatory neuron ratios are thought to be largely invariant across the
neocortex (*14*). Even so, relative proportions of interneuron subtypes do vary and underlie principles of
functional organization in the mouse neocortex (*15*). The small size of the marmoset brain and its near-
lissencephalic neocortex enables quantitative, cell-type-resolved mapping in a primate. We used smFISH
to spatially profile major interneuron types across marmoset striatum and neocortex. In the striatum,
major interneuron types are distributed as medial-lateral gradients. In contrast, the marmoset neocortex
has a much more complex topography of interneuron concentrations that is not explained by a single
spatial axis. Lateral prefrontal areas in particular, which have undergone expansion in primate evolution
(*16*) are typified by higher proportions of *VIP+* and *LAMP5/ID2+* interneurons and lower proportions of
SST+ interneurons relative to all other neocortical areas.

The development of viral tools to effectively deliver transgenes to specific primate brain cell types has
many powerful applications (*17–20*). We used this approach to sparsely label and reconstruct the
morphologies of selected types. We reconstructed a set of molecularly-characterized neurons that had
received systemic delivery of an AAV carrying a reporter (AAV9-hDlx5/6-GFP-fGFP) under an
interneuron-selective regulatory element(*19*). Reconstructions are available for download from the Brain
Image Library (BIL; https://doi.org/10.35077/g.609). As AAV9-hDlx5/6-GFP-fGFP did not label a
recently discovered, primate-specific striatal interneuron type (*3*) w(*3*)This enabled visualization(*3*).
Together, our census of major transcriptomically-defined brain cell types and quantitative mapping of
interneuron biodistributions provides a key resource for the primate neuroscience community and for
comparative studies of cell type evolution.

A transcriptomic census of marmoset brain cell types

We acquired snRNA-seq data from 18 brain regions collected using 10x Genomics 3' 3.1 chemistry
across 10 young adult marmosets (ages 1.5-4; 6 F) as well as a small dataset from PFC of two aged
animals (**Fig. 1, Fig. S1, Table S1**) . The number of donors per brain structure varied (min = 2, max = 10;
Fig. S1), as did the cell sampling rate per brain structure (**Fig. 1A; Fig. S1A, Table S2**). Neocortex was
the most comprehensively sampled in terms of total numbers, donors, and regional dissections. We
acquired data from: cerebellum, brainstem, hypothalamus, thalamus, amygdala, striatum (separate
dissections for caudate, putamen and nucleus accumbens), hippocampus, basal forebrain and neocortex.
Within the neocortex, we separately sampled 8 neocortical locations (prefrontal, temporal pole, S1, M1,
A1, V2, V1, lateral parietal association), and within PFC, from four prefrontal subdivisions (**Fig. S1B-C**).

Nuclei from each brain structure were pooled across donors and analyzed to identify major cell types and
their proportions by brain structure (**Fig. 1B-D**). Using linear discriminant analysis (scPred;(*21*)) trained
on a supervised set of cell class labels, we identified and discarded low quality cells and doublets, and
assigned each nucleus to its probable major type – astrocyte, endothelia, ependyma,
microglia/macrophage, neuron, oligodendrocyte, or oligodendrocyte precursor cell (OPC).

After major cell type assignment, nuclei were clustered by brain structure to reveal subtype diversity
within each major class. We used a previously described clustering pipeline (*6*) based on independent
components analysis (fastICA). Each clustering analysis involved additional curation of doublets and
outlier cells, followed by a second round of sub-clustering of major clusters. At each of these curation

stages, independent components that loaded on individual donors or batches were excluded and
reclustering was performed to attenuate donor and batch effects on clustering results. For each cluster, a
"metacell" was generated by summing transcript counts of individual cells of each type together, followed
by scaling and normalizing to 100,000 transcripts/nucleus. These regional and cell-type resolved
metacells were the starting point for most cross-region analyses. The final snRNA-seq dataset size after
curation contained 2.01 million cells (**Table S2)**.

Hierarchical clustering of neurons
Cell types are identified based on factors including their function, developmental origin, lineage, and
regional context (*22–24*). We studied how transcriptional profiles related to each other across 288
neuronal cell types (metacells) sampled across all 18 marmoset brain structures. Individual replicates and
other variables (age, sex) were generally proportionately represented across resolved cell types with
several exceptions that were likely mainly to differences in dissection across donors (**Fig. S1B-E, Fig S2**).
We used hierarchical clustering to situate the neuronal types on a dendrogram computed using distances
calculated from the top 100 PC scores across neuron types (**Fig. 1C**). Most telencephalic types
(neocortex, amygdala, hippocampus, striatum) clustered distinctly from diencephalic and hindbrain types,
indicating that developmental lineage continues to shape the transcriptional identity of adult primate brain
cell types. However, of the 7 major clades, 4 contained mixtures of cell types at the terminal (leaf level)
originating from distinct cephalic compartments. For example, basal forebrain neuron types intermingled
with hypothalamic types, suggesting closer transcriptional similarity of two structures that occupy distinct
cephalic compartments. The overall dendrogram configuration was broadly conserved when recomputed
using other distance functions (**Fig. S2A**).

Transcription factors are master regulators that determine cell type identity in development through
temporal patterning, suggesting that they may be a key class of genes that determine transcriptional
identity in adults (*8*). Supporting this view, some transcription factors are associated with specific brain
structures or cephalic compartments, such as *FOXG1* in the telencephalon (**Fig. 1D**) and *OTX2* in non-
telencephalic structures (**Fig. 1D**), show expression restricted to specific clades in the dendrogram.
Hierarchical clustering based only on transcription factor genes was highly similar to the original tree
computed over all expressed genes. However, the tree ordering generated by transcription factors alone
did not produce lower tree distances to the original tree than similarly sized sets of randomly selected
expressed genes (**Fig. S2D**). This suggests that though transcription factors undoubtedly play central roles
in determining cell identity in development, they do not determine global transcriptomic similarities
across neuron types in adulthood.

To determine whether a broadly similar transcriptomically-defined dendrogram is conserved across
species, we repeated the analysis using a prior single cell RNA sequencing dataset of regions sampled
across the mouse brain (*6*) (**Fig. S2C,D**). We found a conserved tree structure in the mouse data,
suggesting broad conservation of the features that drive transcriptomic similarities in neurons across adult
primates and rodents. The relative ordering of the major clades was strikingly similar across species: for
example, in both species, telencephalic glutamatergic neurons are most similar to a clade of GABAergic
neurons that includes striatal spiny projection neurons (SPNs) as well as transcriptionally similar types in
amygdala and hypothalamus (**Fig. S2C**). In both species, all telencephalic GABAergic interneurons

formed a large single clade, and cerebellar types were most similar to a mixed clade containing thalamic
and brainstem types.
170
Borrowing from ancestral state reconstruction methods typically used to estimate evolutionary
divergences between genetic sequences or species (*25*), we applied a maximum likelihood-based
approach (fastAnc) to the expression of all genes at each leaf (observed cell type), along with the branch
lengths between adjacent leaves, in order to reconstruct the transcriptomic state (inferred expression
values of genes) at internal nodes of the marmoset dendrogram. We then compared pairwise "ancestral"
expression values of all genes in the parent nodes for each of the 7 major clades depicted in **Fig. 1C, Fig.**
S2E-F). Transcription factors (TFs) were overrepresented in comparisons of internal nodes that contained
cell types stemming from developmentally related brain regions, but were not overrepresented when the
leaf cell types stemmed from multiple regions (**Fig. S2E-F**). While transcription factor expression reflects
the developmental origins of cell types, and while their expression alone recapitulates the tree structure
seen by including all genes (**Fig. S2D** and (*8*)), some cell types are transcriptionally similar despite having
distinct developmental origins. This may reflect convergence in adult transcriptional profiles (*5, 26*).

Many adult neuronal types are imprinted by their developmental origin
We next inspected RNA expression patterns along with details about each cell type's dissection region-of-
origin to assess which brain structures tended to contain highly similar cell types and which had more
dispersed transcriptional profiles. Some tissues, such as the neocortex, gave rise to cell types that
exclusively clustered within their cephalic domain (**Fig. 1C**), a result unlikely to be driven by ambient
RNA contamination since regionally variable genes were not over-enriched in ambient RNA estimates
(**Fig. S5A**). However, within a given cephalic domain, cell types from distinct brain structures were often
more similar to types sampled from other brain structures. For example, while hippocampal cell types
were all found in telencephalic clades in the dendrogram, many individual hippocampal types were more
similar to types in the amygdala or neocortex than they were to other hippocampal types (**Fig. 1C**).
193
Out of 62 neocortical neuron types, only two types joined clades outside of the major GABAergic and
glutamatergic telencephalic branches: (1) the *MEIS2*+ prefrontal GABAergic types (**Fig. S1C; Fig. 1D;**
Fig. S3), which formed a clade most similar to diencephalic and midbrain *OTX2*+ types, and (2)
Neocortical Cajal-Retzius (C-R) neurons, which were more similar to a clade of *LHX9*+ thalamic neurons
despite indications that they predominantly originate from the cortical hem in primates (*27*) (**Fig. 1D**). As
with hippocampal neurons, neocortical GABAergic neurons neighbored other GABAergic neurons from
the striatum, amygdala and hippocampus that tended to express the same marker. For example, all
PVALB+ types sampled across these telencephalic structures grouped together (**Fig. S3**). Unlike
neocortical GABAergic interneurons, glutamatergic neocortical neurons were broadly most similar to
other telencephalic glutamatergic types, but showed almost no mixing at the terminal leaf level (**Fig. 1C**).
Cerebellar neuron types were entirely restricted to a single, unmixed clade in both mouse and marmoset
(**Fig. S2C**), though this relative isolation could be due to the lack of other hindbrain structures in our
datasets (**Fig. 1C**).
207
Consistent with previous work suggesting that mammalian thalamus contains both midbrain-derived and
forebrain-derived GABAergic interneurons (*28*), we observed distinct clades of thalamic neurons that
were most similar to diencephalic or midbrain populations. Interestingly, thalamic GABAergic neurons

that were *OTX2+* were distinct from other thalamic populations (**Fig. 1D, Fig. S3**), and formed a clade
with other *OTX2+* neurons sampled from brainstem, hypothalamus, and basal forebrain. These
populations were *SOX14+*, while thalamic *SOX14-* populations joined mixed diencephalic-telencephalic
clades (**Fig. 1D; Fig. S3**). The dispersion of thalamic neurons to distinct diencephalic and midbrain-
dominated clades supports recent work suggesting multiple developmental origins for primate thalamic
GABAergic neurons (*28*). We did not find thalamic GABAergic populations that were transcriptionally
similar to telencephalic types (*29*), though we note that the next most proximal clade to the thalamic
OTX2+ types were *MEIS2+* GABAergic neocortical neurons (**Fig. 1D**).
219
Amygdala neuron types were distributed to 5 of the 7 major clades (**Fig. 1C**). Consistent with this widely
dispersed cellular profile, the amygdala is composed of loosely associated nuclei with diverse
phylogenetic and developmental origins, as documented by previous studies (*30*, *31*). Specifically, the
basolateral amygdala, which has a high proportion of excitatory neurons, shares properties with cortical
and claustrum neurons, whereas the intercalated nuclei of the amygdala contain inhibitory *FOXP2+*
projection neurons that share developmental origins with some striatal GABAergic populations such as
striatal spiny projection neurons, SPNs (*5*, *32*, *33*). In our dataset, glutamatergic amygdalar neurons were
most similar to neocortical and hippocampal glutamatergic neurons, while *FOXP2+* GABAergic
amygdalar neurons clustered with SPN types (**Fig. S3**), in line with recent lineage tracing studies in mice
(*5*) and with analysis of fetal macaque types (*33*). Amygdala GABAergic interneuron types had highly
divergent transcriptomic profiles; while some were most similar to telencephalic GABAergic neurons,
others clustered with hypothalamic GABAergic neurons. These results underscore the diversity and
developmental complexity of cell types comprising the amygdala nuclei.

Neurotransmitter usage is not strongly associated with transcriptomic identity
The neurotransmitters used by neurons are central to their function, and neurons are often classified by
their primary neurotransmitter. However, synthesis of the major neurotransmitters – glutamate (in
mammals, excitatory) and GABA or glycine (inhibitory) – requires only a modest number of genes, and
GABA itself is synthesized from glutamate (*34*). It is unclear whether many or few other genes are
correlated with GABAergic (or glutamatergic) identity. For instance, whether neurotransmitter utilization
is strongly associated to the identity of neuron types may differ by brain structure: transcriptomically-
defined neuron types in the neocortex and other telencephalic structures are hierarchically grouped into
GABAergic and glutamatergic types (*2*, *7*, *26*), which reflects both their distinct developmental origins
and their distinct neurotransmitter repertoire. In other brain structures such as the hypothalamus, the
distinction between neurons utilizing GABA or glutamate is much less clear at the transcriptional level (*8*,
35) (**Fig. 1C,E,F, Fig. S3**).
245
To determine whether neurotransmitter identity was associated with a general transcriptional identity
across all neurons expressing the same neurotransmitter, we examined expression of genes encoding the
most prevalent vesicular glutamate transporters (*SLC17A6, SLC17A7*) and GABA synthesis enzymes
(*GAD1, GAD2*) (**Fig. 1C-F; Fig. S3**). If primary neurotransmitter usage was strongly associated with the
expression of many other genes, we would expect that neurons belonging to the same neurotransmitter set
(GABAergic or glutamatergic) would preferentially group together regardless of other factors (such as
developmental origin). However, we did not find evidence for strong global transcriptomic identities of
GABAergic and glutamatergic neurons. Neuronal types from each set were distributed across the tree,

suggesting divergent transcriptomic identities of cell types that share a common neurotransmitter **(Fig.**
1C).

In telencephalic structures such as neocortex, hippocampus, and amygdala, glutamatergic neurons all
express *SLC17A7* and segregate from GABAergic telencephalic neurons (**Fig 1C,D**). Telencephalic
GABAergic neurons separated into distinct clades: one for GABAergic interneurons, and another for
GABAergic projection neurons such as SPNs of the striatum (**Fig 1C, Fig S2**). In non-telencephalic
structures such as the hypothalamus, basal forebrain, thalamus, brainstem, and cerebellum, GABAergic
and glutamatergic types were highly intermixed. This pattern held in a mouse scRNA-seq dataset, and
when using different approaches for hierarchical clustering **(Fig. S2)**. Even within the telencephalon,
neurotransmitter identity did not drive global transcriptional similarity between major clades. For
instance, GABAergic projection neurons, such as SPNs of the striatum, were transcriptionally closer to
glutamatergic neurons in the neocortex, amygdala, and hippocampus than they were to telencephalic
GABAergic interneurons **(Fig. 1C; Fig. S3)**.

Although glutamatergic neurons from distinct cephalic origins do not cluster together, maintaining
glutamatergic neurotransmission or associated function could require a common, core set of genes. To
assess how neurotransmitter utilization relates to genome-wide RNA expression patterns, we examined
distributions of gene-gene correlations across cell types (**Fig. 1E**). Surprisingly few genes are strongly
positively correlated with both *SLC17A6* (VGLUT2) and *SLC17A7* (VGLUT1) expression, even those
associated with glutamate synthesis and packaging **(Fig. S3)**. 116 genes had correlated expression to
SLC17A7 (Spearman's tau > 0.5). The median correlation of *SLC17A6* to those 116 genes was centered at
tau $= 0.05$ (**Fig. 1F**). Only a few genes correlated above 0.5 to both *SLC17A6* and *SLC17A7,* including
BDNF, NRN1, and *TAFA1*. Moreover, only 10 genes (*ARPP21, BDNF, CACNA2D1, CHN1, CHST8,*
CPNE4, LDB2, NRN1, PTPRK, TAFA1) are differentially expressed (> 2.5 fold change) in both
SLC17A6+ glutamatergic neurons and *SLC17A7+* glutamatergic neurons relative to *GAD1+* neurons. We
examined the principal component loadings to see if any were associated with primary neurotransmission.
While individual PCs loaded strongly on specific parts of the tree (**Fig. S2B**), none of the top 20 PCs
(accounting for 92% variance) distinguished cell types by their neurotransmission. The bulk of gene
expression that distinguishes neuronal types from each other appears incidental to neurotransmission.

Neocortical expression fingerprints differ between neurons and glia
The mammalian neocortex is partitioned into functionally, connectionally, and cytoarchitectonically
distinct regions, called areas. We examined regional distinctions in proportions and gene expression of
cell types from 8 cortical locations (**Fig. 2A**). Consistent with prior reports (*3*, *6*, *13*, *36*), cell subtypes
identified in one cortical region were generally present in all other cortical regions, though in different
proportions (**Fig. 2B**). There were two main exceptions in neurons. GABAergic *MEIS2+* cells
(GABAergic cluster 6, **Fig. 2B**) were far overrepresented in PFC samples (specifically in dissections of
medial and medio-orbital PFC, **Fig. S1C**), a compositional distinction not observed in mouse (*6*, *36*). The
second exception was a cluster of *RORB+, KCNH8+* glutamatergic neurons in V1 (and to a lesser extent
V2) that diverged from *RORB+* populations found in the other cortical regions (Glutamatergic cluster 2,
Fig. 2B-D). The expansion and divergence of *RORB+* populations in visual cortex is consistent with the
elaboration and sub-specialization of primary V1 layer IV in primates (*37*). One astrocyte type, marked

by high expression of *VCAN*, was highly enriched in V1 and V2 (Astrocyte cluster 3, **Fig. 2D**). Using
smFISH, we validated higher co-localization of *VCAN* and *GFAP* in V1-adjacent white matter compared
with PFC and V1 gray matter (**Fig. S4I**). *VCAN* is also expressed in OPCs. We did not observe higher
VCAN expression in V1 OPCs, suggesting that the regional variation in *VCAN* expression is specific to
astrocytes (**Fig. 2E**).
302
Glutamatergic neurons sampled from different neocortical locations show regionally distinct gene
expression (*36*). In primates, this is also true of neocortical GABAergic neurons (*3*). Astrocytes in mouse
and primate brains are heterogeneous across major subdivisions (**Fig. S4D-H**), but the extent to which
they are locally customized in distinct regions of neocortex is less well understood (*2, 7, 26, 38*)(*4*).
Studies in mice have revealed layer-specific astrocyte subpopulations in the cortex (*39*) and variation
between neocortical and hippocampal astrocytes (*40*).
309
To address whether cortical regional variation in gene expression is shared across cell types, we
performed pairwise comparisons between major clusters of cortical excitatory neurons, inhibitory
neurons, astrocytes and oligodendrocyte lineage types across all eight neocortical locations (**Fig. 2F,**
Table S3). Each of these cell classes displayed regionally differentially expressed genes (rDEGs) across
neocortical regions (**Fig. 2F**), an effect that could not be attributed to ambient RNA contamination (**Fig.**
S5A). Similar to what has been described in the mouse cortex (*39, 40*), astrocytes within the marmoset
cortex exhibited regional transcriptional variation, but overall neurons had more rDEGs than macroglia
(**Fig. 2F; Fig. S5B**). 62% of rDEGs in interneurons were also rDEGs in excitatory neurons. Interestingly,
though cortical astrocytes and oligodendrocytes arise from a common lineage with cortical excitatory
neurons (*41*), they shared a lower percentage of rDEGs in common with excitatory neurons as compared
with interneurons (25% astrocytes, 25% oligodendrocyte lineage). Regionally differentially expressed
genes within a cell class (glutamatergic, GABAergic, astrocyte, oligodendrocyte lineage) tend to be
biased in the same regions across subtypes within that class. However, certain subtypes and regions
accumulated more rDEGs than others. For example, across all cell types and particularly within neurons,
higher-order temporal association cortex and prefrontal cortex tended to be most distinct from V1 and V2
(**Fig. S5B**). To determine the extent to which rDEGs are private to individual donors or represent shared
features of variability within conserved cell types, we analyzed the consistency of neocortical rDEGs
across the three donors that had been sampled at all eight neocortical locations (**Table S3**). Our findings
revealed that 55% and 39% of glutamatergic and GABAergic rDEGs, respectively, were consistent
among donors, while only 24% and 14% of astrocyte and oligodendrocyte rDEGs, respectively, were
shared between donors. These results indicate lower inter-individual consistency in regional gene
expression signatures in non-neuronal cells.

A hypothalamic-origin neuron type in medial amygdala
In mammalian CNS, some cell types migrate long distances from proliferative zones to their mature
destinations (*42, 43*). However, neurons generally tend to respect cephalic boundaries and remain within
the same subdivision as their progenitors (*44*). This tight control over migration potential makes it
difficult to disentangle the persistent influence of developmental origin from potential later influences
arising from shared tissue context that might affect all neurons similarly in a given brain structure.
However, cephalic boundary crossings do exist (*28, 29, 45*). Though rare, such boundary crossing events

can reveal whether cell types that embarked on cross-cephalic migration retain transcriptomic profiles
more in common with their tissues of origin, or more in common with their final destinations.
341
We found a striking example of cross-cephalic migration in the amygdala. First, we observed an
unexpected clustering pattern in our analysis, wherein three amygdala neuron types joined a clade not
with other telencephalic neurons, but instead with *SLC17A6+* hypothalamic and basal forebrain types
(**Fig. 3A**). Moreover, despite expressing *SLC17A6* and lacking expression of *GAD1* and *GAD2,* these
cells showed a closer association with GABAergic rather than glutamatergic types (**Fig. 3B**). They did not
express other genes required for GABAergic transmission such as *SLC32A1* (VGAT) and also lacked the
molecular machinery for non-canonical GABA reuptake or release observed in other cell populations
(**Fig. S3**)(*46*, *47*). These findings suggest that these particular amygdalar neurons exhibit a "cryptic"
transcriptomic identity, in which they are glutamatergic but have transcriptomic profiles that are much
more similar to GABAergic types, relative to other telencephalic neurons.
352
The "cryptic" amygdalar subtypes display additional atypical gene expression features compared to other
amygdala neuronal types, including expression of *OTP* and *SIM1* (**Fig. 3B**). These transcription factors
are typically expressed in neuronal lineages originating from proliferative zones around the third
ventricle. In mice, there is a migratory stream of diencephalic neurons into the telencephalon (*45*, *48*) into
the medial amygdala, and this migration is dependent on *Otp* expression (*45*) (**Fig. 3D**). We generated
mouse amygdala snRNA-seq data from 53,745 nuclei and verified the presence of a *Sim1+* neuron type in
mice that also clusters with GABAergic neurons and expresses *Slc17a6* but not *Gad1* or *Gad2* (**Fig. 3C**).
In mice, these neurons constitute the majority population in the amygdala that express *Adcyap1* (encoding
the protein PACAP) (**Fig. 3C**), a neuropeptide that is extensively (but not exclusively) expressed in
hypothalamic populations (*49*) and associated with energy homeostasis (*50*), stress, anxiety (*51*), and
immune responses (*52*). (In marmosets, *ADCYAP1* is additionally expressed in subsets of *SLC17A7+*
neurons that do not share the "cryptic" phenotype, indicating that *ADCYAP1* has a distinct distribution of
expression in amygdala neurons between mice and marmosets.)
366
We further confirmed the expression of *Sim1+* cells in the medial amygdala in mice using the Allen
Institute ISH atlas (**Fig. 3E**). To precisely locate the specific amygdalar nuclei housing the cryptic
population, we investigated the expression of *SIM1* in neonatal marmosets (*53*, *54*). *SIM1* expression was
highly enriched in the neonatal marmoset medial amygdala (MeA) (https://gene-atlas.brainminds.jp; **Fig.**
3F), which mirrors the migration of *Sim1+* neurons from the diencephalon in mice (28). We verified
SIM1 and *SLC17A6* but not *SIM1* and *GAD1* colocalization in adult marmoset medial amygdala using
single-molecule fluorescence *in situ* hybridization (smFISH) (**Fig. 3G**). These results suggest that the
cryptic amygdala neurons are a conserved population in both mice and primates that likely have
diencephalic origins. Consistent with a "birthplace imprinting" effect, they retain transcriptional identities
more similar to diencephalic types than to the telencephalic types with which they ultimately reside (**Fig.**
3A).

Primate-specific striatal TAC3+ interneurons are similar to specific TAC3+ diencephalic types
Previously, we discovered a *TAC3+, LHX6+* interneuron subtype in the striatum of humans, macaques,
and marmosets that was absent in mice and ferrets (*3*). Compared with other striatal types, they are
transcriptionally most similar to *PVALB+* interneurons and, because they expressed transcription factor

LHX6, we surmised that they likely also arose from the medial ganglion eminences (MGE) (*3*). Consistent
with this inference, a recent study of fetal macaque single nucleus RNA-seq data from ventral
telencephalic progenitor domains found that *TAC3*+ striatal interneurons likely arise from progenitors in
the MGE, and diverge from an ancestral progenitor class that also gives rise to conserved *MAF*+
progenitors that produce *PVALB*+ and *TH*+ striatal interneurons (*33*).

The broader census of brain structures in the current dataset allowed us to compare the transcriptional
identity of the striatal *TAC3*+ type to cell types outside of the striatum. Beyond expression in the striatal
type, *TAC3* is expressed in 20 different neuron types in our dataset, including expression in cortical
GABAergic neurons as well as several amygdala, basal forebrain, thalamic, and hippocampal types (**Fig.
3H; Fig. S3, Fig 6A,B**). *TAC3*+ types did not form a single clade in the dendrogram, and their
transcriptional resemblance to other neuron types largely reflected their tissue or cephalic origin. For
example, thalamic *TAC3*+ subtypes were found in a thalamic-only clade, while hippocampal *TAC3*+
neurons were most similar to other hippocampal and amygdala types. The thalamic types were *SLC17A6*+
while all other *TAC3*+ types were GABAergic.

Unexpectedly, the striatal *TAC3*+ type was most similar not to other striatal interneuron types as we
previously concluded (*3*), but rather to two other *TAC3*+ GABAergic types found in basal forebrain and
hypothalamus (**Fig. 3H**, tissue validation in **Fig. S6A**, **Data S2**). The broader clade containing these three
types (depicted by green arrowhead in **Fig. 3H**) consisted entirely of basal forebrain and hypothalamic
types, with the exception of the striatal *TAC3*+ type. Each of the similar *TAC3*+ populations had distinct
gene expression (such as high expression of OXT and AVP in the hypothalamic type, and DRD2
expression exclusively in the striatal type), ruling out dissection contamination (**Fig. 3I**). The 3 types
remained direct neighbors when the dendrogram was recomputed using other distance metrics (e.g.
correlation-based or PCA scores using transcription factor expression, see **Fig. S2**). As this clade
assignment was unexpected, we omitted the basal forebrain and hypothalamic *TAC3*+ types and
recomputed the dendrogram (retaining 286 of the original 288 neuronal types) to determine whether the
striatal *TAC3*+ was broadly more similar to hypothalamic types than to telencephalic types (**Fig. 6C**). As
striatal interneurons are generally divergent from hypothalamic/basal forebrain types, omitting these two
cell types enabled us to test whether the *TAC3*+ striatal type is broadly more similar to hypothalamic
neurons or to striatal neurons. The global structure of the dendrogram was essentially the same as the
original, except that the striatal *TAC3*+ type now joined the major telencephalic GABAergic clade with
other striatal interneurons (neighboring the striatal *PVALB*+ subtype) (**Fig. 6C**), recapitulating our
original similarity assignment (*3*).

Considering their unexpected transcriptional similarity to both a telencephalic (basal forebrain) and a
diencephalic (hypothalamus) type, the *TAC3*+ striatal type may either arise from a telencephalic
progenitor (*33*) that also gives rise to sister diencephalic (hypothalamic) types, or else shows striking
transcriptional convergence with diencephalic types that have distinct developmental origins. Favoring a
telencephalic origin as suggested by previous report (*33*), the hypothalamic, basal forebrain and striatal
TAC3+ types are all *FOXG1*+, a transcription factor associated with telencephalic origin (**Fig 3I**). They
also express *LHX6*+ and *NKX2-1*+ (**Fig S3**), consistent with a medial ganglionic eminence origin,
however we note that in mice some hypothalamic types also express *Nkx2-1* and *Lhx6* (*9, 55, 56*).

Ultimately, lineage tracing of the striatal *TAC3+* type in a primate would resolve whether a shared
progenitor gives rise to both telencephalic and diencephalic types.

Quantitative maps of interneuron distribution across primate neocortex and striatum
Our analysis of RNA expression in the single nucleus dataset indicates that developmentally-linked
telencephalic GABAergic populations (*42*, *43*) retain globally similar identities in adulthood. For
example, *SST+* striatal interneurons are more similar to *SST+* hippocampal and neocortical interneurons
than they are to other striatal types (**Fig. S3**). Within each brain structure, developmentally linked
populations become functionally specialized and follow distinct spatial rules for their allocation. In the
mouse, quantitative imaging of molecularly identified interneuron types has revealed that densities of
Sst+, *Vip+*, and *Pvalb+* cortical interneurons vary across the cortical mantle; their relative proportions
relate to unique functional and microcircuit properties of different cortical areas (*15*). In primates, overall
neuron densities are more variable by cortical location than in mice: they vary by as much as 5 fold across
the cortical sheet, with highest neuron proportions and densities found in occipital cortex and particularly
in V1 (*57*). However, quantitative mapping of molecularly-defined subtypes of neurons have not been
performed in a primate, and it is not known if they have conserved or distinct spatial distributions
compared to mice.
441
We used smFISH to image the major cortical and striatal interneuron types across sagittal sections in
marmoset (**Fig. 4** and **Fig. S7-9**). We quantified proportions and densities of each type relative to all cells
(DAPI; **Fig. S7**). In the neocortex, we binned these using an areal parcellation of marmoset neocortex
(https://doi.org/10.24475/bma.4520) to determine whether interneuron proportions varied systematically
by brain area. In total, we quantified 377,554 neocortical interneurons across 30 sections by smFISH. In
striatum, we quantified 6,848 interneurons across 32 sections. Each series sampled sagittal sections ~160
µm apart, beginning 1,184-1,584 µm from the midline up to 6,384 µm laterally, including the majority of
the striatum with the exception of the most-lateral portion of the putamen. We used a cell segmentation
algorithm to count positive cells across sagittal sections and expressed interneuron proportions as a
percent of all cells (DAPI+).
452
Neocortical types were identified with probes for *SST, PVALB, CXCL14, VIP,* and *LAMP5* (**Table S4**),
which collectively account for all major cortical interneuron populations (**Fig. 4A**). *CXCL14* is a marker
for caudal ganglionic eminence derived cortical interneurons. It is expressed in most *VIP+* and *LAMP5+*
cortical neurons, as well as a smaller population of *VIP-, LAMP5-* types, some of which are *PAX6+* (and
which correspond to the *SNCG+* population in humans and mice (*2*, *3*))). *VIP+* and *LAMP5+*
interneurons are the two other major CGE-derived populations present in primates and mice. As *LAMP5*
is also expressed in subsets of excitatory neurons, we performed dual labeling smFISH with *GAD1* to
avoid counting glutamatergic types. Major striatal interneuron types were identified with probes for *SST,*
PVALB, CHAT/SLC5A7, TH, CCK, and *TAC3* (**Table S4**). These together account for most major
populations of non-SPN neurons in the striatum, with the exception of a population of *MEIS2+*
GABAergic striatal neurons that cluster together with non-SPN GABAergic neurons (**Fig. 1C, Fig. S3**)
but are difficult to distinguish uniquely as several other markers are also expressed at variable levels in
other striatal cell types.

Neocortical interneuron types have highly focal biodistributions

Quantitative analysis of smFISH revealed highly focal and variable distributions of interneuron subtypes across marmoset neocortex (**Fig. 4B-E, Fig. S8A-D**). In both absolute numbers and relative proportions, *PVALB+* interneurons were strongly enriched in the occipital lobe, particularly along the calcarine sulcus in the medial sections, as well as the occipital pole more laterally (**Fig. 4B-C; Fig. S8A, Data S1**). *SST+* interneurons in the neocortex increase medio-laterally (**Fig. 4B-C, Fig. S8A, Data S1**), but closer inspection revealed this is driven not by a spatial gradient so much as by highly focal enrichments around primary motor area (M1) and primary somatosensory cortex (S3, S1/2), the cingulate cortex, entorhinal cortex and medial prefrontal cortex (**Fig. 4C**). *CXCL14+* neurons are enriched along the calcarine sulcus medially, as well as in ventral aspects of the occipital cortex more laterally. There are higher proportions dorsomedially in the parietal cortex (**Fig. 4C**). *VIP+* neurons were enriched in PFC and also increased laterally at or near somatosensory cortex and posterior parietal cortex (**Fig. 4C**). *LAMP5+* interneurons showed a bias to the top of the cortical layers, consistent with dominant composition of this class as neurogliaform Layer 1 types (*3, 58, 59*), though this class also contains the *LAMP5/LHX6* type that is found in deeper layers (*3, 13*) (**Fig. 4C**).

To better appreciate how these distributions relate to neocortical areas, we used a histologically-based marmoset neocortical parcellation (https://doi.org/10.24475/bma.4520; **Fig. S8B-C**) to bin smFISH interneuron proportions by cortical area (**Fig. 4D-E; Fig. S8B**). Small adjacent areas were merged, resulting in 26 areal groupings (**Fig. S8C**). Overall interneuron proportions relative to all cells varied by 4.5-fold across areal groupings (**Fig. 4D-E, Fig. S8B-C**). As a fraction of all cells, A31 (dorsal posterior cingulate) had the lowest overall proportion of interneurons (3%), while insular cortex (In) had the highest (17%) (**Fig. 4D**).

In mice, quantitative mapping of interneuron densities showed higher *Sst+* densities and lower *Pvalb+* densities in neocortical areas involved in higher cognitive functions, such as medial frontal and lateral association areas (*15*). This local circuit feature follows cortico-cortical connectivity network topography in mouse (*60*). In contrast, lower *Sst/Pvalb* ratios were associated with mouse primary motor and sensory areas, which are associated with less distributed (and more local) cortical connectivity (*15*). To determine whether primate neocortex followed similar rules as mouse neocortex of interneuron allocation, we examined normalized proportions of the four largely mutually exclusive interneuron classes (*PVALB+, SST+, LAMP5+, VIP+*) within our marmoset areal groupings (**Fig. 4D-E; Fig. S8B**). Lateral temporal cortex, including A36, had the highest *SST/PVALB* ratio. Other areas with high *SST/PVALB* ratios included piriform cortex (Pir), M1 and several medial/orbitofrontal areas (A25, A13, A32). This suggests that some marmoset association areas, notably medial frontal area and lateral temporal cortex, have high *SST/PVALB* ratio composition consistent with high *Sst/Pvalb* ratios in higher order association network in mice (*15*) (**Fig S8B**).

Strikingly, polar and lateral prefrontal areas (including A8, A46, A10, A47, A45), which are thought to be the most divergent relative to prefrontal areas in rodents (*61*), and which in primates are characterized by long-range cortico-cortical connectivity to other association areas (*62–64*), do not exhibit high *SST/PVALB* ratios (**Fig. 4D-E; Fig. S8B**). Of all cortical areal groupings we measured, the lowest *SST/PVALB* ratio was found in A45, a higher-order lateral prefrontal area with extensive projections to all lobes of the neocortex (*64*); http://analysis.marmosetbrain.org/). Thus, unlike the high *Sst/Pvalb* ratios observed in frontal areas in mouse (*15*), primate lateral prefrontal areas are characterized not by

exceptional *SST/PVALB* ratios but rather as having some of the highest *VIP+* interneuron proportions
(**Fig. 4D-E; Fig. S8B**). Lateral prefrontal areas also have the highest total fraction of *VIP+* and
LAMP5/GAD1+ interneurons (all above 50% of all interneurons); both of these populations
predominately arise from the caudal ganglionic eminence, a progenitor zone that has expanded in primate
evolution (*65*). These results suggest that primate lateral prefrontal areas do not follow the same local
cortical circuit organizing principles that typify frontal areas in the mouse (*15*), or medial frontal areas in
marmoset (**Fig. 4D-E; Fig. S8B**).

Interneuron proportions in marmoset striatum follow medio-lateral gradients
The striatum, a crucial brain region involved in motor control, reward, and decision-making, displays
complex functional topography and connectivity. In humans, anterio-medial portions containing nucleus
accumbens/ventral striatum are functionally coupled to limbic and higher order cognitive networks, while
lateral subdivisions are more coupled to sensory and motor networks (*66*). Prior work relating bulk gene
expression measurements in human and macaque striatum to cortico-striatal network organization found a
relationship between functional domain (e.g. somato/motor vs limbic network) and gene expression,
including enrichment of *PVALB* in lateral portions of the striatum that are coupled to somatosensory and
motor cortex (*67*). These results suggest that differences in local cell type composition across striatum
may underlie aspects of its functional topography.
528
In mice, striatal interneuron subtypes have different spatial distributions: cholinergic (*Chat+*) neuron
proportions increase dorsally and anteriorly (*68*), *Pvalb+* interneurons are more abundant in dorsolateral
striatum than in dorsomedial striatum, and *Sst+* interneurons are spatially homogeneous (*69*) (**Fig. S9A**).
Primates retain the major populations of striatal interneurons found in mice (*6*, *70*), and additionally have
gained the novel type distinguished by *TAC3* expression (*3*, *33*) described in previous sections (**Fig. 3H,**
4F). A systematic quantification of striatal interneuron types has not been performed comprehensively in
a primate, and it is unknown if they follow similar or distinct spatial distributions observed in mice. We
used single-molecule FISH (smFISH) to investigate distributions of the major types of conserved striatal
interneurons (*SST+*, *PVALB+*, *SLC5A7+/CHAT+*, *TH+*, *CCK+*, *TAC3+*, probes in **Table S4**) in serial
sagittal sections of marmoset striatum (**Fig. 4G-H**).
539
Each striatal interneuron population exhibited a non-uniform distribution across the marmoset striatum,
particularly in the medial-lateral axis. Similar to mice, the proportion of striatal *PVALB+* interneurons
increases in lateral sections, from ~0% to 0.8% of all cells (**Fig. 4G, Data S1**). Unlike mice, marmoset
SST+ interneuron distribution is non-uniform, appearing sparse near the midline and increasing in
proportion (0.1-0.7%; **Fig. 4G, Data S1**). Cholinergic neurons (*CHAT+*) show the opposite medial-lateral
gradient (0.45%-0.1%; **Fig. 4G, Data S1**). Similar to *CHAT+* neurons, *TH+* striatal interneurons, which
are transcriptionally similar to the *PVALB+* type, exhibited a decreasing medial-lateral gradient (**Fig. 4G,**
Data S1). *CCK+* striatal interneurons, which are a minority population in marmoset (*3*) and mouse (*70*)
are enriched close to the midline and become much sparser laterally (**Fig. 4G, Data S1**). The *TAC3+*
interneurons showed an increasing medial-lateral gradient, similar to *CHAT+* neurons (2%-0.5%; **Fig.**
4G, Data S1). No striatal population exhibited anterior-posterior or dorsal-ventral gradients with the
exception of *PVALB* interneurons, which showed a modest dorsal-ventral gradient. Unlike striatal
interneurons, MSNs proportions were largely uniform across the major axes: *DRD1* and *DRD2*, which
distinguish direct and indirect MSNs, respectively, were slightly enriched medially but otherwise

exhibited largely uniform distributions across the striatum (**Fig. S9B**). Overall, these findings suggest that
the spatial distribution of interneuron subtypes within the primate striatum may play a role in shaping its
functional topography and connectivity. While mice and primates share similar striatal interneuron
populations, there are also differences in their spatial distributions.

Morphology of conserved neocortical and striatal interneuron types
The morphology of interneuron types relates essentially to their function and contributions to neural
circuits. While methods such as biocytin filling and Golgi staining are the gold standard for
morphological reconstructions, the administration of low-titer AAVs carrying membrane-bound
fluorescent reporters can be used to sparsely label cells for morphological reconstruction. We utilized a
reporter AAV under the control of the forebrain interneuron-specific mDlx enhancer (*19*) to label
neocortical and striatal interneurons, and then performed smFISH (probes in **Table S4**) on thick sections
(120 µm) to confirm the molecular identity of GFP+ cells. Reconstructions using viral labeling are more
challenging than with single cell filling methods, because GFP expression from neighboring cells and
passing fibers have to be distinguished from signal attributable to the target cell. In some cases, the GFP
signal appears punctate, making it challenging to follow discontinuous processes. For these reasons, our
reconstructions are conservative: as we aimed to avoid reconstructing false positive fibers (fibers
originating from other cells), we may in some cases under-ascertain the full dendritic arborization of
target cells.
572
We imaged 1,203 GFP+ neurons in the striatum and 4,374 GFP+ neurons in the neocortex. We used
NeuTube to reconstruct the top 216 telencephalic neurons that had the best GFP signal, did not have other
cells labeled in the field of view, and were positive for at least one smFISH probe **(Table S5; examples**
in Fig. S10). Raw image stacks and NeuTube reconstructions are available at
https://doi.org/10.35077/g.609. Using combinations of 1-2 probes for marker genes of different types, we
identified GFP+ neocortical and striatal interneurons, respectively, with smFISH based on their type
markers (neocortex: *SST*, *PVALB*, *CXCL14*, *VIP*, *LAMP5*; striatum: *SST*, *PVALB*, *SLC5A7*, *TH*, *CCK*,
Table S4) (**Fig. 5A-B,** *top rows*). We used a second tracing and reconstruction method (Imaris) on the 41
most complete GFP+ cells. Of these, we discarded 5 due to incomplete soma or highly discontinuous
processes in the image stack, retaining 36 cells (**Fig. 5A-B,** *bottom rows*; **Table S6**). Morphological
parameters were measured using the Surface function, which detects surface area and volume based on
the fluorescence of the mDlx-AAV-GFP expression, and the Filament Tracer function, which traces
structural features starting from the soma to the terminal processes based on the diameter of the soma and
the thinnest projection of the cell (**Table S6**). To assess whether there are region-dependent
morphological differences within molecularly similar types, we compared several parameters between a
collection of neocortical and striatal *PVALB*+ GFP+ cells that were reconstructed from the same
marmoset (Cj 17-154; **Fig. 5C-D**) to avoid variable tissue shrinkage arising from different tissue storage
conditions across the animals (see *Methods*). Striatal *PVALB*+ GFP+ cells were larger than cortical
PVALB+ GFP+ cells in terms of length, area, but not soma diameter or the volume of GFP+ fluorescence
(**Fig. 5D**). While the number of branches issued by their somas also did not differ, the striatal *PVALB*+
GFP+ interneurons exhibited more dendritic branch points than their cortical counterparts (**Fig. 5D**),
suggesting that striatal *PVALB*+ interneurons have greater arborization than *PVALB*+ cortical
interneurons.

 Development of a novel TAC3-rAAV-GFP reporter

Given that the *TAC3+* type comprises almost 30% of striatal interneurons in marmoset, we expected a
sizable proportion of striatal GFP+ cells labeled by AAV9-hDlx5/6-GFP-fGFP (**Fig. 5**) to be *TAC3+*.
However, while 55 GFP+ striatal cells were imaged across all smFISH *TAC3*-probe treated sections, we
failed to find any colocalization with *TAC3* transcripts. In these experiments, which were double-labeled
with *TAC3* and *PVALB* probes, 9/55 were *PVALB*+ (16% *PVALB*+, vs expected 21% of interneurons
expected from snRNA-seq proportions), while 0/55 were *TAC3*+ (0% vs 28% expected from snRNA-
seq).

The mDlx enhancer is a regulatory element specific to forebrain interneurons (*71*). Though the regulatory
element, and the forebrain-interneuron expressed genes flanking it, *Dlx5* and *Dlx6*, are highly conserved
in vertebrates, we wondered whether the lack of accessibility of the mDlx locus in the *TAC3* interneurons
could explain our inability to find colocalization of *TAC3* expression and GFP. To assess this, we
generated single nucleus ATAC-seq (snATAC-seq) data (69,808 nuclei) from fresh marmoset striatum
(**Fig. 6A**). We used Signac (*72*) to integrate our previously annotated striatal snRNA-seq data and identify
major striatal types. We then examined accessibility of the marmoset sequence homologous to the mDlx
locus across interneuron types. While other striatal interneuron types (particularly the *SST*+ and *PVALB*+
types) showed accessibility (ATAC-seq peaks, reflecting chromatin accessibility) at the mDlx locus, the
TAC3+ type did not (**Fig. 6B**).

To develop a viral tool that could transduce the striatal *TAC3+* cell type, we next nominated new
candidate regulatory elements specific to the *TAC3+* interneuron type. We used Signac to identify
differentially accessible peaks in the *TAC3+* cluster relative to all others, and filtered the set by fold
change, percent accessibility across the target cell type population, and peak size (**Fig. 6C**). To maximize
the likelihood of obtaining a functional reporter while minimizing the number of marmosets used, and
because several of our top candidates were very short, we selected four top regulatory element candidates
for the *TAC3+* type for tandem packaging (**see *Methods***). The four candidates were on four different
chromosomes and spanned between 94-215 bp. One site was in an exon of *CDH13*, one was in the first
exon of *TAC3*, one was in an exon of *LOC108592287*, and one was intergenic (closest gene 200kb
distance). These four elements were packaged in tandem in an AAV9 vector (AAV9-tandemE-TAC3-
EGFP**)** containing a cytoplasmic GFP reporter (**Fig. 6D**). We delivered the virus via MRI-guided local
injection into the anterior striatum of two marmosets (Cj 19-207 and 17-B111, **Fig. 6E and Fig. S11A,**
Table S1), and imaged coronal sections of striatum for GFP positive cells after 6 & 10 weeks' incubation
time (**Fig. 6F and S11C**). smFISH confirmed colocalization of *TAC3* transcripts in GFP+ neurons in
striatum (**Fig. 6G and Fig. S11B,D**). In one animal the virus diffused beyond the boundary of the
striatum. In this animal we also detected strongly labeled GFP+ cells in a border zone between striatum,
BNST, and the globus pallidus where the viral injection had diffused beyond striatum (**Fig. S11B-E**), and
smFISH confirmed that these too were *TAC3+*.

To determine the broader biodistribution of cells transduced by the tandemE-TAC3 enhancer, we next
packaged the same enhancers in a novel AAV capsid (BI103) capable of efficient transduction of brain
cell types after systemic IV delivery in marmosets (Chan et al, in prep). We delivered this AAV (AAV-
BI103-tandemE-TAC3-EGFP) to one marmoset (Cj 20-214, **Table S1**). After a 4 week incubation, the
brain was perfused, sliced and stained with anti-GFP antibody to amplify GFP signal, and smFISH probes

against *TAC3+* to confirm colocalization. We found GFP+/*TAC3+* colocalized cells in striatum as well as
several extra-striatal locations including hypothalamus, substantia nigra, superior colliculus, brainstem,
and neocortex (**Fig. 6I**). To assess the morphology of *TAC3+* striatal interneurons produced by viral
labeling, we reconstructed several of the most complete cells from both the local injections (AAV9-
tandemE-TAC3-EGFP) and from the systemic injection (AAV-BI103-tandemE-TAC3-EGFP) using
Imaris (**Figs. 6H,J,K** and **Table S6**). The cells tended to have 2-3 thick branches that extended from the
soma, and which bifurcated close to the soma and became thinner thereafter. Reconstructed cells had
median 13.1 µm (+/- 5.1 s.t.d) soma diameter, 2221.3 volume (µm^3; +/- 1192.8 s.t.d) and 16.5 (+/- 6.11)
total dendritic branch points. These experiments show that whereas systemic injections of AAV9 under
the mDlx enhancer could not transduce the striatal *TAC3+* type, presumably due to loss of accessibility at
the mDlx locus (**Fig. 6B**), local injections of AAV9 as well as systemic injections with alternative capsid
with the cell type targeted enhancers were successful. Broadly, the use of cell type specific regulatory
elements, coupled with viral engineering, enables a new horizon for the study of primate brain cell types.

Discussion

In this study, we generated a molecular and cellular census of the adult marmoset brain using single-
nucleus RNA sequencing, quantitative smFISH, and cell type specific AAV labeling. Our study reveals
the complex repertoire of cell types in the marmoset brain. Our snRNA-seq dataset of over 2.4 million
brain cells across 18 brain regions in the marmoset indicates that lineage is an important factor shaping
adult transcriptomic identity of neuronal types, apparently more so than neurotransmitter utilization.
Using quantitative smFISH we revealed, for the first time in a primate, the spatial distributions of
molecularly-resolved GABAergic interneuron types. Using GFP delivered by interneuron-specific AAVs,
we generated morphological reconstructions of all major interneuron types in the neocortex and striatum.
We generated a novel viral genetic tool under the control of cell type specific regulatory elements to
transduce a previously described putative primate-specific striatal interneuron type. These datasets,
generated as part of the BRAIN Inititive Cell Census Network, complement other recent and extensive
cellular profiling studies in other species (*8–11*, *73*) and will enable insights in cell type innovations and
modifications in future comparative studies.

Telencephalic glutamatergic and GABAergic neurons strongly segregate in mammals as well as in
homologous structures in amphibians (*26*), suggesting an evolutionarily conserved distinction. An initial
atlas in mouse indicated that glutamatergic and GABAergic neurons from diencephalic and midbrain
structures also partition almost perfectly by neurotransmitter usage (*7*), suggesting this could be a general
rule. However, more recent transcriptomic censuses in mouse (*8*, *9*) and human (*11*) indicate that across
mammalian species, many non-telencephalic glutamatergic and GABAergic types tend to form highly
intermixed clades that do not separate clearly by neurotransmitter identity. Our results in the adult
marmoset (and mouse, **Fig S2**) concord with the notion that gene expression distinctions between
telencephalic glutamatergic and GABAergic neurons do not hold for neurons in other brain structures
(**Fig. 1; Fig. S2**). This has implications for generalizing transcriptomic associations to other phenotypes.
For example, transcriptomic changes in glutamatergic or GABAergic neurons have been associated to
diseases such as autism and schizophrenia (*74–76*). Such associations may not generalize to GABAergic
or glutamatergic types outside of the sampled brain region (usually neocortex), consistent with
observations that "global" changes to glutamatergic or GABAergic neurons in relation to disease actually
often only surface in a few brain regions (*77*).

A previous analysis, based on shared patterns of several key transcription factors, proposed that
telencephalic GABAergic neurons are developmentally and evolutionarily related to diencephalic
GABAergic neurons (*78*). Our results indicate that when profiled at adulthood, only a limited number of
telencephalic GABAergic types are transcriptionally similar to diencephalic types, some of which may
arise from cephalic boundary crossings. Most neocortical, hippocampal, and some amygdalar and striatal
GABAergic types are so distinct from diencephalic GABAergic types that they share more gene
expression in common with telencephalic glutamatergic types (**Fig. 1C**).

Developmental origin or shared lineage plays a strong role in shaping the adult transcriptomic identity of
neurons, but phenotypic convergence, whereby adult cell types converge on similar transcriptomic
identities despite a non-shared developmental origin, may also drive apparent similarities amongst cell
types. Recent advances in lineage tracing coupled with single cell RNA sequencing demonstrates that
phenotypic convergence is surprisingly common between transcriptomically defined types (*5, 79*). As
such, it is difficult to disambiguate between these possibilities in absence of data that confirm lineage
directly (*5, 26*). For example, in our data the clade of GABAergic projection neurons that contains striatal
SPNs and amygdala and basal forebrain *GAD1+, FOXP2+* neurons also contained several subtypes of
hypothalamic *GAD1+, FOXP2+* neurons (**Fig. 1C; Fig. S2E, Fig. S3**). Either these hypothalamic
FOXP2+ subtypes have a convergent expression identity with long-range GABAergic projection neurons
of the telencephalon, or else they arise from a common lineage.

Our data support known instances of cross-cephalic migrations in thalamus and amygdala, and also
suggest new ones (**Fig. 3**). The similarity of the primate-specific *TAC3+* striatal type to hypothalamic and
basal forebrain *TAC3+* types in particular is unexpected (**Fig. 3H**) and suggests a potential example of
cross-cephalic vesicle migration. Recent work (*33*) proposes that the striatal *TAC3+* type has a ventral
ventricular telencephalic origin, similar to most other GABAergic interneuron types destined for striatum.
While phenotypic convergence remains possible, another possibility is that a ventral telencephalic
progenitor gives rise to both telencephalic and diencephalic types. This is supported by the expression of
FOXG1, a transcription factor necessary for ventral telencephalic fate (*80*), in all three transcriptionally
similar *TAC3+* populations.

Developmentally linked interneuron populations in striatum and neocortex (e.g *PVALB+* types in both
structures) displayed distinct spatial distributions measured by cell counting of smFISH (**Fig. 4**). While
all interneuron subtype distributions in the striatum followed a gradient along a medial-lateral axis, the
interneurons in the neocortex followed much more complex distributions that for the most part were not
captured by simple gradients. While in the mouse, *Sst+/Pvalb+* ratios are a hallmark of higher order
cortex (*15*), we found a different pattern of relative interneuron proportions across much of the higher-
order association cortex in marmosets. Areas in the lateral prefrontal cortex were unique in their relatively
high proportions of *VIP+* neurons (and had unexceptional or even low ratios of *SST+/PVALB+* neurons).
These results suggest that primate lateral prefrontal areas may not follow the same local cortical circuit
organizing principles of association cortex in the mouse.

Morphological characterization of the striatal and neocortical interneuron populations suggests variation
in overall size and dendritic arborization amongst subtypes. To avoid interindividual and technical
variability arising from different sample storage conditions (see Methods), we analyzed several
morphological parameters in a subset of cortical and striatal *PVALB+* interneurons reconstructed from a
single marmoset (**Figs. 5C-D**). Striatal *PVALB+* cells were found to be larger in length, volume, and
surface area and consisted of more dendritic branch points than the cortical cells. Our data altogether

suggest that the striatal *PVALB+* interneurons exhibit higher dendritic complexity compared to their
respective counterparts in the neocortex. These morphological differences could result in differing
electrophysiological properties. For example, it is possible that *PVALB+* cells in the striatum have higher
capacitance and receive more synaptic inputs onto them, thereby impacting local signal integration and
computation, than cortical *PVALB+* interneurons. The functional significance of morphological
differences across the interneuron subtypes identified in our study will ultimately require studying how
these cells affect the circuits in which they reside (i.e., cellular/subcellular targeting biases and functional
properties)(*81–83*).

The development of virally-based tools enables cell-type-specific access in nonhuman primates (*2, 18–*
20). To maximize translatability across species, most approaches nominate candidates using evolutionary
conservation (usually between mice and humans) as a selection criterion. However, the evolutionarily
conserved mDlx enhancer did not drive expression in the novel striatal *TAC3+* type **(Fig. 6B)**. In general,
we observed that the mDlx enhancer selectively under-ascertained several interneuron populations in
marmoset in these experiments. For example, it systematically under-labeled *VIP+* and *SST+* types in the
neocortex (expected 22% and 26% of interneurons, obtained 2% and 3%, respectively), as well as *SST+*
interneurons in the striatum (14%, obtained 1.8%). This underlabeling could in part be due to the titer and
systemic delivery approach we adopted, which was necessary in order to achieve sparse labeling for
morphology. The overall low efficiency in our ability to molecularly characterize GFP positive cells
suggests a need for further optimization for this species and application.

The lack of transduction of *TAC3+* striatal interneurons using the AAV-mDlx virus prompted us to
develop a novel AAV under the control of *TAC3+* interneuron striatal regulatory elements (**Fig. 6D-H**).
The novel reporter virus (AAV9-tandemE-TAC3-EGFP) that we developed to study the striatal *TAC3+*
type was driven by striatal *TAC3+* enhancer elements. When delivered systemically, the virus also labeled
TAC3+ neurons elsewhere in the brain, including in the hypothalamus, neocortex, substantia nigra. As
neocortical *TAC3+* cells bear little resemblance to *TAC3+* striatal interneurons in terms of their global
gene expression profiles (**Fig. 3H**), the widespread transduction of *TAC3+* neurons suggests that the
regulatory elements that we identified in the *TAC3+* striatal type (defined only relative to other striatal
cell types) may endogenously regulate the *TAC3* gene itself, or else a gene whose expression is highly
correlated to *TAC3*.

Our study reveals the complex landscape of transcriptionally-defined cell types in the marmoset brain.
Compared to mouse or other outgroup species, few primate lineage-gained cell types have emerged;
likely more common is the redirection, repurposing, or elaboration of conserved types (*2, 13, 33, 84, 85*).
The persistent fingerprint of developmental origin present in neuronal gene expression underscores
important roles for novel neurogenic niches, developmental patterning and altered mechanisms guiding
proliferation and cell migration in primate brain evolution that could be the subject of future study. For
example, *in vivo* lineage tracing in primates could reveal the developmental origin of the striatal TAC3+
type and its relationship to other TAC3+ populations in other brain structures. We report the largest
transcriptomic cell census of the marmoset brain and uncover the complex biodistribution of telencephalic
interneurons by quantitative smFISH mapping. We present a compendium of morphological
reconstructions of marmoset telencephalic interneurons, and describe new viral reagents for targeting and
visualizing a primate-specific striatal interneuron type. Together, these resources will enable further
comparative studies of the evolution and development of brain cell types.

METHODS

Animals used for study

Marmosets. Marmosets were pair-housed in spacious holding rooms with environmental control of
temperature (23–28°C), humidity (40–72%), and 12 hr light/dark cycle. Their cages were equipped with a
variety of perches and enrichment devices, and they received regular health checks and behavioral
assessment from MIT DCM veterinary staff and researchers. All animal procedures were conducted with
prior approval by the MIT Committee for Animal Care (CAC) and following veterinary guidelines.

Mice. Experimental mice were purchased from The Jackson Laboratory company and housed at the
Mclean hospital animal facility (3–5 mice per cage) on a 12:12 hr light-dark cycle in a temperature-
controlled colony room with unrestricted access to food and water. All procedures were conducted in
accordance with policy guidelines set by the National Institutes of Health and were approved by the
McLean Institutional Animal Care and Use Committee (IACUC).

Tissue processing for single nucleus sequencing and smFISH
Marmoset specimens for snRNA-seq. Marmoset experiments were approved by and in accordance with
Massachusetts Institute of Technology CAC protocol number 051705020. Adult marmosets (1.5–14.5
years old, 12 individuals; **Table S1**) were deeply sedated by intramuscular injection of ketamine (20–40
mg kg–1) or alfaxalone (5–10 mg kg–1), followed by intravenous injection of sodium pentobarbital (10–
30 mg kg–1). When the pedal with-drawal reflex was eliminated and/or the respiratory rate was
diminished, animals were trans-cardially perfused with ice-cold sucrose-HEPES buffer (*3*, *6*). Whole
brains were rapidly extracted into fresh buffer on ice. Sixteen 2-mm coronal blocking cuts were rapidly
made using a custom-designed marmoset brain matrix. Slabs were transferred to a dish with ice-cold
dissection buffer (*3*, *6*). All regions were dissected using a marmoset atlas as reference(*86*), and were
snap-frozen in liquid nitrogen or dry ice-cooled isopentane, and stored in individual microcentrifuge tubes
at –80 °C.
792
793 *Marmoset specimens for snATAC-seq.* Tissue from 1 marmoset (female, 1.5 y.o., **Table S1**) was used for
794 both snRNA-seq and snATAC-seq (**Table S1**). Fresh tissue was dissected from anterior striatum
795 (including caudate and putamen) and used immediately for snATAC-seq.
796
Mouse specimens for snRNA-seq. Three adult (P80-90) male wild-type mice were deeply anesthetized
with isoflurane and sacrificed by decapitation. Brains were quickly excised, washed in ice-cold sterile
0.1M phosphate buffer saline (PBS) and dissected onto an ice-cold glass surface. Amygdala nuclei were
identified and isolated using "The Allen mouse brain atlas" (https://mouse.brain-map.org/static/atlas,
Table S7) as a reference for anatomical landmarks. The basolateral amygdaloid nucleus was exposed by
performing two coronal cuts using the borders of *primary somatosensory cortex* and *primary visual*
cortex as landmarks. Dissected specimens were collected in 1.5ml micro-centrifuge tubes, snap-frozen on
dry ice, and stored at -80 °C until used.
805
Marmoset specimens for smFISH. Two marmosets were deeply sedated by intramuscular injection of
alfaxalone (5–10 mg kg–1) (**Table S1**), followed by intravenous overdose of sodium pentobarbital (10–
30 mg kg–1). When the pedal with-drawal reflex was eliminated and/or the respiratory rate was

diminished, animals were trans-cardially perfused with ice-cold saline. The brain was immediately
removed, embedded in Optimal Cutting Temperature (OCT) freezing medium, and flash frozen in an
isopropyl ethanol-dry ice bath. Samples were cut on a cryostat (Leica CM 1850) at a thickness of 16µm,
adhered to SuperFrost Plus microscope slides (VWR, 48311-703), and stored at -80 °C until use. Portions
of the brain that were not cut were recoated in OCT and stored again for future use. Samples were
immediately fixed in 4% paraformaldehyde and stained on the slide according to the Molecular
Instruments HCR generic sample in solution RNA-FISH protocol (Molecular Instruments,
https://files.molecularinstruments.com/MI-Protocol-RNAFISH-GenericSolution-Rev7.pdf) or the
Advanced Cell Diagnostics RNAscope Multiplex Fluorescent Reagent Kit v2 Assay (ACD, 323100,
https://acdbio.com/sites/default/files/USM-
323100%20Multiplex%20Fluorescent%20v2%20User%20Manual_10282019_0.pdf) protocol (**Table
S7**).
821
Single nucleus RNA-seq, library preparation, sequencing
10x RNA-seq. Unsorted single-nucleus suspensions from frozen marmoset and mouse samples were
generated as in (*87*). GEM generation and library preparation followed the manufacturer's protocol (10X
Chromium single-cell 3′ v.3, protocol version #CG000183_ ChromiumSingleCell3′_v3_UG_Rev-A).
Raw sequencing reads were aligned to the NCBI CJ1700 reference (marmoset) or GRCm38 (mouse).
Reads that mapped to exons or introns were assigned to annotated genes. Libraries were sequenced to a
median read depth of 8 reads per Unique Molecular Identifier (UMI, or transcript), obtaining a median
7,262 UMIs per cell.
830
RNA sequencing data processing, curation, clustering
Processing and alignment steps follow those outlined in: https://github.com/broadinstitute/Drop-seq/
(**Table S7)**. Raw BCL files were processed using IlluminaBasecallsToSam, and reads with a barcode
quality score below Q10 were discarded. Cell barcodes (CBCs) were filtered using the 10X CBC
whitelist, followed by TSO and polyA trimming. Reads were aligned using STAR, and tagged with their
gene mapping (exonic, intronic, or UTR) and function (strand). The reads were then processed through
GATK BQSR, and tabulated in a digital gene expression matrix (DGE) containing all CBCs with at least
20 transcripts aligning to coding, UTR, or intronic regions. Cell selection was performed based on
CellBender remove-background non-empties (*88*), % intronic (% of a CBC's reads that are intronic), and
number of UMIs for a CBC. A new filtered DGE containing only these selected CBCs was then
generated. Finally, a gene-metagene DGE was created by merging the selected CBCs DGE with a
metagene DGE (made by identifying reads from the selected CBCs that have a secondary alignment
mapping to a different gene than its primary alignment).
844
Cell type classification models were trained using our annotations and scPred R package version 1.9.2
(*21*). Detection of cell-cell doublets was performed using a two-step process based on the R package
DoubletFinder (*89*). DoubletFinder implements a nearest-neighbors approach to doublet detection. First,
artificial doublets are simulated from input single-cell data and are co-clustered with true libraries. True
doublet libraries are identified by their relative fraction of artificial doublet nearest-neighbors in gene
expression space. In our workflow this process is run twice, once with high stringency to identify and
remove clear doublets, and again with a lower threshold to identify remaining, subtler doublet libraries.
We used the following parameters for round 1: PN = 0.4, PK = 10 ^ seq(-4, -1, length.out=50),

NUM_PCS = 5. For the second round: PN = 0.45, NUM_PCS = 10, PK = 10 ^ seq(-4, -1, length.out=50).
Because doublet libraries are initially categorized as true libraries in the nearest-neighbor search, we find
this two-step process improves the sensitivity and accuracy of doublet detection.

Clustering was performed using independent component analysis (ICA; R package fastICA)
dimensionality reduction followed by Louvain. Cells assigned to one of the major glial types
(oligodendrocyte lineage, astrocytes, vascular/endothelia, microglia/macrophages) by scPred were
collected across all brain regions and clustered together. Neurons from most telencephalic structures
(neocortex, hippocampus, striatum, amygdala) confidently assigned to the categories "GABAergic" and
"glutamatergic" and so were clustered separately by neurotransmitter usage for each brain structure.
Striatal neuron categories were "SPN" and "GABAergic interneuron". Neurons in non-telencephalic brain
structures were clustered separately by brain structure. All clusterings were performed in two stages: first-
round clustering was based on the top 60 independent components (ICs) and a default resolution (res) of
0.1, nearest neighbors (nn) = 25. Following the process outlined in(6), each was manually reviewed for
skew and kurtosis of gene loadings on factors and cells to identify ICs that loaded on outliers, doublets, or
artifactual signals. These were discarded and reclustering was performed on the remaining ICs. Each
resulting cluster was then subjected to second-round clustering, during which ICs were again curated.
Second-round clustering explored a range of parameters: nn=10,20,30;
res=0.01,0.05,0.1,0.2,0.3,0.4,0.5,1.0. Final parameter values were chosen to optimize concordance, when
possible, between the final number of clusters and the number of included ICs, such that each cluster was
defined by one primary IC. Metacells for each cluster are generated by summing transcript counts for
each cell across all cells in the cluster, normalizing by total number of transcripts, and scaling to counts
per 100,000.

Identification of regionally differentially expressed genes (rDEGs)

We computed rDEGs for neocortical gluatmatergic neurons, GABAergic neurons, astrocytes, and
oligodendrocyte lineage types. We used individual cell level cluster assignments from the initial
neocortical clustering (which contained all regions together) to create per-region metacells for each
cluster. Normalized metacells were log10 transformed and pairwise differences across regions within the
same cluster (cell type) were examined. Genes with > 3 fold difference in the same cluster between two
regions were considered rDEGs. For this analysis we omitted from comparisons any region-cluster
metacell generated from fewer than 50 cells, but retain comparisons between regions that had > 50 cells
in that cluster. Genes that were consistently rDEGs in at least 3 individuals for each cluster-region pair are
reported in **Table S3**.

Ancestral State Reconstruction
Ancestral state reconstruction (ASR) is a method to infer hidden ancestral traits from extant observations.
For example, given a phylogenetic tree of species and genomic sequences thought to be homologous
across those species, the reconstruction takes into account branch lengths to reconstruct the most likely
ancestral sequence. We applied a maximum likelihood-based ASR approach (R package: fastAnc) to the
scaled, normalized metacells of cell types and the dendrogram of their similarity to produce estimates of
expression of each gene at the internal nodes of the tree. This enabled comparisons of leaf nodes to
internal nodes as well as internal nodes to each other. For example, compared to the parent node of

amygdala, basal forebrain, and hypothalamic SPN-like GABAergic projection neurons, the reconstructed
parent node of striatal SPNs had higher expression of known markers of striatal projection neurons such
as *DACH1*. We used these reconstructions of internal node gene expression to compare major clades of
the tree, and used a Chi-square test to determine whether transcription factors were overrepresented
among differentially expressed genes (threshold: 3-fold difference) between pairs of internal nodes.
901

902   **Spatial smFISH experiments**
903   All probes are listed in **Table S4**. All smFISH validation experiments were carried out on distinct
904   biological replicates from those used for snRNA-seq or single-cell ATAC-seq experiments.
905

smFISH tissue processing and quantification. Two marmosets (Cj 18-134 and 19-212) were euthanized
and perfused with PBS (**Table S1**). The brain was removed, embedded rapidly in OCT, and stored in the -
80C freezer. Tissue was then cut to 16µm on a cryostat and stored in the -80C until needed. *In situ*
hybridization was performed for genes of interest (see **Table S4**) with HCR or ACD antisense probes,
incubated with TrueBlack Lipofuscin Autofluorescence Quencher (Biotium, 23007) for 10 seconds at
room temperature to eliminate confounding lipofuscin autofluorescence present in the tissue. Samples
were then coverslipped with ProLong Diamond Antifade Mountant (Invitrogen, P36970). Z-stack serial
images were taken through the whole depth across striatum, hypothalamus and basal forebrain regions,
and several regions of neocortex, on the TissueGnostic TissueFAXS SL slide-scanning, spinning disk
confocal microscope (Hamamatsu Orca Flash 4.0 v3) using a 20×/0.8 NA air objective for ACD stains or
a 40×/1.2 NA water-immersion objective for HCR stains.
917

Series images were segmented using StrataQuest, a software package from TissueGnostics, which enables
the quantification of signals within segmented images (similar to CellProfiler). Nuclei objects were
generated using the DAPI channel, and artifacts were removed based on size and intensity. Exclusion
ROIs were manually drawn to avoid areas of white matter, large artifacts, and autofluorescence before
computing intensity and other statistical and morphological measurements (20 parameters) for each
channel. Specifically, 50 cells were hand-labeled as positive or negative for the markers of interest in
order to identify the appropriate threshold for feature selection using the parameters that best
discriminated this binary. Parameters included mean intensity, maximum intensity, standard deviation of
intensity, and range of intensity, and equivalent diameter. These filters were then applied to the unlabeled
data to identify positive cells.
928

Segmented cells were further analyzed using in-house code (https://github.com/klevando/BICCN-
StrataQuest-Script, **Table S7**). Spatial locations of the cells were visualized by plotting the x-y
coordinates associated with each nuclei. These were then binned into 2-D histograms across the x and y
axis (corresponding to the rostrocaudal plane and to the dorsoventral plane respectively). A size of 100
bins was chosen for the first (medial-most) slice of a series across the x axis and the calculated bin size
was then used across the y axis of the first slice and across other slices in the series. Positive events for a
gene in a given bin were either normalized to the number of detected nuclei in that bin (DAPI) and plotted
in 2-D as a relative heatmap, or simply plotted as a density heatmap without being normalized to DAPI.
Whole slice normalizations across the mediolateral axis (all positive events for a gene in a given slice
relative to DAPI) were plotted as bar graphs. DAPI and marker of interest counts are available in **Data**
S1-S2.

940
Neocortical Areal Proportions
In addition to the mediolateral subdivisions of marmoset neocortex, we further parcellated the neocortex
into subregions referencing the Brain/MINDS 3D Digital Marmoset Atlas
(https://doi.org/10.24475/bma.4520). Slice selection was based upon visual recognition of prominent
landmarks (white matter, striatal boundary as well as DAPI nuclei staining) within our tissue and matched
with the nearest atlas slice. ROIs for each parcellated region were created using the StrataQuest software
in the anterior-posterior and dorsal-ventral axes (**Fig. S8C**). The corresponding feature selection that was
previously set within the full neocortical sections was carried across gene and slice. The parameters used
were mean intensity, maximum intensity, standard deviation of intensity, and range of intensity, and
equivalent diameter. These filters were then applied to the unlabeled data to identify positive cells within
each individual ROI. The parcellated neocortical areas were then analyzed using in-house code
(https://github.com/klevando/BICCN-StrataQuest-Script, **Table S7**). As above, positive events for a gene
in a given ROI were normalized to DAPI and plotted in a (stacked) bar graph. Relative percentages
between genes were calculated and plotted stacked.

Morphology and smFISH experiments

AAV9-hDlx5/6-GFP-fGFP virus was generated as in (*19*). Virus was systemically IV injected (400ul-
700ul at 1.7- 2.4^{10} titer) in 5 marmosets. The virus was allowed to incubate for approximately 2 months.
After systemic IV injection with AAV9-hDlx5/6-GFP-fGFP, marmosets were euthanized and perfused
with saline followed by 4% paraformaldehyde (PFA). Brains were removed and 120μm sections were cut
on a vibratome into PBS-Azide and stored at 4C or moved into 70% ethanol for storage at -20C. 70%
ethanol storage prevents RNA degradation at this temperature without significant tissue shrinkage for
short storage times. Due to lab shutdowns during the pandemic, sections from two marmosets were stored
in 70% ethanol for approximately 4 months. These samples exhibited significant shrinkage, measured by
DAPI-stained nuclei diameters (**Table S1**), therefore we only compared morphology parameters within-
donor. Sections were taken as needed and *in situ* hybridization was performed with HCR antisense
probes, following the generic sample in solution HCR protocol with a 2-fold increase in concentration of
probe to hybridization buffer (Molecular Instruments, https://files.molecularinstruments.com/MI-
Protocol-RNAFISH-GenericSolution-Rev8.pdf), for markers of interest (**Table S4**) that corresponded
with RNA-seq defined clusters. The sections were then stained with anti-GFP antibody (**Table S4**) and a
secondary antibody (**Table S4**) to amplify the endogenous GFP signal
(https://www.protocols.io/view/marmoset-nhp-free-floating-anti-gfp-antibody-stain-3byl47nb2lo5/v1,
Table S4). Sections were incubated in TrueBlack (Biotium, 23007) for 3-5 minutes in order to mask
confounding lipofuscin autofluorescence throughout the section. Sections were then mounted onto a slide
and coverslipped with ProLong Diamond Antifade Mountant (Invitrogen, P36970) for imaging.

Imaging for morphology. Sections prepared for morphology were imaged on a Nikon Ti Eclipse inverted
microscope with an Andor CSU-W1 confocal spinning disc unit and Andor DU-888 EMCCD using a
40×/1.15 NA water-immersion objective, and later on a TissueGnostic TissueFAXS SL slide-scanning,
spinning disk confocal microscope (with Hamamatsu Orca Flash 4.0 v3) using a 40×/1.2 NA water-
immersion objective. With the TissueFAXS, overview images were taken in order to select GFP+ cells
for imaging at 40× and to highlight the exact location of the cell. GFP+ cells were imaged for stained
markers of interest. Selected sections were imaged on an upright confocal laser scanning microscope

(Olympus Fluoview FV3000) using a 40x/0.95 NA air objective or a 60x/1.50 NA oil-immersion
objective and cooled GaAsP PMTs.

Morphological reconstruction and feature quantification - Imaris. Without pre-processing the confocal
images, three-dimensional (3D) reconstruction and surface rendering of striatal and neocortical
interneurons were performed using Imarisx64 9.9 software (Oxford Instruments) based on GFP+ signal.
Surface-rendered images were used to determine the soma diameter, total volume, and total surface area
for each z-stack image. 3D-skeleton diagrams (**Figs. 5A-C** and **6H**), corresponding to each surface-
rendered image (data not shown), were generated using the Filament Tracing wizard in Imaris and then
pseudo-colored in Adobe Illustrator. The total number of primary dendritic branches, dendritic branch
points, area, volume, and length of the 3D-skeleton diagrams were calculated using the AutoPath (no
loops) algorithm in the filament tracing wizard in Imaris. The total number of primary dendritic branches
for each cell is defined by the number of dendrite branches in the filament trace 1 distance value away
from the soma. The distance value is calculated automatically by the AutoPath (no loops) algorithm based
on the diameter of the soma and the diameter of the thinnest cellular process. All data were exported to
CSV, and data collected from exemplar cells (**Fig. 5A-B** and **6G)** were reported in **Table S6**.

A separate dataset containing GFP+ interneurons from one animal (Cj 17-154) was first blinded using a
custom Python script, then reconstructed with Imaris (**Figs. 5C-D**). All results are presented as mean ±
SEM. Comparisons between soma diameter, surface area, surface volume, and the number of primary
branches were carried out using unpaired *t*-tests. Unpaired Mann-Whitney *U* tests were used to
statistically compare the total length, area, volume, and total number of dendritic branch points
measurements. Statistically significant analyses were denoted as follows: *$p < 0.05$; **$p < 0.01$.

1004
Morphological reconstruction and feature quantification - NeuTube. Automatic or semi-automatic tracing
algorithms are challenged by some data, perhaps due to the low SNR of a given image. To overcome this,
we manually reconstructed the sparse neurons via Neutube (*90*) tracing software. With the software, we
1) Create 3D volume rendering of the GFP-AAV marmoset neuron, 2) use the signal transfer function
(e.g., histogram equalization) for overall intensity and opacity values to optimize the signal-to-noise ratio
by manually examining the clearest visualization of the dendrites, 3) Build the neuron skeleton over the
3D volume by tracing the processes, 4) Scan through the 3D volume to make sure no parts of the neuron
are missed, 5) Double check the raw 2D images to see if any of the branches were not presented well in
3D due to volume rendering artifacts, 6) Label axon, dendrites and soma parts of the skeleton model.
Reconstructed neurons were saved as SWC format.
1015
Single nucleus ATAC-seq, library preparation, sequencing
Single nucleus ATAC-seq (snATAC-seq; 10x Genomics Single Cell ATAC v1) was conducted on fresh
marmoset tissue (1 female, 1.5 y.o.) dissected from anterior striatum. Nuclei suspensions were generated
following the 10x recommended protocol (CG000212 Rev B). Library preparation followed 10x
Genomics Single Cell ATAC v1 Guide (CG000168 Rev D). Library was sequenced on an Illumina
NovaSeq (RRID:SCR_016387) using 100 bp paired-end reads to a median per-cell fragment depth of
25,472. Alignment and fragment counting was conducted using Cell Ranger (RRID:SCR_01734), aligned
to marmoset genome cj1700 (https://www.ncbi.nlm.nih.gov/assembly/GCF_009663435.1/;
GCA_009663445.2). snATAC-seq data were integrated with marmoset striatal snRNA-seq data from two
independent animasl (bi005, bi006) using Signac (RRID:SCR_021158) with the following parameters:
integration method: cca; weight.reduction = lsi. Differentially accessible peaks for each cluster were

calculated with a 3-fold change cutoff and the minimum fraction of expressed cells (in target cluster) =
0.2. Enhancer locations were as follows (CJ1700 coordinates): Eh14 = chr1-122696345-122696560 (215
bp); Eh15 = chr17-32919427-32919564 (137 bp); Eh16 = chr20-38770509-38770619 (110 bp); Eh17 =
chr9-60258694-60258788 (94 bp).
1031
TAC3-AAV viral design and production.
AAVs were produced in accordance with the protocol (*91*). 5.7 µg/150 mm dish of construct DNA was
transfected with 22.8 µg/150 mm dish of pAAV9 capsid plasmid and 11.45 µg/150 mm dish of
pAdDeltaF6 helper plasmid using polyethylenimine 25K MW (Polysciences, 23966-1). Collection of
cells and media for AAV harvesting began 72 hours following transfection before iodixanol gradient
ultracentrifugation purification using a Type 70 Ti Fixed-Angle Titanium Rotor (Beckman-Coulter,
337922). Titer was calculated using digital droplet PCR (ddPCR) according to the protocol "ddPCR
Titration of AAV Vectors" (https://www.addgene.org/protocols/aav-ddpcr-titration/) using the QX200
AutoDG Droplet Digital PCR System (BioRad, 1864100). gBlock fragments containing the tandem *TAC3*
enhancers (E14E15E16E17) were synthesized by Integrated DNA Technologies (IDT). The enhancers
were cloned into AAV plasmid vector backbones containing a minimal CMV promoter, the reporter and a
barcode unique to each enhancer sequence using the NEBuilder® HiFi DNA Assembly Cloning Kit
(NEB-E5520S), following standard protocol. 2 uL of Gibson assembly product was used to transform 50
uL of home-made Stbl3 cells following a standard transformation protocol. Mini-preparation of plasmids
was carried out using the NucleoSpin mini kit (Macherey-Nagel). Positive clones were identified by
restriction enzyme digestion and sequencing. The positive clones were grown in 300 mL LB cultures and
the plasmids were extracted using the NucleoBond Xtra Midi EF kit (Macherey-Nagel). The plasmids
were sent out for sequencing again before being packaged into AAVs.

TAC3-AAV local injection procedure

Structural MRI Scanning. In preparation for MR imaging under anesthesia, an animal (2 marmosets, Cj
19-207 and 17-B111, **Table S1**) checked for robust health was fasted overnight with *ad libitum* access to
water. Prior to scanning, the animal was sedated with Alphaxalone (5-10 mg/kg) or a combination of
Alphaxalone (4-8 mg/kg) and Ketamine (5-10 mg/Kg) given intramuscularly. It was then transferred to a
custom designed MRI cradle equipped with a 3D printed nose cone for delivering anesthetic gases. The
animal's head was securely held to minimize motion artifacts and stereotaxically align the rostral-caudal
axis with the MR scanner bore using contrast filled MR compatible ear bars, an adjustable palette bar and
eye bars. During scanning, anesthesia was maintained at 1-2% isoflurane in a mixture of oxygen. A
temperature controlled water circulating heat pad was used for thermal support. Heart rate and blood
oxygen saturation levels were continuously monitored and recorded every 10 min with an MR compatible
monitoring system (Nonin, MN). MR scans consisted of high-resolution 3D T1 mapping using a
Magnetization-Prepared Rapid Gradient-Echo (MPRAGE) sequence (Liu et al., 2011) with repetition
time (TR) = 6000 ms, TE (echo time) = 4.6 ms; flip angle (FA) =12 degrees; FOV = 80, matrix size =
256x256. A high-resolution T2-weighted anatomical scan was also obtained using a RARE pulse
sequence with effective TE = 35.5 ms, TR = 2500 ms, RARE factor = 8, spatial resolution = 156 µm ×
156 µm × 0.5 mm, and matrix size = 256 × 256, RARE factor = 4, 80 mm FOV.
1067
Virus injection surgery. Following sedation with Alphaxalone (5-10 mg/kg), the animal was intubated and
maintained at 1.5-2% isoflurane in a mixture of oxygen throughout the rest of the procedure. An

intravenous (IV) catheter was placed in the saphenous vein for infusion of fluids and drugs during the
surgery and the recovery period. Heart rate, SpO2, ECG, end-tidal CO_2 and rectal temperature were
continuously monitored and logged every 5 minutes throughout. Once the animal acquired a stable plane
of anesthesia, it was placed in a stereotaxic apparatus (Narishige, Japan). A thin layer of sterile eye
lubricant was applied to protect against corneal drying. Animal was also provided with a single bolus
dose of intravenous dexamethasone (0.4 mg/kg) to guard against brain swelling. The scalp and fascia
were removed in layers via blunt dissection to expose the injection sites. Using MRI atlas guidance,
bilateral craniotomies were performed to expose the medial anterior caudate and ventral posterior
putamen region. The tandem enhancer virus AAV9-tandemE-TAC3-EGFP was injected in left Caudate
(high titer, 10^13 low volume, 0.5 ul) and left Putamen (low titer, 10^11, high volume 1.5 ul). Each
injection was delivered with a controlled syringe pump (100 nL/minute). After each injection the needle
was kept in place for 10 minutes then slowly retracted over 1-2 minutes to minimize injection backflow.
Following withdrawal, the cranial openings were covered with bonewax, fascia and scalp were sutured in
layer wise manner with 3-0 vicryl sutures. The suture site was cleaned with warm sterile saline and
covered with hypafix. For recovery the animal was transferred in an incubator and monitored closely till it
was able to move effortlessly and accepted treats before returning to its home cage. Post-op medications
were provided under veterinary advice for pain and inflammation control until full recovery. The animals
were euthanized 6-10 weeks after viral injection and were perfused transcardially with 4% PFA. The
brain was extracted for histology, smFISH, and confocal microscopy.
1089

TAC3-AAV-BI103 systemic injection procedure

One 2-year old female marmoset (**Table S1**) was sedated with Alphaxalone (5-10 mg/kg) given
intramuscularly. A catheter was placed into the tail vein and 200 μl of purified viral particles were
injected at a dose of 7.85×10^{13} vg/kg, followed by injection of 1.5 ml saline to flush the line. The animal
was euthanized 4 weeks after viral injection and was perfused transcardially with 4% PFA. The brain was
extracted for histology, smFISH, and confocal microscopy.

TAC3-AAV histology

For AAV9-tandemE-TAC3-EGFP experiments, coronal sections around the injection sites were cut at
100 μm on a vibratome (Leica, VT1000S) and then stored in PBS-NaN3 at 4 °C until use. For AAV-
BI103-tandemE-TAC3-EGFP experiments, coronal and sagittal sections were processed. Sections were
stained for *TAC3* (Molecular Instruments, PRC843, **Table S4**) using the HCR v.3.0 protocol (Molecular
Instruments) described above and a rabbit anti-GFP antibody (Invitrogen, A11122, **Table S4**) with
subsequent secondary goat anti-rabbit conjugated with AF488 (ThermoFIsher, A-11008). Stained sections
were then incubated 1-3 minutes in TrueBlack to eliminate confounding lipofuscin autofluorescence. The
striatum was imaged for each section using the TissueFAXS SL with a 40×/1.2 NA water-immersion lens
through the whole thickness of the tissue.
1106

References

1.   S. Ma, M. Skarica, Q. Li, C. Xu, R. D. Risgaard, A. T. N. Tebbenkamp, X. Mato-Blanco, R. Kovner,

Ž. Krsnik, X. de Martin, V. Luria, X. Martí-Pérez, D. Liang, A. Karger, D. K. Schmidt, Z. Gomez-
Sanchez, C. Qi, K. T. Gobeske, S. Pochareddy, A. Debnath, C. J. Hottman, J. Spurrier, L. Teo, A. G.
Boghdadi, J. Homman-Ludiye, J. J. Ely, E. W. Daadi, D. Mi, M. Daadi, O. Marín, P. R. Hof, M.-R.
Rasin, J. Bourne, C. C. Sherwood, G. Santpere, M. J. Girgenti, S. M. Strittmatter, A. M. M. Sousa,
N. Sestan, Molecular and cellular evolution of the primate dorsolateral prefrontal cortex. *Science*,
eabo7257 (2022).

2. T. E. Bakken, N. L. Jorstad, Q. Hu, B. B. Lake, W. Tian, B. E. Kalmbach, M. Crow, R. D. Hodge, F.
M. Krienen, S. A. Sorensen, J. Eggermont, Z. Yao, B. D. Aevermann, A. I. Aldridge, A. Bartlett, D.
Bertagnolli, T. Casper, R. G. Castanon, K. Crichton, T. L. Daigle, R. Dalley, N. Dee, N. Dembrow,
D. Diep, S.-L. Ding, W. Dong, R. Fang, S. Fischer, M. Goldman, J. Goldy, L. T. Graybuck, B. R.
Herb, X. Hou, J. Kancherla, M. Kroll, K. Lathia, B. van Lew, Y. E. Li, C. S. Liu, H. Liu, J. D.
Lucero, A. Mahurkar, D. McMillen, J. A. Miller, M. Moussa, J. R. Nery, P. R. Nicovich, S.-Y. Niu,
J. Orvis, J. K. Osteen, S. Owen, C. R. Palmer, T. Pham, N. Plongthongkum, O. Poirion, N. M. Reed,
C. Rimorin, A. Rivkin, W. J. Romanow, A. E. Sedeño-Cortés, K. Siletti, S. Somasundaram, J. Sulc,
M. Tieu, A. Torkelson, H. Tung, X. Wang, F. Xie, A. M. Yanny, R. Zhang, S. A. Ament, M. M.
Behrens, H. C. Bravo, J. Chun, A. Dobin, J. Gillis, R. Hertzano, P. R. Hof, T. Höllt, G. D. Horwitz,
C. D. Keene, P. V. Kharchenko, A. L. Ko, B. P. Lelieveldt, C. Luo, E. A. Mukamel, A. Pinto-Duarte,
S. Preissl, A. Regev, B. Ren, R. H. Scheuermann, K. Smith, W. J. Spain, O. R. White, C. Koch, M.
Hawrylycz, B. Tasic, E. Z. Macosko, S. A. McCarroll, J. T. Ting, H. Zeng, K. Zhang, G. Feng, J. R.
Ecker, S. Linnarsson, E. S. Lein, Comparative cellular analysis of motor cortex in human, marmoset
and mouse. *Nature*. **598**, 111–119 (2021).

3. F. M. Krienen, M. Goldman, Q. Zhang, R. C. H. del Rosario, M. Florio, R. Machold, A. Saunders,
K. Levandowski, H. Zaniewski, B. Schuman, C. Wu, A. Lutservitz, C. D. Mullally, N. Reed, E.
Bien, L. Bortolin, M. Fernandez-Otero, J. D. Lin, A. Wysoker, J. Nemesh, D. Kulp, M. Burns, V.
Tkachev, R. Smith, C. A. Walsh, J. Dimidschstein, B. Rudy, L. S. Kean, S. Berretta, G. Fishell, G.
Feng, S. A. McCarroll, Innovations present in the primate interneuron repertoire. *Nature*. **586**, 262–
269 (2020).

4. J.-P. Lin, H. M. Kelly, Y. Song, R. Kawaguchi, D. H. Geschwind, S. Jacobson, D. S. Reich,
Microenvironment Impacts the Molecular Architecture and Interactivity of Resident Cells in
Marmoset Brain. *bioRxiv* (2021), p. 2021.01.25.426385.

5. R. C. Bandler, I. Vitali, R. N. Delgado, M. C. Ho, E. Dvoretskova, J. S. Ibarra Molinas, P. W. Frazel,
M. Mohammadkhani, R. Machold, S. Maedler, S. A. Liddelow, T. J. Nowakowski, G. Fishell, C.
Mayer, Single-cell delineation of lineage and genetic identity in the mouse brain. *Nature*. **601**, 404–
409 (2022).

6. A. Saunders, E. Z. Macosko, A. Wysoker, M. Goldman, F. M. Krienen, H. de Rivera, E. Bien, M.
Baum, L. Bortolin, S. Wang, A. Goeva, J. Nemesh, N. Kamitaki, S. Brumbaugh, D. Kulp, S. A.
McCarroll, Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. *Cell*.
174, 1015–1030.e16 (2018).

7. A. Zeisel, H. Hochgerner, P. Lönnerberg, A. Johnsson, F. Memic, J. van der Zwan, M. Häring, E.
Braun, L. E. Borm, G. La Manno, S. Codeluppi, A. Furlan, K. Lee, N. Skene, K. D. Harris, J.
Hjerling-Leffler, E. Arenas, P. Ernfors, U. Marklund, S. Linnarsson, Molecular Architecture of the
Mouse Nervous System. *Cell*. **174**, 999–1014.e22 (2018).

8. Z. Yao, C. T. J. van Velthoven, M. Kunst, M. Zhang, D. McMillen, C. Lee, W. Jung, J. Goldy, A.
Abdelhak, P. Baker, E. Barkan, D. Bertagnolli, J. Campos, D. Carey, T. Casper, A. B. Chakka, R.
Chakrabarty, S. Chavan, M. Chen, M. Clark, J. Close, K. Crichton, S. Daniel, T. Dolbeare, L.

Ellingwood, J. Gee, A. Glandon, J. Gloe, J. Gould, J. Gray, N. Guilford, J. Guzman, D. Hirschstein, W. Ho, K. Jin, M. Kroll, K. Lathia, A. Leon, B. Long, Z. Maltzer, N. Martin, R. McCue, E. Meyerdierks, T. N. Nguyen, T. Pham, C. Rimorin, A. Ruiz, N. Shapovalova, C. Slaughterbeck, J. Sulc, M. Tieu, A. Torkelson, H. Tung, N. V. Cuevas, K. Wadhwani, K. Ward, B. Levi, C. Farrell, C. L. Thompson, S. Mufti, C. M. Pagan, L. Kruse, N. Dee, S. M. Sunkin, L. Esposito, M. J. Hawrylycz, J. Waters, L. Ng, K. A. Smith, B. Tasic, X. Zhuang, H. Zeng, A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. *bioRxiv* (2023), p. 2023.03.06.531121.

9. J. Langlieb, N. Sachdev, K. Balderrama, N. Nadaf, M. Raj, E. Murray, J. Webber, C. Vanderburg, V. Gazestani, D. Tward, C. Mezias, X. Li, D. Cable, T. Norton, P. P. Mitra, F. Chen, E. Macosko, The cell type composition of the adult mouse brain revealed by single cell and spatial genomics. *bioRxiv* (2023), doi:10.1101/2023.03.06.531307.

10. K. L. Chiou, X. Huang, M. O. Bohlen, S. Tremblay, D. R. O'Day, C. H. Spurrell, A. A. Gogate, T. M. Zintel, Cayo Biobank Research Unit, M. G. Andrews, M. I. Martinez, L. M. Starita, M. J. Montague, M. L. Platt, J. Shendure, N. Snyder-Mackler, A single-cell multi-omic atlas spanning the adult rhesus macaque brain. *bioRxiv* (2022), p. 2022.09.30.510346.

11. K. Siletti, R. Hodge, A. M. Albiach, L. Hu, K. W. Lee, P. Lönnerberg, T. Bakken, S.-L. Ding, M. Clark, T. Casper, N. Dee, J. Gloe, C. Dirk Keene, J. Nyhus, H. Tung, A. M. Yanny, E. Arenas, E. S. Lein, S. Linnarsson, Transcriptomic diversity of cell types across the adult human brain. *bioRxiv* (2022), p. 2022.10.12.511898.

12. J.-P. Lin, H. M. Kelly, Y. Song, R. Kawaguchi, D. H. Geschwind, S. Jacobson, D. S. Reich, Transcriptomic architecture of nuclei in the marmoset CNS. *Nat. Commun.* **13**, 5531 (2022).

13. R. D. Hodge, T. E. Bakken, J. A. Miller, K. A. Smith, E. R. Barkan, L. T. Graybuck, J. L. Close, B. Long, N. Johansen, O. Penn, Z. Yao, J. Eggermont, T. Hollt, B. P. Levi, S. I. Shehata, B. Aevermann, A. Beller, D. Bertagnolli, K. Brouner, T. Casper, C. Cobbs, R. Dalley, N. Dee, S.-L. Ding, R. G. Ellenbogen, O. Fong, E. Garren, J. Goldy, R. P. Gwinn, D. Hirschstein, C. D. Keene, M. Keshk, A. L. Ko, K. Lathia, A. Mahfouz, Z. Maltzer, M. McGraw, T. N. Nguyen, J. Nyhus, J. G. Ojemann, A. Oldre, S. Parry, S. Reynolds, C. Rimorin, N. V. Shapovalova, S. Somasundaram, A. Szafer, E. R. Thomsen, M. Tieu, G. Quon, R. H. Scheuermann, R. Yuste, S. M. Sunkin, B. Lelieveldt, D. Feng, L. Ng, A. Bernard, M. Hawrylycz, J. W. Phillips, B. Tasic, H. Zeng, A. R. Jones, C. Koch, E. S. Lein, Conserved cell types with divergent features in human versus mouse cortex. *Nature*. **573**, 1–38 (2019).

14. S. Sahara, Y. Yanagawa, D. D. M. O'Leary, C. F. Stevens, The Fraction of Cortical GABAergic Neurons Is Constant from Near the Start of Cortical Neurogenesis to Adulthood. *Journal of Neuroscience*. **32**, 4755–4761 (2012).

15. Y. Kim, G. R. Yang, K. Pradhan, K. U. Venkataraju, M. Bota, L. C. G. del Molino, G. Fitzgerald, K. Ram, M. He, J. M. Levine, P. Mitra, Z. J. Huang, X.-J. Wang, P. Osten, Brain-wide Maps Reveal Stereotyped Cell-Type- Based Cortical Architecture and Subcortical Sexual Dimorphism. *Cell*. **171**, 456–460.e22 (2017).

16. J. Hill, T. Inder, J. Neil, D. Dierker, J. Harwell, D. Van Essen, Similar patterns of cortical expansion during human development and evolution. *Proceedings of the National Academy of Sciences*. **107**, 13135–13140 (2010).

17. D. Goertsen, N. C. Flytzanis, N. Goeden, M. R. Chuapoco, A. Cummins, Y. Chen, Y. Fan, Q. Zhang, J. Sharma, Y. Duan, L. Wang, G. Feng, Y. Chen, N. Y. Ip, J. Pickel, V. Gradinaru, AAV capsid

variants with brain-wide transgene expression and decreased liver targeting after intravenous
delivery in mouse and marmoset. *Nat. Neurosci.* **25**, 106–115 (2022).

18. J. K. Mich, L. T. Graybuck, E. E. Hess, J. T. Mahoney, Y. Kojima, Y. Ding, S. Somasundaram, J. A.
Miller, B. E. Kalmbach, C. Radaelli, B. B. Gore, N. Weed, V. Omstead, Y. Bishaw, N. V.
Shapovalova, R. A. Martinez, O. Fong, S. Yao, M. Mortrud, P. Chong, L. Loftus, D. Bertagnolli, J.
Goldy, T. Casper, N. Dee, X. Opitz-Araya, A. Cetin, K. A. Smith, R. P. Gwinn, C. Cobbs, A. L. Ko,
J. G. Ojemann, C. D. Keene, D. L. Silbergeld, S. M. Sunkin, V. Gradinaru, G. D. Horwitz, H. Zeng,
B. Tasic, E. S. Lein, J. T. Ting, B. P. Levi, Functional enhancer elements drive subclass- selective
expression from mouse to primate neocortex. *Cell Rep.* **34**, 108754 (2021).

19. J. Dimidschstein, Q. Chen, R. Tremblay, S. L. Rogers, G.-A. Saldi, L. Guo, Q. Xu, R. Liu, C. Lu, J.
Chu, J. S. Grimley, A.-R. Krostag, A. Kaykas, M. C. Avery, M. S. Rashid, M. Baek, A. L. Jacob, G.
B. Smith, D. E. Wilson, G. Kosche, I. Kruglikov, T. Rusielewicz, V. C. Kotak, T. M. Mowery, S. A.
Anderson, E. M. Callaway, J. S. Dasen, D. Fitzpatrick, V. Fossati, M. A. Long, S. Noggle, J. H.
Reynolds, D. H. Sanes, B. Rudy, G. Feng, G. Fishell, A viral strategy for targeting and manipulating
interneurons across vertebrate species. *Nat. Neurosci.* **19**, 1743–1749 (2016).

20. D. Vormstein-Schneider, J. D. Lin, K. A. Pelkey, R. Chittajallu, B. Guo, M. A. Arias-Garcia, K.
Allaway, S. Sakopoulos, G. Schneider, O. Stevenson, J. Vergara, J. Sharma, Q. Zhang, T. P.
Franken, J. Smith, L. A. Ibrahim, K. J. M. Astro, E. Sabri, S. Huang, E. Favuzzi, T. Burbridge, Q.
Xu, L. Guo, I. Vogel, V. Sanchez, G. A. Saldi, B. L. Gorissen, X. Yuan, K. A. Zaghloul, O.
Devinsky, B. L. Sabatini, R. Batista-Brito, J. Reynolds, G. Feng, Z. Fu, C. J. McBain, G. Fishell, J.
Dimidschstein, Viral manipulation of functionally distinct interneurons in mice, non-human primates
and humans. *Nat. Neurosci.* **23**, 1–21 (2020).

21. J. Alquicira-Hernandez, A. Sathe, H. P. Ji, Q. Nguyen, J. E. Powell, scPred: accurate supervised
method for cell-type classification from single-cell RNA-seq data. *Genome Biol.* **20**, 264 (2019).

22. H. Zeng, J. R. Sanes, Neuronal cell-type classification: challenges, opportunities and the path
forward. *Nat. Rev. Neurosci.* **18**, 530–546 (2017).

23. D. Arendt, J. M. Musser, C. V. H. Baker, A. Bergman, C. Cepko, D. H. Erwin, M. Pavlicev, G.
Schlosser, S. Widder, M. D. Laubichler, G. P. Wagner, The origin and evolution of cell types. *Nat.
Rev. Genet.* **17**, 744–757 (2016).

24. H. Zeng, What is a cell type and how to define it? *Cell* (2022) (available at
https://www.sciencedirect.com/science/article/pii/S0092867422007838).

25. L. J. Revell, phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution. **3**, 217–223 (2012).

26. J. Woych, A. Ortega Gurrola, A. Deryckere, E. C. B. Jaeger, E. Gumnit, G. Merello, J. Gu, A. Joven
Araus, N. D. Leigh, M. Yun, A. Simon, M. A. Tosches, Cell-type profiling in salamanders identifies
innovations in vertebrate forebrain evolution. *Science*. **377**, eabp9186 (2022).

27. G. Meyer, Building a human cortex: the evolutionary differentiation of Cajal-Retzius cells and the
cortical hem. *J. Anat.* **217**, 334–343 (2010).

28. P. Jager, G. Moore, P. Calpin, X. Durmishi, I. Salgarella, L. Menage, Y. Kita, Y. Wang, D. W. Kim,
S. Blackshaw, S. R. Schultz, S. Brickley, T. Shimogori, A. Delogu, Dual midbrain and forebrain
origins of thalamic inhibitory interneurons. *Elife*. **10** (2021), doi:10.7554/eLife.59272.

29. K. Letinic, P. Rakic, Telencephalic origin of human thalamic GABAergic neurons. *Nat. Neurosci.* **4**,
931–936 (2001).

30. L. W. Swanson, G. D. Petrovich, What is the amygdala? *Trends Neurosci.* **21**, 323–331 (1998).

31. T. Hirata, P. Li, G. M. Lanuza, L. A. Cocas, M. M. Huntsman, J. G. Corbin, Identification of distinct
telencephalic progenitor pools for neuronal diversity in the amygdala. *Nat. Neurosci.* **12**, 141–149
(2009).

32. T. Kaoru, F. C. Liu, M. Ishida, T. Oishi, M. Hayashi, M. Kitagawa, K. Shimoda, H. Takahashi,
Molecular characterization of the intercalated cell masses of the amygdala: implications for the
relationship with the striatum. *NSC.* **166**, 220–230 (2010).

33. M. T. Schmitz, K. Sandoval, C. P. Chen, M. A. Mostajo-Radji, W. W. Seeley, T. J. Nowakowski, C.
J. Ye, M. F. Paredes, A. A. Pollen, The development and evolution of inhibitory neurons in primate
cerebrum. *Nature.* **603**, 871–877 (2022).

34. D. L. Kaufman, C. R. Houser, A. J. Tobin, Two forms of the gamma-aminobutyric acid synthetic
enzyme glutamate decarboxylase have distinct intraneuronal distributions and cofactor interactions.
J. Neurochem. **56**, 720–723 (1991).

35. J. R. Moffitt, D. Bambah-Mukku, S. W. Eichhorn, E. Vaughn, K. Shekhar, J. D. Perez, N. D.
Rubinstein, J. Hao, A. Regev, C. Dulac, X. Zhuang, Molecular, spatial, and functional single-cell
profiling of the hypothalamic preoptic region. *Science.* **362**, eaau5324–80 (2018).

36. B. Tasic, Z. Yao, L. T. Graybuck, K. A. Smith, T. N. Nguyen, D. Bertagnolli, J. Goldy, E. Garren,
M. N. Economo, S. Viswanathan, O. Penn, T. Bakken, V. Menon, J. Miller, O. Fong, K. E.
Hirokawa, K. Lathia, C. Rimorin, M. Tieu, R. Larsen, T. Casper, E. Barkan, M. Kroll, S. Parry, N.
V. Shapovalova, D. Hirschstein, J. Pendergraft, H. A. Sullivan, T. K. Kim, A. Szafer, N. Dee, P.
Groblewski, I. Wickersham, A. Cetin, J. A. Harris, B. P. Levi, S. M. Sunkin, L. Madisen, T. L.
Daigle, L. Looger, A. Bernard, J. Phillips, E. Lein, M. Hawrylycz, K. Svoboda, A. R. Jones, C.
Koch, H. Zeng, Shared and distinct transcriptomic cell types across neocortical areas. *Nature.* **563**,
1–41 (2018).

37. P. Balaram, J. H. Kaas, Towards a unified scheme of cortical lamination for primary visual cortex
across primates: insights from NeuN and VGLUT2 immunoreactivity. *Front. Neuroanat.* **8**, 81
(2014).

38. O. A. Bayraktar, L. C. Fuentealba, A. Alvarez-Buylla, D. H. Rowitch, Astrocyte development and
heterogeneity. *Cold Spring Harb. Perspect. Biol.* **7**, a020362 (2014).

39. O. A. Bayraktar, T. Bartels, S. Holmqvist, V. Kleshchevnikov, A. Martirosyan, D. Polioudakis, L.
Ben Haim, A. M. H. Young, M. Y. Batiuk, K. Prakash, A. Brown, K. Roberts, M. F. Paredes, R.
Kawaguchi, J. H. Stockley, K. Sabeur, S. M. Chang, E. Huang, P. Hutchinson, E. M. Ullian, M.
Hemberg, G. Coppola, M. G. Holt, D. H. Geschwind, D. H. Rowitch, Astrocyte layers in the
mammalian cerebral cortex revealed by a single-cell in situ transcriptomic map. *Nat. Neurosci.* **23**,
500–509 (2020).

40. M. Y. Batiuk, A. Martirosyan, J. Wahis, F. de Vin, C. Marneffe, C. Kusserow, J. Koeppen, J. F.
Viana, J. F. Oliveira, T. Voet, C. P. Ponting, T. G. Belgard, M. G. Holt, Identification of region-
specific astrocyte subtypes at single cell resolution. *Nat. Commun.* **11**, 1220 (2020).

41. A. Kriegstein, A. Alvarez-Buylla, The glial nature of embryonic and adult neural stem cells. *Annu.*
Rev. Neurosci. **32**, 149–184 (2009).

42. C. C. Harwell, L. C. Fuentealba, A. Gonzalez-Cerrillo, P. R. L. Parker, C. C. Gertz, E. Mazzola, M.
T. Garcia, A. Alvarez-Buylla, C. L. Cepko, A. R. Kriegstein, Wide Dispersion and Diversity of
Clonally Related Inhibitory Interneurons. *Neuron*. **87**, 999–1007 (2015).

43. C. Mayer, C. Hafemeister, R. C. Bandler, R. Machold, R. B. Brito, X. Jaglin, K. Allaway, A. Butler,
G. Fishell, R. Satija, Developmental diversification of cortical inhibitory interneurons. *Nature*. **555**,
457–462 (2018).

44. C. Neyt, M. Welch, A. Langston, J. Kohtz, G. Fishell, A short-range signal restricts cell movement
between telencephalic proliferative zones. *J. Neurosci.* **17**, 9194–9203 (1997).

45. F. García-Moreno, M. Pedraza, L. G. Di Giovannantonio, M. Di Salvio, L. López-Mascaraque, A.
Simeone, J. A. De Carlos, A neuronal migratory pathway crossing from diencephalon to
telencephalon populates amygdala nuclei. *Nat. Neurosci.* **13**, 680–689 (2010).

46. A. J. Granger, M. L. Wallace, B. L. Sabatini, ScienceDirect Multi-transmitter neurons in the
mammalian central nervous system. *Curr. Opin. Neurobiol.* **45**, 85–91 (2017).

47. N. X. Tritsch, A. J. Granger, B. L. Sabatini, Mechanisms and functions of GABA co-release. *Nat.*
Genet. **17**, 139–145 (2016).

48. M. Bupesh, I. Legaz, A. Abellán, L. Medina, Multiple telencephalic and extratelencephalic
embryonic domains contribute neurons to the medial extended amygdala. *J. Comp. Neurol.* **519**,
1505–1525 (2011).

49. L. Zhang, V. S. Hernandez, C. R. Gerfen, S. Z. Jiang, L. Zavala, R. A. Barrio, L. E. Eiden,
Behavioral role of PACAP signaling reflects its selective distribution in glutamatergic and
GABAergic neuronal subpopulations. *Elife*. **10** (2021), doi:10.7554/eLife.61718.

50. N. Bozadjieva-Kramer, R. A. Ross, D. Q. Johnson, H. Fenselau, D. L. Haggerty, B. Atwood, B.
Lowell, J. N. Flak, The Role of Mediobasal Hypothalamic PACAP in the Control of Body Weight
and Metabolism. *Endocrinology*. **162** (2021), doi:10.1210/endocr/bqab012.

51. J.-H. Cho, K. Zushida, G. P. Shumyatsky, W. A. Carlezon Jr, E. G. Meloni, V. Y. Bolshakov,
Pituitary adenylate cyclase-activating polypeptide induces postsynaptically expressed potentiation in
the intra-amygdala circuit. *J. Neurosci.* **32**, 14165–14177 (2012).

52. E. Y. Lee, L. C. Chan, H. Wang, J. Lieng, M. Hung, Y. Srinivasan, J. Wang, J. A. Waschek, A. L.
Ferguson, K.-F. Lee, N. Y. Yount, M. R. Yeaman, G. C. L. Wong, PACAP is a pathogen-inducible
resident antimicrobial neuropeptide affording rapid and contextual molecular host defense of the
brain. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021), doi:10.1073/pnas.1917623117.

53. Y. Kita, H. Nishibe, Y. Wang, T. Hashikawa, S. S. Kikuchi, M. U, A. C. Yoshida, C. Yoshida, T.
Kawase, S. Ishii, H. Skibbe, T. Shimogori, Cellular-resolution gene expression profiling in the
neonatal marmoset brain reveals dynamic species- and region-specific differences. *Proc. Natl. Acad.*
Sci. U. S. A. **118** (2021), doi:10.1073/pnas.2020125118.

54. T. Shimogori, A. Abe, Y. Go, T. Hashikawa, N. Kishi, S. S. Kikuchi, Y. Kita, K. Niimi, H. Nishibe,
M. Okuno, K. Saga, M. Sakurai, M. Sato, T. Serizawa, S. Suzuki, E. Takahashi, M. Tanaka, S.

Tatsumoto, M. Toki, M. U, Y. Wang, K. J. Windak, H. Yamagishi, K. Yamashita, T. Yoda, A. C. Yoshida, C. Yoshida, T. Yoshimoto, H. Okano, Digital gene atlas of neonate common marmoset brain. *Neurosci. Res.* **128**, 1–13 (2018).

55. D. P. Orquera, M. B. Tavella, F. S. J. de Souza, S. Nasif, M. J. Low, M. Rubinstein, The Homeodomain Transcription Factor NKX2.1 Is Essential for the Early Specification of Melanocortin Neuron Identity and Activates Pomc Expression in the Developing Hypothalamus. *J. Neurosci.* **39**, 4023–4035 (2019).

56. R. Murcia-Ramón, V. Company, I. Juárez-Leal, A. Andreu-Cervera, F. Almagro-García, S. Martínez, D. Echevarría, E. Puelles, Neuronal tangential migration from Nkx2.1-positive hypothalamus. *Brain Struct. Funct.* **225**, 2857–2869 (2020).

57. C. E. Collins, D. C. Airey, N. A. Young, D. B. Leitch, J. H. Kaas, Neuron densities vary across and within cortical areas in primates. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 15927–15932 (2010).

58. B. Schuman, R. P. Machold, Y. Hashikawa, J. Fuzik, G. J. Fishell, B. Rudy, Four Unique Interneuron Populations Reside in Neocortical Layer 1. *J. Neurosci.* **39**, 125–139 (2019).

59. M. Valero, T. J. Viney, R. Machold, S. Mederos, I. Zutshi, B. Schuman, Y. Senzai, B. Rudy, G. Buzsáki, Sleep down state-active ID2/Nkx2.1 interneurons in the neocortex. *Nat. Neurosci.* **24**, 401–411 (2021).

60. B. Zingg, H. Hintiryan, L. Gou, M. Y. Song, M. Bay, M. S. Bienkowski, N. N. Foster, S. Yamashita, I. Bowman, A. W. Toga, H.-W. Dong, Neural networks of the mouse neocortex. *Cell*. **156**, 1096–1111 (2014).

61. H. B. M. Uylings, H. J. Groenewegen, B. Kolb, Do rats have a prefrontal cortex? *Behav. Brain Res.* **146**, 3–17 (2003).

62. R. L. Buckner, D. S. Margulies, Macroscale cortical organization and a default-like apex transmodal network in the marmoset monkey. *Nat. Commun.* **10**, 1976 (2019).

63. M. K. Lin, Y. S. Takahashi, B.-X. Huo, M. Hanada, J. Nagashima, J. Hata, A. S. Tolpygo, K. Ram, B. C. Lee, M. I. Miller, M. G. Rosa, E. Sasaki, A. Iriki, H. Okano, P. Mitra, A high-throughput neurohistological pipeline for brain-wide mesoscale connectivity mapping of the common marmoset. *Elife*. **8** (2019), doi:10.7554/eLife.40042.

64. P. Majka, S. Bai, S. Bakola, S. Bednarek, J. M. Chan, N. Jermakow, L. Passarelli, D. H. Reser, P. Theodoni, K. H. Worthy, X.-J. Wang, D. K. Wójcik, P. P. Mitra, M. G. P. Rosa, Open access resource for cellular-resolution analyses of corticocortical connectivity in the marmoset monkey. *Nat. Commun.* **11**, 1133 (2020).

65. A. Hladnik, D. Džaja, S. Darmopil, N. Jovanov-Milošević, Z. Petanjek, Spatio-temporal extension in site of origin for cortical calretinin neurons in primates. *Front. Neuroanat.* **8**, 50 (2014).

66. E. Y. Choi, B. T. T. Yeo, R. L. Buckner, The organization of the human striatum estimated by intrinsic functional connectivity. *J. Neurophysiol.* **108**, 2242–2263 (2012).

67. K. M. Anderson, F. M. Krienen, E. Y. Choi, J. M. Reinen, B. T. T. Yeo, A. J. Holmes, Gene expression links functional networks across cortex and striatum. *Nat. Commun.* **9**, 1428 (2018).

68. M. Matamales, J. Götz, J. Bertran-Gonzalez, Quantitative Imaging of Cholinergic Interneurons

Reveals a Distinctive Spatial Organization and a Functional Gradient across the Mouse Striatum.
PLoS One. **11**, e0157682 (2016).

69. E. Fino, M. Vandecasteele, S. Perez, F. Saudou, L. Venance, Region-specific and state-dependent
action of striatal GABAergic interneurons. *Nat. Commun.* **9**, 3339 (2018).

70. A. B. M. Manchado, C. B. Gonzales, A. Zeisel, H. Munguba, B. Bekkouche, N. G. Skene, P.
Lonnerberg, J. Ryge, K. D. Harris, S. Linnarsson, J. Hjerling-Leffler, Diversity of Interneurons in
the Dorsal Striatum Revealed by Single-Cell RNA Sequencing and PatchSeq. *Cell Rep.* **24**, 2179–
2190.e7 (2018).

71. T. Zerucha, T. Stühmer, G. Hatch, B. K. Park, Q. Long, G. Yu, A. Gambarotta, J. R. Schultz, J. L.
Rubenstein, M. Ekker, A highly conserved enhancer in the Dlx5/Dlx6 intergenic region is the site of
cross-regulatory interactions between Dlx genes in the embryonic forebrain. *J. Neurosci.* **20**, 709–
721 (2000).

72. T. Stuart, A. Srivastava, S. Madad, C. A. Lareau, R. Satija, Single-cell chromatin state analysis with
Signac. *Nat. Methods*. **18**, 1333–1341 (2021).

73. N. L. Jorstad, J. H. T. Song, D. Exposito-Alonso, H. Suresh, N. Castro, F. M. Krienen, A. M. Yanny,
J. Close, E. Gelfand, K. J. Travaglini, S. Basu, M. Beaudin, D. Bertagnolli, M. Crow, S.-L. Ding, J.
Eggermont, A. Glandon, J. Goldy, T. Kroes, B. Long, D. McMillen, T. Pham, C. Rimorin, K. Siletti,
S. Somasundaram, M. Tieu, A. Torkelson, K. Ward, G. Feng, W. D. Hopkins, T. Höllt, C. Dirk
Keene, S. Linnarsson, S. A. McCarroll, B. P. Lelieveldt, C. C. Sherwood, K. Smith, C. A. Walsh, A.
Dobin, J. Gillis, E. S. Lein, R. D. Hodge, T. E. Bakken, Comparative transcriptomics reveals human-
specific cortical features. *bioRxiv* (2022), p. 2022.09.19.508480.

74. D. Velmeshev, L. Schirmer, D. Jung, M. Haeussler, Y. Perez, S. Mayer, A. Bhaduri, N. Goyal, D. H.
Rowitch, A. R. Kriegstein, Single-cell genomics identifies cell type–specific molecular changes in
autism. *Science*. **364**, 685–689 (2019).

75. W. B. Ruzicka, S. Mohammadi, J. F. Fullard, J. Davila-Velderrain, S. Subburaju, D. R. Tso, M.
Hourihan, S. Jiang, H.-C. Lee, J. Bendl, G. Voloudakis, V. Haroutunian, G. E. Hoffman, P. Roussos,
M. Kellis, PsychENCODE Consortium, Single-cell multi-cohort dissection of the schizophrenia
transcriptome. *bioRxiv* (2022), , doi:10.1101/2022.08.31.22279406.

76. C. Nagy, M. Maitra, A. Tanti, M. Suderman, J.-F. Théroux, M. A. Davoli, K. Perlman, V. Yerko, Y.
C. Wang, S. J. Tripathy, P. Pavlidis, N. Mechawar, J. Ragoussis, G. Turecki, Single-nucleus
transcriptomics of the prefrontal cortex in major depressive disorder implicates oligodendrocyte
precursor cells and excitatory neurons. *Nat. Neurosci.* **23**, 771–781 (2020).

77. A. Sequeira, F. Mamdani, C. Ernst, M. P. Vawter, W. E. Bunney, V. Lebel, S. Rehal, T. Klempan,
A. Gratton, C. Benkelfat, G. A. Rouleau, N. Mechawar, G. Turecki, Global brain gene expression
analysis links glutamatergic and GABAergic alterations to suicide and major depression. *PLoS One*.
4, e6585 (2009).

78. D. Arendt, P. Y. Bertucci, K. Achim, J. M. Musser, Evolution of neuronal types and families. *Curr.*
Opin. Neurobiol. **56**, 144–152 (2019).

79. R. N. Delgado, D. E. Allen, M. G. Keefe, W. R. Mancia Leon, R. S. Ziffra, E. E. Crouch, A.
Alvarez-Buylla, T. J. Nowakowski, Individual human cortical progenitors can produce excitatory
and inhibitory neurons. *Nature*. **601**, 397–403 (2022).

80.   B. Martynoga, H. Morrison, D. J. Price, J. O. Mason, Foxg1 is required for specification of ventral
1398   telencephalon and region-specific regulation of dorsal telencephalic precursor proliferation and
1399   apoptosis. *Dev. Biol.* **283**, 113–127 (2005).

1400   81.   J. M. Tepper, F. Tecuapetla, T. Koós, O. Ibáñez-Sandoval, Heterogeneity and diversity of striatal
1401   GABAergic interneurons. *Front. Neuroanat.* **4**, 150 (2010).

1402   82.   J. DeFelipe, P. L. López-Cruz, R. Benavides-Piccione, C. Bielza, P. Larrañaga, S. Anderson, A.
1403   Burkhalter, B. Cauli, A. Fairén, D. Feldmeyer, G. Fishell, D. Fitzpatrick, T. F. Freund, G. González-
1404   Burgos, S. Hestrin, S. Hill, P. R. Hof, J. Huang, E. G. Jones, Y. Kawaguchi, Z. Kisvárday, Y.
1405   Kubota, D. A. Lewis, O. Marín, H. Markram, C. J. McBain, H. S. Meyer, H. Monyer, S. B. Nelson,
1406   K. Rockland, J. Rossier, J. L. R. Rubenstein, B. Rudy, M. Scanziani, G. M. Shepherd, C. C.
1407   Sherwood, J. F. Staiger, G. Tamás, A. Thomson, Y. Wang, R. Yuste, G. A. Ascoli, New insights into
1408   the classification and nomenclature of cortical GABAergic interneurons. *Nat. Rev. Neurosci.* **14**,
1409   202–216 (2013).

1410   83.   R. Tremblay, S. Lee, B. Rudy, GABAergic Interneurons in the Neocortex: From Cellular Properties
1411   to Circuits. *Neuron.* **91**, 260–292 (2016).

1412   84.   E. Boldog, T. E. Bakken, R. D. Hodge, M. Novotny, B. D. Aevermann, J. Baka, S. Bordé, J. L.
1413   Close, F. Diez-Fuertes, S.-L. Ding, N. Faragó, Á. K. Kocsis, B. Kovács, Z. Maltzer, J. M.
1414   McCorrison, J. A. Miller, G. Molnár, G. Oláh, A. Ozsvár, M. Rózsa, S. I. Shehata, K. A. Smith, S.
1415   M. Sunkin, D. N. Tran, P. Venepally, A. Wall, L. G. Puskás, P. Barzó, F. J. Steemers, N. J. Schork,
1416   R. H. Scheuermann, R. S. Lasken, E. S. Lein, G. Tamás, Transcriptomic and morphophysiological
1417   evidence for a specialized human cortical GABAergic cell type. *Nat. Neurosci.* **21**, 1185–1195
1418   (2018).

1419   85.   A. M. M. Sousa, Y. Zhu, M. A. Raghanti, R. R. Kitchen, M. Onorati, A. T. N. Tebbenkamp, B.
1420   Stutz, K. A. Meyer, M. Li, Y. I. Kawasawa, F. Liu, R. G. Perez, M. Mele, T. Carvalho, M. Skarica,
1421   F. O. Gulden, M. Pletikos, A. Shibata, A. R. Stephenson, M. K. Edler, J. J. Ely, J. D. Elsworth, T. L.
1422   Horvath, P. R. Hof, T. M. Hyde, J. E. Kleinman, D. R. Weinberger, M. Reimers, R. P. Lifton, S. M.
1423   Mane, J. P. Noonan, State, Matthew W, E. S. Lein, J. A. Knowles, T. Marques-Bonet, C. C.
1424   Sherwood, M. B. Gerstein, N. Šestan, Molecular and cellular reorganization of neural circuits in the
1425   human lineage. *Science.* **358**, 1027–1032 (2017).

1426   86.   G. Paxinos, C. Watson, M. Petrides, M. Rosa, H. Tokuno, *The marmoset brain in stereotaxic
1427   coordinates* (2012).

1428   87.   F. M. Krienen, M. Goldman, Q. Zhang, R. C H Del Rosario, M. Florio, R. Machold, A. Saunders, K.
1429   Levandowski, H. Zaniewski, B. Schuman, C. Wu, A. Lutservitz, C. D. Mullally, N. Reed, E. Bien, L.
1430   Bortolin, M. Fernandez-Otero, J. D. Lin, A. Wysoker, J. Nemesh, D. Kulp, M. Burns, V. Tkachev,
1431   R. Smith, C. A. Walsh, J. Dimidschstein, B. Rudy, L. S Kean, S. Berretta, G. Fishell, G. Feng, S. A.
1432   McCarroll, Innovations present in the primate interneuron repertoire. *Nature.* **586**, 262–269 (2020).

1433   88.   S. J. Fleming, M. D. Chaffin, A. Arduini, A.-D. Akkad, E. Banks, J. C. Marioni, A. A. Philippakis,
1434   P. T. Ellinor, M. Babadi, Unsupervised removal of systematic background noise from droplet-based
1435   single-cell experiments using CellBender. *bioRxiv* (2019), , doi:10.1101/791699.

1436   89.   C. S. McGinnis, L. M. Murrow, Z. J. Gartner, DoubletFinder: Doublet Detection in Single-Cell RNA
1437   Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst.* **8**, 329–337.e4 (2019).

90.   L. Feng, T. Zhao, J. Kim, neuTube 1.0: A New Design for Efficient Neuron Reconstruction Software

Based on the SWC Format. *eNeuro*. **2** (2015), doi:10.1523/ENEURO.0049-14.2014.

91. R. C. Challis, S. Ravindra Kumar, K. Y. Chan, C. Challis, K. Beadle, M. J. Jang, H. M. Kim, P. S.
Rajendran, J. D. Tompkins, K. Shivkumar, B. E. Deverman, V. Gradinaru, Systemic AAV vectors
for widespread and targeted gene delivery in rodents. *Nat. Protoc.* **14**, 379–414 (2019).

Acknowledgements:

We thank Tim Blosser for his early involvement in developing the spatial transcriptomics workflows. We
thank Atsushi Takahashi for assistance with MRI scanning for animals receiving local viral injections.
We thank the MIT veterinarian staff for animal husbandry and for their assistance with surgical
procedures. We thank Monika Burns and Yuanyuan Hou for assistance with AAV IV injections and
animal perfusions.

1449

Funding:
National Institutes of Health grant U01MH114819 (GF, SAM, EB)
The National Institute of Neurological Disorders and Stroke grant UG3NS111689 (BED)
NSF GRFP # 1745302 (MES)
MathWorks Science Fellowship (MES)
Collamore-Rogers Fellowship at MIT (MES)
NSF GRFP # 1122374 (TWS)
Broad Institute's Stanley Center for Psychiatric Research (SAM, GF, BED)
Dean's Innovation Award (Harvard Medical School) (SAM)
Hock E. Tan and K. Lisa Yang Center for Autism Research at MIT (GF)
Poitras Center for Psychiatric Disorders Research at MIT (GF)
McGovern Institute for Brain Research at MIT (GF)

1462
Author Contributions:
RNA/ATAC Data Generation: FMK, MG, AL
Spatial Data Generation: KML, HZ
Data Analysis: FMK, KML, HZ, RCHdR, MES, MG, AL, KXL, VFB-G, TWS, SAM
Data Interpretation: FMK, KML, HZ, RCHdR, MES, MG, KXL, VFB-G, SB, EB, SAM, GF
Tissue Samples/Tissue Processing: FMK, KML, HZ, MG, AL, QZ, GC, SB
AAV design, generation & experiments: MW, CC, QZ, JS, SJV, JD, KC, BED
Morphology Data generation/analysis: KML, HZ, VFB-G, TWS, AM, EB
Software/Data management: FMK, RCHdR, MG, AW, JN, SK
Writing: FMK, KML, HZ, MES, VFB-G, SAM, GF

1473

1474 **Competing interests:** Authors declare that they have no competing interests.

1475

Data and materials availability:
Raw sequence data were produced as part of the BRAIN Initiative Cell Census Network are available for
download from the Neuroscience Multi-omics Archive
(https://assets.nemoarchive.org/dat-1je0mn3) and the Brain Cell Data Center (https://biccn.org/data).
Morphological reconstructions and single molecule FISH of interneuron types are available for download
through the Brain Image Library (https://submit.brainimagelibrary.org/search?grant_number=1-U01-

MH114819-01). The AAV-BI103 rep-cap plasmid will be made available through Addgene upon
publication of the characterization of this capsid and through direct requests to Ben Deverman.

Supplementary Materials:

Figs. S1-S11
Table S1-S7
Data S1-S2

Legends:

FIGURE 1

1489

Figure 1. Single nucleus RNA sequencing of marmoset brain. **(A)** Number of nuclei per brain structure. Colors indicate cell classes. **(B)** Proportions of cell classes across brain structures. **(C)** Neurons in each dissected region are clustered separately, then pseudo-bulked "metacells" of all 288 clusters are arranged in the dendrogram using hierarchical clustering of top 100 PCA scores of expressed genes. Outer rings colored by dissected subregion (1st ring), cephalic vescicle (2nd ring) and major neurotransmitter usage (3rd ring). Seven major "clades" are colored and labeled according to cell type and regional composition of clade. **(D)** Expression of markers for glutamatergic neurons (*SLC17A7*, *SLC17A6*), GABAergic neurons (*GAD1*), and transcription factors (*FOXG1*, *ZFHX4*, *SOX14, DLX1, OTX2, LHX9*) are plotted as heatmaps on dendrogram shown in (C). These transcription factors are largely restricted to specific clades or are associated with particular cephalic vesicles. **(E)** Gene-gene correlation (Spearman tau) distributions across all neuron clusters in (C) for *SLC17A7*, *SLC17A6*, and *CHAT* (marker for cholinergic neurons) as well as background (all genes). The distribution of pairwise correlations to *CHAT* had a lower standard deviation (mean $r = 0.002$, std. dev $= 0.116$) relative to baseline gene-gene correlations (mean $r = 0.02$, std. dev $= 0.199$; $F(5078, 17315809) = 0.33$, p-adj $< 1e\text{-}15$). In contrast, pairwise correlations to *SLC17A7* were much broader relative to the background distribution of all gene-gene correlations. **(F)** Distribution of cross-cell-type correlation to *SLC17A6* of genes most correlated with *SLC17A7* (top 116 genes with Spearman tau > 0.5 to *SLC17A7*). **(G)** Same hierarchical clustering procedure as (C) but for non-neuronal cell types. Outer ring colors indicate major non-neuronal cell class. Inner ring colors indicate region dissection and vesicle; colors as in (C).

FIGURE 2

1509
Figure 2. Regional variation in neocortical cell types and expression patterns. (A) Cortical regions
sampled. **(B)** t-SNE embeddings of neocortical macroglia (astrocytes, oligodendrocyte lineage types) and
neurons (GABAergic interneurons, glutamatergic neurons). Colors represent clusters (numbered). **(C)**

Same as (B) but cells colored by neocortical dissection. **(D)** Regional proportions of each cluster, colors

same as (A). **(E)** *VCAN* expression across astrocyte clusters and oligodendrocyte lineage clusters. Colors

as in (B). **(F)** Venn diagrams showing overlap of neocortical regionally differentially expressed genes

(rDEGs) across GABAergic neurons, glutamatergic neurons, astrocytes and oligodendrocyte lineage cells.

rDEGs are defined as > 3 fold expression difference in homologous cell types across pairs of cortical

regions.

FIGURE 3

1519
Figure 3. Examples suggesting cross-cephalic-boundary cell type migration. (A) Locations of
amygdala clusters in the dendrogram from **Fig. 1C**. **(B)** Clustering of marmoset amygdala cells (n =

44,165 neurons) predicted to be glutamatergic (top row) or GABAergic (bottom row) based on a linear

classifier (scPred) trained on supervised cell type models. t-SNEs for each class are colored by cluster

(first column) or genes (*SLC17A6*, *GAD1, ADCYAP1, OTP, SIM1*). Arrowhead indicates cluster of

neurons that are classified as GABAergic yet do not express *GAD1* and do express *SLC17A6*. **(C)**

snRNA-seq from mouse amygdala neurons. Heatmap shows normalized, scaled expression of

neurotransmitters in neurons from snRNA-seq of adult mouse amygdala (n = 25,930 nuclei, 3 replicates

pooled). Dendrogram shows hierarchical clustering of neuron types. Dotted outline shows presence of 3

Slc17a6+ subtypes that preferentially cluster with GABAergic (*GAD1+/GAD2+*) subtypes and that

express *Sim1* and/or *Adcyap1*. **(D)** Cartoon of embryonic migration of *Otp+* cells that migrate from

proliferative zones around the 3rd ventricle to periventricular hypothalamus (vzPZH) and medial

amygdala (MeA), following data in Garcia-Moreno et al. (*45*). **(E)** ISH for *Otp* in sagittal section of P28

mouse brain. Dotted outline indicates borders of medial amygdala (MeA). **(F)** Marmoset P0 coronal

section showing ISH staining for *OTP*. Data from (https://gene-atlas.brainminds.jp/; (Shimogori et al.

2018; Kita et al. 2021)). **(G)** Cartoon of marmoset amygdala the sagittal plane. FISH staining for *GAD1*

(magenta), *SLC17A6* (cyan), and *SIM1* (yellow) in the medial amygdala (Me). Magenta arrows highlight

GAD1 expressing nuclei. Green arrows highlight *SLC17A6* and *SIM1* dual expressing nuclei. **(H)**

Marmoset neuronal dendrogram shown in **Fig. 1C** with clades colored by *TAC3* expression.

Unexpectedly, a novel primate-specific striatal *TAC3* population (*3*) clusters with hypothalamic and basal

forebrain populations, and not with other telencephalic GABAergic interneurons. Red arrowhead

indicates clade containing three *TAC3+* populations found in striatum, basal forebrain, and hypothalamus.

(I) Heatmap of genes expressed in the clade indicated by red arrowhead in (H). Expression values of each

gene is normalized to its max within the clade. The three *TAC3+* populations are labeled and each show

distinct expression differences.

FIGURE 4

1545
Figure 4. Cell-type-specific distributions of interneurons in primate telencephalon using
quantitative smFISH. (A) t-SNE of GABAergic neocortical interneurons as in **Fig. 2B**, colored by

1548 subclass marker (*PVALB, SST, VIP, LAMP5, CXCL14*). Gray points are the *MEIS2+* population that is

1549 restricted to orbiomedial prefrontal cortex and was not spatially profiled. **(B)** Medial-lateral proportions of

1550 each major class as percentage of all cells (DAPI+). Barplots quantify positive cells as proportion of all

1551 (DAPI+) cells from medial to lateral sections of smFISH performed on 6 thin (16 µm) sagittal sections of

1552 marmoset neocortex, each section being 1600 µm apart and covering 9600 µm of neocortex. Colors as in

1553 (A). **(C)** smFISH for neocortical interneuron subclass markers showing locations of cells positive for each

1554 marker across 6 sagittal sections of the marmoset neocortex. First column shows density of all DAPI+

1555 nuclei per unit area (approximately 387 µm per bin) profiled from one series. Heatmap scale in

1556 subsequent columns show percentage of marker-positive cells relative to DAPI+ cells. Average

1557 proportions across section shown in (B). Med. = medial, Lat. = Lateral. Asterisks in section ii and iii of

1558 the SST series and iv of the LAMP5 series denote regions where tissue or staining artifacts caused loss of

1559 signal. These can be seen in the DAPI-only series of these experiments, which show lower overall cell

1560 counts at these locations (**Fig S7**). **(D)** Quantitation of interneuron proportions by cortical area parcellated

1561 according to **Fig. S8B-C**. **(E)** Relative percentages of interneuron proportions by cortical area shown in

1562 Fig. S8B-C, sorted by max relative proportion of *VIP*+ interneurons. **(F)** t-SNE of striatal, cholinergic

1563 neurons (*SLC5A7*) and GABAergic interneurons (*SST*, *PVALB*, *TH*, *CCK*, *TAC3*). Green points

1564 correspond to the *MEIS2+* population and were not spatially profiled. **(G)** Medial-lateral gradients of

1565 striatal cell type proportions across 4 sagittal sections. Dots = individual replicates. Colors as in (F). **(H)**

1566 smFISH for striatal cholinergic and interneuron subclass markers showing locations of cells positive for

1567 each subclass marker across 4 (16 µm) sagittal sections of the marmoset striatum, each 1600 µm apart

1568 covering 6400 µm in total. First column shows the anatomical context of the four sagittal striatal planes.

1569 Average proportions across each section are shown in (G).

FIGURE 5

1570
Figure 5. Morphological characteristics of marmoset cortical and striatal interneurons. (A-B) *Top*
rows, Examples of AAV9-hDlx5/6-GFP-fGFP labeled neocortical **(A)** and striatal **(B)** interneurons. Insets

show magnified cell nucleus of GFP+ cell along with smFISH staining for interneuron type marker to

confirm molecular identity of labeled cell. Scale Bar = 100 µm. *Bottom rows,* Reconstructed skeletonized

morphology (Imaris) of GFP+ cells depicted in *top rows*. Scale Bar = 50 µm. **(C)** Representative

reconstructed neocortical and striatal *PVALB+* interneurons from one marmoset (Cj 17-154; **Table S1,**

S6). Scale Bar = 50 µm. **(D)** Quantification of morphological characteristics of *PVALB+* cells from Cj 17-

154. Means ± SEM. *$*P < 0.05$* and *$**P < 0.01$*. See the Methods section for detailed statistical

information.

FIGURE 6

A Striatum → snATAC-seq

int_tac3
int_pvalb_th
int_sst

UMAP_2

UMAP_1

B Normalized accessibility (range 0 – 240)

astrocyte
dSPN
endothelia
ependyma
int_chat
int_pvalb_th
int_sst
int_tac3
iSPN
microglia
gad_foxp2
glut_satb2
oligo
OPC

50687000   50689500
chr8 position (bp)

C Normalized accessibility (range 0 – 530)

101132000   101134000
chr4 position (bp)

D Tandem enhancers
ITR WPRE $P_{minCMV2}$ EGFP r.s. WPRE pA ITR

E

F Cd (G) Pu GP

G merge   AAV9-tandemE-TAC3-GFP   TAC3   50µm

H 50µm

I i PFC ii Striatum iii Pons iv Cerebellum
AAV-BI103-tandemE-TAC3-EGFP

J Striatal TAC3   50µm

K Whole cell | Somatic | Neuritic

Total surface area x10³ | Total volume x10³ | Total length | Total area x10³ | Soma size | Num branches | Num 1° branches
TAC3 | TAC3 | TAC3 | TAC3 | TAC3 | TAC3 | TAC3

1580

Figure 6. Development of a novel enhancer-AAV for *TAC3*+ primate-specific striatal interneurons.
(A) snATAC-seq (69,808 nuclei) from fresh marmoset striatum (1 male, Cj 18-153). UMAP shows major
clusters with labels transferred from striatal snRNA-seq data. **(B)** Chromatin accessibility at the locus
corresponding to the mDlx sequence in marmoset shows read pileups in *PVALB*, *SST*, and *CHAT*+
neurons, but not in the *TAC3*+ type. **(C)** Chromatin accessibility of a candidate selective enhancer for the
TAC3+ type. **(D)** Sequence construct design for *TAC3* interneuron specific AAV. Four *TAC3* interneuron-
specific regulatory elements (example in (C)) are packaged in tandem upstream of a minimal promoter
driving EGFP expression. ITR = inverted terminal repeats. WPRE = Woodchuck Hepatitis Virus
Posttranscriptional Regulatory Element. PminCMV2 = minimal CMV2 promoter. EGFP = enhanced
green fluorescent protein. R.S. = 300 bp random sequence. pA = polyadenylation sequence. **(E)** MRI
showing injection location (white arrowheads) of the AAV9-tandemE-TAC3-EGFP virus into bilateral
dorsal striatum (caudate) in one animal (Cj 19-207, **Table S1**). **(F)** Cartoon showing location of cell
shown in (G). **(G)** Main: EGFP antibody-amplified confocal image of a labeled cell (position shown in
(F)). Insets: smFISH for *TAC3* showing colocalization. Scale bar = 50 µm. **(H)** Morphological
reconstruction (Imaris) of cell shown in (G). Scale bar = 50 µm. **(I)** Whole sagittal section (20x image) of
adult marmoset showing cells transduced by the AAV-BI103-tandemE-TAC3-EGFP virus in marmoset
Cj 20-214. GFP+/*TAC3*+ cells were detected sparsely in neocortex as well as striatum, cerebellum,
substantia nigra, superior colliculus, and brainstem. (*i*) Prefrontal cortex, (*ii*) Striatum, (*iii*) Pons, (*iv*)
Cerebellum. Green arrows highlight cells in areas with sparse expression. **(J)** Four reconstructions
(Imaris) of striatal GFP+/*TAC3*+ cells from similar section shown in (I). **(K)** Quantification of
morphological parameters of reconstructed cells from striatal GFP+/*TAC3*+ cells from the local injection
in (E) as well as the systemic injection in (I) (**Table S6**).

Supp S1

Figure S1. Neuron counts by donor across brain regions. (A) Neuronal dendrogram as in **Fig. 1C**, with outer barplots depicting number of nuclei per cell type and replicate. Ring colors are brain regions,

colors in barplots correspond to replicates. **(B)** Proportional per-cluster representation of PFC neurons between young adult donors and aged (n=2) donors. While our snRNA-seq collection focused on post-sexual maturity young adults, we acquired an additional dataset of PFC sampled from 2 aged animals (1 M, 11y0m; 1F, 14y4m, 37,260 cells total; **Table S1**). Individual replicates contributed similar proportions of neurons to each prefrontal neuron subtype, and clusters generally had proportional representation across young adults and aged animals, as well as across males and females (**Fig. S1B-C**), suggesting that these variables do not dramatically impact neuronal ensembles and identities in prefrontal cortex. **(C)** t-SNE embeddings of PFC neurons (*top row*, GABAergic; *bottom row*, glutamatergic) with colors representing different metadata: age (young vs aged), replicate, PFC subregion. There was notable enrichment of *MEIS2+* GABAergic neurons in medial prefrontal and orbital prefrontal dissections **(Fig. S1C)**. Based on their gene expression profiles, these cells likely correspond to the recently described population of LGE-derived *MEIS2+* neurons that populate the olfactory bulb in mice, and which are instead directed to medial prefrontal cortex in macaques and humans (*33*). **(D)** t-SNEs of neurons in each brain structure, with cells colored by replicate (colors as in (A)). Telencephalic neurons are plotted separately by class: GABAergic and glutamatergic classes (neocortex, hippocampus, amygdala), or GABAergic interneurons and spiny projection neurons (striatum). Compared with neocortex, greater cross-donor variability was observed in some subcortical structures such as hypothalamus and thalamus, though this was likely driven more by dissection variability than by donor variability, as the donor-specific clusters tended to be from subregions that were only sampled in one individual (see E). **(E)** t-SNEs of thalamic neurons with cells colored by thalamic subdivision.

Figure S2. Conservation of neuronal hierarchy across species and clustering methods. (A) Comparison of marmoset and mouse neurons (mouse atlas data from (*6*)) using three different distance

calculations for hierarchical clustering: HCA_{corr} distance = gene expression correlations of top 5904 genes

in marmoset or 3528 in mouse (mouse genes include all genes with 1:1 orthologs to the 5904 marmoset

genes and that are expressed in at least 10 transcripts per 100,000 in at least one mouse neuron type);

HCA_{PCA} distance = top 100 PCA scores across the same genes; HCA_{TF-PCA} distance = top 100 PCA scores

using expressed transcription factors only (marmoset = 349 TFs, mouse = 450 TFs). **(B)** Distance between

hierarchical clustering (HC) dendrogram trees computed using different methods. Cyan line = tree

distance (R package TreeDist) between hierarchical clustering using distance = HCA_{corr} and using

distance = HCA_{PCA}. Pink distribution is tree distance scores between HCA_{corr} and shuffled PCA scores (n

= 100 shuffling iterations). Lower values of tree_distance (x-axis) mean higher agreement between

dendrogram tree structures. **(C)** PCA loadings and top genes for PC2-PC4. PC scores are plotted on the

HCA_{PCA} dendrogram. Ranked gene loading plots show top 20 genes per PC. **(D)** Tree distances computed

as in (B) between HCA_{PCA} and HCA_{TF-PCA}. The tree distances between these two trees is low, but not

different from distributions of random, same-sized sets of genes. **(E)** Marmoset dendrogram in **Fig. 1C**

(HCA_{PCA}) indicating major clades compared in (F). **(F)** Ancestral reconstruction (AR; R package

phytools) of gene expression profiles of major clades of marmoset neuron types from dendrogram in (E).

Maximum likelihood estimates of gene expression (fastAnc) were computed for 7 major internal nodes

(gray circles) of the HCA_{PCA} dendrogram. Scatterplots show pairwise comparisons between AR of

internal nodes of major clades. Blue dots = genes with >3 foldchange difference between the two ARs.

Magenta circles = differentially expressed transcription factors (DE-TFs). Chi-square and p-values

describe whether TFs are significantly differentially enriched between the AR pairs.

1650

Supp S3

1651
Figure S3. Gene expression across neural populations. Expression of broad class marker genes and
other genes of interest across all neurons sampled by snRNA-seq. Heatmap colors are scaled to max

1654   normalized expression for each row (gene). Dendrogram ordering and metadata colors as in **Fig. 1C**. Cell

1655   types discussed in the main text are labeled at bottom.

1656

Supp S4 related to Fig 2

1657
Figure S4. Glia diversity across regions. **(A)** t-SNE embeddings of major non-neuronal types colored
1658 by cluster. **(B)** Barplots of glial proportions colored by donor and by sex. **(C)** Non-neuronal nuclei counts.
1659

Colors indicate donor, same as (B). **(D)** Expression of marker genes in non-neurons. Genes as in (*8*).

Heatmap colors are scaled to max normalized expression for each row (gene). Dendrogram and metadata

colors as in **Fig. 1G**. **(E)** Differentially expressed genes in astrocytes across cephalic compartments. **(F)**

Tissue validation for astrocyte differentially expressed genes (*FOXG1, SPARC*) in coronal sections of

marmoset brain. Green dots indicate locations of cells that stain positive for *GFAP* (IHC, mAb **Table S4**)

and *FOXG1* (smFISH). Magenta dots indicate cell positions for *GFAP* (IHC) and *SPARC* (smFISH). **(G)**

Cartoon of coronal section imaged; Red boxes (*i-ii*) correspond to tissue validation in (H). **(H)** (*Left)*

Fields of view from neocortex and thalamus stained for *GFAP* antibody (gray), *FOXG1* (green), and

SPARC (magenta). Green arrows highlight *GFAP* cells colocalized with *FOXG1*, magenta arrows

highlight *GFAP* cells colocalized with *SPARC*. (*Right*) Magnified examples of double positive cells in

neocortex and thalamus. Ctx = cortex, Thal = thalamus. **(I)** (*Left*) Cartoon of sagittal section imaged; red

boxes (*i-iii*) correspond to (*Right*) tissue validation of increased abundance of *VCAN*+ astrocytes in adult

marmoset V1-adjacent white matter (*iii*) compared with PFC-adjacent white matter (*i*) and V1 gray

matter (*ii*). GFAP antibody (gray) combined with smFISH probes against *VCAN* (magenta) and *GFAP*

(green). Green arrows correspond to *GFAP*+ (antibody), *GFAP*+ (smFISH) cells. White arrows

correspond to *GFAP*+ (antibody), *GFAP*+ (smFISH), and *VCAN*+ cells. V1 = visual cortex V1, PFC =

prefrontal cortex, GM = gray matter, WM = white matter. Red arrow highlights locale of *VCAN*+ *GFAP*+

images. Scale bar = 10 µm.

Supp S5 related to Fig 2

1678
1679 Figure S5. Cortical rDEGs are defined across cell types and do not reflect ambient RNA
1680 contamination. (A) *(i)* rDEG ambient score distributions (from CellBender) in temporal cortex samples,

which had amongst the highest numbers of rDEGs compared to other neocortical regions. Despite

glutamatergic neurons being more numerous and having more expressed genes/transcripts per cell, the

median ambient contamination scores for glutamatergic rDEGs were not higher than median

contamination scores for GABAergic rDEGs. rDEGs shared between glutamatergic and GABAergic

neurons had indistinguishable scores compared with rDEGs private to one neuronal class. *(ii)* Ambient

scores in temporal cortex of temporal cortex rDEGs are indistinguishable from V1 rDEGs. *(iii)*

Distributions of temporal cortex rDEG ambient scores by neuron class, again showing no difference

between rDEGs that are shared or private to a neuronal class. **(B)** Numbers of regionally differentially

expressed genes (rDEGs) between pairs of cortical regions for neurons, astrocytes, and oligodendrocyte

lineage types.

1691

Figure S6. Locations of *TAC3*+ cells in marmoset forebrain. (A) smFISH reveals anatomical locations

and expression levels *TAC3*+ types in different brain regions. *(i)* Schematic of *TAC3*+ cells imaged across

cortex, dorsal striatum. Ctx = neocortex, Cd = Caudate. *(ii)* Cartoon close-up of nuclei in striatum, basal

forebrain and hypothalamus. Magenta stars = locations of *TAC3+* cells in lower image panel. Cd =

Caudate, AcbC = nucleus accumbens core, AcbSh = nucleus accumbens shell, SI = Substantia

innominata, B = basal nucleus of Meynert, EAM = extended amygdala, medial, EAC = extended

amygdala, central, ac = anterior commissure, BSTLP = bed nuc st, lateral posterior, BSTLJ = bed nuc st,

juxtacap, BSTLV = bed nuc st, lateral ventral, LH = lateral hypothalamus, VLH = ventrolateral

hypothalamus, LPO = lateral preoptic area. **(B)** Density and location of *TAC3+* cells as proportion of all

DAPI+ cells. Barplots show percentages in bins (approximately 1,290 µm per bin) taken across the

anterior-posterior (top) and dorsal-ventral (left side) axes. **(C)** Effect on placement of the *TAC3+* striatal

type on the neuronal dendrogram when omitting the two *TAC3+* types in hypothalamus and basal

forebrain. When these types are omitted and hierarchical clustering is repeated (using HCA_{PCA}), the

TAC3+ striatal type is most similar to *PVALB+* striatal interneurons, consistent with previous reports that

only compared telencephalic interneurons (*3*).

1707

FIGURE S7

Figure S7. Total cell numbers across marmoset neocortex. (A) Total numbers of DAPI+ cells per unit area (approximately 387 µm per bin) for each of the sections shown in **Fig. 4C**.

Figure S8. Interneuron numbers across marmoset neocortex. (A) smFISH for neocortical interneuron
subclass markers showing locations of cells positive for each marker across 6 sagittal sections of the
marmoset neocortex. Heatmap scale shows absolute density per unit area (approximately 387 µm per
bin). First column shows DAPI and area profiled. **(B)** (*Left, middle)* Quantitation of interneuron
proportions by cortical area in marmoset parcellated according to Fig. S8C. *(Right)* Quantitation of
interneuron proportions by cortical area in mouse reproduced from Kim et al. (2017). (*Left*) Absolute
percentages of *SST, PVALB,* and *VIP* populations. (*Middle*) Same as (*left),* but scaled as proportions to
100%. (*Right*) Schematic describing ratios of major interneuron types (*Sst+, Pvalb+*) in mouse from Kim
et al. (2017). **(C)** Cartoons of cortical areas and areal groupings used to bin smFISH neocortical
interneuron proportions in (B) and **Fig. 4D-E**. Neocortical parcellation from
https://doi.org/10.24475/bma.4520. **(D)** Examples of smFISH images quantitated in **Fig. 4B-E**. Panels for
each marker show example positive cells in anterior, middle, and posterior locations across neocortex.
White arrows indicate positive cells. Scale bar = 20 µm.

FIGURE S9

1726
Figure S9. Interneuron numbers across marmoset striatum. (A) Mouse striatal *in situ* from Allen
Brain Atlas for *Sst, Pvalb, Slc5a7, Cck, Th*. **(B)** Cartoon of marmoset striatum illustrates area profiled,

1729   medial to lateral. Proportions of *DRD1+* (dark blue) and *DRD2+* (dark green) cells across primate
1730   striatum calculated as in **Fig. 4H**.

1731

1732   **Figure S10. Morphology examples using NeuTube reconstructions. (A)** Example morphological

1733   reconstructions of striatal and neocortical interneurons using the NeuTube pipeline. Each cell, along with

1734   associated smFISH staining, is available for download at https://doi.org/10.35077/g.609.

Figure S11

1735
Figure S11. Examples of striatal and peri-striatal *TAC3*+ neurons labeled by AAV-tandemE-TAC3-
1736
EGFP. (A) MRI showing injection location (white arrowheads) of virus into bilateral dorsal striatum

(caudate) in one animal (Cj 17-B111). **(B)** EGFP antibody-amplified confocal image of a labeled cell
(position shown in (C) with smFISH for *TAC3* showing colocalization). **(C)** Cartoon showing location of
positive cells in (B) as well as labeled cells in transition zone (D). **(D)** Examples of extra-striatal labeled
cells from injections in (A). **(E)** Position of cells in (D). Cd = Caudate, SI = Substantia innominata, B =
basal nucleus of Meynert, ac = anterior commissure, BSTLP = bed nuc st, lateral posterior, BSTLJ = bed
nuc st, juxtacap, 3V = third ventricle, LV = lateral ventricle.

Table S1. Marmoset sample information. Table of animals, metadata, and experimental information.

Table S2. snRNA-seq dataset by donor and brain area. Tables of the number of cells per brain
structure and samples per donor.

Table S3. Neocortical rDEGs across three donors. Neocortical rDEGs across three donors with
pairwise comparisons between neocortical locations for major clusters of cortical excitatory neurons,
inhibitory neurons, astrocytes and oligodendrocyte lineage types.

Table S4. Fluorescent *in situ* hybridization probes and antibodies used. A list of all FISH probes
(RNA-Scope and Molecular Instruments) and antibodies used for validation of gene and protein
expression *in situ*.

Table S5. Morphological reconstructions performed with Neutube. A list of all cells reconstructed
with Neutube with their corresponding morphological measurements.

Table S6. Morphological reconstructions performed with Imaris. A list of all cells reconstructed with
Imaris with their corresponding morphological measurements.

Table S7. Links and DOIs. A list of all links and DOIs referenced.

Data S1. Striatal and cortical medio-lateral quantification. Tabular data for striatal and cortical
medio-lateral histograms/bar plots.

Data S2. *TAC3* medio-lateral quantification. Tabular data for *TAC3* medio-lateral histograms and bar
plots.