1   **Title**
2
3   **Type I PRMT inhibitor MS023 promotes *SMN2* exon 7 inclusion and synergizes with**
4   **nusinersen to rescue the phenotype of SMA mice**
5

6   Anna J Kordala[1,3], Nina Ahlskog[2,3], Muhammad Hanifi [2], Amarjit Bhomra[2,3], Jessica
7   Stoodley[2,3], Wooi Fang Lim[2,3], Suzan M Hammond[2], Matthew JA Wood[2,3], Carlo Rinaldi[2,3]
8
9   [1]Department of Physiology Anatomy and Genetics, University of Oxford, Oxford, UK
10   [2]Department of Paediatrics, University of Oxford, Oxford, UK
11   [3]MDUK Oxford Neuromuscular Centre, Oxford, UK
12
13   Address correspondence to: Carlo Rinaldi, Institute of Developmental and Regenerative
14   medicine (IDRM), Old Road Campus, University of Oxford, OX3 7TY Oxford, United
15   Kingdom. Phone: +441865272148; email: carlo.rinaldi@idrm.ox.c.uk.

16
17   The authors have declared that no conflict of interest exists.

Abstract
Spinal muscular atrophy (SMA) is the leading genetic cause of infant mortality. The advent
of approved treatments for this devastating condition has significantly changed SMA
patients' life expectancy and quality of life. Nevertheless, these are not without limitations,
and research efforts are underway to develop new approaches to be used alone and in
combination, to ensure improved and long-lasting benefits for SMA patients. Protein arginine
methyltransferases (PRMT) are emerging as druggable epigenetic targets, with several
small molecule PRMT inhibitors already in clinical trial stage. From a screen of highly potent
and selective next generation epigenetic small molecules, we have identified MS023, a
potent and selective type I PRMT inhibitor, able to promote *SMN2* exon 7 inclusion and
increase SMN protein levels in preclinical SMA model, by inhibiting the binding of splicing
factor hnRNPA1 to *SMN2* pre-mRNA. Treatment of SMA mice with MS023 results in
amelioration of the disease phenotype, with strong synergistic amplification of the positive
effect when delivered in combination with the *SMN2*-targeting antisense oligonucleotide
nusinersen. Moreover, transcriptomic analysis revealed that MS023 treatment has very
minimal off-target effects and that the added benefit of the combination therapy is mainly
attributable to targeting neuroinflammation. Our study warrants further clinical investigation
of PRMT inhibition both as a stand-alone and add-on therapy for SMA patients.

Introduction

Spinal muscular atrophy (SMA) is a genetic neuromuscular condition, affecting 1:8,000–10,000 live births (Pearn 1978) and, to this date, a leading inherited cause of infant mortality worldwide (Crawford and Pardo 1996). SMA is caused by inactivating mutations, mainly homozygous deletions, in the survival motor neuron 1 (*SMN1)* gene on chromosome 5 (Lefebvre et al. 1995). The encoded protein SMN is ubiquitously expressed, localizing both to the cytoplasmic and nuclear compartments within the cell, where it exerts numerous essential functions, including biogenesis of small nuclear ribonucleoproteins (snRNPs), its most widely studied function (Friesen, Massenet, et al. 2001; Pellizzoni, Yong, and Dreyfuss 2002; R. Zhang et al. 2011), 3' processing of histone mRNAs (Tisdale et al. 2013), control of transcription (Strasswimmer et al. 1999; Suraweera et al. 2009; Yanling Zhao et al. 2016), R-loop resolution (Suraweera et al. 2009; Yanling Zhao et al. 2016), RNA trafficking (Akten et al. 2011; Fallini et al. 2011, 2014, 2016; Hubers et al. 2011; Piazzon et al. 2008; Rage et al. 2013; Rossoll et al. 2003; Tadesse et al. 2008) and pre-mRNA splicing (Charroux et al. 1999; Makarov et al. 2012; Pellizzoni et al. 1998; Shafey et al. 2010). The *SMN2* gene is a centromeric copy of telomeric *SMN1*, with a critical C to T substitution in position 6 of exon 7, which creates an exonic splicing silencer site and is recognised by a splicing factor, hnRNPA1 (Kashima et al. 2007; Kashima and Manley 2003). As a consequence, the *SMN2* gene mainly encodes a shorter and rapidly degraded SMN isoform lacking exon 7 (Δ7 SMN), with only 10–15% of *SMN2* transcripts still capable of generating a full length SMN protein (Lorson et al. 1999; Monani et al. 1999). The number of *SMN2* copies varies in the general population and is the main modifier of disease severity identified so far, with higher number of copies being associated with a milder SMA phenotype (Feldkötter et al. 2002; McAndrew et al. 1997). Depending on the age of onset and motor milestones achieved, SMA has been divided into four clinical types (I–IV) (Munsat and Davies 1992), with type I SMA infants showing symptoms before 6 months of age and never gaining the ability to sit unaided.

Loss of SMN leads to degeneration of lower α-motor neurons by molecular mechanisms which are not fully understood. Other neuronal and non-neuronal cell populations are affected in SMA patients and include sensory neurons (Gogliotti et al. 2012; Jablonka et al. 2006; Martinez et al. 2012; Mentis et al. 2011; Rudnik-Schöneborn et al. 2003), skeletal muscle (Arnold et al. 2004; Kim et al. 2020; Martínez-Hernández et al. 2009) heart and vasculature (Araujo, Araujo, and Swoboda 2009; Bevan et al. 2010; Heier et al. 2010; Lipnick et al. 2019; Rudnik-Schöneborn et al. 2010; Somers et al. 2016), liver (Crawford et al. 1999; Deguise et al. 2019), and pancreas (Bowerman et al. 2012), supporting the notion that SMA is a multisystemic condition.

In the last decade, the advent of successful therapeutic approaches, combined with improvements in standard of care, have effectively changed the course of this disease, significantly slowing the progression of all SMA types (Harding et al. 2015; Eugenio Mercuri et al. 2020, 2022). One of such approved strategies entails intrathecal injections of nusinersen, an antisense oligonucleotide (ASO) targeting intronic splicing silencer N1 (ISS-N1), promoting *SMN2* exon 7 inclusion and increasing levels of full length SMN protein. Concomitantly, these treatment opportunities pose new challenges, including their yet-to-be determined long-term effects, the rise of new phenotypes in treated patients, which is particularly relevant for approaches such as nusinersen, solely targeting the central nervous system (CNS), the need for repeated invasive administrations, and high costs. Altogether,

these considerations highlight the urgency for development of therapeutic combinations to
address these important limitations and to provide additional benefit to patients.
Epigenetic regulation of gene expression, which involves covalent and sequence-specific
modifications of histone and non-histone proteins, is a dynamic and reversible process that
establishes normal cellular phenotypes and, when dysregulated, contributes to a wide range
of human diseases, including SMA (Allis et al. 2007; Cao et al. 2016; Hauke et al. 2009;
Murray et al. 2015; Portela and Esteller 2010; Zheleznyakova et al. 2013). In recent years,
key protein families that mediate epigenetic signalling through the acetylation and
methylation of histones and non-histone proteins, including histone deacetylases (HDACs),
protein methyltransferases (PRMT), histone lysine methyltransferases (KMTs) and
demethylases (KDMs), and bromodomain-containing proteins (BRD) have emerged as
attractive druggable targets using small molecules, due to the dynamic nature of disease-
associated epigenetic states (Arrowsmith et al. 2012). Several small molecule inhibitors of
histone deacetylases have been tested in SMA models (Mohseni, Zabidi-Hussin, and
Sasongko 2013), but their lack of specificity, low potency, and poor understanding of their
mechanisms of action have significantly limited their translation into the clinic (Kissel et al.
2014; Krosschell et al. 2018; E. Mercuri et al. 2007). A recent study has shown that
nusinersen, while promoting exon 7 inclusion, also induces a silencing histone mark
H3K9me2 on *SMN2* gene, creating a roadblock to RNA polymerase II elongation. The
histone deacetylase inhibitor valproic acid counteracts the chromatin effects of the ASO,
resulting in higher exon 7 inclusion upon combined treatment compared to nusinersen alone
(Marasco et al. 2022). The primary aim of this project is to identify next generation small
molecules targeting epigenetic proteins able to increase SMN protein and evaluate their
therapeutic potential in SMA animal models alone and as an add-on treatment.
25
Results
Epigenetic screening of SMN2 modulators. In order to identify small molecules that
selectively modulate *SMN2* pre-mRNA splicing to include exon 7, we performed a cell-based
screen in SMA type II-patient derived fibroblasts carrying 3 copies of the *SMN2* gene, using
a collection of 54 chemical probes from the Structural Genomics Consortium (SGC)
collection (https://www.thesgc.org/chemical-probes) (Scheer et al. 2019; Wu et al. 2019)
(Supplementary Table 1). This unique library includes compounds targeting key epigenetic
regulatory proteins with a high degree of potency and selectivity, and a favourable
therapeutic index (Ackloo, Brown, and Müller 2017). The maximum non-toxic concentrations
for each compound, established by a viability assay in these cells, were used in the screen
(Supplementary Figure 1). Of the 54 molecules, only one molecule, selective type I PRMT
inhibitor MS023, was able to promote exon 7 inclusion in *SMN2* pre-mRNA, without affecting
total *SMN2* mRNA levels (Figure 1A and B). MS023 treatment in these fibroblasts also
increased SMN protein levels up to 1.6-fold, as determined by Western blot analysis (Figure
1C). Notably, treatment with PRMT5 inhibitors LLY-283 and GSK591 resulted in reduction of
exon 7 inclusion in *SMN2* pre-mRNA and in SMN protein, respectively (Figure 1A–C),
overall suggesting that protein arginine asymmetric and symmetric dimethylation by different
families of PRMTs exert opposite effects on SMN regulation. Other compounds, including
bromodomain inhibitors (BAY-299, BI-9564, JQ1) and lysine demethylase inhibitors (GSK-
J1, GSK-LSD), also elicited a ≥1.5-fold increase in SMN protein without affecting mRNA
levels, hinting at a direct or indirect effect on SMN protein regulation (Figure 1A-C). PRMTs
are involved in several critical biological functions (Blanc and Richard 2017), and represent a

promising therapeutic target for many human diseases from cancer to neurodegeneration,
with at least eight PRMT inhibitors attaining clinical trial testing in human cancers (Guccione
and Richard 2019; Hwang et al. 2021; Yang and Bedford 2013). Altogether, the direct effect
of PRMT type I inhibition on *SMN2* exon 7 inclusion and the potential for clinical impact of
this class of molecules, prompted us to further investigate MS023 as a therapeutic agent for
SMA.
7
Type I PRMT inhibition promotes exon 7 inclusion in SMN2 *pre-mRNA by decreasing*
hnRNPA1 binding. MS023 is a recently identified potent and selective inhibitor of type I
PRMTs harbouring an ethylendiamino group, a critical moiety for its activity (Figure 2A)
(Eram et al. 2016). Treatment of SMA fibroblasts with MS023 led to a dose-dependent
increase in both *SMN2* exon 7 inclusion and protein levels (Figure 2B–D; Supplementary
Figure 2A). This effect was not observed upon treatment with MS094 (Supplementary Figure
2B and C), an inactive MS023 analogue where the terminal primary amino group in the
ethylenediamino group is replaced with a hydroxyl group (Eram et al. 2016), further
confirming its dependency on PRMT activity. Knock-down of each and any combination of
two of the type I PRMTs (PRMT 1, 3, 4, 6, and 8) resulted in no change in *SMN2* splicing,
suggesting a high degree of functional overlap between these proteins (not shown). In order
to identify the PRMT substrate mediating the effect of MS023 on *SMN2*, we interrogated a
recently published dataset of the arginine methyl proteome in human NB4 cells upon
treatment with MS023 (Fong et al. 2019). Out of 72 responsive targets, we noted that
MS023 determined largely a downregulation of asymmetric dimethylarginine (ADMA) and an
increase in monomethylarginine (MMA) sites in the heterogeneous nuclear ribonucleoprotein
A1 (hnRNPA1) (Fong et al. 2019). hnRNPA1 directly binds *SMN2* pre-mRNA across multiple
sites and is a well-established negative regulator of exon 7 splicing (Bose et al. 2008;
Cartegni et al. 2006; Chen et al. 2008; Hua et al. 2008; Kashima, Rao, and Manley 2007;
Koed Doktor et al. 2011; Singh et al. 2013; Xiao et al. 2012). Given that PRMT-regulated
methylation regulates binding of RNA-binding proteins to RNAs (Blanc and Richard 2017),
we hypothesised that MS023-induced ADMA-to-MMA switch in hnRNPA1 methylation
affects its binding to *SMN2* pre-mRNA, resulting in exon 7 inclusion. In order to test this
hypothesis, we treated SMA fibroblasts with increasing concentrations of MS023 and
confirmed a dose-responsive increase in both symmetric dimethylarginine (SDMA) and MMA
methylation and a concomitant reduction in ADMA levels of hnRNPA1 protein (Figure 2E
and F). Concomitantly, we observed a dose-responsive reduction of the fraction of FL *SMN2*
transcripts bound to hnRNPA1, as determined by a crosslinking and immunoprecipitation
(CLIP) assay (Figure 2G), which validated our working model (Figure 2H). Treatment with
MS023 did not change hnRNPA1 levels nor its subcellular localization, further suggesting
that the treatment-induced change in arginine methylation specifically affects hnRNPA1
binding affinity with its RNA target (Supplementary Figure 2D and E). Taken together, these
findings indicate that MS023 promotes *SMN2* exon 7 inclusion via decreased binding of
hnRNPA1.
42
43 *Oral administration of MS023 improves the phenotype of SMA mice alone and in synergy*
44 *with nusinersen*. We next evaluated the efficacy and tolerability of MS023 treatment *in vivo*
45 in a severe preclinical mouse model of SMA (Hsieh-Li et al. 2000). These mice, which lack
46 the mouse *Smn* gene and only carry a single copy of the human *SMN2* gene, display a
47 phenotype with weight loss and reduced motor activity starting at postnatal day 5 (P5) and

typically reach humane end point by P9. Daily oral administration of MS023 or vehicle (0.5% DMSO in 0.9% saline solution) was performed in SMA mice from P0 until reaching a humane end point (Figure 3A). We tested a range of doses (1, 2, 5, and 40 mg/kg) and found that treatment with both 2 mg/kg and 5 mg/kg resulted in significant increase in survival, with the 2 mg/kg dose achieving the best effect (median: 10 days), compared to vehicle-treated mice (median: 6 days; $P < 0.0001$) (Figure 3B). Mice treated with this regimen also showed an improvement in the disease-associated weight loss (Figure 3C). No further amelioration was observed with 40 mg/kg MS023, hinting that with this dose the therapeutic window has been surpassed (Figure 3B, Supplementary Figure 3A). Notably, we detected an increase in full length (FL) *SMN2* transcript and SMN protein levels in the tissues mostly affected in the disease, spinal cord and skeletal muscle (Eugenio Mercuri et al. 2022), of SMA mice treated with 2 mg/kg MS023 (Figure 3D–F; Supplementary Figure 3B). Since the mechanism of action of the 2'-O-methoxyethyl phosphorothioate-modified drug nusinersen, a currently approved ASO therapy for SMA patients, consists of promoting exon 7 inclusion into *SMN2* pre-mRNA by blocking the recruitment of hnRNP splicing repressors at the ISS-N1 site (Chiriboga et al. 2016; Finkel et al. 2016, 2017; Haché et al. 2016; Eugenio Mercuri et al. 2018), we postulated that a combinatorial treatment with MS023 would synergistically lead to improved therapeutic benefit and allow a cost-effective ASO dosing regimen in SMA. In order to test this hypothesis, SMA mice were treated at P0 with a single subcutaneous administration of 30 mg/kg nusinersen, a minimum effective dose previously shown to extend the SMA mice life span (Hammond et al. 2016), alone or in combination with daily oral administration of 2 mg/kg MS023 from P1 to P6 (Figure 4A). Analysis of tissues collected at P7 showed that the combinatorial treatment was able to further enhance exon 7 inclusion in *SMN2* pre-mRNA (Figure 4B) and increase SMN protein levels (Figure 4C and D; Supplementary Figure 4), both in CNS and peripheral tissues. A further follow-up study designed to assess the effect on survival (Figure 4E) revealed that the combinatorial treatment of nusinersen and MS023 dramatically prolonged the lifespan (median: 97.5 days) compared to nusinersen alone (median: 19.5; $P = 0.02$) (Figure 4F), and improved the body weight of SMA mice (Figure 4G). Overall, these results suggest that oral administration of MS023 synergises with nusinersen to provide a therapeutic benefit in SMA.

Molecular signature of the combinatorial treatment. In order to understand the molecular underpinnings of the added benefit provided by the combinatorial treatment, we performed bulk transcriptomic analysis in spinal cords of symptomatic (P7) SMA mice treated with MS023 only, nusinersen only, and MS023 in combination with nusinersen, compared to wild type and untreated SMA mice, following the treatment paradigm described above (Figure 4A). Out of a total of 5509 significantly dysregulated transcripts in SMA ($P < 0.05$, false discovery rate < 0.05), 2482/2874 (86%) of the upregulated and 2060/2635 (83%) of the downregulated genes were commonly corrected by nusinersen treatment alone and by combinatorial treatment (Figure 5A). Combinatorial treatment was able to exclusively rescue 359 (6.5% out of 5509) additional genes that were not corrected upon treatment with nusinersen alone (Supplementary Table 2). We generated a heatmap of the top significant genes ($P < 0.01$) of this category (Figure 5B) and performed a hallmark analysis, to depict and identify the changes that could explain the beneficial effects of the combinatorial treatment (Figure 5C and D). TNF-α signalling, together with several other immune-related pathways, including interferon response and complement activation, were highly enriched,

overall suggesting that targeting neuroinflammation for therapy is key to achieving a
beneficial effect in SMA. Interestingly, astrocyte dysfunction and chronic microglia activation
have been observed early in SMA and other neurological conditions (Eikelenboom et al.
2002; Heneka, Kummer, and Latz 2014; Sargsyan, Monk, and Shaw 2005; Vukojicic et al.
2019) and their contribution to neuronal dysfunction and death is abundantly supported
(Martin et al. 2017; Mcgivern et al. 2013; Rindt et al. 2015; Zhou, Feng, and Ko 2016). Given
the role of SMN in RNA biogenesis and spliceosomal proteins assembly (Pellizzoni et al.
2001), and the observation of increased mis-splicing events upon SMN depletion (Bäumer et
al. 2009; Custer et al. 2016; Doktor et al. 2017; Huo et al. 2014; Z. Zhang et al. 2008, 2013)
we wondered whether the beneficial effect of the combinatorial treatment was due to
restoration in the splicing profile. We identified 446 mis-splicing events in the spinal cord of
SMA mice, with the vast majority ($n = 353$, 79%) being skipped exons, in accordance with
previous reports (Bäumer et al. 2009; Custer et al. 2016; Doktor et al. 2017; Huo et al. 2014;
Z. Zhang et al. 2008, 2013) (Supplementary Figure 5A and B). Individual treatment with
MS023 or nusinersen were both able to restore the splicing profile almost fully (347/446 and
368/446, respectively), with only 14 unique splicing events corrected exclusively upon the
combinatorial treatment (Figure 5E; Supplementary Table 3), suggesting that mis-splicing
has a low threshold for normalization in SMA and is a poor predictor of treatment response.
19
Discussion
Thousands of SMA patients are treated each year with nusinersen, the first SMA treatment
to receive FDA and EMA approval in 2016 and 2017, respectively. Despite having
enormously changed the life expectancy and quality of life of patients, this approach is not
without limitations: it fails to address the peripheral manifestations of the disease, is costly,
and the intrathecal administrations require patients' hospitalization and well-trained
clinicians, becoming more troublesome with time as the disease progresses (Eugenio
Mercuri et al. 2020). Research efforts are underway to develop both new SMN-dependent
and SMN-independent therapies for use in combination with current therapies, to provide
improved and sustained benefit for all SMA patients.
From a screen of potent and highly selective epigenetic probes, we have identified MS023, a
potent and selective next-generation type I PRMT inhibitor (Eram et al. 2016), able to
elevate SMN protein in SMA models, at levels comparable to recently clinically approved
small molecule risdiplam (Naryshkin et al. 2014) and to synergistically amplify the effects of
nusinersen. Interestingly, type II PRMT-targeting by selective PRMT5 inhibitors GSK591 and
LLY-283 reduced FL *SMN2*, supporting a mechanism of regulation of *SMN2* by this class of
enzymes.
Protein arginine methylation is a prevalent posttranscriptional modification occurring at the
nitrogen of guanidinium group of over 4000 proteins, many of which are RNA-binding
proteins, influencing their stability, function, and interaction with other macromolecules
(Hornbeck et al. 2015). In mammals, arginine methylation of histone and non-histone
proteins is performed by nine PRMT enzymes, divided into three families: type I PRMTs
(PRMT1, PRMT2, PRMT3, PRMT4, PRMT6 and PRMT8) perform monomethylation
followed by an asymmetric dimethylation (ADMA) of arginine, with both methyl groups on a
single guanidino nitrogen atom; type II PRMTs (PRMT5 and PRMT9) catalyse
monomethylation, followed by a symmetric dimethylation (SDMA) where one methyl group is
transferred to each of the nitrogen atoms; and type III PRMT (PRMT7) only performs
monomethylation (MMA) (Guccione and Richard 2019). PRMTs have been shown to

compete over the same substrates, frequently with opposite functional effects (Dhar et al.
2013).

Several studies have highlighted a link between PRMTs and SMN: To ensure fidelity of
loading onto the correct snRNAs, three of the core spliceosomal Sm proteins (SmB, SmD1
and SmD3) undergo symmetric dimethylation by PRMT5, a modification recognised by the
Tudor domain of SMN (Friesen, Paushkin, et al. 2001; Meister et al. 2001). SMN interaction
with senataxin, a DNA-RNA helicase, depends on the dimethylated arginine and is reduced
upon PRMT5 knockdown (Yanling Zhao et al. 2016). Lastly, levels of PRMT4 (CARM1) are
upregulated in SMA mice spinal cord and patients' cells (Sanchez et al. 2015). PRMTs
represent a promising therapeutic target for many human diseases from cancer (Drew et al.
2017; Wang et al. 2016; Wu et al. 2022) to neurodegeneration (Dormann et al. 2012;
Scaramuzzino et al. 2015; Suárez-Calvet et al. 2016), with eight PRMT inhibitors attaining
clinical trial testing in human cancers (Blanc and Richard 2017; Guccione and Richard 2019;
Hwang et al. 2021; Yang and Bedford 2013). Consistently with previous methylome
analyses in cells treated with type I PRMT inhibitors, GSK3368712 (Noto et al. 2020) or
MS023 (Karuppagounder et al. 2016), we propose that MS023 mechanism of action entails
a switch in the arginine methylation profile from ADMA to SDMA/MMA of hnRNPA1, a major
negative regulator of *SMN2* exon 7 inclusion (Bose et al. 2008; Cartegni et al. 2006; Chen et
al. 2008; Hua et al. 2008; Kashima, Rao, and Manley 2007; Koed Doktor et al. 2011; Singh
et al. 2013; Xiao et al. 2012), resulting in reduced binding affinity to *SMN2* pre-mRNA.
Notably, nusinersen also acts by interfering with hnRNPA1 binding to the ISS-N1 sequence,
suggesting a synergistic convergence on this mechanism of regulation. Since splicing is
regulated by several RNA binding proteins, we cannot exclude that methylation of splicing
factors other than hnRNPA1 also contribute, directly or indirectly, to the effect of MS023 on
SMN2 (Ravindra N. Singh et al. 2018).
Hallmark analysis of the 359 genes whose expression levels in spinal cords were only
corrected by the combinatorial treatment, revealed enrichment of immune-related pathways,
such as TNF-α signalling, interferon response and complement activation, overall suggesting
that therapeutic targeting of neuroinflammation is key to achieving optimal and long-lasting
effect in SMA. Interestingly, astrocyte dysfunction and chronic microglia activation have
been observed early in SMA and other neurological conditions (Eikelenboom et al. 2002;
Heneka, Kummer, and Latz 2014; Sargsyan, Monk, and Shaw 2005; Vukojicic et al. 2019)
and play a determinative role in the disease pathogenesis (Martin et al. 2017; Mcgivern et al.
2013; Rindt et al. 2015; Zhou, Feng, and Ko 2016). Transcriptomic data in the preclinical
model indicates that, at the dose employed in this study, MS023 shows a favourable safety
profile, with minimal off-target effects both on gene expression and splicing alterations, in
stark contrast to the pleiotropic effects of other epigenetic modulators such as valproic acid.
Altogether, these promising preclinical results warrant further clinical investigations of MS023
or other selective type I PRMT inhibitors both as a stand-alone and an add-on treatment with
nusinersen in SMA patients.
41

42   **Methods**
43   *Small molecules screening.* The library of small molecules was obtained from the Structural
44   Genomic Consortium (SGC) (Williamoson AR, 2000). Upon reception the molecules were
45   diluted in DMSO to 10 mM, aliquoted and stored at −20°C for the duration of the study. All the
46   compounds, their targets and doses used in the study are listed in Supplementary Table 1.
47

Cell lines and culture. Cells were grown in a humidified incubator at 37°C with 5% CO_2. SMA
type II patient fibroblasts were obtained from Coriell Institute (GM03813). The cells were
maintained in Dulbecco's Modified Eagle Medium GlutaMAX (Gibco) supplemented with 10%
foetal bovine serum (Gibco) and 1× Antibiotic-Antimycotic (Gibco). SMA patient fibroblasts
were plated in triplicates, in 12-well plates at 50,000 cells per well in 500 µL of medium for
RNA or in duplicates in 6-well plates at 100,000 cells per well in 1 mL of medium for protein.
After 6 h, compounds of interest were diluted in medium to 2× final concentration and added
to the cells. RNA was isolated after 48 h incubation, protein after 72 h.
9
Cell viability assay. MTS Cell Proliferation Assay Kit (Abcam) was used to determine the
highest non-toxic concentration of small molecules in SMA patient fibroblasts. Briefly, 3000
cells/wells were plated in triplicates in a 96-well plate in half of the final volume of medium
(100 µL). After 6 h, compounds of interest were diluted to 2× final concentration in 100 µL of
medium and added in a range of concentrations (1 µM to 10 µM). 1 µM of staurosporine
(Sigma) was used as a positive control for cell death. After 48 h, 20 µL of MTS reagent was
added to each well. Four hours later, the absorbance was measured at 490 nm on
CLARIOstar plate reader (BMG Labtech). The concentration was deemed non-toxic if the
absorbance was not significantly different from the control and cells looked viable after visual
inspection. In subsequent experiments, only the highest non-toxic concentration of each
compound was used, unless stated otherwise.
21
RNA/cDNA preparation and RT-qPCR. RNA extraction was performed using Maxwell® RSC
simply RNA Kit (Promega) following the manufacturer's instructions. The concentration was
measured using Nanodrop 1000 spectrophotometer (Thermo Fisher), and cDNA was
generated using an ABI High Capacity cDNA Reverse Transcription Kit (Invitrogen) following
the manufacturer's instructions. A qPCR reaction using Power SYBR Green Master Mix (Life
Technologies) was performed and analysed on an Applied Biosystems StepOnePlus™ real-
time PCR system (Life Technologies). FL *SMN2*, Δ7 *SMN2*, Tot *SMN2*, *PolJ*, *Gapdh* and
GAPDH transcripts were amplified using gene- and species-specific primers (Supplementary
Table 4).
31
Protein extraction and Western blot. Proteins were harvested from ~30 mg of tissue (*in vivo*)
or two 6-well plates (*in vitro*) and homogenised in Ripa buffer with complete mini-proteinase
inhibitors (Roche). Preparation of nuclear and cytoplasmic extracts from human fibroblasts
was performed using a NE-PER Nuclear and Cytoplasmic Extraction Reagents (Thermo
Fisher), as per manufacturer's instructions. Proteins (10–15 µg from cells and 20–30 µg from
tissue) were probed for human SMN protein using anti-SMN, clone SMN-KH monoclonal
IgG1 (Sigma, MABE230), Histone 3 (Cell Signalling, 9715), Vinculin (Sigma, 062M4762),
hnRNPA1 (Santa Cruz, 4B10), MMA (Cell Signalling, 8015), ADMA (Sigma Aldrich, 07-414),
SDMA (Cell Signalling, 13222), β-tubulin (Abcam, 108342) and FAST green total protein
stain and secondary antibody IRDye 800CW goat anti-mouse IgG (LI-COR Biosciences).
Membranes were imaged on a LI- COR Odyssey FC imager and analysed with Image
Studio™ software (LI-COR Biosciences).
44
Crosslinking and immunoprecipitation assay (CLIP). Briefly, for each condition, four 150 cm^2
cell dishes of human patient fibroblasts were treated with 100 nM, 250 nM or 1 µM of MS023
for 48 h. After the incubation, the medium was aspirated and the cells were subjected to 150

mJ/cm^2 254 nm UV light in a Stratalinker UV Crosslinker and pelleted. The cells were then
lysed in NP-40 lysis buffer and hnRNPA1 immunoprecipitation was performed with
Dynabeads Protein G (Thermofisher) and hnRNAP1 antibody (Santa Cruz). Following DNA
and protein digestion, RNA was isolated and cDNA was generated as previously outlined. A
qPCR reaction using Power SYBR Green Master Mix (Life Technologies) was performed
and analysed on an Applied Biosystems StepOnePlus™ real-time PCR system (Life
Technologies). FL *SMN2* and *GAPDH* transcripts were amplified using gene- and species-
specific primers.
9
Protein immunoprecipitation. To detect changes in hnRNPA1 arginine methylation, SMA
patient fibroblasts were either left untreated or treated with increasing concentrations MS023
(100 nM, 250 nM or 1 µM) for 48h. Cell pellets were lysed in NP-40 buffer and hnRNPA1
immunoprecipitation was performed with Dynabeads Protein G (Thermofisher, 10003D) and
hnRNAP1 antibody (Santa Cruz), following the manufacturer's protocol. Subsequently,
Western blots were run according to the protocol described above.
16
Mice. Mice were housed and all the procedures were carried out at the Biomedical Services
Building, University of Oxford and authorised by the UK Home Office in accordance with the
Animals (Scientific Procedures) Act 1986 and by the University of Oxford ethics committee
(PPL no: PDFEDC6F0). All experiments were performed on the SMA mouse strain FVB.Cg-
Smn1^{tm1Hung}Tg(SMN2)2Hung/J – the 'Taiwanese' model (*Smn*$^{-/-}$;*hSMN2*$^{+/-}$), generated and
maintained as previously described (Gogliotti et al. 2010; Hsieh-Li et al. 2000). MS023 was
administered daily orally from P0 or P1 using a Hamilton syringe (Hamilton). Doses of 0, 1,
2, 5 or 40 mg/kg of MS023 were diluted in 0.5% DMSO and 0.9% saline and administered at
a volume of 5 µL/g of body weight. Nusinersen (sequence:
U*sC*sA*sC*sU*sU*sU*sC*sA*sU*sA*sA*sU*sG*sC*sU*sG*sG*s, where 'S' is
phosphorothioate backbone and '*' is a 2'-O-(2-Methoxyethyl)-oligoribonucleotides
chemistry) was diluted in 0.9% saline and given once at 20 µL/g body weight, via
subcutaneous injection at P0, in a dose of 30 mg/kg. Weights were recorded and overall
health was assessed daily. Since the pups were treated daily from P0 and identifying and
marking individual pups between P0 and P7 poses a high risk of misidentifying an animal, in
this study a litter constitutes an experimental unit and all mice in the litter were subjected to
the same treatment. Only litters between 7 and 11 pups were used to correct for average
weight (mice in smaller litters tend to be bigger and live longer and the opposite is true for
litter of 12 and above) and treatment was allocated randomly to a litter before it was born. All
the experimental units treated were included in the analysis. Personnel performing daily
weights and welfare checks for combinatorial therapy were blinded, researcher performing
oral administration, injections and data analysis was not blinded. For survival analysis, the
humane end point was reached upon 15% weight loss from a maximum weight or when the
mouse was not able to right itself for 30 s. Mice were culled by decapitation (if younger than
postnatal day 10) or cervical dislocation (if 10 days old or older). Tissues were harvested at
the indicated postnatal day.
43
RNA sequencing. Transcriptomic analysis was performed by Novogene (UK) company
Limited (https://en.novogene.com/) on P7 spinal cords of mice in the following treatment
groups: untreated SMA mice, SMA mice treated with nusinersen, SMA mice treated with
MS023, SMA mice treated with nusinersen and MS023, untreated controls (4 biological

replicates in each group). RNA quantification and integrity were assessed using the RNA
Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, CA, USA).
mRNA was purified using poly-T oligo-attached magnetic beads. Fragmentation was carried
out using divalent cations under elevated temperature in First Strand Synthesis Reaction
Buffer. First strand cDNA was synthesised using random hexamer primer and M-MuLV
Reverse Transcriptase. Second strand cDNA synthesis was subsequently performed using
DNA Polymerase I and RNase H. Remaining overhangs were converted into blunt ends via
exonuclease/polymerase activities. After adenylation of 3' ends of DNA fragments, adaptor
with hairpin loop structure were ligated to prepare for hybridization. In order to select cDNA
fragments of preferentially 370~420 bp in length, the library fragments were purified with
AMPure XP system (Beckman Coulter, Beverly, USA). PCR products were purified (AMPure
XP system) and library quality was assessed on the Agilent Bioanalyzer 2100 system. The
clustering of the index-coded samples was performed on a cBot Cluster Generation System
using TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the manufacturer's
instructions. After cluster generation, the library preparations were sequenced on an Illumina
Novaseq platform with a coverage of 25 million reads and 150 bp paired-end reads were
generated. Raw data (raw reads) of fastq format were firstly processed through in-house perl
scripts. In this step, clean data (clean reads) were obtained by removing reads containing
adapter, reads 1 containing ploy-N and low-quality reads from raw data. Mus Musculus
(GRCm38/mm10) reference genome was used, index of the reference genome was built
using Hisat2 v2.0.5 and paired-end clean reads were aligned to the reference genome using
Hisat2 v2.0.5. The mapped reads of each sample were assembled by StringTie (v1.3.3b)
(Pertea et al. 2015) in a reference-based approach. Quantification of gene expression level
Feature Counts v1.5.0-p3 was used to count the reads numbers mapped to each gene.
Differential expression analysis was performed using the DESeq2 R package (1.20.0),
Benjamini-Hochberg-adjusted *P*-values reported. Corrected *P*-value of 0.05 and absolute
fold change of 2 were set as the threshold for significantly differential expression. Alternative
splicing analysis rMATS(4.1.0) software was used to analysis the splicing event. We used
the *psiPerEvent* operation of SUPPA to calculate the Ψ values from the transcript
quantifications obtained for all the alternative splicing events generated as described above
with the *generateEvents* module of SUPPA (Alamancos GP, 2014). Data were visualised
with R (R Core Team, 2017).
33
34   *Data and materials availability.* All data needed to evaluate the conclusions in the paper are
35   present in the paper and/or the Supplementary Materials. The RNA-Seq data from wild type
36   and transgenic SMA mice is available for download from Gene Expression Omnibus (GEO)
37   with accession number GSE206400.
38
Statistics. ANOVA tests were used to compare the means between two and three or more
groups, respectively. Statistics of survival times of SMA mice were determined by Kaplan-
Meier estimation, and comparisons were made with the log-rank test. A two-way ANOVA
was conducted to compare the effect of the treatment on weights of the animals using
treatment as a between-subjects factor and time as a within-subjects factor. Power analysis
was performed using G*Power 3.1.9.2 software (Erdfelder et al. 2009). GraphPad Prism
version 8 was used to perform the statistical analyses (GraphPad, La Jolla, CA). A *P*-value
less than 0.05 was set as statistically significant.
47

1   *Study approval*
2   All animal procedures were authorised by the UK Home Office in accordance with the
3   Animals (Scientific Procedures) Act 1986 and by the University of Oxford ethics committee
4   (PPL no: PDFEDC6F0).

Author contributions
AJK, SMH, MJAW, and CR conceived the study. AJK, NA, MH, AB, JS, and WFL performed the experiments. AJK and CR analysed the data and wrote the manuscript. All authors contributed to the discussion and writing of the manuscript.

Acknowledgments
This study was supported by a Career Development Fellowship grant from the Wellcome Trust to CR (205162/Z/16/Z). AJK was supported by the Medical Research Council programme grant (BRT00040). WFL was supported by the Medical Research Council programme grant to M.J.A.W. (MR/N024850/1). The authors would like to thank the Structural Genomics Consortium for providing the library of small molecules.

References

Ackloo, Suzanne, Peter J. Brown, and Susanne Müller. 2017. "Chemical Probes Targeting Epigenetic Proteins: Applications beyond Oncology." *Epigenetics* 12(5): 378–400.

Akten, Bikem et al. 2011. "Interaction of Survival of Motor Neuron (SMN) and HuD Proteins with MRNA Cpg15 Rescues Motor Neuron Axonal Deficits." *Proceedings of the National Academy of Sciences of the United States of America* 108(25): 10337–42.

Allis, C. David et al. 2007. "New Nomenclature for Chromatin-Modifying Enzymes." *Cell* 131(4): 633–36.

Araujo, Alexandra prufer de Queiroz Campos, Mario Araujo, and Kathryn J. Swoboda. 2009. "Vascular Perfusion Abnormalities in Infants with Spinal Muscular Atrophy." *The Journal of Pediatrics* 155(2): 292–94.

Arnold, Anne Sophie et al. 2004. "Reduced Expression of Nicotinic AChRs in Myotubes from Spinal Muscular Atrophy I Patients." *Laboratory investigation; a journal of technical methods and pathology* 84(10): 1271–78.

Arrowsmith, Cheryl H. et al. 2012. "Epigenetic Protein Families: A New Frontier for Drug Discovery." *Nature reviews. Drug discovery* 11(5): 384–400.

Bäumer, Dirk et al. 2009. "Alternative Splicing Events Are a Late Feature of Pathology in a Mouse Model of Spinal Muscular Atrophy." *PLoS Genet* 5(12): e1000773.

Bevan, Adam K. et al. 2010. "Early Heart Failure in the SMNDelta7 Model of Spinal Muscular Atrophy and Correction by Postnatal ScAAV9-SMN Delivery." *Human molecular genetics* 19(20): 3895–3905.

Blanc, Roméo S., and Stéphane Richard. 2017. "Arginine Methylation: The Coming of Age." 65(1): 8–24.

Bose, Jayarama Krishnan et al. 2008. "TDP-43 Overexpression Enhances Exon 7 Inclusion during the Survival of Motor Neuron Pre-MRNA Splicing." *The Journal of biological chemistry* 283(43): 28852–59.

Bowerman, Melissa et al. 2012. "Glucose Metabolism and Pancreatic Defects in Spinal Muscular Atrophy." *Annals of Neurology* 72(2): 256–68.

Cao, Yan Yan et al. 2016. "Association between SMN2 Methylation and Disease Severity in Chinese Children with Spinal Muscular Atrophy." *Journal of Zhejiang University. Science. B* 17(1): 76–82.

Cartegni, Luca et al. 2006. "Determinants of Exon 7 Splicing in the Spinal Muscular Atrophy Genes, SMN1 and SMN2." *American journal of human genetics* 78(1): 63–77.

Charroux, Bernard et al. 1999. "Gemin3: A Novel DEAD Box Protein That Interacts with SMN, the Spinal Muscular Atrophy Gene Product, and Is a Component of Gems." *Journal of Cell Biology* 147(6): 1181–93.

Chen, Hung-Hsi et al. 2008. "The RNA Binding Protein HnRNP Q Modulates the Utilization of Exon 7 in the Survival Motor Neuron 2 (SMN2) Gene." *Molecular and Cellular Biology* 28(22): 6929–38.

Chiriboga, Claudia A. et al. 2016. "Results from a Phase 1 Study of Nusinersen (ISIS-SMN(Rx)) in Children with Spinal Muscular Atrophy." *Neurology* 86(10): 890–97.

Crawford, Thomas O. et al. 1999. "Abnormal Fatty Acid Metabolism in Childhood Spinal Muscular Atrophy." *Annals of Neurology* 45(3): 337–43.

Crawford, Thomas O., and Carlos A. Pardo. 1996. "The Neurobiology of Childhood Spinal Muscular Atrophy." *Neurobiology of Disease* 3(2): 97–110.

Custer, Sara K. et al. 2016. "Altered MRNA Splicing in SMN-Depleted Motor Neuron-like Cells." *PloS one* 11(10): e0163954.

Deguise, Marc Olivier et al. 2019. "Abnormal Fatty Acid Metabolism Is a Core Component of Spinal Muscular Atrophy." *Annals of Clinical and Translational Neurology* 6(8): 1519–32.

Dhar, Surbhi et al. 2013. "Loss of the Major Type I Arginine Methyltransferase PRMT1 Causes Substrate Scavenging by Other PRMTs." *Scientific reports* 3: 1311.

Doktor, Thomas Koed et al. 2017. "RNA-Sequencing of a Mouse-Model of Spinal Muscular Atrophy Reveals Tissue-Wide Changes in Splicing of U12-Dependent Introns." *Nucleic*

Acids Research 45(1): 395–416.

Dormann, Dorothee et al. 2012. "Arginine Methylation next to the PY-NLS Modulates Transportin Binding and Nuclear Import of FUS." *EMBO Journal* 31(22): 4258–75.

Drew, Allison E. et al. 2017. "Identification of a CARM1 Inhibitor with Potent in Vitro and in Vivo Activity in Preclinical Models of Multiple Myeloma." *Scientific reports* 7(1): 17993.

Eikelenboom, P. et al. 2002. "Neuroinflammation in Alzheimer's Disease and Prion Disease." *Glia* 40(2): 232–39.

Eram, Mohammad S. et al. 2016. "A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein Arginine Methyltransferases." *ACS Chemical Biology* 11(3): 772–81.

Erdfelder, Edgar, Franz FAul, Axel Buchner, and Albert Georg Lang. 2009. "Statistical Power Analyses Using G*Power 3.1: Tests for Correlation and Regression Analyses." *Behavior research methods* 41(4): 1149–60.

Fallini, Claudia et al. 2011. "The Survival of Motor Neuron (SMN) Protein Interacts with the MRNA-Binding Protein HuD and Regulates Localization of Poly(A) MRNA in Primary Motor Neuron Axons." *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31(10): 3914–25.

———. 2014. "Dynamics of Survival of Motor Neuron (SMN) Protein Interaction with the MRNA-Binding Protein IMP1 Facilitates Its Trafficking into Motor Neuron Axons." *Developmental Neurobiology* 74(3): 319–32.

———. 2016. "Deficiency of the Survival of Motor Neuron Protein Impairs MRNA Localization and Local Translation in the Growth Cone of Motor Neurons." *The Journal of neuroscience : the official journal of the Society for Neuroscience* 36(13): 3811–20.

Feldkötter, Markus et al. 2002. "Quantitative Analyses of SMN1 and SMN2 Based on Real-Time LightCycler PCR: Fast and Highly Reliable Carrier Testing and Prediction of Severity of Spinal Muscular Atrophy." *American Journal of Human Genetics* 70(2): 358–68.

Finkel, Richard S. et al. 2016. "Treatment of Infantile-Onset Spinal Muscular Atrophy with Nusinersen: A Phase 2, Open-Label, Dose-Escalation Study." *Lancet (London, England)* 388(10063): 3017–26.

———. 2017. "Nusinersen versus Sham Control in Infantile-Onset Spinal Muscular Atrophy." *New England Journal of Medicine* 377(18): 1723–32.

Fong, Jia Yi et al. 2019. "Therapeutic Targeting of RNA Splicing Catalysis through Inhibition of Protein Arginine Methylation." *Cancer Cell* 36(2): 194-209.e9.

Friesen, Westley J., Sergey Paushkin, et al. 2001. "The Methylosome, a 20S Complex Containing JBP1 and PICln, Produces Dimethylarginine-Modified Sm Proteins." *Molecular and Cellular Biology* 21(24): 8289–8300.

Friesen, Westley J, Severine Massenet, et al. 2001. "SMN, the Product of the Spinal Muscular Atrophy Gene, Binds Preferentially to Dimethylarginine-Containing Protein Targets Sufficient for SMN Binding." *Molecular cell* 7(5): 1111–17.

Gogliotti, Rocky G. et al. 2012. "Motor Neuron Rescue in Spinal Muscular Atrophy Mice Demonstrates That Sensory-Motor Defects Are a Consequence, Not a Cause, of Motor Neuron Dysfunction." *The Journal of neuroscience : the official journal of the Society for Neuroscience* 32(11): 3818–29.

Gogliotti, Rocky G, Suzan M Hammond, Cathleen Lutz, and Christine J Didonato. 2010. "Molecular and Phenotypic Reassessment of an Infrequently Used Mouse Model for Spinal Muscular Atrophy." *Biochemical and Biophysical Research Communications* 391(1): 517–22.

Guccione, Ernesto, and Stéphane Richard. 2019. "The Regulation, Functions and Clinical Relevance of Arginine Methylation." *Nature reviews. Molecular cell biology* 20(10): 642–57.

Haché, Manon et al. 2016. "Intrathecal Injections in Children with Spinal Muscular Atrophy: Nusinersen Clinical Trial Experience." *Journal of child neurology* 31(7): 899–906.

Hammond, Suzan M et al. 2016. "Systemic Peptide-Mediated Oligonucleotide Therapy

Improves Long-Term Survival in Spinal Muscular Atrophy." *Proceedings of the National Academy of Sciences* 113(39): 10962–67.

Harding, Brian N. et al. 2015. "Spectrum of Neuropathophysiology in Spinal Muscular Atrophy Type I." *Journal of neuropathology and experimental neurology* 74(1): 15–24.

Hauke, Jan et al. 2009. "Survival Motor Neuron Gene 2 Silencing by DNA Methylation Correlates with Spinal Muscular Atrophy Disease Severity and Can Be Bypassed by Histone Deacetylase Inhibition." *Human Molecular Genetics* 18(2): 304–17.

Heier, Christopher R., Rosalba Satta, Cathleen Lutz, and Christine J. Didonato. 2010. "Arrhythmia and Cardiac Defects Are a Feature of Spinal Muscular Atrophy Model Mice." *Human molecular genetics* 19(20): 3906–18.

Heneka, Michael T., Markus P. Kummer, and Eicke Latz. 2014. "Innate Immune Activation in Neurodegenerative Disease." *Nature reviews. Immunology* 14(7): 463–77.

Hornbeck, Peter V. et al. 2015. "PhosphoSitePlus, 2014: Mutations, PTMs and Recalibrations." *Nucleic acids research* 43(Database issue): D512–20.

Hsieh-Li, Hsiu Mei et al. 2000. "A Mouse Model for Spinal Muscular Atrophy." *Nature Genetics* 24(1): 66–70.

Hua, Yimin et al. 2008. "Antisense Masking of an HnRNP A1/A2 Intronic Splicing Silencer Corrects SMN2 Splicing in Transgenic Mice." *American journal of human genetics* 82(4): 834–48.

Hubers, Lisa et al. 2011. "HuD Interacts with Survival Motor Neuron Protein and Can Rescue Spinal Muscular Atrophy-like Neuronal Defects." *Human Molecular Genetics* 20(3): 553–79.

Huo, Qing et al. 2014. "Splicing Changes in SMA Mouse Motoneurons and SMN-Depleted Neuroblastoma Cells: Evidence for Involvement of Splicing Regulatory Proteins." *RNA Biology* 11(11): 1430–46.

Hwang, Jee Won et al. 2021. "Protein Arginine Methyltransferases: Promising Targets for Cancer Therapy." *Experimental & molecular medicine* 53(5): 788–808.

Jablonka, Sibylle et al. 2006. "Distinct and Overlapping Alterations in Motor and Sensory Neurons in a Mouse Model of Spinal Muscular Atrophy." *Human molecular genetics* 15(3): 511–18.

Karuppagounder, Saravanan S. et al. 2016. "Therapeutic Targeting of Oxygen-Sensing Prolyl Hydroxylases Abrogates ATF4-Dependent Neuronal Death and Improves Outcomes after Brain Hemorrhage in Several Rodent Models." *Science Translational Medicine* 8(328).

Kashima, Tsuyoshi, and James L. Manley. 2003. "A Negative Element in SMN2 Exon 7 Inhibits Splicing in Spinal Muscular Atrophy." *Nature Genetics* 34(4): 460–63.

Kashima, Tsuyoshi, Nishta Rao, Charles J. David, and James I. Manley. 2007. "HnRNP A1 Functions with Specificity in Repression of SMN2 Exon 7 Splicing." *Human Molecular Genetics* 16(24): 3149–59.

Kashima, Tsuyoshi, Nishta Rao, and James L. Manley. 2007. "An Intronic Element Contributes to Splicing Repression in Spinal Muscular Atrophy." *Proceedings of the National Academy of Sciences of the United States of America* 104(9): 3426–31.

Kim, Jeong Ki et al. 2020. "Muscle-Specific SMN Reduction Reveals Motor Neuron-Independent Disease in Spinal Muscular Atrophy Models." *The Journal of clinical investigation* 130(3): 1271–87.

Kissel, John T. et al. 2014. "SMA Valiant Trial: A Prospective, Double-Blind, Placebo-Controlled Trial of Valproic Acid in Ambulatory Adults with Spinal Muscular Atrophy." *Muscle & nerve* 49(2): 187–92.

Koed Doktor, Thomas et al. 2011. "SMN2 Exon 7 Splicing Is Inhibited by Binding of HnRNP A1 to a Common ESS Motif That Spans the 3' Splice Site." *Human Mutation* 32(2): 220–30.

Krosschell, Kristin J. et al. 2018. "Clinical Trial of L-Carnitine and Valproic Acid in Spinal Muscular Atrophy Type I." *Muscle & nerve* 57(2): 193–99.

Lefebvre, Suzie et al. 1995. "Identification and Characterization of a Spinal Muscular Atrophy-Determining Gene." *Cell* 80: 155–65.

Lipnick, Scott L. et al. 2019. "Systemic Nature of Spinal Muscular Atrophy Revealed by Studying Insurance Claims." *PLOS ONE* 14(3): e0213680.

Lorson, C, E Ahnen, Elliot J. Androphy, and Brunhilde Wirth. 1999. "A Single Nucleotide in the SMN Gene Regulates Splicing and Is Responsible for Spinal Muscular Atrophy." *Proceedings of the National Academy of Sciences* 96(May): 6307–11.

Makarov, Evgeny M., Nicholas Owen, Andrew Bottrill, and Olga V. Makarova. 2012. "Functional Mammalian Spliceosomal Complex E Contains SMN Complex Proteins in Addition to U1 and U2 SnRNPs." *Nucleic Acids Research* 40(6): 2639–52.

Marasco, Luciano E. et al. 2022. "Counteracting Chromatin Effects of a Splicing-Correcting Antisense Oligonucleotide Improves Its Therapeutic Efficacy in Spinal Muscular Atrophy." *Cell* 185(12): 2057-2070.e15.

Martin, Jasmin E. et al. 2017. "Decreased Motor Neuron Support by SMA Astrocytes Due to Diminished MCP1 Secretion." *The Journal of neuroscience : the official journal of the Society for Neuroscience* 37(21): 5309–18.

Martínez-Hernández, Rebeca et al. 2009. "The Developmental Pattern of Myotubes in Spinal Muscular Atrophy Indicates Prenatal Delay of Muscle Maturation." *Journal of neuropathology and experimental neurology* 68(5): 474–81.

Martinez, Tara L. et al. 2012. "Survival Motor Neuron Protein in Motor Neurons Determines Synaptic Integrity in Spinal Muscular Atrophy." *Journal of Neuroscience* 32(25): 8703–15.

McAndrew, P. E. et al. 1997. "Identification of Proximal Spinal Muscular Atrophy Carriers and Patients by Analysis of SMNT and SMNC Gene Copy Number." *American journal of human genetics* 60(6): 1411–22.

Mcgivern, Jered V. et al. 2013. "Spinal Muscular Atrophy Astrocytes Exhibit Abnormal Calcium Regulation and Reduced Growth Factor Production." *Glia* 61(9): 1418–28.

Meister, Gunter et al. 2001. "Methylation of Sm Proteins by a Complex Containing PRMT5 and the Putative U SnRNP Assembly Factor PICln." *Current Biology* 11(24): 1990–94.

Mentis, George Z. et al. 2011. "Early Functional Impairment of Sensory-Motor Connectivity in a Mouse Model of Spinal Muscular Atrophy." *Neuron* 69(3): 453–67.

Mercuri, E. et al. 2007. "Randomized, Double-Blind, Placebo-Controlled Trial of Phenylbutyrate in Spinal Muscular Atrophy." *Neurology* 68(1): 51–55.

Mercuri, Eugenio et al. 2018. "Nusinersen versus Sham Control in Later-Onset Spinal Muscular Atrophy." 378(7): 625–35.

———. 2020. "Spinal Muscular Atrophy — Insights and Challenges in the Treatment Era." *Nature Reviews Neurology* 16(12): 706–15.

———. 2022. "Spinal Muscular Atrophy." *Nature Reviews Disease Primers* 8(1): 1–16.

Mohseni, Jafar, Z. A M H Zabidi-Hussin, and Teguh Haryo Sasongko. 2013. "Histone Deacetylase Inhibitors as Potential Treatment for Spinal Muscular Atrophy." *Genetics and Molecular Biology* 36(3): 299–307.

Monani, Umrao R. et al. 1999. "A Single Nucleotide Difference That Alters Splicing Patterns Distinguishes the SMA Gene SMN1 from the Copy Gene SMN2." *Human molecular genetics* 8(7): 1177–83.

Munsat, Theodore L., and Kay E. Davies. 1992. "International SMA Consortium Meeting (26-28 June 1992, Bonn, Germany)." *Neuromuscular Disorders* 2(5–6): 423–28.

Murray, Lyndsay M et al. 2015. "Transcriptional Profiling of Differentially Vulnerable Motor Neurons at Pre-Symptomatic Stage in the Smn 2b/-Mouse Model of Spinal Muscular Atrophy." *Acta Neuropathologica Communications* Sep 15(3): 55.

Naryshkin, Nikolai A. et al. 2014. "SMN2 Splicing Modifiers Improve Motor Function and Longevity in Mice with Spinal Muscular Atrophy." *Science* 345(6197): 688–93.

Noto, Paul B. et al. 2020. "Identification of HnRNP-A1 as a Pharmacodynamic Biomarker of Type I PRMT Inhibition in Blood and Tumor Tissues." *Scientific Reports* 10(1): 1–19.

Pearn, John. 1978. "Incidence, Prevalence and Gene Frequency Studies of Chronic Childhood Spinal Muscular Atrophy." *Journal of Medical Genetics* Dec;15(6): 409–13.

Pellizzoni, Livio, Jennifer Baccon, Bernard Charroux, and Gideon Dreyfuss. 2001. "The Survival of Motor Neurons (SMN) Protein Interacts with the SnoRNP Proteins Fibrillarin and GAR1." *Current Biology* 11(14): 1079–88.

Pellizzoni, Livio, Naoyuki Kataoka, Bernard Charroux, and Gideon Dreyfuss. 1998. "A Novel Function for SMN, the Spinal Muscular Atrophy Disease Gene Product, in Pre-MRNA Splicing." *Cell* 95(5): 615–24.

Pellizzoni, Livio, Jeongsik Yong, and Gideon Dreyfuss. 2002. "Essential Role for the SMN Complex in the Specificity of SnRNP Assembly." *Science* 298(5599): 1775–79.

Pertea, Mihaela et al. 2015. "StringTie Enables Improved Reconstruction of a Transcriptome from RNA-Seq Reads." *Nature biotechnology* 33(3): 290–95.

Piazzon, Nathalie et al. 2008. "In Vitro and in Cellulo Evidences for Association of the Survival of Motor Neuron Complex with the Fragile X Mental Retardation Protein." *Journal of Biological Chemistry* 283(9): 5598–5610.

Portela, Anna, and Manel Esteller. 2010. "Epigenetic Modifications and Human Disease." *Nature biotechnology* 28(10): 1057–68.

Rage, Florence et al. 2013. "Genome-Wide Identification of MRNAs Associated with the Protein SMN Whose Depletion Decreases Their Axonal Localization." *RNA* 19(12): 1755–66.

Ravindra N. Singh, Singh and Natalia N., Ravindra N. Singh, and Natalia N. Singh. 2018. 20 Advances in Neurobiology *Mechanism of Splicing Regulation of Spinal Muscular Atrophy Genes*. Springer New York LLC.

Rindt, Hansjörg et al. 2015. "Astrocytes Influence the Severity of Spinal Muscular Atrophy." *Human molecular genetics* 24(14): 4094–4102.

Rossoll, Wilfried et al. 2003. "Smn, the Spinal Muscular Atrophy-Determining Gene Product, Modulates Axon Growth and Localization of β-Actin MRNA in Growth Cones of Motoneurons." *Journal of Cell Biology* 163(4): 801–12.

Rudnik-Schöneborn, Sabine et al. 2003. "Classical Infantile Spinal Muscular Atrophy with SMN Deficiency Causes Sensory Neuronopathy." *Neurology* 60(6): 983–87.

———. 2010. "Digital Necroses and Vascular Thrombosis in Severe Spinal Muscular Atrophy." *Muscle & nerve* 42(1): 144–47.

Sanchez, Gabriel et al. 2015. "A Novel Role for CARM1 in Promoting Nonsense-Mediated MRNA Decay: Potential Implications for Spinal Muscular Atrophy." *Nucleic Acids Research* 44(6): 2661–76.

Sargsyan, Siranush A., Peter N. Monk, and Pamela J. Shaw. 2005. "Microglia as Potential Contributors to Motor Neuron Injury in Amyotrophic Lateral Sclerosis." *Glia* 51(4): 241–53.

Scaramuzzino, Chiara et al. 2015. "Protein Arginine Methyltransferase 6 Enhances Polyglutamine-Expanded Androgen Receptor Function and Toxicity in Spinal and Bulbar Muscular Atrophy." *Neuron* 85(1): 88–100.

Scheer, Sebastian et al. 2019. "A Chemical Biology Toolbox to Study Protein Methyltransferases and Epigenetic Signaling." *Nature communications* 10(1): 19.

Shafey, Dina, Justin G. Boyer, Kunal Bhanot, and Rashmi Kothary. 2010. "Identification of Novel Interacting Protein Partners of SMN Using Tandem Affinity Purification." *Journal of Proteome Research* 9(4): 1659–69.

Singh, Natalia N. et al. 2013. "An Intronic Structure Enabled by a Long-Distance Interaction Serves as a Novel Target for Splicing Correction in Spinal Muscular Atrophy." *Nucleic acids research* 41(17): 8144–65.

Somers, Eilidh et al. 2016. "Vascular Defects and Spinal Cord Hypoxia in Spinal Muscular Atrophy." *Annals of Neurology* 79(2): 217–30.

Strasswimmer, John et al. 1999. "Identification of Survival Motor Neuron as a Transcriptional Activator-Binding Protein." *Human Molecular Genetics* 8(7): 1219–26.

Suárez-Calvet, Marc et al. 2016. "Monomethylated and Unmethylated FUS Exhibit Increased Binding to Transportin and Distinguish FTLD-FUS from ALS-FUS." *Acta neuropathologica* 131(4): 587–604.

Suraweera, Amila et al. 2009. "Functional Role for Senataxin, Defective in Ataxia Oculomotor Apraxia Type 2, in Transcriptional Regulation." *Human Molecular Genetics* 18(18): 3384–96.

Tadesse, Helina, Julie Deschênes-Furry, Sophie Boisvenue, and Jocelyn Côté. 2008. "KH-Type Splicing Regulatory Protein Interacts with Survival Motor Neuron Protein and Is Misregulated in Spinal Muscular Atrophy." *Human Molecular Genetics* 17(4): 506–24.

Tisdale, Sarah et al. 2013. "SMN Is Essential for the Biogenesis of U7 Small Nuclear Ribonucleoprotein and 3'-End Formation of Histone MRNAs." *Cell Reports* 5(5): 1187–95.

Vukojicic, Aleksandra et al. 2019. "The Classical Complement Pathway Mediates Microglia-Dependent Remodeling of Spinal Motor Circuits during Development and in SMA." *Cell reports* 29(10): 3087-3100.e7.

Wang, Lu et al. 2016. "CARM1 Methylates Chromatin Remodeling Factor BAF155 to Enhance Tumor Progression and Metastasis." *Cancer cell* 30(1): 179–80.

Wu, Qin et al. 2019. "A Chemical Toolbox for the Study of Bromodomains and Epigenetic Signaling." *Nature communications* 10(1): 1915.

———. 2022. "PRMT Inhibition Induces a Viral Mimicry Response in Triple-Negative Breast Cancer." *Nature Chemical Biology* 18(8): 821–830.

Xiao, Rui et al. 2012. "Nuclear Matrix Factor HnRNP U/SAF-A Exerts a Global Control of Alternative Splicing by Regulating U2 SnRNP Maturation." *Molecular cell* 45(5): 656–68.

Yang, Yanzhong, and Mark T. Bedford. 2013. "Protein Arginine Methyltransferases and Cancer." *Nature Reviews Cancer* Jan;13(1): 37–50.

Yanling Zhao, Dorothy et al. 2016. "SMN and Symmetric Arginine Dimethylation of RNA Polymerase II C-Terminal Domain Control Termination." *Nature* 529(7584): 48–53.

Zhang, Rundong et al. 2011. "Structure of a Key Intermediate of the SMN Complex Reveals Gemin2's Crucial Function in SnRNP Assembly." *Cell* 146(3): 384–95.

Zhang, Zhenxi et al. 2008. "SMN Deficiency Causes Tissue-Specific Perturbations in the Repertoire of SnRNAs and Widespread Defects in Splicing." *Cell* 133(4): 585.

———. 2013. "Dysregulation of Synaptogenesis Genes Antecedes Motor Neuron Pathology in Spinal Muscular Atrophy." *Proceedings of the National Academy of Sciences of the United States of America* 110(48): 19348–53.

Zheleznyakova, Galina Y. et al. 2013. "Genome-Wide Analysis Shows Association of Epigenetic Changes in Regulators of Rab and Rho GTPases with Spinal Muscular Atrophy Severity." *European Journal of Human Genetics* Sep;21(9): 988–93.

Zhou, Chunyi, Zhihua Feng, and Chien Ping Ko. 2016. "Defects in Motoneuron-Astrocyte Interactions in Spinal Muscular Atrophy." *The Journal of neuroscience : the official journal of the Society for Neuroscience* 36(8): 2543–53.

Figure 1

A

B

C

1
2
Figure 1. Screening of epigenetic small molecules in SMA fibroblasts. SMA type II
patient-derived fibroblasts were treated with the indicated small molecule at the appropriate
maximum tolerated dose (range: 1–10 µM) (n = 3–4). Cells were harvested for RNA (**A**–**B**)
and protein (**C**) quantification 48 and 72 hours post treatment, respectively. **A**, Full length
(FL) *SMN2* transcript levels relative to Δ7 *SMN2* are expressed as fold change compared to
untreated SMA fibroblasts, normalised to one (dashed line). FL *SMN2*/Δ7 *SMN2* ratios was
significantly increased by MS023 and decreased by LLY-283 treatment. **B**, Tot *SMN2*
transcript levels relative to *GAPDH* are expressed as fold change compared to untreated
SMA fibroblasts, normalised to one (dashed line). **A**–**B**, Each dot represents a biological
replicate (n = 3–4). Grey bar indicates values within the 0.5-1.5 range. **C**, SMN protein levels
relative to total protein are expressed as fold change compared to untreated SMA
fibroblasts, normalised to one (dashed line). Each dot represents a biological replicate (n =
2–4). Values ≥1.5 and ≤0.5 are depicted in red and blue, respectively. **A-C**, Data are

1 represented as mean ± s.e.m. and compared with a one-way ANOVA test with multiple
2 comparisons (**$P \leq 0.01$; ****$P \leq 0.0001$).

Figure 2

1
2
3 **Figure 2. MS023 promotes exon 7 inclusion into *SMN2* transcript via hnRNPA1**

4 **binding inhibition. A**, Chemical structure depiction of MS023 (PubChem CID: 92136227).

5 **B-D** SMA type II patient-derived fibroblasts were treated with the indicated concentration of

6 MS023 (range: 100 nm–10 μM), (n = 3–4). Cells were harvested for RNA (**B**) and protein (**C,**

D) quantification 48 and 72 hours post treatment, respectively, **B.** Full length (FL) *SMN2*
transcript levels relative to Δ7 *SMN2* are expressed as fold change compared to untreated
SMA fibroblasts, normalised to one. Each dot represents a biological replicate (*n* = 3–4). **C**,
Western blot showing SMN protein levels upon treatment with increasing MS023
concentrations (top). A representative section of total protein stain, used for protein
normalization, is shown (bottom). Size in kilodalton is indicated on the right. **D**, Quantification
of SMN protein levels relative to total protein is shown. Each dot represents a biological
replicate (*n* = 3). **E,** Arginine monomethylation (MMA) and symmetric dimethylation (SDMA)
of hnRNPA1 are increased and asymmetric dimethylation (ADMA) of hnRNPA1 is
decreased upon MS023 treatment, as shown by hnRNPA1 immunoprecipitation followed by
a Western blot. SMA type II patient-derived fibroblasts were treated with the indicated
concentration of MS023 (range: 100 nm–1 µM). hnRNPA1 was immunoprecipitated and
Western blots were performed using and anti-MMA, anti-SDMA, anti-ADMA, anti-hnRNPA1
and anti-tubulin antibodies. **F**, Quantification of arginine methylation changes normalized to
total hnRNPA1 is shown. **G**, Ratio of FL *SMN2* transcripts bound to hnRNPA1 protein and
relative to *GAPDH* was assayed using crosslinking immunoprecipitation (CLIP) in SMA
patient fibroblasts treated with increasing concentration of MS023. IgG antibody and MS094
were used controls, MS094 is a negative control for MS023 lacking PRMT inhibition
property. Each dot represents a biological replicate (*n* = 2-4). **B, D, G** Data are represented
as mean ± s.e.m. and compared with a one-way ANOVA test with multiple comparisons (*$*P$
≤ 0.05; ***P$ ≤ 0.01; ****P$ ≤ 0.001). **H**, Representation of the mechanism of exon 7 inclusion
into the *SMN2* transcript by MS023. Blue circles, purple squares and red triangles represent
asymmetric dimethylarginine (ADMA), symmetric dimethylarginine (SDMA)
monomethylarginine (MMA) respectively.
25

Figure 3

1
2
Figure 3. Oral administration of MS023 improves the phenotype of SMA mice. A,
Diagram of the study design; mice were treated daily with oral administrations of MS023 or
vehicle (0.5% DMSO in saline) from postnatal day 0 (P0) using a Hamilton syringe. **B**,
Kaplan-Meier survival estimation of mice treated with vehicle (n = 14), 1 mg/kg MS023 (n =
4), 2 mg/kg MS023 (n = 26), 5 mg/kg MS023 (n = 4), or 40 mg/kg MS023 (n = 4) SMA mice.
C, Body weights of MS023- and vehicle-treated SMA mice from postnatal day 0 are shown.
D, FL *SMN2* transcript levels relative to Δ7 *SMN2* in spinal cord and skeletal muscle of
treated mice compared to vehicle-treated mice, normalised to one. Each dot represents a
biological replicate (n = 3–9). **E**, Western blot showing SMN protein levels upon MS023
treatment in spinal cords and skeletal muscles of treated mice (top). A representative section

of total protein stain, used for protein normalization, is also shown (bottom). Size in

kilodalton is indicated on the right. **F**, Quantification of SMN protein levels in the spinal cord

and skeletal muscle relative to total protein is shown. Each dot represents a biological

replicate (n = 3–8). **B–D**, and **F**, Data are represented as mean ± s.e.m. and compared with

Mantel-Cox test (*P ≤ 0.05; **P ≤ 0.01) (**B**) or a one-way ANOVA test with multiple

comparisons (**P ≤ 0.01; ***P ≤ 0.001, ****P ≤ 0.0001) (**C, D, F**)**.**

Figure 4

1
2
Figure 4. Combinatorial treatment with MS023 and ASO exerts synergistic effects in
SMA mice. A, Diagrams of the study design. Study plan #2: At P0 SMA mice were injected
subcutaneously with *SMN2*-targeting ASO (30 mg/kg) or saline; from P1 mice were treated
daily with oral administrations of MS023 (2 m/kg) or vehicle (0.5% DMSO in saline) using a
Hamilton syringe until P6. At P7, mice were sacrificed and tissues collected for analysis. **B**,
FL *SMN2* transcript levels relative to Δ7 *SMN2* in spinal cord, brain, skeletal muscle, kidney
and liver of treated SMA mice compared to vehicle-treated SMA mice, normalised to one.
Each dot represents a biological replicate (*n* = 10–12). **C**, Western blot showing SMN protein
levels upon the indicated treatments in spinal cords of SMA mice (top). A representative
section of total protein stain, used for protein normalization, is also shown (bottom). Size in
kilodalton is indicated on the right. **D**, Quantification of SMN protein levels relative to total
protein is shown. Each dot represents a biological replicate (*n* = 7–13). **E**, Study design #3:

At P0 mice were injected subcutaneously with *SMN2*-targeting ASO (30 mg/kg) or saline.
From P1 mice were treated daily with oral administrations of MS023 (2 m/kg) or vehicle
(0.5% DMSO in saline) using a Hamilton syringe until P12. Mice were weighed daily until
they reached their humane end point. **F**, Kaplan-Meier survival estimation of SMA mice
treated with *SMN2*-targeting ASO ($n = 15$), MS023 ($n = 14$), *SMN2*-targeting ASO and
MS023 ($n = 10$), and vehicle ($n = 23$). **G**, Body weights of SMA mice from postnatal day 0
are shown. **B, D, F,** and **G**, Data are represented as mean ± s.e.m. and compared with a
one-way ANOVA test with multiple comparisons (*$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$, ****$P \leq$
0.0001) or (**F**) Mantel-Cox test (*$P \leq 0.05$; ****$P \leq 0.0001$).

Figure 5

A

Genes upregulated in SMA

192 125
75
2482

Total=2874

Genes downregulated in SMA

278
167
130
2060

Total=2635

- Corrected both by ASO and ASO + MS023
- Only corrected by ASO
- Only corrected by ASO + MS023
- Non corrected

B

Heatmap

SMA   ASO   ASO + MS023   Wild type

-0.5   0.0   0.5

C

Pathways upregulated in SMA and only corrected by ASO + MS023

TNFα signalling
Coagulation
Complement
Inflammatory response
Oxydative phosphorylation
Epithelial mesench. transition
Xenobiotic metabolism
DNA repair
Adipogenesis
P53 pathway
MTORC1 signalling
Glycolysis

0.0   0.5   1.0   1.5   2.0   2.5
NES

D

Pathways downregulated in SMA and only corrected by ASO + MS023

G2M checkpoint
Mitotic spindle
Interferon response
Heme metabolism
Apoptosis
Estrogen response
KRAS signalling UP
Apical junction
KRAS signalling DOWN

-2.0   -1.5   -1.0   -0.5   0.0
NES

E

Analysis of splicing events

Corrected events: 347/446

Wild type [Ψ]

0.00 0.25 0.50 0.75 1.00
MS023 [Ψ]

Corrected events: 368/446

Wild type [Ψ]

0.00 0.25 0.50 0.75 1.00
nusinersen [Ψ]

Corrected events: 370/446

Wild type [Ψ]

Zfp420
Nin *Nrcam*
Pfkp *Mgat1*
Eml4 *Usp40*
Camk2d *Apc*
Klf12 *Mtmr1*
Nsmf
Brsk2
Gsap

0.00 0.25 0.50 0.75 1.00
MS023 + nusinersen [Ψ]

1
2
3 **Figure 5. Combinatorial treatment with MS023 and ASO results in improved correction**
4 **of the SMA transcriptomic signature compared to ASO alone. A,** Pie charts showing the

proportion of transcripts normalised upon the treatments, among the genes that are
upregulated (left) and downregulated (right) in SMA mice. **B**, Normalised counts were used
to generate the hierarchical clustering heatmap. Upregulated and downregulated genes are
displayed in yellow and blue, respectively. **C**, Hallmark pathway analysis of genes
upregulated and **D**, downregulated in SMA corrected by the combinatorial treatment only. **E**,
Plot charts show the distribution of splicing events (Ψ) in SMA mice upon treatment relative
to wild type littermates. The red dotted lines mark the ± 15% normalization range, corrected
values are indicated in green.
9