In depth characterization of an archaeal virus-host system reveals numerous virus exclusion mechanisms

Coraline Mercier[1], Daniela Thies[1], Ling Zhong[2], Mark J. Raftery[2], Ricardo Cavicchioli[3], Susanne Erdmann*[1,3]

[1]Max Planck Institute for Marine Microbiology, Archaeal Virology, Celsiusstrasse 1, 28359, Bremen, Germany.

[2] Bioanalytical Mass Spectrometry Facility, The University of New South Wales, Sydney, New South Wales, 2052, Australia.

[3] School of Biotechnology and Biomolecular Sciences, The University of New South Wales, Sydney, New South Wales, 2052, Australia.

* To whom correspondence should be addressed: Susanne Erdmann. Tel: +49 421 2028-7340; Email: serdmann@mpi-bremen.de

Authors Contributions

 C.M. and D.T. performed a majority of the experimental work. S.E. performed some of the experiments, conceived and led the study, and performed the primary writing of the manuscript. R.C. led the initial phase of the study. L.Z and M.J.R performed mass spectrometry and analyzed data. All authors participated in the analysis and interpretation of the data and contributed to the writing of the manuscript.

Competing Interest Statement:

The authors declare no competing interests.

Classification:

Biological Sciences, Microbiology

Keywords: Archaea, archaeal virus, CRISPR, virus defense mechanisms

Abstract

Deep Lake, Antarctica, is a hypersaline lake characterized by a very low microbial diversity, dominated by archaea, and a lack of grazing organisms, making viruses important players in regulating community structure. Here we present an in-depth characterization of an archaeal virus-host system isolated from Deep Lake. Upon purifying the environmental virus lysate through plaque assays, with the intention to generate a clean and stable virus-host system, we select for a virus variant with an exclusively lytic life cycle (HRTV-DL1). The host *Halorubrum lacusprofundi* exhibits a large arsenal of virus exclusion mechanisms, indicating a long ongoing arms race with viruses. However, we uncover that the majority of this arsenal was lost spontaneously in a strain grown under non-challenging laboratory conditions. By challenging both the parental strain and the sensitive strain with HRTV-DL1, we discovered a number of putative virus exclusion mechanisms that are only activated in the sensitive strain upon the lack of defense systems present in the parental strain. We identify one of two S-layer proteins as primary receptor for HRTV-DL1 and conclude that the presence of two different S-layer proteins in one strain provides a strong advantage in the arms race with viruses. We propose the involvement of origin recognition proteins (ORC1/CDC6) in virus exclusion. While *Hrr. lacusprofundi* and HRTV-DL1 represent a great archaeal model system to observe the arms race between viruses and hosts, our work also demonstrates that an isolated model virus-host systems does not reflect the entire spectrum of interactions as they occur in the environment.

Significance statement

Archaeal head-tailed viruses appear, at first sight, very similar to head-tailed bacteriophages simply due to their morphological similarities and similar life cycles. However, they encounter host cells that are very different from bacteria and share characteristics, that greatly influence virus life cycles, with eukaryotes. Our study presents an in-depth characterization of the virus-host interactions between an archaeal head-tailed virus and its host, uncovering a number of previously undescribed putative virus exclusion mechanisms and reflecting the differences between bacterial and archaeal head-tailed viruses. By bringing this system into the laboratory and isolating a pure model system, we generated a virus-host pair with reduced genomes that barely reflects this virus-host pair in the environment, demonstrating the disadvantages of laboratory model systems.

Introduction

Deep Lake is a cold, marine-derived hypersaline lake in Antarctica with temperatures ranging between -20°C in the bulk of the lake and up to +12°C at the surface [1]. The lake is globally isolated and is characterized by a very limited microbial diversity [2]. The community is dominated by four halophilic archaeal species (haloarchaea) accounting for ~72% of the population, with *Hst. litchfieldiae* tADL accounting for 43.5%, halophilic archaeon DL31 for 18.2%, *Halorubrum lacusprofundi* for 9.9% and *Halobacterium* DL1 for 0.9% [3]. These species, except one (*Halohasta litchfieldae* tADL), exhibit one main chromosome and one or several secondary replicons (secondary chromosomes or megaplasmids). The secondary replicons, that are also common in other haloarchaea [4], were shown to be extremely diverse even between strains of the same species, in particular in *Hrr. lacusprofundi* [5]. While some of these secondary replicons strongly interact with the main chromosomes and exhibit some genes that are essential [6], others have been lost and shown to be non-essential [7]. Nevertheless, the majority of genes encoded on secondary replicons are thought to allow adaptation to changing environmental conditions, but are possibly not essential under stable laboratory conditions. *Hrr. lacusprofundi* exhibts three replicons, the main chromosome CHR1, a secondary chromosome CHR2 and the megaplasmid pHLAC01. One ribosomal rRNA operon is located on CHR2 and a number of virus defense systems, including two CRISPR-Cas systems, are found on both CHR2 and pHLAC01. A metagenomics study on Antarctic salt lakes revealed that the main chromosome of *Hrr. lacusprofundi* is highly conserved between species, but the secondary chromosomes and megaplasmids are very diverse and responsible for the genetic diversity of the species [5]. A plasmid mobilized between *Hrr. lacusprofundi* strains in specialized membrane vesicles was proposed to be involved in generating this diversity [8].

Owing to its low-complexity community and the closed nature of the system, Deep Lake is ideal for studying virus-host interactions and the impact of viruses on a microbial community. Only one putative primary producer, the green alga *Dunaliella* sp., was identified in the lake, however, no grazing organisms were detected. Therefore, while viruses are important regulators of community structure in all environments [9], we suggest that they play a particular important role in Deep Lake as one of the major players influencing microbial community structure. Several omics-based studies on Deep Lake revealed an ongoing arms race between the Deep Lake haloarchaea and viruses, and suggest that viruses are one of the drivers of genetic exchange in the community [3, 5, 10]. However, in depth characterization of virus-host interactions, including the life cycle of viruses, the host range and the influence on host viability, is only feasible on an isolated virus-host system. Even though the major microbial players have been isolated, no virus was isolated from Deep Lake so far. In an effort to understand the influence of viruses on Deep Lake haloarchaea we envisaged to isolate and characterize representative virus-host systems, subsequently allowing the assessment of the respective relationships in their natural environment using metagenomics data from several years and different seasons .

While studies on archaeal viruses in general are still significantly underrepresented, the majority of isolated viruses to date have been described for members of Halobacteriota, Methanobacteriota and Thermoproteota phyla. Haloarchaea (belonging to the Halobacteriota) are infected by diverse viruses including head-tailed viruses, pleomorphic viruses, spherical viruses and spindle shaped viruses [11]. A number of haloarchaeal viruses exhibit lysogenic and chronic life cycles, however, the majority of isolated viruses exhibit lytic life cycles, reflecting a methodical bias caused by using plaque assays for isolation, selecting mostly for lytic viruses [12]. A recent study described a number of new haloarchaeal

virus genomes and proposed a new classification of archaeal tailed viruses (arTV) [13]. Even though arTV appear morphological similar to head-tailed bacteriophages, they encounter cells that are dramatically different, in particular with respect to the cell surface, replication, transcription and translation mechanisms. Nevertheless, to date only very few arTV infecting haloarchaea have been characterized in more depth [11, 13].

Recent studies, using bioinformatics approaches, identified an enormous diversity of previously unrecognised bacterial and archaeal virus exclusion mechanisms. While some of these defence systems have been characterized in bacteria [14], their activity in archaea remains to be elucidatedAdditionally, archaeal viruses are in general significantly more diverse than isolated bacterial viruses, raising the question as to which extent the currently known defence systems are active against which range of archaeal viruses. Moreover, the fact that archaea share characteristics with eukaryotes potentially expands the range of potential virus exclusion mechanisms, that are not present in bacteria, to be discovered.

Here we present an in-depth characterisation of an isolated virus-host system from Deep Lake including the characterisation of the virus life cycle, determination of the virus host range, analysis of the host transcriptional response to virus infection and the characterisation of virus escape mutants. We uncover the genomic plasticity of both the host and the virus. We determine the host receptor for the virus and discover a number of new potential virus exclusion mechanisms. Establishing this isolated virus-host system indeed allows to uncover many molecular aspects of virus–host interactions that are not extractable from 'omics' data. However, we also realized that the isolated system only allows a glimpse into the possible interactions ongoing in the natural environment, which perhaps can be elucidated using 'omics' data.

Materials and Methods

Isolation of HRTV-DL, strains and culture conditions

HRTV-DL, formerly described as DLHTHV [5], was isolated from the supernatant of a colony that lysed upon propagation in liquid culture. Briefly, a sample, taken from Deep Lake (Antarctica) [1] in 2014, was used to generate enrichment cultures in liquid medium (culture medium and conditions see below). Enrichment cultures were plated on solid media and a single colonies picked into liquid media. The colony was identified as *Halorubrum lacusprofundi* by 16s rDNA sequencing using universal primers (21F and 1510R [15]). After lysis was observed in the culture, cells and cell debris were removed by centrifugation (4,500 x *g*, 30 min), the supernatant of the culture (lysate) was filtered through 0.2 um filters and used for infection of *Halorubrum lacusprofundi* ACAM34. *Halorubrum lacusprofundi* ACAM34 (DSM 5036), hereafter named ACAM34 _DSMZ, was obtained from the German Collection of Microorganisms and Cell Cultures (DSMZ). The parental strain of *Halorubrum lacusprofundi* ACAM34_UNSW was provided by either Peter Franzmann or John Bowman (not traceable anymore) and subsequently cultured in the laboratory (Ricardo Cavicchioli, School of Biotechnology and Biomolecular Sciences, The University of New South Wales, Sydney) from glycerol stocks stored at -80 °C. The strain went through several rounds of culturing, plating and -80 °C storage in three different laboratories (UNSW and University of Technology, Sydney, Australia; Max Planck Institute for Marine Microbiology, Bremen, Germany) prior to re-sequencing. The number of generations that the strain had undergone during this time is not traceable. *Halohasta litchfieldiae* tADL (DSM 22187) was provided by the DSMZ. Halophilic archaeon DL31 and *Halobacterium* DL, and *Hrr.*

lacusprofundi R1S1 and DL18 were isolated previously [3, 8]. All strains were grown in DBCM2 media
[16], with 5 g peptone and 1 g yeast extract added per liter. Incubation temperature was 28 °C unless
stated otherwise. Cultures were incubated in glass flasks aerobically at 120 rpm. Solid ager plates
contained 16 g agar per liter and top layer agar contained 6 g agar per liter. Plaque assays were
performed by mixing virus and host with 10 ml of top layer agar that was subsequently poured on solid
plates and incubated at 28 °C until growth was visible.

Isolation and purification of HRTV-DL1 particles

For virus production, *Hrr. lacusprofundi* was grown in liquid culture to mid exponential phase, with an
optical density at 600 nm (OD_{600}) of 0.5. Cells were harvested (4,500 x g for 45 min), mixed with viral
suspension with a multiplicity of infection (MOI) of 5-10 and incubated for 2 hours at room temperature
to allow viral adsorption. Treated cells were inoculated into 500 ml liquid cultures that were monitored
by measuring optical density changes (OD_{600}). After lysis occurred, cultures were centrifuged at 4,500
x g for 45 minutes to pellet the cells. The supernatant was recovered and viruses were subsequently
precipitated with polyethylene glycol (PEG) 6,000 (10 % w/v final concentration) and incubation at 4
°C overnight. Then, viral preparations were collected by centrifugation (30,000 x g, 45 min, 4 °C).
Pellets were resuspended in DBCM2 salt solution (DBCM2 media without nutrient sources pyruvate,
trypton and yeast extract), sterile filtered (pore size 0.2 µm) one time when used for plaque assay and
twice when used in liquid culture infection assays. Virus solutions were stored at 4 °C and were active
for a minimum of 1 year. For downstream analyses such as genome sequencing and mass spectrometry
the virus particles were further purified by Cesium chloride (CsCl) density gradient centrifugation. The
virus solution was treated with 200 U of DNase I and 50 ug/ml of RNase A to reduce host genomic
contamination prior to loading on a CsCl density gradient (0.45g CsCl /ml in DBCM2 salt solution) and
centrifuged at 38,000 rpm for 22 hours, 4 °C (SW41 Ti Swinging-Bucket rotor, Beckman & Coulter).
Bands containing virus particles were extracted with a syringe, diluted in three volumes of DBCM2 salt
solution and re-precipitated with PEG (final concentration 10 %, 4 °C, overnight). After centrifugation
at 30,000 x g for 30 min the resulting pellet was washed twice with DBCM2 salt solution and stored at
-20 °C before further processing.

Imaging

191 For Transmission electron microscopy (TEM), virus containing solution was adsorbed for 5 min to
192 carbon coated copper grids and stained for 1 min with 2 % uranyl acetate (w/v in water). Electron
193 micrographs were generated using JEM2100 Plus at 200 kV acceleration voltage. For light microcopy,
194 cells were fixed with 1 % glutaraldehyde for 1 hour at room temperature and then stored at 4 °C until
195 imaging with a Zeiss AxioPhot microscope with AxioCam MRm.

DNA extraction, manipulation and genome sequencing.

Genomic DNA of all samples, if not otherwise stated, was extracted using genomic DNA extraction kit
(Bioline, London, UK) according to the manufacturer's instructions. Plasmid extraction was performed
with ISOLATE II Plasmid Mini Kit (Bioline, London, UK) according to the manufacturer's
instructions. Nuclease digestions using specified restriction enzymes (New England Biolabs [NEB], 20
units) or exonuclease III (NEB, 5 units) were performed on 2 – 3 µg of DNA extracted from virus
particles for 1 h at 37 °C. PCR reactions were performed with the Q5® High-Fidelity DNA Polymerase
(NEB) and contained 0.02 units/µL of DNA polymerase, primer concentration of 0.1 µM for both
forward and reverse, 1X of Q5 Reaction Buffer and 1X Q5 High GC Enhancer. The following program
was used: 5 min at 95 °C, followed by 35 cycles of 30 s at 95 °C, 30 s at 68 °C for annealing and 30 s

at 72 °C for elongation. Digested DNA and PCR products were separated on 1 % agarose gels in Tris-
borate-EDTA buffer and stained with SYBR™ Safe DNA stain (Invitrogen). For HRTV-DL, library
preparation (Nextera XT) and sequencing (MiSeq) was performed at the Ramaciotti Centre for
Genomics (UNSW, Sydney, Australia). For HRTV-DL1, UNSW_1.7, UNSW_2.14 and UNSW_3.3
library preparation (FS DNA Library, NEBNext® Ultra™) and sequencing (Illumina HiSeq3000, 2 x
150 bp, 1 Gigabase per sample) was performed at the Max Planck-Genome-Centre Cologne (Cologne,
Germany). For PacBio sequencing of ACAM34_UNSW and ACAM34_DSMZ, DNA extraction,
library preparation, sequencing (Pacific Biosciences Sequel, 4 samples on one SMRT cell) and
assembly ('Microbial Assembly' function in PacBio SMRT® Link v8.0 for ACAM34_DSMZ and flye
assembly tool v2.6 [17] for ACAM34_UNSW) was performed at the Max Planck-Genome-Centre
Cologne (Cologne, Germany).

Genome assembly, annotation and phylogenetic analysis

After assembly with SPAdes [18], the HRTV-DL genome was manually closed to one contig using
primers listed in Supplementary table 1, followed by Sanger sequencing (MICROMON DNA
Sequencing Facility, Monash University, Victoria, Australia) of PCR products. The HRTV-DL1
genome was assembled using metaviralSPAdes [19]. Genome annotation for all genomes was done
using Prokka [20] followed by manual corrections using conserved domain based searches [21] or
hidden Markov model (HMM) based searches [22]. For genome comparison HRTV-DL and HRTV-
DL1 nucleotide sequences were compared using BlastN (standard settings) and comparison was
visualized using Artemis comparison tool [23]. ORFs of HRTV-DL and HRTV-DL1 were compared
using BlastP (standard settings) (Supplementary table 2). Phylogenetic tree reconstructions from protein
sequences was done with VICTOR under optimal settings (formula VICTOR d6), as implemented at
the DSMZ webserver https://victor.dsmz.de. Prediction of termini was done using PhageTerm [24].

Protein content analysis

Protein content analysis of HRTV-DL1 particles was performed by mass spectrometry on purified virus
pellets as described previously for plasmid vesicles and membrane vesicles [8]. Peak lists were
generated using Mascot Daemon/Mascot Distiller (Matrix Science, London, England) and initially
submitted to the database search program Mascot (Matrix Science). Search parameters were: Precursor
tolerance 4 ppm and product ion tolerances ± 0.05 Da; Met(O) carboxyamidomethyl-Cys specified as
variable modification; enzyme specificity was trypsin; 1 missed cleavage was possible; customized
databases searched: *Hrr. lacusprofundi* ACAM34 and HRTV-DL. After genome sequencing of
ACAM34_UNSW, results presented in Supplementary Table 4 were obtained using the SEQUEST
search algorithm with Thermo Proteome Discoverer™ 2.5.0.400 (Thermo Fisher Scientific) with the
same settings searching customized databases *Hrr. lacusprofundi* ACAM34_UNSW and HRTV-DL1.

Virus infectivity and kinetics

To study the life cycle, cultures of *Hrr. lacusprofundi* (all strains) were synchronized using an
adaptation of the "Stationary phase method" [25]. The strain was scratched from a -80 °C glycerol stock
into liquid culture and grown up to an $OD_{600} \sim 1$. Then a 20-fold dilution step in fresh media was
performed (final $OD_{600} = 0.05$) and cultures were then regrown up to $OD_{600} \sim 1$. Iterative dilution and
growth of the culture were repeated twice before considering a culture synchronized. For infection with
HRTV-DL1, cells from synchronized cultures ($OD_{600} \sim 1$) were collected by centrifugation,
resuspended in 1 ml of fresh media and infected with HRTV-DL1 virus with a MOI of 5. After
incubation (2 h, room temperature), cells were transferred into liquid cultures and growth was monitored

by optical density (OD$_{600}$). Long-term cultures were established by continuously diluting infected
cultures at OD$_{600}$ ~ 1 to OD$_{600}$ ~ 0.05. Infection was confirmed by PCR using primers HRTV-DLF and
HRTV-DLR (Supplementary table 1).Viral titer in culture supernatants was quantified by plaque assay
after removal of cells by centrifugation (11,000 x g, 10 min, room temperature [RT]). Intracellular virus
titers and host chromosome copy numbers were quantified by qPCR. Briefly, samples of 2 ml culture
in biological replicates were collected and pelleted (11,000 x g, 10 min, RT). Cell pellets were washed
two times with 1 ml of fresh media and stored at -20 °C upon DNA or RNA extraction. Quantification
of *Hrr. lacusprofundi* and HRTV-DL1 genome copy number were carried out using a CFX96 Touch
Real-Time PCR (Bio-Rad Laboratories, Inc., Hercules, CA, USA) and the software CFX Manager™
Software. Primers are listed in Supplementary table 1. Each reaction (10 µL) contained 1X
SsoAdvanced Universal SYBR™ Green Supermix (Bio-Rad) and primer concentrations as stated in
Supplementary table 1. The following amplification thermal cycling program was used for both primer
sets: 5 min at 95 °C, followed by 40 cycles of 30 s at 95 °C and 30 s at annealing temperature stated in
Supplementary table 1, with readings taken between each cycle. Efficiencies of the assays were 95–
102 %, with R^2 values ≥0.99 for all assays. The specificity of the qPCR was confirmed by unique
signals in melting curves and gel electrophoresis of PCR products.

Adsorption assays and host range assessment of HRTV-DL1

For adsorption assays, 5 ml of cells at OD$_{600}$ = 0.5 were harvested by centrifugation (4,500 × g, 30 min,
RT) and resuspended in 1ml fresh medium. The cells were subsequently infected using a MOI of 5. At
defined time intervals the adsorption was stopped by immediate centrifugation (10,000 × g, 5 min, RT).
The number of remaining free virus particles in the supernatant was determined by plaque assay. A cell-
free control (only media) served as control. Three different strains of haloarchaea (*Halohasta*
litchfieldiae tADL, halophilic archaeon DL31, and *Halobacterium* DL1 [3]) and two additional strains
of *Hrr. lacusprofundi* (R1S1 and DL18 [8]) were tested to determine the host range of HRTV-DL1.
CRISPR matches to HRTV-DL1 were identified by searching for CRISPR loci using CRISPRs finder
[26] and blasting (BlastN) identified spacer against the HRTV-DL1 genome. Adsorption of HRTV-
DL1 to the different strains was determined by adsorption assay as described above and the absence of
HRTV-DL1 DNA within cells was confirmed by PCR on cell pellets 1 day and 5 days post infection.
278
Transcriptomic analyses

RNA extraction of frozen cell pellets with 3 biological replicates for the infected and 2 biological
replicates for the controls, was performed with the Direct-zol™ RNA miniprep Kit (R2051, Zymo
Research). RNA concentration and integrity were assessed using Nanodrop DS-11 Spectrophotometer
(DeNovix) according to the manufacturer's instructions. Library preparation and sequencing was done
at the Max Planck-Genome-Centre Cologne (Cologne, Germany). Briefly, ribosomal RNA were
depleted prior to sequencing using the rRNA depletion Kit riboPOOL™, for *Haloferax volcanii*
(88.36% identity to 16s rDNA sequences of *Hrr. lacusprofundi*), siTOOLs Biotech®. Libraries were
prepared with library kit NEBNext® Ultra™ II RNA Library Prep Kit for Illumina and sequencing was
performed on an Illumina HiSeq3000 sequencer, following a 1 x 150 run. Read trimming and mapping
was performed with the 'Map to reference' function (Mapper 'Geneious RNA') with medium-low
sensitivity within Geneious Prime® 2022.2.1. Expression values (FPKM values) were calculated using
standard settings and comparison of expression levels were performed using DeSeq2 within Geneious
Prime® 2022.2.1 using default settings. Genes with p-values < 0.01 and a fold change of at least two
times (log2FC ≥ 1 or ≤ -1) were considered to be differentially expressed (DE).

Analysis of ACAM34_UNSW escape mutants

For the isolation of ACAM34_UNSW HRTV-DL1 escape mutants we established long-term infected
cultures as described above (Virus infectivity and kinetics), in triplicates. Cells were only infected with
an MOI of ~1 to allow a recovery of infected cultures. After lysis of ACAM34_UNSW the cultures
were maintained to allow the growth of HRTV-DL1 resistant clones. Cells from the recovered cultures
were plated on agar plates to obtain single colonies. HRTV-DL1 infection was determined by PCR as
described above. HRTV-DL1 life cycle in escape mutants and adsorption of HRTV-DL1 to escape
mutants was determined as described above. Isolation of genomic DNA and genome sequencing is
described above. For genomic analysis of escape mutants, reads were mapped against
ACAM34_UNSW and HRTV-DL1 using 'geneious mapper' with medium-low sensitivity and default
settings and variants were called using the Geneious function 'Find Variations/SNPs' with default
settings in Geneious Prime® 2022.2.1. Variants with a coverage below 150 and a variant frequency
below 75% were excluded from the analysis. HRTV-DL1_1.7 was assembled using MetaviralSPades
[19].

Results and Discussion

Isolation of a lytic virus that infects Halorubrum lacusprofundi ACAM34 (DSM 5036)

A virus with head-tailed morphology (Figure 1b) was obtained from the culture supernatant of a
Halorubrum lacusprofundi strain, that was isolated from a sample taken in 2014 from Deep Lake,
Antarctica [1]. Analytic restriction digest of DNA isolated from the viral lysate revealed a dsDNA
genome of approximately 60 kb (Figure 1a). The lysate produced plaques on *Hrr. lacusprofundi* strain
ACAM34 [27] that was isolated from a Deep Lake sample in 1988, while it remains elusive when the
sample was actually obtained from Deep Lake. Lysis was also observed in liquid cultures of this strain.
The viral lysate was formerly describe as DLHTHV [5], however, we rename it hereby as HRTV-DL
(**H**alo**R**ubrum **T**ailed **V**irus-**D**eep **L**ake), according to the nomenclature that was previously used for
archaeal tailed viruses (arTV) [13]. HRTV-DL is the first virus infecting *Hrr. lacusprofundi* that has
been isolated from Deep Lake and, interestingly, even though the virus was isolated at least 26 years
later than the host, it is still able to generate a successful infection. *Hrr. lacusprofundi* ACAM34 was
used to purify the virus through several rounds of plaque assays. When analysing genomic DNA of
virus particles propagated from single plaques, we observed a remarkable high genomic variability
(Supplementary figure 1). Such a genomic variation has previously been observed for *Halobacterium*
salinarium virus ØH [28]. After two rounds of plaque purification, we isolated a variant that appeared
to have a stable genome through several series of infection (P2V1, Supplementary figure 1c) and we
choose this variant, HRTV-DL1, for further characterization. Virus particles analysed by electron
microscopy exhibit a head-tailed morphology, with a head diameter of about 50 nm and a non-
contractile tail of approximately 80 nm-100 nm in length (Figure 1b, Supplementary figure 2).
331

Genome comparison of the original virus lysate and the isolated variant genome reveals genome plasticity of the virus and unintentional selection for a lytic virus variant

334 DNA isolated from HTRV-DL1 particles appeared sensitive to Exonuclease treatment, while the
335 majority of virus DNA isolated from infected cells was shown to be circular (Supplementary figure 3),
336 indicative of circularisation of the viral genome after injection into the host. Analytic digest of virus
337 and host DNA with *Dpn*I, showed that both the host and the virus genome are dam methylated.

Methylation of the virus genome likely supports successful proliferation of the virus within its host, allowing the virus to escape host Restriction-modification (R-M) defence systems. Both the original lysate (HRTV-DL) and the isolated variant HRTV-DL1 were subjected to genome sequencing. The genome of HRTV-DL assembled in a number of contigs, that could not be computationally connected, and the genome of HRTV-DL1 assembled into a single circular contig of 37.7 kb. Since the HRTV-DL1 genome was shown to be linear within the virus particle but assembles as a circular contig, with no indication in the sequence data for discrete ends, we conclude that the genome is circularly permuted and terminally redundant. Contigs of HRTV-DL could manually be connected by PCR into one contig of 66.2kb, similar to what has been estimated by analytic restriction digest (Figure 1a). We assume that we were able to close it to only one contig, because the genome is circularly permuted and terminally redundant, however, it would never appear as one complete genome in one virus particle. Comparison of the two genomes revealed that HRTV-DL contains several duplicated regions and a few regions that are unique (Supplementary figure 4). This included 10 hypothetical proteins, 2 integrases and one adenine-specific DNA methylase. Additionally, the phage tail tape measure protein is 28aa shorter in HRTV-DL1 (Supplementary table 2). We suggest that upon initial isolation of HRTV-DL, the viral lysate contained a mixture of different virus variants that shared the majority of their genome. Co-infection allowed the recombination of the different genomic regions and the generation of diverse virus variants as observed during plaque purification (Supplementary figure 1). Interestingly, none of the integrases that are present in the HRTV-DL lysate was found in the isolated variant HRTV-DL1 that exclusively exhibits a lytic cycle. By using plaque assay for purification, we selected for a virus variant with a lytic cycle and thereby lost variants that would be able to integrate into the host genome and exhibit a lysogenic life cycle. After obtaining the genome of HRTV-DL we established that the virus is present in Deep Lake by blasting the virus genome against available metagenomes from 2006, 2008, and 2013-15 [5] in the Integrated Microbial Genomes and Microbiomes (IMG) database [29]. *Hrr. lacusprofundi* was also detected in Deep Lake in the same metagenomes [5], confirming that both host and virus have been co-existing over several years.

Genomic features of HRTV-DL1 and proteins associated with the virus particles

The genome of HRTV-DL1 encodes 49 putative open reading frames (ORFs) (Figure 2, Supplementary table 3). Using Hidden Markov models (HMM) [22] and domain prediction [21] we can predict the function for about 40 % of the annotated open reading frames (ORF) (Supplementary table 3). Two ORFs were identified that we propose to be involved in virus genome replication, both are also encoded by arTV CGphi46 (Figure 2). ORF48 is predicted as primase-helicase, probably generating a short primer that is then elongated by the host DNA polymerase, and ORF41, predicted as DNA polymerase sliding clamp, binding the DNA polymerase and preventing its dissociation from the template DNA. Two ORFs could be involved in transcription. ORF43, a predicted transcription initiation factor IIB, might be responsible for recruiting the host RNA Polymerase and ORF32, with weak hits to antitermination protein NusA, could associate with the host RNA polymerase elongation complex. ORF29 is a predicted adenine-specific methyltransferase and 100 % identical to a host protein, probably recruited by the virus to protect its genome from host R-M-systems. ORF7 and ORF8 were identified as small and large subunit of the terminase, responsible for genome packaging into the virus particle.

Proteins building the virus particle were identified by mass spectrometry of purified virus particles and confirm our predictions (Supplementary table 3 and 4). The major capsid protein belongs to the HK97-like fold and is most similar to that of arTV BJ1 (45 % identity). Additionally, we could identify the phage portal protein, two capsid proteins with similarity to *Natrialba* phage PhiCh1 capsid proteins, the putative head-tail adaptor, the tail tape measure protein and the tail terminator protein. ORF18 and

ORF25, identified in the virus particle, did not show any similarities to other viral proteins, therefore,
their role within the virus particle remains enigmatic. Two proteins that could be involved in host
attachment were identified. ORF24, that is present in virus particle, exhibits a immunglobulin-like fold
(fibronectin type III domain) at the N-terminus. This domain is commonly found in bacterial phage tails
and has been shown to exhibit membrane binding activity [30-32]. In the C-terminus we identified a
sialidase domain. Sialidase (neuraminidase) is known to cleave glycosidic linkages and is essential for
some eukaryotic viruses [33]. It could be involved in destabilizing the glycosylated S-layer and
facilitating access to the host membrane. ORF25, also detected in virus preparations, shows similarities
to a phage receptor binding protein with a high probability (P= 99.06). ORF4, with significant
similarities to DNA double-strand break repair protein MRE11, and ORF43, annotated as transcription
initiation factor, were also detected in virus particles. Both could be associated with the virus genome.
ORF4 might facilitate circularization of the virus genome within the host, while ORF43 could recruit
the host RNA polymerase ensuring timely transcription of the viral genome after injection.

Finally, we attempted to identify proteins involved in cell lysis. The mechanism of egress by complete
membrane disruption, as it is known for head-tailed viruses of bacteria, has so far not been elucidated
for arTV [34]. Lysis in gram negative bacteria is usually induced by two proteins. A holin that
permeabilizes the lipid layer and an endolysin that subsequently degrades the cell wall [35].
Haloarchaea don not exhibit a peptidoglycan cell wall, but are protected by a protein shell, the S-layer.
Specific proteases degrading the S-layer have not been found yet [34] and we could also not identify a
protease on the HRTV-DL1 genome. However, glycosylation plays a significant role for the stability
of the S-layer in haloarchaea [36] and degradation of the glycan can lead to a destabilisation. ORF26
shows similarity (using Hidden Markov models [22]) to a levanase and other glycosyl hydrolases with
a high probability (P=99.12, 98.84 respectively) and could be responsible for glycan degradation and
S-layer destabilization. Identifying holins is difficult due to their sequence diversity and only a few
putative holins have been detected in archaea, but none on arTV genomes [37]. Holins are typically
small proteins (75-174 aa) with 1-4 transmembrane domains (TM-domain). We can identify 3 small
proteins with TM domains on the HRTV-DL1 genome, ORF6 (100 aa, two TM domains), ORF20 (67
aa, one TM domain) and ORF27 (222 aa, one Signal peptide, one TM domain). ORF6 and ORF27 hit
other arTVs with low identity, however, none of the 3 ORFs show any similarities to proteins in public
databases. Most holins are encoded by the gene adjacent to the endolysin gene in bacteria, making
ORF27 to a potential candidate for a holin. Protein structure prediction of ORF27 [38] and comparison
with the PDB database did not reveal any significants hits, however, only one holin structure has been
determined so far [39]. Therefore, a potential involvement of ORF26 and ORF27 in cell lysis will need
experimental investigation.

Genome comparison with 63 arTV (archaeal tailed viruses) [13] and phylogenetic tree construction
(Supplementary figure 5), suggests HRTV-DL1 to be closest related to arTV BJ1 [40] and CGphi46
(NC_021537) (Figure 2) that have been classified into the *Flexireviridae* family. However, HRTV-DL1
is sufficiently different to be classified into a different genera, that we propose to name *Deelavirus* in
accordance with its origin (**Dee**p **L**ake).

Life cycle of HRTV-DL1

To gain insights into the initial stages of HRTV-DL1 entry, we followed the adsorption of HRTV-DL1
to ACAM34 host cells. The adsorption was very rapid, with ~75 % of virions being bound to cells
within the first 30 seconds of infection. Further adsorption plateaus after 15 min with 99 % of particles
bound to host cells (Supplementary figure 6). Such a rapid adsorption is rather uncommon for

haloarchaeal arTV. Haloarcula hispanica tailed virus 1 (HHTV-1; *Siphoviridae*), binds extremely slowly with virions adsorbed only after 3h and binding of *Halorubrum* virus HRTV-1 was only detected after 25min [41]. Such diversity in adsorption rates indicates differences in adsorption mechanisms and receptors. The rapid adsorption of HRTV-DL1 suggests that the receptor is easily accessible and highly abundant on the cell surface. After successful infection, host cell lysis occurred between 56 and 68 hours post infection (p.i.) at 28 °C. The virus-to-host ratio (VHR) was determined by qPCR as 95 viral genome copies per host main chromosome copy number a few hours prior lysis (Figure 3). Interestingly, we observed changes to the cell morphology of infected *Hrr. lacusprofundi* at the onset of cell lysis. While the majority of the uninfected cells retained their rod-shaped morphology, infected cells tended to round up in the late stages of infection (Supplementary figure 7). However, a significant increase in cell size, as shown for another archaeal virus [42], was not observed.

Both *Hrr. lacusprofundi* and HRTV-DL1 were isolated from a lake that experiences very low temperatures [1]. To determine whether low temperatures have an influence on the virus life cycle, we performed growth experiments at different temperature between 4 °C and 30 °C. Host growth rates change at different temperatures, consequently, lysis was observed at different time points post infection. However, lysis always occurred in early exponential phase at all temperatures (Supplementary figure 8). Therefore, we conclude that temperature does not influence the life cycle of HRTV-DL1 under the laboratory growth conditions tested. However, the presence of integrases in the HRTV-DL lysate indicate that other virus variants could exhibit a lysogenic life cycle, and this life cycle could well be influenced by temperature in its natural environment.

HRTV-DL1 exhibits a very narrow host range that uncovers genetic differences in the same ACAM34 strain from different sources

HRTV-DL1 was isolated from Deep Lake, that is dominated by 4 different haloarchaea (*Halohasta litchfieldiae* tADL, *Hrr. lacusprofundi*, halophilic archaeon DL31, and *Halobacterium* DL1) [3]. CRISPR spacers against HRTV-DL1 were detected in the CRISPR loci of all except 1 species (*Halobacterium* DL1) (Supplementary table 5). The presence of spacers against HRTV-DL1 in the CRISPR loci of *Hrr. lacusprofundi* indicate that the virus, or variants of it, have been present in Deep Lake already before 1988. CRISPR spacers of all three different Deep Lake haloarchaea target different positions on the virus genome, indicating that they were acquired independently and not by gene transfer. Therefore, DL31 and tADL could be potential host organisms for HRTV-DL1. We tested all 4 species, one other *Hrr. lacusprofundi* strains from Deep Lake (DL18) and *Hrr. lacusprofundi* R1S1 from Rauer Lake 1, Antarctica [5, 8], but only *Hrr. lacusprofundi* ACAM34 was susceptible to infection. Adsorption assays revealed that attachment of the virus to all tested strains is impaired (Supplementary figure 9), showing that infection is already compromised at the adsorption stage. We assume that the attachment site of HRTV-DL1 is highly diverse, even in different strains of the same species. Modifications of cell surface proteins, in particular the S-layer that was found to be highly distinct, has already been proposed as a virus exclusion mechanisms of Deep Lake haloarchaea [5, 10].

When continuing characterisation of HRTV-DL1 in a new laboratory, we used a fresh stock of *Hrr. lacusprofundi* ACAM34 provided by the DSMZ. Surprisingly, while the virus successfully adsorbed to the host cell and injected its genome (Supplementary figure 9), HRTV-DL1 was not able to complete a lytic life cycle in the DSMZ strain. No lysis was observed in cultures and virus DNA could not be detected within the host at 56 hours p.i. (Supplementary figure 10). We subsequently assumed that the HRTV-DL1 susceptible strain, from now on referred to as ACAM34_UNSW (for University of New

477 South Wales, Australia, the location of the laboratory the strain originated) had experienced genomic
478 changes while being grown over a long time in the laboratory.
479

Genome comparison of the ACAM34 laboratory strain with the ACAM34 strain from a
culture collection reveals dramatic genomic rearrangements in the laboratory strain

To determine genetic differences between the DSMZ strain (ACAM34_DSMZ) and the HRTV-DL1
sensitive strain (ACAM34_UNSW), we re-sequenced the genome of both strains.

ACAM34_DSMZ assembled into 4 contigs. Contig 1 is circular with a size of 2,735,247 bp, (coverage
1080) representing the main chromosome (CHR1) with 99.99 % identity to the published genome.
Contig 2 is circular with a size of 525,899 bp (coverage 1167), representing the secondary replicon
(CHR2) with 99.99 % identity to the published sequence. Contig 3 is circular with a size of 431,344 bp
(coverage 1137), representing the plasmid pHLAC01 with 99.99 % identity to the published sequence.
Contig 4 is linear with the size of 6,176 bp matching a short region on CHR2 covering Hlac_3232 to
Hlac_3234 with an insertion of a transposon within Hlac_3234. This contig has a lower coverage (432)
than contig 2 and likely represents a variant of this region in the secondary replicon of the strain. In
conclusion, the genome of ACAM34_DSMZ is to 99.99 % identical with the published sequence.

Surprisingly, ACAM34_UNSW assembled into only one single circular contig of 3,058,421 bp.
Genome comparison with the published genome (Figure 4 and Figure 7) revealed that the entire plasmid
pHLAC01 and about 38 % (173 kb) of CHR2 is not present, while the remaining 61.11 % of CHR2
(Hlac_2782 - Hlac_3106) is integrated into CHR1 between Hlac_1757 and Hlac_1759, two copies of
ISH3 family transposase ISHla1 [43] with 100 % sequence identity. Hlac_2781 and Hlac_3106 flanking
the CHR2 region integrated into CHR1 are also ISHla1 transposases that are 100 % identical to
Hlac_1757 and Hlac_1759. We suggest that the integration of CHR2 occurred by homologues
recombination between the copies of ISHla1 and at the same time caused the loss of the remaining
CHR2. The insertion of a large genomic fragment into the main chromosome due to transposition
activity has already been reported for a *Halobacterium salinarium* laboratory strain [44], and the
insertion of a large plasmid into the main chromosome has been reported for *Haloferax volcanii* [7].
Apart from the insertion of CHR2, CHR1 remains highly conserved with an identity of 99.999 %. One
minor change is found in an intergenic region between Hlac_0599 and Hlac_0600 that is missing a
stretch of 23 nucleotides. The integrated remains of CHR2 have three significant changes. First, a
transposase was inserted into Hlac_2793, a predicted ADP-ribosylglycohydrolase, and destroyed the
gene. Second, Hlac_2835, with a MarR-like HTH domain within the N-terminus, has a number of
changes on amino acid level within the C-terminus. And third, one of the two 16s ribosomal RNA genes
has a number of nucleotide changes that make up 3% of the sequence. It remains unknown how the
plasmid pHLAC01 was lost in ACAM34_UNSW. The secondary replicons are thought to be important
for adaptation to varying environmental conditions. However, when grown under defined laboratory
conditions the plasmid does not seem to be essential and is therefore likely too costly to be maintained.
pHLAC01 exhibits the only CRISPR locus on the ACAM34 genome. This CRISPR locus contains 10
spacers that give a 100% match to the HRTV-DL1 genome. We therefore assume, that the resistance of
ACAM34_DSMZ against HRTV_DL1 infection could be caused by the intact CRISPR system on
pHLAC01, that is lost in ACAM34_UNSW. Nevertheless, there are also a number of other putative
virus defense systems encoded on pHLAC01 and the region of CHR2 that is lost in ACAM34_UNSW,
such as restriction modification systems (R-M-systems), Argonaut proteins and a BREX system
(Supplementary table 6). These could also cause ACAM34_DSMZ resistance to HRTV-DL1 infection.

Only two putative defense mechanism, one Argonaut protein (ACAM34UNSW_01791, Hlac_2785)
[45], is still present in ACAM34_UNSW, leaving the strain basically unprotected against viral infection
without any known virus exclusion mechanism.
526
527

Response of ACAM34_DSMZ to HRTV-DL1 infection does not show increased activity of any of the predicted virus exclusion mechanisms

To determine the mechanism underlying the resistance of ACAM34_DSMZ to HRTV-DL1 infection,
we analysed transcriptional changes at two time points 32 hours post infection (middle phase, after
infection is usually established in ACAM34_UNSW) and 56 hours post infection (late phase,
immediately before ACAM34_UNSW is usually lysed).

The ACAM34_DSMZ transcriptome recovered no reads for the virus genome in both samples, which
is consistent with our results that the virus genome copy number is already very low at 32 hours p.i.
(time point 1 =T1) and not detectable at 56 hours p.i. (time point 2 =T2) (Supplementary figure 10).
However, when comparing uninfected controls with infected samples, we still detect some changes to
the host transcription. At T1 (32 hours p.i.) 468 genes are upregulated more than 2 fold (log2>1) and
none are downregulated, and at T2 (56 hours p.i.) we detect 98 genes up- and 5 downregulated
(Supplementary table 7). When analysing the functional potential of the upregulated genes at both time
points, the majority falls into the categories transcription, replication and repair mechanisms,
posttranslational modification, nucleotide metabolism and energy production and conversion (Figure
5). Interestingly, within the 30 most upregulated genes at T1, we identified 4 ORFs annotated as cold-
shock proteins (CSPs) (Hlac_1630, Hlac_0048, Hlac_1986 and Hlac_1312). However, none of them is
upregulated at T2, so this appears to be an early stress response to virus infection. In contrast, within
the 30 most upregulated genes at T2 we identified three ORFs annotated as universal stress proteins
(UspA) (Hlac_0044, Hlac_1730, Hlac_1573), possibly representing a late stress response to virus
infection. The most upregulated gene at T1 is Hlac_2176 a predicted ferredoxin. Ferredoxins are small
electron transfer proteins that are ubiquitous in biological redox systems [46] and upregulation of might
be a response to oxidative stress caused by the virus infection. However, Hlac_2176 shows 87%
sequence identity to a ferredoxin in *Hbt. salinarium* (OE_4217R), that was shown to be a coenzyme of
alpha-ketoacid oxidoreductases [47], suggesting a possible involvement in energy conversion. Second
most upregulated gene is Hlac_1324, that contains a winged helix DNA-binding domain and is likely a
transcriptional regulator. Other genes within the 30 most upregulated genes are some ribosomal proteins
and two other putative transcriptional regulators. While there is some basal expression, no significant
transcriptional changes (log2>1) were observed for the integrated provirus Hlac-Pro1, suggesting that
the integrated virus does not respond to superinfection, while such a response has been shown for
proviruses in *H. volcanii* [12].
559
Only one gene categorized as defence mechanisms is upregulated (Hlac_0998), however, the gene is
not predicted to be involved in virus defence mechanisms (Na -driven multidrug efflux pump). Despite
the presence of a number of spacers against HRTV-DL1 in ACAM34_DSMZ CRISPR loci
(Supplementary table 5), none of the CRISPR-Cas genes showed upregulation (log2>1) at the two time
points. Cas6b (Hlac_3333) is slightly upregulated with log2=0.5 at T1 and Cas6d (Hlac_3572) with
log2=0.75 at T2, which could indicate some activity of the CRISPR-Cas system against HRTV-DL1.
None of the other predicted virus defence mechanism (Supplementary table 6) show significant
upregulation (log2>1). Perhaps, the basal activity of the CRISPR-Cas system, or basal activity of any
of the other systems, is sufficient to exclude HRTV-DL1 (Figure 6). Alternatively, upregulation might

have only been detectable shortly after virus genome injection (before 32 hour p.i.), even though other
studies observed upregulation over an extended period during virus infection [48]. However, it is also
possible that neither the CRISPR-Cas system, nor the other systems, are the main driver of HRTV-DL1
exclusion.
573
Therefore, we endeavoured to identify other potential HRTV-DL1 exclusion mechanisms. We also
screened for upregulated genes that are not present on the genome of susceptible strain
ACAM34_UNSW. Six genes were found to be upregulated on the portion of CHR2 that is absent in
ACAM34_UNSW and 23 genes are upregulate on pHLAC01 (Supplementary table 8). The respective
ORFs include two transposases, two predicted transcriptional regulators, a cdc6/orc1 replication
initiation protein, a predicted universal stress protein, a few predicted RNA binding proteins and a
number of small hypothetical proteins. Transposons are known to be activated upon virus infection [49]
and co-localize with defense islands [50], they also have been found on virus genomes [13] and might
inactivate viruses by insertion into essential virus genes. The universal stress protein and the
transcriptional regulators are likely a host response to the stress condition rather than a virus exclusion
mechanisms. A possible involvement of CDC6 in virus exclusion is discussed below (*Characterisation*
of virus escape mutants shed light on virus-host interactions and reveal additional virus exclusion
mechanisms). The remaining RNA binding proteins and small hypothetical proteins could represent so
far undetected virus exclusion mechanisms, such as toxin-antitoxin (TA) systems or proteins
influencing viral RNA stability. However, this would need to be investigated experimentally. In
conclusion, our approach did not allow us to identify with certainty the virus exclusion mechanisms
that are responsible for HRTV-DL1 exclusion in ACAM34_DSMZ.
591

HRTV-DL1 infection induces the re-mobilization of CHR2 in ACAM34_UNSW and uncovers alternative virus exclusion mechanisms

We analysed the host response to HRTV-DL1 infection by determining transcriptional changes at two
time points 32 hours p. i. (middle phase, after infection has been established, time point 1 = T1) and 56
hours p.i. (late phase, immediately before lysis, time point 2 = T2). Reads mapping the HRTV-DL1
genome represented 0.2% of all mapped reads 32 hours p.i. and 41% of all mapped reads at 56 hours
p.i., consistent with the virus genome copy numbers increasing over time (Figure 3).

The transcriptional profile at time point 1 is very similar to that of T1 in ACAM34_DSMZ with 371
genes upregulated (log2>1) and none downregulated, and functional categories transcription,
replication and repair mechanisms, posttranslational modification, nucleotide metabolism and energy
production and conversion being the most upregulated (Figure 5). Additionally, we find the same genes
among the 30 most upregulated genes, encoding for cold-shock proteins, ribosomal proteins, the
ferredoxin Hlac_2176 and transcriptional regulators, being upregulated in ACAM34_UNSW and
ACAM34_DSMZ, indicating that both strains exhibit a very similar response to early virus infection.

However, T2 shows a different profile, with 201 genes upregulated and 333 genes downregulated. The
most upregulated functional category is 'replication and repair', consistent with a takeover of HRTV-
DL1 and the degradation of the host genome (Supplementary figure 11). Genes involved in metabolic
processes seem to be subjected to downregulation. Amongst the 30 most upregulated genes, only two
were also detected as upregulated in ACAM34_DSMZ upon virus infection (Hlac_0148, Hlac_0919).
Surprisingly, the majority of genes on former CHR2 (ACAM34UNSW_01788 to
ACAM34UNSW_02100) was upregulated (Supplementary figure 12, Supplementary Table 9).
Therefore, we assumed that this region might have been re-mobilized during virus infection and that

the upregulation is due to an increased copy number of the secondary replicon. Indeed when comparing copy numbers of CHR1 with former CHR2 by qPCR in the same samples, we detect a slightly higher copy number for CHR2 in the late stage of infected samples (1.3x the copy number of the main chromosome) (Supplementary figure 11). We conclude that CHR2 is mobilized only in a portion of cells in the population, because CHR2 usually has a copy number of 1.5-2 copies per CHR1 in ACAM34_DSMZ. When normalizing the expression values to the copy numbers, only 30 of 109 genes are still upregulated (log2>1), correcting the total number of upregulated genes from 201 to 122. As already observed in ACAM34_DSMZ, we do not detect differential expression of the integrated provirus.

Within the 30 most downregulated genes at T2 we find genes encoding for ribosomal proteins, for proteins involved in amino acid metabolism, proteins involved in energy production and conversion and proteins involved in oxidative stress response (Supplementary table 9), being a typical signature for a virus driven takeover of the host cell metabolism [51, 52].

Only one gene, predicted to be potentially involved in virus defence (ACAM34UNSW_01791), an Argonaut related nuclease (PADLOC [53]) that is still present in ACAM34_UNSW, is slightly upregulated (log2=0.9) in ACAM34_UNSW, but not in ACAM34_DSMZ (Hlac_2785) at any time point. We suggest that ACAM34UNSW_01791 is activated in ACAM34_UNSW due to the lack of other defence mechanisms, however, its activity does not seem to interrupt the lytic cycle of HRTV-DL1. Whether it has an impact on the efficiency of infection or virus production remains to be determined. To identify other potential virus exclusion mechanisms we analysed the 30 most upregulated genes. We propose that some of these genes might be implicated in virus exclusion of ACAM34_UNSW, that has lost all currently known virus defence systems (Figure 6). The predicted functions of the upregulated genes are presented in table 1 and some of particular interest, are discussed below.

Surprisingly, we find an entire genomic region being upregulated (ACAM34UNSW_00934 - _00942/ Hlac_0915 – Hlac_0923), with 8 of 9 genes within the 30 most upregulated genes. The region includes a predicted integrase (Hlac_0916), a couple of small hypothetical proteins and is enclosed by a tRNA gene and a transposase (Hlac_0923), indicating that it might have been part of a mobile genetic element that is activated upon virus infection. The region includes both partners of a two-component regulatory system, that is known to sense environmental changes and mediate the cellular response [54]. Whether the system is beneficial towards the host or high-jacked by the virus [55] remains to be elucidated.

ACAM34UNSW_01107 (Hlac_1086), that is originally annotated as transposase ORFB (conserved domain), also shows significant similarities (HMM) to Cas14 proteins that have been shown to exhibit RNA-guided nuclease activity [56]. However, since the ACAM34_UNSW does not contain any CRISPR arrays, we suggest that Hlac_1086 activity is more similar to the Transposon-associated TnpB, that is guided by an RNA derived from a sequence upstream of the TnpB [57]. Interestingly, we do find sequence similarities of the upstream region of Hlac_1086 with the PATE (see paragraph *HRTV-DL1 gene expression during infection reveals presence of a PATE with strong activity*) encoded on the virus and host genome. In absence of any other virus defence mechanism in ACAM34_UNSW, this gene might play an important role as a virus exclusion mechanism.

Two proteins (ACAM34UNSW_01618 and ACAM34UNSW_01620) within the 30 most upregulated genes, enclose a predicted chromosome maintenance protein (SMC), ACM34UNSW_01619 (log2=1.1). Additionally, another predicted SMC, ACM34UNSW_01796, is also upregulated

(log2=1.1). SMC complexes were found to interfere with virus replication in humans [58, 59].
Additionally, SMC-like proteins were found to be associated with the recently discovered Wadjet
system, that has been shown to exclude circular foreign DNA in bacteria [60, 61].
ACAM34UNSW_01618, ACAM34UNSW_01619 and ACAM34UNSW_01620 could potentially
represent an archaeal derivative of this system.

Finally, another interesting candidate for a virus exclusion mechanism is ACAM34UNSW_02085
(Hlac_3088) (within the 50 most upregulated genes when normalized by 1.3). The gene is a passenger
gene of a transposon with transposase ACAM34UNSW_02084 (Hlac_3078), and could be mobilized
similarly to other virus exclusion mechanisms [50] or antibiotic resistance genes [62]. Hlac_3088
exhibits a PGF-TERM domain, an archaeal protein-sorting motif recognized by an archaeosortase [63],
and is predicted to be a non-cytoplasmic protein with a TM domain. HMM search showed significant
similarities with the S-layer protein of *H. volcanii,* however, Hlac_3088 is too small to be an S-layer
protein. When comparing Hlac_3088 expression levels with those of the S-layer protein (Hlac_2976),
Hlac_3088 has usually lower expression levels compared to the S-layer. However, at T2 it actually
exceeds the expression levels of the S-layer gene (Supplementary table 10). We suggest that Hlac_3088
is located at the surface of the cell and could protect the cell by hiding the receptor and interfering with
the attachment of the virus, similar to TraT, encoded by the F plasmid of *Escherichia coli*, that interacts
with OmpA and blocks phage adsorption [64]. Alternatively, it could stabilize the cell and prevent or
at least delay cell lysis.

While none of the proposed mechanisms are fully effective against HRTV-DL1, because the virus is
completing its lytic cycle, they might prevent the lysis of the entire population and allow virus escape
mutants to evolve.

HRTV-DL1 gene expression during infection reveals presence of a PATE with strong
activity

Transcriptomic data from HRTV-DL1 infected ACAM34_UNSW cultures revealed that all annotated
HRTV_DL1 ORFs were expressed, while ORF3 was, by far, the most transcribed ORF (Supplementary
table 11). When examining the reads mapping ORF3, we discovered that there is a peak that expands
upstream of the annotated gene covering a region of approximately 300 bp and the majority of reads
mapped antisense to the predicted coding sequence. Upon blasting the sequence against public
databases, we discovered that it matched 99 % (1 mismatch, 1 gap) to a region within the hosts genome
and a region that was previously described in plasmid pR1SE infecting the same host [8]. Further
inspection revealed that this region represents a PATE (Palindrome-Associated Transposable Element),
with a length of 489 bp that expands over the gap between ORF2 and ORF3. Only the C-terminal end
of the PATE shows high expression, which includes the N-terminus of predicted ORF3, and expression
occurs almost exclusively in reverse direction. We assume that ORF3 is a pseudogene and expression
within this ORF is due to the PATE activity. In *Halobacterium salinarium* a PATE has been shown to
be active as ncRNA [65] and we imagine a similar function for this PATE. However, it is encoded on
both the host and the virus, so it is difficult to predict whether its activity is beneficial for the host or
the virus, and whether the virus acquired it from the host or vice versa.

ACAM34_UNSW can develop resistance to HRTV-DL1 infection

To observe adaptation to HRTV-DL1 infection, we established long-term infected cultures, in
triplicates, of both ACAM34_UNSW and ACAM34_DSMZ with an MOI of only 1 to allow a recovery

of infected cultures. ACAM34_DSMZ remained virus free over several dilution cycles and no new CRISPR spacers could be detected by PCR of the leader adjacent region of the CRISPR locus (Supplementary Fig. 13). For ACAM34_UNSW lysis was observed as expected between 45 and 70 hours post infection and the cultures were maintained to allow the growth of HRTV-DL1 resistant clones. The cultures slowly recovered over a time frame of app. 200 hours. After transfer of the recovered cells in fresh media, the cultures experienced another long lag phase of about 120 hours before establishing a normal growth (Supplementary Fig. 14). Cells from the recovered cultures were plated on agar plates to obtain single colonies and for each biological replicate we tested 24 colonies for infection with HRTV-DL1 by PCR. In 14 of the 72 tested colonies we could still detect the virus by PCR (Supplementary Fig. 15), indicating that the majority of the cells were able to exclude the virus.

Ten virus-free colonies were then chosen from each replicate to test re-infection with HRTV-DL1, and surprisingly, each biological replicate showed a different behaviour. Ten colonies chosen from replicate 1 could all be re-infected with HRTV-DL1, indicating that virus exclusion was only temporarily and likely due the activity of an existing virus exclusion mechanisms. All tested colonies were lysed identical to the original ACAM34_UNSW, except one colony ACAM34_UNSW_1.7 showing a reduced lysis (Supplementary Fig. 16). In contrast, none of the colonies that were isolated from replicate 2 could be re-infected with HRTV-DL1 and showed normal growth behaviour, apart from one ACAM34_UNSW_2.10 that grew faster than all the other tested colonies (Supplementary Fig. 16). Finally, colonies retrieved from replicate 3 showed a mixed behaviour. 9 out of 10 could be re-infected with the virus, however, for 8 out of the 9 the PCR signal for HRTV-DL1 was significantly weaker than in the control and no lysis could be observed in any of the cultures (Supplementary Fig. 16). We conclude that the majority of escape mutants retrieved from replicate 2 and 3, in contrast to replicate 1, had undergone some genetic changes to exclude HRTV-DL1.

Out of each replicate we chose one particularly interesting colony for further characterisation. ACAM34_UNSW_1.7 for showing reduced lysis, ACAM34_UNSW_2.14 as a representative for a fully resistant clone and ACAM34_UNSW_3.3 for showing a reduced infection without cell lysis.

Characterisation of virus escape mutants shed light on virus-host interactions and reveal additional virus exclusion mechanisms

In-depth characterization of virus escape mutants (details presented in Supplementary Results and Discussion), including genome sequencing, revealed further insights into the interactions between HRTV-DL1 and ACAM34_UNSW. A graphic summary of the results is presented in Figure 6.

ACAM34_UNSW_1.7 is partially resistant to HRTV-DL1 infection, both HRTV-DL1 adsorption and virus particle production are significantly reduced (Supplementary Fig. 17 and 18). Genome sequencing uncovered that a variant of HRTV-DL1 (HRTV-DL_1.7), that is incapable of independent replication and virus particle production, is integrated into the UNSW_1.7 genome. We infer that HRTV-DL_1.7 is responsible for the partial resistance of UNSW_1.7 to HRTV-DL1 infection by a superinfection exclusion mechanisms as previously described for another arTV [28]. We propose that HRTV-DL_1.7 interferes with virus adsorption by causing changes to the cell surface of the host. However, how the integrated virus impacts virus particle production remains to be elucidated.

ACAM34_UNSW_2.14 is fully resistant to HRTV-DL1 infection (Supplementary Figure 18). Infection is already impaired at the adsorption stage (Supplementary Fig. 17), indicating that the receptor for HRTV-DL1 might have undergone changes in this mutant. Genome sequencing revealed that

UNSW_2.14 experienced a large genomic rearrangement of former CHR2 that was caused by transposase activity and led to the loss of about 200 kb. The deletion includes one of two S-layer genes (Hlac_2976) encoded by *Hrr. lacusprofundi*. While the strain is growing slightly slower than the parental strain (Supplementary Figure 18), we did not observe any morphological changes to the cells (Supplementary figure 23) and conclude that the second S-layer gene is sufficient to maintain cell morphology and stability. However, we detected also an insertion of two amino acids into the second S-layer gene (Hlac_0412) and therefore conclude that the S-layer is the primary receptor for HRTV-DL1. Variability of S-layer genes of ACAM34 was previously detected by two 'omics'-based studies of Deep Lake, and suggested to be driven by an arms race between viruses and host [5, 10].

Adsorption of HRTV-DL1 to escape mutant ACAM34_UNSW_3.3 is greatly reduced, but not fully impaired (Supplementary Fig. 17). While the virus genome can be detected in host cells, no cell lysis can be observed and no virus particles are detected in culture supernatants (Supplementary Fig. 18). We conclude that adsorption is partially impaired, but virus particle production and cell lysis is inhibited. The genome of UNSW_3.3 reveals an insertion of 12 amino acids into one of the S-layer genes (Hlac_2976) in 90% of the population, being the cause for the reduced adsorption and confirming the S-layer as primary receptor for HRTV-DL1. Only one other genomic change is detected in UNSW_3.3, the deletion of an origin of replication (ORI), that could possibly be responsible for preventing virus particle production and cell lysis. ORIs are binding sites for CDC6/ORC1 proteins and the deletion of an ORI could redirect CDC6 causing the same effect as the upregulation of CDC6 proteins that we detect in infected ACAM34_UNSW cells. Studies have shown that CDC6 can bind dsDNA without sequence specificity [66], that dysregulation of CDC6 expression can lead to inhibition of replication [67] and that CDC6 is also able to recruit the RNA polymerase I for rDNA transcription initiation [68]. Therefore, we suggest that CDC6 could affect HRTV-DL1 replication and transcription.

Conclusion

In this study we present an in-depth characterisation of the virus-host interactions between the archaeal head-tailed virus HRTV-DL1 and its host *Hrr. lacusprofundi* ACAM34. HRTV-DL1 was isolated from a sample taken in 2014 from the same environment (Deep Lake, Antarctica) as *Hrr. lacusprofundi* ACAM34, that was isolated in 1988. HRTV-DL1 was retrieved from a lysate (HRTV-DL) that was likely a mixture of different virus variants, and most importantly, HRTV-DL exhibited integrase genes that were almost entirely lost upon isolation of the pure variant HRTV-DL1. By using plaque assays we forced a selection for a variant with a lytic life cycle. We conclude that the virus has a high genome plasticity and our work demonstrates that working with an isolated model virus-host systems does not reflect the entire spectrum of the respective virus-host interactions as they are occurring in their natural environment.

Despite the fact that the time of isolation of virus and host is at least 26 years apart, the virus was able to generate a successful infection on ACAM34. However, the ability of HRTV-DL1 to infect the laboratory strain (ACAM34_UNSW), in contrast to its inability to complete a full lytic cycle in the same strain obtained from the DSMZ (ACAM34_DSMZ), uncovered that the laboratory strain had undergone a massive genomic rearrangement and lost an entire megaplasmid, demonstrating genome plasticity also for the host and demonstrating that the lost genomic information is not essential when grown under laboratory conditions. ACAM34_UNSW lost a number of virus defence mechanisms including a CRISPR-Cas system, upon growth under non-challenging laboratory conditions. This is a great example to demonstrate, that resequencing laboratory strains should be a good common practice.

We were not able to determine with certainty the mechanisms that is responsible for HRTV-DL1
exclusion from ACAM34_DSMZ. However, having both strains in hand, allowed us to uncover a
number of new putative virus exclusion mechanism, that were only activated in ACAM34_UNSW in
absence of other defence mechanisms. Analysis of the HRTV-DL1 infected transcriptome revealed the
activity of a TnpB/Cas14-like nuclease and a transposon passenger gene that could potentially prevent
the virus from attaching to the host cell. Additionally, we propose that a predicted SMC protein, in
complex with at least two other proteins, might represent a derivative of the bacterial Wadjet system.
Isolation of escape mutants, uncovered that one of two S-layer protein is the preferred virus receptor,
indicating that exhibiting two S-layer proteins could be a strategy of virus exclusion. HRTV-DL1 can
provide partial superinfection exclusion when integrated into the host genome. Additionally, we
propose that the origin binding protein CDC6 can be involved in virus exclusion by a so far unknown
mechanisms. We observed that both the host and the virus actively undergo genomic rearrangements
when interacting with each other, allowing both host an virus to adapt to each other.

Our study uncovers numerous virus exclusion mechanisms within one single organisms, that have likely
been acquired over a very long time by a continuously ongoing arms race between viruses and *Hrr.*
lacusprofundi. The establishment and characterization of this virus-host system will allow to observe
this ongoing race both in the laboratory and in the environment for which metagenomics data are
available. The new virus exclusion mechanisms discovered in this study can be investigated in detail
and possibly be identified in other organisms. ACAM34_UNSW itself can be used as model to study
virus exclusion mechanisms. It is susceptible to a variety of viruses, including HRTV-DL1 and others
[8, 12, 16, 69, 70], and virus exclusion mechanisms can be re-introduced [71] to test their activity
against diverse viruses.

Data availability

The genomes of HRTV-DL, HRTV-DL1 and HRTV-DL1_1.7 are available on NCBI under the with
accession numbers OP630574, OP630575, OP630576. RNA sequencing raw data were submitted to
ENA-EMBL under project number: PRJEB56294. Raw data from genome sequencing of UNSW_1.7,
UNSW_2.14 and UNSW_3.3 are available in the Sequence Read Archive under BioProject ID:
PRJNA887929. The genome of ACAM34_UNSW is associated with the manuscript as supplementary
data file.
829

830
Acknowledgments

We thank the Max Planck-Genome-Centre Cologne (Cologne, Germany) for assistance with DNA and
RNA sequencing. We thank Ingrid Kunze (MPI for Marine Microbiology, Bremen, Germany) for
assistance with some of the experiments. We thank Tomas Alarcón-Schumacher and Bernhard
Tschitschko for help with some of the figures. We thank the entire Archaeal Virology group at the Max-
Planck-Institute for Marine Microbiology and Friedhelm Pfeiffer for critically reviewing the
manuscript. We also acknowledge Friedhelm Pfeiffer and Mike Dyall-Smith for identifying the PATE
and helpful discussions. Part of this work was supported by the Australian Research Council
[DP170102576]. Mass spectrometry results were obtained at the Bioanalytical Mass Spectrometry
Facility (BMSF) within the Analytical Centre of the University of New South Wales. Subsidized access

to the BMSF is gratefully acknowledged. Finally, we want to thank the Max-Planck-Institute for Marine Microbiology and the Max-Planck-Society for continuous support.

References

1. Gibson, J.A.E., *The meromictic lakes and stratified marine basins of the Vestfold Hills, East Antarctica.* Antarctic Science, 1999. **11**(2): p. 175-192.
2. Bowman, J.P., et al., *The microbial composition of three limnologically disparate hypersaline Antarctic lakes.* FEMS Microbiology Letters, 2000. **183**(1): p. 81-88.
3. DeMaere, M.Z., et al., *High level of intergenera gene exchange shapes the evolution of haloarchaea in an isolated Antarctic lake.* Proc Natl Acad Sci U S A, 2013. **110**(42): p. 16939-44.
4. Ausiannikava, D., et al., *Evolution of Genome Architecture in Archaea: Spontaneous Generation of a New Chromosome in Haloferax volcanii.* Mol Biol Evol, 2018. **35**(8): p. 1855-1868.
5. Tschitschko, B., et al., *Genomic variation and biogeography of Antarctic haloarchaea.* Microbiome, 2018. **6**(1): p. 113.
6. Wang, H., et al., *Archaeal extrachromosomal genetic elements.* Microbiol Mol Biol Rev, 2015. **79**(1): p. 117-52.
7. Hawkins, M., et al., *Accelerated growth in the absence of DNA replication origins.* Nature, 2013. **503**(7477): p. 544-547.
8. Erdmann, S., et al., *A plasmid from an Antarctic haloarchaeon uses specialized membrane vesicles to disseminate and infect plasmid-free cells.* Nature Microbiology, 2017. **2**(10): p. 1446-1455.
9. Rohwer, F., D. Prangishvili, and D. Lindell, *Roles of viruses in the environment.* Environmental Microbiology, 2009. **11**(11): p. 2771-2774.
10. Tschitschko, B., et al., *Antarctic archaea–virus interactions: metaproteome-led analysis of invasion, evasion and adaptation.* The ISME Journal, 2015. **9**(9): p. 2094-2107.
11. Luk, A.W., et al., *Viruses of haloarchaea.* Life (Basel), 2014. **4**(4): p. 681-715.
12. Alarcón-Schumacher, T., et al., *Isolation of a virus causing a chronic infection in the archaeal model organism Haloferax volcanii reveals antiviral activities of a provirus.* Proc Natl Acad Sci U S A, 2022. **119**(35): p. e2205037119.
13. Liu, Y., et al., *Diversity, taxonomy, and evolution of archaeal viruses of the class Caudoviricetes.* PLOS Biology, 2021. **19**(11): p. e3001442.
14. Bernheim, A. and R. Sorek, *The pan-immune system of bacteria: antiviral defence as a community resource.* Nature Reviews Microbiology, 2020. **18**(2): p. 113-119.
15. Takahashi, S., et al., *Development of a prokaryotic universal primer for simultaneous analysis of Bacteria and Archaea using next-generation sequencing.* PLoS One, 2014. **9**(8): p. e105592.
16. Dyall-Smith, M., *The Halohandbook: Protocols for Halobacterial Genetics. .* https://haloarchaea.com/wp-content/uploads/2018/10/Halohandbook_2009_v7.3mds.pdf, 2021.
17. Kolmogorov, M., et al., *Assembly of long, error-prone reads using repeat graphs.* Nat Biotechnol, 2019. **37**(5): p. 540-546.
18. Bankevich, A., et al., *SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing.* J Comput Biol, 2012. **19**(5): p. 455-77.
19. Antipov, D., et al., *MetaviralSPAdes: assembly of viruses from metagenomic data.* Bioinformatics, 2020. **36**(14): p. 4126-4129.

20. Seemann, T., *Prokka: rapid prokaryotic genome annotation.* Bioinformatics, 2014. **30**(14): p.
2068-9.
21. Blum, M., et al., *The InterPro protein families and domains database: 20 years on.* Nucleic
Acids Research, 2020. **49**(D1): p. D344-D354.
22. Söding, J., *Protein homology detection by HMM–HMM comparison.* Bioinformatics, 2004.
21(7): p. 951-960.
23. Carver, T.J., et al., *ACT: the Artemis comparison tool.* Bioinformatics, 2005. **21**(16): p. 3422-
3423.
24. Garneau, J.R., et al., *PhageTerm: a tool for fast and accurate determination of phage termini
and packaging mechanism using next-generation sequencing data.* Scientific Reports, 2017.
7(1): p. 8292.
25. Cutler, R.G. and J.E. Evans, *Synchronization of bacteria by a stationary-phase method.* J
Bacteriol, 1966. **91**(2): p. 469-76.
26. Grissa, I., G. Vergnaud, and C. Pourcel, *CRISPRFinder: a web tool to identify clustered
regularly interspaced short palindromic repeats.* Nucleic Acids Res, 2007. **35**(Web Server
issue): p. W52-7.
27. Franzmann, P.D., et al., *Halobacterium lacusprofundi sp. nov., a Halophilic Bacterium
Isolated from Deep Lake, Antarctica.* Systematic and Applied Microbiology, 1988. **11**(1): p.
20-27.
28. Schnabel, H., et al., *Halobacterium halobium phage øH.* The EMBO Journal, 1982. **1**(1): p. 87-
92.
29. Chen, I.-M.A., et al., *The IMG/M data management and analysis system v.6.0: new tools and
advanced capabilities.* Nucleic Acids Research, 2020. **49**(D1): p. D751-D763.
30. Fraser, J.S., et al., *Ig-like domains on bacteriophages: a tale of promiscuity and deceit.* J Mol
Biol, 2006. **359**(2): p. 496-507.
31. Pell, L.G., et al., *The solution structure of the C-terminal Ig-like domain of the bacteriophage
λ tail tube protein.* J Mol Biol, 2010. **403**(3): p. 468-79.
32. Kageyama, Y., et al., *Observation of the membrane binding activity and domain structure of
gpV, which comprises the tail spike of bacteriophage P2.* Biochemistry, 2009. **48**(42): p.
10129-35.
33. von Itzstein, M., *The war against influenza: discovery and development of sialidase
inhibitors.* Nature Reviews Drug Discovery, 2007. **6**(12): p. 967-974.
34. Quemin, E.R.J. and T.E.F. Quax, *Archaeal viruses at the cell envelope: entry and egress.*
Frontiers in microbiology, 2015. **6**: p. 552-552.
35. Young, R., *Phage lysis: do we have the hole story yet?* Curr Opin Microbiol, 2013. **16**(6): p.
790-7.
36. Tamir, A. and J. Eichler, *N-Glycosylation Is Important for Proper Haloferax volcanii S-Layer
Stability and Function.* Appl Environ Microbiol, 2017. **83**(6).
37. Saier, M.H., Jr. and B.L. Reddy, *Holins in bacteria, eukaryotes, and archaea: multifunctional
xenologues with potential biotechnological and biomedical applications.* J Bacteriol, 2015.
197(1): p. 7-17.
38. Jumper, J., et al., *Highly accurate protein structure prediction with AlphaFold.* Nature, 2021.
596(7873): p. 583-589.
39. Krieger, I.V., et al., *The Structural Basis of T4 Phage Lysis Control: DNA as the Signal for Lysis
Inhibition.* Journal of Molecular Biology, 2020. **432**(16): p. 4623-4636.
40. Pagaling, E., et al., *Sequence analysis of an Archaeal virus isolated from a hypersaline lake in
Inner Mongolia, China.* BMC Genomics, 2007. **8**: p. 410.
41. Kukkaro, P. and D.H. Bamford, *Virus-host interactions in environments with a wide range of
ionic strengths.* Environ Microbiol Rep, 2009. **1**(1): p. 71-7.
42. Liu, J., et al., *Virus-induced cell gigantism and asymmetric cell division in archaea.*
Proceedings of the National Academy of Sciences, 2021. **118**(15): p. e2022578118.

43. Siguier, P., et al., *ISfinder: the reference centre for bacterial insertion sequences.* Nucleic
Acids Res, 2006. **34**(Database issue): p. D32-6.
44. Pfeiffer, F., et al., *Evolution in the laboratory: The genome of Halobacterium salinarum strain*
R1 compared to that of strain NRC-1. Genomics, 2008. **91**(4): p. 335-346.
45. Willkomm, S., K.S. Makarova, and D. Grohmann, *DNA silencing by prokaryotic Argonaute*
proteins adds a new layer of defense against invading nucleic acids. FEMS Microbiol Rev,
2018. **42**(3): p. 376-387.
46. Lodeyro, A.F., et al., *The importance of flavodoxin for environmental stress tolerance in*
photosynthetic microorganisms and transgenic plants. Mechanism, evolution and
biotechnological potential. FEBS Letters, 2012. **586**(18): p. 2917-2924.
47. Kerscher, L. and D. Oesterhelt, *Ferredoxin is the coenzyme of alpha-ketoacid oxidoreductases*
in Halobacterium halobium. FEBS Lett, 1977. **83**(2): p. 197-201.
48. Quax, T.E., et al., *Massive activation of archaeal defense genes during viral infection.* J Virol,
2013. **87**(15): p. 8419-28.
49. Macchietto, M.G., R.A. Langlois, and S.S. Shen, *Virus-induced transposable element*
expression up-regulation in human and mouse host cells. Life Sci Alliance, 2020. **3**(2).
50. Broecker, F. and K. Moelling, *Evolution of Immune Systems From Viruses and Transposable*
Elements. Front Microbiol, 2019. **10**: p. 51.
51. Karlsson, F., et al., *Genome-wide comparison of phage M13-infected vs. uninfected*
Escherichia coli. Can J Microbiol, 2005. **51**(1): p. 29-35.
52. Howard-Varona, C., et al., *Phage-specific metabolic reprogramming of virocells.* Isme j, 2020.
14(4): p. 881-895.
53. Payne, L.J., et al., *PADLOC: a web server for the identification of antiviral defence systems in*
microbial genomes. Nucleic Acids Research, 2022.
54. Koretke, K.K., et al., *Evolution of Two-Component Signal Transduction.* Molecular Biology and
Evolution, 2000. **17**(12): p. 1956-1970.
55. Zeng, Q. and S.W. Chisholm, *Marine viruses exploit their host's two-component regulatory*
system in response to resource limitation. Curr Biol, 2012. **22**(2): p. 124-8.
56. Harrington, L.B., et al., *Programmed DNA destruction by miniature CRISPR-Cas14 enzymes.*
Science, 2018. **362**(6416): p. 839-842.
57. Karvelis, T., et al., *Transposon-associated TnpB is a programmable RNA-guided DNA*
endonuclease. Nature, 2021. **599**(7886): p. 692-696.
58. Gibson, R.T. and E.J. Androphy, *The SMC5/6 Complex Represses the Replicative Program of*
High-Risk Human Papillomavirus Type 31. Pathogens, 2020. **9**(10).
59. Han, C., et al., *KSHV RTA antagonizes SMC5/6 complex-induced viral chromatin compaction*
by hijacking the ubiquitin-proteasome system. PLOS Pathogens, 2022. **18**(8): p. e1010744.
60. Doron, S., et al., *Systematic discovery of antiphage defense systems in the microbial*
pangenome. Science, 2018. **359**(6379).
61. Liu, H.W., et al., *DNA-measuring Wadjet SMC ATPases restrict smaller circular plasmids by*
DNA cleavage. bioRxiv, 2022: p. 2022.10.04.510915.
62. Babakhani, S. and M. Oloomi, *Transposons: the agents of antibiotic resistance in bacteria.* J
Basic Microbiol, 2018. **58**(11): p. 905-917.
63. Abdul Halim, M.F., et al., *Haloferax volcanii archaeosortase is required for motility, mating,*
and C-terminal processing of the S-layer glycoprotein. Molecular Microbiology, 2013. **88**(6):
p. 1164-1175.
64. Achtman, M., N. Kennedy, and R. Skurray, *Cell--cell interactions in conjugating Escherichia*
coli: role of traT protein in surface exclusion. Proc Natl Acad Sci U S A, 1977. **74**(11): p. 5104-
8.
65. Gomes-Filho, J.V., et al., *Sense overlapping transcripts in IS1341-type transposase genes are*
functional non-coding RNAs in archaea. RNA Biol, 2015. **12**(5): p. 490-500.

66. Feng, L., et al., *Identification and Characterization of Saccharomyces cerevisiae Cdc6 DNA-*
binding Properties. Molecular Biology of the Cell, 2000. **11**(5): p. 1673-1685.
67. Kundu, L.R., et al., *Deregulated Cdc6 inhibits DNA replication and suppresses Cdc7-mediated*
phosphorylation of Mcm2-7 complex. Nucleic Acids Res, 2010. **38**(16): p. 5409-18.
68. Huang, S., et al., *DNA replication initiator Cdc6 also regulates ribosomal DNA transcription*
initiation. J Cell Sci, 2016. **129**(7): p. 1429-40.
69. Porter, K., et al., *SH1: A novel, spherical halovirus isolated from an Australian hypersaline*
lake. Virology, 2005. **335**(1): p. 22-33.
70. Nuttall, S.D. and M.L. Dyall-Smith, *HF1 and HF2: novel bacteriophages of halophilic archaea.*
Virology, 1993. **197**(2): p. 678-84.
71. Liao, Y., et al., *Developing a genetic manipulation system for the Antarctic archaeon,*
Halorubrum lacusprofundi: investigating acetamidase gene function. Scientific Reports,
2016. **6**(1): p. 34639.
72. Gilchrist, C.L.M. and Y.H. Chooi, *Clinker & clustermap.js: Automatic generation of gene*
cluster comparison figures. Bioinformatics, 2021.
73. Kurtz, S., et al., *Versatile and open software for comparing large genomes.* Genome Biology,
2004. **5**(2): p. R12.
74. Krzywinski, M., et al., *Circos: an information aesthetic for comparative genomics.* Genome
Res, 2009. **19**(9): p. 1639-45.
75. Nagel, C., et al., *Several One-Domain Zinc Finger μ-Proteins of Haloferax Volcanii Are*
Important for Stress Adaptation, Biofilm Formation, and Swarming. Genes (Basel), 2019.
10(5).
76. Esquivel, R.N., R. Xu, and M. Pohlschroder, *Novel Archaeal Adhesion Pilins with a Conserved*
N Terminus. Journal of Bacteriology, 2013. **195**(17): p. 3808-3818.
77. Nußbaum, P., et al., *An Oscillating MinD Protein Determines the Cellular Positioning of the*
Motility Machinery in Archaea. Curr Biol, 2020. **30**(24): p. 4956-4972.e4.
78. Lutkenhaus, J., *Assembly Dynamics of the Bacterial MinCDE System and Spatial Regulation of*
the Z Ring. Annual Review of Biochemistry, 2007. **76**(1): p. 539-562.

1039 **Figures and Tables**

1040

1041

Figure 1. Nucleic acid content and particle of HRTV-DL. (A) Total DNA of HRTV-DL undigested
(-) and digested with *Eco*RV (+EcoRV). MW size marker is shown to the left of the gel (GeneRuler 1
kb Plus DNA Ladder, Thermo Fisher Scientific). DNA was separated on 1% agarose gels and stained
with SYBR™ Safe DNA stain. **(B)** Electron micrograph of HRTV-DL particle. Sample was negatively
stained with 2% uranyl acetate. Size bar: 50nm. Original images have been modified by cropping to
improve visual presentation.

1048

1049

1050

1051

Figure 2: Comparisons of genomes HRTV-DL1 with BJ1 and CGphi46 belonging to the *Flexireviridae.* Protein functions are indicated above or below the corresponding ORFs. Genes encoding virus morphogenesis–related proteins are colored in green, whereas replication-related genes are colored in red. Homologous genes shared between viruses are connected by shadings of different degrees of blue based on the amino acid sequence identity. Figure was generated with clinker & clustermap.js [72].

1066

Figure 3: Life cycle of HRTV-DL1. Growth curve of uninfected control and HRTV-DL1 infected *Hrr. lacusprofundi* ACAM34_UNSW. Free virus particles in 10ul culture supernatant were determined by plaque assay. **Inlet:** Virus-host ratio determined by comparing copy numbers of host main chromosome and virus genome. Copy numbers were determined per ml cell culture by qPCR and samples were taken at time points encircled in the growth curve in the main figure. Graphs represent one of three biological replicates.

1080

1081

Figure 4: Genome comparison of ACAM34_UNSW with the three replicons CHR1, CHR2 and pHLAC01 (NC_012029.1, NC_012028.1 and NC_012030.1) of the reference genome from NCBI (ATCC 49239). Interconnecting lines highlight regions present in both genomes (red/blue colors indicates same/reverse orientation on the two genomes). The figure highlights that the ACAM34_UNSW genome consists of a single replicon comprising the full length primary replicon (NC_012029.1) and parts of one secondary replicon (NC_012028.1) of the reference genome, while pHLAC01 (NC_012030.1) of the reference genome is lost. Genomic regions shared between the two genomes were identified using NUCmer [73] and visualized with Circos [74] wrapped within the script mummer2circos.py (https://github.com/metagenlab/mummer2circos).

1091

1092

1093

Figure 5. Functional profile of differentially expressed genes in ACAM34_DSMZ and ACAM34_UNSW under infection with HRTV-DL1. Bars represent the number of differentially expressed genes assigned to each particular functional category at a given time point (T1 = 32 hours post infection, T2 = 56 hours post infection). Functional classification of the *Hrr. lacusprofundi* genomes was performed using the cluster of orthologous genes (COG) database. The colors of the bars indicate if genes are upregulated (red) or downregulated (blue) respectively. Genes that were not assigned to any functional category are not displayed on the plot (for detailed information see Supplementary Table 7 and 9).

1102

1103

Figure 6. Summary of interactions observed between *Hrr. lacusprofundi* ACAM34 and HRTV-DL1. *Hrr. lacusprofundi* ACAM34_DSMZ allows adsorption and genome injection of HRTV-DL1. However, The genome is either degraded immediately after injection or replication is inhibited by one or several virus exclusion mechanisms present in the strain. ACAM34_DSMZ lost pHLAC01 and a portion of CHR2, while the remaining CHR2 is integrated into CHR1. All virus exclusion mechanisms potentially active against HRTV-DL1 in ACAM34_DSMZ are lost. However, other virus exclusion mechanism are activated allowing the emergence of escape mutants. Escape mutant UNSW_1.7 includes an integrated HRTV-DL1 variant, that provides superinfection exclusion. Escape mutant UNSW_2.14 lost one of two S-layer genes while the second S-layer gene experienced an insertion of two amino acids. This leads to a change of the cell surface and prevents HRTV-DL1 adsorption. Escape mutant UNSW_3.3 experienced an insertion of 12 amino acids into one of the two S-layer genes in 90% of the population, preventing adsorption of HRTV-DL1 to the majority of cells. The deletion of an origin of replication (ORI) and the subsequent repurposing of CDC6 is hypothesized to prevent virus particle production by a so far unknown mechanisms. Figure created with BioRender.com.

CDS number ACAM34_UNSW	CDS number ACAM34_ DSMZ	Differential Expression Log2 Ratio	conserved domains/structural homologies (HMM search)	Predicted function
ACAM34UNSW_ 00279	Hlac_0269	2.42	Small zinc finger protein	Stress response [75]
ACAM34UNSW_ 00405	Hlac_0390	1.98	Archaeal type IV pilus N-terminal domain	Inducing biofilm formation to protect against viral infection. [76]
ACAM34UNSW_ 03039	Hlac_2674	1.94	MinD-like	Enhancing motility or inhibiting cell division [77, 78]
ACAM34UNSW_ 00280	Hlac_0270	1.90	small hypothetical protein	Same operon as Hlac_0269 (above). No functional prediction.
ACAM34UNSW_ 03040	Hlac_2675	1.77	coil	Same operon as Hlac_2674 (MinD-like). No functional prediction.
ACAM34UNSW_ 00927	Not annotated	1.67	small hypothetical protein	No functional prediction.
ACAM34UNSW_ 00941	Hlac_0922	1.56	hypothetical	Within upregulated region that includes an integrase. Mobile genetic element.
ACAM34UNSW_ 01637	Hlac_0148	1.53	Iron-sulphur (FeS) cluster	Oxidative stress response
ACAM34UNSW_ 01620	Hlac_1593	1.48	Small hypothetical protein	Upstream of Hlac_1592 (log2=1.1), predicted SMC (structural maintenance of chromosomes). Possible functionally related to SMC and Hlac_1591 (see below ACM34UNSW_01618). Protecting host DNA or inhibiting viral replication [58, 59] , possibly similar to the Wadjet system [60].
ACAM34UNSW_ 00939	Hlac_0920	1.47	Sensor protein, two-component system	Signal transduction (with Hlac_0919). Within upregulated region that includes an integrase. Mobile genetic element.

ACAM34UNSW_00938	Hlac_0919	1.47	Sensor kinase, two-component system	Signal transduction (with Hlac_0920). Within upregulated region that includes an integrase. Mobile genetic element.
ACAM34UNSW_01618	Hlac_1591	1.43	hypothetical	Downstream of Hlac_1592 (log2=1.1), predicted SMC and Hlac_1593 (see above). Possible functionally related to SMC and protecting host DNA or inhibiting viral replication [58, 59] , possibly similar to the Wadjet system [60].
ACAM34UNSW_01210	Hlac_1190	1.42	ISH9-type transposase (ISHla3)	Transposase
ACAM34UNSW_00937	Hlac_0918	1.42	Small hypothetical protein	Within upregulated region that includes an integrase. Mobile genetic element.
ACAM34UNSW_01107	Hlac_1086	1.40	CRISPR-associated protein Cas14a (99.95% probability)	RNA-guided DNA endonuclease [56],[57]. Degradation of virus genome intermediates.
ACAM34UNSW_01789	Hlac_2783	1.38	KTSC domain, RNA binding	No functional prediction.
ACAM34UNSW_01086	Hlac_1065	1.37	Glycosyl transferase	Cell surface modification.
ACAM34UNSW_01670	Hlac_1641	1.35	Small hypothetical protein, DNA-binding response regulator (Probability 87%, E-value 1.6)	Between to Hlac_1640 (log2= 1.26) and Hlac_1642 (log2= 1.16), both homing endonucleases. Mobile genetic element.
ACAM34UNSW_00935	Hlac_0916	1.33	Resolvase, site specific recombinase	Integrase within upregulated region. Mobile genetic element.
ACAM34UNSW_01692	Hlac_1663	1.33	Small hypothetical protein	No functional prediction.
ACAM34UNSW_00942	Hlac_0923	1.31	IS6 family transposase	End of upregulated region that includes an integrase. Mobile genetic element.
ACAM34UNSW_02065	Hlac_3065	1.30 (normalized by 1.3)	Chromosome-partitioning protein ParB (95% probability)	On former CHR2, possibly activated upon re-mobilization of CHR2
ACAM34UNSW_02231	Hlac_1874	1.27	Transmembrane protein	No functional prediction.

ACAM34UNSW_00934	Hlac_0915	1.26	creatinine amidohydrolase	Transport and metabolism, Secondary metabolites biosynthesis. Within upregulated region that includes an integrase. Mobile genetic element.
ACAM34UNSW_01217	Hlac_1197	1.26	HD domain, putative phosphohydrolase	Nucleic acid metabolism, signal transduction
ACAM34UNSW_01669	Hlac_1640	1.26	Homing endonuclease	Adjacent to Hlac_1641 (above) and Hlac_1642 (log2= 1.16), another homing endonucleases. Mobile genetic element.
ACAM34_UNSW_02089	Hlac_3092	1.26 (normalized by 1.3)	YTH (YT521-B homology) domain	Stabilizing mRNA.
ACAM34UNSW_00936	Hlac_0917	1.26	Small hypothetical protein	Within upregulated region that includes an integrase. Mobile genetic element.
ACAM34UNSW_01549	Hlac_1524	1.25	orc1/cdc6 family replication initiation protein	Virus exclusion (see paragraph '*Characterisation of virus escape mutants shed light on virus-host interactions and reveal additional virus exclusion mechanisms*')
ACAM34_UNSW_01787	Hlac_1757	1.24	ISH3-type transposase	Transposase

Table 1: Top 30 of upregulated genes in ACAM34_UNSW upon HRTV-DL1 infection.