Title: D1R^CaMKII in claustrum mediates ACE-induced anxiety and electro-

acupuncture therapy

Authors: Liying Chen[1,2,5], Zhaoyu Liu[1,5], Ziheng Zhao[1,5], Hao Guo[1], Demin Du[1],

Weichao Pan[1], Xiaoyan Wei[1], Jiaxun Nie[1], Feifei Ge[1], Jianhua Ding[3], Yu Fan[1,*],

Hee Young Kim[4,*], Xiaowei Guan[1,*]

Affiliations:

[1]Department of Human Anatomy and Histoembryology, Nanjing University of
Chinese Medicine, Nanjing, China

[2]Department of Critical Care Medicine, Jiangxi Province Hospital of Integrated
Chinese and Western Medicine, Nanchang, China

[3]Department of Pharmacology, Nanjing Medical University, Nanjing, China

[4]Department of Physiology, Yonsei University College of Medicine, Seoul,
South Korea

[5]These authors equally contribute to this work

Running title: Claustrum D1R^CaMKII in ACE anxiety and EA

* Correspondence authors: guanxw918@njucm.edu.cn (XG),
yfan@njucm.edu.cn (YF), vet202001@yuhs.ac (HYK)

Phone number: +86-25-85811387

Abstract

Adolescent cocaine exposure (ACE) increases risk of developing psychiatric problems such as anxiety in their later life. Here, ACE animal models were established by exposing to cocaine during adolescent period in male mice, followed by electro-acupuncture (EA) at acupoints of Baihui (GV 20) and Yintang (GV 29) with stimuli of a mixed frequency of 2/100 Hz for 28 withdrawal days. ACE mice of adulthood exhibited anxiety-like behaviors, accompanying with obviously activated CaMKII-positive neurons and increased dopamine receptor 1 (D1R) on CaMKII-positive neurons (D1R^{CaMKII}) in claustrum. *In vivo*, local blocking D1R with SCH23390 or knocking-down of D1R^{CaMKII} with specific virus efficiently reduced claustrum activation, and ultimately suppressed anxiety in ACE mice of adulthood. EA treatment alleviated ACE-induced anxiety by suppressing D1R^{CaMKII} and claustrum activation. Our findings identified a novel role of claustrum in ACE-induced anxiety, and put new insight into the function of claustral D1R^{CaMKII}. Targeting claustral D1R^{CaMKII}, EA obviously alleviates ACE-induced anxiety in mice of adulthood.

Introduction:

Adolescents present higher vulnerability to substance abuse (Chambers, Taylor, & Potenza, 2003; Laviola, Adriani, Terranova, & Gerra, 1999). Adolescent experience of drugs increase the risk of developing psychiatric disorders that even persists into adulthood (Moss, Chen, & Yi, 2014; Rutherford, Mayes, & Potenza, 2010). Anxiety is a prominent psychiatric symptom of adolescent drug abuse, which is generally believed as critical factor for driving relapse to drugs in later life (Erb, 2010; Stopponi et al., 2014). Previously, we showed that adolescent cocaine-exposed (ACE) mice exhibit more anxiety in their adulthood (Shi et al., 2019; Zhu et al., 2018), however, the involved brain region remains much to be clarified.

The claustrum is a thin stripped nucleus located between insular cortex and putamen, being architected in anterior-posterior and dorsal-ventral organization. Above 90% neurons in claustrum are glutamatergic (CaMKII-positive) neurons (Druga, Smythies, Edelstein, & Ramachandran, 2014; Hinova-Palova et al., 2014). Anatomically, the claustrum has extensive reciprocal connectivity with all neocortical regions (Wang et al., 2017) and communicates densely with limbic structure and thalamus (Brennan et al., 2021; Markowitsch, Irle, Bang-Olsen, & Flindt-Egebak, 1984), implying that claustrum may be a hub for integrating and relaying "executive process" of cortex. At present, the functions of claustrum remain poorly understood. Emerging studies demonstrate that claustrum may be key nucleus involved in consciousness (Stiefel, Merrifield, & Holcombe, 2014), attention (Bayat et al., 2018) and cognition (Madden et al., 2022). The claustrum has been implicated in numerous brain disorders, such as seizures (Kurada, Bayat, Joshi, & Koubeissi, 2019), autism (Wegiel et al.,

2014) and Parkinson Disease (Kalaitzakis, 2014), but it has not been evidently

as the core structure of any specific disease. In marijuana addicts (Acheson et

al., 2015; Yanes et al., 2018) or internet game addicts (Zarrindast & Khakpai,

2015), the claustrum shows greater activity. Recently, Terem et al. (Terem et al.,

2020) found that the D1 dopamine receptor (D1R)-positive neurons in

claustrum is required for acquisition of cocaine-preferred behaviors, and

stimulating these neurons could drive context-reinstatement of cocaine

preference. In addition, the other recent study shows that the claustrum

mediates stress-induced anxiety in mice (Niu et al., 2022). These studies

demonstrate that claustrum is involved in both addiction and anxiety, but its role

in drugs-related anxiety is unknown.

Currently, there is few pharmacotherapy options for adolescent substance

use-induced anxiety, partially due to serious side-effects caused by anxiolytic

medicine. As a non-invasive therapeutic approach, acupuncture has been

considered as an efficient intervention on substance use, especially for

treatment of the protract abstinence syndrome (such as anxiety) (Chae et al.,

2008; Dong Hyeon Kim et al., 2018), but its pharmacological targets remain

largely unclear. The electro-acupuncture (EA), a kind of peripheral acupuncture

with electric stimulation, has been shown to improve anxiety without adverse

effects (Amorim et al., 2018; Zeng, Tao, Hou, Zong, & Yu, 2018). Previously, we

found that EA at acupoints of Baihui (GV 20) and Yintang (GV 29) effectively

reduced the anxiety in ACE mice of adulthood (Nie et al., 2020). Early in 2006,

Shapo et al. (Shapo, Ma, & Liu, 2006) reported that acupuncture at Weizhong

points (BL40) could significantly evoke claustrum in humans. Whether

claustrum is one of key targets for EA to treat ACE-related anxiety, and its

potential pharmacological mechanism, has not been explored yet.

In the present study, male mice were exposed to cocaine during adolescent

period from postnatal day 28 (P28) to P42 (Spear, 2000), followed by

subsequent EA treatment. When growing up to adult age, the mice were

subjected to anxiety behavioral tests to explore the role of claustral D1R,

especially expressed on CaMKII-positive neurons (D1R^{CaMKII}), in ACE-induced

anxiety and EA therapy.

Results:

ACE enhances anxiety with evoked claustrum and D1R^{CaMKII} in mice of

adulthood

To examine anxiety-like behaviors following ACE, mice were subjected to

behavioral tests of EPM, OFT and MBT in their adulthood (Figure 1A). As

shown in Figure 1B, ACE mice of adulthood exhibited less time spent and less

entries into the open arms, but similar distances traveled in open arms of the

EPM apparatus than those of adolescent saline exposure (ASE) mice of

adulthood. During the OFT (Figure 1C), ACE mice spent less time in the central

area of box, while had similar entries into the central area and total distances

traveled in the OFT box than ASE controls. In the MBT, the buried beads by

ACE mice had no significant difference at each time point and in total when

compared to ASE controls (Figure 1D).

The c-Fos and CaMKII were used as markers to detect the neuronal activity

and glutamatergic neurons, respectively. The total protein level of c-Fos was

higher in ACE than in ASE mice of adulthood (Figure 1E). Both the percentage

and number of c-Fos-positive & CaMKII-positive neurons were higher in ACE

mice than those in ASE mice of adulthood (Figure 1F).

On CaMKII-positive neurons, the expression of D1R ($D1R^{CaMKII}$) was much

higher in ACE than in ASE mice of adulthood (Figure 1G). The total levels of

mRNA (Figure 1H) and protein (Figure 1I) of claustral D1R was higher, but not

that of D2R and D3R in ACE than in ASE mice of adulthood. In parallel, the

protein levels of p-$ERK_{1/2}$, ratio of p-$ERK_{1/2}$ to $ERK_{1/2}$, p-CREB, ratio of p-CREB

to CREB and BDNF were higher in ACE group mice than controls, but that of

$ERK_{1/2}$ and CREB had no significant difference between ACE and ASE mice of

adulthood (Figure 1J, 1K).

Figure 1

Figure 1. ACE enhances anxiety with evoked claustrum and D1R^{CaMKII} in

mice of adulthood. A, Experimental timeline. **B**, EPM test. Two-tailed unpaired

t test. Left: t = 2.292, df = 22, *p* = 0.0319. Middle: t = 2.120, df = 22, *p* = 0.0455.

Right: t = 1.497, df = 22, *p* = 0.1486. ASE group, n = 13; ACE group, n = 11. **C**,

OFT. Two-tailed unpaired t test. Left: t = 2.308, df = 22, *p* = 0.0308. Middle: t =

1.488, df = 22, *p* = 0.1509. Right: t = 0.008965, df = 22, *p* = 0.9929. ASE group,

n = 13; ACE group, n = 11. **D**, MBT. Left: two-way RM ANOVA, $F_{(6,132)}$ = 0.3929，

p = 0.8826. Right: two-tailed unpaired t test, t = 0.8578, df = 22, *p* = 0.4003.

ASE group, n = 13; ACE group, n = 11. **E**, Total protein levels of c-Fos. Two-

tailed unpaired t test, t = 2.811, df = 4, *p* = 0.0483. ASE group, n = 3; ACE group,

n = 3. **F**, The percentage and number of c-Fos-positive & CaMKII-positive

neurons. Two-tailed unpaired t test. Upper: t = 2.466, df = 9, *p* = 0.0358. Lower:

t = 3.106, df = 9, *p* = 0.0126. ASE group, n = 6; ACE group, n = 5. Scale bar,

300 μm/50 μm. **G,** The expression of $D1R^{CaMKII}$. Two-tailed unpaired t test, t =

3.177, df = 4, *p* = 0.0336. ASE group, n = 3; ACE group, n = 3. Scale bar, 200

μm/50 μm. **H**, The mRNA levels of *D1R, D2R and D3R*. Two-tailed unpaired t

test. D1R: t = 2.583, df = 7, *p* = 0.0363, ASE group, n = 6; ACE group, n = 3.

D2R: t = 0.2676, df = 9, *p* = 0.7950, ASE group, n = 5; ACE group, n = 6. D3R:

t = 0.3458, df = 9, *p* = 0.7374, ASE group, n = 5; ACE group, n = 6. **I,** The protein

levels of D1R, D2R and D3R. Two-tailed unpaired t test. D1R: t = 7.389, df = 4,

p = 0.0018, ASE group, n = 3; ACE group, n = 3. D2R: t = 0.1621, df = 4, *p* =

0.8791. ASE group, n = 3; ACE group, n = 3. D3R: t = 1.159, df = 4, *p* = 0.2035,

ASE group, n = 3; ACE group, n = 3. **J,** Levels of p-ERK$_{1/2}$ and ERK$_{1/2}$. Two-

tailed unpaired t test. p-ERK$_{1/2}$: t = 4.905, df = 4, *p* = 0.0080. Ratio of p-ERK$_{1/2}$

to ERK$_{1/2}$: t = 3.239, df = 4, *p* = 0.0317. ERK$_{1/2}$: t = 0.9475, df = 4, *p* = 0.3970.

ASE group, n = 3; ACE group, n = 3. **K,** Levels of p-CREB, CREB and BDNF.

Two-tailed unpaired t test. p-CREB: t = 5.871, df = 4, p = 0.0042. Ratio of p-

CREB to CREB: t = 3.502, df = 4, p = 0.0248. BDNF: df = 4, t = 2.940, p =

0.0424. ASE group, n = 3; ACE group, n = 3. ASE, adolescent saline exposure;

ACE, adolescent cocaine exposure. N.S., $p > 0.05$, *, $p < 0.05$, **, $p < 0.01$ vs

ASE.

Blocking D1R within claustrum suppresses ACE-increased anxiety and

claustrum activation in mice of adulthood

To evaluate the role of claustral D1R in ACE-induced anxiety, the selective D1R

antagonist SCH-23390 was bilaterally injected into local claustrum 20 min prior

to the tests of EPM, OFT and NSFT (Figure 2A). Compared with vehicle-treated

mice, local SCH-23390 treatment significantly enhanced the time spent in open

arms, and did not affect the entries into the open arms and the distances

traveled in open arms of EPM apparatus in ACE mice of adulthood (Figure 2B).

SCH-23390 treatment had no influence in the time spent, the entries in the

central area, and the total distances traveled by ACE mice in the OFT box,

when compared with the vehicle-treated mice (Figure 2C). Similarly, there is no

difference on the latency and intake of feeding in the NSFT between vehicle-

treated and SCH-23390-treated ACE mice (Figure 2D).

As shown in Figure 2E, local SCH-23390 treatment significantly attenuated

the number of c-Fos-positive neurons, when compared with vehicle-treated

mice.

Figure 2

Figure 2. Blocking D1R within claustrum suppresses ACE-increased anxiety and claustrum activation in mice of adulthood. A, Experimental timeline and local injections of SCH-23390 into claustrum. Scale bar, 400/100 μm. **B**, EPM test. Two-tailed unpaired t test. Left: t = 2.322, df = 17, p = 0.0329. Middle: t = 1.085, df = 17, p = 0.2930. Right: t = 0.5363, df = 17, p = 0.5987. Vehicle group, n = 11; SCH-23390 group, n = 8. **C**, OFT. Two-tailed unpaired t test. Left: t = 1.688, df = 17, p = 0.2847. Middle: t = 1.105, df = 17, p = 0.2847. Right: t = 1.676, df = 17, p = 0.1121. Vehicle group, n = 11; SCH-23390 group, n = 8. **D**, NSFT. Two-tailed unpaired t test. Left: t = 0.6239, df = 17, p = 0.5410. Right: t = 0.9492, df = 17, p = 0.3558. Vehicle group, n = 11; SCH-23390 group, n = 8. **E,** The number of c-Fos-positive neurons. Two-tailed unpaired t test, t = 3.097, df = 10, p = 0.0113. Vehicle group, n = 6; SCH-23390 group, n = 6. Scale bar, 100 μm. Vehicle, vehicle-treated ACE mice; SCH-23390, SCH-23390-treated ACE mice. N.S., p > 0.05, *, p < 0.05, **, p < 0.01 vs Vehicle.

Knocking-down D1R^{CaMKII} of claustrum suppresses ACE-increased anxiety and claustrum activation in mice of adulthood

Specific knockdown of D1R^{CaMKII} was performed in ACE mice of adulthood (Figure 3A). Adeno-associated virus of either *CaMKIIap-EGFP-MIR155(shDrd1)-SV40 PolyA* (ACE$_{KD}$) or *CaMKIIap-EGFP-MIR155(MCS)-SV40 PolyA* (ACE$_{Ctrl}$) was bilaterally infused into clautrum, and about 55% CaMKII-positive neurons within clautrum were transfected with virus (Figure 3B). Compared with ACE$_{Ctrl}$ mice, the levels of D1R protein (Figure 3C) was decreased in CL of ACE$_{KD}$ mice.

ACE$_{KD}$ mice exhibited more time spent, and did not affect the entries into the

open arms, and similar distance traveled in the EPM apparatus, when

compared with ACE_Ctrl mice (Figure 3D). There were no differences on the time

spent in central area, the entries in the central area, the total distances traveled

by ACE mice in the OFT box (Figure 3E). Moreover, the latency and intake of

feeding did not change in the NSFT (Figure 3F) between ACE_KD and ACE_Ctrl

mice. In parallel, the percentage of c-Fos-positive neurons in EGFP-

transfected neurons and the mean gray value of c-Fos in EGFP-transfected

neurons were much lower in ACE_KD mice than that in ACE_Ctrl mice (Figure 3G).

Figure 3

Figure 3. Knocking-down D1R^{CaMKII} of claustrum suppresses ACE-increased anxiety and claustrum activation in mice of adulthood. A, Experimental design and timeline. **B**, Local injections of virus into claustrum. Two-tailed unpaired t test, t = 0.8496, df = 4, *p* = 0.4434. ACE$_{Ctrl}$ group, n = 3; ACE$_{KD}$ group, n = 3. Scale bar, 300 μm. **C**, The D1R protein level. Two-tailed unpaired t test, t = 3.465, df = 4, *p* = 0.0257. ACE$_{Ctrl}$ group, n = 3; ACE$_{KD}$ group, n = 3. **D**, EPM test. Two-tailed unpaired t test. Left: t = 2.152, df = 14, *p* = 0.0493. Middle: t = 1.134, df = 14, *p* = 0.2759. Right: t = 1.831, df = 14, *p* = 0.0885. ACE$_{Ctrl}$ group, n = 7; ACE$_{KD}$ group, n = 9. **E**, OFT. Two-tailed unpaired t test. Left: t = 1.500, df = 14, *p* = 0.1559. Middle: t = 0.1574, df = 14, *p* = 0.8772. Right: t = 0.1926, df = 14, *p* = 0.8500. ACE$_{Ctrl}$ group, n = 7; ACE$_{KD}$ group, n = 9. **F**, NSFT. Two-tailed unpaired t test. Left: t = 0.02965, df = 14, *p* = 0.9768. Right: t = 1.088, df = 14, *p* = 0.2950. ACE$_{Ctrl}$ group, n = 7; ACE$_{KD}$ group, n = 9. **G**, The

percentage of c-Fos-positive neurons in EGFP- transfected neurons and the

mean gray value of c-Fos in EGFP-transfected neurons. Two-tailed unpaired t

test. Upper: t = 3.207, df = 4, p = 0.0327. Lower: t = 3.006, df = 4, p = 0.0397.

ACE_{Ctrl} group, n = 3; ACE_{KD} group, n = 3. Scale bar, 400 μm/50 μm. ACE,

adolescent cocaine exposure; ACE_{Ctrl}, ACE mice injected with control virus;

ACE_{KD}, ACE mice injected with knockdown virus. N.S., $p > 0.05$, *, $p < 0.05$, **,

$p < 0.01$ vs ACE_{Ctrl}.

EA treatment rescues ACE-increased anxiety in mice of adulthood by suppressing claustrum activation and $D1R^{CaMKII}$

EA treatments at acupoints of GV20 and GV29 during P42-P70 were carried

out in ACE mice (ACE_{EA}), and placebo acupoints were used as sham EA

treatment (ACE_{SEA}) (Figure 4A, 4B). Compared with ACE_{SEA} mice, ACE_{EA} mice

spent more time, similar entries into the open arms and similar distances

traveled in open arms of the EPM apparatus (Figure 4C). There were no

differences in the time spent, the entries into the central area and the total

distances traveled in the OFT box between ACE_{EA} and ACE_{SEA} mice (Figure

4D). In the NSFT (Figure 4E), ACE_{EA} mice intake more food, and spent similar

latency of feeding, when compared with ACE_{SEA} mice.

Compared with ACE_{SEA} mice, both the percentage and the number of c-Fos-

positive & CaMKII-positive neurons were much lower in ACE_{EA} mice (Figure 4F).

In parallel, both levels of total D1R protein in CL (Figure 4G) and $D1R^{CaMKII}$

staining (Figure 4H) were lower in ACE_{EA} mice than that in ACE_{SEA} mice.

As shown in Figure 4I-J, there were no differences on protein levels p-$ERK_{1/2}$,

$ERK_{1/2}$, CREB and ratio of p-$ERK_{1/2}$ to $ERK_{1/2}$ in claustrum between ACE_{EA} and

242   ACE$_{SEA}$ mice. While, the levels of p-CREB, BDNF and radio of p-CREB to

243   CREB were lower in ACE$_{EA}$ mice, when compared with ACE$_{SEA}$ mice.

Figure 4

Figure 4. EA treatment rescues ACE-increased anxiety in mice of

adulthood by suppressing claustrum activation and D1R^{CaMKII}. A,

Experimental design and timeline. **B**, Schematic diagram of EA and SEA

treatments in mice. **C**, EPM test. Two-tailed unpaired t test. Left: t = 2.084, df =

21, p = 0.0496. Middle: t = 1.536, df = 21, p = 0.1394. Right: t = 1.186, df = 21,

p = 0.0837. ACE$_{SEA}$ group, n = 10; ACE$_{EA}$, n = 13. **D**, OFT. Two-tailed unpaired

t test. Left: t = 0.1918, df = 21, p = 0.8498. Middle: t = 0.08086, df = 21, p =

0.9363. Right: t = 0.3799, df = 21, p = 0.7078. ACE$_{SEA}$ group, n = 10; ACE$_{EA}$, n

= 13. **E**, NSFT. Two-tailed unpaired t test. Left: t = 2.532, df = 21, p = 0.0194.

Right: t = 0.6190, df = 21, p = 0.5426. ACE$_{SEA}$ group, n = 10; ACE$_{EA}$, n = 13. **F**,

The percentage and number of c-Fos-positive & CaMKII-positive neurons. Two-

tailed unpaired t test. Upper: t = 4.246, df = 10, p = 0.0017. Lower: t = 3.076, df

= 4, p = 0.0117. ACE$_{SEA}$ group, n = 6; ACE$_{EA}$, n = 6. Scale bar, 300 μm/50 μm.

G, The D1R protein level. Two-tailed unpaired t test. t = 4.396, df = 4, p = 0.0117.

ACE$_{SEA}$ group, n = 3; ACE$_{EA}$, n = 3. **H**, The staining of D1R^{CaMKII}. Two-tailed

unpaired t test. t = 4.874, df = 4, p = 0.0082. ACE$_{SEA}$ group, n = 3; ACE$_{EA}$, n =

3. Scale bar, 200 μm/50 μm. **I**, Levels of p-ERK$_{1/2}$ and ERK$_{1/2}$. Two-tailed

unpaired t test. p-ERK$_{1/2}$: t = 0.2242, df = 4, p = 0.8336. Ratio of p-ERK$_{1/2}$ to

ERK$_{1/2}$: t = 0.3008, df = 4, p = 0.7786. ERK$_{1/2}$: t = 0.7388, df = 4, p = 0.5010.

ACE$_{SEA}$ group, n = 3; ACE$_{EA}$, n = 3. **J**, Levels of p-CREB, CREB and BDNF.

Two-tailed unpaired t test. p-CREB: t = 4.464, df = 4, p = 0.0111. Ratio of p-

CREB to CREB: t = 3.191, df = 4, p = 0.0332. BDNF: df = 4, t = 2.957, p =

0.0417. ACE$_{SEA}$ group, n = 3; ACE$_{EA}$, n = 3. ACE$_{SEA}$, sham electro-acupuncture-

treated ACE mice; ACE$_{EA}$: electro-acupuncture (GV 20 and GV 29)-treated

ACE mice. N.S., $p > 0.05$, *, $p < 0.05$, **, $p < 0.01$ vs ACE$_{SEA}$.

Discussion:

Adolescent cocaine exposure (ACE) produces obvious anxiety in mice of

adulthood. In parallel, ACE evokes CaMKII-positive neurons in claustrum and

increases D1R expression on CaMKII-positive neurons (D1R^{CaMKII}). Blocking

D1R or knocking-down D1R^{CaMKII} within claustrum efficiently rescues ACE-

increased claustrum activation and D1R^{CaMKII}, and ultimately suppresses ACE-

induced anxiety in mice. EA at acupoints of Baihui (GV 20) and Yintang (GV 29)

alleviates ACE-induce anxiety by targeting claustral D1R^{CaMKII} in mice of

adulthood (Figure 5).

Figure 5

Figure 5. Schematic summary of present study. Adolescent cocaine

exposure (ACE) enhances anxiety-like behaviors in mice of adult, accompanied

with triggered activation of CaMKII-positive neurons and more dopamine

receptor 1 (D1R) on CaMKII-positive neurons (D1R^{CaMKII}) in claustrum. Either

blocking D1R with SCH23390 or knocking-down D1R^{CaMKII} with virus in the

claustrum efficiently reduce claustrum activation, and ultimately suppresses

ACE-induced anxiety in mice. By targeting claustral D1R^{CaMKII}, electro-

acupuncture (EA) at acupoints of GV20 and GV29 rescues ACE-increased

anxiety in mice of adulthood. In the process, the p-CREB and BDNF may be

the potential subsequent signals to D1R^{CaMKII}.

Consistent with human and animal studies on ACE (Estelles, Lluch,

Rodríguez-Arias, Aguilar, & Miñarro, 2007; Lopez, Turner, & Saavedra, 2005;

Nie et al., 2020; Shi et al., 2019), including ours, here we found that ACE mice

exhibited more anxiety in their adulthood. Further, we found that CaMKII-

positive neurons in claustrum seem to be involved in the process. Recently, Niu

et.al (Niu et al., 2022) report that claustrum is crucial encoding stress-induced

anxiety, as indicated by attenuation of stress-induced anxiety with silencing

CaMKII-positive neurons in claustrum. Claustrum receives inputs and sends

back axonal outputs to the prefrontal cortex (PFC), a critical region coding

cocaine-related anxiety (Jackson, Karnani, Zemelman, Burdakov, & Lee, 2018;

Zingg, Dong, Tao, & Zhang, 2018). In our previous study, we found that ACE

altered the function of local circuit from GABAergic interneuron to pyramidal

neurons in mPFC, which mediates ACE-increased anxiety in adult mice (Shi et

al., 2019). Due to the robust communications between mPFC and claustrum,

ACE-triggered claustrum should contribute to the anxiety in mice of adulthood.

Firstly, we explored the potential molecules that underlie the evoked claustrum

by ACE. A recent study by Terem et al. (Terem et al., 2020) found that D1R-

positive (D1R$^+$) neurons in claustrum, especially those connected with PFC are

required for acquisition and reinstatement of cocaine preference. They focus

the cell subgroups of D1R$^+$ and D1R$^-$ in cocaine-preferred behaviors. Here, our

findings show that it is D1R^{CAMKII} that were elevated by ACE in claustrum of adult mice, and we focus on the role of D1R in activation of CaMKII-positive neurons and ACE-related anxiety. Moreover, locally blocking D1R or knocking-down D1R^{CaMKII} in claustrum reduced ACE-evoked claustrum and subsequent anxiety. These findings indicate that activity of claustrum CaMKII-positive neurons encodes ACE-increased anxiety, which is mediated at least partially by D1R^{CaMKII}. In other D1R-rich brain regions, such as PFC and hippocampus, cocaine abuse could alter p-ERK (Siemsen, Lombroso, & McGinty, 2018), p-CREB (Valzachi, Teodorov, Marcourakis, Bailey, & Camarini, 2013) and BDNF (Caffino, Giannotti, Messa, Mottarlini, & Fumagalli, 2019; Caffino et al., 2020). Here, we found that protein levels of p-ERK, p-CREB and BDNF were significantly increased in claustrum of ACE mice, implying that they are potential downstream signals for claustral D1R^{CaMKII} in the process of ACE-induced anxiety.

At present, neither D1R antagonists nor specific virus-regulated D1R could be clinically used to treat emotional disorders such as ACE-induced anxiety in humans. EA has been widely applied in management of substance abuse, especially for drugs-caused withdrawal symptoms, including anxiety (D. H. Kim et al., 2018; Nie et al., 2020; Zeng et al., 2018). Shao et al (Shao et al., 2022) reported that EA at Zusanli (ST36) and Kunlun (BL60) could reduce anxiety-like behaviors in animals by suppressing neuropeptide Y in anterior cingulate cortex. Wu et al. (Wu et al., 2022) found that EA at ST36 and Sanyinjiao (SP6) alleviated anxiety by activating D1R or antagonizing D2R in the basolateral amygdala. Previously, we found that EA with a stimulus of a mixed frequency of 2/100 Hz could alleviate ACE-induced anxiety in mice (Nie et al., 2020). With

the findings of claustrum here, we speculated that claustral D1R^{CaMKII} might be

one pharmacological targets of EA treatment. As expected, we found that EA at

acupoints of GV20 and GV29 reversed ACE-enhanced claustrum activation by

suppressing claustral D1R^{CaMKII}, which ultimately ameliorated ACE-induced

anxiety.

Collectively, our findings identified a novel role of claustrum in ACE-induced

anxiety, and put new insight into the function of claustral D1R^{CaMKII}. By targeting

claustral D1R^{CaMKII}, EA obviously alleviates ACE-induced anxiety in mice of

adulthood.

Materials and methods:

Animals

Male C57BL/6 wild type (WT) mice weighing 12-16 g were used. All animals

were housed at constant humidity (40~60%) and temperature (24 ± 2^{o}C) with a

12-hour light/dark cycle (lights on at 8 a.m.) and allowed free access to food

and water. Handling of the animals was carried out for three days before onset

of experiments. All procedures were carried out in accordance with the National

Institutes of Health Guide for the Care and Use of Laboratory Animals and

approved by the Institutional Animal Care and Use Committee (IACUC) at

Nanjing University of Chinese Medicine.

Drug treatment

On postnatal day 28 (P28), male C57BL/6 WT mice were assigned to receive

a single daily injection of cocaine hydrochloride (15 mg/kg, dissolved in saline,

i.p., Qinghai Pharmaceutical, China) or saline (0.2 mL, i.p.) once daily at 10 a.m.

for 14 consecutive days. From P42-P70, these mice were kept at their home

cage (four mice one cage). On P71、P73 and P75, the related behavioral tests

were performed. On P79, the mice were sacrificed after stimulus of elevated

plus maze (EPM) procedure. The brain tissue was collected for qPCR, western

blot and immunofluorescence.

Electro-acupuncture treatment

During P42 to P70, two acupuncture needles (0.16 × 7 mm, Hwato, China) were

inserted gently in a depth of 2–3 mm at the acupoints of GV20 (Baihui, the

midpoint between the auricular apices) and GV29 (Yintang, the midpoint

between the medial ends of the two eyebrows) in each mouse, the acupoints

were electric stimuli of 2/100 Hz at an intensity of 1 mA (HANS Electronic

Apparatus, China) for 30 min once a day at acupoints of GV20 and GV29 in EA

mice, while at "placebo-point" (at 1 cm from base of the tail) in Sham EA mice

(SEA).

Behavioral tests

EPM, open field test (OFT), marble burying test (MBT) and novelty suppressed

feeding test (NSFT) were performed to assess anxiety-like behaviors. All

behavioral tests were conducted during light cycle. In the current study, four

separated cohorts of the adolescent cocaine exposure mice were used to

assess anxiety-like behaviors. All data were recorded and analyzed by TopScan

H system (TopScan, USA). The mice were kept at their home cage until

experiments.

The EPM apparatus (TopScan, Kansas City, MO, USA) consists of 4 elevated

arms (52 cm above the floor) arranged in a cross pattern with two 1-cm walled

arms (open arms) and two 40-cm walled arms (closed arms). Each mouse was

placed in the center portion of the EPM with its head facing an open arm and

allowed freely to explore the maze for 5 min. The time spent, the entries and

the distance traveled in the open arms were recorded.

In OFT, mice were placed gently in the center of the open field apparatus (50

× 50 × 50 cm) individually, and the time spent and the entries into the central

area (16× 16 × 16 cm) of the box and the total distance traveled in the box were

recorded for 5 min with a camera positioned overhead.

In MBT, mice were placed individually in a clean cage (25 × 25 × 50 cm) with

5 cm corncob bedding. Sixteen marbles (14 mm diameter) were distributed

evenly in the cages in four rows of four on top of cage bedding. Mice were left

undisturbed for 30 min, and the number of marbles buried (>50% of marble

covered by bedding material) was recorded.

In NSFT, mice were placed in a square acrylic box (50 × 50 × 50 cm) with the

floor covered by 3 cm of corncob bedding. Twenty-four hours before the test,

all animals were food deprived. On the day of the test a single regular chow

pellet was placed in a white platform located in the middle of the box. Each

animal was placed in one of the apparatus corners, and the latency to start

feeding in the new environment was measured with a cutoff period of 5 min.

The stopwatch was immediately stopped when the mouse bit the chow, using

its forepaws sitting on its haunches. After the test, all animals returned to their

home cages, and the amount of food consumed in a total period of 5 min was

measured.

Western blot

Total protein was extracted from mice CL (the middle region) using RIPA lysis

buffer (Beijing ComWin Biotech Co., China) according to the manufacturer's

instructions. Protein samples (15 μg) was separated by 10% SDS–PAGE, 12%

SDS–PAGE and electrophoretically transferred onto PVDF membranes. The

transferred membranes were blocked with 5% non-fat dry milk or 5% BCA and

0.1% Tween 20 in 10 mM Tris–HCl (TBST buffer, pH 7.5) for 1.5 h at room

temperature, then subsequently incubated with the following primary antibodies:

D1R (1:1000, Rabbit, RRID:AB_10598308, Proteintech, USA), D2R (1:1000,

Rabbit, ER1907-70, Hua An Biotechnology Company, China), D3R (1:1000,

Mouse, ab155098, Abcam, UK), ERK$_{1/2}$ (1:1000, Rabbit, BM4326, Boster,

China), p-ERK$_{1/2}$ (1:1000, Rabbit, BM4326, Boster, China), CREB (1:1000,

Rabbit, RRID:AB_10699020, Cell Signaling Technology, USA), p-CREB

(1:1000, Rabbit, RRID:AB_2561044, Cell Signaling Technology, USA), c-Fos

(1:2000, Rabbit, RRID:AB_2247211, Cell Signaling Technology, USA), BDNF

(1:1000, Mouse, RRID:AB_2881675, Proteintech, USA). The next day, the

membranes were washed three times in Tris-buffered saline with Tween 20

(TBS-T) and incubated with horseradish peroxidase (HRP)-conjugated

secondary antibody goat anti-rabbit or goat anti mouse (1:5000, Beijing

ComWin Biotech Co., China) at room temperature for 1 h. The blots were

visualized by the ECL kit (Beyotime, China) and the signal was visualized by

imaging system (Tanon-5200, Shanghai, China). The blots were washed with

stripping buffer (Beyotime Institute of Biotechnology) to allow reprobing with

other antibodies. In this study, GAPDH or β-actin was used as the loading

control. Values for target protein levels were calculated using Image J software

(NIH, USA). The relative level of each protein expression (blot volume) was

normalized to GAPDH or β-actin.

RNA extraction and quantitative real-time PCR

Mice were decapitated 45 min following EPM behavioral tests on P79. CL were

collected immediately. Total RNA was extracted from the middle of CL using

FastPure Cell/Tissue Total RNA Isolation Kit (Vazyme, China), according to the

manufacturer's instructions. All samples were tested for concentration and

purified using a 4200 TapeStation (Agilent Technologies, Santa Clara, USA).

After RNA extraction, the amount of RNA was normalized across samples, and

cDNA was created using iScript (1708891, Bio-Rad, Hercules, USA). Actin was

used as the internal control. The primers of *D1R, D2R, D3R* and Actin were as

follows: *D1R* (forward, ATGGCTCCTAACACTTCTACCA; reverse,

GGGTATTCCCTAAGAGAGTGGAC); *D2R* (forward,

ACCTGTCCTGGTACGATGATG; reverse,

GCATGGCATAGTAGTTGTAGTGG); *D3R* (forward,

AGACAACATGGAGAGCTGAAACGCT; reverse,

TTCAGGGCTGTGGATAACCTGCC); Actin (forward,

GGCTGTATTCCCCTCCATCG; reverse, CCAGTTGGTAACAATGCCATGT).

D1R, D2R and *D3R* mRNA levels were normalized to Actin mRNA levels. The

relative mRNA level was calculated by the comparative CT method ($2^{-\Delta\Delta Ct}$).

Immunofluorescence

453 Mice brains were collected within 90 min after EPM procedure and perfused

454 with 4% paraformaldehyde (PFA). The coronal brain sections (30 μm) were cut

455 on a cryostat (Leica, Germany). The sections were incubated in 0.3% (v/v)

Triton X-100 for 0.5 h, blocked with 5% donkey serum for 1.5 h at room temperature, and incubated overnight at 4°C with the following primary antibodies: rabbit anti-c-Fos (1:2000, RRID: AB_2247211, Cell Signalling Technology, USA) , mouse anti-NeuN (1:800, RRID: AB_2298772, Millipore, USA), mouse anti-GAD67 (1:800, RRID: AB_2278725, Millipore, USA), mouse anti-PV (1:250, BM1339, Boster Biotechnology, China), mouse anti-SOM (1:50, RRID: AB_2271061, Santa Cruz, USA) and mouse anti-CaMKIIα (1:100, RRID: AB_626789, Santa Cruz, USA), followed by the corresponding fluorophore-conjugated secondary antibodies for 1 h at room temperature. The following secondary antibodies were used here: Alexa Fluor 555-labeled donkey anti-rabbit secondary antibody (1:500, RRID: AB_162543, Invitrogen, USA), Alexa Fluor 488-labeled donkey anti-mouse secondary antibody (1:500, RRID: AB_141607, Invitrogen, USA). Fluorescence signals were visualized using a Leica DM6B upright digital research microscope (Leica, Germany) or Leica TCS SP8 (Leica, Germany).

Local administration of SCH-23390 in claustrum

The cannula (O.D. 0.41 mm, C.C 3.0 mm, Cat 62004, RWD, China) were planted into claustrum (AP + 0.75 mm, ML ± 3.30 mm, DV – 2.50 mm) of ACE mice on P64 before anxiety-like behaviors test. After surgery, mice were maintained at home cage about 1 week. From P71, mice were subjected to EPM, OFT, NSFT. The D1 receptor antagonist R(+)-7-chloro-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride (SCH-23390, Selleck chemicals, China) was dissolved in 0.9% saline and stored -80°C. A volume of 70 nL of SCH-23390 were bilaterally injected into claustrum at a rate of 10 nL/min at 20 min prior to behavioral tests.

Specific knockdown of D1R on CaMKII-positive neurons of claustrum

On P50, mice were anesthetized with 2% isoflurane in oxygen, and were fixed in a stereotactic frame (RWD, Shenzhen, China). A heating pad was used to maintain the core body temperature of the mice at 36°C. The coordinates of claustrum were defined as AP + 0.62 mm, ML ± 3.05 mm, DV –3.50 mm. A volume of 70 nL virus of *CaMKIIap-EGFP-MIR155(shDrd1)-SV40 PolyA* (KD group, Titer: 8.89E+13v.g/ml, GeneChem, Shanghai, China) and *CaMKIIap-EGFP-MIR155(MCS)-SV40 PolyA* (Ctrl group, Titer: 1E+13v.g/ml, GeneChem, Shanghai, China) were injected bilaterally into claustrum at a rate of 10 nL/min. After surgery, mice were kept at home-cage about 3 weeks.

Statistical analysis

Statistical analysis was carried out using GraphPad Prism 8.0.2 software. Data are presented as the Mean ± SEM. The MBT results was analyzed by Two-Way ANOVA followed by Šídák's multiple comparisons test. Other data was analyzed by unpaired t-tests. Statistical significance was set as $p < 0.05$.

Acknowledgments:

This work is supported by National Natural Science Foundation of China (82211540400, 82271531 and 82071495), Natural Science Foundation of Jiangsu Province, China (BK20201398) and Natural Science Foundation of the Higher Education Institutions of Jiangsu Province, China (21KJB360007).

Author contribution:

Chen L, Liu Z and Zhao Z have equal contribution to the manuscript. Chen L, Liu Z and Zhao Z performed behavioral tests, morphological tests and virus-related experiment. Guo H, Du D, Pan W, Wei X and Nie J assist the experiment.

Fan Y, Ge F and Ding J performed data analysis. Guan X and Fan Y wrote the

manuscript. Kim HY edited the manuscript. Guan X developed the overall

Competing interests:

The authors have no conflict of interests to declare.

Data availability statement:

The data that support the findings of this study are available from the

corresponding author upon reasonable request.

Reference

Acheson, A., Ray, K. L., Hines, C. S., Li, K., Dawes, M. A., Mathias, C. W., . . . Laird, A. R. (2015). Functional activation and effective connectivity differences in adolescent marijuana users performing a simulated gambling task. *J Addict, 2015*, 783106. doi: 10.1155/2015/783106

Amorim, D., Amado, J., Brito, I., Fiuza, S. M., Amorim, N., Costeira, C., & Machado, J. (2018). Acupuncture and electroacupuncture for anxiety disorders: A systematic review of the clinical research. *Complement Ther Clin Pract, 31*, 31-37. doi: 10.1016/j.ctcp.2018.01.008

Bayat, A., Joshi, S., Jahan, S., Connell, P., Tsuchiya, K., Chau, D., . . . Koubeissi, M. Z. (2018). A pilot study of the role of the claustrum in attention and seizures in rats. *Epilepsy Res, 140*, 97-104. doi: 10.1016/j.eplepsyres.2018.01.006

Brennan, E. K. W., Jedrasiak-Cape, I., Kailasa, S., Rice, S. P., Sudhakar, S. K., & Ahmed, O. J. (2021). Thalamus and claustrum control parallel layer 1 circuits in retrosplenial cortex. *Elife, 10*, e62207. doi: 10.7554/eLife.62207

Caffino, L., Giannotti, G., Messa, G., Mottarlini, F., & Fumagalli, F. (2019). Repeated cocaine exposure dysregulates BDNF expression and signaling in the mesocorticolimbic pathway of the adolescent rat. *World J Biol Psychiatry, 20*(7), 531-544. doi: 10.1080/15622975.2018.1433328

Caffino, L., Mottarlini, F., Mingardi, J., Zita, G., Barbon, A., & Fumagalli, F. (2020). Anhedonic-like behavior and BDNF dysregulation following a single injection of cocaine during adolescence. *Neuropharmacology, 175*, 108161. doi: 10.1016/j.neuropharm.2020.108161

Chae, Y., Yeom, M., Han, J.-H., Park, H.-J., Hahm, D.-H., Shim, I., . . . Lee, H. (2008). Effect of acupuncture on anxiety-like behavior during nicotine withdrawal and relevant mechanisms. *Neurosci Lett, 430*(2), 98-102. doi: https://doi.org/10.1016/j.neulet.2007.10.026

Chambers, R. A., Taylor, J. R., & Potenza, M. N. (2003). Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *Am J Psychiatry, 160*(6), 1041-1052. doi: 10.1176/appi.ajp.160.6.1041

Druga, R., Smythies, J., Edelstein, L., & Ramachandran, V. (2014). The Claustrum: Structural, Functional, and Clinical Neuroscience. *The Claustrum: Structural, Functional and Clinical Neuroscience, eds Smythies J. R., Edelstein L. R., Ramachandran V. S.*, 29-84.

Erb, S. (2010). Evaluation of the relationship between anxiety during withdrawal and stress-induced reinstatement of cocaine seeking. *Prog Neuropsychopharmacol Biol Psychiatry, 34*(5), 798-807. doi: 10.1016/j.pnpbp.2009.11.025

Estelles, J., Lluch, J., Rodríguez-Arias, M., Aguilar, M. A., & Miñarro, J. (2007).

Cocaine exposure during adolescence affects anxiety in adult mice. *Brain Res Bull, 71*(4), 393-403. doi: 10.1016/j.brainresbull.2006.10.008

Hinova-Palova, D. V., Edelstein, L., Landzhov, B., Minkov, M., Malinova, L., Hristov, S., . . . Ovtscharoff, W. (2014). Topographical distribution and morphology of NADPH-diaphorase-stained neurons in the human claustrum. *Front Syst Neurosci, 8*, 96. doi: 10.3389/fnsys.2014.00096

Jackson, J., Karnani, M. M., Zemelman, B. V., Burdakov, D., & Lee, A. K. (2018). Inhibitory Control of Prefrontal Cortex by the Claustrum. *Neuron, 99*(5), 1029-1039.e1024. doi: 10.1016/j.neuron.2018.07.031

Kalaitzakis, M. E. (2014). Chapter 12 - Parkinson's Disease and the Claustrum. In J. R. Smythies, L. R. Edelstein & V. S. Ramachandran (Eds.), *The Claustrum* (pp. 277-297). San Diego: Academic Press.

Kim, D. H., Kim, N. J., Zhao, R. J., Kim, D. H., Yang, C. H., Kim, H. Y., . . . Lee, B. H. (2018). Effects of acupuncture on the anxiety-like behavior induced by withdrawal from chronic morphine use. *Neurosci Lett, 664*, 38-42. doi: https://doi.org/10.1016/j.neulet.2017.11.008

Kim, D. H., Kim, N. J., Zhao, R. J., Kim, D. H., Yang, C. H., Kim, H. Y., . . . Lee, B. H. (2018). Effects of acupuncture on the anxiety-like behavior induced by withdrawal from chronic morphine use. *Neurosci Lett, 664*, 38-42. doi: 10.1016/j.neulet.2017.11.008

Kurada, L., Bayat, A., Joshi, S., & Koubeissi, M. Z. (2019). The Claustrum in Relation to Seizures and Electrical Stimulation. *Front Neuroanat, 13*, 8. doi: 10.3389/fnana.2019.00008

Laviola, G., Adriani, W., Terranova, M. L., & Gerra, G. (1999). Psychobiological risk factors for vulnerability to psychostimulants in human adolescents and animal models. *Neurosci Biobehav Rev, 23*(7), 993-1010. doi: 10.1016/s0149-7634(99)00032-9

Lopez, B., Turner, R. J., & Saavedra, L. M. (2005). Anxiety and risk for substance dependence among late adolescents/young adults. *J Anxiety Disord, 19*(3), 275-294. doi: 10.1016/j.janxdis.2004.03.001

Madden, M. B., Stewart, B. W., White, M. G., Krimmel, S. R., Qadir, H., Barrett, F. S., . . . Mathur, B. N. (2022). A role for the claustrum in cognitive control. *Trends Cogn Sci, 26*(12), 1133-1152. doi: 10.1016/j.tics.2022.09.006

Markowitsch, H. J., Irle, E., Bang-Olsen, R., & Flindt-Egebak, P. (1984). Claustral efferents to the cat's limbic cortex studied with retrograde and anterograde tracing techniques. *Neuroscience, 12*(2), 409-425. doi: 10.1016/0306-4522(84)90062-9

Moss, H. B., Chen, C. M., & Yi, H. Y. (2014). Early adolescent patterns of alcohol, cigarettes, and marijuana polysubstance use and young adult substance use outcomes in a nationally representative sample. *Drug Alcohol Depend, 136*, 51-62. doi: 10.1016/j.drugalcdep.2013.12.011

Nie, J., Wei, X., Xu, X., Li, N., Li, Y., Zhao, Y., . . . Guan, X. (2020). Electro-acupuncture alleviates adolescent cocaine exposure-enhanced anxiety-like behaviors in adult mice by attenuating the activities of PV

interneurons in PrL. *FASEB J, 34*(9), 11913-11924. doi: 10.1096/fj.202000346RR

Niu, M., Kasai, A., Tanuma, M., Seiriki, K., Igarashi, H., Kuwaki, T., . . . Hashimoto, H. (2022). Claustrum mediates bidirectional and reversible control of stress-induced anxiety responses. *Sci Adv, 8*(11), eabi6375. doi: doi:10.1126/sciadv.abi6375

Rutherford, H. J., Mayes, L. C., & Potenza, M. N. (2010). Neurobiology of adolescent substance use disorders: implications for prevention and treatment. *Child Adolesc Psychiatr Clin N Am, 19*(3), 479-492. doi: 10.1016/j.chc.2010.03.003

Shao, F., Du, J., Wang, S., Cerne, R., Fang, J., Shao, X., . . . Fang, J. (2022). Electroacupuncture Alleviates Anxiety-like Behavior in Pain Aversion Rats by Attenuating the Expression of Neuropeptide Y in Anterior Cingulate Cortex. *Clinical Complementary Medicine and Pharmacology, 2*(2), 100019. doi: https://doi.org/10.1016/j.ccmp.2022.100019

Shapo, G. R., Ma, L., & Liu, C. (2006). [Effects of acupuncture at Weizhong (BL40) on brain function with PET/CT]. *Zhongguo Zhong Xi Yi Jie He Za Zhi, 26*(11), 969-972.

Shi, P., Nie, J., Liu, H., Li, Y., Lu, X., Shen, X., . . . Guan, X. (2019). Adolescent cocaine exposure enhances the GABAergic transmission in the prelimbic cortex of adult mice. *FASEB J, 33*(7), 8614-8622. doi: 10.1096/fj.201802192RR

Siemsen, B. M., Lombroso, P. J., & McGinty, J. F. (2018). Intra-prelimbic cortical inhibition of striatal-enriched tyrosine phosphatase suppresses cocaine seeking in rats. *Addict Biol, 23*(1), 219-229. doi: 10.1111/adb.12504

Spear, L. P. (2000). The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev, 24*(4), 417-463. doi: 10.1016/s0149-7634(00)00014-2

Stiefel, K. M., Merrifield, A., & Holcombe, A. O. (2014). The claustrum's proposed role in consciousness is supported by the effect and target localization of Salvia divinorum. *Front Integr Neurosci, 8*, 20. doi: 10.3389/fnint.2014.00020

Stopponi, S., Soverchia, L., Ubaldi, M., Cippitelli, A., Serpelloni, G., & Ciccocioppo, R. (2014). Chronic THC during adolescence increases the vulnerability to stress-induced relapse to heroin seeking in adult rats. *Eur Neuropsychopharmacol, 24*(7), 1037-1045. doi: 10.1016/j.euroneuro.2013.12.012

Terem, A., Gonzales, B. J., Peretz-Rivlin, N., Ashwal-Fluss, R., Bleistein, N., Del Mar Reus-Garcia, M., . . . Citri, A. (2020). Claustral Neurons Projecting to Frontal Cortex Mediate Contextual Association of Reward. *Curr Biol, 30*(18), 3522-3532.e3526. doi: 10.1016/j.cub.2020.06.064

Valzachi, M. C., Teodorov, E., Marcourakis, T., Bailey, A., & Camarini, R. (2013). Enhancement of Behavioral Sensitization, Anxiety-Like Behavior, and Hippocampal and Frontal Cortical CREB Levels Following Cocaine

Abstinence in Mice Exposed to Cocaine during Adolescence. *PLoS One, 8*(10), e78317. doi: 10.1371/journal.pone.0078317

Wang, Q., Ng, L., Harris, J. A., Feng, D., Li, Y., Royall, J. J., . . . Zeng, H. (2017). Organization of the connections between claustrum and cortex in the mouse. *J Comp Neurol, 525*(6), 1317-1346. doi: 10.1002/cne.24047

Wegiel, J., Moryś, J., Kowiański, P., Ma, S. Y., Kuchna, I., Nowicki, K., . . . Wisniewski, T. (2014). Chapter 8 - Delayed Development of the Claustrum in Autism. In J. R. Smythies, L. R. Edelstein & V. S. Ramachandran (Eds.), *The Claustrum* (pp. 225-235). San Diego: Academic Press.

Wu, M., Chen, Y., Shen, Z., Zhu, Y., Xiao, S., Zhu, X., . . . Shao, X. (2022). Electroacupuncture Alleviates Anxiety-Like Behaviors Induced by Chronic Neuropathic Pain via Regulating Different Dopamine Receptors of the Basolateral Amygdala. *Mol Neurobiol*. doi: 10.1007/s12035-022-02911-6

Yanes, J. A., Riedel, M. C., Ray, K. L., Kirkland, A. E., Bird, R. T., Boeving, E. R., . . . Sutherland, M. T. (2018). Neuroimaging meta-analysis of cannabis use studies reveals convergent functional alterations in brain regions supporting cognitive control and reward processing. *Journal of Psychopharmacology, 32*(3), 283-295. doi: 10.1177/0269881117744995

Zarrindast, M. R., & Khakpai, F. (2015). The Modulatory Role of Dopamine in Anxiety-like Behavior. *Arch Iran Med, 18*(9), 591-603.

Zeng, L., Tao, Y., Hou, W., Zong, L., & Yu, L. (2018). Electro-acupuncture improves psychiatric symptoms, anxiety and depression in methamphetamine addicts during abstinence: A randomized controlled trial. *Medicine (Baltimore), 97*(34), e11905. doi: 10.1097/md.0000000000011905

Zhu, W., Ge, X., Gao, P., Li, M., Guan, Y., & Guan, X. (2018). Adolescent cocaine exposure induces prolonged synaptic modifications in medial prefrontal cortex of adult rats. *Brain Struct Funct, 223*(4), 1829-1838. doi: 10.1007/s00429-017-1590-0

Zingg, B., Dong, H. W., Tao, H. W., & Zhang, L. I. (2018). Input-output organization of the mouse claustrum. *J Comp Neurol, 526*(15), 2428-2443. doi: 10.1002/cne.24502