2   **Challenges and advances in methods for measuring phenotypic convergence**

4   David M. Grossnickle[1]*, William H. Brightly[2], Lucas N. Weaver[3], Kathryn E. Stanchak[1], Rachel A.

Roston[4], Spencer K. Pevsner[5], C. Tristan Stayton[6], P. David Polly[7], Chris J. Law[1,8]

[1]University of Washington, Department of Biology

8   [2]University of Sheffield, School of Biosciences

[3]University of Michigan, Department of Ecology and Evolutionary Biology and Museum of Paleontology

10   [4]University of Washington, School of Dentistry, Department of Oral Health Sciences

[5]University of Oxford, Department of Earth Sciences

12   [6]Bucknell University, Department of Biology

[7]Indiana University, Department of Earth and Atmospheric Sciences

14   [8]University of Texas, Austin, Department of Integrative Biology

16   *D. M. G., dmgrossn@uw.edu

ABSTRACT

20   Tests of phenotypic convergence can provide evidence of adaptive evolution, and the

popularity of such studies has grown in recent years due to the development of novel,

22   quantitative methods for identifying and measuring convergence. These methods include the

commonly applied $C1$–$C4$ measures of Stayton (2015), which measure morphological distances

24   between lineages, and Ornstein-Uhlenbeck (OU) evolutionary model-fitting analyses to test

whether lineages convergently evolved toward adaptive peaks. We test the performance of C-

26   measures and other convergence measures under various evolutionary scenarios. We reveal a

critical issue with *C*-measures: they often misidentify divergent lineages as convergent. We

28 address this issue by developing novel convergence measures (*Ct*1–*Ct*4-measures) that

measure distances between lineages at specific points in time, minimizing the possibility of

30 misidentifying divergent taxa as convergent. *Ct*-measures are most appropriate when putatively

convergent lineages are of the same or similar geologic ages (e.g., extant taxa), meaning that

32 the lineages' evolutionary histories include considerable overlap in time. Beyond *C*-measures,

we find that all convergence measures are influenced by the position of focal taxa in

34 morphospace, with morphological outliers often statistically more likely to be measured as

strongly convergent by chance. Further, we mimic scenarios in which researchers assess

36 convergence using OU models with *a priori* regime assignments (e.g., classifying taxa by

ecological traits), and we find that multiple-regime OU models with phenotypically divergent

38 lineages assigned to a shared selective regime often outperform simpler models. This highlights

that model support for these multiple-regime OU models should not be assumed to reflect

40 convergence among focal lineages of a shared regime. Our new *Ct*1–*Ct*4-measures provide

researchers with an improved comparative tool, but we emphasize that all available

42 convergence measures are imperfect, and researchers should recognize the limitations of these

methods and use multiple lines of evidence when inferring and measuring convergence.

46 **INTRODUCTION**

Phenotypic convergence among distantly related taxa is commonly associated with adaptive

48 evolution (e.g., Darwin 1859, Losos 2011), but it can also occur stochastically (Stayton 2008) or

as a byproduct of shared developmental constraints (Losos 2011, Speed and Arbuckle 2016).

50   Evidence for adaptive convergence is strengthened when the magnitude of convergence is

      greater than expected by chance, and also when the convergent phenotypes are shown to be

52   tied to similar ecological or functional roles. Thus, quantitative examinations of phenotypic

      convergence are important; they assist researchers in identifying adaptive phenotypes that are

54   driven by shared selective pressures and/or developmental constraints. Novel methods for

      identifying and measuring convergence have recently been developed (Mahler et al. 2013,

56   Arbuckle et al. 2014, Ingram and Mahler 2013, Stayton 2015A, Speed and Arbuckle 2017,

      Castiglione et al. 2019), and these methods are often accompanied by statistical tests for

58   comparing the measured convergence to that which is expected from a null model-fitting

      hypothesis or random data permutations. This has increased the accessibility of quantitative

60   tests for phenotypic convergence, leading to a flood of recent studies on that topic (e.g.,

      Friedman et al. 2016, Zelditch et al. 2017, Baliga and Mehta 2018, Da Silva et al. 2018, Arbour

62   and Zanno 2020, Button and Zanno 2020, Grossnickle et al. 2020, Martinez et al. 2020, Serio et

      al. 2020, Rincon-Sandoval et al. 2020, Spear and Williams 2020, Baumgart et al. 2021, Huie et al.

64   2021, Rovinsky et al. 2021, Tamagnini et al. 2021, Alfieri et al. 2022, Bennion et al. 2022, Canale

      et al. 2022, Law 2022).

66   Phenotypic convergence is often defined as lineages evolving to be more similar to

      one another than were their ancestors (Losos 2011, Stayton 2015A, Mahler et al. 2017), and we

68   follow that definition here. Thus, a signature of convergence is phylogenetic tips that are

      phenotypically more similar to one another than expected based on assumptions of random

70   change over time; the degree of this similarity of tips is often quantified by convergence

      measures (Speed and Arbuckle 2017). However, a confounding issue is that multiple types of

72   evolutionary trajectories can result in lineages that are more similar to one another than

expected by chance but are not convergent (as defined above). This includes lineages that

74   retain a shared ancestral phenotype (see discussion on 'conservatism' below) and lineages that

have parallel evolutionary trajectories from a similar ancestral trait condition.

76   The $C1$–$C4$ measures (hereafter, 'C-measures') developed by Stayton (2015A) have

emerged as an especially popular means of quantifying morphological convergence (e.g.,

78   Friedman et al. 2016, Zelditch et al. 2017, Baliga and Mehta 2018, Da Silva et al. 2018, Arbour

and Zanno 2020, Button and Zanno 2020, Grossnickle et al. 2020, Martinez et al. 2020, Rincon-

80   Sandoval et al. 2020, Spear and Williams 2020, Baumgart et al. 2021, Huie et al. 2021, Rovinsky

et al. 2021, Tamagnini et al. 2021, Alfieri et al. 2022, Bennion et al. 2022, Canale et al. 2022,

82   Law 2022). C-measures are calculated using geometric distances in phenotypic space between

focal lineages, relying on ancestral reconstructions for morphologies at ancestral nodes. The

84   underlying feature of the C-measures is the comparison of two measurements: the maximum

phenotypic distance between lineages at any points in their evolutionary histories (D_{max}) and

86   the phenotypic distance between phylogenetic tips (D_{tip}). More specifically, D_{max} is the greatest

distance between any two points along the lineages in phylomorphospace, with candidate

88   distances including any points between the lineages' most recent common ancestor and the

tips (Fig. 1A). $C1$ is the primary C-measure and calculated as $1 - (D_{tip}/D_{max})$, with the resulting

90   value representing "the proportion of the maximum distance between two lineages that has

been 'closed' by subsequent evolution" (Stayton 2015A). In our conceptual illustration (Fig. 1A),

92   two focal lineages have convergently evolved such that their tips are 70% closer to each other

than their D_{max}, resulting in a $C1$ score of 0.7.

Figure 1. Conceptual illustrations of two methods for assessing phenotypic convergence of focal
lineages (maroon): *A*, *C*1 of Stayton (2015A) and, *B*, multiple-regime Ornstein-Uhlenbeck (OU)
model-fitting. The *C*1 score of 0.7 indicates that lineages have evolved toward each other to cover
70% of the maximum distance (D_{max}) between their lineages. D_{max} can be measured at any point
along the evolutionary histories, including the dashed branches in *A*, and D_{tip} is the morphological
distance between phylogenetic tips. Although time is a variable in the univariate illustration in *A*, the
C-measures do not incorporate time. *B*, OU models include fitting a trait optimum parameter that is
often interpreted as the location of an adaptive peak and an 'attraction' parameter that is
commonly interpreted as the strength of selection. Abbreviation: MRCA, most recent common
ancestor.

108   One reason for the popularity of C-measures is that they can distinguish between

convergence and conservatism, which both result in distantly-related phylogenetic tips with

110   similar phenotypes. The key difference between convergence and conservatism centers on the

ancestral morphologies of the lineages. Whereas convergence involves ancestors that were less

112   morphologically similar to each other than their descendant tips are to one another (Losos

2011, Stayton 2015A, Mahler et al. 2017), conservatism is the lack of substantial phenotypic

114   divergence from ancestral morphologies relative to what is expected from random processes

(Losos 2008, Moen et al. 2013, McLean et al. 2018). The 'blue' lineages in Figure 1B could be

116   considered an example of conservatism; they have not evolved far from the ancestral

morphology. C-measures account for ancestral patterns via the D_{max} measurement (Fig. 1A).

118   Alternative distance-based methods for testing for convergence (e.g., Wheatsheaf index,

Arbuckle et al. 2014, Arbuckle and Minter 2015; θ, Castiglione et al. 2019) cannot adequately

120   differentiate between convergence and conservatism (or parallelism) because phenotypic

distances between ancestral morphologies are not considered or, in the case of θ, only partially

122   integrated (Castiglione et al. 2019).

        In addition to distance-based measures, researchers often use evolutionary model-

124   fitting analyses to assess convergence, relying on strong fits of multiple-regime Ornstein-

Uhlenbeck (OU) models (Hansen 1997, Butler and King 2004) to morphological data for

126   evidence of convergence (e.g., Mahler et al. 2013, Ingram and Mahler 2013, Friedman et al.

2016, Mahler et al. 2017, Baliga and Mehta 2018, Grossnickle et al. 2020, Martinez et al. 2020).

128   An OU process involves 'attraction' toward an 'attractor' or trait optimum (commonly

interpreted as the location of an adaptive peak in morphospace), and this attraction and any

130   resulting convergence is often assumed to be due to selective pressures toward adaptive peaks

(Fig. 1B). Convergence is sometimes assumed when the best-supported model indicates that

132   two or more focal lineages have independently evolved toward the same trait optimum.

Implementations of these multiple-regime OU models either classify focal taxa into regimes *a*

134   *priori* (e.g., based on shared ecological or behavioral traits) or examine whether focal taxa are

independently identified (i.e., without *a priori* classifications) as shifting toward shared adaptive

136   peaks. The latter method can be implemented using functions in the *SURFACE R* package

(Ingram and Mahler 2013; but see Adams and Collyer 2018 for a critique), which was developed

138   specifically for identifying convergence. Other *R* packages have been designed to identify peak

shifts without *a priori* classifications (e.g., *Bayou*, Uyeda and Harmon 2014; *l1ou*, Khabbazian et

140   al. 2016; *phyloEM*, Bastide et al. 2018), although they are not specifically designed to identify

convergence. However, an OU process does not require that independent lineages shifting

142   toward a peak come from more dissimilar ancestors and thus lineages independently evolving

to a shared peak (as evidenced by support for OU models) may not meet the criteria of

144   convergence used in this paper. In addition, OU models may be prone to overfitting to data,

and model support should thus be interpreted with caution (Ho and Ané 2014, Cooper et al.

146   2016, Adams and Collyer 2018; but see Grabowski et al. 2023). Further, in some scenarios it

may be challenging to use OU model-fitting analyses to differentiate convergence from

148   conservatism; focal taxa may share a given morphology with their common ancestor, but

lacking information about ancestral states, it may be difficult to determine whether this is

150   because the taxa retained the ancestral morphology (conservatism) or have converged on this

ancestral morphology from diverse ancestors which did not exhibit that morphology.

152   Nonetheless, a benefit of OU model-fitting analyses is that the magnitude of the attraction

parameter allows an estimate of selective strength toward adaptive peaks, therefore providing

154   valuable information about the evolutionary process that may be driving convergence.

Here, we test the performances of *C*-measures and other means of assessing

156   convergence by applying those methods to simulated data in which a subset of lineages are

modeled as truly convergent or truly divergent. Our results highlight a critical concern with the

158   *C*-measures: in many circumstances, they may lead to misclassification of divergent lineages as

convergent. Further, we find that the magnitude of convergence measured by distance-based

160   metrics is influenced by the position of focal lineages in morphospace. *C*-measures and θ (i.e.,

the angle between phenotypic vectors; Castiglione et al. 2019) measure greater convergence in

162   morphological outliers, whereas the Wheatsheaf index (Arbuckle et al. 2014, Arbuckle and

Minter 2015) measures greater convergence among lineages that retain their shared ancestral

164   morphology. We also find that in specific scenarios in which OU model regimes are defined *a*

priori by traits independent of morphology (e.g., shared ecological or behavioral traits), OU

166   models treating divergent lineages as having a shared optimum (via a shared regime

assignment) outperform simpler models; such *a priori* approaches are prevalent, highlighting

168   that researchers should use caution when using model-fitting analyses to assess convergence.

Finally, we present an improved method for calculating *C*-measures that minimizes the

170   possibility of erroneously measuring divergent lineages as convergent.

174 **METHODS**

Evolutionary simulations

176 We generated a series of simulated trait datasets to ascertain how frequently convergence

measures correctly identify *convergent* lineages and misclassify *divergent* lineages as

178 convergent. Simulated datasets are intended to reflect typical empirical datasets, but we

include some extreme morphological outliers to help identify trends. Thus, we simulated traits

180 for 201 species included in a recent phylogenetic treatment of extant mammals, resulting in

taxon sampling and phylogenetic structure comparable to many recent comparative studies

182 (Chen and Wilson 2015, Grossnickle et al. 2020, Weaver and Grossnickle 2020, Pevsner et al.

2022). We obtained 1000 randomly chosen phylogenetic trees from the posterior distribution

184 of Upham et al.'s (2019) 'completed trees' analysis. We then used *TreeAnnotator* (Drummond

et al. 2012) to generate a maximum clade credibility tree, which was pruned to the species in

186 our sample. The sample includes 13 gliding-mammal species representing five independent

evolutionary origins of gliding behavior. We treated the gliders as the focal lineages (*sensu*

188 Grossnickle et al. 2020); they were the subject of manipulation in our simulations (as such, we

refer to those simulated glider data as 'gliders'). The five 'glider' clades are spread across the

190 mammalian phylogeny and have varying evolutionary origin ages, making them ideal for

representing typical empirical datasets.

192 Each simulated dataset included six morphological traits. In the non-glider (i.e., 'base')

parts of the tree, all six traits were simulated to evolve via Brownian motion (BM). In 'glider'

194 clades, between three to six of these traits were simulated to evolve via an OU process (except

for drift-based divergence simulations in which all traits were BM-evolved; see descriptions

196   below), with optima varying to produce convergent or divergent trajectories. The BM

component of the simulation ('base simulation') was performed using the

198   *SimulateContinuousTraitsOnTree* function in the *Phylogenetics for Mathematica* package (Polly

2019). The ancestral value for each trait was arbitrarily set to 0.0 and the step rate, σ^2, was set

200   at 1.0 per million years. (Note that the phylogenetic tree was the same for each simulation, but

the BM trait values were re-simulated with every 'base simulation.') In 'gliders,' traits evolving

202   via an OU process toward trait optima (see below) were simulated using the *LineageEvolution*

function in *Phylogenetics for Mathematica* (ancestral trait values were those generated via BM

204   at the base of each clade). Phylogenetic branches of 'gliders' were those tipped by one of the

13 'glider' species, plus the subtending branches below clades whose tips were all 'gliders.'

206   We generated four types of simulations: 1) convergence of 'gliders' via selection (OU-

evolved traits), 2) divergence of 'gliders' via drift (BM-evolved traits), 3) 'constrained

208   divergence' of 'gliders' via selection (OU-evolved traits, limited to positive values), and 4)

'unconstrained divergence' of focal taxa via selection (OU-evolved traits; no limits on traits).

210   The following text describes these four types of simulations, and Figure S1 provides a graphical

summary.

212   *Convergence simulations.* For the convergence simulations, we systematically altered 1)

the number of traits (of the six total traits) that were subjected to convergent selection (three

214   through six) and 2) their optimum values (or 'targets') (Fig. S1). Convergent traits were evolved

via selection (i.e., OU-evolved) toward the same trait optimum (target). The convergent traits

216   were subjected to directional OU-selection for their full branch lengths, which allowed most

'glider' tips (besides those with the shortest branches) to arrive at the adaptive peak. Traits not

218   subject to convergent evolution continued to evolve by BM. Note that the OU-parameters for

        all simulations were the same, and the 'strength' of convergence was represented by the

220   number of traits subject to convergence rather than the strength of selection per se. Six of six

        traits being convergent thus represents very strong convergence, whereas three of six traits

222   being convergent represents weaker convergence. Further, we alter the number of convergent

        traits because it is likely that in empirical datasets some traits are convergent whereas other

224   traits are not convergent (e.g., see the discussion on long bone lengths versus long bone widths

        in Grossnickle et al. 2020).

226   By changing the values of trait optima ('targets') we systematically controlled the

        morphological distance of 'gliders' from their ancestral condition, allowing us to test how

228   morphospace position influences the results of applied convergence measures (Fig. S1). We

        used a series of nine trait optima positioned at successively greater distances from the

230   ancestral point in morphospace, starting at zero (convergence toward the ancestral trait values)

        and successively increasing by 10 to a distance of 80 trait units from the ancestral value. For

232   example, in a simulation with four convergent traits and an optimum of 30, the first four traits

        all evolved toward an optimum trait value of 30 and the two remaining traits evolved by BM.

234   The range of tip values in the 'base' portion of the tree (BM-evolved 'non-gliders') has a radius

        of about 15 trait units, so the first two optima in this iteration (zero and 10) lie within the

236   morphospace occupied by 'non-glider' taxa and the last seven lie increasingly outside the range

        of morphology of 'non-gliders.' The farthest tested optima (i.e., ~60–80) result in the 'gliders'

238   being very far from 'non-gliders' in morphospace and thus unlikely to reflect most empirical

        scenarios, but we include them here to help infer trends in how morphospace position

240   influences convergence measures. Each simulation with all of its iterations was repeated with

15 unique 'base simulations,' and we report results for the means and standard errors of these

242   15 replicates.

Divergence simulations. To simulate divergence via drift (i.e., BM-evolved 'gliders'), we

244   simulated six traits using the *fastBM* function of the *phytools* package (Revell 2012) for *R* (*R*

Core Team 2020). Ancestral trait values were set at zero, and, to mimic natural variation, the

246   rate parameter (σ^2) was sampled from a log-normal distribution with log-mean and standard

deviation 0 and 0.75, respectively. This was repeated to produce 15 replicate datasets. In

248   contrast to all other simulations, gliders in these datasets did not possess any OU-evolved

traits.

250   For the selection-based (OU-evolved) methods for generating divergence, we simulated

divergence as selection of the individual 'glider' lineages each toward a different trait optimum.

252   We used a procedure that is as similar as possible to that used in the convergence simulations.

Between three and six traits were selected toward the clade-specific trait optimum with a

254   series of target distances ranging from 30 trait units from the ancestral morphology (which

extends the lineages past the periphery of the base BM tree and thus ensures that the targets

256   are divergent) to 100 units in steps of 10. This choice, however, means that the divergence

simulations are limited to cases in which the lineages are morphological outliers, whereas the

258   drift-based divergence simulations include non-outlying lineages. A different target was

randomly selected for each 'glider' clade by choosing a random trait value for each of the traits

260   under selection with the condition that their sum of squares equal the squared target distance

(i.e., that the target lies at a distance of 30, 40, etc. units from the ancestral trait values). The

262   selected lineages are allowed to fully reach their trait optima.

For our primary selection-based divergence simulations, we limited the OU-evolved

264   'glider' traits to positive values. Choosing only positive trait values ensures that the trait optima

('targets') are divergent yet lie within the same quadrant (or hyper-region) of multidimensional

266   space. This constriction of adaptive peaks to one region of morphospace helps to better mimic

empirical datasets in which lineages exhibit some morphological similarities but are still

268   geometrically divergent (e.g., Collar et al. 2014, Grossnickle et al. 2020). Further, researchers

are most likely to apply convergence measures to lineages with shared ecological or behavioral

270   traits, which often result in morphological similarities, making our simulations more applicable

to empirical research. Nonetheless, we also simulated 'unconstrained divergence' for a smaller

272   subset of evolutionary scenarios (see discussion on subset analyses below) in which the 'glider'

trait optimum values were not limited to positive values.

274   In total, we generated and analyzed 960 simulated datasets: 540 that simulated trait

convergence and 420 that simulated trait divergence.

C-measures

278   We applied the *C*-measures (Stayton 2015A) to focal lineages ('gliders') in the simulated

datasets. The primary measure, $C1$, is calculated as $1 - (D_{tip}/D_{max})$, with D_{tip} representing the

280   distance between phylogenetic tips of focal taxa and D_{max} representing the maximum distance

between any tips or ancestral nodes of those lineages. The $C1$ value is one for complete

282   convergence zero for divergence (i.e., D_{max} is D_{tip}). $C2$ is the difference between D_{max} and D_{tip},

and it captures the absolute magnitude of convergent change. C3 and C4 are standardized

284   versions of C2 that are calculated by dividing C2 by the phenotypic change along branches

leading to the focal taxa (C3) or the total amount of phenotypic change in the entire clade (C4).

286   See Stayton (2015A) for full descriptions of C1–C4. To calculate C-measure scores, we used

functions in the R script from Zelditch et al. (2017), which are computationally faster than the

288   functions in v1.0 through v1.3 of the *convevol R* package (Stayton 2015A, Stayton 2018). C1–C4

scores were calculated for all simulated datasets, but due to computational limits of analyzing a

290   large number of simulated datasets, we only calculated simulation-based p-values for a smaller

subset of datasets used for subsequent analyses (see the following subsection).

292   *C*-measures quantify phenotypic convergence between individual phylogenetic tips, not

between clades with multiple tips (Stayton 2015A). Thus, we calculated average phenotypes for

294   taxa of focal clades. For example, one glider clade includes six flying squirrel species, so for each

of the six simulated traits we calculated mean values for the six species. The averages were

296   then used as the representative flying squirrel lineage. Thus, C-measures were measured for

five 'glider' lineages, each representing an independent evolution of gliding. The species' traits

298   were not averaged prior to the other types of convergence analyses described below.

300   **Additional measures of convergence**

Subset of simulated datasets. We applied additional measures of convergence (OU model-

302   fitting, θ, and Wheatsheaf index) to a smaller subset of 165 simulated datasets. This subset only

includes datasets in which four of six traits were simulated to converge on a specific trait

304   optimum or diverge toward multiple optima, with the remaining two traits evolved by BM (see

Evolutionary simulations subsection for more details). We did not use datasets in which all six

306 traits are convergent because this leads to nearly complete convergence on a trait optimum,

and complete convergence appears to be very rare among empirical analyses (e.g., Grossnickle

308 et al. 2020, Law et al. 2022). Nonetheless, four of six traits being convergent on an optimum

often results in strong convergence (i.e., statistically significant distance-based measures of

310 convergence and strong fits of multiple-peak OU models) among lineages, especially when trait

optima are outliers in morphospace (see Results & Discussion). For convergence simulations,

312 we used the 15 simulated datasets each from the sets of simulations where trait optima were

set at 0, 20, 40, 60, and 80. These represent simulations in which focal lineages evolve toward

314 the ancestral morphology (optimum = 0), evolve just beyond the outer edge of the

morphospace region of BM-evolved lineages (optimum = 20), and evolve far into outlying

316 morphospace (optima = 40, 60, and 80). For selection-based divergence simulations (both

'constrained' and 'unconstrained'), we used the 15 simulated datasets with optima of 40, 60,

318 and 80. (Using trait optima of 0 or 20 could mistakenly simulate convergence toward ancestral

morphologies.) Thus, the subset of datasets includes 75 convergence simulations (15 datasets

320 each for five trait optima) and 45 simulations each for unconstrained and constrained

divergence simulations (15 datasets each for three optima). The following methods were only

322 applied to this subset of datasets.

 Evolutionary model-fitting analyses. We performed evolutionary model-fitting analyses

324 using both multivariate models via functions in the *mvMORPH R* package (Clavel et al. 2015)

and univariate models via functions in the *OUwie R* package (Beaulieu et al. 2012). The

326 multivariate *mvMORPH* models offer the benefit of incorporating all data, whereas the

univariate *OUwie* models can be set to allow rate (σ^2) and selection (α) parameters to vary

328   between regimes and therefore can better represent the underlying simulated data, which

include both BM-evolved and OU-evolved lineages.

330   We fit four multivariate models to all six simulated traits using functions within the

mvMORPH R package. The first three models were a single-rate multivariate BM model

332   (mvBM1) that assumes trait variance accumulates stochastically but proportionally to

evolutionary time, a single-optimum Ornstein-Uhlenbeck model (mvOU1) that modifies the BM

334   model to constrain each trait to evolve toward a single optimum, and an early burst model

(mvEB) that modifies the BM model to decrease evolutionary rate with time. Note that all of

336   these models do not model 'glider' lineages as evolving to a distinct adaptive peak and thus are

not expected to reflect convergence of 'gliders' on an adaptive peak (e.g., Fig. 1B). In all model-

338   fitting analyses, the mvEB model collapsed to a mvBM model (i.e., the change-in-rate

parameter was zero). Thus, we did not report the mvEB results, and we did not include it when

340   calculating Akaike weights.

        We then fit a multivariate OU model with two selective regimes (mvOU2) that allowed

342   'gliders' and 'non-gliders' to exhibit different trait optima (θ). Support for mvOU2 would

provide evidence of convergence by indicating that selective forces are driving 'glider' lineages

344   to a shared adaptive peak (Fig. 1B). Note that in the simulations 'non-gliders' evolved via BM,

and thus any support for the mvOU2 model is likely to be driven by the 13 'glider' lineages.

346   Relative support for each of the three models was assessed through computation of small-

sample corrected Akaike weights (AICcW; Akaike 1974; Hurvich and Tsai 1989). We calculated

348   AICcW for each of the 15 base simulations, and we report the mean values. Prior to fitting the

mvOU2 model, we mapped the gliding regime character state onto the 'glider' branches of the

350   phylogeny (including the subtending branches below 'glider' clades) using the *paintSubTree*

function in the *phytools R* package (Revell 2012).

352   In addition to fitting models to multivariate data, we fit models to univariate data (PC1

scores) using functions in the *OUwie R* package (Beaulieu et al. 2012). Although fitting

354   evolutionary models to PC scores may generate biased results (Uyeda et al. 2015, Adams and

Collyer 2018), we performed these analyses because they allowed us to fit multiple-regime OU

356   models with varying evolutionary rates (σ) and/or attractions (α) to optima between regimes,

which is not a feature of the multivariate *mvMORPH* models. By allowing parameters to vary

358   between regimes, these univariate models are expected to better fit the underlying simulated

data compared to the *mvMORPH* models. We fit seven univariate evolutionary models to the

360   subset of simulated datasets, including single-regime models (BM1 and OU1) and a suite of

two-regime OU models (i.e., 'OUM' models of Beaulieu et al. 2012). The OU2 model keeps α

362   and σ constant for both 'glider' and 'non-glider' regimes, the OU2A model allows α to vary

between regimes, the OU2V model allows σ^2 to vary between regimes, and the OU2VA model

364   allows both σ and α to vary between regimes. See the Supplemental Methods for additional

discussion on the univariate model-fitting analyses.

366   *Additional distance-based convergence measures*. We applied two other measures of

convergence to the subset of 30 simulated datasets (using all six traits): Wheatsheaf index,

368   which was implemented via the *R* package *windex* (Arbuckle et al. 2014, Arbuckle and Minter

2015), and θ_{real}, which was implemented using the *RRphylo* package (Castiglione et al. 2018,

370   Castiglione et al. 2019). The Wheatsheaf index measures pairwise morphological distances

between putatively convergent taxa, with distances corrected for the degree of phylogenetic

372   relatedness of lineages. These distances are compared to pairwise distances between other

lineages in the sample to determine whether putatively convergent lineages are more similar to

374   each other than expected. The θ measurement is the angle between the phenotypic vectors of

putatively convergent lineages (note that this θ is different from the θ parameter of OU

376   models), and it is based upon phylogenetic ridge regression. We report the angle obtained by

all pairwise comparisons between putatively convergent clades (θ_{real}), standardized by the

378   phylogenetic distance separating them (i.e., expected divergence under a BM model).

Significance tests compare standardized θ_{real} values of putatively convergent taxa to values

380   computed for randomly selected tip pairs.

382   **RESULTS & DISCUSSION**

C-measure issues

384   Our analysis of the *C*-measure calculations reveals that the measures do not always perform as

intended (Stayton 2015A), especially when putatively convergent lineages are outliers in

386   morphospace. This critical problem can manifest in at least three ways, which we illustrate in

Figure 2. First, the more outlying the morphologies are in phylomorphospace (and all else being

388   equal), the greater the *C* scores, indicating stronger convergence. We demonstrate this with a

conceptual illustration in Figure 2A (note that D_{tip} remains constant in both scenarios). This does

390   not align with our working definition of convergence and in Stayton (2015A); the distances

between ancestral nodes and the distances between descendants are unchanged between the

392   scenarios, and thus we could expect *C* scores to be the same for both scenarios. The pattern of

greater convergence in outliers is also demonstrated by results of applying *C*-measures to

394   evolutionary simulations (Figs. 3 and S2); although this pattern only occurs when some

simulated traits are evolved by BM (see the *Relatively greater convergence in morphological*

396   *outliers* subsection below).

Figure 2. Conceptual illustrations of *C*-measure issues. *C*1 scores are greater than zero for the

404   divergent (*B*) and parallel (*C*) focal lineages ('maroon'), incorrectly indicating that the lineages are

convergent. See the main text and Figure 1 for more information on *C*1, D_{max}, and D_{tip}. The *C*-

406   measure issues highlighted here also apply to evolutionary model-fitting analyses. The distribution

curves in univariate illustrations could represent adaptive peaks, and OU model-fitting analyses are

408   more likely to identify unique shared adaptive peaks when a peak is farther from the ancestral

morphology (Table 1).

Figure 3. PCA phylomorphospaces for example datasets that simulate convergence (*A*) and

416   divergence (*B*) of all six traits of five focal clades ('gliders'). Traits were selected toward optima via

an OU process, with all traits of convergent taxa selected toward a value of 20 and traits of

418   divergent taxa selected toward varying values that result in evolutionary change equal to that of

convergent taxa (see Methods). (*C*) *C*1 scores for simulated convergent lineages (solid lines) and

420   divergent lineages (dashed lines), using datasets in which focal taxa have varying numbers of

convergent/divergent traits (of six total) and trait optima positions. Any traits and lineages not

422   selected to be convergent/divergent were evolved by Brownian motion (BM). Focal taxa evolved

toward trait optima after they originated. Divergent trait optima are randomized, but they are

424   limited to being positive numbers (whereas BM-evolved traits can be positive or negative), resulting

in divergent lineages evolving in the same direction along PC1 (e.g., *B*) but otherwise being divergent

426   (unless convergence occurs by chance). *C*1 values above zero indicate convergence, and a value of

one would reflect complete convergence (i.e., phenotypically identical tips). We did not simulate

428   divergence to trait optima of 0, 10, and 20 because the simulations might mistakenly generate

convergent lineages near the middle of morphospace. As a second means of simulating divergence,

430   we allowed the focal lineages to evolve via BM, and these results are displayed as a box-and-whisker

plot in *C*. *C*1 results are means and standard errors of 15 simulated datasets.

434   The second and third issues with the C-measures are more problematic: divergent and

parallel lineages can have $C1$ scores that are greater than zero, incorrectly indicating that the

436   lineages are convergent (Fig. 2B, C). In Figure 2B we illustrate lineages that are diverging

morphologically (in univariate and multivariate morphospace), but which have a $C1$ score of

438   0.3, incorrectly suggesting that the lineages have experienced substantial convergence (i.e.,

closing about 30% of the maximum distance between lineages). To further test this issue, we

440   measured $C1$ in lineages simulated to have divergent traits (Fig. 3B), and $C1$ values are

consistently greater than zero (and statistically significant) when applied to the 'constrained

442   divergence' simulations (Fig. 3C; Table 1), incorrectly indicating convergence instead of

divergence. The $C1$ scores for divergent focal lineages remain around 0.3 at all morphospace

444   positions (Fig. 3B), although see the Supplemental Results and Figure S4 for discussion on how

these results for different morphospace positions may be sensitive to methodological choices.

Table 1. Tests of convergence among focal lineages ('gliders') of the simulated datasets using distance-based measures. Results are means of the 15 simulated datasets for each

optimum. For θ results, we report θ$_{real}$ standardized to phylogenetic distance between clades. Note that relatively smaller θ$_{real}$ values (i.e., smaller angles between phenotypic

vectors) suggest greater convergence, whereas relatively larger Wheatsheaf index, $C1$, and $Ct1$ values indicate greater convergence. Statistical significance (*, $p \leq 0.05$; **, $p \leq 0.01$;

***, $p \leq 0.001$) for $C1$ and $Ct1$ is based on comparisons to results of 100 simulations via a BM model, and for the Wheatsheaf index it is based on bootstrapping with 1000 replicates.

Significance of standardized θ$_{real}$ values is based on bootstrapping with 1000 replicates for each pairwise comparison between the five 'glider' clades (except the monospecific clade).

In all cases, the reported significance is based on means of 15 analyses for a given trait optimum. The 'constrained divergence' simulations limited the simulated traits of 'gliders'

to be positive values, thus constraining them to one region of morphospace (see Methods).

Simulations	Convergence measure	Trait optimum				
		0	20	40	60	80
Convergence	θ (*RRphylo*)	0.393	0.126*	0.071**	0.044**	0.040**
	Wheatsheaf index	1.976***	1.715***	1.314	0.905	0.626
	*C*1	0.350*	0.526***	0.685***	0.751***	0.769***
	*Ct*1	0.176*	0.233**	0.367***	0.497***	0.483***
Divergence ('constrained')	θ (*RRphylo*)	–	–	0.220	0.242	0.229
	Wheatsheaf index	–	–	0.826	0.581	0.434
	*C*1	–	–	0.363**	0.325**	0.310**
	*Ct*1	–	–	0.035	0.011	-0.003

470 Similarly, outlying lineages evolving along parallel phylomorphospace trajectories from a

similar ancestral condition have extremely strong $C1$ scores (Fig. 2C). This is unexpected

472 because the ancestral nodes of both lineages are the same morphological distance from one

another as the distance between tips; this is not convergence according to the definition of

474   convergence adopted here or in Stayton 2015A (see Introduction; Losos 2011, Mahler et al.

2017).

476   The second and third issues with C-measures may commonly occur when at least one

focal lineage has diverged rapidly from other taxa, which is a scenario that can often generate

478   morphological outliers (Fig. 2). However, these issues are not limited to outliers. For instance,

the scenarios for focal ('maroon') lineages in Figures 2B and 2C could occur in any region of

480   morphospace. Thus, these issues have major implications for many empirical studies (see

discussion of examples below); divergent or parallel lineages may often be incorrectly

482   interpreted to be convergent, no matter their position in morphospace.

       The possibility of diverging and parallel lineages having $C1$ scores that incorrectly

484   indicate convergence (i.e., are greater than zero) stems from the D_{max} measurement (as defined

by Stayton 2015A and calculated in versions 1.0 through 1.3 of the *convevol R* package), which

486   can be erroneously inflated, especially when lineages are morphological outliers. D_{max} can be

measured between ancestral nodes (e.g., see the illustration in Figure 1A), between tips, or

488   between a node and a tip (which is the case in all examples in Figure 2). For converging

lineages, D_{max} is expected to be longer than D_{tip} (Stayton 2015A). For diverging lineages, in

490   contrast, D_{max} is expected to be the morphological distance between the tips, meaning that

D_{max} equals D_{tip} (and $C1 = 0$). However, this is not always the case; divergent lineages can have a

492   D_{max} length that is not between tips, as illustrated in Figure 2B. Thus, D_{max} can be greater than

D_{tip} (indicating convergence) even when lineages are divergent. Although we illustrate this issue

494   using diverging phylogenetic tips (Fig. 2B), the problem could also arise if there are internal

nodes that are similarly divergent and outlying in morphospace (and branching lineages from

496   those nodes do not converge on other focal lineages); thus, this issue is not solely due to

        allowing D_{max} to be measured to tips.

        Other measures of convergence

500   To test whether other convergence measures also experience similar issues as those of the C-

        measures, we applied two other 'distance-based' metrics (Wheatsheaf index [Arbuckle et al.

502   2014, Arbuckle and Minter 2015] and θ [Castiglione et al. 2019]) and OU model-fitting analyses

        to a subset of simulated datasets (Table 1).

504   *Distance-based convergence measures*. Consistent with the C-measures, the $θ_{real}$

        results (standardized to phylogenetic distance between clades) indicate greater convergence in

506   morphological outliers (Table 1). That is, the angle between phenotypic vectors, $θ_{real}$, decreases

        when lineages evolve toward optima that are farther from the ancestral morphology. This is

508   unsurprising because a relatively farther trait optimum results in greater trait values in the

        lineages simulated to be convergent, and, all other variables being equal, greater trait values

510   should result in smaller angles between phenotypic vectors. However, unlike the C-measures,

        $θ_{real}$ does not consistently identify simulated divergent lineages as convergent (Tables 1 and S),

512   although many 'constrained divergence' simulations (~40%) still returned significant results. In

        contrast to the C-measures and standardized $θ_{real}$, the Wheatsheaf index measures less

514   convergence in outliers relative to non-outliers; values decrease when convergent taxa are

        farther from the ancestral morphology in morphospace.

516   *Evolutionary model-fitting analyses*. When 'gliders' are simulated to be morphologically

        convergent, multiple-regime OU models (with 'gliders' assigned to one regime) are the best-

518   fitting models (Table 2), correctly suggesting that the 'glider' lineages are convergently evolving

toward a shared adaptive peak (Fig. 1B). However, we obtain similar results when 'gliders' are

520   simulated to be divergent – multiple-regime OU models outperform simpler models even

though the 'gliders' are not evolving toward a shared adaptive peak (Table 2). The following

522   text explains these results in greater detail.

544 **Table 2.** Evolutionary model-fitting analyses in which multivariate models (via functions in the *mvMORPH R* package) were fitted to all six simulated traits and univariate models (via

546 functions in the *OUwie R* package) were fitted to PC1 scores. Model-fitting results for each trait optimum are the mean AICcWs of 15 simulated datasets. Model support for the two-

548 regime models (mvOU2 and all OU2 models) could be interpreted as evidence for convergence because these are modeling evolution of focal lineages toward a shared

550 adaptive peak (although see the Results & Discussion for cautionary notes on this assumption). The 'constrained divergence' simulations limited the simulated traits of focal

552 taxa ('gliders') to be positive values, thus constraining them to one region of morphospace (see Methods). See the Supplemental Results for 'unconstrained divergence' results, and

554 see the Methods for descriptions of the models. Abbreviations: AICcW, small-sample corrected Akaike weights; mvBM, multivariate Brownian motion model; mvOU, multivariate

556 Ornstein-Uhlenbeck model.

Simulations		Model	Trait optimum of focal taxa				
			0	20	40	60	80
Multivariate (via *mvMORPH*)	**Convergence**	mvBM1	1.000	0.000	0.000	0.000	0.000
		mvOU1	0.000	0.000	0.000	0.000	0.000
		mvOU2	0.000	1.000	1.000	1.000	1.000
	Divergence ('constrained')	mvBM1	–	–	0.000	0.000	0.000
		mvOU1	–	–	0.000	0.000	0.000
		mvOU2	–	–	1.000	1.000	1.000
Univariate (via *OUwie*)	**Convergence**	BM1	0.087	0.051	0.000	0.000	0.000
		OU1	0.032	0.022	0.000	0.000	0.000
		OU2	0.045	0.023	0.000	0.000	0.000
		OU2A	0.024	0.010	0.001	0.000	0.000
		OU2V	0.038	0.040	0.000	0.000	0.000
		OU2VA	0.773	0.854	0.999	1.000	1.000
	Divergence ('constrained')	BM1	–	–	0.000	0.000	0.000
		OU1	–	–	0.000	0.000	0.000
		OU2	–	–	0.000	0.000	0.000
		OU2A	–	–	0.105	0.079	0.214
		OU2V	–	–	0.046	0.000	0.000
		OU2VA	–	–	0.849	0.921	0.786

       Our simulations and analyses mimic an empirical scenario in which researchers test an *a*

560   *priori* hypothesis that specific focal taxa (in this case, mammalian 'glider' species; *sensu*

Grossnickle et al. 2020) are evolving toward, and possibly converging on, an adaptive peak. We

562   have assigned the 'gliders' to one selective regime and the 'non-gliders' to a second regime,

and then use model support for multiple-regime OU models as evidence for convergence on

564   adaptive peaks (Fig. 1B). In this type of analysis, support for multiple-regime OU models could

suggest that the focal lineages (assigned to a single regime) have convergently shifted toward a

566   shared adaptive peak for that regime. This procedure is prevalent in the literature (e.g., Collar

et al. 2014, Friedman et al. 2016, Baliga and Mehta 2018, Grossnickle et al. 2020, Rincon-

568   Sandoval et al. 2020, Law 2022), although other studies assess convergence without *a priori*

hypotheses for lineages using alternative model-fitting procedures (e.g., using functions in the

570   *SURFACE R* package; Ingram and Mahler 2013), which is not examined here (see Adams and

Collyer 2018).

572   The simulated datasets used in model-fitting analyses include lineages that were

evolved by both BM ('non-gliders') and by a mix of OU and BM processes ('gliders'; two traits

574   BM-evolved and four traits OU-evolved), meaning that we do not expect any of the tested

multivariate *mvMORPH* models (mvBM1, mvOU1, mvEB, mvOU2) to be good fits to the data

576   because they do not include a mix of BM and OU processes. Nonetheless, we fit these models

because the multivariate models enable us to incorporate all data into the analyses. Conversely,

578   the univariate *OUwie* models are expected to be relatively better fits to the data because they

allow parameters to vary between regimes, but they are only fit to PC1 scores and thus do not

580   model all data.

For the convergence simulations, the univariate OU model-fitting analyses, which

582   include models with varying rate and attraction parameters between regimes, correctly identify

convergence of focal lineages at any optima values (i.e., at any positions in morphospace): the

584   OU2VA model is the best-fitting model to PC1 scores and it allows the selection and rate

parameters to vary between regimes (Table 2). The results correctly identify different

586   underlying evolutionary processes for the two regimes, with 'gliders' showing stronger

evidence of convergence via an OU process (i.e., greater selection and rate parameters

588   compared to those of the non-glider regime). However, for the multivariate model-fitting

analyses in which rate and attraction parameters are kept constant between regimes, the

590   analyses struggle to identify convergence when focal lineages converge on a morphology that is

similar to the ancestral morphology (i.e., in the middle of morphospace), with mvBM1 out-

592   performing mvOU2 (Table 2). The results correctly identify different underlying evolutionary

processes for the two regimes, with 'gliders' showing stronger evidence of convergence via an

594   OU process (i.e., greater selection and rate parameters compared to those of the non-glider

regime). Thus, for robust analyses of convergence via model-fitting analyses, we recommend

596   that researchers include models that allow all model parameters to vary among regimes.

Our results for 'constrained divergence' simulations highlight a potential pitfall of the

598   assumption that model support for multiple-regime OU models is indicative of convergence

(Fig. 1B). Contrary to this assumption, the two-regime OU models (both multivariate and

600   univariate) outperformed simpler, single-regime models when fit to data in which lineages of

both regimes ('gliders' and 'non-gliders') are divergent (via the 'constrained divergence'

602   simulations; Table 2). Although the divergent focal lineages ('gliders') are partially constrained

to one region of morphospace in the simulations (see Methods and further discussion below),

604   they still show considerable divergence in phylomorphospace (Fig. 3C) rather than convergent

or parallel attraction toward a shared optimum (a presumptive adaptive peak), which is an

606   assumption of the OU process (Fig. 1B). Further, taxa representing the second regime ('non-

gliders') were evolved by BM, not an OU process. Thus, taxa of neither selective regime are

608   expected to be well-fit by by an OU model, and yet the two-regime OU model is a substantially

better fit to the data than the null, BM1 or mvBM1 model in both the univariate and

610   multivariate analyses (Table 2). For the univariate analyses, the best-fitting model is OU2VA,

which allows parameters to vary between 'glider' and 'non-glider' regimes – this is the model

612   that is most appropriate considering the generating processes of the simulations (i.e., divergent

BM and OU processes). However, OU2VA is still a multiple-regime OU model, and its

614   performance could be interpreted as evidence for the presence of two adaptive peaks, one for

'gliders' and one for 'non-gliders,' even though lineages of each regime were not simulated as

616   evolving toward single, regime-specific adaptive peaks. This result is consistent with the model-

fitting results of Collar et al. (2014) for eel morphology – durophagous eels appear to be

618   divergent in phylomorphospace, but two-regime OU models (with durophagous taxa in one

regime and non-durophagous taxa in a second regime) were the best-fitting models.

620   That the two-regime OU model is the best fit to our divergence data, despite the

expectation that none of our fitted models fit the simulated data, simply reflects that two-

622   regime OU models are the best-fitting of bad-fitting models. Knowing the underlying processes

that generated the simulated data, we recognize that the multiple-regime OU models are not a

624   good representation of the underlying data. However, we reiterate that we are mimicking

specific empirical scenarios in which researchers use *a priori* regime assignments (e.g., based on

626   gliding behavior) and do not know the evolutionary processes that generated their trait data.

Our results here highlight how researchers using OU model-fitting analyses in the same manner

628   should be cautious when interpreting results of model-fitting analyses – if multiple-regime OU

models are best-fitting to data, it does not necessarily indicate that focal lineages experienced

630   convergence.

The use of OU model-fitting analyses to test for convergence can offer a valuable

632   complement to distance-based metrics, but researchers should be cautious both when

choosing models to fit to data and when interpreting results, especially considering that

634   empirical datasets often include complex evolutionary patterns and small sample sizes for some

regimes (Ho and Ané 2014, Cooper et al. 2016, Grabowski et al. 2023). Although not explored in

636   this study, multiple-regime BM (BMM) models may offer alternative options that complement

multiple-regime OU models (e.g., Law et al. 2018, Grossnickle et al. 2020, Martinez et al. 2020,

638   Rincon-Sandoval et al. 2020). BMM models allow varying phylogenetic means among regimes

and can be fit using functions within some *R* packages, including *mvMORPH* (Clavel et al. 2015).

640   Because BMM models do not model selection toward an optima, support for BMM models over

OUM models may suggest that there is limited or no convergence among lineages of interest

642   (e.g., Grossnickle et al. 2020), and in some cases a BMM model might serve as a more

appropriate null model than BM1.

644   An additional OU model-fitting approach involves testing for convergent shifts toward

        shared adaptive peaks without *a priori* regime classifications for focal lineages (e.g., using

646   functions in the *SURFACE R* package; Ingram and Mahler 2013). We did not apply this method in

        part because results are challenging to interpret across many simulated datasets, especially

648   when peak shifts are likely to be identified in taxa that evolved via BM (Adams and Collyer

        2018), and because the method may be less appropriate than our method (with *a priori* regime

650   classifications for 'gliders' and 'non-gliders') for testing the specific hypothesis of this study.

        Nonetheless, this method provides a valuable research tool for examining convergence,

652   especially for analyses that are exploratory (i.e., without hypotheses for specific lineages), and

        we recommend applying it to empirical datasets as a supplement to other convergence

654   measures (e.g., Grossnickle et al. 2020, Rincon-Sandoval et al. 2020).

        Our goal with divergence simulations was to mimic empirical samples with lineages that

656   might share some similarities and thus be tested for convergence. Therefore, in our primary

        divergence simulations (i.e., 'constrained divergence'), the divergent focal lineages ('gliders')

658   were partially constrained to one region of morphospace by limiting the randomized optima to

        be positive values for the four OU-evolved divergent traits (the other two traits were evolved

660   by BM and could be negative; see Methods). This methodological decision helps to explain the

        relatively strong fits of the two-regime OU models to the simulated divergence datasets.

662   Because 'glider' lineages are shifting in similar directions (e.g., all divergent lineages are in

        negative PC2 space in Figure 3B), the 'gliders' may be modeled as evolving toward a broad

664   adaptive peak that occupies a large region of one side of morphospace, especially because an

        OU process does not require that independent lineages shifting toward a peak come from more

666   dissimilar ancestors. Figure 2B provides a conceptual illustration of this scenario; the focal

       lineages ('maroon') are diverging but still evolved toward a broad adaptive peak, which is

668   broader than the adaptive peak of the ancestral ('blue') lineages. This could be a similar

       scenario to the divergent outlier lineages in our simulated dataset (Fig. 3C). The durophagous

670   and non-durophagous eels examined by Collar et al. (2014) are best fit by a two-peak model,

       and they may provide a possible empirical example of this scenario; durophagous eels are

672   morphologically divergent but appear to be 'constrained' to one region of morphospace, a

       scenario similar to our simulations.

674   For 'unconstrained divergence' simulations, we allowed the 'gliders' to diverge in all

       directions in morphospace, and all convergence measures, including OU model-fitting analyses,

676   correctly identify the taxa as being divergent (Supplemental Results; Table S2). Thus, the central

       issue highlighted here for OU model-fitting analyses (i.e., that multiple-regime OU models are

678   best-fitting to divergent lineages) is likely limited to scenarios in which divergent focal taxa

       share some morphological similarities.

680   *Summary of issues with convergence measures.* We found that all convergence

       measures, including both distance-based and model-fitting approaches, correctly identified

682   convergence among lineages simulated to be convergent (Tables 1 and 2, Figs. 3C and 5A).

       However, we uncovered a critical issue with *C*-measures: they often misidentify divergent

684   lineages as being convergent (Fig. 2B, Fig. 3C, Table 1). In addition, all convergence measures

       appear to be influenced by the positions of focal taxa in morphospace. *C*-measures, θ_{real}, and

686   some OU model-fitting analyses (those in which rate and attraction parameters are not allowed

       to vary between regimes) all result in stronger measures of convergence when simulated

688   convergent taxa evolve toward relatively farther trait optima. In contrast, the Wheatsheaf index

shows weaker convergence in outliers (Table 1). Further, model support for multiple-regime OU

690   models, which is often interpreted as support for convergent shifts toward shared adaptive

peaks, should be considered with caution because in some scenarios these models may be the

692   best fits to divergent lineages (Table 2). To help address this issue, we recommend that

researchers fit a broad range of models that allow parameters to vary among regimes and

694   perform model-fitting analyses in conjunction with distance-based convergence measures.

696   **Measuring convergence through time via *Ct*-measures**

Despite any shortcomings, *C*-measures have benefits over other convergence measures,

698   including the ability to distinguish between convergence and conservatism (Stayton 2015A).

Thus, our objective is not to discourage the use of distance-based metrics like *C*-measures but

700   rather to identify issues and encourage the development of improved measures.

We help to address the *C*-measure issues by presenting novel distance-based

702   convergence measures that are derived from the *C*-measures. The new measures are calculated

using the same equations as those for *C*1–*C*4 (except with a change to *C*4; see below and

704   Supplemental Methods), but we limit the candidate D_{max} measurements to distances between

lineages at synchronous 'time slices' coinciding with internal phylogenetic nodes (Fig. 4). We

706   refer to the new measures as *Ct*-measures (or *Ct*1–*Ct*4) and D_{max} as $D_{max.t}$ because time (*t*) is

incorporated when measuring morphological distances between lineages, unlike the *C*-

708   measures. *Ct*1 scores can be interpreted in the same way as *C*1 scores were intended to be

interpreted (Stayton 2015A): positive *Ct*1 scores represent a proportion of the maximum

710   morphological distance between lineages that has been closed by convergent evolution, with a

value of one representing complete convergence. Like *C*-measures, statistical significance for

712   *Ct*-measures is based on comparison with expectations for evolution proceeding entirely on a

BM model, with simulations used to generate the expectations.

Figure 4. Conceptual illustration of our new $Ct1$ convergence measure, which is calculated

726   like $C1$ of Stayton (2015A) but candidate $D_{max.t}$ measurements are limited to 'time slices' at

internal phylogenetic nodes. The plots on the right show the three candidate $D_{max.t}$

728   measurements and the distance between lineages at the tips (D_{tip}). The scenarios in A and B

are the same as those in Figures 2A and 2B, respectively. In contrast to $C1$, $Ct1$ correctly

730   identifies divergence (negative score) in the scenario in B. Although the $Ct1$ score is greater

when lineages are outliers (A), note that the $Ct1$ scores (0.3 for non-outliers and 0.4 for

732   outliers) are more similar to each than are the $C1$ scores in the same scenarios (0.4 and 0.7;

Fig. 2A), indicating that Ct-measures are less influenced by positions of taxa in morphospace

734   compared to C-measures.

By limiting the candidate $D_{max.t}$ measurements to time slices, the Ct-measures minimize

the possibility of $D_{max.t}$ being erroneously inflated by divergent tips. This is conceptually

illustrated in Figures 4A and 4B, which show the same scenarios as in Figure 2A and 2B.

Whereas the $C1$ score in Figure 2B incorrectly indicates convergence (i.e., $C1$ is greater than

zero), the $Ct1$ score in Figure 4B correctly indicates divergence (i.e., the value is negative; unlike

the C-measures, the Ct-measures allow divergence results to be negative).

Unlike the D_{max} measurement, the D_{tip} measurement has not been altered from its

original implementation in C-measures (Stayton 2015A) and is not limited to a synchronous

time slice, thus allowing for distances between tips to be compared even if the tips vary in

geologic age (e.g., comparison of an extinct taxon and an extant taxon). However, unlike the C-

measures, the Ct-measures do not allow $D_{max.t}$ to be measured between tips (i.e., $D_{max.t}$ cannot

equal D_{tip}). This means that divergent taxa will have negative Ct scores, whereas C-measures (as

they were initially intended) will measure divergent taxa as having scores of zero (i.e., D_{max}

equals D_{tip}). See the Supplemental Methods for more information on the Ct-measures.

In addition to developing the Ct-measures, we added several new features to the

convevol R package (Stayton 2018). This includes allowing C and Ct-measures to compare clades

that contain multiple lineages, whereas the original routines were limited to comparisons of

individual lineages (see Methods; Stayton 2015A). Clade comparisons are enabled by 1)

excluding pairwise comparisons between within-clade lineages (e.g., two flying squirrel species)

and 2) weighting of C and Ct scores and p-values based on the number of pairwise comparisons

between focal clades (see Supplemental Methods). Further, C and Ct-measures can be

measured using single traits (the original routines only permitted measures of multivariate

distances, although they were adapted for univariate analyses in some studies; Spear and

762   Williams 2020, Law 2022), and we updated the $C4$ (and $Ct4$) calculation to better match the

original description of that measure. See the Supplemental Methods for additional information

764   on these updates. We used an R script written by Jonathan S. Mitchell and published in Zelditch

et al. (2017) as a foundation for the updated functions. The run times for the revised R

766   functions (*convrat.t* and *convratsig.t*) are approximately ten times faster than the original

functions of Stayton (2015A) when using our simulated datasets. We did not revise $C5$, which is

768   a frequency-based convergence measure that tallies the number of times lineages enter a

region of morphospace (Stayton 2015A), because it is not influenced by the issues highlighted

770   here.

        We have also developed a new R function, *plot.C*, that produces a plot of the distances

772   between lineages through time. This type of plot is conceptually illustrated in Figure 4, and

Figure 5B includes a phylogeny and plot produced by the *plot.C* function for an example dataset

774   from our convergence simulations, showing pairwise distances between three 'glider' lineages.

An additional example output of *plot.C* is provided in Figure S7 for the 'twig' ecomorphotype

776   lineages of anoles, although we separated convergent and non-convergent pairwise

comparisons for ease of interpretation. These plots allow researchers to visualize when the

778   measured $D_{max.t}$ occurred during the evolutionary history of the lineages, and they may be

useful for applications beyond studies of convergence. The candidate $D_{max.t}$ measurements at

780   time slices are provided as an output of the *calcConvCt* function.

Figure 5. (*A*) *Ct*1 scores for simulated convergent lineages (top results in plot) and divergent

784   lineages (bottom) under varying evolutionary scenarios. See the Methods and Figure 3

caption for more information. Although in some cases the divergence *Ct*1 results are greater

786   than zero (indicating convergence), these results were not statistically significant when we

calculated simulation-based *p*-values for a subset of datasets (Table 1). *C*1 results are means

788   and standard errors of 15 simulated datasets. (*B*) An example output from the *plot.C R*

function that shows the pairwise distances between lineages with time. Note that although

790   only three 'glider' lineages are highlighted in the plot, five lineages were used for *Ct1*

measurements.

We tested the performance of *Ct*-measures by applying them to the simulated datasets

796   using the same methodology as that for *C*-measures. Unlike the *C*-measures (Fig. 3C), the *Ct*-

measures do not consistently misidentify divergent lineages as being convergent (Figs. 5A and

798   S3); most of the simulated divergence datasets (via both drift/BM and selection/OU) exhibit *Ct*1

scores that are negative, correctly indicating divergence. Although in some cases the Ct1 score

800   is greater than zero when applied to divergence datasets (incorrectly indicating convergence;

Fig. 5A), these Ct1 scores are not statistically significant (Table 1), which contrasts with the

802   strongly significant C1 scores for divergence simulations. Further, the greater-than-zero Ct1

results could be due in part to convergence occasionally occurring by chance in our simulated-

804   divergence datasets (e.g., BM-evolved 'glider' lineages evolving toward each other by chance).

This indicates the importance of researchers considering the p-values associated with Ct-

806   measures when evaluating convergence in their samples.

Although both C1 and Ct1 have a maximum value of one (complete convergence), the

808   two measures scale differently and thus their values should not be directly compared. For

instance, C1 values for 'constrained' divergence simulations are approximately 3.0 (Fig. 3C),

810   whereas Ct1 values for the same divergence simulations are approximately 0.0 (Fig. 5A). Thus,

positive C1 values reflect both convergence and some degree of divergence (due to the D_{max}

812   issue highlighted in this study), and their values are expected to be greater than those of Ct1 for

the same scenario. Further, limiting $D_{max.t}$ to specific time slices may dampen the convergence

814   signal (see discussion on this topic below).

816   **Relatively greater convergence in morphological outliers**

Like C-measures and θ_{real}, the Ct-measures often measure more convergence when focal

818   lineages are morphological outliers (Figs. 4A, 5A, and S8). This pattern appears to be more

pronounced in C-measures compared to Ct-measures (Figs. 2A, 3C, and S8), likely due to the C-

820   measure issues highlighted in this study (Fig. 2). Further, this trend may occur in Ct-measures

(despite the *C*-measure-related issues being eliminated) in part because values can be inflated

822   when focal taxa have different origination ages, and this issue is expected to be more

pronounced in morphological outliers than non-outliers (see Supplemental Results and Figure

824   S5). To help address this issue for *Ct*-measures, we added an optional feature to the *calcConvCt*

function that limits candidate $D_{max.t}$ measurements to the time prior to the evolution of the

826   focal lineages (e.g., prior to the evolution of the earliest glider clade). We recommend that

researchers use this option as a supplement to regular *Ct*-measures when their clades of

828   interest have very different origination ages (see Supplemental Results).

The trend of greater convergence in outliers largely disappears for *C*- and *Ct*-measures

830   when all six simulated traits are OU-evolved to trait optima (i.e., all six traits are convergent) –

the *C*1 and *Ct*1 values remain relatively constant at any optimum values (Figs. 3C and 5A).

832   Conversely, the trend is more distinct when a greater number of traits are BM-evolved. This

indicates that the pattern of greater convergence in outliers is influenced by the types of

834   simulated traits (BM-evolved or OU-evolved to optima) and not just the aforementioned issues

with *C*- and *Ct*-measures. We believe that trait type influences this pattern because the relative

836   amount of morphological evolution in OU-evolved traits (vs BM-evolved traits) varies

depending on the value of trait optima. For instance, when OU-evolved traits are evolved

838   toward optima near the ancestral morphology, they may undergo less morphological change

relative to the BM-evolved traits. Thus, there is a relatively greater influence of BM-evolved

840   traits (i.e., divergent traits) on *C*- and *Ct*-measures at smaller optima values, resulting in weaker

convergence signals via *C*- and *Ct*-measures. Conversely, OU-evolved traits (i.e., convergent

842   traits) of outliers have undergone relatively greater morphological change, evolving farther

from the ancestral morphology, and therefore the OU-evolved traits have a relatively larger

844   influence on *C*- and *Ct*-measures. This pattern is likely to reflect empirical scenarios because the

strength of natural selection is expected to vary among traits (e.g., Grossnickle et al. 2020).

846   Thus, when different traits of taxa have undergone varying amounts of adaptive morphological

change, researchers should expect to observe relatively stronger evidence of convergence in

848   outliers when using most convergence measures.

850   **Empirical examples – *C*1 vs *Ct*1**

The *C*-measure issues highlighted here are relevant to the many studies that have employed (or

852   will employ) the *C*-measures. In many cases, erroneous *C*-measure results may have led

researchers to either infer convergence in lineages that are divergent or infer inflated degrees

854   of convergence. For instance, Grossnickle et al. (2020) tested for convergence among gliding

mammal lineages using limb measurements and observed conflicting results. Statistically

856   significant *C*-measure scores indicated strong convergence, but other analyses (i.e.,

evolutionary model-fitting, morphological disparity, phylomorphospace trajectories) suggested

858   parallel evolutionary patterns. The authors concluded that the conflicting lines of evidence

indicated weak, incomplete convergence. But considering the issues highlighted here, the

860   strong *C*-measure results in Grossnickle et al. (2020) are probably misleading. For instance, the

C-measure scores were likely inflated due to the outlying morphologies of some gliders (e.g.,

862   dermopterans) because outliers produce greater *C* scores (Figs. 2A and 3C), meaning that the

gliders are probably less convergent than the authors concluded. We re-analyzed the data from

864   Grossnickle et al. (2020) using the *Ct*-measures, and in contrast to strong *C*-measure scores, we

found that all glider comparisons have negative $Ct1$ scores, indicating divergence instead of

866   convergence. In some instances, the $Ct1$ scores are only slightly negative and have significant p-

values (e.g., $Ct1$ = -0.01 and $p < 0.01$ for the comparison of scaly-tailed squirrels and flying

868   squirrels), which is congruent with the other lines of evidence examined in Grossnickle et al.

(2020) that suggested parallel evolutionary changes rather than convergence for most glider

870   groups.

Huie et al. (2021) and Stayton (2015A; using data from Mahler et al. 2013)

872   independently analyzed *Anolis* lizard morphologies using distinct datasets, and both found that

the ecomorphotypes with the greatest $C1$ scores are those in the outermost regions of

874   morphospace ('crown-giant,' 'grass-bush,' and 'twig'; see Figure 3 of Huie et al. 2021). The $C1$

values for these ecomorphotypes ranged from 0.31 to 0.43 in these studies, whereas other,

876   non-outlying ecomorphotypes had $C1$ values ranging from 0.09 to 0.25 (Stayton 2015A, Huie et

al. 2021). The relatively large $C1$ scores of outlying ecomorphotypes, in addition to the positive

878    $C1$ scores for all pairwise comparisons, may be due in part to the biases in the C-measures. We

evaluated this possibility by applying Ct-measures to one of the outlying ecomorphotypes

880   ('twig') from the anole dataset of Mahler et al. (2013; ten standardized measurements of body

dimensions, e.g., tail length). We found that, although the overall $Ct1$ score was statistically

882   significant, it was near zero (Table S1), in contrast to the $C1$ score being 0.36 (Stayton 2015A).

Interestingly, there was considerable disparity in the pairwise Ct results for the five twig

884   lineages, with $Ct1$ scores ranging from 0.346 (*A. paternus* vs. *A. valencienni*) to -0.763 (*A.*

occultus vs. *A. paternus*) and six of ten pairwise comparisons not significant. (See Figure S7 and

886   Table S1 for full results and plotted pairwise distances through time.) Thus, these results

highlight not only the issues with the *C*-measures, namely the inflation of *C* scores among

888   outliers, but also the importance of considering pairwise comparisons when evaluating

convergence among multiple focal lineages.

Ct-measures – recommendations and limitations

892   In contrast to *C*-measures, the *Ct*-measures are influenced by the timing of evolutionary change

because they limit candidate $D_{max.t}$ measurements to specific time slices. This feature should be

894   considered by researchers who apply the *Ct*-measures because it may alter expectations about

the degree of measured convergence. For instance, if different focal lineages evolve toward a

896   specific morphology (or adaptive peak) at different points in time, then the $D_{max.t}$ measurement

may not measure the morphologically farthest distances between the lineages, possibly

898   resulting in lower-than-expected *Ct* scores. Conversely, and as noted above, if the putatively

convergent taxa evolve toward outlying regions of morphospace, then the asynchronous origins

900   of the clades could inflate the *Ct*-measures (Supplemental Results; Fig. S5). To help mitigate this

issue, we recommend that researchers generate and assess phylomorphospace and distances-

902   between-lineages-through-time plots, and compare default *Ct* results to those generated when

using the alternative option of the *calcConvCt* function that limits candidate $D_{max.t}$

904   measurements to the period in which lineages of interest overlap in time (see Supplemental

Methods).

906   The *Ct*-measures can be applied to taxa of varying geologic ages (assuming there is an

internal node for a $D_{max.t}$ measurement), but they may perform poorly when the tips of focal

908   taxa are very different in geologic age because candidate $D_{max.t}$ measurements are restricted to

the period in which the evolutionary histories of the lineages overlap in time. For instance, the

910   evolutionary histories of ichthyosaurs and dolphins overlap from their most recent common

ancestor (MRCA; early amniotes) to the ichthyosaur tips, providing considerable time (more

912   than 200 million years) for candidate $D_{max.t}$ measurements. However, there are no candidate

$D_{max.t}$ measurements between the extinction of ichthyosaurs and extant dolphins, and if this is

914   the period in which the ancestral dolphin lineages shows the greatest divergence from

ichthyosaurs, then $D_{max.t}$ will not capture the maximum divergence of the lineages. That is,

916   much of the evolutionary history of dolphins (and placental mammals more broadly) would be

excluded by Ct-measures, likely resulting in smaller-than-expected Ct scores. (Note, however,

918   that D_{tip} ignores time and would measure the morphological distance between ichthyosaur and

dolphin tips.) In these types of cases, alternative methods for measuring convergence (e.g., OU

920   model-fitting analyses) may be more appropriate.

The restriction of candidate $D_{max.t}$ measurements to coincide with internal nodes

922   exacerbates an issue inherent to many phylogenetic comparative methods: the reliance on

inferred ancestral states. $D_{max.t}$ is the critical value that enables the Ct-measures to diagnose

924   convergence, and it is drawn entirely from ancestral state data, which are estimated from tip

values assuming a BM model of evolution. The consequence is that ancestral reconstructions

926   are likely to reflect average morphologies of the sampled taxa, decreasing the chance of

measuring convergence via the Ct-measures because $D_{max.t}$ estimates may be artificially shorter

928   than the 'real' $D_{max.t}$ values. For instance, despite simulating extremely strong convergence on a

trait optimum for all six traits, the greatest Ct1 scores are around 0.7, indicating that about 70%

930   of $D_{max.t}$ has been closed by convergent evolution. Based on the simulation methods, we

expected $Ct1$ values to be closer to 1.0. We tested this issue by applying $Ct1$ (and $C1$) measures

932 to simulations in which we used the generated ('true') ancestral traits. $Ct1$ scores are

consistently greater when 'true' ancestral states are used in calculations (Fig. S8), helping to

934 confirm that the convergence signal of Ct-measures might often be diluted due to the

assumption of a BM model of evolution. Note that because C-measures rely on the same

936 ancestral state reconstructions to infer morphology at internal nodes, this issue is not unique to

the Ct-measures (Fig. S8).

938 The aforementioned issue is likely to be exacerbated under conditions where there are

relatively few intervening nodes between putatively convergent lineages (i.e., there is a small

940 sample of candidate $D_{max.t}$ measurements), when those putatively convergent lineages are

subtended by long branches (i.e., distances from which to draw $D_{max.t}$ are biased toward deeper

942 nodes), and when only contemporary tips are included (i.e., there is a lack of fossil data

informing reconstructions at internal nodes). Therefore, the Ct-measures may be most

944 appropriate for well-sampled study systems that include a substantial number of internal nodes

and relatively few long branches, and researchers should include fossil taxa whenever possible

946 to improve ancestral reconstructions at internal nodes.

The number of phenotypic traits used to assess convergence is likely of increased

948 importance when using Ct-measures. In multivariate datasets, some traits may be convergent

and others non-convergent (i.e., divergent, parallel, or conservative). While including a greater

950 number of non-convergent traits in analyses is expected to decrease the overall convergence

signal of any convergence measure, it may also exacerbate the Ct-related issues raised in this

952 section. In general, adding traits increases the measured distances between tips and internal

nodes. However, ancestral inference via BM tends to average variation at internal nodes; thus,

954   D_{tip} typically increases at a higher rate than $D_{max.t}$ for each non-convergent trait that is added to

a dataset. This pattern is illustrated in Figure S6, highlighting that increasing the number of BM-

956   evolved traits (which are expected to be mostly divergent) in simulations results in relatively

greater increases of D_{tip} scores compared to $D_{max.t}$ scores. Therefore, an increased number of

958   traits in analyses (with all else equal) could result in a relative decrease in Ct scores compared

to datasets with fewer traits, unless the additional traits are strongly convergent. We

960   recommend that researchers carefully choose traits based on the specific hypothesis being

tested, and analyze individual traits or subsets of traits whenever feasible to tease apart unique

962   patterns among traits.

964   **Summary**

The *C*-measures are a popular means of identifying and quantifying phenotypic convergence, in

966   part because they can differentiate between convergence and conservatism. However, we

highlight a critical issue with *C*-measures: they can misidentify divergent lineages as convergent

968   (Figs. 2 and 3, Table 1). To help address this issue, we developed improved convergence

measures (*Ct*-measures) that quantify distances between lineages at time slices at internal

970   phylogenetic nodes, minimizing the possibility of divergent taxa being mistakenly measured as

convergent. We have also developed new features (available in the *convevol R* package), such

972   as a function to produce distances-between-lineages-through-time plots and the ability to

compare clades that include multiple taxa. Although *Ct*-measures improve on *C*-measures,

974   researchers should recognize their limitations. For instance, *Ct*-measures may be unreliable if

convergent evolutionary change is asynchronous between lineages of interest (e.g., fossils of

976   very different geologic ages), especially when lineages are morphological outliers. More

broadly, we find that multiple methods (including *Ct*-measures) are biased by the location of

978   taxa in morphospace, with most methods measuring greater convergence in morphological

outliers. OU model-fitting analyses are a valuable tool for investigating convergence, but we

980   urge caution when applying and interpreting the results of the analyses. Model support for

multiple-regime OU models is not always indicative of convergent shifts toward shared

982   adaptive peaks, especially when all of the fitted models are poor fits to the data (Table 2).

Because all available methods for identifying and measuring phenotypic convergence are

984   imperfect, we recommend that researchers use multiple methods. Researchers should also

recognize the benefits and drawbacks of their chosen methods when interpreting results, and

986   they should rely on methods that are most appropriate for their dataset and the hypotheses

being tested.

ACKNOWLEDGEMENTS

For helpful discussion and feedback, we thank two anonymous reviewers, Miriam Zelditch,

Gregory Wilson Mantilla, Eli Amson, Kenneth-Dieter Benton, Allison Bormet, Richard Meyn,

Sharlene Santana, and members of the 'Tempo & Mode in Isolation' discussion group. Jonathan

S. Mitchell wrote the *R* script (published in Zelditch et al. 2017) that we adapted for the

updated *convevol* functions, and we greatly appreciate his permission to use the code. Funding

for D. M. G. was provided by National Science Foundation (NSF) grants DBI-1812126 and IOS-

2017738. C. J. L. was funded by an Early Career Provost Fellowship at the University of Texas at

Austin, the NSF (DBI–1906248 and DBI–2128146), the Gerstner Family Foundation, the Richard Gilder Graduate School, and the Department of Mammalogy at the American Museum of Natural History. L. N. W. was funded by the NSF Graduate Research Fellowship and NSF grant EAR-PF 2052992.

LITERATURE CITED

992   Adams, D. C., and M. L. Collyer. 2018. Multivariate phylogenetic comparative methods: Evaluations, comparisons, and recommendations. Syst. Biol. 67:14-31.

994   Akaike, H. 1974. A new look at the statistical model identification. IEEE Transactions on Automatic Control 19:716–723.

996   Alfieri, F., L. Botton-Divet, J. A. Nyakatura, and E. Amson. 2022. Integrative approach uncovers new patterns of ecomorphological convergence in slow arboreal xenarthrans. J. Mamm.

998   Evol. 29:283–312.

Arbour, V. M., and L. E. Zanno. 2020. Tail Weaponry in Ankylosaurs and Glyptodonts: An

1000   Example of a Rare but Strongly Convergent Phenotype. Anat. Rec. 303:988–998.

Arbuckle, K., C. M. Bennett, and M. P. Speed. 2014. A simple measure of the strength of

1002   convergent evolution. Methods Ecol. Evol. 5:685–693.

Arbuckle, K., and Minter, A. 2015. Windex: Analyzing convergent evolution using the

1004   Wheatsheaf index in R. Evol. Bioinform. 11:EBO-S20968.

Baliga, V. B., and R. S. Mehta. 2018. Phylo-allometric analyses showcase the interplay between

1006   life-history patterns and phenotypic convergence in cleaner wrasses. Am. Nat. 191:E129-E143.

1008   Bastide, P., C. Ané, S. Robin, and M. Mariadassou. 2018. Inference of Adaptive Shifts for

Multivariate Correlated Traits. Syst. Biol. 67:662–680.

1010   Beaulieu, J. M., D.-C. Jhwueng, C. Boettiger, and B. C. O'Meara. 2012. Modeling stabilizing

selection: Expanding the Ornstein-Uhlenbeck model of adaptive evolution. Evolution

1012   66:2369–2383.

Bennion, R. F., J. A. MacLaren, J. A., E. J. Coombs, F. G. Marx, O. Lambert, and V. Fischer. 2022.

1014   Convergence and constraint in the cranial evolution of mosasaurid reptiles and early

cetaceans. Paleobiology (First View, https://doi.org/10.1017/pab.2022.27).

1016   Butler, M. A., and A. A. King. 2004. Phylogenetic Comparative Analysis: A Modeling Approach

for Adaptive Evolution. Am. Nat. 164:683–695.

1018   Button, D. J., and L. E. Zanno. 2020. Repeated evolution of divergent modes of herbivory in non-

avian dinosaurs. Curr. Biol. 30:158–168.

1020   Baumgart, S. L., P. C. Sereno, and M. W. Westneat. 2021. Wing shape in waterbirds:

morphometric patterns associated with behavior, habitat, migration, and phylogenetic

1022   convergence. Integr. Org. Biol. 3:obab011.

Canale, J. I., Apesteguía, S., Gallina, P.A., Mitchell, J., Smith, N.D., Cullen, T.M., Shinya, A.,

1024   Haluza, A., Gianechini, F.A., Makovicky, P.J. 2022. New giant carnivorous dinosaur reveals

convergent evolutionary trends in theropod arm reduction. Curr. Biol. 32:3195–3202.

1026   Castiglione, S., G. Tesone, M. Piccolo, M. Melchionna, A. Mondanaro, C. Serio, M. De Febbraro,

and P. Raia. 2018. A new method for testing evolutionary rate variation and shifts in

1028   phenotypic evolution. Methods. Ecol. Evol. 9:974–983.

Castiglione, S., C. Serio, D. Tamagnini, M. Melchionna, A. Mondanaro, M. De Febbraro, A.

1030   Profico, P. Piras, F. Barattolo, P. Raia. 2019. A new, fast method to search for morphological

convergence with shape data. PLOS One 16:e0252264.

1032   Chen, M., and G. P. Wilson. 2015. A multivariate approach to infer locomotor modes in

Mesozoic mammals. Paleobiology 41:280–312.

1034   Clavel, J., G. Escarguel, and G. Merceron. 2015. mvMORPH: an R package for fitting multivariate

evolutionary models to morphometric data. Methods Ecol. Evol. 6:1311–1319.

1036   Collar, D. C., J. S. Reece, M. E. Alfaro, P. C. Wainwright, and R. S. Mehta. 2014. Imperfect

morphological convergence: variable changes in cranial structures underlie transitions to

1038   durophagy in moray eels. Am. Nat. 183:E168–E184.

Cooper, N., Thomas, G. H., Venditti, C., Meade, A., & Freckleton, R. P. (2016). A cautionary note

1040   on the use of Ornstein Uhlenbeck models in macroevolutionary studies. Biol. J. Linn. Soc.

Lond. 118:64–77.

1042   Da Silva, F. O., A. C. Fabre, Y. Savriama, J. Ollonen, K. Mahlow, A. Herrel, J. Müller, and N. Di-Poï,

N. 2018. The ecological origins of snakes as revealed by skull evolution. Nat. Commun. 9:1–

1044   11.

Darwin, C. 1859. *On the Origin of Species by Means of Natural Selection*. John Murray, London.

1046   Drummond, A. J., M. A. Suchard, D. Xie, and A. Rambaut, 2012. Bayesian phylogenetics with

BEAUti and the BEAST 1.7. Mol. Biol. Evol. 29:1969–1973.

1048   Friedman, S. T., S. A. Price, A. S. Hoey, and P. C. Wainwright. 2016. Ecomorphological

convergence in planktivorous surgeonfishes. J. Evol. Biol. 29:965–978.

1050   Grabowksi, et al. 2023. A cautionary note on "A cautionary note on the use of Ornstein

        Uhlenbeck models in macroevolutionary studies". Syst. Biol. syad012,

1052   https://doi.org/10.1093/sysbio/syad012.

        Grossnickle, D. M. 2020. Feeding ecology has a stronger evolutionary influence on functional

1054   morphology than on body mass in mammals. Evolution 74:610–628.

        Hansen, T. F. 1997. Stabilizing selection and the comparative analysis of adaptation. Evolution

1056   51:1341–1351.

        Ho, L. S. T., and Ané, C. 2014. Intrinsic inference difficulties for trait evolution with Ornstein-

1058   Uhlenbeck models. Methods Ecol. Evol. 5:1133-1146.

        Huie, J. M., I. Prates, R. C. Bell, and K. de Queiroz. 2021. Convergent patterns of adaptive

1060   radiation between island and mainland *Anolis* lizards. Biol. J. Linn. Soc. Lond. 134:85–110.

        Hurvich, C. M., and C. L. Tsai. 1989. Regression and time series model selection in small

1062   samples. Biometrika 76:297–307.

        Ingram, T., and D. L. Mahler. 2013. SURFACE: detecting convergent evolution from comparative

1064   data by fitting Ornstein-Uhlenbeck models with stepwise Akaike Information Criterion.

        Methods Ecol. Evol. 4:416–425.

1066   Khabbazian, M., Kriebel, R., Rohe, K. and Ané, C. 2016. Fast and accurate detection of

        evolutionary shifts in Ornstein-Uhlenbeck models. Methods Ecol. Evol. 7:811–824.

1068   Law, C. J., Slater G. J., and Mehta, R. S. 2018. Shared extremes by ectotherms and endotherms:

        Body elongation in mustelids is associated with small size and reduced limbs. Evolution.

1070   73:735–749.

Law, C. J. 2022. Different evolutionary pathways lead to incomplete convergence of elongate

1072   body shapes in carnivoran mammals. Syst. Biol. 71:788-796.

Losos, J. B. 2011. Convergence, adaptation, and constraint. Evolution 65:1827–1840.

1074   Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence on the

macroevolutionary landscape in island lizard radiations. Science 341:292–295.

1076   Mahler, D. L., M. G. Weber, C. E. Wagner, and T. Ingram. 2017. Pattern and process in the

comparative study of convergent evolution. Am. Nat. 190:S13–S28.

1078   Martinez, Q., J. Clavel, J. A. Esselstyn, A. S. Achmadi, C. Grohé, N. Pirot, and P. H. Fabre. 2020.

Convergent evolution of olfactory and thermoregulatory capacities in small amphibious

1080   mammals. Proc. Natl. Acad. Sci. 117:8958–8965.

McLean, B. S., K. M. Helgen, H. T. Goodwin, and J. A. Cook. 2018. Trait-specific processes of

1082   convergence and conservatism shape ecomorphological evolution in ground-dwelling

squirrels. Evolution 72:473–489.

1084   Pevsner, S. K., D. M. Grossnickle, and Z. X. Luo. 2022. The functional diversity of marsupial limbs

is influenced by both ecology and developmental constraint. Biol. J. Linn. Soc. Lond.

1086   135:569–585.

Polly, P. D. 2019. Phylogenetics for Mathematica. Version 6.5. Department of Earth and

1088   Atmospheric Sciences, Indiana University: Bloomington, Indiana.

https://pollylab.indiana.edu/software.html.

1090   R Core Team 2020. R: A language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing.

1092   Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

        things). Methods Ecol. Evol. 2:217–223.

1094   Rincon-Sandoval, et al. (2020). Evolutionary determinism and convergence associated with

        water-column transitions in marine fishes. Proc. Natl. Acad. Sci. 117:33396–33403.

1096   Rovinsky, D. S., A. R. Evans, and J. W. Adams. 2021. Functional ecological convergence between

        the thylacine and small prey-focused canids. BMC Ecol. Evol. 21:1–17.

1098   Serio, C., P. Raia, and C. Meloro. 2020. Locomotory adaptations in 3D humerus geometry of

        Xenarthra: testing for convergence. Front. Ecol. Evol. 8:139.

1100   Spear, J. K., and S. A. Williams. 2020. Mosaic patterns of homoplasy accompany the parallel

        evolution of suspensory adaptations in the forelimb of tree sloths (Folivora: Xenarthra).

1102   Zool. J. Linn. Soc. zlaa154.

        Speed, M. P., and K. Arbuckle. 2017. Quantification provides a conceptual basis for convergent

1104   evolution. Biol. Rev. Camb. Philos. Soc. 92:815–829.

        Stayton, C. T. 2008. Is convergence surprising? An examination of the frequency of convergence

1106   in simulated datasets. J. Theor. Biol. 252:1–14.

        Stayton, C. T. 2015A. The definition, recognition, and interpretation of convergent evolution,

1108   and two new measures for quantifying and assessing the significance of convergence.

        Evolution 69:2140–2153.

1110   Stayton, C. T. 2015B. What does convergent evolution mean? The interpretation of

        convergence and its implications in the search for limits to evolution. Interface Focus

1112   5:20150039.

Stayton C. T. 2018. *Convevol: quantifies and assesses the significance of convergent evolution.* R

1114   package version 1.3. https:cran.r-project.org/package=convevol.

Tamagnini, D., C. Meloro, P. Raia, and L. Maiorano. 2021. Testing the occurrence of

1116   convergence in the craniomandibular shape evolution of living carnivorans. Evolution

75:1738–1752.

1118 Upham, N., J. A. Esselstyn, and W. Jetz. 2019. Inferring the mammal tree: species-level sets of

phylogenies for questions in ecology, evolution, and conservation. PLOS Biol. 17:e3000494.

1120 Uyeda, J. C. and L. J. Harmon. 2014. A novel Bayesian method for inferring and interpreting the

dynamics of adaptive landscapes from phylogenetic comparative data. Syst. Biol. 63:902–

1122   918.

Uyeda, J. C., D. S. Caetano, and M. W. Pennell. et al. 2015. Comparative analysis of principal

1124   components can be misleading. Syst. Biol. 64:677-689.

Weaver, L. N., and D. M. Grossnickle. 2020. Functional diversity of small-mammal postcrania is

1126   linked to both substrate preference and body size. Curr. Zool. 66:539–553.

Zelditch, M. L., J. Ye, J. S. Mitchell, and D. L. Swiderski. 2017. Rare ecomorphological

1128   convergence on a complex adaptive landscape: Body size and diet mediate evolution of jaw

shape in squirrels (Sciuridae). Evolution 71:633–649.