Spatial Transcriptomics of Intraductal Papillary Mucinous Neoplasms of The Pancreas Identifies NKX6-2 Expression as a Driver of Gastric Differentiation and Indolent Biological Potential

Marta Sans[1,2], Yuki Makino[1,2], Jimin Min[1,2], Kimal I. Rajapakshe[1,2], Michele Yip-Schneider[3], C Max Schmidt[3], Mark W. Hurd[2], Jared K. Burks[4], Javier A. Gomez[4], Fredrik Thege[1,2], Johannes Fahrmann[5], Robert A. Wolff[6], Michael P. Kim[7], Paola A. Guerrero[1,2], Anirban Maitra[1, 2]

[1] Department of Translational Molecular Pathology, The University of Texas MD Anderson Cancer Center, Houston, TX, 77030, USA

[2] Ahmed Center for Pancreatic Cancer Research, The University of Texas MD Anderson Cancer Center, Houston, TX, 77030, USA

[3] Department of Surgery, Indiana University School of Medicine, Indianapolis, IN, 46202, USA

[4] Department of Leukemia, The University of Texas MD Anderson Cancer Center, Houston, TX, 77030, USA

[5] Department of Clinical Cancer Prevention, The University of Texas MD Anderson Cancer Center, Houston, TX, 77030, USA

[6] Department of Gastrointestinal Medical Oncology, University of Texas MD Anderson Cancer Center, Houston, TX, 77030, USA

[7] Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX, 77030, USA

Running Title: NKX6-2 and Gastric IPMNs

Conflict of Interest

A.M. receives royalties for a pancreatic cancer biomarker test from Cosmos Wisdom Biotechnology, and this financial relationship is managed and monitored by the UTMDACC Conflict of Interest Committee. A.M. is also listed as an inventor on a patent that has been licensed by Johns Hopkins University to ThriveEarlier Detection. A.M. serves as a consultant for Freenome and Tezcat Biotechnology. M.K. is a consultant for PanTher Therapeutics.

Funding

A.M. is supported by the MD Anderson Pancreatic Cancer Moon Shot Program, P50CA221707, and U54CA274371. A.M. and R.A.W. are supported by the Sheikh Khalifa Bin Zayed Al-Nahyan Foundation. A.M., M.Y.S, C.M.S, are supported by the U01CA200468. J.F. is supported by the U01CA239522. M.S. is supported by the CPRIT Training Award (RP210028) and the Anne Eastland Spears Endowed Fellowship. This work was also supported by the Pre Cancer Atlas through the MD Anderson Strategic Initiative Development (STRIDE) program. The COMET imaging was performed in the Flow Cytometry & Cellular Imaging Core Facility, which is supported in part by the National Institutes of Health through M. D. Anderson's Cancer Center Support Grant CA016672 and the NCI's Research Specialist 1 R50 CA243707-01A1. Bulk RNA sequencing was conducted by the Advanced Technology Genomics Core (ATCG) at the MD Anderson Cancer Center, funded by the NCI's Core grant CA016672.

Acknowledgements: The authors would like to thank Drs. Vittorio Branchi, Maria Monberg, Sonja Woermann, Cara Haymaker, and Mr. Bret M. Stephens for helpful discussions.

Abstract

Intraductal Papillary Mucinous Neoplasms (IPMNs) of the pancreas are *bona fide* precursor lesions of pancreatic ductal adenocarcinoma (PDAC). The most common subtype of IPMNs harbor a gastric foveolar type lining epithelium, and these low-grade mucinous neoplasms are harbingers of IPMNs with high-grade dysplasia and cancer. The molecular underpinning of gastric differentiation in IPMNs is unknown, although identifying drivers of this indolent phenotype might enable opportunities for intercepting progression to high-grade IPMN and cancer. We conducted spatial transcriptomics on a cohort of patient IPMNs, followed by orthogonal and cross species validation studies, which established the transcription factor NKX6-2 as a key determinant of gastric cell identity in low-grade IPMNs. Loss of *NKX6-2* expression is a consistent feature of IPMN progression, while restitution of *NKX6-2* in murine IPMN lines recapitulates the aforementioned gastric transcriptional program. Our study identifies NKX6-2 as a previously unknown transcription factor driving indolent gastric differentiation in IPMN pathogenesis.

Significance

Improved understanding of the molecular features driving IPMN development and differentiation is critical to prevent cancer progression and enhance risk stratification. Our study employed spatial profiling technologies to characterize the epithelium and microenvironment of IPMN, which revealed a previously unknown link between *NKX6-2* expression and gastric differentiation in IPMNs, the latter associated with an indolent biological potential.

Introduction

Pancreatic Ductal Adenocarcinoma (PDAC) is projected to become the second leading cause of cancer death by the year 2040 (1). Significantly higher survival rates are observed in patients diagnosed with localized (42%) compared to regional or distant disease (14% and 3%, respectively), emphasizing the need for early detection and interception strategies to improve patient outcomes (2). Intraductal Papillary Mucinous Neoplasms (IPMNs) are the most common category of mucin-producing pancreatic cysts and can arise from either the main pancreatic duct, or from one or more of the branch ducts. IPMNs are *bona fide* precursor lesions of PDAC, with ~5-10% of non-invasive cystic lesions progressing to invasive adenocarcinoma (3,4). Based on the degree of atypia, the lining epithelium of IPMNs is classified into either low-grade (LG) or high-grade (HG) dysplasia, with the latter harboring a greater propensity for progression to invasive neoplasia (5). In addition, based on morphological features and parallels to the epithelial cells from other gastrointestinal organs, the lining epithelium of IPMNs has also been categorized into gastric, intestinal, or pancreato-biliary subtypes (6). The gastric subtype is by far the most common subtype of IPMNs, mostly arising within the branch ducts, and is typically associated with LG dysplasia. In contrast, IPMNs of the intestinal and pancreato-biliary subtypes can arise in either the main or branch ducts, and usually correspond to HG IPMNs. The phylogeny of intestinal and pancreato-biliary subtypes continues to be debated, with some studies speculating that gastric IPMNs represent a common precursor to both categories of HG lesions (7,8), and others postulating that gastric and pancreato-biliary IPMNs share a developmental trajectory that is distinct from the intestinal subtype (9,10). Irrespective, the deregulated transcriptional programs that characterize the most common category of IPMNs, the gastric subtype, remain mostly an enigma. Identifying drivers of gastric differentiation within the IPMN epithelium would not only facilitate insights into the pathogenesis of these early cystic lesions but would also provide the basis for cataloging alterations that accompany further progression to HG IPMNs, and eventually cancer.

Newly emergent spatial technologies for transcriptome and protein profiling offer the opportunity to evaluate the levels of gene transcripts and proteins directly from tissue samples while preserving their localization and distribution (11,12). In this study, we utilized the Visium spatial transcriptomics (ST) platform to profile snap-frozen archival samples from a panel of IPMNs with varying grades of dysplasia and epithelial morphology. With this approach, we identified distinct epithelial patterns of gene expression that were associated with IPMN grade and subtype. Using single-cell deconvolution algorithms, the cellular composition of the corresponding IPMN-associated microenvironment was also characterized. Notably, we identified that elevated levels of transcripts encoding the transcription factor *NKX6-2* (NK6 homeobox 2) were localized to the epithelium of LG IPMNs and were associated with a

gene signature recapitulating the gastric pit and isthmus epithelium, which we have termed gastric isthmus-pit (GIP) signature. We validated the GIP signature associated with elevated *NKX6-2* expression using multiple orthogonal approaches, including RNA sequencing performed on archival microdissected IPMN samples and cross-species spatial transcriptomics performed in a genetically engineered model of IPMNs (13,14). Moreover, using a quantitative high-plex platform for protein profiling on archival material (Lunaphore COMET assay), we confirmed that elevated nuclear NKX6-2 was a feature of the gastric subtype of IPMNs, wherein this transcription was co-localized with markers of gastric foveolar mucin (TFF1 and MUC5AC), and with nuclear GATA6, a prototypal marker of so-called "classical" differentiation within pancreatic neoplasms (15). Lastly, the GIP signature was recapitulated with overexpression of NKX6-2 in cells derived from a genetically engineered model of IPMNs, supporting a hitherto unknown role for NKX6-2 in driving a gastric-type transcriptional profile and gastric foveolar differentiation in the most commonly observed subtype of IPMNs.

Results:

Spatially resolved cell-type specific expression profiles of IPMN samples

The archival snap-frozen IPMN cohort analyzed in our series was comprised of seven LG IPMNs, three HG IPMNs, and three IPMN-associated cancers (**Table S1**). Of note, all LG cases in the dataset were categorized as gastric subtype, while HG cases included pancreato-biliary and intestinal subtypes of IPMN. ST analyses on the thirteen specimens yielded spatially resolved expression profiles for up to 8,000 genes per each of the 12,685 spots covering the tissue samples (**Fig. S1**). Utilizing the scMC (16) processing and clustering pipeline, spots were grouped into 14 clusters according to their gene expression profiles (**Fig. 1A, Table S2**). The majority of clusters included spots from a multitude of samples, while others were predominantly contributed by one or two samples (**Fig. S2**). Visualization of the spatial localization of the spots for each of the clusters showed correlations with histological features. Cell-type proportions for each of the spots were predicted using the robust cell type deconvolution (RCTD) algorithm (17), with the overall highest proportions per spot attributed to epithelial cells and fibroblasts. Not unexpectedly, spots enriched in epithelial cells corresponded to IPMN epithelium while spots containing fibroblasts were in areas of high stromal density (**Table S3**). On the other hand, most immune cell types garnered lower percentage scores per spot, which was expected due to the overall sparsity of such cell types within the IPMN microenvironment. As shown in **Fig. 1B**, trends in cell composition were observed for each of the clusters, which also aligned with the histological assignments. For example, clusters 7 and 11 included the highest average epithelial cell percentages, expressing elevated levels of *EPCAM* transcripts, and were found to be localized to the epithelial cells in the IPMN and associated cancer samples (**Fig. 1C**). High proportion of fibroblasts

were found in clusters 3, 9, 13 and 14, which corresponded to stromal tissue areas with high expression of collagen genes (*COL1A1, COL3A1, COL1A2*). Spots located in tissue areas containing lymphocyte aggregates in cluster 2 were comprised of the highest per spot percentages in naïve lymphocyte cells and expressed elevated levels of T and B cell markers, such as *CR2* and *CD52* (**Fig. 1D**). High percentage of plasma cells (up to 65%) were found in clusters 4 and 6, which were derived from the stroma adjacent to IPMN epithelium and showed elevated expression of various immunoglobulin transcripts (i.e., *IGHA1, IGLC2*) (**Fig. 1E**). Overall, unsupervised characterization of the transcriptional profiles obtained from ST analyses from whole tissue samples, together with deconvolution approaches, provided a correlation between gene expression and tissue histology.

Region-specific differential expression of transcripts and cell types associated with IPMN progression

Clinically, HG IPMNs are considered as high-risk lesions that mandate surgical resection, with or without an associated invasive cancer. Therefore, for purposes of further analyses, the three HG IPMNs and three IPMN-associated cancers in our ST cohort were combined into one "high-risk" (HR) category for comparison *versus* the LG IPMNs. To evaluate region-specific changes in gene expression and cell composition between the LG IPMNs *versus* HR IPMNs, we selected spots overlapping with the neoplastic epithelium ("epilesional", n=755); the immediately adjacent microenvironment, corresponding to two layers of spots (~200 μm) surrounding the lining epithelium ("juxtalesional", n=1142); and an additional two layers of spots located further distal to the juxtalesional region ("perilesional", n=1030) (**Fig. 2A**). Comparisons between the transcriptomic profiles of LG IPMN *versus* HR IPMNs resulted in the identification of 946, 705, 272 transcripts that were significantly differentially expressed in the epilesional, juxtalesional, and perilesional areas, respectively, between the two categories of IPMNs (**Table S4-S6, Fig. 2B-D**). Interestingly, expression of genes associated with gastric mucus-secreting cells, such as *MUC6*, *NKX6-2*, *PGC*, *PSCA*, and *TFF2* were found to be elevated in the LG IPMN epithelium when compared to HR IPMN. Similarly, higher expression levels of *IGLC2* and *IGKC*, markers of activated B cells and plasma cells, were detected in the LG IPMN epilesional areas, suggesting infiltration of plasma cells in these lesions. On the other hand, transcripts previously described as upregulated in "conventional" PDAC, such as *CEACAM6* and *CLDN4* were found to be expressed at higher levels in HR IPMN samples when compared to LG IPMN (18,19). Elevated levels of transcripts associated with intestinal differentiation (*CDX2* and *MUC2*) were observed in epilesional regions of an intestinal type IPMN and an invasive "colloid" carcinoma, which arose in the backdrop of an intestinal IPMN (**Fig. S3**). In the juxtalesional region, higher levels of various immunoglobulin transcripts, including those also observed in the epilesional areas (*IGLC2, IGKC*) and others unique to this compartment (*IGHM, IGHG2*) were observed in LG IPMN samples compared to

HR IPMNs, reiterating the role of an active humoral immune response in early *versus* late stage cystic lesions. Transcripts corresponding to cancer associated fibroblasts, like *COL10A1*, were found to be significantly higher expressed in the juxtalesional regions surrounding HR IPMNs, compared to those surrounding LG IPMN. A lower number of differentially expressed genes were present when comparing the perilesional areas, suggesting that changes in the microenvironment between LG and HR IPMN are more prominent in the areas closest to the lesions.

Utilizing the cell-type proportions assigned by RCTD, we then evaluated the cell composition in the three selected compartments in our IPMN cohort. As expected, higher proportion of fibroblast and immune cells were assigned in the jutxalesional and perilesional regions compared to the epilesional areas, but immune infiltration was still predicted in the latter compartment (**Fig. 3A**). Trends in cell proportions were observed when comparing the three regions between LG IPMN *versus* HR IPMN, even though large variance was observed between spots and patients, suggesting heterogeneous spatial distribution of cell types (**Table S7, Fig. S4A**). Of note, significantly higher proportions of plasma and mast cells were assigned in the jutxalesional areas of LG IPMN compared to HR IPMNs (**Fig. 3B**). By RCTD, juxtalesional spots were predicted to contain an average of 10% plasma cells in LG IPMN cells, while an average of 5% was observed in HR IPMN samples, with a single case (HG_2) contributing to most of the plasma cell counts (**Fig. 3C**). Spots with high levels of plasma cells were also characterized by expression of immunoglobulin genes, such as *IGKC* (**Fig. S4B**). Mast cells were predicted to comprise as much as 10% of the cells in the juxtalesional spots, suggesting a significant infiltration of such cells in the IPMN microenvironment (**Fig. 3B, D**). Spots with mast cell contributions showed high levels of *TPSAB1*, a transcript encoding for mast-cell specific tryptase (**Fig. S4C**).

To further evaluate the presence of plasma and mast cells in the IPMN microenvironment, we quantified the expression of CD27 (plasma cells) and TPSAB1 (mast cells) in the IPMN-adjacent microenvironment by cyclic IF (COMET) in an independent cohort of archival formal-fixed paraffin embedded (FFPE) IPMNs (**Fig. 3E, Table S8**). The COMET analysis confirmed that a higher percentage of plasma cells in LG *versus* HG IPMN was observed, validating the difference observed on ST, while similar mast cell percentages were observed in the adjacent microenvironment from both LG and HG IPMNs (**Table S9**). Nonetheless, when comparing the adjacent microenvironment between synchronous LG and HG IPMN areas in the same tissue section, a higher percentage of both plasma and mast cells was observed surrounding the LG (11.2% and 4.4%, respectively) compared to the HG lesions (6.0% and 1.1%, respectively) (**Fig. S5A**). Utilizing the additional markers in the COMET panel, other immune cell phenotypes, including macrophages, T cells, and B cells were quantified in the

adjacent microenvironment. Although we observed trends between LG and HG IPMN samples with similar directionality as ST analysis, none of these comparisons were found to be statistically significant (**Table S9, Fig. S5B**). Thus, given the broad heterogeneity in cell type composition of the juxtalesional microenvironment observed in the IPMN lesions, which has been previously described by others (20), we decided to focus on further dissecting the epilesional compartment of LG and HG IPMN lesions to identify potential markers of grade and subtype.

Elevated epithelial NKX6-2 expression is associated with gastric differentiation in IPMNs

While the gastric subtype is the most commonly observed variant of IPMNs in the pancreas, the putative transcriptional drivers underlying this pattern of differentiation are not known. In our ST dataset, all LG IPMNs were uniformly comprised of the gastric subtype, while HG IPMNs were either intestinal or pancreato-biliary, and thus the terms "LG" and "gastric" are used interchangeably. As previously noted, expression of genes associated with gastric mucus-secreting cells were found to be elevated in the epilesional regions of LG IPMN *versus* HR IPMNs (**Fig. 2B-C**). Of these, *MUC6* was the highest differentially expressed transcript, which was not surprising given the encoded apomucin is a prototype gastric mucin. The second highest differentially expressed gene in the epilesional regions of LG IPMNs versus HR IPMNs was *NKX6-2* (**Fig. 2B, Fig. 4A-B**), which encodes for a homeobox domain containing transcription factor. *NKX6-2* is expressed in pancreatic endocrine cells, but not in normal ductal cells, and is known to promote beta-cell differentiation (21,22). *NKX6-2* is also highly expressed in human stomach, particularly in isthmus and pit cells (23,24). Moreover, a homolog to *NKX6-2*, *Nkx2-1,* has been implicated in regulating gastric differentiation in the lung (25). To validate NKX6-2 expression in IPMN, we performed quantitative analysis of nuclear NKX6-2 expression using the COMET platform on an independent cohort of archival IPMN samples, which also demonstrated a significantly higher percentage of nuclei staining positive for NKX6-2 in LG (all gastric subtype) compared to HG IPMN samples, which included both intestinal and pancreato-biliary subtypes (**Fig. 4C-D, Table S8, Table S10**). Notably, in cases with synchronous LG and HG IPMN foci in the same section, a higher proportion of NKX6-2-expressing nuclei were observed in the LG nuclei (25%) *versus* HG IPMN (1%) (**Fig. 4E**). Agreements between *NKX6-2* levels determined by ST and the percent of nuclei positive for NKX6-2 in the COMET images was also observed for the single patient with tissue analyzed by both methods (**Fig. S6**). Despite the overall significantly higher levels of expression of this transcription factor in LG IPMNs compared to HG IPMNs, we nonetheless observed inter- and intra-patient heterogeneity in expression of both *NKX6-2* transcripts and protein within the LG IPMN epithelium. In instances of reduced expression of nuclear NKX6-2 in LG IPMN, this pattern was usually localized to transitory

states from gastric to pancreato-biliary IPMN, underscoring that loss of this transcription factor might be an accompaniment of progression (**Fig. S6**).

To interrogate molecular changes associated with *NKX6-2* expression, epilesional spots in the ST dataset corresponding to the LG IPMN epithelium were separated based on *NKX6-2* levels into "low", "mid", and "high", and differential expression analyses were performed (**Fig. S7A-B, Table S11**). Importantly, the three categories included spots from almost all samples, allowing us to perform both inter and intra-patient comparisons. Higher levels of transcripts comprising a gastric isthmus-pit (GIP) signature were detected in epilesional spots of high and mid versus low NKX6-2 spots, including *TFF1, LEFTY1, CST3, S100P, TSPAN8, PLA2G10, MUC5AC,* and *GKN2*, among others (**Fig. 4F, S7C**). On the other hand, higher expression of *CXCL2, CD74, CLDN4, LY6E, and SSP1*, many of which have been associated with PDAC progression and neoplastic progression of IPMN (18,26-28), was observed in epilesional spots with minimal *NKX6-2* expression (**Fig. 4F, S7D)**. Expression of transcripts associated with the classical subtype of PDAC, such as *GATA6*, was also higher in epilesional spots with high *NKX6-2* expression (15,29). Moreover, GATA6 and NKX6-2 protein co-localization was observed in the nuclei of LG IPMN using the COMET platform, with overall significantly higher median intensity of GATA6 observed in NKX6-2 positive versus negative nuclei (**Fig. 4G-H, Table S12**). On the contrary, LG epilesional spots with low *NKX6-2* expression had lower levels of classical-like genes, and a percentage of those (~25%) expressed prototypal markers of the basal subtype, including *KRT17 (30)* (**Fig. S7E**). Gene set enrichment analysis (GSEA) revealed increased expression of genes associated with epithelial-to-mesenchymal transition (EMT) in areas with loss of NKX6-2 (**Fig. S7F-G**).

Additional GSEA analyses using cell type signatures showed enrichment of gene sets obtained from gastric pit and isthmus cells from previously published single cell RNA sequencing datasets of human normal stomach (**Fig. 5A, Table S13**) in epilesional spots expressing *NKX6-2* (high and mid) compared to spots with low expression in the LG IPMN ST dataset (24). These results further corroborated the association of *NKX6-2* expression with the GIP signature. Gene sets corresponding to gastric REG3A positive cells, fetal pancreas acinar cells, and pancreas ductal cells were enriched in the transcriptional profiles obtained from spots with minimal NKX6-2 expression (**Fig. S8A**). To further evaluate tissue level expression of NKX6-2 together with markers of the GIP signature, we interrogated TFF1 and MUC5AC co-expression with NKX6-2 using the COMET cyclic IF panel, and observed localization of both markers in the cytoplasm of LG IPMN epithelium, preferentially in areas with NKX6-2 positively stained nuclei (**Fig. 5B**). Quantification of per sample protein expression in the IPMN epithelium showed

moderately strong positive correlations between percentage of positive NKX6-2 nuclei with MUC5AC (cor = 0.54, p-val = 0.02) and TFF1 mean intensities (cor = 0.55, p-val = 0.02) (**Table S12**).

To validate whether *NKX6-2* expression in IPMNs correlates with a gastric transcriptional program, we analyzed an independent dataset of RNA sequencing obtained following laser-capture microdissection (LCM) of 23 archival IPMNs, conducted as part of the NCI Cancer Moonshot PreCancer Atlas Pilot Project (PCAPP) (14). Remarkably, majority of the transcripts that comprise the aforementioned GIP signature (including *TFF1*, *GKN1*, *GKN2*, and *MUC5AC*, amongst several others), as well as the prototypal classical-like transcript *GATA6* were also positively correlated with *NKX6-2* expression in the PCAPP dataset (**Fig. 5C, Fig. S8B**). GSEA also revealed enrichment of the gastric pit and isthmus gene sets with *NKX6-2* expression in the PCAPP cohort similar to what was observed in the ST dataset (**Fig. 5D**). Thus, in two independent datasets of archival IPMNs profiled by disparate approaches (ST and RNA-sequencing), elevated *NKX6-2* was consistently associated with the GIP signature.

Status of NKX6-2 expression in IPMNs correlates with alterations in regulatory networks

We applied SCENIC (31,32) to evaluate regulatory networks that might be associated with *NKX6-2* expression utilizing the ST dataset. Using this approach, 6,266 genes were identified as targets of the *NKX6-2* transcription factor. The targets identified included 119 out of the 134 genes selected as significantly higher expressed in *NKX6-2* positive *versus NKX6-2* negative epilesional spots in LG IPMN from the ST analyses, with markers of the GIP signature, including *TFF1, CAPN8, CA2, SULT1C2*, *S100P*, and *PLA2G10,* selected among the targets receiving the highest weights (**Table S14**). We then applied *RcisTarget* to identify "regulons", defined as gene modules that are co-regulated by a main transcription factor (**Table S15**). We identified the GATA6 regulon presenting the highest positive fold change in activity between *NKX6-2* positive versus *NKX6-2* negative spots, reiterating the link between NKX6-2 and classical differentiation. *FOXA3* was among the other regulons with the highest score positively correlated with NKX6-2 expression (**Fig. 5E**). Higher levels of the *FOXA3* transcriptional network have been reported in the pancreatic progenitor subtype of PDAC, which expresses a foregut-related differentiation program, compared to the more mesenchymal/squamous or basal subtypes (33). Interestingly, Foxa1 and Foxa2, homologs to Foxa3, have been found to physically interact with lineage specific homeobox transcription factors, such as Nkx2-1 in the pulmonary epithelium, to drive a gastric differentiation program in lung carcinomas (25,34). *KLF*-related regulons, such as *KLF4*, *KLF3*, and *KLF2*, were also observed at higher activities in *NKX6-2* expressing spots. Gastric epithelial proliferation and differentiation, including that of pit cells, is largely controlled by *KLF4*, which provides

essential functions to regulate normal gastric epithelial homeostasis (35). Of note, *NKX6-2* itself is predicted as a target for many of the transcription factors of the regulons with higher activity in *NKX6-2+* areas, such as *FOXA3*, *NR2F6*, *GATA6*, and *KLF4* (**Table S14**).

Among the highest scored regulons negatively associated with *NKX6-2* expression, *ONECUT-2 and SPDEF* regulons were assigned higher activities in spots with no NKX6-2 expression. Both have been linked to PDAC pathogenesis via the non-cystic pancreatic intraepithelial neoplasia (PanIN) pathway and their overexpression is associated with adverse prognosis in invasive cancers (36,37). SPDEF is of particular interest because it has recently been implicated in the development of PDAC with classical differentiation and mucin production, two features that are also observed with elevated NKX6-2 in LG IPMNs (38). Nonetheless, the anti-correlation between NKX6-2 and SPDEF levels suggests that this transcription factor "switch" might be one of the prerequisites for progression of indolent LG lesions to higher grade lesions and cancer.

Cross species validation of recapitulation of a gastric transcriptional program by Nkx6-2

To evaluate *Nkx6-2* expression in a murine model of IPMN, we applied the Visium technology to analyze cystic lesions collected from two *Kras;Gnas* mice. Data was processed and 7 cluster assignments were identified (**Fig. S9A**). Based on gene expression and histologic assessment of the areas covered by the spots, clusters 0, 1, 3, and 6 were selected for further analyses of the epithelial compartments from pancreatic murine lesions (**Fig. S9B, Table S16**). Spots in these clusters were then grouped by *Nkx6-2* expression (*Nkx6-2* positive *versus Nkx6-2* negative) (**Fig. 6A**). Notably, higher expression of *Gkn1, Gkn2, Mucl3, Tff1, Tff2, and Muc5ac,* among others, which are markers of the NKX6-2-associated GIP signature identified in human IPMN samples, was observed in areas with *Nkx6-2* expression compared to those without detectable expression in the murine samples (**Fig. 6B, Table S17**). Gene set analyses of the differentially expressed transcripts between *Nkx6-2* positive *versus Nkx6-2* negative spots also showed enrichment of the human gastric pit-like cell profiles in areas with *Nkx6-2* expression (**Fig. S10A**). Expression of markers included in the GIP signature, such as *Tff1* and *Muc5ac*, have also been recently reported from single cell RNA sequencing analyses of murine pancreatic intraepithelial neoplasia (PanIN) cells in *Kras*[G12D] induced acinar-to-ductal metaplasia models (36,37). On the other hand, *Reg1* and *Reg3b* levels were higher in areas with no detectable *Nkx6-2* expression. Genes encoding for pro-inflammatory proteins, such as *Cd74*, *C3*, and *C4b* were also found to be elevated in *Nkx6-2*-negative spots. Enrichment of these features have also been observed at later stages of progression in both acinar-derived (*Reg* genes) and ductal-derived cells (*Cd74, C3, C4b*) in *Kras*[G12D] models compared to earlier time points (37). Interestingly, expression of

Cldn3, Igfbp7, Mmp7, and *Ly6a*, which have been proposed as markers of a "pre-tumor" stage in murine lesions (36,37), were all detected at higher levels in areas of no *Nkx6-2* expression. Of note, many of these genes were also found to be negatively correlated to *NKX6-2* expression in the human ST IPMN dataset (**Fig. S10B**).

To investigate gene expression changes that may be driven by *Nkx6-2*, we overexpressed *Nkx6-2* in a cell line isolated from the *Kras;Gnas* mice. The parental cell line, derived from an established murine IPMN-associated cancer, lacks discernible *Nkx6-2* expression. After lentivirus transduction, *Nkx6-2* expression was confirmed by qPCR and western blot analyses, and RNA-seq was performed in triplicate (**Fig. S11A-B**). A total of 7654 genes were found to be significantly differentially expressed by RNA-seq between the *Nkx6-2* overexpressing (Kras;Gnas$^{Nkx6-2+}$) and control lines (**Table S18**). Remarkably, enrichment of gene sets corresponding to gastric isthmus and mature pit cells was also found by GSEA in the Kras;Gnas$^{Nkx6-2+}$ cells, recapitulating our findings on the human ST datasets (**Fig. 6C**). *Gkn2, Tff1, Mucl3, and Gkn1* were the four top ranked genes increased in *Nkx6-2* overexpressing cells. When comparing the differentially expressed transcripts between Kras;Gnas$^{Nkx6-2+}$ and control cell lines to the transcriptomic profile associated with *Nkx6-2* expression in the murine ST dataset, 209 gene transcripts were found to exhibit overlapping trends, including higher levels of markers of the GIP signature (*Muc5ac, Tff1, Psca, Gkn2, Mucl3, Gkn1, Klf4,* and others) with *Nkx6-2* overexpression (**Table S19, Fig. 6D, Fig. S11D**). A substantial overlap (n = 86 genes) was also observed between the GIP signature associated with *NKX6-2* expression in the human ST dataset, and the significantly altered transcripts driven by *Nkx6-2* overexpression in the *Kras;Gnas* cell lines (**Table S20, Fig. S11C,E**). Overall, our cumulative cross species results support the role of *Nkx6-2* as a pivotal transcription factor in driving a gastric-like transcriptional profile in IPMNs, whose sustained expression might be a requirement for the observed indolent biological behavior of most gastric IPMN subtypes.

Discussion

In the current study, ST analysis paired with single cell deconvolution provided novel insights into histologically annotated gene expression patterns in our cohort of patient derived archival IPMN samples. By using a region-specific approach of "epilesional", "juxtalesional" and "perilesional" spots centered on proximity to the neoplastic epithelium, we were able to discern alterations in expression profiles that occur with IPMN progression within the neoplastic cells, as well as the surrounding milieu. While several microenvironment-specific alterations in immune cell composition were observed on ST, the one consistent change we observed on validation studies performed with the COMET cyclic IF platform on an independent cohort of archival IPMNs was a reduction in the infiltration of plasma cells

with progression. This suggests that humoral immunity may be an important mediator of immune surveillance in early stages of multistep IPMN progression, and this function wanes with subsequent progression to HG dysplasia and cancer. As recently shown by Mirlekar and colleagues, the balance between plasma cells and immune suppressive regulatory B cells governs immune surveillance in established PDAC, with our analysis supporting a role for the former even in non-invasive disease (39). Mast cells were also found to be an abundant component of the IPMN microenvironment (varying between 5-10% of the immune cells), albeit the COMET platform did not show a significant attenuation in mast cell numbers with IPMN progression. Mast cells have been implicated in the development and growth of "conventional" PDAC, but their role in cystic precursor lesions, especially in early disease, remains to be explored (40).

In this ST analysis of a panel of IPMNs, we identified the homeobox transcription factor *NKX6-2* to be a pivotal driver of a so-called "gastric isthmus-pit" (GIP) signature in LG IPMNs of the gastric subtype, the most common subtype of mucinous cystic lesion of the pancreas. NKX6-2 protein levels were also quantified by COMET cyclic IF in an independent cohort of IPMN archival samples, showing increased levels of nuclear NKX6-2 in the epithelium of LG IPMN compared to HG IPMN, together with co-expression of GIP markers MUC5AC and TFF1. Leveraging the spot-level and full-transcriptome information provided by Visium, we investigated the transcriptomic profiles associated with *NKX6-2* expression, which resulted in the identification of a highly enriched signature resembling that of gastric isthmus and pit cells of the stomach, which we termed as GIP signature. Even though gastric IPMN is morphologically similar to gastric epithelium, the remarkable similarity between the *NKX6-2* expressing LG gastric IPMN and the gastric stomach isthmus-pit cell transcriptome was striking. The *NKX6-2* associated GIP signature was also observed in an independent RNA sequencing cohort of archival IPMN samples conducted as part of the NCI Cancer Moonshot Precancer Atlas Pilot project(14). Moreover, *Nkx6-2* expression was also observed in IPMN lesions collected from a genetically engineered IPMN model (13), with enrichment of the GIP signature also detected in spots of high *Nkx6-2* expression. To investigate the role of *NKX6-2* in driving the observed gastric IPMN phenotype, we overexpressed *Nkx6-2* in an IPMN-derived cancer cell line generated from the *Kras;Gnas* mouse model. Remarkably, this single alteration alone was able to recapitulate the GIP signature observed with *NKX6-2* in both human and murine IPMN tissue datasets.

Although *NKX6-2* expression is consistently higher in LG IPMNs *versus* HG IPMNs and invasive cancer, including in cases with synchronous lesions of distinct histological grades, we did observe heterogeneity of *NKX6-2* expression within LG IPMNs. Previous single cell RNA sequencing analyses performed by our group have also identified intra-lesion heterogeneity in IPMN, with a minor

subpopulation of epithelial cells in LG IPMN demonstrating transcriptomic overlap with HG IPMNs or even cancer (41). This leads us to hypothesize that while elevated *NKX6-2* expression is an accompaniment of the earliest steps of IPMN development, loss of NKX6-2 within the LG epithelium might be a harbinger of progression to HG IPMN or cancer. The assessment of differentially expressed genes and transcriptional regulatory networks between *NKX6-2* positive and negative spots within LG IPMNs were quite revealing in this regard. For example, epilesional spots with minimal *NKX6-2* expression in LG IPMNs showed higher levels of PDAC-associated markers, such as *CD74*, *CXCL2*, *LY6E*, and *CLDN4*, as well as *KRT17*, a marker of the aggressive basal subtype of PDAC, pointing to a transcriptional reprogramming away from the indolent NKX6-2-expressing epithelium (18,26,27,30). On the other hand, enrichment of the classical-like signature was observed in spots with high NKX6-2 expression, with higher levels of GATA6 staining quantified in NKX6-2-expressing nuclei by cyclic IF (15,29). Similarly, SCENIC analysis for aberrant regulons found that even in LG IPMNs, loss of *NKX6-2* leads to higher predicted activities of ONECUT-2 and SPDEF regulatory networks, both markers of PDAC progression and aggressive biology (36-38). The cumulative assessment from this data leads us to believe that *NKX6-2* serves as an "epithelial rheostat" in IPMN pathogenesis, and its loss acts as a transcription factor switch, driving an otherwise indolent mucin producing epithelium on a path towards invasive neoplasia.

Our study has certain limitations. First, it is important to note that all IPMN cases included in this study were surgically resected and thus, by definition, the natural history of the lesions was disrupted. Therefore, determining the impact of longitudinal alterations in *NKX6-2* expression within the cyst epithelium on clinical outcome was not feasible. In future studies, such effect could be evaluated by potentially monitoring methylation levels of *NKX6-2* DNA in cystic fluid of LG IPMNs that are undergoing surveillance. Loss of *NKX6-2* expression has been previously attributed to DNA methylation events, and detection of methylated *NKX6-2* DNA in biofluids has been proposed as an early detection biomarker for various cancer types, such as in bladder and renal carcinomas (42,43). We are also pursuing experiments on modulating *Nkx6-2* expression in genetically engineered models of IPMNs, however those are outside the scope of the current study. Second, the Visium technology used in this study did not permit the capture of transcripts at single cell resolution, but the application of deconvolution approaches as well as targeted selection of tissue areas allowed us to capitalize on the available histological information to circumvent this challenge. Moreover, the use of the innovative cyclic IF COMET technology enabled the quantification and spatial localization of selected epilesional and juxtalesional markers at the protein level. Finally, an important unanswered question in our study

remains the upstream regulators of *NKX6-2* expression in gastric IPMNs, and whether frequently mutated genomic drivers like *KRAS* or *GNAS* may be playing a role in this regard.

Given the heterogeneity of IPMN lesions in the pancreas, efforts directed at accurately defining the molecular events dictating biological potential and propensity for progression to cancer are paramount for improving their clinical management. The establishment of molecular markers associated with an indolent biology will prevent the overtreatment of otherwise benign lesions, while encouraging increased surveillance and treatment of higher risk lesions. We posit that NKX6-2 is not only a driver of the unique gastric transcriptional profile and histological phenotype that is observed in early IPMNs, but its sustained expression in the IPMN epithelium portends an indolent biological potential in this cystic precursor lesion.

Materials and Methods:

Patient and murine sample collection for ST analyses

Archival snap frozen (**Table S1**) tissue samples from 13 IPMN resections at either MDACC or Indiana University School of Medicine were sectioned, stained with hematoxylin & eosin (H&E) and evaluated by a pancreas pathologist (A.M) to identify areas containing IPMN lesions and cancer. The analyzed samples were comprised of 7 LG IPMN (all gastric), 3 HG IPMN, and 3 PDAC arising in the backdrop of IPMN. Samples were cryo-sectioned onto Visium Spatial Gene Expression slides for ST analysis. For the murine samples, pancreata were collected from p48-Cre; LSL-KrasG12D; Rosa26R-LSL-rtTA-TetO-GnasR201C mice (*Kras*;*Gnas* mice) at 15 and 25 weeks on doxycycline diet. Samples were flash frozen in liquid nitrogen, cryo-sectioned and H&E stained. After sectioning onto the Visium slides, sample processing was performed as described in the Visium 10x Genomics protocol for fresh-frozen tissue samples (15 minutes were used for permealization time). After library preparation, samples were pooled and sequenced using a NextSeq 550 (Ilumina, CA, USA). The sequencing outputs for all Visium datasets were preprocessed using SpaceRanger and mapped to either the latest human or mouse reference genome and the filtered matrices were converted to Seurat objects in R and used for downstream analyses.

ST data analysis

The single-cell data integration method to match and compare (scMC) (16) pipeline was used for initial processing and clustering of the human IPMN ST dataset. To infer single cell composition from the ST dataset, we applied the robust cell type deconvolution (RCTD) algorithm (17) using full-mode

and a previously collected and annotated single-cell RNA sequencing pancreatic cancer dataset as reference (44), which provided cell type proportions for each of the spots. For histologically directed analysis, the Loupe Browser software was used to select spots covering specific areas in the tissue samples. ST data for the two *Kras;Gnas* samples was SCT normalized and merged. Differential expression analyses of ST data were performed using the FindMarkers and FindAllMarkers functions in Seurat, filtering based on Log2FC, minimum percentage of spots expressing that gene, and adjusted p-value (log2FC > 0.25, min.pct > 0.25, p-val_adj < 0.05). GSEA analyses were conducted using the GSEA 4.2.3 software and the ranked features from differential expression. To evaluate changes in cell composition in the three selected compartments, log2FC and p-values of the per-spot cell-type proportions between LG and HG+PDAC samples were calculated. For regulon analyses, we applied the Single-cell rEgulatory Network Inference and Clustering (SCENIC) (32) method to evaluate regulatory networks that might be associated with NKX6-2 expression. *GENIE3* was first used to identify genes co-expressed with transcription factors based on the transcriptional profiles obtained from epilesional areas in the LG IPMN samples (31). These co-expression modules were then evaluated using *RcisTarget* based on cis-regulatory motif analyses to identify regulons. The activity of each regulon was weighted per each spot and compared between the areas with and without detected *NKX6-2* expression. To evaluate differences in regulon activity associated to *NKX6-2* expression, regulons were scored based on the fold change in activity between the two groups and average activity value per group.

COMET cyclic IF protein profiling

We performed immunofluorescence staining of 19 protein markers, which were selected based on transcriptomic changes observed from the ST dataset (**Table S21**), on an independent cohort of 26 archival FFPE IPMN tissue samples (**Table S8**) using the COMET platform (Lunaphore, Switzerland) (*see supporting methods*). Prior to imaging and staining, antigen retrieval was performed using EZ antigen retrieval buffer at 107°C for 15 minutes in an EZ-Retriever Microwave System (BioGenex, CA, USA). Images were imported into Visiopharm software, where IPMN areas were selected, and nuclei and cell segmentation were performed for quantification (*see supporting methods*).

Overexpression of Nkx6-2 in a murine Kras;Gnas cell line

We used a cell line derived from a murine IPMN-associated PDAC in a 12-month-old male *Kras;Gnas* mouse (13) (p48-*Cre*;LSL-Kras^{G12D};Rosa26R-LSL-rtTA-TetO-Gnas^{R201C}, on 0.0060% doxycycline diet). A Nkx6-2 vector: pLV[Exp]-Puro-CMV>mNkx6-2[NM_183248.3](ns)/3xFLAG/Myc:T2A:EGFP or a control vector (pLV[Exp]-EGFP:T2A:Puro-

EF1A>mCherry) (VectorBuilder Inc, IL, USA) were transduced in the cell line, followed by puromycin (Thermo Fisher Scientific, Inc., MA, USA) selection. To obtain protein extracts, cell pellets were lysed by RIPA Lysis Buffer (Rockland Immunochemicals, Philadelphia, PA, USA) containing protease/phosphatase inhibitor cocktail. Electrophoresis was performed to separate proteins in 4%–12% of Bis-Tris gels (Thermo Fisher Scientific, Waltham, MA, USA). The samples were subsequently transferred onto a nitrocellulose membrane. After blocking by 5% dry milk, membranes were incubated with anti-NKX6.2 (1:1000 dilution) (Sigma Aldrich, St. Louis, MO, USA), anti-Myc-Tag (1:1000)(Cell Signaling Technology, Beverly, MA, USA), or HRP-conjugated anti-β-actin (1:200) (Santa Cruz Biotechnology, Dallas, TX, USA) primary antibodies at 4°C overnight. Blots for NKX6.2 and Myc-Tag were thereafter incubated with an HRP-conjugated anti-rabbit secondary antibody (Cell Signaling Technology). Clarity ECL Western Blotting Substrates (Bio-Rad, Hercules, CA) were used to visualize the blots. For RNA collection, GFP-positive cells were sorted from the cells transduced with a Nkx6-2 vector or a control vector using a CytoFLEX SRT (Beckman Coulter Life Sciences, CA, USA). 24 hours after plating, cells were treated with 100 ng/mL doxycycline for 24 hours before RNA collection. Total RNA was extracted using the RNeasy Mini Kit (QIAGEN, Venlo, Limburg, Netherlands), according to the manufacturer's protocol. cDNA was subsequently produced by reverse transcription. Gene expression was measured by real-time RT-PCR using the delta-delta Ct method. Following primers for PCR were purchased from Integrated DNA Technologies (IDT) (Coralville, IA, US); mouse NKX6-2 forward (5'-AAGTCTGCCCCGTCTCAAC-3'), mouse NKX6-2 reverse (5'-GGTCTGCTCGAAAGTCTTCTC-3'), mouse actin forward (5'-CACAGCTTCTTTGCAGCTCCTT-3'), and mouse actin reverse (5'-CGTCATCCATGGCGAACTG-3'). Bulk RNA sequencing was performed in triplicate on a NextSeq500 (Illumina, CA, USA). Sequencing reads were mapped using STAR (45) against the Mus musculus genome (UCSC mm10), and gene level quantification was performed with featureCounts (46) using the mm10 genome transcriptome annotation. Differential expression analysis was performed with DESeq2 (47).

FIGURES

Figure 1. Spatially resolved transcriptomic analyses of IPMN tissue samples. A) Uniform Manifold Approximation and Projection (UMAP) showing clustering of spots based on gene expression profiles for all 13 samples. B) Dot plot including cell type scores per cluster obtained with the robust cell type decomposition (RCTD) algorithm. Color represents the normalized average cell proportion and dot size represents the maximum cell proportion per cluster. Representative spatial maps showing correlation between cell composition and localization provided by RCTD, spatial gene expression, cluster assignments, and histology. C) Mapping of epithelial cells, EPCAM expression, and localization of spots from cluster 11 for a HG IPMN sample. D) Localization and proportion of naïve CD4T and naïve B cells, and visualization of cluster 2 spots from a HG IPMN sample with co-occurring cancer. A tissue area corresponding to a lymphocyte aggregate is outlined and a zoomed-in image is provided. (E) Plasma cells, IGLC2 expression, and cluster 4 shown for a LG IPMN sample. IPMN areas are outlined in black in the optical images from H&E stained tissues.

Figure 2. Histologically directed spot selection reveals area-specific gene expression changes associated with IPMN grade and PDAC. A) Selection of epilesional, juxtalesional, and perilesional areas shown in a LG IPMN sample. B) Heatmaps plotting expression of genes differentially expressed between LG and HR (high-risk) in epilesional (left) and juxtalesional (right) spots. Transcripts exclusively showing differential expression in the jutxalesional and not the epilesional compartments were included in the juxtalesional heatmap. Violin plot showing gene expression levels for selected features for spots in C) epilesional and D) juxtalesional areas.

Figure 3. Spatial organization of the tumor microenvironment in IPMN samples. A) Violin plot of predicted cell proportions per spot grouped by region. B) Violin plot showing per spot proportions of plasma and mast cells grouped by sample code and colored by type. Representative spatial maps highlighting the proportion and localization of epithelial, plasma (C), and mast(D) cells in the IPMN samples. On the right side, spots are colored according to their corresponding region (epilesional, jutxalesional, and perilesional). E) Percentage of cells staining positive for CD27 (left) and TPSAB1(right) in the adjacent microenvironment to the IPMN lesions in the samples analyzed by the COMET imaging system. A dash line connects the LG and HG IPMN lesions that correspond to the same patient and tissue section. F) COMET images showing co-staining of CD27 (top) and TPSAB1 (bottom) with EPCAM and DAPI in LG and HG IPMN samples.

Figure 4. *NKX6-2 expression is elevated in LG IPMN and associated with a distinct transcriptional profile*. A) Spatial maps of *NKX6-2* expression and localization from ST analyses of four representative samples. Optical images for the corresponding H&E images are provided below, with the neoplastic epithelium outlined. B) Violin plot showing *NKX6-2* expression in the IPMN epilesional spots grouped by sample. C) Percentage of NKX6-2 positive nuclei in IPMN epithelium determined by COMET cyclic IF staining (p = 0.0039). D) Localization of NKX6-2 in the nuclei of LG IPMN epithelium showed by co-staining with EPCAM, DAPI, and NKX6-2. An H&E image of the area stained is provided in the bottom left. E) NKX6-2 staining of LG and HG IPMN lesions from the same tissue section showing higher NKX6-2 levels in the LG IPMN nuclei. H&E images of the areas stained are provided below. F) Representative gene transcripts observed at significantly higher levels in spots of high *NKX6-2* (left panel) or low *NKX6-2* levels (right panel) in the ST dataset from LG IPMN. G) Co-

staining of GATA6 and NKX6-2 showing co-expression in the LG IPMN nuclei. H) Median GATA6 intensity obtained from the COMET staining in NKX6-2+ versus NKX6-2- nuclei (p = 0.03).

Figure 5. Enrichment of the GIP signature in *NKX6-2* expressing IPMN. A) Enrichment of gastric pit and isthmus gene sets in the *NKX6-2* associated signature obtained from the ST LG IPMN dataset. B) Representative COMET images showing co-staining of NKX6-2, MUC5AC, and TFF1 in three different LG IPMN samples. C) Heatmap showing expression of *NKX6-2* and the GIP signature in 23 archival IPMN tissue areas from the NCI Cancer Moonshot PCAPP dataset (14). Samples are ranked by *NKX6-2* expression from right to left. D) Dot plot showing normalized enrichment scores and size for gastric pit and isthmus gene sets in both the PCAPP and ST dataset by GSEA. E) Regulon activities scored for *NKX6-2*+ (green) and *NKX6-2*- (light yellow) spots based on SCENIC analyses.

Figure 6. *Nkx6-2* expression in *a genetically engineered model* of IPMN. *ST analyses of murine cystic lesions collected from Kras;Gnas mice.* A) From top to bottom: H&E stained tissue samples. Spots corresponding to clusters of high epithelial contribution were selected and colored according to *Nkx6-2* expression, red = Nkx6-2+, blue = Nkx6-2 –. Zoomed-in images of areas presenting with high percentage of Nkx6-2 + spots. B) Expression of gene transcripts associated with *Nkx6-2* in the Kras;Gnas ST dataset shown in a violin plot. *Enforced expression of Nkx6-2 in a cell line derived from Kras;Gnas mice.* C) Enrichment of the gastric isthmus and pit cell gene sets in the *Nkx6-2* overexpressing cell line (Kras;Gnas^Nkx6-2+). D) Heatmap showing expression of genes differentially expressed in the Kras;Gnas^Nkx6-2+ versus control cell line that were also found to be differentially expressed in *Nkx6-2* + vs *Nkx6-2* – spots in the ST dataset from Kras;Gnas mice. Expression data is provided for three RNA-seq replicates. Note that genes with average raw counts < 10 from were not included in the heatmap.

References

1. Rahib L, Wehner MR, Matrisian LM, Nead KT. Estimated Projection of US Cancer Incidence and Death to 2040. JAMA Netw Open **2021**;4(4):e214708 doi 10.1001/jamanetworkopen.2021.4708.
2. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA Cancer J Clin **2022**;72(1):7-33 doi 10.3322/caac.21708.
3. Singhi AD, Koay EJ, Chari ST, Maitra A. Early Detection of Pancreatic Cancer: Opportunities and Challenges. Gastroenterology **2019**;156(7):2024-40 doi 10.1053/j.gastro.2019.01.259.
4. Tanaka M. Thirty years of experience with intraductal papillary mucinous neoplasm of the pancreas: from discovery to international consensus. Digestion **2014**;90(4):265-72 doi 10.1159/000370111.
5. Basturk O, Hong SM, Wood LD, Adsay NV, Albores-Saavedra J, Biankin AV*, et al.* A Revised Classification System and Recommendations From the Baltimore Consensus Meeting for Neoplastic Precursor Lesions in the Pancreas. Am J Surg Pathol **2015**;39(12):1730-41 doi 10.1097/Pas.0000000000000533.
6. Kloppel G, Basturk O, Schlitter AM, Konukiewitz B, Esposito I. Intraductal neoplasms of the pancreas. Semin Diagn Pathol **2014**;31(6):452-66 doi 10.1053/j.semdp.2014.08.005.
7. Omori Y, Ono Y, Kobayashi T, Motoi F, Karasaki H, Mizukami Y*, et al.* How does intestinal-type intraductal papillary mucinous neoplasm emerge? CDX2 plays a critical role in the process of intestinal differentiation and progression. Virchows Arch **2020**;477(1):21-31 doi 10.1007/s00428-020-02806-8.
8. Noe M, Brosens LAA. Gastric- and intestinal-type IPMN: two of a kind? Virchows Arch **2020**;477(1):17-9 doi 10.1007/s00428-020-02827-3.
9. Liffers ST, Godfrey L, Frohn L, Haeberle L, Yavas A, Vesce R*, et al.* Molecular heterogeneity and commonalities in pancreatic cancer precursors with gastric and intestinal phenotype. Gut **2022** doi 10.1136/gutjnl-2021-326550.
10. Iyer MK, Shi C, Eckhoff AM, Fletcher A, Nussbaum DP, Allen PJ. Spatial Transcriptome Profiling of Intraductal Papillary Mucinous Neoplasms Establishes a Molecular Framework for Risk Stratification. bioRxiv **2022**:2022.09.08.507112 doi 10.1101/2022.09.08.507112.
11. Stahl PL, Salmen F, Vickovic S, Lundmark A, Navarro JF, Magnusson J*, et al.* Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. Science **2016**;353(6294):78-82 doi 10.1126/science.aaf2403.
12. Francisco-Cruz A, Parra ER, Tetzlaff MT, Wistuba II. Multiplex Immunofluorescence Assays. Biomarkers for Immunotherapy of Cancer: Methods and Protocols **2020**;2055:467-95 doi 10.1007/978-1-4939-9773-2_22.
13. Ideno N, Yamaguchi H, Ghosh B, Gupta S, Okumura T, Steffen DJ*, et al.* GNAS(R201C) Induces Pancreatic Cystic Neoplasms in Mice That Express Activated KRAS by Inhibiting YAP1 Signaling. Gastroenterology **2018**;155(5):1593-+ doi 10.1053/j.gastro.2018.08.006.
14. Semaan A, Bernard V, Wong J, Makino Y, Swartzlander D, Rajapakshe KI*, et al.* Integrated Molecular Characterization of Intraductal Papillary Mucinous Neoplasms: An NCI Cancer Moonshot Precancer Atlas Pilot Project. bioRxiv **2022**:2022.10.14.512148 doi 10.1101/2022.10.14.512148.
15. O'Kane GM, Grunwald BT, Jang GH, Masoomian M, Picardo S, Grant RC*, et al.* GATA6 Expression Distinguishes Classical and Basal-like Subtypes in Advanced Pancreatic Cancer. Clin Cancer Res **2020**;26(18):4901-10 doi 10.1158/1078-0432.CCR-19-3724.
16. Zhang L, Nie Q. scMC learns biological variation through the alignment of multiple single-cell genomics datasets. Genome Biol **2021**;22(1):10 doi 10.1186/s13059-020-02238-2.
17. Cable DM, Murray E, Zou LS, Goeva A, Macosko EZ, Chen F*, et al.* Robust decomposition of cell type mixtures in spatial transcriptomics. Nat Biotechnol **2022**;40(4):517-26 doi 10.1038/s41587-021-00830-w.
18. Tsutsumi K, Sato N, Cui L, Mizumoto K, Sadakari Y, Fujita H*, et al.* Expression of claudin-4 (CLDN4) mRNA in intraductal papillary mucinous neoplasms of the pancreas. Mod Pathol **2011**;24(4):533-41 doi 10.1038/modpathol.2010.218.
19. Pandey R, Zhou M, Islam S, Chen B, Barker NK, Langlais P*, et al.* Carcinoembryonic antigen cell adhesion molecule 6 (CEACAM6) in Pancreatic Ductal Adenocarcinoma (PDA): An integrative analysis of a novel therapeutic target. Sci Rep **2019**;9(1):18347 doi 10.1038/s41598-019-54545-9.
20. Hernandez S, Parra ER, Uraoka N, Tang X, Shen Y, Qiao W*, et al.* Diminished Immune Surveillance during Histologic Progression of Intraductal Papillary Mucinous Neoplasms Offers a Therapeutic

Opportunity for Cancer Interception. Clin Cancer Res **2022**;28(9):1938-47 doi 10.1158/1078-0432.CCR-21-2585.

21. Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, *et al.* Proteomics. Tissue-based map of the human proteome. Science **2015**;347(6220):1260419 doi 10.1126/science.1260419.

22. Nelson SB, Schaffer AE, Sander M. The transcription factors Nkx6.1 and Nkx6.2 possess equivalent activities in promoting beta-cell fate specification in Pdx1+ pancreatic progenitor cells. Development **2007**;134(13):2491-500 doi 10.1242/dev.002691.

23. Dai J, Peng T, Yu X. NK6 Homeobox 2 Regulated Gastrokin-2 Suppresses Gastric Cancer Cell Proliferation and Invasion via Akt Signaling Pathway. Cell Biochemistry and Biophysics **2021**;79(1):123-31 doi 10.1007/s12013-020-00948-9.

24. Busslinger GA, Weusten BLA, Bogte A, Begthel H, Brosens LAA, Clevers H. Human gastrointestinal epithelia of the esophagus, stomach, and duodenum resolved at single-cell resolution. Cell Rep **2021**;34(10):108819 doi 10.1016/j.celrep.2021.108819.

25. Snyder EL, Watanabe H, Magendantz M, Hoersch S, Chen TA, Wang DG, *et al.* Nkx2-1 represses a latent gastric differentiation program in lung adenocarcinoma. Mol Cell **2013**;50(2):185-99 doi 10.1016/j.molcel.2013.02.018.

26. Steele CW, Karim SA, Leach JDG, Bailey P, Upstill-Goddard R, Rishi L, *et al.* CXCR2 Inhibition Profoundly Suppresses Metastases and Augments Immunotherapy in Pancreatic Ductal Adenocarcinoma. Cancer Cell **2016**;29(6):832-45 doi 10.1016/j.ccell.2016.04.014.

27. Zhang JF, Tao LY, Yang MW, Xu DP, Jiang SH, Fu XL, *et al.* CD74 promotes perineural invasion of cancer cells and mediates neuroplasticity via the AKT/EGR-1/GDNF axis in pancreatic ductal adenocarcinoma. Cancer Lett **2021**;508:47-58 doi 10.1016/j.canlet.2021.03.016.

28. Asundi J, Crocker L, Tremayne J, Chang P, Sakanaka C, Tanguay J, *et al.* An Antibody-Drug Conjugate Directed against Lymphocyte Antigen 6 Complex, Locus E (LY6E) Provides Robust Tumor Killing in a Wide Range of Solid Tumor Malignancies. Clin Cancer Res **2015**;21(14):3252-62 doi 10.1158/1078-0432.CCR-15-0156.

29. Moffitt RA, Marayati R, Flate EL, Volmar KE, Loeza SG, Hoadley KA, *et al.* Virtual microdissection identifies distinct tumor- and stroma-specific subtypes of pancreatic ductal adenocarcinoma. Nat Genet **2015**;47(10):1168-78 doi 10.1038/ng.3398.

30. Roa-Pena L, Leiton CV, Babu S, Pan CH, Vanner EA, Akalin A, *et al.* Keratin 17 identifies the most lethal molecular subtype of pancreatic cancer. Sci Rep **2019**;9(1):11239 doi 10.1038/s41598-019-47519-4.

31. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring regulatory networks from expression data using tree-based methods. Plos One **2010**;5(9) doi 10.1371/journal.pone.0012776.

32. Aibar S, Gonzalez-Blas CB, Moerman T, Huynh-Thu VA, Imrichova H, Hulselmans G, *et al.* SCENIC: single-cell regulatory network inference and clustering. Nat Methods **2017**;14(11):1083-6 doi 10.1038/nmeth.4463.

33. Bailey P, Chang DK, Nones K, Johns AL, Patch AM, Gingras MC, *et al.* Genomic analyses identify molecular subtypes of pancreatic cancer. Nature **2016**;531(7592):47-52 doi 10.1038/nature16965.

34. Camolotto SA, Pattabiraman S, Mosbruger TL, Jones A, Belova VK, Orstad G, *et al.* FoxA1 and FoxA2 drive gastric differentiation and suppress squamous identity in NKX2-1-negative lung cancer. Elife **2018**;7 doi 10.7554/eLife.38579.

35. Katz JP, Perreault N, Goldstein BG, Actman L, McNally SR, Silberg DG, *et al.* Loss of Klf4 in mice causes altered proliferation and differentiation and precancerous changes in the adult stomach. Gastroenterology **2005**;128(4):935-45 doi 10.1053/j.gastro.2005.02.022.

36. Ma Z, Lytle NK, Chen B, Jyotsana N, Novak SW, Cho CJ, *et al.* Single-Cell Transcriptomics Reveals a Conserved Metaplasia Program in Pancreatic Injury. Gastroenterology **2022**;162(2):604-20 e20 doi 10.1053/j.gastro.2021.10.027.

37. Schlesinger Y, Yosefov-Levi O, Kolodkin-Gal D, Granit RZ, Peters L, Kalifa R, *et al.* Single-cell transcriptomes of pancreatic preinvasive lesions and cancer reveal acinar metaplastic cells' heterogeneity. Nat Commun **2020**;11(1):4516 doi 10.1038/s41467-020-18207-z.

38. Tonelli C, Yordanov GN, Hao Y, Deschênes A, Klingbeil O, Ting H-C, *et al.* SPDEF promotes the classical subtype of pancreatic ductal adenocarcinoma. bioRxiv **2022**:2022.03.18.484951 doi 10.1101/2022.03.18.484951.

39. Mirlekar B, Wang Y, Li SR, Zhou M, Entwistle S, De Buysscher T, *et al.* Balance between immunoregulatory B cells and plasma cells drives pancreatic tumor immunity. Cell Rep Med **2022**;3(9) doi ARTN 100744

10.1016/j.xcrm.2022.100744.

40. Chang DZ, Ma Y, Ji B, Wang H, Deng D, Liu Y, *et al.* Mast cells in tumor microenvironment promotes the in vivo growth of pancreatic ductal adenocarcinoma. Clin Cancer Res **2011**;17(22):7015-23 doi 10.1158/1078-0432.CCR-11-0607.

41. Bernard V, Semaan A, Huang J, San Lucas FA, Mulu FC, Stephens BM, *et al.* Single-Cell Transcriptomics of Pancreatic Cancer Precursors Demonstrates Epithelial and Microenvironmental Heterogeneity as an Early Event in Neoplastic Progression. Clin Cancer Res **2019**;25(7):2194-205 doi 10.1158/1078-0432.CCR-18-1955.

42. Chung W, Bondaruk J, Jelinek J, Lotan Y, Liang SD, Czerniak B, *et al.* Detection of Bladder Cancer Using Novel DNA Methylation Biomarkers in Urine Sediments. Cancer Epidem Biomar **2011**;20(7):1483-91 doi 10.1158/1055-9965.Epi-11-0067.

43. Slater AA, Alokail M, Gentle D, Yao M, Kovacs G, Maher ER, *et al.* DNA methylation profiling distinguishes histological subtypes of renal cell carcinoma. Epigenetics **2013**;8(3):252-67 doi 10.4161/epi.23817.

44. Lee JJ, Bernard V, Semaan A, Monberg ME, Huang J, Stephens BM, *et al.* Elucidation of Tumor-Stromal Heterogeneity and the Ligand-Receptor Interactome by Single-Cell Transcriptomics in Real-world Pancreatic Cancer Biopsies. Clin Cancer Res **2021**;27(21):5912-21 doi 10.1158/1078-0432.CCR-20-3925.

45. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, *et al.* STAR: ultrafast universal RNA-seq aligner. Bioinformatics **2013**;29(1):15-21 doi 10.1093/bioinformatics/bts635.

46. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics **2014**;30(7):923-30 doi 10.1093/bioinformatics/btt656.

47. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol **2014**;15(12):550 doi 10.1186/s13059-014-0550-8.